

AROUND COFIN

ANDRZEJ ROŚLANOWSKI AND SAHARON SHELAH

ABSTRACT. We show the consistency of “there is a nice σ -ideal \mathcal{I} on the reals with $\text{add}(\mathcal{I}) = \aleph_1$ which cannot be represented as the union of a strictly increasing sequence of length ω_1 of σ -subideals”. This answers Borodulin–Nadzieja and Głab [3, Problem 6.2(ii)].

1. INTRODUCTION

Borodulin–Nadzieja and Głab [3] studied generalizations of the Moko-bodzki ideal and they showed that those σ -ideals do not have Borel bases of bounded Borel complexity. In [3, Section 5] they noticed that the unbounded Borel complexity of bases implies that the additivity of the σ -ideal under consideration is \aleph_1 . This observation exposed the heart of a result of Cichoń and Pawlikowski [5, Corollary 2.4] and showed the importance of the existence of a strictly increasing ω_1 -sequence of σ -subideals which add up to the whole ideal.

Therefore Borodulin–Nadzieja and Głab introduced a new cardinal invariant $\text{cofin}(\mathcal{I})$ associated with non-trivial σ -ideals \mathcal{I} : the minimal length of a strictly increasing sequence of σ -subideals with union \mathcal{I} (see Definition 2.1). They showed that the additivity of the σ -ideal \mathcal{I} is not larger than $\text{cofin}(\mathcal{I})$ (see [3, Proposition 5.2] or Theorem 2.2 here) and in [3, Problem 6.2(ii)] they asked if the two invariants can be different. In the present paper we answer this question in positive.

In the second section we define the relevant cardinal invariants and we point out situations when $\text{cofin}(\mathcal{I}) < \text{cof}(\mathcal{I})$ for the meager and the null ideals. In Section 3 we introduce a nicely definable σ -ideal \mathcal{I}_f with a Borel bases consisting of Π_2^0 sets. Then we show that, consistently, $\text{add}(\mathcal{I}_f) = \aleph_1$ while $\text{cofin}(\mathcal{I}_f) = \aleph_2$ (Corollary 3.15).

Notation Most of our notation is standard and compatible with that of classical textbooks (like Bartoszyński and Judah [1]). However, in forcing we keep the older convention that *a stronger condition is the larger one*.

- Ordinals will be denoted with initial letters of the Greek alphabet (α – ζ) and integers (finite ordinals) will be denoted by i, j, k, ℓ, m, n . Letters κ, λ, μ will denote uncountable cardinals.

- By a *sequence* we mean a function whose domain is a set of ordinals. Sequences will be denoted by letters $\eta, \nu, \rho, \sigma, \varsigma, \varphi, \psi$ (with possible indices).

Date: December 06, 2013.

2010 Mathematics Subject Classification. Primary 03E17; Secondary 03E35.

Key words and phrases. cardinal invariants, Borel ideals, cofin , forcing.

For two sequences η, ν we write $\nu \triangleleft \eta$ whenever ν is a proper initial segment of η , and $\nu \leq \eta$ when either $\nu \triangleleft \eta$ or $\nu = \eta$. The length of a sequence η is the order type of its domain and it is denoted by $\ell g(\eta)$.

- The power set of a set X is denoted by $\mathcal{P}(X)$ and the collection of all subsets of X of size m is called $[X]^m$ and the collection of all finite subsets of X is denoted by $[X]^{<\aleph_0}$.

- The Cantor space ${}^\omega 2$ is the space of all functions from ω to 2 , equipped with the product topology generated by sets of the form $\{\eta \in {}^\omega 2 : \nu \triangleleft \eta\}$ for $\nu \in {}^\omega > 2$.

- A family \mathcal{I} of subsets of X which is closed under finite unions and taking subsets is called an *ideal on X* . It is a *proper ideal* if $X \notin \mathcal{I}$ (i.e., $\mathcal{I} \neq \mathcal{P}(X)$) and it is a σ -ideal if it is closed under countable unions. The σ -ideal of meager subsets of the Cantor space ${}^\omega 2$ is called \mathcal{M} and the σ -ideal of Lebesgue null sets is \mathcal{N} .

- For a forcing notion \mathbb{P} , all \mathbb{P} -names for objects in the extension via \mathbb{P} will be denoted with a tilde below (e.g. $\tilde{A}, \tilde{\eta}$). The canonical name for a \mathbb{P} -generic filter over \mathbf{V} is denoted $\tilde{G}_{\mathbb{P}}$. The Cohen forcing for adding κ many Cohen reals in ${}^\omega 2$ is called \mathbb{C}_κ (so a condition in \mathbb{C}_κ is a finite function $p : \text{dom}(p) \rightarrow 2$ with $\text{dom}(p) \subseteq \kappa \times \omega$ and the order of \mathbb{C}_κ is the inclusion). The forcing \mathbb{C} is \mathbb{C}_1 .

2. COFIN AND \mathcal{M}, \mathcal{N}

Definition 2.1. Let \mathcal{I} be an ideal on X . We define the following cardinal characteristics of \mathcal{I} :

- (1) $\text{add}(\mathcal{I}) = \min\{|\mathcal{A}| : \mathcal{A} \subseteq \mathcal{I} \ \& \ \bigcup \mathcal{A} \notin \mathcal{I}\}$;
- (2) $\text{cof}(\mathcal{I}) = \min\{|\mathcal{B}| : \mathcal{B} \subseteq \mathcal{I} \ \& \ (\forall A \in \mathcal{I})(\exists B \in \mathcal{B})(A \subseteq B)\}$;
- (3) $\text{cofin}(\mathcal{I})$ is the minimal limit ordinal γ for which there exists a sequence $\bar{\mathcal{I}} = \langle \mathcal{I}_\alpha : \alpha < \gamma \rangle$ such that
 - (a) $\mathcal{I} = \bigcup_{\alpha < \gamma} \mathcal{I}_\alpha$ and
 - (b) $\mathcal{I}_\alpha \not\subseteq \mathcal{I}_\beta$ for $\alpha < \beta < \gamma$, and
 - (c) each \mathcal{I}_α is a σ -ideal,
 (or ∞ if there is no sequence $\bar{\mathcal{I}}$ as above);
- (4) $\text{cofin}^-(\mathcal{I})$ and $\text{cofin}^*(\mathcal{I})$ are defined similarly to $\text{cofin}(\mathcal{I})$, but clause (c) is replaced by (c)⁻ and (c)^{*}, respectively, where
 - (c)⁻ each \mathcal{I}_α is an ideal;
 - (c)^{*} each \mathcal{I}_α is closed under taking subsets (i.e., $B \subseteq A \in \mathcal{I}_\alpha$ implies $B \in \mathcal{I}_\alpha$);
- (5) $\text{cofin}^+(\mathcal{I})$ is the minimal limit ordinal γ for which there exists a sequence $\langle \mathcal{I}_\alpha : \alpha < \gamma \rangle$ such that clauses (a),(b) and (c) of (3) above are satisfied and
 - (d) all singletons belong to \mathcal{I}_0 .

If \mathcal{I} is a non-principal ideal (i.e., $\text{cof}(\mathcal{I}) \geq \aleph_0$), then $\text{cofin}^-(\mathcal{I})$ is a well defined cardinal and $\text{cofin}^-(\mathcal{I}) \leq \text{cof}(\mathcal{I})$. To see this, pick a basis $\{B_\zeta : \zeta < \text{cof}(\mathcal{I})\} \subseteq \mathcal{I}$ for \mathcal{I} . Let ζ_0 be the first ordinal $\zeta \leq \text{cof}(\mathcal{I})$ such that for some set $B \in \mathcal{I}$ every member of \mathcal{I} can be covered by finitely many elements of

$\{B_\varepsilon : \varepsilon < \zeta\} \cup \{B\}$. Necessarily, ζ_0 is a limit ordinal. Let $B^* \in \mathcal{I}$ be such that $\{B_\varepsilon : \varepsilon < \zeta_0\} \cup \{B^*\}$ generates \mathcal{I} , i.e., every set in \mathcal{I} can be covered by B^* and finitely many sets B_ε with $\varepsilon < \zeta_0$. For $\zeta < \zeta_0$ let \mathcal{I}_ζ be the ideal generated by $\{B_\varepsilon : \varepsilon < \zeta\} \cup \{B^*\}$. Then $\mathcal{I} = \bigcup_{\zeta < \zeta_0} \mathcal{I}_\zeta$ and, by the minimality of ζ_0 , the sequence $\langle \mathcal{I}_\zeta : \zeta < \zeta_0 \rangle$ does not stabilize. Consequently, we may choose an increasing sequence $\langle \zeta_\alpha : \alpha < \text{cf}(\zeta_0) \rangle$ cofinal in ζ_0 and such that $\langle \mathcal{I}_{\zeta_\alpha} : \alpha < \text{cf}(\zeta_0) \rangle$ is a strictly increasing sequence of ideals with the union \mathcal{I} .

The cardinal invariant cofin was introduced by Borodulin–Nadzieja and Głab in [3, Section 5]. It has the flavour of the altitude of Boolean algebras (see van Douwen, Monk and Rubin [9, p. 236]), but the two cardinal coefficients seem to be unrelated.

Theorem 2.2 (Borodulin–Nadzieja and Głab [3, Section 5]). *Let \mathcal{I} be a non-principal ideal of subsets of X . Then*

$$\text{add}(\mathcal{I}) \leq \text{cofin}^*(\mathcal{I}) \leq \text{cofin}^-(\mathcal{I}) \leq \text{cof}(\mathcal{I}) \quad \text{and} \\ \text{cofin}^-(\mathcal{I}) \leq \text{cofin}(\mathcal{I}) \leq \text{cofin}^+(\mathcal{I}).$$

Proposition 2.3. *Let $\kappa = \kappa^{\aleph_0}$ be an uncountable cardinal.*

(1) *The Cohen algebra \mathbb{C}_κ for adding κ many Cohen reals forces that*

$$\text{add}(\mathcal{M}) = \text{cofin}(\mathcal{M}) = \text{cofin}^+(\mathcal{M}) = \aleph_1 \leq \text{cof}(\mathcal{M}) = \kappa = 2^{\aleph_0}.$$

(2) *The Solovay algebra \mathbb{B}_κ for adding κ many random reals forces that*

$$\text{add}(\mathcal{N}) = \text{cofin}(\mathcal{N}) = \text{cofin}^+(\mathcal{N}) = \aleph_1 \leq \text{cof}(\mathcal{N}) = \kappa = 2^{\aleph_0}.$$

Proof. (2) In both cases the proof is essentially the same, so let us argue for the Solovay algebra only. Represent κ as the disjoint union $\kappa = \bigcup_{\varepsilon < \omega_1} K_\varepsilon$ where each K_ε is of size κ . For $\varepsilon < \omega_1$ set $\alpha_\varepsilon = \min(K_\varepsilon)$ and $A_\varepsilon = \bigcup_{\zeta < \varepsilon} K_\zeta$.

Suppose that $\bar{r} = \langle r_\alpha : \alpha < \kappa \rangle$ is a \mathbb{B}_κ -generic over \mathbf{V} , so $r_\alpha \in {}^\omega 2$ are random reals, and let us argue in $\mathbf{V}[\bar{r}]$. For each $\varepsilon < \omega_1$ let \mathcal{I}_ε be the σ -ideal generated by singletons and the family of all Borel null sets coded in $\mathbf{V}[r_\alpha : \alpha \in A_\varepsilon]$. Then $\langle \mathcal{I}_\varepsilon : \varepsilon < \omega_1 \rangle$ is an increasing sequence of σ -ideals, \mathcal{I}_0 contains all singletons and $\mathcal{N} = \bigcup_{\varepsilon < \omega_1} \mathcal{I}_\varepsilon$. Moreover, for each $\varepsilon < \omega_1$,

$$B \stackrel{\text{def}}{=} \{x \in {}^\omega 2 : (\forall n < \omega)(x(2n) = r_{\alpha_\varepsilon}(2n))\} \in \mathcal{I}_{\varepsilon+1} \setminus \mathcal{I}_\varepsilon.$$

Why? Clearly, B is a Borel null set coded in $\mathbf{V}[r_\alpha : \alpha \in A_{\varepsilon+1}]$, so $B \in \mathcal{I}_{\varepsilon+1}$. Suppose B_i are Borel null sets coded in $\mathbf{V}[r_\alpha : \alpha \in A_\varepsilon]$ and $x_i \in ({}^\omega 2)^{\mathbf{V}[\bar{r}]}$ (for $i < \omega$). Choose $x^* \in \{x \in {}^\omega 2 \cap \mathbf{V} : (\forall n < \omega)(x(2n) = 0)\} \setminus \{x_i + r_{\alpha_\varepsilon} : i < \omega\}$. Then $x^* + r_{\alpha_\varepsilon}$ is a random real over $\mathbf{V}[r_\alpha : \alpha \in A_\varepsilon]$, so $x^* + r_{\alpha_\varepsilon} \in B \setminus (\bigcup_{i < \omega} B_i \cup \{x_i : i < \omega\})$. Thus we may conclude that $B \notin \mathcal{I}_\varepsilon$. \square

Definition 2.4 (Rosłanowski and Shelah [6, Definition 3.4]). Let \mathcal{I} be an ideal of subsets of a space X and α^*, β^* be limit ordinals. An $\alpha^* \times \beta^*$ -base for \mathcal{I} is an indexed family $\mathcal{B} = \{B_{\alpha, \beta} : \alpha < \alpha^* \ \& \ \beta < \beta^*\}$ of sets from \mathcal{I} such that

modified: 2013-12-11

1022 revision: 2013-12-11

- (i) \mathcal{B} is a basis for \mathcal{I} , i.e., $(\forall A \in \mathcal{I})(\exists B \in \mathcal{B})(A \subseteq B)$, and
(ii) for each $\alpha_0, \alpha_1 < \alpha^*$, $\beta_0, \beta_1 < \beta^*$ we have

$$B_{\alpha_0, \beta_0} \subseteq B_{\alpha_1, \beta_1} \quad \Leftrightarrow \quad \alpha_0 \leq \alpha_1 \ \& \ \beta_0 \leq \beta_1.$$

If follows from results of Bartoszyński and Kada [2] (for the meager ideal) and Burke and Kada [4] (for the null ideal) that for any cardinals κ and λ of uncountable cofinality we may force that \mathcal{M} has a $\kappa \times \lambda$ -basis, and we may also force that \mathcal{N} has a $\kappa \times \lambda$ -basis. In [6, Theorem 3.7] we constructed a model in which both ideals have $\kappa \times \lambda$ -bases.

Proposition 2.5. *Let κ, λ be regular uncountable cardinals, $\kappa \leq \lambda$,*

- (1) *If \mathcal{I} is a σ -ideal on a space X and \mathcal{I} has a $\kappa \times \lambda$ -base, then*

$$\kappa = \text{add}(\mathcal{I}) = \text{cofin}(\mathcal{I}) \quad \text{and} \quad \text{cof}(\mathcal{I}) = \lambda.$$

- (2) *There is a ccc forcing notion \mathbb{P} forcing that $2^{\aleph_0} = \lambda^{\aleph_0}$ and*

- (i) *the σ -ideal \mathcal{N} has a $\kappa \times \lambda$ -base $\{A_{\alpha, \beta} : \alpha < \kappa, \beta < \lambda\}$ with the property that*

$$\alpha_0 > \alpha_1 \ \vee \ \beta_0 > \beta_1 \quad \Rightarrow \quad |A_{\alpha_0, \beta_0} \setminus A_{\alpha_1, \beta_1}| = 2^{\aleph_0},$$

and

- (ii) *the σ -ideal \mathcal{M} has a $\kappa \times \lambda$ -base $\{B_{\alpha, \beta} : \alpha < \kappa, \beta < \lambda\}$ with the property that*

$$\alpha_0 > \alpha_1 \ \vee \ \beta_0 > \beta_1 \quad \Rightarrow \quad |B_{\alpha_0, \beta_0} \setminus B_{\alpha_1, \beta_1}| = 2^{\aleph_0}.$$

In particular,

$$\Vdash_{\mathbb{P}} \text{ " } \text{add}(\mathcal{M}) = \text{add}(\mathcal{N}) = \text{cofin}^+(\mathcal{M}) = \text{cofin}^+(\mathcal{N}) = \kappa \quad \text{and} \\ \text{cof}(\mathcal{M}) = \text{cof}(\mathcal{N}) = \lambda. \text{ "}$$

Proof. (1) Assume that $\{B_{\alpha, \beta} : \alpha < \kappa, \beta < \lambda\}$ is a $\kappa \times \lambda$ -base for \mathcal{I} . It should be clear that then $\kappa = \text{add}(\mathcal{I})$ and $\text{cof}(\mathcal{I}) = \lambda$.

Let us argue that $\text{cofin}(\mathcal{I}) \leq \kappa$. For $\zeta < \kappa$ let \mathcal{I}_{ζ} be the σ -ideal generated by the family $\{B_{\alpha, \beta} : \alpha \leq \zeta \ \& \ \beta < \lambda\}$. Plainly, $\langle \mathcal{I}_{\zeta} : \zeta < \kappa \rangle$ is an increasing sequence of σ -ideals such that $\mathcal{I} = \bigcup_{\zeta < \kappa} \mathcal{I}_{\zeta}$. We claim that $B_{\zeta+1, 0} \in \mathcal{I}_{\zeta+1} \setminus \mathcal{I}_{\zeta}$.

Suppose that $I \subseteq (\zeta+1) \times \lambda$ is countable. Then we may choose $\beta^* < \lambda$ such that $I \subseteq (\zeta+1) \times \beta^*$ and consequently $\bigcup\{B_{\alpha, \beta} : (\alpha, \beta) \in I\} \subseteq B_{\zeta, \beta^*}$. But $B_{\zeta+1, 0} \not\subseteq B_{\zeta, \beta^*}$ and so $B_{\zeta+1, 0} \not\subseteq \bigcup\{B_{\alpha, \beta} : (\alpha, \beta) \in I\}$. Now we may conclude now that $B_{\zeta+1, 0} \notin \mathcal{I}_{\zeta}$.

- (2) The forcing notion $\mathbb{Q}^{\kappa, \lambda}$ constructed in the proof of [6, Theorem 3.7] has the desired properties, see [6, Remark 3.8]. \square

3. COFIN AND \mathcal{I}_f

We introduce here a nicely definable Borel ideal \mathcal{I}_f for which, consistently, $\text{add}(\mathcal{I}_f) < \text{cofin}(\mathcal{I}_f)$. The proof of the consistency will resemble Shelah [8, Chapter II, Theorem 4.6] (and thus also [7]). The respective forcing notion is obtained by means of FS iteration of ccc forcing notions, however the iteration itself is forced too.

Context 3.1. Let us fix two strictly increasing functions $f, g : \omega \rightarrow \omega$ such that for each $n < \omega$ we have

$$2 < g(n) < f(n) \quad \text{and} \quad \frac{g(n)}{f(n)} \leq \frac{1}{n+1}.$$

Definition 3.2. (1) A null slalom below f is a function $\varphi \in \prod_{n < \omega} \mathcal{P}(f(n))$

such that $\lim_{n \rightarrow \infty} \frac{|\varphi(n)|}{f(n)} = 0$.

(2) Let \mathcal{S}_f be the collection of all null slaloms below f and let $\mathcal{X}_f = \prod_{n < \omega} f(n)$ be equipped with the natural product topology (so \mathcal{X}_f is a Polish space).

(3) For $\varphi \in \mathcal{S}_f$ we define

$$[\varphi] = \{x \in \mathcal{X}_f : (\exists^\infty n < \omega)(x(n) \in \varphi(n))\}.$$

Observation 3.3. Let $\varphi_i \in \mathcal{S}_f$ (for $i < \omega$).

(1) $[\varphi_0] \subseteq [\varphi_1]$ if and only if $(\forall^\infty n < \omega)(\varphi_0(n) \subseteq \varphi_1(n))$.

(2) There is $\psi \in \mathcal{S}_f$ such that $\bigcup_{i < \omega} [\varphi_i] \subseteq [\psi]$.

Definition 3.4. Let \mathcal{I}_f be the σ -ideal of subsets of \mathcal{X}_f generated by all sets $[\varphi]$ for $\varphi \in \mathcal{S}_f$. Thus, by Observation 3.3,

$$\mathcal{I}_f = \{A \subseteq \mathcal{X}_f : (\exists \varphi \in \mathcal{S}_f)(A \subseteq [\varphi])\}.$$

We will construct a forcing notion \mathbb{P} forcing that $\text{add}(\mathcal{I}_f) < \text{cofin}(\mathcal{I}_f)$, but first we need several technical ingredients.

Definition 3.5. For a cardinal κ we define a forcing notion \mathbb{Q}_0^κ :

A condition p in \mathbb{Q}_0^κ is a finite function such that $\text{dom}(p) \subseteq \kappa$ and for some $n = n^p < \omega$, for all $\varepsilon \in \text{dom}(p)$ we have $p(\varepsilon) \in \prod_{i < n} [f(i)]^{g(i)}$.

The order $\leq_{\mathbb{Q}_0^\kappa}$ of \mathbb{Q}_0^κ is defined by letting $p \leq q$ if and only if

$$\text{dom}(p) \subseteq \text{dom}(q) \quad \text{and} \quad (\forall \varepsilon \in \text{dom}(p))(p(\varepsilon) \trianglelefteq q(\varepsilon)).$$

For $\varepsilon < \kappa$, a \mathbb{Q}_0^κ -name $\nu(\varepsilon)$ is defined by

$$\Vdash_{\mathbb{Q}_0^\kappa} \nu(\varepsilon) = \bigcup \{p(\varepsilon) : \varepsilon \in \text{dom}(p) \ \& \ p \in \mathbb{G}_{\mathbb{Q}_0^\kappa}\}.$$

Observation 3.6. (1) The forcing notion \mathbb{Q}_0^κ is equivalent to \mathbb{C}_κ , the forcing adding κ many Cohen reals.

(2) $\Vdash_{\mathbb{Q}_0^\kappa}$ “for every $\varepsilon < \kappa$ we have $\nu(\varepsilon) \in \prod_{i < \omega} [f(i)]^{g(i)} \subseteq \mathcal{S}_f$ ”.

Definition 3.7. Let μ be an infinite cardinal and $\bar{\varphi} = \langle \varphi_\zeta : \zeta < \mu \rangle$ be a sequence of null slaloms below f (so $\varphi_\zeta \in \mathcal{S}_f$ for $\zeta < \mu$). We define a forcing notion $\mathbb{Q}_\mu^*(\bar{\varphi})$.

A condition in $\mathbb{Q}_\mu^*(\bar{\varphi})$ is a tuple $p = (k^p, m^p, u^p, \sigma^p) = (k, m, u, \sigma)$ such that

(a) $k, m < \omega$, $\emptyset \neq u \in [\mu]^{< \aleph_0}$, $\sigma \in \prod_{i < k} \mathcal{P}(f(i))$, and

(b) for each $\ell \geq k$ and $\zeta \in u$ we have $|\varphi_\zeta(\ell)| < \frac{f(\ell)}{m \cdot |u|}$.

The order $\leq = \leq_{\mathbb{Q}_\mu^*(\bar{\varphi})}$ of $\mathbb{Q}_\mu^*(\bar{\varphi})$ is defined by

$p \leq q$ if and only if $(p, q \in \mathbb{Q}_\mu^*(\bar{\varphi})$ and) $k^p \leq k^q$, $m^p \leq m^q$, $u^p \subseteq u^q$, $\sigma^p \leq \sigma^q$ and for each $\ell \in [k^p, k^q)$ we have

$$|\sigma^q(\ell)| \leq \frac{f(\ell)}{m^p} \quad \text{and} \quad \bigcup \{ \varphi_\zeta(\ell) : \zeta \in u^p \} \subseteq \sigma^q(\ell).$$

We also define a $\mathbb{Q}_\mu^*(\bar{\varphi})$ -name ζ by

$$\Vdash_{\mathbb{Q}_\mu^*(\bar{\varphi})} \zeta = \bigcup \{ \sigma^p : p \in \mathcal{G}_{\mathbb{Q}_\mu^*(\bar{\varphi})} \}.$$

Proposition 3.8. *Let μ be an infinite cardinal and $\bar{\varphi} = \langle \varphi_\zeta : \zeta < \mu \rangle \subseteq \mathcal{S}_f$. Then $\mathbb{Q}_\mu^*(\bar{\varphi})$ is a well defined ccc forcing notion of size μ and*

$$\Vdash_{\mathbb{Q}_\mu^*(\bar{\varphi})} \text{ “ } \zeta \in \mathcal{S}_f \ \& \ \bigcup_{\zeta < \mu} [\varphi_\zeta] \subseteq [\zeta] \in \mathcal{I}_f \text{ ”.}$$

Proof. First note that if $p \in \mathbb{Q}_\mu^*(\bar{\varphi})$ and $m = m^p$, $k = k^p + 1$, $u = u^p$ and $\sigma = \sigma^p \setminus \langle \bigcup_{\zeta \in u} \varphi_\zeta(k^p) \rangle$, then $(k, m, u, \sigma) \in \mathbb{Q}_\mu^*(\bar{\varphi})$ is a condition stronger than p . Hence we may conclude that $\Vdash_{\mathbb{Q}_\mu^*(\bar{\varphi})} \zeta \in \prod_{i < \omega} \mathcal{P}(f(i))$.

Also, if $p \in \mathbb{Q}_\mu^*(\bar{\varphi})$ and $m > m^p$, then we may find $k > k^p$ such that $|\varphi_\zeta(\ell)| < \frac{f(\ell)}{m \cdot |u^p|}$ for all $\zeta \in u^p$ and $\ell \geq k$. Let $u = u^p$ and $\sigma \in \prod_{i < k} \mathcal{P}(f(i))$ be such that $\sigma(\ell) = \sigma^p(\ell)$ for $\ell < k^p$ and $\sigma(\ell) = \bigcup_{\zeta \in u} \varphi_\zeta(\ell)$ for $\ell \in [k^p, k)$.

Then $(k, m, u, \sigma) \in \mathbb{Q}_\mu^*(\bar{\varphi})$ is a condition stronger than p and it forces that $|\zeta(\ell)| \leq \frac{f(\ell)}{m}$ for all $\ell \geq k$. Hence we may conclude that $\Vdash_{\mathbb{Q}_\mu^*(\bar{\varphi})} \zeta \in \mathcal{S}_f$.

It follows from the definition of the order of $\mathbb{Q}_\mu^*(\bar{\varphi})$ that

$$p \Vdash_{\mathbb{Q}_\mu^*(\bar{\varphi})} (\forall \ell \geq k^p) (\forall \zeta \in u^p) (\varphi_\zeta(\ell) \subseteq \zeta(\ell)),$$

and hence easily $\Vdash_{\mathbb{Q}_\mu^*(\bar{\varphi})} \bigcup_{\zeta < \mu} [\varphi_\zeta] \subseteq [\zeta]$.

Let us argue now that $\mathbb{Q}_\mu^*(\bar{\varphi})$ satisfies the ccc. Suppose $\langle p_\varepsilon : \varepsilon < \omega_1 \rangle \subseteq \mathbb{Q}_\mu^*(\bar{\varphi})$. For each $\varepsilon < \omega_1$, we may find $K^\varepsilon > k^{p_\varepsilon}$ such that

$$(\oplus)_1 (\forall \ell \geq K^\varepsilon) (\forall \zeta \in u^{p_\varepsilon}) (|\varphi_\zeta(\ell)| < \frac{f(\ell)}{2 \cdot |u^{p_\varepsilon}| \cdot m^{p_\varepsilon}})$$

and define $\rho^\varepsilon \in \prod_{i < K^\varepsilon} \mathcal{P}(f(i))$ so that $\rho^\varepsilon(\ell) = \sigma^{p_\varepsilon}(\ell)$ for $\ell < k^{p_\varepsilon}$ and $\rho^\varepsilon(\ell) =$

$\bigcup_{\zeta \in u^{p_\varepsilon}} \varphi_\zeta(\ell)$ for $\ell \in [k^{p_\varepsilon}, K^\varepsilon)$. Then we may find an uncountable set $S \subseteq \omega_1$

and K^*, m^*, ρ^*, ℓ^* such that for all $\varepsilon \in S$:

$$(\oplus)_2 K^* = K^\varepsilon, m^* = m^{p_\varepsilon}, \rho^* = \rho^\varepsilon \text{ and } |u^\varepsilon| = \ell^*.$$

Consider distinct $\varepsilon_0, \varepsilon_1 \in S$: letting $u^* = u^{\varepsilon_0} \cup u^{\varepsilon_1}$ we get a condition $(K^*, m^*, u^*, \rho^*) \in \mathbb{Q}_\mu^*(\bar{\varphi})$ stronger than both p_{ε_0} and p_{ε_1} . \square

Definition 3.9. Let $\kappa < \lambda$ be uncountable regular cardinals.

- (1) A Y -iteration for κ, λ is a finite support iteration $\langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \alpha \rangle$ of ccc forcing notions such that the following demands $(\otimes)_1$ – $(\otimes)_3$ are satisfied.

- (\otimes)₁ $0 < \alpha \leq \lambda$ and $\mathbb{Q}_0 = \mathbb{Q}_0^\kappa$ is the forcing notion adding κ Cohen reals as represented in Definition 3.5 with \mathbb{Q}_0^κ -names $\nu(\varepsilon)$ (for $\varepsilon < \kappa$) as defined there.
- (\otimes)₂ For each $\beta < \alpha$ we have $\Vdash_{\mathbb{P}_\beta} |\mathbb{Q}_\beta| \leq \lambda$.
- (\otimes)₃ Let $n < \omega$. Suppose that $\langle p_\zeta : \zeta < \kappa \rangle \subseteq \mathbb{P}_\alpha$ and $\langle \delta_\zeta : \zeta < \kappa \rangle \subseteq \kappa$ and $\delta_\zeta \neq \delta_{\zeta'}$ for $\zeta < \zeta' < \kappa$. Then there are $q \in \mathbb{P}_\alpha$, $m > n$, $v \subseteq \kappa$, and A_ζ (for $\zeta \in v$) such that
 - (i) $|v| \geq \frac{f(m)}{2 \cdot g(m)}$,
 - (ii) $p_\zeta \leq q$ for all $\zeta \in v$,
 - (iii) $A_\zeta \in [f(m)]g(m)$ (for $\zeta \in v$) are pairwise disjoint sets,
 - (iv) $q \Vdash_{\mathbb{P}_\alpha}$ “ $(\forall \zeta \in v) (\nu(\delta_\zeta)(m) = A_\zeta)$ ”.
- (2) The collection of all (dense subsets of) Y -iterations for κ, λ of length $< \lambda$ which belong to $\mathcal{H}(\mathfrak{I}_\lambda^+)$ is denoted by $\mathbb{Y}_\kappa^\lambda$. It is ordered by the end-extension of iterations \trianglelefteq .

The condition 3.9(1)(\otimes)₃ implies that the null slaloms added at the first step of a Y -iteration provide a family of sets whose union is not included in any null slalom. Note that in 3.9(1)(\otimes)₃ necessarily $|v| \leq \frac{f(m)}{g(m)}$.

Lemma 3.10. *Assume $\kappa < \lambda$ are regular uncountable cardinals. Suppose that $\langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \alpha \rangle$ is a Y -iteration for κ, λ . Then $\Vdash_{\mathbb{P}_\alpha} \text{add}(\mathcal{I}_f) \leq \kappa$.*

Proof. We know that for each $\varepsilon < \kappa$ we have $\Vdash_{\mathbb{P}_\alpha} \nu(\varepsilon) \in \mathcal{S}_f$ (remember Observation 3.6) and therefore $\Vdash_{\mathbb{P}_\alpha} \{[\nu(\varepsilon)] : \varepsilon < \kappa\} \subseteq \mathcal{I}_f$. We are going to argue that

$$\Vdash_{\mathbb{P}_\alpha} \bigcup \{[\nu(\varepsilon)] : \varepsilon < \kappa\} \notin \mathcal{I}_f.$$

Suppose towards contradiction that this is not the case. Then we may pick $p \in \mathbb{P}_\alpha$ and a \mathbb{P}_α -name φ such that

$$p \Vdash_{\mathbb{P}_\alpha} \varphi \in \mathcal{S}_f \ \& \ (\forall \varepsilon < \kappa) (\forall^\infty n < \omega) (\nu(\varepsilon)(n) \subseteq \varphi(n))$$

(remember Observation 3.3). Now for each $\varepsilon < \kappa$ we pick a condition $p_\varepsilon \geq p$ and an integer $n_\varepsilon < \omega$ such that

$$p_\varepsilon \Vdash_{\mathbb{P}_\alpha} (\forall n \geq n_\varepsilon) (\nu(\varepsilon)(n) \subseteq \varphi(n) \ \& \ \frac{|\varphi(n)|}{f(n)} < 1/4).$$

For some $n^* < \omega$ the set $S = \{\varepsilon < \kappa : n_\varepsilon = n^*\}$ is of size κ . Apply 3.9(1)(\otimes)₃ to $\langle p_\varepsilon : \varepsilon \in S \rangle \subseteq \mathbb{P}_\alpha$ and $\langle \varepsilon : \varepsilon \in S \rangle \subseteq \kappa$ and $n = n^*$ to find $q \in \mathbb{P}_\alpha$, $m > n^*$, $v \subseteq S$, and A_ε (for $\varepsilon \in v$) such that conditions (i)–(iv) there hold. Then

$$q \Vdash_{\mathbb{P}_\alpha} \text{“ } \bigcup_{\varepsilon \in v} A_\varepsilon = \bigcup_{\varepsilon \in v} \nu(\varepsilon)(m) \subseteq \varphi(m) \ \& \ |\varphi(m)| < \frac{f(m)}{4} \text{”}.$$

But $|\bigcup_{\varepsilon \in v} A_\varepsilon| = |v| \cdot g(m) \geq \frac{f(m)}{2}$, a contradiction. \square

Context 3.11. For the rest of this section we fix uncountable regular cardinals $\kappa < \lambda$ such that $\lambda^\kappa = \lambda$. Also, instead of “ Y -iteration for κ, λ ” we will just say “ Y -iteration”.

Lemma 3.12. (1) $\langle \mathbb{P}_0, \mathbb{Q}_0^\kappa \rangle$ is a Y -iteration (of length 1).

- (2) Assume that $\langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \alpha \rangle$ is a Y -iteration of length $\alpha < \lambda$ and \mathbb{Q} is a \mathbb{P}_α -name for a ccc forcing notion of size $< \kappa$ (i.e., $\Vdash_{\mathbb{P}_\alpha} |\mathbb{Q}| < \kappa$). Then $\langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \alpha \rangle \frown \langle \mathbb{P}_\alpha, \mathbb{Q} \rangle$ is a Y -iteration of length $\alpha + 1$. In particular, $\langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \alpha \rangle \frown \langle \mathbb{P}_\alpha, \mathbb{C} \rangle$ is a Y -iteration.
- (3) If $\langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \alpha \rangle$ is a Y -iteration and \mathbb{Q} is a \mathbb{P}_α -name for a σ -centered forcing, then $\langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \alpha \rangle \frown \langle \mathbb{P}_\alpha, \mathbb{Q} \rangle$ is a Y -iteration.
- (4) If $\gamma \leq \lambda$ is a limit ordinal and $\langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \gamma \rangle$ is an FS iteration such that $\langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \alpha \rangle$ is a Y -iteration for every $\alpha < \gamma$, then $\langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \gamma \rangle$ is a Y -iteration.
- (5) $(\mathbb{Y}_\kappa^\lambda, \sqsubseteq)$ is a $< \lambda$ -complete forcing notion (i.e., all chains of length $< \lambda$ have an upper bound in $\mathbb{Y}_\kappa^\lambda$).

Proof. In all cases the only demand of Definition 3.9(1) that needs to be verified is $(\otimes)_3$.

(1) Let \mathbb{Q}_0^κ be the forcing notion adding κ Cohen reals as described in Definition 3.5. Let $n < \omega$, $\delta_\zeta \in \kappa$ and $p_\zeta \in \mathbb{Q}_0^\kappa$ (for $\zeta < \kappa$) satisfy the assumptions of 3.9(1)(\otimes)₃. By making conditions p_ζ stronger and possibly passing to a subsequence, we may assume also that:

- (*)₁ $\delta_\zeta \in \text{dom}(p_\zeta)$ for all $\zeta < \kappa$,
- (*)₂ for some $m > n + 2$, for all $\zeta < \kappa$, we have $n^{p_\zeta} = m$ (so $p_\zeta(\varepsilon) \in \prod_{i < m} [f(i)]^{g(i)}$ for $\varepsilon \in \text{dom}(p_\zeta)$),
- (*)₃ the family $\{\text{dom}(p_\zeta) : \zeta < \kappa\}$ forms a Δ -system of finite sets and for all $\zeta, \zeta' < \kappa$ the conditions $p_\zeta, p_{\zeta'}$ are compatible.

Pick any $v \subseteq \kappa$ of size $\lceil \frac{f(m)}{2 \cdot g(m)} \rceil$. Since

$$\lceil \frac{f(m)}{2 \cdot g(m)} \rceil \cdot g(m) \leq \frac{f(m)}{2} + g(m) \leq \frac{f(m)}{2} + \frac{f(m)}{m+1} < f(m),$$

we may choose pairwise disjoint sets $A_\zeta \in [f(m)]^{g(m)}$ (for $\zeta \in v$). Now define a condition $q \in \mathbb{Q}_0^\kappa$ so that $\text{dom}(q) = \bigcup \{\text{dom}(p_\zeta) : \zeta \in v\}$, $n^q = m+1$ and for $\varepsilon \in \text{dom}(p_\zeta)$ the sequence $q(\varepsilon)$ extends $p_\zeta(\varepsilon)$ and $q(\delta_\zeta)(m) = A_\zeta$ (for $\zeta \in v$).

(2) Without loss of generality, for some ordinal $\gamma^* < \kappa$ we have $\Vdash_{\mathbb{P}_\alpha}$ “the set of conditions in \mathbb{Q} is γ^* ”. Let $n < \omega$ and $p_\zeta \in \mathbb{P}_{\alpha+1}$, $\delta_\zeta \in \kappa$ (for $\zeta < \kappa$) satisfy the assumptions of 3.9(1)(\otimes)₃. We may make our conditions stronger and we may pass to a subsequence, so we may assume that $\alpha \in \text{dom}(p_\zeta)$ and $p_\zeta(\alpha) = \gamma < \gamma^*$ is an actual object, not a name (for $\zeta < \kappa$). Apply the assumption of 3.9(1)(\otimes)₃ for $\langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \alpha \rangle$ to $n, p_\zeta \restriction \alpha, \delta_\zeta$ (for $\zeta < \kappa$) and choose $m > n$, $q^* \in \mathbb{P}_\alpha$, $v \subseteq \kappa$ and pairwise disjoint sets $A_\zeta \subseteq f(m)$ each of size $g(m)$ (for $\zeta \in v$) such that

- $|v| \geq \frac{f(m)}{2 \cdot g(m)}$ and
- q^* is stronger than all $p_\zeta \restriction \alpha$ for $\zeta \in v$ and it forces that $\nu(\delta_\zeta)(m) = A_\zeta$ (for $\zeta \in v$).

Let $q \in \mathbb{P}_{\alpha+1}$ be such that $q \restriction \alpha = q^*$ and $q(\alpha) = \gamma$.

(3) Assume that $\Vdash_{\mathbb{P}_\alpha}$ “ \mathbb{Q} is a σ -centered forcing notion” and fix a \mathbb{P}_α -name \check{F} such that

$\Vdash_{\mathbb{P}}$ “ $\check{F} : \mathbb{Q} \longrightarrow \omega$ is a function satisfying:

if $x_0, \dots, x_k \in \mathbb{Q}$, $k < \omega$, and $\check{F}(x_0) = \dots = \check{F}(x_k) = m$,

then the conditions x_0, \dots, x_k have a common upper bound in \mathbb{Q} ”.

Suppose that $n < \omega$ and $p_\zeta \in \mathbb{P}_{\alpha+1}$, $\delta_\zeta \in \kappa$ (for $\zeta < \kappa$) satisfy the assumptions of 3.9(1)(\otimes)₃. By making the conditions stronger and passing to a subsequence we may demand that $\alpha \in \text{dom}(p_\zeta)$ and for some $M < \omega$ we also have $p_\zeta \restriction \alpha \Vdash_{\mathbb{P}_\alpha}$ “ $\check{F}(p_\zeta(\alpha)) = M$ ”. Use the assumption of 3.9(1)(\otimes)₃ for $\langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \alpha \rangle$ for $n, p_\zeta \restriction \alpha, \delta_\zeta$ (for $\zeta < \kappa$) to find $m > n$, $q^* \in \mathbb{P}_\alpha$, $v \subseteq \kappa$ and pairwise disjoint sets $A_\zeta \in [f(m)]^{g(m)}$ (for $\zeta \in v$) such that

- $|v| \geq \frac{f(m)}{2 \cdot g(m)}$ and
- q^* is stronger than all $p_\zeta \restriction \alpha$ for $\zeta \in v$ and it forces that $\nu(\delta_\zeta)(m) = A_\zeta$ (for $\zeta \in v$).

Then also the condition q^* forces that $\check{F}(p_\zeta(\alpha)) = M$ for all $\zeta \in v$, and thus we may pick a \mathbb{P}_α -name \check{q}_α such that $q^* \Vdash$ “ \check{q}_α is a condition stronger than all $p_\zeta(\alpha)$ for $\zeta \in v$ ”. Define $q \in \mathbb{P}_{\alpha+1}$ by $q \restriction \alpha = q^*$ and $q(\alpha) = \check{q}_\alpha$.

(4) Let n, p_ζ, δ_ζ (for $\zeta < \kappa$) be as in the assumptions of 3.9(1)(\otimes)₃. By passing to a subsequence we may also demand that $\{\text{dom}(p_\zeta) : \zeta < \kappa\}$ is a Δ -system of finite subsets of γ with root D . Pick $\alpha < \gamma$ such that $D \subseteq \alpha$. Since $\langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \alpha \rangle$ is a Y -iteration, we may apply 3.9(1)(\otimes)₃ to n, δ_ζ and $p_\zeta \restriction \alpha$ (for $\zeta < \kappa$). This will give us q^*, v and A_ζ (for $\zeta \in v$) satisfying (i)–(iv) there (with $p_\zeta \restriction \alpha$ in place of p_ζ and q^* in place of q). Let $q \in \mathbb{P}_\gamma$ be such that $\text{dom}(q) = \text{dom}(q^*) \cup \bigcup \{\text{dom}(p_\zeta) : \zeta \in v\}$ and $q \restriction \alpha = q^*$ and $q(\beta) = p_\zeta(\beta)$ whenever $\zeta \in v$, $\beta \in \text{dom}(p_\zeta) \setminus \alpha$.

(5) Follows from (4). □

Lemma 3.13. *Assume that*

- (a) $\aleph_0 \leq \mu \leq \kappa$ is a regular cardinal, $\alpha < \lambda$ is a limit ordinal of cofinality μ and $\langle \alpha(\zeta) : \zeta < \mu \rangle$ is a strictly increasing sequence cofinal in α ,
- (b) $\langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \alpha \rangle$ is a Y -iteration,
- (c) $\bar{\varphi} = \langle \varphi_\zeta : \zeta < \mu \rangle$ is a $\mathbb{P}_{\alpha(0)}$ -name for a μ -sequence of null slaloms below f (so $\Vdash \varphi_\zeta \in \mathcal{S}_f$),
- (d) for each $\zeta < \mu$ we have $\Vdash_{\mathbb{P}_{\alpha(\zeta)}} \mathbb{Q}_{\alpha(\zeta)} = \mathbb{C}$ with \check{c}_ζ being the $\mathbb{P}_{\alpha(\zeta)+1}$ -name for the Cohen real in ${}^\omega 2$ added by $\mathbb{Q}_{\alpha(\zeta)}$,
- (e) $\check{\tau}_\zeta$ is a \mathbb{P}_α -name for an element of 2 (for $\zeta < \mu$),
- (f) for $\zeta < \mu$, $\check{\psi}_\zeta$ is a \mathbb{P}_α -name for a null slalom below f such that

$$\Vdash_{\mathbb{P}_\alpha} \quad \check{\psi}_\zeta(i) = \begin{cases} \varphi_\zeta(i) & \text{if } \check{c}_\zeta(i) = \check{\tau}_\zeta, \\ \emptyset & \text{if } \check{c}_\zeta(i) = 1 - \check{\tau}_\zeta \end{cases} \quad \text{for each } i < \omega \text{ ”},$$

and $\bar{\psi} = \langle \check{\psi}_\zeta : \zeta < \mu \rangle$ is the resulting \mathbb{P}_α -name for a μ -sequence of null slaloms below f .

Then $\langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \alpha \rangle \frown \langle \mathbb{P}_\alpha, \mathbb{Q}_\mu^*(\bar{\psi}) \rangle$ is a Y -iteration of length $\alpha + 1$.

Proof. First we consider the case when $\mu = \kappa$ and let us argue that conditions 3.9(1)(\otimes)₃ holds for $\mathbb{P}_{\alpha+1}$.

Let $n < \omega$, $p_\zeta \in \mathbb{P}_{\alpha+1}$ and $\delta_\zeta < \kappa$ (for $\zeta < \kappa$) be such that $\delta_\zeta \neq \delta_{\zeta'}$ for $\zeta < \zeta' < \kappa$. For each $\zeta < \kappa$ pick a condition $p'_\zeta \in \mathbb{P}_{\alpha+1}$ stronger than p_ζ and such that

- (*)₁ $\alpha \in \text{dom}(p'_\zeta)$ and for some $k^\zeta, m^\zeta, u^\zeta$ and σ^ζ (objects, not names) we have $p'_\zeta \upharpoonright \alpha \Vdash_{\mathbb{P}_\alpha} "p'_\zeta(\alpha) = (k^\zeta, m^\zeta, u^\zeta, \sigma^\zeta)"$.

Choose conditions $p''_\zeta \in \mathbb{P}_{\alpha+1}$ stronger than p'_ζ (so also $p''_\zeta \geq p_\zeta$) and such that $p''_\zeta(\alpha) = p'_\zeta(\alpha)$ and for all ζ :

- (*)₂ for some (objects, not names) $\mathfrak{t}_\varepsilon^\zeta$ for $\varepsilon \in u^\zeta$ we have $p''_\zeta \upharpoonright \alpha \Vdash_{\mathbb{P}_\alpha} " \mathcal{I}_\varepsilon = \mathfrak{t}_\varepsilon^\zeta "$,
- (*)₃ for some $i^\zeta < \omega$ for all $\varepsilon \in u^\zeta$ we have that

$$\alpha(\varepsilon) \in \text{dom}(p''_\zeta) \quad \text{and} \quad p''_\zeta(\alpha(\varepsilon)) \in {}^{i^\zeta}2 \text{ are actual objects, not names.}$$

Since each φ_ε is a $\mathbb{P}_{\alpha(0)}$ -name, we may decide the initial segments of φ_ε by strengthening $p''_\zeta \upharpoonright \alpha(0)$ only (i.e., without changing $p''_\zeta \upharpoonright [\alpha(0), \alpha]$). Therefore, after using a procedure similar to that in the proof of 3.8, for each $\zeta < \kappa$ we may find a condition $p^*_\zeta \in \mathbb{P}_{\alpha+1}$, $K^\zeta > k^\zeta + i^\zeta$ and a sequence $\rho^\zeta \in \prod_{i < K^\zeta} \mathcal{P}(f(i))$ such that

- (*)₄ $p_\zeta \leq p''_\zeta \leq p^*_\zeta$, and $p''_\zeta \upharpoonright [\alpha(0), \alpha] = p^*_\zeta \upharpoonright [\alpha(0), \alpha]$, and
- (*)₅ $p^*_\zeta \Vdash_{\mathbb{P}_\alpha} "p^*_\zeta(\alpha) = (K^\zeta, m^\zeta, u^\zeta, \rho^\zeta)"$.

Next we may find a set $S \subseteq \kappa$ of size κ and K^*, m^*, ρ^*, i^* and ℓ^* such that

- (*)₆ $K^* = K^\zeta$, $m^* = m^\zeta$, $\rho^* = \rho^\zeta$, $|u^\zeta| = \ell^*$ and $i^\zeta = i^*$ for all $\zeta \in S$,
- (*)₇ $\{u^\zeta : \zeta \in S\}$ is a Δ -system of finite subsets of κ with root U ,
- (*)₈ $\{\text{dom}(p^*_\zeta) : \zeta \in S\}$ is a Δ -system of finite subsets of $\alpha + 1$ with root D ,
- (*)₉ for some $\varepsilon^* < \kappa$ we have $D \setminus \{\alpha\} \subseteq \alpha(\varepsilon^*)$ and $U = u^\zeta \cap \varepsilon^*$ for all $\zeta \in S$.

Since $\langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \alpha(\varepsilon^*) \rangle$ is a Y-iteration, we may apply 3.9(1)(\otimes)₃ to $\langle p^*_\zeta \upharpoonright \alpha(\varepsilon^*), \delta_\zeta : \zeta \in S \rangle$ and n . This will give us $v \subseteq S$, $q_0 \in \mathbb{P}_{\alpha(\varepsilon^*)}$, $m > n$ and $A_\zeta \in [f(m)]^{g(m)}$ for $\zeta \in v$ such that

- (*)₁₀ • $|v| \geq \frac{f(m)}{2 \cdot g(m)}$ and $p^*_\zeta \upharpoonright \alpha(\varepsilon^*) \leq q_0$ for all $\zeta \in v$, and
- $A_\zeta \cap A_{\zeta'} = \emptyset$ for distinct $\zeta, \zeta' \in v$, and
- $q_0 \Vdash_{\mathbb{P}_{\alpha(\varepsilon^*)}} "(\forall \zeta \in v)(\nu(\delta_\zeta)(m) = A_\zeta)"$.

Next, since φ_ε are $\mathbb{P}_{\alpha(0)}$ -names, we may pick $q_1 \in \mathbb{P}_{\alpha(\varepsilon^*)}$, $q_1 \geq q_0$, $K > K^* \geq i^*$ and $\rho_\varepsilon \in \prod_{i < K} \mathcal{P}(f(i))$ (for $\varepsilon \in U$) such that $q_1 \Vdash_{\mathbb{P}_{\alpha(\varepsilon^*)}} "(\forall \varepsilon \in U)(\varphi_\varepsilon \upharpoonright K = \rho_\varepsilon)"$ and

$$q_1 \Vdash_{\mathbb{P}_{\alpha(\varepsilon^*)}} "(\forall j \geq K)(\forall \zeta \in v)(\forall \varepsilon \in u^\zeta)(|\varphi_\varepsilon(j)| < \frac{f(j)}{|v| \cdot \ell^* \cdot m^*})"$$

Define $q \in \mathbb{P}_{\alpha+1}$ so that

- $\text{dom}(q) = \text{dom}(q_1) \cup \bigcup \{\text{dom}(p^*_\zeta) : \zeta \in v\}$,

- $q \upharpoonright \alpha(\varepsilon^*) = q_1$,
- if $\zeta \in v$ and $\beta \in \text{dom}(p_\zeta^*) \setminus (\alpha(\varepsilon^*) \cup \{\alpha(\varepsilon) : \varepsilon \in u^\zeta\})$, then $q(\beta) = p_\zeta^*(\beta)$,
- if $\zeta \in v$ and $\varepsilon \in u^\zeta \setminus \varepsilon^*$, then $q(\alpha(\varepsilon)) \in {}^K 2$ is such that $p_\zeta''(\alpha(\varepsilon)) = p_\zeta^*(\alpha(\varepsilon)) \triangleleft q(\alpha(\varepsilon))$ and for $i \in [i^*, K)$ we have $q(\alpha(\varepsilon))(i) = 1 - t_\varepsilon^\zeta$,
- $q(\alpha) = (K, m^*, u^+, \sigma^+)$, where $u^+ = \bigcup \{u^\zeta : \zeta \in v\}$ and $\sigma^+ \in \prod_{i < K} \mathcal{P}(f(i))$ is such that $\rho^* \triangleleft \sigma^+$ and $\sigma^+(i) = \bigcup_{\varepsilon \in U} \rho_\varepsilon(i)$ for $i \in [K^*, K)$.

The rest, when $\mu = \kappa$, should be clear.

Let us assume now that $\mu < \kappa$ and again, to argue for 3.9(1)(\otimes)₃, suppose that $n < \omega$, $p_\zeta \in \mathbb{P}_{\alpha+1}$ and $\delta_\zeta < \kappa$ (for $\zeta < \kappa$) are such that $\delta_\zeta \neq \delta_{\zeta'}$ for $\zeta < \zeta' < \kappa$. Passing to stronger conditions we may assume that, for each $\zeta < \kappa$,

$$\alpha \in \text{dom}(p_\zeta) \quad \text{and} \quad p_\zeta \upharpoonright \alpha \Vdash_{\mathbb{P}_\alpha} \text{“ } p_\zeta(\alpha) = (k^\zeta, m^\zeta, u^\zeta, \sigma^\zeta) \text{”}$$

(where $k^\zeta, m^\zeta, u^\zeta, \sigma^\zeta$ are actual objects). For some $\varepsilon^* < \mu$ and k, m, u, σ the set

$$S = \{\zeta < \kappa : \text{dom}(p_\zeta) \subseteq \alpha(\varepsilon^*) \cup \{\alpha\} \ \& \ (k^\zeta, m^\zeta, u^\zeta, \sigma^\zeta) = (k, m, u, \sigma)\}$$

is of size κ . Like before, $\langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \alpha(\varepsilon^*) \rangle$ is a Y-iteration, so we may find $v \subseteq S$, $q_0 \in \mathbb{P}_{\alpha(\varepsilon^*)}$, $m > n$ and $A_\zeta \in [f(m)]^{g(m)}$ for $\zeta \in v$ such that demands listed in $(*)_{10}$ are satisfied. Let $q \in \mathbb{P}_{\alpha+1}$ be such that $\text{dom}(q) = \text{dom}(q_0) \cup \{\alpha\}$ and $q \upharpoonright \alpha \Vdash q(\alpha) = (k, m, u, \sigma)$. \square

Theorem 3.14. *Assume $\kappa < \lambda$ are uncountable regular cardinals such that $\lambda^\kappa = \lambda$. Let $H \subseteq \mathbb{Y}_\kappa^\lambda$ be generic over \mathbf{V} and let $\mathbb{Q} = \langle \mathbb{P}_\alpha, \mathbb{Q}_\alpha : \alpha < \lambda \rangle = \bigcup H \in \mathbf{V}[H]$ and $\mathbb{P}_\lambda = \text{lim}(\mathbb{Q})$. Then \mathbb{P}_λ is a ccc forcing notion with a dense subset of size λ and*

$$\Vdash_{\mathbb{P}_\lambda} \text{“ } \mathbf{MA}_{<\kappa}(\text{ccc}) \text{ and } \mathbf{MA}(\sigma\text{-centered}) \text{ and } \text{add}(\mathcal{I}_f) = \kappa \text{ and } \text{cofin}^-(\mathcal{I}_f) \geq \kappa^+ \text{ and } 2^{\aleph_0} = \lambda \text{”}.$$

Proof. First note that the forcing with $\mathbb{Y}_\kappa^\lambda$ does not add sequences of ordinals of length $< \lambda$ (by Lemma 3.12(5)). Hence in $\mathbf{V}[H]$ we still have that κ, λ are regular cardinals and $\lambda^\kappa = \lambda$.

Let us work in $\mathbf{V}[H]$.

Clearly \mathbb{Q} is a Y-iteration for κ, λ of length λ . Hence \mathbb{P}_λ is a ccc forcing notion, it has a dense subset of size λ and forces that $2^{\aleph_0} = \lambda$ (remember 3.9(1)(\otimes)₂, 3.12(2)). A canonical \mathbb{P}_λ -name η for a real in $\prod_{n < \omega} \mathcal{Z}_n$ (where $\langle \mathcal{Z}_n : n < \omega \rangle \in \mathbf{V}$, $\mathcal{Z}_n \neq \emptyset$) is a sequence $\langle A_n, \pi_n : n < \omega \rangle$ such that each A_n is a maximal antichain in \mathbb{P}_λ , $\pi_n : A_n \rightarrow \mathcal{Z}_n$ and $q \Vdash_{\mathbb{P}_\lambda} \text{“ } \eta(n) = \pi_n(q) \text{”}$ for $q \in A_n$, $n < \omega$. For every \mathbb{P}_λ -name ρ for an element of $\prod_{n < \omega} \mathcal{Z}_n$ there is a canonical name η such that $\Vdash \eta = \rho$. Also, if η is a canonical \mathbb{P}_λ -name for a real, then it is a \mathbb{P}_α -name for some $\alpha < \lambda$.

Let us argue that $\Vdash_{\mathbb{P}_\lambda} \text{cofin}^-(\mathcal{I}_f) \geq \kappa^+$. If not, then for some infinite regular cardinal $\mu \leq \kappa$ and \mathbb{P}_λ -names $\mathcal{I}_\zeta, \varphi_\zeta$ (for $\zeta < \mu$) we have

$$(\otimes)_1 \Vdash_{\mathbb{P}_\lambda} \text{“ } \varphi_\zeta \in \mathcal{S}_f \text{ and } \mathcal{I}_\zeta \subseteq \mathcal{I}_f \text{ is an ideal ”},$$

and for some $p \in \mathbb{P}_\lambda$

$$(\otimes)_2 p \Vdash_{\mathbb{P}_\lambda} \text{“ } \bigcup_{\zeta < \mu} \mathcal{I}_\zeta = \mathcal{I}_f \text{ and } (\forall \zeta < \mu)([\varphi_\zeta] \notin \mathcal{I}_\zeta) \text{”}.$$

We may assume that all φ_ζ are canonical \mathbb{P}_{α_0} -names for some $\alpha_0 < \lambda$.

Suppose now that $\zeta < \mu$ and ϱ_ζ is a canonical \mathbb{P}_λ -name for a real in ${}^\omega 2$. Let $\psi_\zeta^0, \psi_\zeta^1$ be \mathbb{P}_λ -names for elements of \mathcal{S}_f such that for each $n < \omega, i < 2$ we have

$$\Vdash_{\mathbb{P}_\lambda} \text{“ } \psi_\zeta^i(n) = \begin{cases} \varphi_\zeta(n) & \text{if } \varrho_\zeta(n) = i, \\ \emptyset & \text{if } \varrho_\zeta(n) = 1 - i \end{cases} \text{ for each } n < \omega \text{”}.$$

Then $\Vdash_{\mathbb{P}_\lambda} [\varphi_\zeta] = [\psi_\zeta^0] \cup [\psi_\zeta^1]$, so $p \Vdash_{\mathbb{P}_\lambda} \text{“} [\psi_\zeta^0] \notin \mathcal{I}_\zeta \text{ or } [\psi_\zeta^1] \notin \mathcal{I}_\zeta \text{”}$. Let $\tau = \tau(\zeta, \varrho_\zeta)$ be a canonical \mathbb{P}_λ -name for a member of $\{0, 1\}$ such that $p \Vdash \text{“} [\psi_\zeta^\tau] \notin \mathcal{I}_\zeta \text{”}$.

Claim 3.14.1. *For some sequence $\langle \alpha(\zeta), \varrho_\zeta, \psi_\zeta : \zeta < \mu \rangle$ we have*

- (i) $\langle \alpha(\zeta) : \zeta < \mu \rangle \subseteq \lambda$ is strictly increasing with $\alpha_0 \leq \alpha(0)$, and for each $\zeta < \mu$:
- (ii) $\Vdash_{\mathbb{P}_{\alpha(\zeta)}} \mathbb{Q}_{\alpha(\zeta)} = \mathbb{C}$ and ϱ_ζ is the canonical $\mathbb{P}_{\alpha(\zeta)+1}$ -name for the Cohen real in ${}^\omega 2$ added by $\mathbb{Q}_{\alpha(\zeta)}$, and $\tau(\zeta, \varrho_\zeta)$ is a $\mathbb{P}_{\alpha(\zeta+1)}$ -name (for a member of $\{0, 1\}$),
- (iii) ψ_ζ is a $\mathbb{P}_{\alpha(\zeta+1)}$ -name for an element of \mathcal{S}_f such that

$$\Vdash_{\mathbb{P}_{\alpha(\zeta+1)}} \text{“ } \psi_\zeta(n) = \begin{cases} \varphi_\zeta(n) & \text{if } \varrho_\zeta(n) = \tau(\zeta, \varrho_\zeta), \\ \emptyset & \text{if } \varrho_\zeta(n) = 1 - \tau(\zeta, \varrho_\zeta) \end{cases} \text{ for all } n < \omega \text{”},$$

- (iv) if $\alpha^* = \sup\{\alpha(\zeta) : \zeta < \mu\}$, then $\Vdash_{\mathbb{P}_{\alpha^*}} \mathbb{Q}_{\alpha^*} = \mathbb{Q}_\mu^*(\bar{\psi})$, where $\bar{\psi} = \langle \psi_\zeta : \zeta < \mu \rangle$.

Proof of the Claim. We move back to \mathbf{V} and we use a density argument in $\mathbb{Y}_\kappa^\lambda$ above $P = \langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \alpha_0 + 1 \rangle \in \mathbb{Y}_\kappa^\lambda$. Let \mathcal{T} be a $\mathbb{Y}_\kappa^\lambda$ -name for the function $\tau(\cdot, \cdot)$ introduced (in $\mathbf{V}[H]$) earlier. Note that if ϱ is a canonical \mathbb{P}_γ^* -name, $Q^* = \langle \mathbb{P}'_\beta, \mathbb{Q}'_\beta : \beta < \gamma \rangle \in \mathbb{Y}_\kappa^\lambda$ and $\zeta < \mu$, then Q^* forces that (ζ, ϱ) belongs to the domain of \mathcal{T} and $\mathcal{T}(\zeta, \varrho)$ is a $\mathbb{Y}_\kappa^\lambda$ -name for an element of \mathbf{V} .

Suppose that $Q = \langle \mathbb{P}'_\beta, \mathbb{Q}'_\beta : \beta < \alpha \rangle \in \mathbb{Y}_\kappa^\lambda$ is a condition stronger than P (so $\alpha_0 + 1 \leq \alpha$ and $\mathbb{Q}'_\beta = \mathbb{Q}_\beta$ for $\beta \leq \alpha_0$).

By induction on $\zeta < \mu$ we build a sequence $\langle Q_\zeta, \alpha(\zeta), \varrho_\zeta : \zeta < \mu \rangle$ such that

- (\boxtimes)₁ $Q_\zeta = \langle \mathbb{P}'_\beta, \mathbb{Q}'_\beta : \beta \leq \alpha(\zeta) \rangle \in \mathbb{Y}_\kappa^\lambda$ (so $\text{lg}(Q_\zeta) = \alpha(\zeta) + 1 < \lambda$),
- (\boxtimes)₂ for $\zeta < \varepsilon < \mu$ we have

$$Q \leq_{\mathbb{Y}_\kappa^\lambda} Q_\zeta \leq_{\mathbb{Y}_\kappa^\lambda} Q_\varepsilon \quad \text{and} \quad \alpha < \alpha(\zeta) < \alpha(\varepsilon) < \lambda,$$

- (\boxtimes)₃ $\Vdash_{\mathbb{P}'_{\alpha(\zeta)}} \mathbb{Q}'_{\alpha(\zeta)} = \mathbb{C}$ and ϱ_ζ is the canonical $\mathbb{P}'_{\alpha(\zeta)+1}$ -name for the Cohen real in ${}^\omega 2$ added by $\mathbb{Q}'_{\alpha(\zeta)}$,

(\boxtimes)₄ $Q_{\zeta+1}$ decides the value of $\underline{T}(\zeta, \zeta_\zeta)$ and forces (in $\mathbb{Y}_\kappa^\lambda$) that it is a $\mathbb{P}'_{\alpha(\zeta+1)}$ -name.

The construction is clearly possible by Lemma 3.12(2,4). Then letting $\alpha^* = \sup(\alpha(\zeta) : \zeta < \mu)$ we have that $Q_\mu = \langle \mathbb{P}_\beta, \mathbb{Q}'_\beta : \beta < \alpha^* \rangle \in \mathbb{Y}_\kappa^\lambda$ is a condition stronger than all Q_ζ (for $\zeta < \mu$); remember 3.12(4) again. Moreover, if names ψ_ζ are defined as in clause (iii), and τ_ζ is the value forced to $\underline{T}(\zeta, \zeta_\zeta)$ by $Q_{\zeta+1}$ (see (\boxtimes)₄ above) and ζ_ζ are as described in (\boxtimes)₃, then the assumptions of Lemma 3.13 are satisfied. Therefore $Q^* = Q_\mu \wedge \langle \mathbb{P}_{\alpha^*}, \mathbb{Q}_\mu^*(\bar{\psi}) \rangle \in \mathbb{Y}_\kappa^\lambda$ is a condition stronger than Q . This condition forces in $\mathbb{Y}_\kappa^\lambda$ that $\langle \alpha(\zeta), \zeta_\zeta, \psi_\zeta : \zeta < \mu \rangle$ satisfies the demands (i)–(iv). \square

Let $\alpha(\zeta), \zeta_\zeta, \psi_\zeta$ (for $\zeta < \mu$) and α^* be in Claim 3.14.1(i–iv), so in particular $\Vdash_{\mathbb{P}_{\alpha^*}} \mathbb{Q}_{\alpha^*} = \mathbb{Q}_\mu^*(\bar{\psi})$. Let ζ be a \mathbb{P}_{α^*+1} -name for the null slalom added by \mathbb{Q}_{α^*} (see Definition 3.7). It follows from Proposition 3.8 that

$$\Vdash_{\mathbb{P}_\lambda} \bigcup_{\zeta < \mu} [\psi_\zeta] \subseteq [\zeta] \in \mathcal{I}_f,$$

and hence, by (\otimes)₂, $p \Vdash_{\mathbb{P}_\lambda} (\exists \varepsilon < \mu) (\bigcup_{\zeta < \mu} [\psi_\zeta] \in \mathcal{I}_\varepsilon)$. Pick $\varepsilon^* < \mu$ and a condition $q \in \mathbb{P}_\lambda$ stronger than p such that $q \Vdash_{\mathbb{P}_\lambda} \bigcup_{\zeta < \mu} [\psi_\zeta] \in \mathcal{I}_{\varepsilon^*}$. Then also $q \Vdash [\psi_{\varepsilon^*}] \in \mathcal{I}_{\varepsilon^*}$ (remember (\otimes)₁), but this contradicts the choice of $\tau(\varepsilon^*, \zeta_{\varepsilon^*})$ and ψ_{ε^*} .

To argue that $\Vdash_{\mathbb{P}_\lambda} \mathbf{MA}_{<\kappa}(\text{ccc})$ note that every \mathbb{P}_λ -name \mathbb{Q} for a ccc forcing notion on some $\gamma^* < \kappa$ is actually a \mathbb{P}_α -name for some $\alpha < \lambda$. Therefore by the standard density argument in $\mathbb{Y}_\kappa^\lambda$, for unboundedly many $\beta < \lambda$ we have $\Vdash_{\mathbb{P}_\beta} \mathbb{Q}_\beta = \mathbb{Q}$ (remember 3.12(2)). Similarly we may justify that $\Vdash_{\mathbb{P}_\lambda} \mathbf{MA}(\sigma\text{-centered})$.

It follows from Lemma 3.10 that $\Vdash_{\mathbb{P}_\lambda} \text{add}(\mathcal{I}_f) \leq \kappa$. Since $\Vdash_{\mathbb{P}_\lambda} \mathbf{MA}_{<\kappa}(\text{ccc})$ we easily see that the equality is forced. \square

Corollary 3.15. *It is consistent that $\text{add}(\mathcal{I}_f) = \aleph_1$ and $\text{cofin}^-(\mathcal{I}_f) = \text{cofin}(\mathcal{I}_f) = \aleph_2$.*

4. OPEN PROBLEMS

Can we get a result parallel to Corollary 3.15 for the null and/or meager ideals? Or even better:

Problem 4.1. Let \mathcal{I} be either the meager ideal \mathcal{M} or the null ideal \mathcal{N} . Is it consistent that

$$\text{add}(\mathcal{I}) < \text{cofin}(\mathcal{I}) < \text{cof}(\mathcal{I}) \quad ?$$

The method used in the proof of 3.14, 3.15 gives the consistency of $\text{add}(\mathcal{I}_f) \leq \kappa$ & $\kappa^+ \leq \text{cofin}^-(\mathcal{I}_f)$. Can the gap be bigger?

Problem 4.2. Is it consistent that $\text{add}(\mathcal{I}_f) = \aleph_\alpha < \aleph_{\alpha+\omega} \leq \text{cofin}^-(\mathcal{I}_f)$?

modified:2013-12-11

1022 revision:2013-12-11

The cardinal invariant cofin introduced by Borodulin–Nadzieja and Głab has several natural relatives (or variants), some were listed in Definition 2.1. Are those coefficients distinct or they are equivalent within the realm of nice σ -ideals?

Problem 4.3. Is it consistent that for some Borel σ -ideal \mathcal{I} on ${}^\omega 2$ we have $\text{cofin}^*(\mathcal{I}) < \text{cofin}^-(\mathcal{I})$? Or $\text{cofin}^-(\mathcal{I}) < \text{cofin}(\mathcal{I})$? Or $\text{cofin}(\mathcal{I}) < \text{cofin}^+(\mathcal{I})$?

Acknowledgements. Both authors acknowledge support from the United States-Israel Binational Science Foundation (Grant no. 2010405). This is publication 1022 of the second author.

REFERENCES

- [1] Tomek Bartoszyński and Haim Judah. *Set Theory: On the Structure of the Real Line*. A K Peters, Wellesley, Massachusetts, 1995.
- [2] Tomek Bartoszyński and Masaru Kada. Hechler’s theorem for the meager ideal. *Topology Appl.*, 146/147:429–435, 2005.
- [3] Piotr Borodulin-Nadzieja and Szymon Głab. Ideals with bases of unbounded Borel complexity. *MLQ Math. Log. Q.*, 57:582–590, 2011.
- [4] Maxim R. Burke and Masaru Kada. Hechler’s theorem for the null ideal. *Arch. Math. Logic*, 43:703–722, 2004.
- [5] Jacek Cichoń and Janusz Pawlikowski. On Ideals of Subsets of the Plane and on Cohen Reals. *The Journal of Symbolic Logic*, 51:560–569, 1986.
- [6] Andrzej Roslanowski and Saharon Shelah. Monotone hulls for $\mathcal{N} \cap \mathcal{M}$. *Periodica Mathematica Hungarica*, accepted. arXiv:1007.5368.
- [7] Saharon Shelah. Whitehead groups may not be free, even assuming CH. II. *Israel Journal of Mathematics*, 35:257–285, 1980.
- [8] Saharon Shelah. *Proper and improper forcing*. Perspectives in Mathematical Logic. Springer, 1998.
- [9] Eric K. van Douwen, J. Donald Monk, and Matatyahu Rubin. Some questions about Boolean algebras. *Algebra Universalis*, 11:220–243, 1980.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEBRASKA AT OMAHA, OMAHA, NE 68182-0243, USA

E-mail address: roslanow@member.ams.org

URL: <http://www.unomaha.edu/logic>

INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM, 91904 JERUSALEM, ISRAEL, AND DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW BRUNSWICK, NJ 08854, USA

E-mail address: shelah@math.huji.ac.il

URL: <http://www.math.rutgers.edu/~shelah>