

# ON THE COFINALITY OF THE SPLITTING NUMBER

ALAN DOW AND SAHARON SHELAH

ABSTRACT. The splitting number  $\mathfrak{s}$  can be singular. The key method is to construct a forcing poset with finite support matrix iterations of ccc posets introduced by Blass and the second author [*Ultrafilters with small generating sets*, Israel J. Math., **65**, (1989)]

## 1. INTRODUCTION

The cardinal invariants of the continuum discussed in this article are very well known (see [5, van Douwen, p111]) so we just give a brief reminder. They deal with the mod finite ordering of the infinite subsets of the integers. A set  $S \subset \omega$  is *unsplit* by a family  $\mathcal{Y} \subset [\omega]^{\aleph_0}$  if  $S$  mod finite is contained in one member of  $\{Y, \omega \setminus Y\}$  for each  $Y \in \mathcal{Y}$ . The splitting number  $\mathfrak{s}$  is the minimum cardinal of a family  $\mathcal{Y}$  for which there is no infinite set unsplit by  $\mathcal{Y}$  (equivalently every  $S \in [\omega]^{\aleph_0}$  is *split* by some member of  $\mathcal{Y}$ ). It is mentioned in [2] that it is currently unknown if  $\mathfrak{s}$  can be a singular cardinal.

**Proposition 1.1.** *The cofinality of the splitting number is not countable.*

*Proof.* Assume that  $\theta$  is the supremum of  $\{\kappa_n : n \in \omega\}$  and that there is no splitting family of cardinality less than  $\theta$ . Let  $\mathcal{Y} = \{Y_\alpha : \alpha < \theta\}$  be a family of subsets of  $\omega$ . Let  $S_0 = \omega$  and by induction on  $n$ , choose an infinite subset  $S_{n+1}$  of  $S_n$  so that  $S_{n+1}$  is not split by the family  $\{Y_\alpha : \alpha < \kappa_n\}$ . If  $S$  is any pseudointersection of  $\{S_n : n \in \omega\}$ , then  $S$  is not split by any member of  $\mathcal{Y}$ .  $\square$

---

*Date:* October 29, 2017.

*1991 Mathematics Subject Classification.* 03E15 .

*Key words and phrases.* splitting number, cardinal invariants of the continuum, matrix forcing.

The research of the first author was supported by the NSF grant No. NSF-DMS 1501506. The research of the second author was supported by the United States-Israel Binational Science Foundation (BSF Grant no. 2010405), and by the NSF grant No. NSF-DMS 1101597.

One can easily generalize the previous result and proof to show that the cofinality of the splitting number is at least  $\mathfrak{t}$ . In this paper we prove the following.

**Theorem 1.2.** *If  $\kappa$  is any uncountable regular cardinal, then there is a  $\lambda > \kappa$  with  $\text{cf}(\lambda) = \kappa$  and a ccc forcing  $\mathbb{P}$  satisfying that  $\mathfrak{s} = \lambda$  in the forcing extension.*

To prove the theorem, we construct  $\mathbb{P}$  using matrix iterations.

## 2. A SPECIAL SPLITTING FAMILY

**Definition 2.1.** *Let us say that a family  $\{x_i : i \in I\} \subset [\omega]^\omega$  is  $\theta$ -Luzin (for an uncountable cardinal  $\theta$ ) if for each  $J \in [I]^\theta$ ,  $\bigcap\{x_i : i \in J\}$  is finite and  $\bigcup\{x_i : i \in J\}$  is cofinite.*

Clearly a family is  $\theta$ -Luzin if every  $\theta$ -sized subfamily is  $\theta$ -Luzin. We leave to the reader the easy verification that for a regular uncountable cardinal  $\theta$ , each  $\theta$ -Luzin family is a splitting family. A poset being  $\theta$ -Luzin preserving will have the obvious meaning. For example, any poset of cardinality less than a regular cardinal  $\theta$  is  $\theta$ -Luzin preserving.

**Lemma 2.2.** *If  $\theta$  is a regular uncountable cardinal then any ccc finite support iteration of  $\theta$ -Luzin preserving posets is again  $\theta$ -Luzin preserving.*

*Proof.* We prove this by induction on the length of the iteration. Fix any  $\theta$ -Luzin family  $\{x_i : i \in I\}$  and let  $\langle\langle \mathbb{P}_\alpha : \alpha \leq \gamma \rangle\rangle, \langle\langle \dot{\mathbb{Q}}_\alpha : \alpha < \gamma \rangle\rangle$  be a finite support iteration of ccc posets satisfying that  $\mathbb{P}_\alpha$  forces that  $\dot{\mathbb{Q}}_\alpha$  is ccc and  $\theta$ -Luzin preserving, for all  $\alpha < \gamma$ .

If  $\gamma$  is a successor ordinal  $\beta + 1$ , then for any  $\mathbb{P}_\beta$ -generic filter  $G_\beta$ , the family  $\{x_i : i \in I\}$  is a  $\theta$ -Luzin family in  $V[G_\beta]$ . By the hypothesis on  $\dot{\mathbb{Q}}_\beta$ , this family remains  $\theta$ -Luzin after further forcing by  $\dot{\mathbb{Q}}_\beta$ .

Now we assume that  $\alpha$  is a limit. Let  $\dot{J}_0$  be any  $\mathbb{P}_\gamma$ -name of a subset of  $I$  and assume that  $p \in \mathbb{P}_\gamma$  forces that  $|\dot{J}_0| = \theta$ . We must produce a  $q < p$  that forces that  $\dot{J}_0$  is as in the definition of  $\theta$ -Luzin. There is a set  $J_1 \subset I$  of cardinality  $\theta$  satisfying that, for each  $i \in J_1$ , there is a  $p_i < p$  with  $p_i \Vdash i \in \dot{J}_0$ . The case when the cofinality of  $\alpha$  not equal to  $\theta$  is almost immediate. There is a  $\beta < \alpha$  such that  $J_2 = \{i \in J_1 : p_i \in \mathbb{P}_\beta\}$  has cardinality  $\theta$ . There is a  $\mathbb{P}_\beta$ -generic filter  $G_\beta$  such that  $J_3 = \{i \in J_2 : p_i \in G_\beta\}$  has cardinality  $\theta$ . By the induction hypothesis, the family  $\{x_i : i \in I\}$  is  $\theta$ -Luzin in  $V[G_\beta]$  and so we have that  $\bigcap\{x_i : i \in J_3\}$  is finite and  $\bigcup\{x_i : i \in J_3\}$  is co-finite. Choose any  $q < p$  in  $G_\beta$  and a name  $\dot{J}_3$  for  $J_3$  so that  $q$  forces this

property for  $\dot{J}_3$ . Since  $q$  forces that  $\dot{J}_3 \subset \dot{J}_0$ , we have that  $q$  forces the same property for  $\dot{J}_0$ .

Finally we assume that  $\alpha$  has cofinality  $\theta$ . Naturally we may assume that the collection  $\{\text{dom}(p_i) : i \in J_1\}$  forms a  $\Delta$ -system with root contained in some  $\beta < \alpha$ . Again, we may choose a  $\mathbb{P}_\beta$ -generic filter  $G_\beta$  satisfying that  $J_2 = \{i \in J_1 : p_i \upharpoonright \beta \in G_\beta\}$  has cardinality  $\theta$ . In  $V[G_\beta]$ , let  $\{J_{2,\xi} : \xi \in \omega_1\}$  be a partition of  $J_2$  into pieces of size  $\theta$ . For each  $\xi \in \omega_1$ , apply the induction hypothesis in the model  $V[G_\beta]$ , and so we have that  $\bigcap\{x_i : i \in J_{2,\xi}\}$  is finite and  $\bigcup\{x_i : i \in J_{2,\xi}\}$  is co-finite. For each  $\xi \in \omega_1$  let  $m_\xi$  be an integer large enough so that  $\bigcap\{x_i : i \in J_{2,\xi}\} \subset m_\xi$  and  $\bigcup\{x_i : i \in J_{2,\xi}\} \supset \omega \setminus m_\xi$ . Let  $m$  be any integer such that  $m_\xi = m$  for uncountably many  $\xi$ . Choose any condition  $\bar{p} \in \mathbb{P}_\alpha$  so that  $\bar{p} \upharpoonright \beta \in G_\beta$ . We prove that for each  $n > m$  there is a  $\bar{p}_n < \bar{p}$  so that  $\bar{p}_n \Vdash n \notin \bigcap\{x_i : i \in \dot{I}\}$  and  $\bar{p}_n \Vdash n \in \bigcup\{x_i : i \in \dot{I}\}$ . Choose any  $\xi \in \omega_1$  so that  $m_\xi = m$  and  $\text{dom}(p_i) \cap \text{dom}(\bar{p}) \subset \beta$  for all  $i \in J_{2,\xi}$ . Now choose any  $i_0 \in J_{2,\xi}$  so that  $n \notin x_{i_0}$ . Next choose a distinct  $\xi'$  with  $m_{\xi'} = m$  so that  $\text{dom}(p_i) \cap (\text{dom}(\bar{p}) \cup \text{dom}(p_{i_0})) \subset \beta$  for all  $i \in J_{2,\xi'}$ . Now choose  $i_1 \in J_{2,\xi'}$  so that  $n \in x_{i_1}$ . We now have that  $\bar{p} \cup p_{i_0} \cup p_{i_1}$  is a condition that forces  $\{i_0, i_1\} \subset \dot{I}$ .  $\square$

Next we introduce a  $\sigma$ -centered poset that will render a given family non-splitting.

**Definition 2.3.** For a filter  $\mathfrak{D}$  on  $\omega$ , we define the Laver style poset  $\mathbb{L}(\mathfrak{D})$  to be the set of trees  $T \subset \omega^{<\omega}$  with the property that  $T$  has a minimal branching node  $\text{stem}(T)$  and for all  $\text{stem}(T) \subseteq t \in T$ , the branching set  $\{k : t \hat{\smallfrown} k \in T\}$  is an element of  $\mathfrak{D}$ . If  $\mathfrak{D}$  is a filter base for a filter  $\mathfrak{D}^*$ , then  $\mathbb{L}(\mathfrak{D})$  will also denote  $\mathbb{L}(\mathfrak{D}^*)$ .

The name  $\dot{L} = \{(k, T) : (\exists t) t \hat{\smallfrown} k \subset \text{stem}(T)\}$  will be referred to as the canonical name for the real added by  $\mathbb{L}(\mathfrak{D})$ .

If  $\mathfrak{D}$  is a principal (fixed) ultrafilter on  $\omega$ , then  $\mathbb{L}(\mathfrak{D})$  has a minimum element and so is forcing isomorphic to the trivial poset. If  $\mathfrak{D}$  is principal but not an ultrafilter, then  $\mathbb{L}(\mathfrak{D})$  is isomorphic to Cohen forcing. If  $\mathfrak{D}$  is a free filter, then  $\mathbb{L}(\mathfrak{D})$  adds a dominating real and has similarities to Hechler forcing. As usual, for a filter (or filter base)  $\mathfrak{D}$  of subsets of  $\omega$ , we use  $\mathfrak{D}^+$  to denote the set of all subsets of  $\omega$  that meet every member of  $\mathfrak{D}$ .

**Definition 2.4.** If  $E$  is a dense subset of  $\mathbb{L}(\mathfrak{D})$ , then a function  $\rho_E$  from  $\omega^{<\omega}$  into  $\omega_1$  is a rank function for  $E$  if  $\rho_E(t) = 0$  if and only if  $t = \text{stem}(T)$  for some  $T \in E$ , and for all  $t \in \omega^{<\omega}$  and  $0 < \alpha \in \omega_1$ ,  $\rho_E(t) \leq \alpha$  providing the set  $\{k \in \omega : \rho_E(t \hat{\smallfrown} k) < \alpha\}$  is in  $\mathfrak{D}^+$ .

modified:2017-10-29

revision:2017-10-09

1127

When  $\mathfrak{D}$  is a free filter, then  $\mathbb{L}(\mathfrak{D})$  has cardinality  $\mathfrak{c}$ , but nevertheless, if  $\mathfrak{D}$  has a base of cardinality less than a regular cardinal  $\theta$ ,  $\mathbb{L}(\mathfrak{D})$  is  $\theta$ -Luzin preserving.

**Lemma 2.5.** *If  $\mathfrak{D}$  is a free filter on  $\omega$  and if  $\mathfrak{D}$  has a base of cardinality less than a regular uncountable cardinal  $\theta$ , then  $\mathbb{L}(\mathfrak{D})$  is  $\theta$ -Luzin preserving.*

*Proof.* Let  $\{x_i : i \in \theta\}$  be a  $\theta$ -Luzin family with  $\theta$  as in the Lemma. Let  $\dot{J}$  be a  $\mathbb{L}(\mathfrak{D})$ -name of a subset of  $\theta$ . We prove that if  $\bigcap\{x_i : i \in \dot{J}\}$  is not finite, then  $\dot{J}$  is bounded in  $\theta$ . By symmetry, it will also prove that if  $\bigcup\{x_i : i \in \dot{J}\}$  is not cofinite, then  $\dot{J}$  is bounded in  $\theta$ . Let  $\dot{y}$  be the  $\mathbb{L}(\mathfrak{D})$ -name of the intersection, and let  $T_0$  be any member of  $\mathbb{L}(\mathfrak{D})$  that forces that  $\dot{y}$  is infinite. Let  $M$  be any  $< \theta$ -sized elementary submodel of  $H((2^{\mathfrak{c}})^+)$  such that  $T_0, \mathfrak{D}, \dot{J}$ , and  $\{x_i : i \in \theta\}$  are all members of  $M$  and such that  $M \cap \mathfrak{D}$  contains a base for  $\mathfrak{D}$ . Let  $i_M = \sup(M \cap \theta)$ . If  $x \in M \cap [\omega]^\omega$ , then  $I_x = \{i \in \theta : x \subset x_i\}$  is an element of  $M$  and has cardinality less than  $\theta$ . Therefore, if  $i \in \theta \setminus i_M$ , then  $x_i$  does not contain any infinite subset of  $\omega$  that is an element of  $M$ . We prove that  $x_i$  is forced by  $T_0$  to also not contain  $\dot{y}$ . This will prove that  $\dot{J}$  is bounded by  $i_M$ . Let  $T_1 < T_0$  be any condition in  $\mathbb{L}(\mathfrak{D})$  and let  $t_1 = \text{stem}(T_1)$ . We show that  $T_1$  does not force that  $x_i \supset \dot{y}$ . We define the relation  $\Vdash_w$  on  $T_0 \times \omega$  to be the set

$$\{(t, n) \in T_0 \times \omega : \text{there is no } T \leq T_0, \text{stem}(T) = t, \text{s.t. } T \Vdash_w n \notin \dot{y}\}.$$

For convenience we may write, for  $T \leq T_0$ ,  $T \Vdash_w n \in \dot{y}$  providing  $(\text{stem}(T), n)$  is in  $\Vdash_w$ , and this is equivalent to the relation that  $T$  has no stem preserving extension forcing that  $n$  is not in  $\dot{y}$ . Let  $T_2 \in M$  be any extension of  $T_0$  with stem  $t_1$ . Let  $L$  denote the set of  $\ell \in \omega$  such that  $T_2 \Vdash_w \ell \in \dot{y}$ . If  $L$  is infinite, then, since  $L \in M$ , there is an  $\ell \in L \setminus x_i$ . This implies that  $T_1$  does not force  $x_i \supset \dot{y}$ , since  $T_2 \Vdash_w j \in \dot{y}$  implies that  $T_1$  fails to force that  $\ell \notin \dot{y}$ .

Therefore we may assume that  $L$  is finite and let  $\ell$  be the maximum of  $L$ . Define the set  $E \subset \mathbb{L}(\mathfrak{D})$  according to  $T \in E$  providing that either  $t_1 \notin T$  or there is a  $j > \ell$  such that  $T \Vdash_w j \in \dot{y}$ . Again this set  $E$  is in  $M$  and is easily seen to be a dense subset of  $\mathbb{L}(\mathfrak{D})$ . By the choice of  $\ell$ , we note that  $\rho_E(t_1) > 0$ . If  $\rho_E(t_1) > 1$ , then the set  $\{k \in \omega : 0 < \rho_E(t_1 \hat{\ } k) < \rho_E(t_1)\}$  is in  $\mathfrak{D}^+$  and so there is a  $k_1$  in this set such that  $t_1 \hat{\ } k_1 \in T_1 \cap T_2$ . By a finite induction, we can choose an extension  $t_2 \supseteq t_1$  so that  $t_2 \in T_1 \cap T_2$  and  $\rho_E(t_2) = 1$ . Now, there is a set  $D \in \mathfrak{D} \cap M$  contained in  $\{k : t_2 \hat{\ } k \in T_1 \cap T_2\}$  since  $M$  contains a base for  $\mathfrak{D}$ . Also,  $D_E = \{k \in D : \rho_E(t_2 \hat{\ } k) = 0\}$  is in  $\mathfrak{D}^+$ . For each  $k \in D_E$ , choose the minimal  $j_k$  so that  $T_2 \hat{\ } k \Vdash_w j_k \in \dot{y}$ . The set

$\{j_k : k \in D_E\}$  is an element of  $M$ . This set is not finite because if it were then there would be a single  $j$  such that  $\{k \in D_E : j_k = j\} \in \mathcal{D}^+$ , which would contradict that  $\rho_E(t_2) > 0$ . This means that there is a  $k \in D_E^+$  with  $j_k \notin x_i$ , and again we have shown that  $T_1$  fails to force that  $x_i$  contains  $\dot{y}$ .  $\square$

### 3. MATRIX ITERATIONS

The terminology “matrix iterations” is used in [3], see also forthcoming preprint (F1222) from the second author. The paper [3] nicely expands on the method of matrix iterated forcing first introduced in [1].

Let us recall that a poset  $(P, <_P)$  is a complete suborder of a poset  $(Q, <_Q)$  providing  $P \subset Q$ ,  $<_P \subset <_Q$ , and each maximal antichain of  $(P, <_P)$  is also a maximal antichain of  $(Q, <_Q)$ . Note that it follows that incomparable members of  $(P, <_P)$  are still incomparable in  $(Q, <_Q)$ , i.e.  $p_1 \perp_P p_2$  implies  $p_1 \perp_Q p_2$ . We use the notation  $(P, <_P) \triangleleft (Q, <_Q)$  to abbreviate the complete suborder relation, and similarly use  $P \triangleleft Q$  if  $<_P$  and  $<_Q$  are clear from the context. An element  $p$  of  $P$  is a reduction of  $q \in Q$  if  $r \not\perp_Q q$  for each  $r <_P p$ . If  $P \subset Q$ ,  $<_P \subset <_Q$ ,  $\perp_P \subset \perp_Q$ , and each element of  $Q$  has a reduction in  $P$ , then  $P \triangleleft Q$ . The reason is that if  $A \subset P$  is a maximal antichain and  $p \in P$  is a reduction of  $q \in Q$ , then there is an  $a \in A$  and an  $r$  less than both  $p$  and  $a$  in  $P$ , such that  $r \not\perp_Q q$ .

**Definition 3.1.** *We will say that an object  $\mathbf{P}$  is a matrix iteration if there is an infinite cardinal  $\kappa$  and an ordinal  $\gamma$  (thence a  $(\kappa, \gamma)$ -matrix iteration) such that  $\mathbf{P} = \langle \langle \mathbb{P}_{i,\alpha}^{\mathbf{P}} : i \leq \kappa, \alpha \leq \gamma \rangle, \langle \dot{\mathbb{Q}}_{i,\alpha}^{\mathbf{P}} : i \leq \kappa, \alpha < \gamma \rangle \rangle$  where, for each  $(i, \alpha) \in \kappa + 1 \times \gamma$  and each  $j < i$ ,*

- (1)  $\mathbb{P}_{j,\alpha}^{\mathbf{P}}$  is a complete suborder of the poset  $\mathbb{P}_{i,\alpha}^{\mathbf{P}}$  (i.e.  $\mathbb{P}_{j,\alpha}^{\mathbf{P}} \triangleleft \mathbb{P}_{i,\alpha}^{\mathbf{P}}$ ),
- (2)  $\dot{\mathbb{Q}}_{i,\alpha}^{\mathbf{P}}$  is a  $\mathbb{P}_{i,\alpha}^{\mathbf{P}}$ -name of a ccc poset,  $\mathbb{P}_{i,\alpha+1}^{\mathbf{P}}$  is equal to  $\mathbb{P}_{i,\alpha}^{\mathbf{P}} * \dot{\mathbb{Q}}_{i,\alpha}^{\mathbf{P}}$ ,
- (3) for limit  $\delta \leq \gamma$ ,  $\mathbb{P}_{i,\delta}^{\mathbf{P}}$  is equal to the union of the family  $\{\mathbb{P}_{i,\beta}^{\mathbf{P}} : \beta < \delta\}$
- (4)  $\mathbb{P}_{\kappa,\alpha}^{\mathbf{P}}$  is the union of the chain  $\{\mathbb{P}_{j,\alpha}^{\mathbf{P}} : j < \kappa\}$ .

When the context makes it clear, we omit the superscript  $\mathbf{P}$  when discussing a matrix iteration. Throughout the paper,  $\kappa$  will be a fixed uncountable regular cardinal

**Definition 3.2.** *A sequence  $\vec{\lambda}$  is  $\kappa$ -tall if  $\vec{\lambda} = \langle \mu_\xi, \lambda_\xi : \xi < \kappa \rangle$  is a sequence of pairs of regular cardinals satisfying that  $\mu_0 = \omega < \kappa < \lambda_0$  and, for  $0 < \eta < \kappa$ ,  $\mu_\eta < \lambda_\eta$  where  $\mu_\eta = (2^{\sup\{\lambda_\xi : \xi < \eta\}})^+$ .*

Also for the remainder of the paper, we fix a  $\kappa$ -tall sequence  $\vec{\lambda}$  and  $\lambda$  will denote the supremum of the set  $\{\lambda_\xi : \xi \in \kappa\}$ . For simpler notation, whenever we discuss a matrix iteration  $\underline{\mathbf{P}}$  we shall henceforth assume that it is a  $(\kappa, \gamma)$ -matrix iteration for some ordinal  $\gamma$ . We may refer to a forcing extension by  $\underline{\mathbf{P}}$  as an abbreviation for the forcing extension by  $\mathbb{P}_{\kappa, \gamma}^{\underline{\mathbf{P}}}$ .

For any poset  $P$ , any  $P$ -name  $\dot{D}$ , and  $P$ -generic filter  $G$ ,  $\dot{D}[G]$  will denote the valuation of  $\dot{D}$  by  $G$ . For any ground model  $x$ ,  $\check{x}$  denotes the canonical name so that  $\check{x}[G] = x$ . When  $x$  is an ordinal (or an integer) we will suppress the accent in  $\check{x}$ . A  $P$ -name  $\dot{D}$  of a subset of  $\omega$  will be said to be *nice* or *canonical* if for each integer  $j \in \omega$ , there is an antichain  $A_j$  such that  $\dot{D} = \bigcup \{\{j\} \times A_j : j \in \omega\}$ . We will say that  $\dot{\mathcal{D}}$  is a nice  $P$ -name of a family of subsets of  $\omega$  just to mean that  $\dot{\mathcal{D}}$  is a collection of nice  $P$ -names of subsets of  $\omega$ . We will use  $(\dot{\mathcal{D}})_P$  if we need to emphasize that we mean the  $P$ -name. Similarly if we say that  $\dot{\mathcal{D}}$  is a nice  $P$ -name of a filter (base) we mean that  $\dot{\mathcal{D}}$  is a nice  $P$ -name such that, for each  $P$ -generic filter, the collection  $\{\dot{D}[G] : \dot{D} \in \dot{\mathcal{D}}\}$  is a filter (base) of infinite subsets of  $\omega$ .

Following these conventions, the following notation will be helpful.

**Definition 3.3.** For a  $(\kappa, \gamma)$ -matrix  $\underline{\mathbf{P}}$  and  $i < \kappa$ , we let  $\mathbb{B}_{i, \gamma}^{\underline{\mathbf{P}}}$  denote the set of all nice  $\mathbb{P}_{i, \gamma}^{\underline{\mathbf{P}}}$ -names of subsets of  $\omega$ . We note that this then is the nice  $\mathbb{P}_{i, \gamma}^{\underline{\mathbf{P}}}$ -name for the power set of  $\omega$ . As usual, when possible we suppress the  $\underline{\mathbf{P}}$  superscript.

For a nice  $\underline{\mathbf{P}}$ -name  $\dot{\mathcal{D}}$  of a filter (or filter base) of subsets of  $\omega$ , we let  $(\dot{\mathcal{D}})^+$  denote the set of all nice  $\underline{\mathbf{P}}$ -names that are forced to meet every member of  $\dot{\mathcal{D}}$ . It follows that  $(\dot{\mathcal{D}})^+$  is the nice  $\underline{\mathbf{P}}$ -name for the usual defined notion  $(\dot{\mathcal{D}})^+$  in the forcing extension by  $\underline{\mathbf{P}}$ . We let  $\langle \dot{\mathcal{D}} \rangle$  denote the nice  $\underline{\mathbf{P}}$ -name of the filter generated by  $\dot{\mathcal{D}}$ . We use the same notational conventions if, for some poset  $\mathbb{P}$ ,  $\dot{\mathcal{D}}$  is a nice  $\mathbb{P}$ -name of a filter (or filter base) of subsets of  $\omega$ .

The main idea for controlling the splitting number in the extension by  $\underline{\mathbf{P}}$  will involve having many of the subposets being  $\theta$ -Luzin preserving for  $\theta \in \{\lambda_\xi : \xi \in \kappa\}$ . Motivated by the fact that posets of the form  $\mathbb{L}(\dot{\mathcal{D}})$  (our proposed iterands) are  $\theta$ -Luzin preserving when  $\dot{\mathcal{D}}$  is sufficiently small we adopt the name  $\vec{\lambda}$ -thin for this next notion.

**Definition 3.4.** For a  $\kappa$ -tall sequence  $\vec{\lambda}$ , we will say that a  $(\kappa, \gamma)$ -matrix-iteration  $\underline{\mathbf{P}}$  is  $\vec{\lambda}$ -thin providing that for each  $\xi < \kappa$  and  $\alpha \leq \gamma$ ,  $\mathbb{P}_{\xi, \alpha}^{\underline{\mathbf{P}}}$  is  $\lambda_\xi$ -Luzin preserving.

Now we combine the notion of  $\vec{\lambda}$ -thin matrix-iteration with Lemma 2.2. We adopt Kunen's notation that for a set  $I$ ,  $\text{Fn}(I, 2)$  denotes the usual poset for adding Cohen reals (finite partial functions from  $I$  into 2 ordered by superset).

**Lemma 3.5.** *Suppose that  $\mathbf{P}$  is a  $\vec{\lambda}$ -thin  $(\kappa, \gamma)$ -matrix iteration for some  $\kappa$ -tall sequence  $\vec{\lambda}$ . Further suppose that  $\dot{Q}_{i,0}$  is the  $\mathbb{P}_{i,0}$ -name of the poset  $\text{Fn}(\lambda_\xi, 2)$  for each  $\xi \in \kappa$ , and therefore  $\mathbb{P}_{\kappa,1}$  is isomorphic to  $\text{Fn}(\lambda, 2)$ . Let  $\dot{g}$  denote the generic function from  $\lambda$  onto 2 added by  $\mathbb{P}_{\kappa,1}$  and, for  $i < \lambda$ , let  $\dot{x}_i$  be the canonical name of the set  $\{n \in \omega : \dot{g}(i+n) = 1\}$ . Then the family  $\{\dot{x}_i : i < \lambda\}$  is forced by  $\mathbf{P}$  to be a splitting family.*

*Proof.* Let  $G_{\kappa,\gamma}$  be a  $\mathbb{P}_{\kappa,\gamma}$ -generic filter. For each  $\xi \in \kappa$  and  $\alpha \leq \gamma$ , let  $G_{\xi,\alpha} = G_{\kappa,\gamma} \cap \mathbb{P}_{\xi,\alpha}$ . Let  $\dot{y}$  be any nice  $\mathbb{P}_{\kappa,\gamma}$ -name for a subset of  $\omega$ . Since  $\dot{y}$  is a countable name, we may choose a  $\xi < \kappa$  so that  $\dot{y}$  is a  $\mathbb{P}_{\xi,\gamma}$ -name. It is easily shown, and very well-known, that the family  $\{\dot{x}_i : i < \lambda_\xi\}$  is forced by  $\mathbb{P}_{\xi,1}$  (i.e.  $\text{Fn}(\lambda_\xi, 2)$ ) to be a  $\lambda_\xi$ -Luzin family. By the hypothesis that  $\mathbf{P}$  is  $\vec{\lambda}$ -thin, we have, by Lemma 2.2, that  $\{\dot{x}_i : i < \lambda_\xi\}$  is still  $\lambda_\xi$ -Luzin in  $V[G \cap \mathbb{P}_{\xi,\gamma}]$ . Since  $\dot{y}$  is a  $\mathbb{P}_{\xi,\gamma}$ -name, there is an  $i < \lambda_\xi$  such that  $\dot{y}[G_{\xi,\gamma}] \cap \dot{x}_i[G_{\xi,\gamma}]$  and  $\dot{y}[G_{\xi,\gamma}] \setminus \dot{x}_i[G_{\xi,\gamma}]$  are infinite.  $\square$

#### 4. THE CONSTRUCTION OF $\mathbf{P}$

When constructing a matrix-iteration by recursion, we will need notation and language for extension. We will use, for an ordinal  $\gamma$ ,  $\mathbf{P}^\gamma$  to indicate that  $\mathbf{P}^\gamma$  is a  $(\kappa, \gamma)$ -matrix iteration.

**Definition 4.1.** (1) *A matrix iteration  $\mathbf{P}^\gamma$  is an extension of  $\mathbf{P}^\delta$  providing  $\delta \leq \gamma$ , and, for each  $\alpha \leq \delta$  and  $i \leq \kappa$ ,  $\mathbb{P}_{i,\alpha}^{\mathbf{P}^\delta} = \mathbb{P}_{i,\alpha}^{\mathbf{P}^\gamma}$ . We can use  $\mathbf{P}^\gamma \upharpoonright \delta$  to denote the unique  $(\kappa, \delta)$ -matrix iteration extended by  $\mathbf{P}^\gamma$ .*

(2) *If, for each  $i < \kappa$ ,  $\dot{Q}_{i,\gamma}$  is a  $\mathbb{P}_{i,\gamma}^{\mathbf{P}}$ -name of a ccc poset satisfying that, for each  $i < j < \kappa$ ,  $\mathbb{P}_{i,\gamma} * \dot{Q}_{i,\gamma}$  is a complete subposet of  $\mathbb{P}_{j,\gamma} * \dot{Q}_{j,\gamma}$ , then we let  $\mathbf{P} * \langle \dot{Q}_{i,\gamma} : i < \kappa \rangle$  denote the  $(\kappa, \gamma + 1)$ -matrix  $\langle \langle \mathbb{P}_{i,\alpha} : i \leq \kappa, \alpha \leq \gamma + 1 \rangle, \langle \dot{Q}_{i,\alpha} : i \leq \kappa, \alpha < \gamma + 1 \rangle \rangle$ , where  $\dot{Q}_{\kappa,\gamma}$  is the  $\mathbf{P}$ -name of the union of  $\{\dot{Q}_{i,\gamma} : i < \kappa\}$  and, for  $i \leq \kappa$ ,  $\mathbb{P}_{i,\gamma} = \mathbb{P}_{i,\gamma}^{\mathbf{P}}$ ,  $\mathbb{P}_{i,\gamma+1} = \mathbb{P}_{i,\gamma}^{\mathbf{P}} * \dot{Q}_{i,\gamma}$ , and for  $\alpha < \gamma$ ,  $(\mathbb{P}_{i,\alpha}, \dot{Q}_{i,\alpha}) = (\mathbb{P}_{i,\alpha}^{\mathbf{P}}, \dot{Q}_{i,\alpha}^{\mathbf{P}})$ .*

The following, from [3, Lemma 3.10], shows that extension at limit steps is canonical.

**Lemma 4.2.** *If  $\gamma$  is a limit and if  $\{\mathbf{P}^\delta : \delta < \gamma\}$  is a sequence of matrix iterations satisfying that for  $\beta < \delta < \gamma$ ,  $\mathbf{P}^\delta \upharpoonright \beta = \mathbf{P}^\beta$ , then there is a unique matrix iteration  $\mathbf{P}^\gamma$  such that  $\mathbf{P}^\gamma \upharpoonright \delta = \mathbf{P}^\delta$  for all  $\delta < \gamma$ .*

*Proof.* For each  $\delta < \gamma$  and  $i < \kappa$ , we define  $\mathbb{P}_{i,\delta}^{\mathbf{P}^\gamma}$  to be  $\mathbb{P}_{i,\delta}^{\mathbf{P}^\delta}$  and  $\dot{\mathbb{Q}}_{i,\delta}^{\mathbf{P}^\gamma}$  to be  $\dot{\mathbb{Q}}_{i,\delta}^{\mathbf{P}^{\delta+1}}$ . It follows that  $\dot{\mathbb{Q}}_{i,\delta}^{\mathbf{P}^\gamma}$  is a  $\mathbb{P}_{i,\delta}^{\mathbf{P}^\gamma}$ -name. Since  $\gamma$  is a limit, the definition of  $\mathbb{P}_{i,\gamma}^{\mathbf{P}^\gamma}$  is required to be  $\bigcup\{\mathbb{P}_{i,\delta}^{\mathbf{P}^\gamma} : \delta < \gamma\}$  for  $i < \kappa$ . Similarly, the definition of  $\mathbb{P}_{\kappa,\gamma}^{\mathbf{P}^\gamma}$  is required to be  $\bigcup\{\mathbb{P}_{i,\gamma}^{\mathbf{P}^\gamma} : i < \kappa\}$ . Let us note that  $\mathbb{P}_{\kappa,\gamma}^{\mathbf{P}^\gamma}$  is also required to be the union of the chain  $\bigcup\{\mathbb{P}_{\kappa,\delta}^{\mathbf{P}^\gamma} : \delta < \gamma\}$ , and this holds by assumption on the sequence  $\{\mathbf{P}^\delta : \delta < \gamma\}$ .

To prove that  $\mathbf{P}^\gamma$  is a  $(\kappa, \gamma)$ -matrix it remains to prove that for  $j < i \leq \kappa$ , and each  $q \in \mathbb{P}_{i,\gamma}^{\mathbf{P}^\gamma}$ , there is a reduction  $p$  in  $\mathbb{P}_{j,\gamma}^{\mathbf{P}^\gamma}$ . Since  $\gamma$  is a limit, there is an  $\alpha < \gamma$  such that  $q \in \mathbb{P}_{i,\alpha}^{\mathbf{P}^\alpha}$  and, by assumption, there is a reduction,  $p$ , of  $q$  in  $\mathbb{P}_{j,\alpha}^{\mathbf{P}^\alpha}$ . By induction on  $\beta$  ( $\alpha \leq \beta \leq \gamma$ ) we note that  $q \in \mathbb{P}_{i,\beta}^{\mathbf{P}^\beta}$  and that  $p$  is a reduction of  $q$  in  $\mathbb{P}_{j,\beta}^{\mathbf{P}^\beta}$ . For limit  $\beta$  it is trivial, and for successor  $\beta$  it follows from condition (1) in the definition of matrix iteration.  $\square$

We also will need the next result taken from [3, Lemma 13], which they describe as well known, for stepping diagonally in the array of posets.

**Lemma 4.3.** *Let  $\mathbb{P}, \mathbb{Q}$  be partial orders such that  $\mathbb{P}$  is a complete suborder of  $\mathbb{Q}$ . Let  $\dot{\mathbb{A}}$  be a  $\mathbb{P}$ -name for a forcing notion and let  $\dot{\mathbb{B}}$  be a  $\mathbb{Q}$ -name for a forcing notion such that  $\Vdash_{\mathbb{Q}} \dot{\mathbb{A}} \subset \dot{\mathbb{B}}$ , and every  $\mathbb{P}$ -name of a maximal antichain of  $\dot{\mathbb{A}}$  is also forced by  $\mathbb{Q}$  to be a maximal antichain of  $\dot{\mathbb{B}}$ . Then  $\mathbb{P} * \dot{\mathbb{A}} < \circ \mathbb{Q} * \dot{\mathbb{B}}$*

Let us also note if  $\dot{\mathbb{B}}$  is equal to  $\dot{\mathbb{A}}$  in Lemma 4.3, then the hypothesis and the conclusion of the Lemma are immediate. On the other hand, if  $\dot{\mathbb{A}}$  is the  $\mathbb{P}$ -name of  $\mathbb{L}(\dot{\mathcal{D}})$  for some  $\mathbb{P}$ -name of a filter  $\dot{\mathcal{D}}$ , then the  $\mathbb{Q}$ -name of  $\mathbb{L}(\dot{\mathcal{D}})$  is not necessarily equal to  $\dot{\mathbb{A}}$ .

**Lemma 4.4** ([6, 1.9]). *Suppose that  $\mathbb{P}, \mathbb{Q}$  are posets with  $\mathbb{P} < \circ \mathbb{Q}$ . Suppose also that  $\dot{\mathcal{D}}_0$  is a  $\mathbb{P}$ -name of a filter on  $\omega$  and  $\dot{\mathcal{D}}_1$  is a  $\mathbb{Q}$ -name of a filter on  $\omega$ . If  $\Vdash_{\mathbb{Q}} \dot{\mathcal{D}}_0 \subseteq \dot{\mathcal{D}}_1$  then  $\mathbb{P} * \mathbb{L}(\dot{\mathcal{D}}_0)$  is a complete subposet of  $\mathbb{Q} * \mathbb{L}(\dot{\mathcal{D}}_1)$  if either of the two equivalent conditions hold:*

- (1)  $\Vdash_{\mathbb{Q}} ((\dot{\mathcal{D}}_0)^+)_{\mathbb{P}} \subseteq \dot{\mathcal{D}}_1^+$ ,
- (2)  $\Vdash_{\mathbb{Q}} \dot{\mathcal{D}}_1 \cap V^{\mathbb{P}} \subseteq \langle \dot{\mathcal{D}}_0 \rangle$  (where  $V^{\mathbb{P}}$  is the class of  $\mathbb{P}$ -names).

*Proof.* Let  $\dot{E}$  be any  $\mathbb{P}$ -name of a maximal antichain of  $\mathbb{L}(\dot{\mathcal{D}}_0)$ . By Lemma 4.3, it suffices to show that  $\mathbb{Q}$  forces that every member of



$\mathbb{L}(\dot{\mathcal{D}}_1)$  is compatible with some member of  $\dot{E}$ . Let  $G$  be any  $\mathbb{Q}$ -generic filter and let  $E$  denote the valuation of  $\dot{E}$  by  $G \cap \mathbb{P}$ . Working in the model  $V[G \cap \mathbb{P}]$ , we have the function  $\rho_E$  as in Lemma 2.4. Choose  $\delta \in \omega_1$  satisfying that  $\rho_E(t) < \delta$  for all  $t \in \omega^{<\omega}$ . Now, working in  $V[G]$ , we consider any  $T \in \mathbb{L}(\dot{\mathcal{D}}_1)$  and we find an element of  $E$  that is compatible with  $T$ . In fact, by induction on  $\alpha < \delta$ , one easily proves that for each  $T \in \mathbb{L}(\dot{\mathcal{D}}_1)$  with  $\rho_E(\text{stem}(T)) \leq \alpha$ ,  $T$  is compatible with some member of  $E$ .  $\square$

**Definition 4.5.** For a  $(\kappa, \gamma)$ -matrix-iteration  $\underline{\mathbf{P}}$ , and ordinal  $i_\gamma < \kappa$ , we say that an increasing sequence  $\langle \dot{\mathcal{D}}_i : i < \kappa \rangle$  is a  $(\underline{\mathbf{P}}, \vec{\lambda}(i_\gamma))$ -thin sequence of filter bases, if for each  $i < j < \kappa$

- (1)  $\dot{\mathcal{D}}_i$  is a subset of  $\mathbb{B}_{i,\gamma}$  (hence a nice  $\mathbb{P}_{i,\gamma}^{\underline{\mathbf{P}}}$ -name)
- (2)  $\Vdash_{\mathbb{P}_{i,\gamma}} \dot{\mathcal{D}}_i$  is a filter with a base of cardinality at most  $\mu_{i_\gamma}$ ,
- (3)  $\Vdash_{\mathbb{P}_{j,\gamma}} \langle \dot{\mathcal{D}}_j \rangle \cap \mathbb{B}_{i,\gamma} \subseteq \langle \dot{\mathcal{D}}_i \rangle$ .

Notice that a  $(\underline{\mathbf{P}}, \vec{\lambda}(i_\gamma))$ -thin sequence of filter bases can be (essentially) eventually constant. Thus we will say that a sequence  $\langle \dot{\mathcal{D}}_i : i \leq j \rangle$  (for some  $j < \kappa$ ) is a  $(\underline{\mathbf{P}}, \vec{\lambda}(i_\gamma))$ -thin sequence of filter bases if the sequence  $\langle \dot{\mathcal{D}}_i : i < \kappa \rangle$  is a  $(\underline{\mathbf{P}}, \vec{\lambda}(i_\gamma))$ -thin sequence of filter bases where  $\dot{\mathcal{D}}_i$  is the  $\mathbb{P}_{i,\gamma}$ -name for  $\mathbb{B}_{i,\gamma} \cap \langle \dot{\mathcal{D}}_j \rangle$  for  $j < i \leq \kappa$ . When  $\underline{\mathbf{P}}$  is clear from the context, we will use  $\vec{\lambda}(i_\gamma)$ -thin as an abbreviation for  $(\underline{\mathbf{P}}, \vec{\lambda}(i_\gamma))$ -thin.

**Corollary 4.6.** For a  $(\kappa, \gamma)$ -matrix-iteration  $\underline{\mathbf{P}}$ , ordinal  $i_\gamma < \kappa$ , and a  $(\underline{\mathbf{P}}, \vec{\lambda}(i_\gamma))$ -thin sequence of filter bases  $\langle \dot{\mathcal{D}}_\xi : i < \kappa \rangle$ ,  $\underline{\mathbf{P}} * \langle \dot{\mathcal{Q}}_{i,\gamma} : i \leq \kappa \rangle$  is a  $\gamma + 1$ -extension of  $\underline{\mathbf{P}}$ , where, for each  $i \leq i_\gamma$ ,  $\dot{\mathcal{Q}}_{i,\gamma}$  is the trivial poset, and for  $i_\gamma \leq i < \kappa$ ,  $\dot{\mathcal{Q}}_{i,\gamma}$  is  $\mathbb{L}(\dot{\mathcal{D}}_i)$ .

**Definition 4.7.** Whenever  $\langle \dot{\mathcal{D}}_i : i < \kappa \rangle$  is a  $(\underline{\mathbf{P}}, \vec{\lambda}(i_\gamma))$ -thin sequence of filter bases, let  $\underline{\mathbf{P}} * \mathbb{L}(\langle \dot{\mathcal{D}}_i : i_\gamma \leq i < \kappa \rangle)$  denote the  $\gamma + 1$ -extension described in Corollary 4.6.

This next corollary is immediate.

**Corollary 4.8.** If  $\underline{\mathbf{P}}$  is a  $\vec{\lambda}$ -thin  $(\kappa, \gamma)$ -matrix and if  $\langle \dot{\mathcal{D}}_i : i < \kappa \rangle$  is a  $(\underline{\mathbf{P}}, \vec{\lambda}(i_\gamma))$ -thin sequence of filter bases, then  $\underline{\mathbf{P}} * \mathbb{L}(\langle \dot{\mathcal{D}}_i : i_\gamma \leq i < \kappa \rangle)$  is a  $\vec{\lambda}$ -thin  $(\kappa, \gamma + 1)$ -matrix.

We now describe a first approximation of the scheme,  $\mathcal{K}(\vec{\lambda})$ , of posets that we will be using to produce the model.

**Definition 4.9.** For an ordinal  $\gamma > 0$  and a  $(\kappa, \gamma)$ -matrix iteration  $\underline{\mathbf{P}}$ , we will say that  $\underline{\mathbf{P}} \in \mathcal{K}(\vec{\lambda})$  providing for each  $0 < \alpha < \gamma$ ,

- (1) for each  $i \leq \kappa$ ,  $\mathbb{P}_{i,1}^{\mathbf{P}}$  is  $\text{Fn}(\lambda_i, 2)$ , and  
(2) there is an  $i_\alpha = i_\alpha^{\mathbf{P}} < \kappa$  and a  $(\mathbf{P} \upharpoonright \alpha, \vec{\lambda}(i_\alpha))$ -thin sequence  $\langle \dot{\mathcal{D}}_i^\alpha : i < \kappa \rangle$  of filter bases, such that  $\mathbf{P} \upharpoonright \alpha + 1$  is equal to  $\mathbf{P} \upharpoonright \alpha * \mathbb{L}(\langle \dot{\mathcal{D}}_i^\alpha : i_\alpha \leq i < \kappa \rangle)$ .

For each  $0 < \alpha < \gamma$ , we let  $\dot{\mathcal{D}}_\kappa^\alpha$  denote the  $\mathbf{P} \upharpoonright \alpha$ -name of the union  $\bigcup \{ \dot{\mathcal{D}}_i^\alpha : i_\alpha \leq i < \kappa \}$ , and we let  $\dot{L}_\alpha$  denote the canonical  $\mathbf{P} \upharpoonright \alpha + 1$ -name of the subset of  $\omega$  added by  $\mathbb{L}(\dot{\mathcal{D}}_\kappa^\alpha)$ .

Let us note that each  $\mathbf{P} \in \mathcal{K}(\vec{\lambda})$  is  $\vec{\lambda}$ -thin. Furthermore, by Lemma 3.5, this means that each  $\mathbf{P} \in \mathcal{K}(\vec{\lambda})$  forces that  $\mathfrak{s} \leq \lambda$ . We begin a new section for the task of proving that there is a  $\mathbf{P} \in \mathcal{K}(\vec{\lambda})$  that forces that  $\mathfrak{s} \geq \lambda$ .

It will be important to be able to construct  $(\mathbf{P}, \vec{\lambda}(i_\gamma))$ -thin sequences of filter bases, and it seems we will need some help.

**Definition 4.10.** For an ordinal  $\gamma > 0$  and a  $(\kappa, \gamma)$ -matrix iteration  $\mathbf{P}$  we will say that  $\mathbf{P} \in \mathcal{H}(\vec{\lambda})$  if  $\mathbf{P}$  is in  $\mathcal{K}(\vec{\lambda})$  and for each  $0 < \alpha < \gamma$ , if  $i_\alpha = i_\alpha^{\mathbf{P}} > 0$  then  $\omega_1 \leq \text{cf}(\alpha) \leq \mu_{i_\alpha}$  and there is a  $\beta_\alpha < \alpha$  such that

- (1) for  $\beta_\alpha \leq \xi < \alpha$ ,  $i_\xi \in \{0, i_\alpha\}$ ,  
(2) if  $\beta_\alpha \leq \eta < \alpha$ ,  $i_\eta > 0$  and  $\xi = \eta + \omega_1 \leq \alpha$ , then  $\dot{L}_\eta \in \dot{\mathcal{D}}_{i_\xi}^\xi$ , and  $\mathbb{P}_{i_\xi, \xi} \Vdash \dot{\mathcal{D}}_{i_\xi}^\alpha$  has a descending mod finite base of cardinality  $\omega_1$ ,  
(3) if  $\beta_\alpha < \xi \leq \alpha$ ,  $i_\xi > 0$ , and  $\eta + \omega_1 < \xi$  for  $\eta < \xi$ , then  $\{ \dot{L}_\eta : \beta_\alpha \leq \eta < \alpha, \text{cf}(\eta) \geq \omega_1 \}$  is a base for  $\dot{\mathcal{D}}_{i_\xi}^\xi$ .

## 5. PRODUCING $\vec{\lambda}$ -THIN FILTER SEQUENCES

In this section we prove this main lemma.

**Lemma 5.1.** Suppose that  $\mathbf{P}^\gamma \in \mathcal{H}(\vec{\lambda})$  and that  $\mathcal{Y}$  is a set of fewer than  $\lambda$  nice  $\mathbf{P}^\gamma$ -names of subsets of  $\omega$ , then there is a  $\delta < \gamma + \lambda$  and an extension  $\mathbf{P}^\delta$  of  $\mathbf{P}^\gamma$  in  $\mathcal{H}(\vec{\lambda})$  that forces that the family  $\mathcal{Y}$  is not a splitting family.

The main theorem follows easily.

*Proof of Theorem 1.2.* Let  $\theta$  be any regular cardinal so that  $\theta^{<\lambda} = \theta$  (for example,  $\theta = (2^\lambda)^+$ ). Construct  $\mathbf{P}^\theta \in \mathcal{H}(\vec{\lambda})$  so that for all  $\mathcal{Y} \subset \mathbb{B}_{\kappa, \theta}$  with  $|\mathcal{Y}| < \lambda$ , there is a  $\gamma < \delta < \theta$  so that  $\mathcal{Y} \subset \mathbb{B}_{\kappa, \gamma}$  and, by applying Lemma 5.1, such that  $\mathbf{P}^\theta \upharpoonright \delta$  forces that  $\mathcal{Y}$  is not a splitting family.  $\square$

We begin by reducing our job to simply finding a  $(\mathbf{P}, \vec{\lambda}(i_\gamma))$ -thin sequence.

**Definition 5.2.** For a  $(\kappa, \gamma)$ -matrix-iteration  $\underline{\mathbf{P}}^\gamma$ , we say that a subset  $\mathcal{E}$  of  $\mathbb{B}_{\kappa, \gamma}$  is  $(\underline{\mathbf{P}}^\gamma, \vec{\lambda}(i_\gamma))$ -thin filter subbase if,  $i_\gamma < \kappa$ ,  $|\mathcal{E}| \leq \mu_{i_\gamma}$ , and the sequence  $\langle \langle \mathcal{E} \cap \mathbb{B}_{i_\gamma} \rangle : i < \kappa \rangle$  is a  $(\underline{\mathbf{P}}^\gamma, \vec{\lambda}(i_\gamma))$ -thin sequence of filter bases.

**Lemma 5.3.** For any  $\underline{\mathbf{P}}^\gamma \in \mathcal{H}(\vec{\lambda})$ , and any  $(\underline{\mathbf{P}}^\gamma, \vec{\lambda}(i_\gamma))$ -thin filter base  $\mathcal{E}$ , there is an  $\alpha \leq \gamma + \mu_{i_\gamma} + 1$  and extensions  $\underline{\mathbf{P}}^\alpha, \underline{\mathbf{P}}^{\alpha+1}$  of  $\underline{\mathbf{P}}^\gamma$  in  $\mathcal{H}(\vec{\lambda})$ , such that,  $\underline{\mathbf{P}}^{\alpha+1} = \underline{\mathbf{P}}^\alpha * \mathbb{L}(\langle \dot{\mathcal{D}}_i^\alpha : i_\alpha \leq i < \kappa \rangle)$  and  $\underline{\mathbf{P}}^\alpha$  forces that  $\mathcal{E} \cap \mathbb{B}_{i_\gamma}$  is a subset of  $\dot{\mathcal{D}}_i^\alpha$  for all  $i < \kappa$ .

*Proof.* The case  $i_\gamma = 0$  is trivial, so we assume  $i_\gamma > 0$ . There is no loss of generality to assume that  $\mathcal{E} \cap \mathbb{B}_{i_\gamma, \gamma}$  has character  $\mu_{i_\gamma}$ . Let  $\{\dot{E}_\xi : \xi < \mu_{i_\gamma}\} \subset \mathcal{E} \cap \mathbb{B}_{i_\gamma, \gamma}$  enumerate a filter base for  $\langle \mathcal{E} \rangle \cap \mathbb{B}_{i_\gamma, \gamma}$ . We can assume that this enumeration satisfies that  $\dot{E}_\xi \setminus \dot{E}_{\xi+1}$  is forced to be infinite for all  $\xi < \mu_{i_\gamma}$ . Let  $\mathcal{A}$  be any countably generated free filter on  $\omega$  that is not principal mod finite. By induction on  $\xi < \mu_{i_\gamma}$  we define  $\underline{\mathbf{P}}^{\gamma+\xi}$  by simply defining  $i_{\gamma+\xi}$  and the sequence  $\langle \dot{\mathcal{D}}_i^{\gamma+\xi} : i_{\gamma+\xi} \leq i \leq \kappa \rangle$ . We will also recursively define, for each  $\xi < \mu_{i_\gamma}$ , a  $\underline{\mathbf{P}}^{\gamma+\xi}$ -name  $\dot{D}_\xi$  such that  $\underline{\mathbf{P}}^{\gamma+\xi}$  forces that  $\dot{D}_\xi \subset \dot{E}_\xi$ . An important induction hypothesis is that  $\{\dot{D}_\eta : \eta < \xi\} \cup \{\dot{E}_\zeta : \zeta < \mu_{i_\gamma}\} \cup \mathcal{E}$  is forced to have the finite intersection property.

For each  $\xi < \gamma + \omega_1$ , let  $i_\xi = 0$  and  $\dot{\mathcal{D}}_i^\xi$  be the  $\underline{\mathbf{P}}^\xi$ -name  $\langle \mathcal{A} \rangle \cap \mathbb{B}_{i, \xi}$  for all  $i \leq \kappa$ . The definition of  $\dot{D}_0$  is simply  $\dot{E}_0$ . By recursion, for each  $\eta < \omega_1$  and  $\xi = \eta + 1$ , we define  $\dot{D}_\xi$  to be the intersection of  $\dot{D}_\eta$  and  $\dot{E}_\xi$ . For limit  $\xi < \omega_1$ , we note that  $\mathbb{P}_{i_\gamma, \xi}$  forces that  $\mathbb{L}(\langle \mathcal{A} \rangle)$  is isomorphic to  $\mathbb{L}(\langle \{\dot{D}_\eta \cap \dot{E}_\xi : \eta < \xi\} \rangle)$ . Therefore, we can let  $\dot{D}_\xi$  be a  $\underline{\mathbf{P}}^{\xi+1}$ -name for the generic real added by  $\mathbb{L}(\langle \{\dot{D}_\eta \cap \dot{E}_\xi : \eta < \xi\} \rangle)$ . A routine density argument shows that this definition satisfies the induction hypothesis.

The definition of  $i_{\gamma+\omega_1}$  is  $i_\gamma$  and the definition of  $\dot{\mathcal{D}}_{i_\gamma}^{\gamma+\omega_1}$  is the filter generated by  $\{\dot{D}_\xi : \xi < \omega_1\}$ . The definition of  $\dot{D}_{\omega_1}$  is  $\dot{L}_{\gamma+\omega_1}$ .

Let  $S$  denote the set of  $\eta < \mu_{i_\gamma}$  with uncountable cofinality. We now add additional induction hypotheses:

- (1) if  $\zeta = \sup(S \cap \xi) < \xi$  and  $\xi = \nu + 1$ , then  $\dot{D}_\xi = \dot{D}_\nu \cap \dot{E}_\xi$ , and  $i_\xi = 0$  and  $\dot{\mathcal{D}}_i^{\gamma+\xi} = \langle \mathcal{A} \rangle$  for all  $i \leq \kappa$
- (2) if  $\zeta = \sup(S \cap \xi) < \xi$  and  $\xi$  is a limit of countable cofinality, then  $i_\xi = 0$  and  $\dot{\mathcal{D}}_i^{\gamma+\xi} = \langle \mathcal{A} \rangle$  for all  $i \leq \kappa$ , and  $\dot{D}_\xi$  is forced by  $\underline{\mathbf{P}}^{\gamma+\xi+1}$  to be the generic real added by  $\mathbb{L}(\langle \{\dot{D}_\eta \cap \dot{E}_\xi : \zeta \leq \eta < \xi\} \rangle)$ ,
- (3) if  $\zeta = \sup(S \cap \xi)$  and  $\xi = \zeta + \omega_1$ , then  $i_\xi = i_\gamma$ ,  $\dot{\mathcal{D}}_{i_\xi}^{\gamma+\xi}$  is the filter generated by  $\{\dot{E}_\zeta \cap \dot{D}_\eta : \zeta \leq \eta < \xi\}$  and  $\dot{D}_\xi$  is  $\dot{L}_{\gamma+\xi}$ ,

- (4) if  $S \cap \xi$  is cofinal in  $\xi$  and  $\text{cf}(\xi) > \omega$ , then  $i_\xi = i_\gamma$  and  $\dot{\mathcal{D}}_{i_\xi}^{\gamma+\xi}$  is the filter generated by  $\{\dot{D}_{\gamma+\eta} : \eta \in S \cap \xi\}$  and  $\dot{D}_\xi = \dot{L}_{\gamma+\xi}$ ,
- (5) if  $S \cap \xi$  is cofinal in  $\xi$  and  $\text{cf}(\xi) = \omega$ , then  $i_\xi = 0$  and  $\dot{\mathcal{D}}_{i_\xi}^{\gamma+\xi} = \langle \mathcal{A} \rangle$  for all  $i \leq \kappa$ , and  $\dot{D}_\xi$  is forced by  $\mathbf{P}^{\gamma+\xi+1}$  to be the generic real added by  $\mathbb{L}(\{\dot{D}_{\eta_n} \cap \dot{E}_\xi : n \in \omega\})$ , where  $\{\eta_n : n \in \omega\}$  is some increasing cofinal subset of  $S \cap (\gamma, \xi)$ .

It should be clear that the induction continues to stage  $\mu_{i_\gamma}$  and that  $\mathbf{P}^{\gamma+\xi} \in \mathcal{H}(\vec{\lambda}(i_\gamma))$  for all  $\xi \leq \mu_{i_\gamma}$ , with  $\beta_{\gamma_\xi} = \gamma$  being the witness to Definition 4.10 for all  $\xi$  with  $\text{cf}(\xi) > \omega$ .

The final definition of the sequence  $\langle \dot{\mathcal{D}}_i^\delta : i_\delta = i_\gamma \leq i \leq \kappa \rangle$ , where  $\delta = \gamma + \mu_{i_\gamma}$  is that  $\dot{\mathcal{D}}_{i_\gamma}^\delta$  is the filter generated by  $\{\dot{L}_{\gamma+\xi} : \text{cf}(\xi) > \omega\}$ , and for  $i_\gamma < i \leq \kappa$ ,  $\dot{\mathcal{D}}_i^\delta$  is the filter generated by  $\dot{\mathcal{D}}_{i_\gamma}^\delta \cup (\mathcal{E} \cap \mathbb{B}_{i,\gamma})$ .  $\square$

**Lemma 5.4.** *Suppose that  $\mathcal{E}$  is a  $(\mathbf{P}^\gamma, \vec{\lambda}(i_\gamma))$ -thin filter base. Also assume that  $i < \kappa$  and  $\alpha \leq \gamma$  and  $\mathcal{E}_1 \subset \mathbb{B}_{i,\alpha}$  is a  $(\mathbf{P}^\alpha, \vec{\lambda}(i_\gamma))$ -thin filter base satisfying that  $\langle \mathcal{E} \rangle \cap \mathbb{B}_{i,\alpha} \subset \langle \mathcal{E}_1 \rangle$ , then  $\mathcal{E} \cup \mathcal{E}_1$  is a  $(\mathbf{P}^\gamma, \vec{\lambda}(i_\gamma))$ -thin filter subbase.*

*Proof.* Let  $\mathcal{E}_2$  be equal to  $\mathcal{E} \cup \mathcal{E}_1$ . The fact that each member of the sequence  $\langle \dot{\mathcal{D}}_j = \langle \mathcal{E}_2 \cap \mathbb{B}_{j,\gamma} \rangle : j < \kappa \rangle$  is a name of a filter base with character at most  $\mu_{i_\gamma}$  is immediate. Now we verify that if  $j_1 < j_2 < \kappa$ , then  $\Vdash_{\mathbb{P}_{j_2,\gamma}} \dot{\mathcal{D}}_{j_2} \cap \mathbb{B}_{j_1,\gamma} \subset \dot{\mathcal{D}}_{j_1}$ . Let  $\dot{b} \in \mathbb{B}_{j_2,\gamma}$  and suppose there are  $p \in \mathbb{P}_{j_2,\gamma}$ ,  $\dot{E}_0 \in \mathcal{E} \cap \mathbb{B}_{j_2,\gamma}$ , and  $\dot{E}_1 \in \mathcal{E}_1$  such that  $p \Vdash \dot{b} \cap \dot{E}_0 \cap \dot{E}_1$ . It suffices to produce an  $\dot{E} \in \langle \mathcal{E}_2 \rangle \cap \mathbb{B}_{j_1,\gamma}$  satisfying that  $p \Vdash \dot{b} \cap \dot{E} = \emptyset$ . First, using that  $\mathcal{E}$  is  $(\mathbf{P}^\gamma, \vec{\lambda}(i_\gamma))$ -thin, choose  $\dot{E}_2 \in \langle \mathcal{E} \rangle \cap \mathbb{B}_{j_1,\gamma}$  such that  $p \Vdash (\dot{b} \setminus \dot{E}_0) \cap \dot{E}_2 = \emptyset$ . Equivalently, we have that  $p \Vdash (\dot{b} \cap \dot{E}_2) \subset \dot{E}_0$ , and therefore  $p \Vdash (\dot{b} \cap \dot{E}_2) \cap \dot{E}_1 = \emptyset$ . Since  $\dot{E}_1$  is a  $\mathbb{P}_{j_2,\alpha}$ -name, there is a  $\mathbb{P}_{j_1,\alpha}$ -name (which we can denote as)  $(\dot{b} \cap \dot{E}_2) \upharpoonright \alpha$  satisfying that  $p \Vdash \dot{E}_2 \cap (\dot{b} \cap \dot{E}_2) \upharpoonright \alpha$  is empty and that  $p \Vdash (\dot{b} \cap \dot{E}_2) \subset (\dot{b} \cap \dot{E}_2) \upharpoonright \alpha$ . Now using that  $\mathcal{E}_1$  is  $(\mathbf{P}^\alpha, \vec{\lambda}(i_\gamma))$ -thin, choose  $\dot{E}_3 \in \langle \mathcal{E}_1 \rangle \cap \mathbb{B}_{j_1,\alpha}$  so that  $p \Vdash \dot{E}_3 \cap (\dot{b} \cap \dot{E}_2) \upharpoonright \alpha$  is empty. Naturally we have that  $p \Vdash \dot{E}_3 \cap (\dot{b} \cap \dot{E}_2)$  is also empty. This completes the proof since  $\dot{E}_2 \cap \dot{E}_3$  is in  $\langle \mathcal{E}_2 \rangle \cap \mathbb{B}_{j_1,\gamma}$ .  $\square$

Let  $\mathbf{P}^\gamma \in \mathcal{H}(\vec{\lambda})$  and let  $\dot{y} \in \mathbb{B}_{\kappa,\gamma}$ . For a family  $\mathcal{E} \subset \mathbb{B}_{\kappa,\gamma}$  and condition  $p \in \mathbf{P}^\gamma$  say that  $p$  forces that  $\mathcal{E}$  measures  $\dot{y}$  if  $p \Vdash_{\mathbf{P}^\gamma} \{\dot{y}, \omega \setminus \dot{y}\} \cap \langle \mathcal{E} \rangle \neq \emptyset$ . Naturally we will just say that  $\mathcal{E}$  measures  $\dot{y}$  if 1 forces that  $\mathcal{E}$  measures  $\dot{y}$ .

Given Lemma 5.3, it will now suffice to prove.

**Lemma 5.5.** *If  $\mathcal{Y} \subset \mathbb{B}_{\kappa,\gamma}$  for some  $\mathbf{P}^\gamma \in \mathcal{H}(\vec{\lambda})$  and  $|\mathcal{Y}| \leq \mu_{i_\gamma}$  for some  $i_\gamma < \kappa$ , then there is a  $(\mathbf{P}^\gamma, \vec{\lambda}(i_\gamma))$ -thin filter  $\mathcal{E} \subset \mathbb{B}_{\kappa,\gamma}$  that measures every element of  $\mathcal{Y}$ .*

In fact, to prove Lemma 5.5, it is evidently sufficient to prove:

**Lemma 5.6.** *If  $\mathbf{P}^\gamma \in \mathcal{H}(\vec{\lambda})$ ,  $\dot{y} \in \mathbb{B}_{\kappa,\gamma}$ , and if  $\mathcal{E}$  is a  $(\mathbf{P}^\gamma, \vec{\lambda}(i_\gamma))$ -thin filter, then there is a family  $\mathcal{E}_1 \supset \mathcal{E}$  measuring  $\dot{y}$  that is also a  $(\mathbf{P}^\gamma, \vec{\lambda}(i_\gamma))$ -thin filter.*

*Proof.* Throughout the proof we suppress mention of  $\mathbf{P}^\gamma$  and refer instead to component member posets  $\mathbb{P}_{i,\alpha}, \mathbb{Q}_{i,\alpha}$  of  $\mathbf{P}^\gamma$ . Let  $i_{\dot{y}}$  be minimal such that  $\dot{y}$  is in  $\mathbb{B}_{i_{\dot{y}},\gamma}$ . Proceeding by induction, we can assume that the lemma holds for all  $\dot{x} \in \mathbb{B}_{j,\gamma}$  and all  $j < i_{\dot{y}}$ .

We can replace  $\dot{y}$  by any  $\dot{x} \in \mathbb{B}_{i_{\dot{y}},\gamma}$  that has the property that  $1 \Vdash \dot{x} \in \{\dot{y}, \omega \setminus \dot{y}\}$  since if we measure  $\dot{x}$  then we also measure  $\dot{y}$ . With this reduction then we can assume that no condition forces that  $\omega \setminus \dot{y}$  is in the filter generated by  $\mathcal{E}$ .

*Fact 1.* If  $i_{\dot{y}} \leq i_\gamma$ , then there is a  $\dot{E} \in \mathbb{B}_{i_{\dot{y}},\gamma}$  such that  $\mathcal{E} \cup \{\dot{E}\}$  is contained a  $(\mathbf{P}^\gamma, \vec{\lambda}(i_\gamma))$ -thin filter that measures  $\dot{y}$ .

*Proof of Fact 1.* It is immediate that  $\langle \{\dot{y}\} \cup (\mathbb{B}_{i_{\dot{y}},\gamma} \cap \mathcal{E}) \rangle$  is a  $(\mathbf{P}^\gamma, \vec{\lambda}(i_\gamma))$ -thin filter. Therefore, by Lemma 5.4,  $\mathcal{E} \cup \{\dot{y}\}$  is a  $(\mathbf{P}^\gamma, \vec{\lambda}(i_\gamma))$ -thin filter subbase.  $\square$

We may thus assume that  $0 < i_{\dot{y}}$  and that the Lemma has been proven for all members of  $\mathbb{B}_{i,\gamma}$  for all  $i < i_{\dot{y}}$ . Similarly, let  $\alpha_{\dot{y}}$  be minimal so that  $\dot{y} \in \mathbb{B}_{i_{\dot{y}},\alpha_{\dot{y}}}$ , and assume that the Lemma has been proven for all members of  $\mathbb{B}_{i_{\dot{y}},\beta}$  for all  $\beta < \alpha_{\dot{y}}$ . We skip proving the easy case when  $\alpha_{\dot{y}} = 1$  and henceforth assume that  $1 < \alpha_{\dot{y}}$ . Notice also that  $\alpha_{\dot{y}}$  has countable cofinality since  $\mathbb{P}_{i_{\dot{y}},\gamma}$  is ccc.

Now choose an elementary submodel  $M$  of  $H((2^{\lambda_\gamma})^+)$  containing  $\vec{\lambda}, \mathbf{P}^\gamma, \mathcal{E}, \dot{y}$  and so that  $M$  has cardinality equal to  $\mu_{i_\gamma}$  and, by our cardinal assumptions,  $M^{\lambda_j} \subset M$  for all  $j < i_\gamma$ . Naturally this implies that  $M^\omega \subset M$ .

By the inductive assumption we may assume that there is an  $\mathcal{E}_1 \supset \mathcal{E}$  that is  $(\mathbf{P}^\gamma, \vec{\lambda}(i_\gamma))$ -thin and measures every element of  $M \cap \mathbb{B}_{j,\gamma}$  for  $j < i_{\dot{y}}$  as well as every element of  $M \cap \mathbb{B}_{i_{\dot{y}},\beta}$  for all  $\beta \in M \cap \alpha_{\dot{y}}$ . Moreover, it is easily checked that we can assume that  $\mathcal{E}_1$  is a subset of  $M$ . Furthermore, we may assume that  $\mathcal{E}_1$  contains a maximal family of subsets of  $M \cap \mathbb{B}_{i_{\dot{y}},\alpha_{\dot{y}}}$  that forms a  $(\mathbf{P}^\gamma, \vec{\lambda}(i_\gamma))$ -thin filter subbase.

*Fact 2.* There is a maximal antichain  $A \subset \mathbb{P}_{i_{\dot{y}},\gamma}$  and a subset  $A_1 \subset A$  such that

- (1) each  $p \in A_1$  forces that  $\mathcal{E}_1$  measures  $\dot{y}$ ,
- (2) for each  $p \in A \setminus A_1$ ,  $p$  forces that there is an  $i_p < i_{\dot{y}}$  such that  $\mathbb{B}_{i_p, \gamma} \cap \langle \mathcal{E}_1 \cup \{\dot{y}\} \rangle$  is not generated by the elements in  $M$ ,
- (3) for each  $p \in A \setminus A_1$ ,  $p$  forces that there is a  $j_p < i_{\dot{y}}$  such that  $i_p \leq j_p$  and  $\mathbb{B}_{j_p, \gamma} \cap \langle \mathcal{E}_1 \cup \{\omega \setminus \dot{y}\} \rangle$  is not generated by the elements in  $M$ .

*Proof of Fact 2.* Suppose that  $p \in \mathbb{P}_{i_{\dot{y}}, \gamma}$  forces that the conclusion (2) fails. We have already arranged that  $p \Vdash_{\mathbb{P}_{i_{\dot{y}}, \gamma}} \dot{y} \in \langle \mathcal{E}_1 \cap \mathbb{B}_{i_{\dot{y}}, \gamma} \rangle^+$ . Define  $\dot{E} \in \mathbb{B}_{i_{\dot{y}}, \gamma}$  so that  $p$  forces  $\dot{E} = \dot{y}$  and each  $q \in \mathbb{P}_{i_{\dot{y}}, \gamma} \cap p^\perp$  forces that  $\dot{E} = \omega$ . It is easily checked that  $\mathbb{B}_{i_{\dot{y}}, \gamma} \cap \langle \mathcal{E}_1 \cup \{\dot{E}\} \rangle$  is then  $(\mathbf{P}^\gamma, \vec{\lambda}(i_\gamma))$ -thin and that  $p$  forces that it measures  $\dot{y}$ . This condition ensures that  $p$  is compatible with an element of  $A_1$ .

If (2) holds but (3) fails, then by a symmetric argument as in the previous paragraph we can again define  $\dot{E}$  so that  $\mathbb{B}_{i_{\dot{y}}, \gamma} \cap \langle \mathcal{E}_1 \cup \{\dot{E}\} \rangle$  is then  $(\mathbf{P}^\gamma, \vec{\lambda}(i_\gamma))$ -thin and that  $p$  forces that it measures  $\omega \setminus \dot{y}$ .  $\square$

If by increasing  $M$  we can enlarge  $A_1$  we simply do so. Since  $\mathbf{P}^\gamma$  is ccc we may assume that this is no longer possible, and therefore we may also assume that  $A$  is a subset of  $M$ . Now we choose any  $p \in A \setminus A_1$ . It suffices to produce an  $\dot{E}_p \in \mathbb{B}_{i_{\dot{y}}, \gamma}$  that can be added to  $\mathcal{E}_1$  that measures  $\dot{y}$  and satisfies that  $q \Vdash \dot{E}_p = \omega$  for all  $q \in p^\perp$ . This is because we then have that  $\mathcal{E}_1 \cup \{\dot{E}_p : p \in A \setminus A_1\}$  is contained in a  $\vec{\lambda}(i_\gamma)$ -thin filter that measures  $\dot{y}$ .

*Fact 3.* There is an  $\alpha$  such that  $\alpha_{\dot{y}} = \alpha + 1$ .

*Proof of Fact 3.* Otherwise, let  $j = i_p$  and for each  $r < p$  in  $\mathbb{P}_{i_{\dot{y}}, \alpha_{\dot{y}}}$ , choose  $\beta \in M \cap \alpha_{\dot{y}}$  such that  $r \in \mathbb{P}_{i_{\dot{y}}, \beta}$ , and define a name  $\dot{y}[r]$  in  $M \cap \mathbb{B}_{j, \gamma}$  according to  $(\ell, q) \in \dot{y}[r]$  providing there is a pair  $(\ell, p_\ell) \in \dot{y}$  such that  $q <_j p_\ell$  and  $q \upharpoonright \beta$  is in the set  $M \cap \mathbb{P}_{j, \beta} \setminus (r \wedge p_\ell \upharpoonright \beta)^\perp$ . This set, namely  $\dot{y}[r]$ , is in  $M$  because  $\mathbb{P}_{j, \beta}$  is ccc and  $M^\omega \subset M$ .

We prove that  $r$  forces that  $\dot{y}[r]$  contains  $\dot{y}$ . Suppose that  $r_1 < r$  and there is a pair  $(\ell, p_\ell) \in \dot{y}$  with  $r_1 < p_\ell$ . Choose an  $r_2 \in \mathbb{P}_{j, \gamma}$  so that  $r_2 <_j r_1$ . It suffices to show  $r_2 \Vdash \ell \in \dot{y}[r]$ . Let  $q <_j p_\ell$  with  $q \in M$ . Then  $r_2 \not\leq_j p_\ell$  implies  $r_2 \not\leq_j q$ . Since  $r_2$  was any  $<_j$ -projection of  $r_1$  we can assume that  $r_2 < q$ . Since  $r_2 \upharpoonright \beta$  is in  $(\mathbb{P}_{j, \beta} \cap (r \wedge p_\ell \upharpoonright \beta)^\perp)^\perp$ , it follows that  $q \upharpoonright \beta \notin (r \wedge p_\ell \upharpoonright \beta)^\perp$ . This implies that  $(\ell, q) \in \dot{y}[r]$  and completes the proof that  $r_2 \Vdash \ell \in \dot{y}[r]$ .

Now assume that  $\beta < \alpha_{\dot{y}}$  and  $r \Vdash \dot{b} \cap \dot{E} \cap \dot{y}$  is empty for some  $r < p$  in  $\mathbb{P}_{i_{\dot{y}}, \beta}$ ,  $\dot{b} \in \mathbb{B}_{j, \gamma}$ , and  $\dot{E} \in \mathcal{E}_1 \cap \mathbb{B}_{i_{\dot{y}}, \gamma}$ . Let  $\dot{x} = (\dot{E} \cap \dot{y})[r]$  (defined as above for  $\dot{y}[r]$ ). We complete the proof of Fact 3 by proving that  $r \Vdash \dot{b} \cap \dot{x}$  is

empty. Since each are in  $\mathbb{B}_{j,\gamma}$ , we may choose any  $r_1 <_j r$ , and assume that  $r_1 \Vdash \ell \in \dot{b} \cap \dot{x}$ . In addition we can suppose that there is a pair  $(\ell, q) \in \dot{x}$  such that  $r_1 < q$ . The fact that  $(\ell, q) \in \dot{x}$  means there is a  $p_\ell$  with  $(\ell, p_\ell)$  in the name  $\dot{E} \cap \dot{y}$  such that  $q <_j p_\ell$ . Since  $r_1 \in \mathbb{P}_{j,\gamma}$  and  $r_1 < q$ , it follows that  $r_1 \not\leq p_\ell$ . Now it follows that  $r_1$  has an extension forcing that  $\ell \in \dot{b} \cap (\dot{E} \cap \dot{y})$  which is a contradiction.  $\square$

*Fact 4.*  $i_{\dot{y}} = i_\alpha$  and so also  $i_p < i_\alpha$ .

*Proof of Fact 4.* Since  $\mathbb{P}_{i,\alpha+1} = \mathbb{P}_{i,\alpha}$  for  $i < i_\alpha$ , we have that  $i_\alpha \leq i_{\dot{y}}$ . Now assume that  $i_\alpha < i_{\dot{y}}$  and we proceed much as we did in Fact 3 to prove that  $i_p$  does not exist. Assume that  $r < p$  (in  $\mathbb{P}_{i_{\dot{y}},\alpha+1}$ ) and  $r \Vdash \dot{b} \cap (\dot{E} \cap \dot{y})$  is empty for some  $\dot{E} \in M \cap \langle \mathcal{E}_1 \rangle \cap \mathbb{B}_{i_{\dot{y}},\gamma}$  and  $\dot{b} \in \mathbb{B}_{i_p,\gamma}$ . It follows from Lemma 5.4 that we can simply assume that  $\dot{E} \in \mathcal{E}_1 \cap \mathbb{B}_{i_{\dot{y}},\alpha+1}$ , and similarly that  $\dot{b} \in \mathbb{B}_{i_p,\alpha+1}$ .

Let  $\dot{T}_\alpha$  be the  $\mathbb{P}_{i_{\dot{y}},\alpha}$ -name such that  $r \restriction \alpha \Vdash r(\alpha) = \dot{T}_\alpha \in \mathbb{L}(\mathcal{D}_{i_{\dot{y}}}^\alpha)$ . We may assume that there is a  $t_\alpha \in \omega^{<\omega}$  such that  $r \restriction \alpha \Vdash t_\alpha = \text{stem}(\dot{T}_\alpha)$ .

Choose any  $M \cap \mathbb{P}_{i_\alpha,\alpha}$ -generic filter  $\bar{G}$  such that  $r \restriction \alpha \in \bar{G}^+$ . Since  $\mathbb{P}_{i_\alpha,\alpha}$  is ccc and  $M^\omega \subset M$ , it follows that  $M[\bar{G}]$  is closed under  $\omega$ -sequences in the model  $V[\bar{G}]$ .

In this model, define an  $\mathbb{L}(\mathcal{D}_{i_\alpha}^\alpha)$ -name  $\dot{x}$ . A pair  $(\ell, T_\ell) \in \dot{x}$  if  $t_\alpha \leq \text{stem}(T_\ell) \in T_\ell \in \mathbb{L}(\mathcal{D}_{i_\alpha}^\alpha)$  and for each  $\text{stem}(T_\ell) \leq t \in T_\ell$ , there is a pair  $(\ell, q_{\ell,t}) \in M$  in the name  $(\dot{y} \cap \dot{E})$  such that  $q_{\ell,t} \restriction \alpha \in \bar{G}^+$ ,  $q_{\ell,t} \restriction \alpha \Vdash t = \text{stem}(q_{\ell,t}(\alpha))$ , and  $(q_{\ell,t} \restriction \alpha \wedge r \restriction \alpha)$  does not force (over the poset  $\bar{G}^+$ ) that  $t \notin \dot{T}_\alpha$ . We will show that  $r$  forces over the poset  $\bar{G}^+$  that  $\dot{x}$  contains  $\dot{E} \cap \dot{y}$  and that  $\dot{x} \cap \dot{b}$  is empty. This proves that  $p$  forces that  $\langle \mathcal{E}_1 \rangle \cap \mathbb{B}_{i_p,\alpha+1}$  generates  $\langle \mathcal{E}_1 \cup \{\dot{y}\} \rangle \cap \mathbb{B}_{i_p,\alpha+1}$  since  $\dot{x}$  must be forced to be in  $\langle \mathcal{E}_1 \rangle$ . It then follows from Lemma 5.4 that  $\mathcal{E}_1 \cap \mathbb{B}_{i_p,\gamma}$  generates  $\langle \mathcal{E}_1 \cup \{\dot{y}\} \rangle \cap \mathbb{B}_{i_p,\gamma}$ , contradicting the assumption on  $i_p$ .

To prove that  $r$  forces that  $\dot{x}$  contains  $\dot{y} \cap \dot{E}$ , we consider any  $r_\ell < r$  that forces over  $\bar{G}^+$  that  $\ell \in \dot{y} \cap \dot{E}$ . We may choose  $(\ell, p_\ell) \in M$  in the name  $(\dot{E} \cap \dot{y})$  such that (wlog)  $r_\ell < p_\ell$ . We may assume that  $r_\ell \restriction \alpha$  forces a value  $t$  on  $\text{stem}(r_\ell(\alpha))$  and that this equals  $\text{stem}(p_\ell(\alpha))$ . Now show there is a  $T_\ell \in \mathbb{L}(\mathcal{D}_{i_\alpha}^\alpha)$ . In fact, assume  $t \in T_\ell$  with  $q_{\ell,t}$  as the witness. Let  $L^- = \{k : t \frown k \notin T_\ell\}$ ; it suffices to show that  $L^- \notin (\mathcal{D}_{i_\alpha}^\alpha)^+$ .

By assumption that  $q_{\ell,t}$  is the witness, there is an  $r_t < (q_{\ell,t} \restriction \alpha \wedge r \restriction \alpha)$  such that  $r_t \Vdash t \in \dot{T}_\alpha$  and  $r_t \Vdash t = \text{stem}(q_{\ell,t}(\alpha))$ . By strengthening  $r_t$  we can assume that  $r_t$  forces a value  $\dot{D} \in \mathcal{D}_{i_{\dot{y}}}^\alpha$  on  $\{k : t \frown k \in \dot{T}_\alpha \cap q_{\ell,t}(\alpha)\}$ . But now, it follows that  $r_t$  forces that  $\dot{D}$  is disjoint from  $L^-$  since if  $r_{t,k} \Vdash k \in \dot{D}$  for some  $r_{t,k} < r_t$ ,  $r_{t,k}$  is the witness to  $(\ell, q_{\ell,t \frown k})$  is in  $(\dot{y} \cap \dot{E})$  etc., where  $q_{\ell,t \frown k} \restriction \alpha = q_{\ell,t} \restriction \alpha$  and  $q_{\ell,t \frown k}(\alpha) = (q_{\ell,t}(\alpha))_{t \frown k}$ .

Since some condition forces that  $L^-$  is not in  $(\dot{\mathcal{D}}_{i_y}^\alpha)^+$  it follows that  $L^-$  is not in  $(\dot{\mathcal{D}}_{i_\alpha}^\alpha)^+$

Finally we must show that  $r$  forces over  $\bar{G}^+$  that  $\dot{b}$  is disjoint from  $\dot{x}$ . Since each are  $\mathbb{P}_{i_p, \alpha+1}$ -names, it suffices to assume that  $\bar{r} \in \bar{G}^+$  is some  $\mathbb{P}_{i_p, \alpha+1}$ -reduct of  $r$  that forces some  $\ell$  is in  $\dot{b} \cap \dot{x}$ , and to then show that  $r$  fails to force that  $\ell \notin \dot{b} \cap (\dot{E} \cap \dot{y})$ . Choose  $(\ell, q_{\ell, t}) \in (\dot{y} \cap \dot{E})$  witnessing that  $\bar{r} \Vdash \ell \in \dot{x}$ . That is, we may assume that  $\bar{r} \upharpoonright \alpha \Vdash t = \text{stem}(\bar{r}(\alpha))$ , that  $q_{\ell, t} \upharpoonright \alpha \in \bar{G}^+$ , and  $(q_{\ell, t} \wedge r \upharpoonright \alpha)$  does not force over  $\bar{G}^+$  that  $t \notin \dot{T}_\alpha$ . Of course this means that the condition  $\bar{r} \wedge r \wedge [[t \in \dot{T}_\alpha]] \wedge q_{\ell, t}$  is not 0. This condition forces that  $\ell$  is in  $\dot{b} \cap (\dot{E} \cap \dot{y})$  as required.  $\square$

*Fact 5.* The character of  $\mathcal{D}_{i_\alpha}^\alpha$  is greater than  $\mu_{i_\gamma}$ .

*Proof of Fact 5.* We know that  $\mathcal{D}_{i_\alpha}^\alpha$  is forced to have an  $\omega$ -closed base (in fact, descending mod finite with uncountable cofinality). Even more,  $\mathbb{P}_{i_\alpha, \alpha}$  forces that for all  $T \in \mathbb{L}(\mathcal{D}_{i_\alpha}^\alpha)$ , there is a  $D \in \mathcal{D}_{i_\alpha}^\alpha$  such that the condition  $([D]^{<\omega})_{\text{stem}(T)}$  is below  $T$ . Let  $\chi_\alpha$  be the cofinality of  $\alpha$  and fix a list  $\{\dot{D}_\beta : \beta < \chi_\alpha\} \in M$  (closed under mod finite changes) of  $\mathbb{P}_{i_\alpha, \alpha}$ -names of elements of  $\dot{\mathcal{D}}_{i_\alpha}^\alpha$  that is forced to be a base.

Now, suppose that  $\dot{b} \in \mathbb{B}_{i_p, \alpha+1} = \mathbb{B}_{i_p, \alpha}$  and there is an  $\dot{E} \in \mathcal{E}_1$  and an  $r < p$  forcing that  $\dot{b} \cap (\dot{E} \cap \dot{y})$  is empty. We prove there is an  $\dot{x} \in \mathcal{E}_1$  and an  $r_2 < r \upharpoonright \alpha$  in  $\mathbb{P}_{i_\alpha, \alpha}$  such that  $r_2 \Vdash \dot{b} \cap \dot{x}$  is empty. We may assume that  $r_2$  forces a value  $t$  on  $\text{stem}(r(\alpha))$  and that, for some  $\beta < \chi_\alpha$ ,  $r_2 \Vdash (\dot{D}_\beta^{<\omega})_t < r(\alpha)$ . Let

$$\dot{x} = \{(\ell, q_\ell \upharpoonright \alpha) : (\ell, q_\ell) \in (\dot{E} \cap \dot{y}) \text{ and } q_\ell \upharpoonright \alpha \Vdash q_\ell(\alpha) \leq (\dot{D}_\beta^{<\omega})_t\}.$$

It is immediate that  $\dot{x} \in M$  and that  $(r_2 \wedge r) \Vdash_{\mathbb{P}_{i_\alpha, \alpha+1}} \dot{x} \supseteq (\dot{E} \cap \dot{y})$ . Since  $\dot{E} \cap \dot{y}$  is forced to be in  $\mathcal{E}_1^+$ , it follows that  $\dot{x}$  is forced by  $r_2$  to be in  $\langle \mathcal{E}_1 \rangle$ . Now we verify that  $r_2 \Vdash \dot{b} \cap \dot{x}$  is empty. Assume that  $r_3 < r_2$  in  $\mathbb{P}_{i_\alpha, \alpha}$  and that  $r_3 \Vdash \ell \in \dot{b} \cap \dot{x}$ . We may assume there is  $(\ell, q_\ell \upharpoonright \alpha) \in \dot{x}$  such that  $r_3 < q_\ell \upharpoonright \alpha$ . But now  $r_2 \Vdash q_\ell(\alpha) \leq r(\alpha)$  and so  $r_2 \wedge r \Vdash \ell \in \dot{b} \cap (\dot{E} \cap \dot{y})$  – a contradiction.

The conclusion now follows from Lemma 5.4.  $\square$

**Definition 5.7.** For each  $t \in \omega^{<\omega}$ , define that  $\mathbb{P}_{i_\alpha, \alpha}$ -name  $\dot{E}_t$  according to the rule that  $r \Vdash \ell \in \dot{E}_t$  providing  $r \in \mathbb{P}_{i_\alpha, \alpha}$  forces that there is a  $\dot{T}$  with  $r \Vdash \dot{T} \in \mathbb{L}(\dot{\mathcal{D}}_{i_\alpha}^\alpha)$ ,  $r \Vdash t = \text{stem}(\dot{T})$ , and  $r \cup \{(\alpha, \dot{T})\} \Vdash \ell \notin \dot{y}$ .

*Fact 6.* There is a  $\dot{T} \in \mathbb{L}(\dot{\mathcal{D}}_{i_\alpha}^\alpha) \cap M$  such that  $p \upharpoonright \alpha$  forces the statement:  $\dot{E}_t \in \mathcal{E}_1$  for all  $t$  such that  $\text{stem}(\dot{T}) \leq t \in \dot{T}$ .



*Proof of Fact 6.* By elementarity, there is a maximal antichain of  $\mathbb{P}_{i_\alpha, \alpha}$  each element of which decides if there is a  $\dot{T}$  with  $\dot{E}_t \in \mathcal{E}_1$  for all  $t \in \dot{T}$  above  $\text{stem}(\dot{T})$ . Since  $p \in A \setminus A_1$  it follows that there is an  $i_p < i_\alpha$  as in condition (2) of Fact 2. Let  $t_0 \in \omega^{<\omega}$  so that  $p \upharpoonright \alpha \Vdash t_0 = \text{stem}(p(\alpha))$ . By the maximum principle, there is a  $\dot{b} \in \mathbb{B}_{i_p, \gamma}$  and a  $\dot{E}_0 \in \mathcal{E}_1$  satisfying that  $p \Vdash \dot{b} \cap \dot{E}_0 \cap \dot{y}$  is empty, while  $p \Vdash \dot{b} \cap \dot{E}$  is infinite for all  $\dot{E} \in \langle \mathcal{E}_1 \rangle$ . This means that  $p$  forces that  $\dot{b} \cap \dot{E}_0$  is an element of  $\langle \mathcal{E}_1 \rangle^+$  that is contained in  $\omega \setminus \dot{y}$ . As in the proof of Lemma 5.4, there is an  $\dot{E}_2 \in \langle \mathcal{E}_1 \rangle \cap \mathbb{B}_{i_p, \gamma}$  such that  $p$  forces that  $\dot{b} \cap \dot{E}_2$  is contained in  $\dot{E}_0$ . We also have that  $(\dot{b} \cap \dot{E}_2) \upharpoonright \alpha$  is forced to be contained in  $\omega \setminus \dot{y}$ . It now follows that  $p \upharpoonright \alpha$  forces that for all  $t_0 \leq t \in p(\alpha)$ ,  $p \upharpoonright \alpha$  forces that  $\dot{E}_t$  contains  $(\dot{b} \cap \dot{E}_2) \upharpoonright \alpha$  and so is in  $\langle \mathcal{E}_1 \rangle^+$ . Since  $\dot{E}_t$  is also measured by  $\mathcal{E}_1$ , we have that  $p \upharpoonright \alpha$  forces that such  $\dot{E}_t$  are in  $\mathcal{E}_1$ . This completes the proof.  $\square$

Now we show how to extend  $\mathcal{E}_1 \cap \mathbb{B}_{i_\alpha, \gamma}$  so as to measure  $\dot{y}$ . Let  $\beta = \text{sup}(M \cap \alpha)$ . By Fact 5,  $\beta < \alpha$  and by the definition of  $\mathcal{H}(\vec{\lambda})$ ,  $M \cap \dot{\mathcal{D}}_{i_\alpha}^\alpha$  is a subset of  $\langle \dot{\mathcal{D}}_{i_\beta}^\beta \rangle$ ,  $\dot{L}_\beta \in \dot{\mathcal{D}}_{i_\alpha}^\alpha$ , and  $i_\beta = i_\alpha$ . We also have that the family  $\{\dot{L}_\xi : \text{cf}(\xi) \geq \omega_1 \text{ and } \beta_\alpha \leq \xi \in M \cap \beta\}$  is a base for  $\dot{\mathcal{D}}_{i_\beta}^\beta$ . For convenience let  $q <_M p$  denote the relation that  $q$  is an  $M \cap \mathbb{P}_{i_\alpha, \alpha+1}$ -reduct of  $p$ . Let  $\bar{p}$  be any condition in  $\mathbb{P}_{i_\beta, \beta+1}$  satisfying that  $\bar{p} \upharpoonright \beta = p \upharpoonright \alpha$  and  $\bar{p} \upharpoonright \beta \Vdash \text{stem}(\bar{p}(\beta)) = t_\alpha$  (recall that  $p \upharpoonright \alpha \Vdash t_\alpha = \text{stem}(p(\alpha))$ ).

Let us note that for each  $q \in M \cap \mathbb{P}_{i_\alpha, \alpha+1}$ ,  $q \upharpoonright \alpha = q \upharpoonright \beta$  and  $q \upharpoonright \beta \Vdash q(\alpha)$  is also a  $\mathbb{P}_{\beta, i_\beta}$ -name of an element of  $\mathbb{L}(\dot{\mathcal{D}}_{i_\beta}^\beta)$ . Let  $\dot{x}$  be the following  $\mathbb{P}_{i_\beta, \beta+1}$ -name

$$\dot{x} = \{(\ell, q \upharpoonright \beta \cup \{(\beta, q(\beta))\}) : (\ell, q) \in \dot{y} \cap M \text{ and } q <_M p\}.$$

We will complete the proof by showing that there is an extension of  $p$  that forces that  $\mathcal{E}_1 \cup \{\omega \setminus (\dot{x}[\dot{L}_\beta])\}$  measures  $\dot{y}$  and that 1 forces that  $\langle \mathcal{E}_1 \cup \{\omega \setminus (\dot{x}[\dot{L}_\beta])\} \rangle \cap \mathbb{B}_{i_\beta, \beta+1}$  is  $\vec{\lambda}(i_\gamma)$ -thin. Here  $\dot{x}[\dot{L}_\beta]$  abbreviates the  $\mathbb{P}_{i_\beta, \beta+1}$ -name

$$\{(\ell, r) : (\exists q) (\ell, q) \in \dot{x}, q \upharpoonright \beta = r \upharpoonright \beta, \text{ and } r \Vdash \text{stem}(q(\beta)) \in \dot{L}_\beta^{<\omega}\}.$$

The way to think of  $\dot{x}[\dot{L}_\beta]$  is that if  $\bar{p}$  is in some  $\mathbb{P}_{i_\alpha, \alpha}$ -generic filter  $G$ , then  $\dot{y}[G]$  is now an  $\mathbb{L}(\dot{\mathcal{D}}_{i_\alpha}^\alpha)$ -name,  $L_\beta^{<\omega} = (\dot{L}_\beta[G])^{<\omega}$  is in  $\mathbb{L}(\dot{\mathcal{D}}_{i_\alpha}^\alpha)$ , and  $(\dot{x}[\dot{L}_\beta])[G]$  is equal to  $\{\ell : L_\beta^{<\omega} \not\Vdash \ell \notin \dot{y}\}$ . We will use the properties of  $\dot{x}$  to help show that  $\mathcal{E}_1 \cup \{\omega \setminus (\dot{x}[\dot{L}_\beta])\}$  is  $\vec{\lambda}(i_\gamma)$ -thin. This semantic description of  $\dot{x}[\dot{L}_\beta]$  makes clear that  $\bar{p} \cup \{(\alpha, (\dot{L}_\beta)^{<\omega})\} \in \mathbb{P}_{i_\alpha, \alpha+1}$  forces that  $\dot{x}[\dot{L}_\beta]$  contains  $\dot{y}$ . This implies that  $\mathcal{E}_1 \cup \{\omega \setminus (\dot{x}[\dot{L}_\beta])\}$  measures  $\dot{y}$ .

Claim: It is forced by  $\bar{p}$  that  $\omega \setminus \dot{x}$  is not measured by  $\mathcal{E}_1$ .

Each element of  $\mathcal{E}_1$  is in  $M$  and simple elementarity will show that for any condition in  $q$  in  $M$  that forces  $\dot{E} \cap (\omega \setminus \dot{y})$  is infinite, the corresponding  $\bar{q} = q \upharpoonright \alpha \cup \{(\beta, q(\alpha))\}$  will also force that  $\dot{E} \cap (\omega \setminus \dot{x})$  is infinite.

It follows from Fact 5, with  $\omega \setminus \dot{x}$  playing the role of  $\dot{y}$ , that  $\mathcal{E}_1 \cup \{\omega \setminus \dot{x}\}$  is  $\vec{\lambda}(i_\gamma)$ -thin. Recall that  $q \Vdash \dot{x} = \emptyset$  for all  $q \perp \bar{p}$ . Now to prove that  $\mathcal{E}_1 \cup \{\omega \setminus (\dot{x}[\dot{L}_\beta])\}$  is also  $\vec{\lambda}(i_\gamma)$ -thin, we prove that

$$\langle \mathcal{E}_1 \cup \{\omega \setminus \dot{x}\} \rangle \cap \mathbb{B}_{i,\alpha} = \langle \mathcal{E}_1 \cup \{\omega \setminus (\dot{x}[\dot{L}_\beta])\} \rangle \cap \mathbb{B}_{i,\alpha}$$

for all  $i < i_\alpha$ . In fact, first we prove

$$\langle \mathcal{E}_1 \cup \{\omega \setminus \dot{x}\} \rangle \cap \mathbb{B}_{i,\beta} = \langle \mathcal{E}_1 \cup \{\omega \setminus (\dot{x}[\dot{L}_\beta])\} \rangle \cap \mathbb{B}_{i,\beta}$$

for all  $i < i_\alpha$ .

We begin with this main Claim.

**Claim 1.** If  $\dot{b} \in \mathbb{B}_{i,\beta}$  ( $i < i_\beta$ ) and there is an  $\dot{E} \in \mathcal{E}_1 \cap \mathbb{B}_{i_\alpha,\beta}$  and a  $\bar{p} \geq q \in \mathbb{P}_{i_\beta,\beta+1}$  such that  $q \Vdash \dot{b} \cap (\dot{E} \setminus \dot{x}) = \emptyset$  then  $q \upharpoonright \beta \Vdash (\exists \dot{E} \in \mathcal{E}_1) \dot{b} \cap \dot{E} = \emptyset$ .

*Proof of Claim:* We may assume that  $q \upharpoonright \beta$  forces a value  $t$  on  $\text{stem}(q(\beta))$ . Recall that  $q \upharpoonright \beta$  forces the statement: there is a  $\dot{D} \in M \cap \dot{\mathfrak{D}}_{i_\alpha}^\alpha$  such that  $(\dot{D}^{<\omega})_t \leq q(\beta)$ . The definition of  $\dot{x}$  ensures that  $q \upharpoonright \beta \cup \{(\alpha, (\dot{D}^{<\omega})_t)\} \Vdash \dot{b} \cap (\dot{E} \setminus \dot{y})$  is empty. There is a  $\mathbb{P}_{i_\alpha,\alpha}$ -name  $\dot{E}_1 \in M$  such that  $q \upharpoonright \alpha \Vdash \dot{E}_1 = \{\ell : (\dot{D}^{<\omega})_t \Vdash \ell \notin (\dot{E} \setminus \dot{y})\}$ . By assumption  $q \upharpoonright \alpha \Vdash \dot{E}_1 \in \langle \mathcal{E}_1 \rangle$ . Since  $\dot{b}$  is also a  $\mathbb{P}_{i_\alpha,\alpha}$ -name, we have that  $q \upharpoonright \alpha \Vdash \dot{b} \cap \dot{E}_1 = \emptyset$ .  $\square$

Now assume that  $\dot{b} \in \mathbb{B}_{i_\beta,\beta}$  and  $q \Vdash \dot{b} \cap (\dot{E} \cap (\omega \setminus (\dot{x}[\dot{L}_\beta])))$  is empty for some  $q < \bar{p}$  in  $\mathbb{P}_{i_\beta,\beta+1}$ . By Lemma 5.4 it suffices to assume that  $\dot{E} \in \mathbb{B}_{i_\beta,\beta}$ . To prove that  $q$  forces that  $\dot{b} \notin \langle \mathcal{E}_1 \rangle^+$ , it suffices to prove that there is some  $\dot{E}_1 \in \mathcal{E}_1$  such that  $q \Vdash \dot{b} \cap (\dot{E}_1 \cap (\omega \setminus \dot{x}))$  is finite. We proceed by contradiction.

We may again assume that  $q \upharpoonright \beta$  forces that  $q(\beta)$  is  $(\dot{D}^{<\omega})_t$  for some  $t \supset t_\alpha$  and some  $\dot{D} \in \dot{\mathfrak{D}}_{i_\alpha}^\alpha \cap M$ . Let  $H$  be the range of  $t$ . Let, for the moment,  $G$  be a  $\mathbb{P}_{i_\alpha,\alpha}$ -generic filter with  $q \in G$ . Now in  $M[G]$  we have the value  $L_\beta$  of  $\dot{L}_\beta$  and  $H \subset L_\beta$ . We can also let  $E$  denote the value of  $\dot{E}[G]$ . Recall that for each  $s \in H^{<\omega}$ ,  $E_s$  denotes the set of  $\ell \in E$  such that there is some  $T \in \mathbb{L}(\dot{\mathfrak{D}}_{i_\alpha}^\alpha)$  with  $s = \text{stem}(T)$  and  $T \Vdash \ell \notin \dot{y}$ . We have shown in Fact 6 that there is a  $T \in \mathbb{L}(\dot{\mathfrak{D}}_{i_\alpha}^\alpha) \cap M$  such that  $E_s \in \mathcal{E}_1$  for all  $s \in T$  above  $\text{stem}(T)$ . This means that there is an  $\ell \in \dot{b} \cap E$  such that  $\ell \in E_s$  for each of the finitely many suitable  $s$ . For each  $s$ , choose  $T_s \subset T$  witnessing  $\ell \in E_s$ . As before, and since there are only finitely many  $s$  involved, we can assume that  $\dot{T}_s = (\dot{D}^{<\omega})_s$  for

some  $H \subset \dot{D} \in \dot{\mathfrak{D}}_{i_\alpha}^\alpha \cap M$  and we then define an extension  $q$  of  $q'$  so that  $q'(\beta) = (\dot{D}^{<\omega})_{t_\alpha}$  ensures that  $(\dot{L}_\beta^{<\omega})_s < T_s$  for each  $s$ . Note that such a condition  $q'$  we have that  $q' \cup \{(\alpha, (\dot{L}_\beta^{<\omega})_s)\}$  forces that  $\ell \notin \dot{y}$ . But then it should be clear that  $q'$  that forces  $\ell \notin \dot{x}[\dot{L}_\beta]$ . This contradicts that  $q$  forces  $\ell \notin \dot{b} \cap (\dot{E} \cap (\omega \setminus (\dot{x}[\dot{L}_\beta])))$ .  $\square$

## REFERENCES

- [1] Andreas Blass and Saharon Shelah, *Ultrafilters with small generating sets*, Israel J. Math. **65** (1989), no. 3, 259–271, DOI 10.1007/BF02764864. MR1005010 (90e:03057)
- [2] Jörg Brendle, *The almost-disjointness number may have countable cofinality*, Trans. Amer. Math. Soc. **355** (2003), no. 7, 2633–2649 (electronic), DOI 10.1090/S0002-9947-03-03271-9. MR1975392 (2004c:03062)
- [3] Jörg Brendle and Vera Fischer, *Mad families, splitting families and large continuum*, J. Symbolic Logic **76** (2011), no. 1, 198–208, DOI 10.2178/jsl/1294170995. MR2791343 (2012d:03113)
- [4] Jörg Brendle and Saharon Shelah, *Ultrafilters on  $\omega$ —their ideals and their cardinal characteristics*, Trans. Amer. Math. Soc. **351** (1999), no. 7, 2643–2674, DOI 10.1090/S0002-9947-99-02257-6. MR1686797 (2000m:03111)
- [5] Kenneth Kunen and Jerry E. Vaughan (eds.), *Handbook of set-theoretic topology*, North-Holland Publishing Co., Amsterdam, 1984. MR776619 (85k:54001)
- [6] Saharon Shelah, *The character spectrum of  $\beta(\mathbb{N})$* , Topology Appl. **158** (2011), no. 18, 2535–2555, DOI 10.1016/j.topol.2011.08.014. MR2847327

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH CAROLINA AT CHARLOTTE, CHARLOTTE, NC 28223  
*E-mail address:* adow@uncc.edu

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, HILL CENTER, PISCATAWAY, NEW JERSEY, U.S.A. 08854-8019  
*Current address:* Institute of Mathematics, Hebrew University, Givat Ram, Jerusalem 91904, Israel  
*E-mail address:* shelah@math.rutgers.edu