

Postscript to SHELAH & FREMLIN P90

D.H.FREMLIN

University of Essex, Colchester

25 September 1991

1. The statement (‡) of SHELAH & FREMLIN P90, 2G, can be strengthened, as follows. Write (‡*) for the statement

there is a closed negligible set $Q \subseteq [0, 1]$ such that $(\mu_L)_*(Q^{-1}[D]) \geq \mu_L^*D$ for every $D \subseteq [0, 1]$.

Proposition If \mathbb{P} is a partially ordered set as in SHELAH & FREMLIN P90, then $\mathbb{1}_{\mathbb{P}} \Vdash_{\mathbb{P}} (\ddagger^*)$.

proof (a) Take X and \mathbb{P} and μ and Ψ as in SHELAH & FREMLIN P90, §1. Then if $D \subseteq X$ and $\epsilon > 0$, there is a closed set $F \subseteq X$, with $\mu F \geq \mu^*D - \epsilon$, such that

$$\mathbb{1}_{\mathbb{P}} \Vdash_{\mathbb{P}} \Psi \cap \ulcorner F \urcorner \subseteq \ulcorner R^{\neg-1}[D] \urcorner,$$

writing $\ulcorner F \urcorner$ for the \mathbb{P} -name for a closed subset of X corresponding to F .

P Choose k'_0 such that $2^{-k'_0} \leq \frac{1}{4}\epsilon$, and a closed set F_0 such that $\mu F_0 = \mu^*(D \cap F_0) \geq \mu^*D - \frac{1}{2}\epsilon$. Set

$$F = \{x : \forall k \geq k'_0, \mu^*(D \cap \{w : w|k = x|k\}) > 2^{-k+1}\mu\{w : w|k = x|k\}\}.$$

Then F is closed, and

$$F_0 \setminus F \subseteq \{x : x \in F_0, \exists k \geq k'_0, \mu(F_0 \cap \{w : w|k = x|k\}) \leq 2^{-k+1}\mu\{w : w|k = x|k\}\}$$

has measure at most $2^{-k'_0+1} \leq \frac{1}{2}\epsilon$, so $\mu F \geq \mu^*D - \epsilon$.

Now suppose that σ is a \mathbb{P} -name such that

$$\mathbb{1}_{\mathbb{P}} \Vdash_{\mathbb{P}} \sigma \in \Psi \cap \ulcorner F \urcorner.$$

Set $r = 1$, $L_k = n_k$ for every $k \in \mathbb{N}$ and follow the argument of Lemma 1R of SHELAH & FREMLIN P90 down to the end of part (c), but insisting at the beginning that $k_0 \geq k'_0$.

Observe that

$$p_3 \Vdash_{\mathbb{P}} \sigma \supseteq s,$$

where $s = \langle H_i(\mathbf{v}_i^*) \rangle_{i < k_1}$, as in part (d) of the proof of Lemma 1R. Also $\#(\tilde{J}_k) \leq 2^{-k}n_k$ for all $k \geq k_0$, so that

$$\mu\{x : s \subseteq x, x(k) \notin \tilde{J}_k \ \forall k \geq k_1\} > (1 - 2^{-k_1+1})\mu\{x : s \subseteq x\};$$

but as

$$p_3 \Vdash_{\mathbb{P}} \sigma \in \ulcorner F \urcorner, s \subseteq \sigma,$$

we must have

$$\mu^*(D \cap \{x : s \subseteq x\}) > 2^{-k_1+1}\mu\{x : s \subseteq x\},$$

and there is a $\tilde{z} \in D$ such that $s \subseteq \tilde{z}$ and $\tilde{z} \notin \tilde{J}_k$ for every $k \geq k_1$.

Now continue the argument as in (e)-(i) of the proof of Lemma 1R to get $p_5 \leq p_3$ such that $p_5 \Vdash_{\mathbb{P}} (\sigma, \tilde{z}) \in \ulcorner R \urcorner$. **Q**

(b) Now we find that

$$\mathbb{1}_{\mathbb{P}} \Vdash_{\mathbb{P}} \forall D \subseteq \ulcorner X \urcorner, \ulcorner \mu^{\neg} * (\ulcorner R^{\neg-1}[D] \urcorner) \urcorner \geq \ulcorner \mu^{\neg} * D \urcorner.$$

P Let Δ_0 be a \mathbb{P} -name for a subset of X , and ϵ, ϵ' (ground-model) rationals such that

$$\mathbb{1}_{\mathbb{P}} \Vdash_{\mathbb{P}} \ulcorner \mu^{\neg} * \Delta_0 \urcorner > \epsilon > \epsilon'.$$

Take $\beta < \kappa$ such that whenever Γ is a \mathbb{P}_β -name for a closed subset of X and

$$\mathbb{1}_{\mathbb{P}_\beta} \Vdash_{\mathbb{P}_\beta} \ulcorner \mu^{\neg} \Gamma \urcorner > 1 - \epsilon$$

then there is a \mathbb{P}_β -name for a member of $\Gamma \cap \Delta_0$. Now taking Δ to be a \mathbb{P}_β -name for the subset of X consisting of those members of Δ_0 which can be represented by \mathbb{P}_β -names, we see that

$$\mathbb{1}_{\mathbb{P}_\beta} \Vdash_{\mathbb{P}_\beta} \ulcorner \mu^{\neg} * \Delta \urcorner \geq \epsilon.$$

Using part (a) in $V^{\mathbb{P}_\beta}$, we have a \mathbb{P}_β -name Γ for a closed subset of X such that

$$\mathbb{1}_{\mathbb{P}_\beta} \Vdash_{\mathbb{P}_\beta} (\ulcorner \mu^{\neg} \Gamma \urcorner \geq \epsilon' \ \& \ \mathbb{1}_{\mathbb{P}'} \Vdash_{\mathbb{P}'} \Psi^{(\beta)} \cap \Gamma \subseteq \ulcorner R^{\neg-1}[\Delta] \urcorner),$$

expressing \mathbb{P} as an iteration $\mathbb{P}_\beta * \mathbb{P}'$ as in the proof of Theorem 1S of SHELAH & FREMLIN P90. But this must mean that

$$\mathbb{1}_{\mathbb{P}} \Vdash_{\mathbb{P}} \ulcorner \mu^{\neg} * (\ulcorner R^{\neg-1}[\Delta_0] \urcorner) \urcorner \geq \epsilon',$$

and as ϵ' was arbitrary this proves the claim. **Q**

(c) This proves the result for (X, μ) rather than for $([0, 1], \mu_L)$. But as remarked in 2G of SHELAH & FREMLIN P90, we have a continuous inverse-measure-preserving $f : X \rightarrow [0, 1]$ such that $\mu^*f^{-1}[D] = \mu_L^*[D]$

for every $D \subseteq [0, 1]$; so that setting $Q = \{(f(x), f(y)) : (x, y) \in \overline{R}\}$ we obtain the result for Lebesgue measure, as stated.

2. In fact we can go a little further: in $V^{\mathbb{P}}$, Q has the property that

whenever $D \subseteq [0, 1]$ and E is a measurable set such that $\mu_*(E \setminus D) = 0$, then $\mu(E \setminus Q^{-1}[D]) = 0$.

To see this, follow the arguments above, observing that it is enough to consider closed E , and that the set F of part (a) of the proof can be taken to be a subset of E .

Reference

Shelah S. & Fremlin D.H. [p90] 'Pointwise compact and stable sets of measurable functions', to appear in J. Symbolic Logic.