

The Ehrenfeucht-Fraïssé-game of length ω_1

by

Alan Mekler

Department of Mathematics
Simon Fraser University
Burnaby, B.C.
Canada

Saharon Shelah

Institute of Mathematics
Hebrew University
Jerusalem, Israel

Jouko Väänänen

Department of Mathematics
University of Helsinki
Helsinki, Finland

October 6, 2003

Abstract

Let \mathcal{A} and \mathcal{B} be two first order structures of the same vocabulary. We shall consider the *Ehrenfeucht-Fraïssé-game of length ω_1 of \mathcal{A} and \mathcal{B}* which we denote by $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$. This game is like the ordinary Ehrenfeucht-Fraïssé-game of $L_{\omega\omega}$ except that there are ω_1 moves. It is clear that $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ is determined if \mathcal{A} and \mathcal{B} are of cardinality $\leq \aleph_1$. We prove the following results:

Theorem 1 *If $V=L$, then there are models \mathcal{A} and \mathcal{B} of cardinality \aleph_2 such that the game $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ is non-determined.*

Theorem 2 *If it is consistent that there is a measurable cardinal, then it is consistent that $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ is determined for all \mathcal{A} and \mathcal{B} of cardinality $\leq \aleph_2$.*

Theorem 3 *For any $\kappa \geq \aleph_3$ there are \mathcal{A} and \mathcal{B} of cardinality κ such that the game $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ is non-determined.*

1 Introduction.

Let \mathcal{A} and \mathcal{B} be two first order structures of the same vocabulary L . We denote the domains of \mathcal{A} and \mathcal{B} by A and B respectively. All vocabularies are assumed to be relational.

The *Ehrenfeucht-Fraïssé-game of length γ of \mathcal{A} and \mathcal{B}* denoted by $\mathcal{G}_\gamma(\mathcal{A}, \mathcal{B})$ is defined as follows: There are two players called \forall and \exists . First \forall plays x_0 and then \exists plays y_0 . After this \forall plays x_1 , and \exists plays y_1 , and so on. If $\langle (x_\beta, y_\beta) : \beta < \alpha \rangle$ has been played and $\alpha < \gamma$, then \forall plays x_α after which \exists plays y_α . Eventually a sequence $\langle (x_\beta, y_\beta) : \beta < \gamma \rangle$ has been played. The rules of the game say that both players have to play elements of $A \cup B$. Moreover, if \forall plays his x_β in A (B), then \exists has to play his y_β in B (A). Thus the sequence $\langle (x_\beta, y_\beta) : \beta < \gamma \rangle$ determines a relation $\pi \subseteq A \times B$. Player \exists wins this round of the game if π is a partial isomorphism. Otherwise \forall wins. The notion of winning strategy is defined in the usual manner. We say that a player *wins* $\mathcal{G}_\gamma(\mathcal{A}, \mathcal{B})$ if he has a winning strategy in $\mathcal{G}_\gamma(\mathcal{A}, \mathcal{B})$.

Recall that

$$\begin{aligned} \mathcal{A} \equiv_{\omega\omega} \mathcal{B} &\iff \forall n < \omega (\exists \text{ wins } \mathcal{G}_n(\mathcal{A}, \mathcal{B})) \\ \mathcal{A} \equiv_{\infty\omega} \mathcal{B} &\iff \exists \text{ wins } \mathcal{G}_\omega(\mathcal{A}, \mathcal{B}). \end{aligned}$$

In particular, $\mathcal{G}_\gamma(\mathcal{A}, \mathcal{B})$ is determined for $\gamma \leq \omega$. The question, whether $\mathcal{G}_\gamma(\mathcal{A}, \mathcal{B})$ is determined for $\gamma > \omega$, is the subject of this paper. We shall concentrate on the case $\gamma = \omega_1$.

The notion

$$\exists \text{ wins } \mathcal{G}_\gamma(\mathcal{A}, \mathcal{B}) \tag{1}$$

can be viewed as a natural generalization of $\mathcal{A} \equiv_{\infty\omega} \mathcal{B}$. The latter implies isomorphism for countable models. Likewise (1) implies isomorphism for models of cardinality $|\gamma|$:

Proposition 1 *Suppose \mathcal{A} and \mathcal{B} have cardinality $\leq \kappa$. Then $\mathcal{G}_\kappa(\mathcal{A}, \mathcal{B})$ is determined: \exists wins if $\mathcal{A} \cong \mathcal{B}$, and \forall wins if $\mathcal{A} \not\cong \mathcal{B}$.*

Proof. If $f : \mathcal{A} \cong \mathcal{B}$, then the winning strategy of \exists in $\mathcal{G}_\kappa(\mathcal{A}, \mathcal{B})$ is to play in such a way that the resulting π satisfies $\pi \subseteq f$. On the other hand, if $\mathcal{A} \not\cong \mathcal{B}$, then the winning strategy of \forall is to systematically enumerate $A \cup B$ so that the final π will satisfy $A = \text{dom}(\pi)$ and $B = \text{rng}(\pi)$. \square

For models of arbitrary cardinality we have the following simple but useful criterion of (1), namely in the terminology of [15] that they are “potentially isomorphic”. We use $\text{Col}(\lambda, \kappa)$ to denote the notion of forcing which collapses $|\lambda|$ to κ (with conditions of cardinality less than κ).

Proposition 2 *Suppose \mathcal{A} and \mathcal{B} have cardinality $\leq \lambda$ and κ is regular. Player \exists wins $\mathcal{G}_\kappa(\mathcal{A}, \mathcal{B})$ if and only if $\Vdash_{\text{Col}(\lambda, \kappa)} \mathcal{A} \cong \mathcal{B}$.*

Proof. Suppose τ is a winning strategy of \exists in $\mathcal{G}_\kappa(\mathcal{A}, \mathcal{B})$. Since $\text{Col}(\lambda, \kappa)$ is $< \kappa$ -closed,

$$\Vdash_{\text{Col}(\lambda, \kappa)} \text{“}\tau \text{ is a winning strategy of } \exists \text{ in } \mathcal{G}_\kappa(\mathcal{A}, \mathcal{B})\text{”}.$$

Hence $\Vdash_{\text{Col}(\lambda, \kappa)} \mathcal{A} \cong \mathcal{B}$ by Proposition 1. Suppose then $p \Vdash \tilde{f} : \mathcal{A} \cong \mathcal{B}$ for some $p \in \text{Col}(\lambda, \kappa)$. While the game $\mathcal{G}_\kappa(\mathcal{A}, \mathcal{B})$ is played, \exists keeps extending the condition p further and further. Suppose he has extended p to q and \forall has played $x \in A$. Then \exists finds $r \leq q$ and $y \in B$ with $r \Vdash \tilde{f}(x) = y$. Using this simple strategy \exists wins. \square

Proposition 3 *Suppose T is an ω -stable first order theory with NDOP. Then $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ is determined for all models \mathcal{A} of T and all models \mathcal{B} .*

Proof. Suppose \mathcal{A} is a model of T . If \mathcal{B} is not $L_{\infty\omega_1}$ -equivalent to \mathcal{A} , then \forall wins $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ easily. So let us suppose $\mathcal{A} \equiv_{\infty\omega_1} \mathcal{B}$. We may assume A and B are of cardinality $\geq \aleph_1$. If we collapse $|A|$ and $|B|$ to \aleph_1 , T will remain ω -stable with NDOP, and \mathcal{A} and \mathcal{B} will remain $L_{\infty\omega_1}$ -equivalent. So \mathcal{A} and

\mathcal{B} become isomorphic by [18, Chapter XIII, Section 1]. Now Proposition 2 implies that \exists wins $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$. \square

Hyttinen [10] showed that $\mathcal{G}_\gamma(\mathcal{A}, \mathcal{B})$ may be non-determined for all γ with $\omega < \gamma < \omega_1$ and asked whether $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ may be non-determined. Our results show that $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ may be non-determined for \mathcal{A} and \mathcal{B} of cardinality \aleph_3 (Theorem 17), but for models of cardinality \aleph_2 the answer is more complicated.

Let $F(\omega_1)$ be the free group of cardinality \aleph_1 . Using the combinatorial principle \square_{ω_1} we construct an abelian group G of cardinality \aleph_2 such that $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ is non-determined (Theorem 4). On the other hand, we show that starting with a model with a measurable cardinal one can build a forcing extension in which $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ is determined for all models \mathcal{A} and \mathcal{B} of cardinality $\leq \aleph_2$ (Theorem 14).

Thus the free abelian group $F(\omega_1)$ has the remarkable property that the question

Is $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ determined for all G ?

cannot be answered in ZFC alone. Proposition 3 shows that no model of an \aleph_1 -categorical first order theory can have this property.

We follow Jech [11] in set theoretic notation. We use S_n^m to denote the set $\{\alpha < \omega_m : \text{cf}(\alpha) = \omega_n\}$. Closed and unbounded sets are called cub sets. A set of ordinals is λ -closed if it is closed under supremums of ascending λ -sequences $\langle \alpha_i : i < \lambda \rangle$ of its elements. A subset of a cardinal is λ -stationary if it meets every λ -closed unbounded subset of the cardinal. The closure of a set A of ordinals in the order topology of ordinals is denoted by \overline{A} . The free abelian group of cardinality κ is denoted by $F(\kappa)$.

2 A non-determined $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ with G a group of cardinality \aleph_2 .

In this section we use \square_{ω_1} to construct a group G of cardinality \aleph_2 such that the game $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ is non-determined (Theorem 4). For background on almost free groups the reader is referred to [4]. However, our presentation does not depend on special knowledge of almost free groups. All groups below are assumed to be abelian.

By \square_{ω_1} we mean the principle, which says that there is a sequence $\langle C_\alpha : \alpha < \omega_2, \alpha = \cup \alpha \rangle$ such that

1. C_α is a cub subset of α .
2. If $\text{cf}(\alpha) = \omega$, then $|C_\alpha| = \omega$.
3. If γ is a limit point of C_α , then $C_\gamma = C_\alpha \cap \gamma$.

Recall that \square_{ω_1} follows from $V = L$ by a result of R. Jensen. For a sequence of sets C_α as above we can let $E_\beta = \{\alpha \in S_0^2 : \text{the order type of } C_\alpha \text{ is } \beta\}$. For some $\beta < \omega_1$ the set E_β has to be stationary. Let us use E to denote this E_β . Then E is a so called *non-reflecting* stationary set, i.e., if $\gamma = \cup \gamma$ then $E \cap \gamma$ is non-stationary on γ . Indeed, then some final segment D_γ of the set of limit points of C_γ is a cub subset of γ disjoint from E . Moreover, $\text{cf}(\alpha) = \omega$ for all $\alpha \in E$.

Theorem 4 *Assuming \square_{ω_1} , there is a group G of cardinality \aleph_2 such that the game $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ is non-determined.*

Proof. Let \mathbb{Z}^{ω_2} denote the direct product of ω_2 copies of the additive group \mathbb{Z} of the integers. Let x_α be the element of \mathbb{Z}^{ω_2} which is 0 on coordinates $\neq \alpha$ and 1 on the coordinate α . Let us fix for each $\delta \in S_0^2$ an ascending cofinal sequence $\eta_\delta : \omega \rightarrow \delta$. For such δ , let

$$z_\delta = \sum_{n=0}^{\infty} 2^n x_{\eta_\delta(n)}.$$

Let $\langle C_\alpha : \alpha = \cup \alpha < \omega_2 \rangle$, $\langle D_\alpha : \alpha = \cup \alpha < \omega_2 \rangle$ and $E = E_\beta$ be obtained from \square_{ω_1} as above. We are ready to define the groups we need for the proof: Let G be the smallest pure subgroup of \mathbb{Z}^{ω_2} which contains x_α for $\alpha < \omega_2$ and z_δ for $\delta \in E$, let G_α be the smallest pure subgroup of \mathbb{Z}^{ω_2} which contains x_γ for $\gamma < \alpha$ and z_δ for $\delta \in E \cap \alpha$, let $F (= F(\omega_2))$ be the subgroup of \mathbb{Z}^{ω_2} generated freely by x_α for $\alpha < \omega_2$, and finally, let F_α be the subgroup of \mathbb{Z}^{ω_2} generated freely by x_γ for $\gamma < \alpha$.

The properties we shall want of G_α are standard but for the sake of completeness we shall sketch proofs. We need that each G_α is free and for any $\beta \notin E$ any free basis of G_β can be extended to a free basis of G_α for all $\alpha > \beta$.

The proof is by induction on α . For limit ordinals we use the fact that E is non-reflecting. The case of successors of ordinals not in E is also easy. Assume now that $\delta \in E$ and the induction hypothesis has been verified up to δ . By the induction hypothesis for any $\beta < \delta$ such that $\beta \notin E$, there is n_0 so that

$$G_\delta = G_\beta \oplus H \oplus K$$

where K is the group freely generated by $\{x_{\eta_\delta(n)} : n_0 \leq n\}$ and $x_{\eta_\delta(m)} \in G_\beta$ for all $m < n_0$. Then

$$G_{\delta+1} = G_\beta \oplus H \oplus K'$$

where K' is freely generated by

$$\left\{ \sum_{m=n}^{\infty} 2^{m-n} x_{\eta_\delta(m)} : n_0 \leq n \right\}.$$

On the other hand, if $\delta \in E$ and $\{x_{\eta_\delta(n)} : n < \omega\} \subseteq B$, where B is a subgroup of G such that $z_\delta \notin B$, then G/B is non-free, as $z_\delta + B$ is infinitely divisible by 2 in G/B .

Claim 1 \exists does not win $\mathcal{G}_{\omega_1}(F, G)$.

Suppose τ is a winning strategy of \exists . Let $\alpha \in E$ such that the pair (G_α, F_α) is closed under the first ω moves of τ , that is, if \forall plays his first ω moves inside $G_\alpha \cup F_\alpha$, then τ orders \exists to do the same. We shall play $G_{\omega_1}(F, A)$ pointing out the moves of \forall and letting τ determine the moves of \exists . On his move number $2n$ \forall plays the element $x_{\eta_\alpha(n)}$ of G_α . On his move number $2n + 1$ \forall plays some element of F_α . Player \forall plays his moves in F_α in such a way that during the first ω moves eventually some countable direct summand K of F_α as well as some countable $B \subseteq G_\alpha$ are enumerated. Let J be the smallest pure subgroup of G containing $B \cup \{z_\alpha\}$. During the next ω moves of $G_{\omega_1}(F, A)$ player \forall enumerates J and \exists responds by enumerating some $H \subseteq F$. Since τ is a winning strategy, H has to be a subgroup of F . But now H/K is free, whereas J/B is non-free, so \forall will win the game, a contradiction.

Claim 2 \forall does not win $\mathcal{G}_{\omega_1}(F, G)$.

Suppose τ is a winning strategy of \forall . If we were willing to use CH, we could just take α of cofinality ω_1 such that (F_α, G_α) is closed under τ , and derive a contradiction from the fact that $F_\alpha \cong G_\alpha$. However, since we do not want to assume CH, we have to appeal to a longer argument.

Let $\kappa = (2^\omega)^{++}$. Let \mathcal{M} be the expansion of $\langle H(\kappa), \in \rangle$ obtained by adding the following structure to it:

- (H1) The function $\delta \mapsto \eta_\delta$.
- (H2) The function $\delta \mapsto z_\delta$.
- (H3) The function $\alpha \mapsto C_\alpha$.
- (H4) A well-ordering $<$ of the universe.
- (H5) The winning strategy τ .

Let $\mathcal{N} = \langle N, \in, \dots \rangle$ be an elementary submodel of \mathcal{M} such that $\omega_1 \subseteq N$ and $N \cap \omega_2$ is an ordinal α of cofinality ω_1 .

Let $D_\alpha = \{\beta_i : i < \omega_1\}$ in ascending order. Since $C_{\beta_i} = C_\alpha \cap \beta_i$, every initial segment of C_α is in N . By elementarity, $G_{\beta_i} \in N$ for all $i < \omega_1$. Let ϕ be an isomorphism $G_\alpha \rightarrow F_\alpha$ obtained as follows: ϕ restricted to G_{β_0} is the $<$ -least isomorphism between the free groups G_{β_0} and F_0 . If ϕ is defined on all G_{β_j} , $j < i$, then ϕ is defined on G_{β_i} as the $<$ -least extension of $\bigcup_{j < i} \phi_{\beta_j}$ to an isomorphism between G_{β_i} and F_i . Recall that by our choice of D_α $G_{\beta_{i+1}}/G_{\beta_i}$ is free, so such extensions really exist.

We derive a contradiction by showing that \exists can play ϕ against τ for the whole duration of the game $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$. To achieve this we have to show that, when \exists plays his canonical strategy based on ϕ the strategy τ of \forall directs \forall to go on playing elements which are in N , that is, elements of $G_\alpha \cup F_\alpha$.

Suppose a sequence $s = \langle (x_\gamma, y_\gamma) : \gamma < \mu \rangle, \mu < \omega_1$, has been played. It suffices to show that $s \in N$. Choose β_i so that the elements of s are in $G_{\beta_i} \cup F_{\beta_i}$. Now s is uniquely determined by $\phi \upharpoonright G_{\beta_i}$ and τ . Note that because $C_{\beta_i} = C_\alpha \cap \beta_i$, $\phi \upharpoonright G_{\beta_i}$ can be defined inside N similarly as ϕ was defined above, using C_{β_i} instead of C_α . Thus $s \in N$ and we are done.

We have proved that $\mathcal{G}_{\omega_1}(F, G)$ is nondetermined. This clearly implies $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ is nondetermined. \square

Remark. R. Jensen [14, p. 286] showed that if \square_{ω_1} fails, then ω_2 is Mahlo in L . Therefore, if $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ is determined for all almost free groups

\mathcal{A} and \mathcal{B} of cardinality \aleph_2 , then ω_2 is Mahlo in L . If we start with \square_κ , we get an almost free group A of cardinality κ^+ such that $\mathcal{G}_{\omega_1}(F(\omega_1), A)$ is nondetermined.

3 $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ can be determined for all G .

In this section all groups are assumed to be abelian. It is easy to see that \exists wins $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ for any uncountable free group G , so in this exposition $F(\omega_1)$ is a suitable representative of all free groups. In the study of determinacy of $\mathcal{G}_{\omega_1}(F(\omega_1), \mathcal{A})$ for various \mathcal{A} it suffices to study groups \mathcal{A} , since for other \mathcal{A} player \forall easily wins the game.

Starting from a model with a Mahlo cardinal we construct a forcing extension in which $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ is determined, when G is any group of cardinality \aleph_2 . This can be extended to groups G of any cardinality, if we start with a supercompact cardinal.

In the proof of the next results we shall make use of *stationary logic* $L(aa)$. For the definition and basic facts about $L(aa)$ the reader is referred to [1]. This logic has a new quantifier $aa s$ quantifying over variables s ranging over countable subsets of the universe. A cub set of such s is any set which contains a superset of any countable subset of the universe and which is closed under unions of countable chains. The semantics of $aa s$ is defined as follows:

$$aa s \phi(s, \dots) \iff \phi(s, \dots) \text{ holds for a cub set of } s.$$

Note that a group of cardinality \aleph_1 is free if and only if it satisfies

$$aa s aa s'(s \subseteq s' \rightarrow s'/s \text{ is free}). \quad (2)$$

Proposition 5 *Let G be a group. Then the following conditions are equivalent:*

- (1) \exists wins $\mathcal{G}_{\omega_1}(F(\omega_1), G)$.
- (2) G satisfies (2).
- (3) G is the union of a continuous chain $\langle G_\alpha : \alpha < \omega_2 \rangle$ of free subgroups with $G_{\alpha+1}/G_\alpha$ \aleph_1 -free for all $\alpha < \omega_2$.

Proof. (1) implies (2): Suppose \exists wins $\mathcal{G}_{\omega_1}(F(\omega_1), G)$. By Proposition 2 we have $\Vdash_{Col(|G|, \omega_1)}$ “ G is free.” Using the countable completeness of $Col(|G|, \omega_1)$ it is now easy to construct a cub set S of countable subgroups of G such that if $A \in S$ then for all $B \in S$ with $A \subseteq B$ we have B/A free. Thus G satisfies (2). (2) implies (3) quite trivially. (3) implies (1): Suppose a continuous chain as in (3) exists. If we collapse $|G|$ to \aleph_1 , then in the extension the chain has length $< \omega_2$. Now we use Theorem 1 of [8]:

If a group A is the union of a continuous chain of $< \omega_2$ free subgroups $\{A_\alpha : \alpha < \gamma\}$ of cardinality $\leq \aleph_1$ such that each $A_{\alpha+1}/A_\alpha$ is \aleph_1 -free, then A is free.

Thus G is free in the extension and (1) follows from Proposition 2. \square

Let us consider the following principle:

(*) For all stationary $E \subseteq S_0^2$ and countable subsets a_α of $\alpha \in E$ such that a_α is cofinal in α and of order type ω there is a closed $C \subseteq \omega_2$ of order type ω_1 such that $\{\alpha \in E : a_\alpha \setminus C \text{ is finite}\}$ is stationary in C .

Lemma 6 *The principle (*) implies that $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ is determined for all groups G of cardinality \aleph_2 .*

Proof. Suppose G is a group of cardinality \aleph_2 . We may assume the domain of G is ω_2 . Let us assume G is \aleph_2 -free, as otherwise \forall easily wins. If we prove that G satisfies (2), then Proposition 5 implies that \exists wins $\mathcal{G}_{\omega_1}(F(\omega_1), G)$.

To prove (2), assume the contrary. By Proposition 5 we may assume that G can be expressed as the union of a continuous chain $\langle G_\alpha : \alpha < \omega_2 \rangle$ of free groups with $G_{\alpha+1}/G_\alpha$ non- \aleph_1 -free for $\alpha \in E$, $E \subseteq \omega_2$ stationary. By Fodor’s Lemma, we may assume $E \subseteq S_0^2$. Also we may assume that for all α , every ordinal in $G_{\alpha+1} \setminus G_\alpha$ is greater than every ordinal in G_α . Finally by intersecting with a closed unbounded set we may assume that for all $\alpha \in E$ the set underlying G_α is α . Choose for each $\alpha \in E$ some countable subgroup b_α of $G_{\alpha+1}$ with $b_\alpha + G_\alpha/G_\alpha$ non-free. Let $c_\alpha = b_\alpha \cap G_\alpha$. We will choose a_α so that any final segment generates a subgroup containing c_α . Enumerate c_α as $\{g_n : n < \omega\}$ such that each element is enumerated infinitely often. Choose an increasing sequence $(\alpha_n : n < \omega)$ cofinal in α so that for all n , $g_n \in G_{\alpha_n}$. Finally, for each n , choose $h_n \in G_{\alpha_{n+1}} \setminus G_{\alpha_n}$. Let $a_\alpha = \{h_n : n < \omega\} \cup \{h_n + g_n : n < \omega\}$. It is now easy to check that a_α is a

sequence of order type ω which is cofinal in α and any subgroup of G which contains all but finitely many of the elements of a_α contains c_α .

By (*) there is a continuous C of order type ω_1 such that $\{\alpha \in C : a_\alpha \setminus C \text{ is finite}\}$ is stationary in C . Let $D = \langle C \cup \sum_{\alpha \in C} b_\alpha \rangle$. Since $|D| \leq \aleph_1$, D is free.

For any $\alpha \in C$, let

$$D_\alpha = \langle (C \cap \alpha) \cup \left(\sum_{\beta \in (C \cap \alpha)} b_\beta \right) \rangle.$$

Note that $D = \bigcup_{\alpha \in C} D_\alpha$, each D_α is countable and for limit point δ of C , $D_\delta = \bigcup_{\alpha \in (C \cap \delta)} D_\alpha$. Hence there is an $\alpha \in C \cap E$ such that $a_\alpha \setminus C$ is finite and D/D_α is free. Hence $b_\alpha + D_\alpha/D_\alpha$ is free. But

$$b_\alpha + D_\alpha/D_\alpha \cong b_\alpha/b_\alpha \cap D_\alpha = b_\alpha/b_\alpha \cap G_\alpha,$$

which is not free, a contradiction. \square

For the next theorem we need a lemma from [6]. A proof is included for the convenience of the reader.

Lemma 7 [6] *Suppose λ is a regular cardinal and \mathbb{Q} is a notion of forcing which satisfies the λ -c.c. Suppose \mathcal{I} is a normal λ -complete ideal on λ and $\mathcal{I}^+ = \{S \subseteq \lambda : S \notin \mathcal{I}\}$. For all sets $S \in \mathcal{I}^+$ and sequences of conditions $\langle p_\alpha : \alpha \in S \rangle$, there is a set C with $\lambda \setminus C \in \mathcal{I}$ so that for all $\alpha \in C \cap S$,*

$$p_\alpha \Vdash_{\mathbb{Q}} \text{“}\{\beta : p_\beta \in \tilde{G}\} \in \mathcal{J}^+, \text{ where } \mathcal{J} \text{ is the ideal generated by } \mathcal{I}\text{”}.$$

Proof. Suppose the lemma is false. So there is an \mathcal{I} -positive set $S' \subseteq S$ such that for all $\alpha \in S'$ there is an extension r_α of p_α and a set $I_\alpha \in \mathcal{I}$ (note: I_α is in the ground model) so that

$$r_\alpha \Vdash \{\beta : p_\beta \in \tilde{G}\} \subseteq I_\alpha.$$

Let I be the diagonal union of $\{I_\alpha : \alpha \in S'\}$.

Suppose now that $\alpha < \beta$ and $\alpha, \beta \in (S' \setminus I)$. Since $\beta \notin I$, $r_\alpha \Vdash p_\beta \notin \tilde{G}$. Hence $r_\alpha \Vdash r_\beta \notin \tilde{G}$. So r_α, r_β are incompatible. Hence $\{r_\alpha : \alpha \in S' \setminus I\}$ is an antichain which, since S' is \mathcal{I} -positive, is of cardinality λ . This is a contradiction. \square

Theorem 8 *Assuming the consistency of a Mahlo cardinal, it is consistent that $(*)$ holds and hence that $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ is determined for all groups G of cardinality \aleph_2 .*

Proof. By a result of Harrington and Shelah [7] we may start with a Mahlo cardinal κ in which every stationary set of cofinality ω reflects, that is, if $S \subseteq \kappa$ is stationary, and $\text{cf}(\alpha) = \omega$ for $\alpha \in S$, then $S \cap \lambda$ is stationary in λ for some inaccessible $\lambda < \kappa$.

For any inaccessible λ let \mathbb{P}_λ be the Levy-forcing for collapsing λ to ω_2 . The conditions of \mathbb{P}_λ are countable functions $f : \lambda \times \omega_1 \rightarrow \lambda$ such that $f(\alpha, \beta) < \alpha$ for all α and β and each f is increasing and continuous in the second coordinate. It is well-known that \mathbb{P}_λ is countably closed and satisfies the λ -chain condition [11, p. 191].

Let $\mathbb{P} = \mathbb{P}_\kappa$. Suppose $p \in \mathbb{P}$ and

$p \Vdash \text{“}\tilde{E} \subseteq S_0^2 \text{ is stationary and } \forall \alpha \in \tilde{E} (\tilde{a}_\alpha \subseteq \alpha \text{ is cofinal in } \alpha \text{ and of order type } \omega)\text{”}$.

Let

$$S = \{\alpha < \kappa : \exists q \leq p (q \Vdash \alpha \in \tilde{E})\}.$$

For any $\alpha \in S$ let $p_\alpha \leq p$ such that $p_\alpha \Vdash \alpha \in \tilde{E}$. Since \mathbb{P} is countably closed, we can additionally require that for some countable $a_\alpha \subseteq \alpha$ we have $p_\alpha \Vdash \tilde{a}_\alpha = a_\alpha$.

The set S is stationary in κ , for if $C \subseteq \kappa$ is cub, then $p \Vdash C \cap \tilde{E} \neq \emptyset$, whence $C \cap S \neq \emptyset$. Also $\text{cf}(\alpha) = \omega$ for $\alpha \in S$. Let λ be inaccessible such that $S \cap \lambda$ is stationary in λ . We may choose λ in such a way that $\alpha \in S \cap \lambda$ implies $p_\alpha \in \mathbb{P}_\lambda$. By Lemma 7 there is a $\delta \in S \cap \lambda$ such that

$$p_\delta \Vdash_{\mathbb{P}_\lambda} \text{“}\tilde{E}_1 = \{\alpha < \lambda : p_\alpha \in \tilde{G}\} \text{ is stationary.”}$$

Let \mathbb{Q} be the set of conditions $f \in \mathbb{P}$ with $\text{dom}(f) \subseteq (\kappa \setminus \lambda) \times \omega_1$. Note that $\mathbb{P} \cong \mathbb{P}_\lambda \otimes \mathbb{Q}$. Let G be \mathbb{P} -generic containing p_δ and $G_\lambda = G \cap \mathbb{P}_\lambda$ for any inaccessible $\lambda \leq \kappa$. Then G_λ is \mathbb{P}_λ -generic and ω_2 of $V[G_\lambda]$ is λ . Let us work now in $V[G_\lambda]$. Thus λ is the current ω_2 , $E_1 = \{\alpha < \lambda : p_\alpha \in G_\lambda\}$ is stationary, and we have the countable sets $a_\alpha \subseteq \alpha$ for $\alpha \in E_1$. Since \mathbb{Q} collapses λ there is a name \tilde{f} such that

$$\Vdash_{\mathbb{Q}} \text{“}\tilde{f} : \omega_1 \rightarrow \lambda \text{ is continuous and cofinal.”}$$

More precisely \tilde{f} is the name for the function f defined by $f(\alpha) = \beta$ if and only if there is some $g \in G$ so that $g(\lambda, \alpha) = \beta$. Let \tilde{C} denote the range of \tilde{f} . We shall prove the following statement:

Claim: $\Vdash_{\mathbb{Q}} \{ \alpha < \lambda : a_\alpha \setminus \tilde{C} \text{ is finite} \}$ is stationary in \tilde{C} .

Suppose $q \in \mathbb{Q}$ so that $q \Vdash \text{“}\tilde{D} \subseteq \omega_1 \text{ is a cub.}”$ Let \mathcal{M} be an appropriate expansion of $\langle H(\kappa), \in \rangle$ and $\langle \mathcal{N}_i : i < \lambda \rangle$, $\mathcal{N}_i = \langle N_i, \in, \dots \rangle$, a sequence of elementary submodels of \mathcal{M} such that:

- (i) Everything relevant is in N_0 .
- (ii) If $\alpha_i = N_i \cap \lambda$, then $\alpha_i < \alpha_j$ for $i < j < \lambda$.
- (iii) N_{i+1} is closed under countable sequences.
- (iv) $|N_i| = \omega_1$.
- (v) $N_i = \bigcup_{j < i} N_j$ for i a limit ordinal.

Choose $\gamma = \alpha_i \in E_1$ and let $\langle i_n : n < \omega \rangle$ be a sequence of successor ordinals such that $\gamma = \sup\{\alpha_{i_n} : n < \omega\}$. Let $q_0 \leq q$ and $\beta_0 \in \omega_1$ such that $q_0, \beta_0 \in N_{i_0}$,

$$q_0 \Vdash \text{“}\beta_0 \in \tilde{D}\text{”}$$

and q_0 decides the value of $\tilde{f}''\beta_0$ (which will by elementarity necessarily be a subset of α_{i_0}).

If q_n and β_n are defined we choose $q_{n+1} \leq q_n$ and $\beta_{n+1} \in \omega_1$ such that $q_{n+1}, \beta_{n+1} \in N_{i_{n+1}}$,

$$q_{n+1} \Vdash \text{“}\beta_{n+1} \in \tilde{D} \text{ and } a_\gamma \cap (\alpha_{i_{n+1}} \setminus \alpha_{i_n}) \subseteq \tilde{f}''\beta_{n+1} \subseteq \alpha_{i_{n+1}}\text{”}$$

and q_{n+1} decides $\tilde{f}''\beta_{n+1}$. Finally, let $q_\omega = \bigcup\{q_n : n < \omega\}$ and $\beta = \bigcup\{\beta_n : n < \omega\}$. Then

$$q_\omega \Vdash \text{“}\beta \in \tilde{D} \text{ and } a_\gamma \setminus \tilde{f}''\beta \text{ is finite.}”$$

The claim, and thereby the theorem, is proved. \square

Corollary 9 *The statement that $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ is determined for every structure \mathcal{A} of cardinality \aleph_2 and every uncountable free group \mathcal{B} , is equiconsistent with the existence of a Mahlo cardinal.*

Remark. If $\mathcal{G}_{\omega_1}(A, F(\omega_1))$ is determined for all groups A of cardinality κ^+ , κ singular, then \square_κ fails. This implies that the Covering Lemma fails for the Core Model, whence there is an inner model for a measurable cardinal. This shows that the conclusion of Theorem 8 cannot be strengthened to arbitrary G . However, by starting with a larger cardinal we can make this extension:

Theorem 10 *Assuming the consistency of a supercompact cardinal, it is consistent that $\mathcal{G}_{\omega_1}(F(\omega_1), G)$ is determined for all groups G .*

Proof. Let us assume that the stationary logic $L_{\omega_1\omega}(aa)$ has the Löwenheim-Skolem property down to \aleph_1 . This assumption is consistent relative to the consistency of a supercompact cardinal [2]. Let G be an arbitrary \aleph_2 -free group. Let H be an $L(aa)$ -elementary submodel of G of cardinality \aleph_1 . Thus H is a free group. The group H satisfies the sentence (2), whence so does G . Now the claim follows from Proposition 5. \square

Corollary 11 *Assuming the consistency of a supercompact cardinal, it is consistent that $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ is determined for every structure \mathcal{A} and every uncountable free group \mathcal{B} .*

4 $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ can be determined for all \mathcal{A} and \mathcal{B} of cardinality \aleph_2 .

We prove the consistency of the statement that $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ is determined for all \mathcal{A} and \mathcal{B} of cardinality $\leq \aleph_2$ assuming the consistency of a measurable cardinal. Actually we make use of an assumption that we call $I^*(\omega)$ concerning stationary subsets of ω_2 . This assumption is known to imply that ω_2 is measurable in an inner model. It follows from the previous section that some large cardinal axioms are needed to prove the stated determinacy.

Let $I^*(\omega)$ be the following assumption about ω_1 -stationary subsets of ω_2 :

$I^*(\omega)$ Let \mathcal{I} be the ω_1 -nonstationary ideal NS_{ω_1} on ω_2 . Then \mathcal{I}^+ has a σ -closed dense subset K .

Hodges and Shelah [9] define a principle $I(\omega)$, which is like $I^*(\omega)$ except that \mathcal{I} is not assumed to be the ω_1 -nonstationary ideal. They use $I(\omega)$ to prove

the determinacy of an Ehrenfeucht-Fraïssé-game played on several boards simultaneously.

Note that $I^*(\omega)$ implies \mathcal{I} is precipitous, so the consistency of $I^*(\omega)$ implies the consistency of a measurable cardinal [12].

Theorem 12 ([12]) *The assumption $I^*(\omega)$ is consistent relative to the consistency of a measurable cardinal.*

We shall consider models \mathcal{A}, \mathcal{B} of cardinality \aleph_2 , so we may as well assume they have ω_2 as universe. For such \mathcal{A} and $\alpha < \omega_2$ we let \mathcal{A}_α denote the structure $\mathcal{A} \cap \alpha$. Similarly \mathcal{B}_α .

Lemma 13 *Suppose \mathcal{A} and \mathcal{B} are structures of cardinality \aleph_2 . If \forall does not have a winning strategy in $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$, then*

$$S = \{\alpha : \mathcal{A}_\alpha \cong \mathcal{B}_\alpha\}$$

is ω_1 -stationary.

Proof. Let $C \subseteq \omega_2$ be ω_1 -closed and unbounded. Suppose $S \cap C = \emptyset$. We derive a contradiction by describing a winning strategy of \forall : Let $\pi : \omega_1 \rightarrow \omega_1 \times \omega_1 \times 2$ be onto with $\alpha, \beta, d \leq \pi(\alpha, \beta, d)$ for all $\alpha, \beta < \omega_1$ and $d < 2$. If $\alpha < \omega_2$, let $\theta_\alpha : \omega_1 \rightarrow \alpha$ be onto. Suppose the sequence $\langle (x_i, y_i) : i < \alpha \rangle$ has been played. Here x_i denotes a move of \forall and y_i a move of \exists . During the game \forall has built an ascending sequence $\{c_i : i < \alpha\}$ of elements of C . Now he lets c_α be the smallest element of C greater than all the elements $x_i, y_i, i < \alpha$. Suppose $\pi(\alpha) = (i, \gamma, d)$. Now \forall will play $\theta_{c_i}(\gamma)$ as an element of \mathcal{A} , if $d = 0$, and as an element of \mathcal{B} if $d = 1$.

After all ω_1 moves of $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ have been played, some \mathcal{A}_α and \mathcal{B}_α , where $\alpha \in C$, have been enumerated. Since $\alpha \notin S$, \forall has won the game. \square

Theorem 14 *Assume $I^*(\omega)$. The game $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ is determined for all \mathcal{A} and \mathcal{B} of cardinality $\leq \aleph_2$.*

Proof. Suppose \forall does not have a winning strategy. By Lemma 13 the set $S = \{\alpha : \mathcal{A}_\alpha \cong \mathcal{B}_\alpha\}$ is ω_1 -stationary. Let I and K be as in $I^*(\omega)$. If $\alpha \in S$, let $h_\alpha : \mathcal{A}_\alpha \cong \mathcal{B}_\alpha$. We describe a winning strategy of \exists . The idea of this strategy is that \exists lets the isomorphisms h_α determine his moves. Of course,

different h_α may give different information to \exists , so he has to decide which h_α to follow. The key point is that \exists lets some h_α determine his move only if there are stationarily many other h_β that agree with h_α on this move.

Suppose the sequence $\langle (x_i, y_i) : i < \alpha \rangle$ has been played. Again x_i denotes a move of \forall and y_i a move of \exists . Suppose \forall plays next x_α and this is (say) in A . During the game \exists has built a descending sequence $\{S_i : i < \alpha\}$ of elements of K with $S_0 \subseteq S$. The point of the sets S_i is that \exists has taken care that for all $i < \alpha$ and $\beta \in S_i$ we have $y_i = h_\beta(x_i)$ or $x_i = h_\beta(y_i)$ depending on whether \forall played x_i in A or B . Now \exists lets $S'_\alpha \subseteq \bigcap_{i < \alpha} S_i$ so that $S'_\alpha \in K$ and $\forall i \in S'_\alpha (x_\alpha < i)$. For each $i \in S'_\alpha$ we have $h_i(x_\alpha) < i$. By normality, there are an $S_\alpha \subseteq S'_\alpha$ in K and a y_α such that $h_i(x_\alpha) = y_\alpha$ for all $i \in S_\alpha$. This element y_α is the next move of \exists . Using this strategy \exists wins. \square

5 A non-determined $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ with \mathcal{A} and \mathcal{B} of cardinality \aleph_3 .

We construct directly in ZFC two models \mathcal{A} and \mathcal{B} of cardinality \aleph_3 with $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ non-determined. It readily follows that such models exist in all cardinalities $\geq \aleph_3$. The construction uses a square-like principle (Lemma 16), which is provable in ZFC.

Lemma 15 [17, 19] *There is a stationary $X \subseteq S_1^3$ and a sequence $\langle D_\alpha : \alpha \in X \rangle$ such that*

1. D_α is a cub subset of α for all $\alpha \in X$.
2. The order type of D_α is ω_1 .
3. If $\alpha, \beta \in X$ and $\gamma < \min\{\alpha, \beta\}$ is a limit of both D_α and D_β , then $D_\alpha \cap \gamma = D_\beta \cap \gamma$.
4. If $\gamma \in D_\alpha$, then γ is a limit point of D_α if and only if γ is a limit ordinal.

Proof. We shall sketch, for completeness, a proof of this given by Burke and Magidor [3, Lemma 7.7].

Let $<^*$ be a well-ordering of $H(\omega_3)$. For each $\alpha \in S_1^3$, let $\langle N_\delta^\alpha : \delta < \omega_2 \rangle$ be a continuously increasing chain of elementary submodels of $\langle H(\omega_3), \in, <^* \rangle$ such that

(N1) $(\omega_1 + 1) \cup \{\omega_2, \alpha\} \subseteq N_0^\alpha$.

(N2) $|N_\delta^\alpha| \leq \omega_1$.

(N3) $N_\delta^\alpha \cap \omega_2 \in \omega_2$.

(N4) $\overline{N_\delta^\alpha \cap \omega_3} \in N_{\delta+1}^\alpha$.

Let $A_\delta^\alpha = N_\delta^\alpha \cap \alpha$ for each $\alpha \in S_1^3$. Since, $\alpha \in N_\delta^\alpha$, A_δ^α is cofinal in α . Let $X \subseteq S_1^3$ be stationary such that for some $\delta, \rho < \omega_2$ and for all $\alpha \in X$ we have

1. $\delta =$ least ordinal of cofinality ω_1 with $N_\delta^\alpha \cap \omega_2 = \delta$.
2. The order type of $\overline{A_\delta^\alpha}$ is $\rho + 1$.

Let $f : \omega_1 \rightarrow \rho$ be cofinal and continuous. Let $g : \rho + 1 \cong \overline{A_\delta^\alpha}$ such that gf maps successors to successors. Let D_α be the image of ω_1 under gf . \square

Lemma 16 *There are sets S, T and C_α for $\alpha \in S$ such that the following hold:*

1. $S \subseteq S_0^3 \cup S_1^3$ and $S \cap S_1^3$ is stationary.
2. $T \subseteq S_0^3$ is stationary and $S \cap T = \emptyset$.
3. If $\alpha \in S$, then $C_\alpha \subseteq \alpha \cap S$ is closed and of order-type $\leq \omega_1$.
4. If $\alpha \in S$ and $\beta \in C_\alpha$, then $C_\beta = C_\alpha \cap \beta$.
5. If $\alpha \in S \cap S_1^3$, then C_α is cub on α .

Proof. Let S and $\langle D_\alpha : \alpha \in S \rangle$ be as in Lemma 15. Let $S' = X \cup Y$, where Y consists of ordinals which are limit points $< \alpha$ of some $D_\alpha, \alpha \in X$. If $\alpha \in X$, we let C_α be the set of limit points $< \alpha$ of D_α . If $\alpha \in Y$, we let C_α be the set of limit points $< \alpha$ of $D_\beta \cap \alpha$, where $\beta > \alpha$ is chosen arbitrarily from X .

Now claims 1,3,4 and 6 are clearly satisfied.

Let $S_0^3 = \bigcup_{i < \omega_2} T_i$ where the T_i are disjoint stationary sets. Since $|\overline{C_\alpha}| \leq \omega_1$, there is $i_\alpha < \omega_2$ such that $i \geq i_\alpha$ implies $\overline{C_i} \cap T_i = \emptyset$. Let $S'' \subseteq S'$ be stationary such that $\alpha \in S''$ implies i_α is constant i . Let $T = T_i$. Finally, let $S = S'' \cup \bigcup \{C_\alpha : \alpha \in S''\}$. Claim 2 is satisfied, and the Lemma is proved. \square

Theorem 17 *There are structures \mathcal{A} and \mathcal{B} of cardinality \aleph_3 with one binary predicate such that the game $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ is non-determined.*

Proof. Let S, T and $\langle C_\alpha : \alpha \in S \rangle$ be as in Lemma 16. We shall construct a sequence $\{M_\alpha : \alpha < \omega_3\}$ of sets and a sequence $\{G_\alpha : \alpha \in S\}$ of functions such that the conditions (M1)–(M6) below hold. Let W_α be the set of all mappings

$$G_{\gamma_0}^{d_0} \dots G_{\gamma_n}^{d_n},$$

where $\gamma_0, \dots, \gamma_n \in S \cap \alpha$, $d_i \in \{-1, 1\}$, G_γ^1 means G_γ and G_γ^{-1} means the inverse of G_γ . Let $W = W_{\omega_3}$. (Note that W consists of a set of partial functions.)

The conditions on the M_α 's and the G_α 's are:

- (M1) $M_\alpha \subseteq M_\beta$ if $\alpha < \beta$, and $M_\alpha \subset M_{\alpha+1}$ if $\alpha \in S$.
- (M2) $M_\nu = \bigcup_{\alpha < \nu} M_\alpha$ for limit ν .
- (M3) G_α is a bijection of $M_{\alpha+1}$ for $\alpha \in S$.
- (M4) If $\beta \in S$ and $\alpha \in C_\beta$, then $G_\alpha \subseteq G_\beta$.
- (M5) If for some β , $G_\beta(a) = b$ and for some $w \in W$, $w(a) = b$, then there is some γ so that $w \subseteq G_\gamma$. Furthermore if β is the minimum ordinal so that $G_\beta(a) = b$ then $\gamma = \beta$ or $\beta \in C_\gamma$.

In order to construct the set $M = \bigcup_{\alpha < \omega_3} M_\alpha$ and the mappings G_α we define an oriented graph with M as the set of vertices. We use the terminology of Serre [16] for graph-theoretic notions. If x is an edge, the origin of x is denoted by $o(x)$ and the terminus by $t(x)$. Our graph has an inverse edge \bar{x} for each edge x . Thus $o(\bar{x}) = t(x)$ and $t(\bar{x}) = o(x)$. Some edges are called *positive*, the rest are called *negative*. An edge is positive if and only if its inverse is negative. For each edge x of M there is a set $L(x)$ of labels. The set of possible labels for positive edges is $\{g_\alpha : \alpha < \omega_3\}$. The negative edges

can have elements of $\{g_\alpha^{-1} : \alpha < \omega_3\}$ as labels. The labels are assumed to be given in such a way that a positive edge gets g_α as a label if and only if its inverse gets the label g_α^{-1} . During the construction the sets of labels will be extended step by step.

The construction is analogous to building an acyclic graph on which a group acts freely. The graph then turns out to be the Cayley graph of the group. The labelled graph we will build will be the ‘‘Cayley graph’’ of W which will be as free as possible given (M1)–(M4). Condition (M5) is a consequence of the freeness of the construction.

Let us suppose the sets $M_\beta, \beta < \alpha$, of vertices have been defined. Let $M_{<\alpha} = \bigcup_{\beta < \alpha} M_\beta$. Some vertices in $M_{<\alpha}$ have edges between them and a set $L(x)$ of labels has been assigned to each such edge x .

If α is a limit ordinal, we let $M_\alpha = M_{<\alpha}$. So let us assume $\alpha = \beta + 1$. If $\beta \notin S$, $M_\alpha = M_\beta$. So let us assume $\beta \in S$. Let $\gamma = \sup(C_\beta)$. Notice that since S consists entirely of limit ordinals and $C_\beta \subseteq S$, either $\gamma = \beta$ or $\gamma + 1 < \beta$.

Case 1. $\gamma = \beta$: We extend M_β to M_α by adding new vertices $\{P_z : z \in \mathbb{Z}\}$ and for each $z \in \mathbb{Z}$ a positive edge $x_\alpha^{P_z}$ with $o(x_\alpha^{P_z}) = P_z$ and $t(x_\alpha^{P_z}) = P_{z+1}$. We also let $L(x_\alpha^{P_z}) = \{g_\beta\} \cup \{g_\delta : \beta \in C_\delta\}$.

Case 2. $\gamma + 1 < \beta$: We extend M_β to M_α by adding new vertices $\{P'_z : z \in \mathbb{Z} \setminus \{0\}\}$ for each $P \in M_\beta \setminus M_{\gamma+1}$. For notational convenience let $P'_0 = P$. Now we add for each $P \in M_\beta \setminus M_{\gamma+1}$ new edges as follows. For each $z \in \mathbb{Z}$ we add a positive edge $x_\alpha^{P'_z}$ with

$$o(x_\alpha^{P'_z}) = P'_z, t(x_\alpha^{P'_z}) = P'_{z+1}, L(x_\alpha^{P'_z}) = \{g_\beta\} \cup \{g_\delta : \beta \in C_\delta\}$$

This determines completely the inverse of $x_\alpha^{P'_z}$.

This ends the construction of the graph. In the construction each vertex P in $M_{\alpha+1}, \alpha \in S$, is made the origin of a unique edge x_α^P with $g_\alpha \in L(x_\alpha^P)$. We define

$$G_\alpha(P) = t(x_\alpha^P).$$

The construction of the sets M_α and the mappings G_α is now completed. It follows immediately from the construction that each $G_\alpha, \alpha \in S$, is a bijection of $M_{\alpha+1}$. So (M1)–(M3) hold. (M4) holds, because g_α is added to the labels of any edge with g_β , where $\beta \in C_\alpha$, as a label. Finally, (M5) is a consequence of the fact that the graph is circuit-free.

Let us fix $a_0 \in M_1$ and $b_0 = G_{\beta_0}(a_0)$, where $\beta_0 \in C_\alpha$ for all $\alpha \in S$. Note that we may assume, without loss of generality, the existence of such a β_0 .

If $a_0, a_1 \in M$, let

$$R_{(a_0, a_1)} = \{(a'_0, a'_1) \in M^2 : \exists w \in W(w(a_0) = a'_0 \wedge w(a_1) = a'_1)\}.$$

We let

$$\mathcal{M} = \langle M, (R_{(a_0, a_1)})_{(a_0, a_1) \in M^2} \rangle$$

$$\mathcal{A} = \langle \mathcal{M}, a_0 \rangle$$

$$\mathcal{B} = \langle \mathcal{M}, b_0 \rangle$$

and show that $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ is non-determined.

The reduction of the language of \mathcal{A} and \mathcal{B} to one binary predicate is easy. One just adds a copy of ω_3 , together with its ordering, and a copy of $M \times M$ to the structures with the projection maps. Then fix a bijection ϕ from ω_3 to M^2 . Add a new binary predicate R to the language and interpret R to be contained in $\omega_3 \times M^2$ such that $R(\beta, (a, b))$ holds if and only if $R_{\phi(\alpha)}(a, b)$ holds. We can now dispense with the old binary predicates. We have replaced our structure by one in a finite language without making any difference to who wins the game $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$. The extra step of reducing to a single binary predicate is standard.

An important property of these models is that if $\alpha \in S \cap S_1^3$, then $G_\alpha \upharpoonright M_\alpha$ is an automorphism of the restriction of \mathcal{M} to M_α and takes a_0 to b_0 .

Claim 3 \forall does not win $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$.

Suppose \forall has a winning strategy τ . Again, there is a quick argument which uses CH: Find $\alpha \in S$ such that M_α is closed under τ and $\text{cf}(\alpha) = \omega_1$. Now C_α is cub on α , whence G_α maps M_α onto itself. Using G_α player \exists can easily beat τ , a contradiction.

In the following longer argument we need not assume CH. Let κ be a large regular cardinal. Let \mathcal{H} be the expansion of $\langle H(\kappa), \in \rangle$ obtained by adding the following structure to it:

(H1) The function $\alpha \mapsto M_\alpha$.

(H2) The function $\alpha \mapsto G_\alpha$.

(H3) The function $\alpha \mapsto C_\alpha$.

(H4) A well-ordering $<^*$ of the universe.

(H5) The winning strategy τ .

(H6) The sets S and T .

Let $\mathcal{N} = \langle N, \in, \dots \rangle$ be an elementary submodel of \mathcal{H} such that $\alpha = N \cap \omega_3 \in S \cap S_1^3$.

Now C_α is a cub of order-type ω_1 on α and G_α maps M_α onto M_α . Moreover, G_α is a partial isomorphism from \mathcal{A} into \mathcal{B} . Provided that τ does not lead \forall to play his moves outside M_α , \exists has an obvious strategy: he lets G_α determine his moves. So let us assume a sequence $\langle (x_\xi, y_\xi) : \xi < \gamma \rangle$ has been played inside M_α and $\gamma < \omega_1$. Let $\beta \in C_\alpha$ such that M_β contains the elements x_ξ, y_ξ for $\xi < \gamma$. The sequence $\langle y_\xi : \xi < \gamma \rangle$ is totally determined by G_β and τ . Since $G_\beta \in N$, $\langle y_\xi : \xi < \gamma \rangle \in N$, and we are done.

Claim 4 \exists does not win $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$.

Suppose \exists has a winning strategy τ . Let \mathcal{H} be as above and $\mathcal{N} = \langle N, \in, \dots \rangle$ be an elementary submodel of \mathcal{H} such that $\alpha = N \cap \omega_3 \in T$. We let \forall play during the first ω moves of $\mathcal{G}_{\omega_1}(\mathcal{A}, \mathcal{B})$ a sequence $\langle a_n : n < \omega \rangle$ in \mathcal{A} such that if α_n is the least α_n with $a_n \in M_{\alpha_n}$, then the sequence $\langle \alpha_n : n < \omega \rangle$ is ascending and $\sup\{\alpha_n : n < \omega\} = \alpha$. Let \exists respond following τ with $\langle b_n : n < \omega \rangle$. As his move number ω player \forall plays some element $a_\omega \in M \setminus M_\alpha$ in \mathcal{A} and \exists answers according to τ with b_ω .

For all $i \leq \omega$, $R_{(a_0, a_i)}(a_0, a_i)$ holds. Hence $R_{(a_0, a_i)}(b_0, b_i)$ holds. So there is w_i such that $w_i(a_0) = b_0$ and $w_i(a_i) = b_i$. Since $G_{\beta_0}(a_0) = b_0$, by (M5), for each i there is β_i so that $G_{\beta_i}(a_i) = b_i$. We can assume that β_i is chosen to be minimal. Notice that for all i , $\beta_i > \alpha_i$ and for $i < \omega$, $\beta_i \in \mathcal{N}$. So $\sup\{\beta_i : i < \omega\} = \alpha$.

Also, by the same reasoning as above, for each $i < \omega$, $R_{(a_i, a_\omega)}(b_i, b_\omega)$ holds. Applying (M5), we get that $G_{\beta_\omega}(a_i) = b_i$. Using (M5) again and the minimality of β_i , for all $i < \omega$, $\beta_i \in C_{\beta_\omega}$. Thus α is a limit of elements of C_{β_ω} , contradicting $\alpha \in T$. \square

References

- [1] J. Barwise, M. Kaufmann and M. Makkai, Stationary logic, *Ann. Math. Logic* 13 (1978), 171–224.
- [2] S. Ben-David, On Shelah’s compactness of cardinals, *Israel J. of Mathematics* 31 (1978), 34–56.
- [3] M. Burke and M. Magidor, Shelah’s pcf theory and its applications, manuscript.
- [4] P. Eklof and A. Mekler, *Almost free modules*, North-Holland, Amsterdam, 1990.
- [5] F. Galvin, T. Jech and M. Magidor, An ideal game, *Journal of Symbolic Logic* 43 (1978), 285–292.
- [6] M. Gitik and S. Shelah, Cardinal preserving ideals, *Journal of Symbolic Logic*, to appear.
- [7] L. Harrington and S. Shelah, Some exact equiconsistency results in set theory, *Notre Dame Journal of Formal Logic* 26 (1985), 178–188.
- [8] P. Hill, New criteria for freeness in abelian groups II, *Trans. Amer. Math. Soc.* **196** (1974), 229–308.
- [9] W. Hodges and S. Shelah, Infinite games and reduced products, *Ann. Math. Logic* 20 (1981), 77–108.
- [10] T. Hyttinen, Games and infinitary languages, *Ann. Acad. Sci. Fenn. Ser. A I Math. Dissertationes*, vol. 64, 1987, pp. 1–32.
- [11] T. Jech, *Set Theory*, Academic Press, 1978.
- [12] T. Jech, M. Magidor, W. Mitchell and K. Prikry, Precipitous ideals, *Journal of Symbolic Logic* 45 (1980), 1–8.
- [13] T. Jech and S. Shelah, Full reflection of stationary sets below \aleph_ω , *Journal of Symbolic Logic* 55 (1990), 822–830.
- [14] R. Jensen, The fine structure of the constructible hierarchy, *Ann. Math. Logic* 4, 229–308.

- [15] M. Nadel and J. Stavi, $L_{\infty\lambda}$ -equivalence, isomorphism and potential isomorphism. Trans. Amer. Math. Soc **236** (1978), 51–74.
- [16] J.-P. Serre, Trees, Springer, 1980.
- [17] S. Shelah, Reflecting stationary sets and successors of singular cardinals, Arch. Math. Logic 30 (1990), 1–29.
- [18] S. Shelah, Classification Theory, revised edition, North-Holland, 1990.
- [19] S. Shelah, Cardinal Arithmetic, in preparation.