

Killing Luzin and Sierpinski sets

by

H. Judah

Department of Mathematics and Computer Science
Bar-Ilan University
Ramat-Gan 52900, Israel

and

S. Shelah*

Institute of Mathematics
Hebrew University of Jerusalem
Givat Ram, Jerusalem
Israel
and
Rutgers University

ABSTRACT. We will kill the old Luzin and Sierpinski sets in order to build a model where $U(\mathcal{M}) = U(\mathcal{N}) = \aleph_1$ and there are neither Luzin nor Sierpinski sets. Thus we answer a question of J. Steprans, communicated by S. Todorcevic on route from Evans to MSRI.

modified:1996-03-11

478 revision:1996-03-11

The authors are partially supported by the Basic Research Foundation, Israel Academy of Science.

*publication number 478

In this note we will build a model where there are non-measurable sets and non-meager sets of size \aleph_1 and there are neither Luzin nor Sierpinski sets. All our notation is standard and can be found in [Ku], [BJ1]. Let us start with the basic concept, underlying this work.

Let $U(\mathcal{M})$ be the minimal cardinal of a non-meager set.

Let $U(\mathcal{N})$ be the minimal cardinal of a non-null set.

We say that a set of reals X is a Luzin set if X is uncountable and $X \cap M$ is countable for every meager set M . We say that a set X is a Sierpinski set if X is uncountable and $X \cap N$ is countable, for every null set N .

Fact. (a) If there is a Luzin set, then $U(\mathcal{M}) = \aleph_1$.

(b) If there is a Sierpinski set then $U(\mathcal{N}) = \aleph_1$. \square

In [Sh] it was proved that if ZF is consistent, then there is a model where there are no Luzin sets and $U(\mathcal{M}) = \aleph_1$. In [BGJS] it was proved that if there is a Sierpinski set, then there is a non-measurable meager filter on ω . It was natural to ask if from $U(\mathcal{N}) = \aleph_1$ we can get such a filter. Clearly it will be enough to answer positively the following question.

(Strepan's) Does $U(\mathcal{N}) = \aleph_1$ imply the existency of a Sierpinski set?

We give a negative answer to this question by proving the following

Theorem. *Cons(ZF) \rightarrow Cons(ZFC + $U(\mathcal{M}) = U(\mathcal{N}) = \aleph_1$ + there are neither Luzin nor Sierpinski sets).*

We will prove this theorem by iterating with countable support iteration Miller reals (rational perfect forcing). We will use the machinery produced by “preservation theorems” to show that the old reals are neither non-meager nor non-measurable sets. We will show that Miller reals kill Luzin and Sierpinski sets from the ground model.

The reader can find a complete analysis of Luzin and Sierpinski sets in [BJ2].

1. *Definition.* Let $P = \{T : T \subseteq \omega^{<\omega} \ \& \ T \text{ is a tree} \ \& \ (\forall s \in T)(s \text{ is increasing}) \ \& \ (\forall s \in T)(\exists t \in T)(\exists^\infty n)(s \subset t \wedge \langle n \rangle \in T)\}$.

Let \leq be defined by $T \leq S$ iff $S \subseteq T$. \square

$\langle P, \leq \rangle$ is called rational perfect forcing ([Mi]) and if $G \subseteq P$ is generic, then $\mathbf{m} = \cap G \in \omega^\omega$ is called a Miller real. From our assumption we have that \mathbf{m} is increasing.

2. *Definition.* Let $r \in \omega^\omega$ be increasing. We define the following set

$$B(r) = \bigcup_{j < \omega} B_j(r)$$

where

$$B_j(r) = \{\eta \in 2^\omega : (\forall i > j) (\eta \upharpoonright [r(i), r(i) + 10(i + 1)]) \text{ is not identically zero}\}$$

3. *Fact.* $\mu(B_j(r)) \geq 1 - \frac{1}{j+1}$ and $B_j(r)$ is closed. \square

Therefore $\mu(B(r)) = 1$.

4. **Lemma.** Let A be a set of reals such that $\mu^*(A) > 0$. Let \mathbf{m} be the canonical name for the Miller real. Then

$$\Vdash_P \text{“} A - B(\mathbf{m}) \text{ is uncountable”}.$$

Proof. Let $p \in P$, $p \Vdash_P \text{“} \tilde{A}' = A \setminus B(\mathbf{m}) \text{ is countable”}$. As P is proper, w.l.o.g. for some countable set $A^* \subseteq A$ and $q \geq p$ we have $q \Vdash_P \text{“} \tilde{A}' \subseteq A^* \text{”}$. Let $N \prec \langle H((2^{\aleph_0})^*), \in \rangle$ be countable, $q \in N$, $A \in N$, $A^* \in N$. As $\mu^*(A) > 0$ there is $\eta \in A$, η random over N . Therefore $\eta \notin A^*$. Let $t \in q$ be such that $mc_q(t) = \{n : t \wedge \langle n \rangle \in q\}$ is infinite. Let us write this set as

$$mc_q(t) = \{k_\ell^t : \ell < \omega\},$$

where $k_\ell^t < k_{\ell+1}^t$. Let $i_t = |t|$. For $n < \omega$, we define

$$E_t^n = \{x \in 2^\omega : (\forall \ell \geq n)(x \upharpoonright [k_\ell^t, k_\ell^t + 10(i_t + 1)]) \text{ is not identically zero}\}.$$

5. *Fact.* $\mu(E_t^n) = 0$. \square

Therefore $E_t = \bigcup_n E_t^n$ is null and $E_t \in N$. Therefore $\eta \notin E_t$. Therefore

$$D_t = \{k_\ell^t : \eta \upharpoonright [k_\ell^t, k_\ell^t + 10(i_t + 1)] \text{ is identically zero}\}$$

is infinite.

Now using this we can define, inductively, $q' \geq q$ satisfying

$$\text{if } t \in q' \text{ and } mc_q(t) \text{ is infinite, then } mc_{q'}(t) = D_t.$$

Therefore $q' \Vdash_P \text{“}\eta \notin B(\mathbf{m})\text{”}$, a contradiction. \square

6. Corollary. *If $Y \in V$ is a Sierpinski set, then Y is not a Sierpinski set in any extension of V containing a Miller real over V .* \square

7. *Remark.* The same result can be obtained if you replace Miller real by Laver real.

8. *Definition.* Let $r \in \omega^\omega$ be increasing. We define the following set

$$T(r) = \bigcup_{j < \omega} [T_j(r)]$$

where $T_j(r)$ is the tree defined by

$$\eta \in [T_j(r)] \text{ iff } \eta \in 2^\omega \ \& \ (\forall i > j)(\eta(r(i)) = 0).$$

We say that for a tree T , $[T]$ is the set of ω -branches of T . \square

9. *Fact.* $[T_j(r)]$ is a closed nowhere dense set. \square

Therefore $T(r)$ is a meager set.

10. Lemma. *Let A be a non-meager set of reals. Let \mathbf{m} be the canonical name for the Miller real. Then*

$$\Vdash_P \text{“}A \cap T(\mathbf{m}) \text{ is uncountable”}.$$

Proof. Let $p \in P$ and let $N \prec (H((2^{\aleph_0})^+), \in)$ be countable such that $p \in N$. Then there is $\eta \in A$ such that η is Cohen over N . We will find q such that

$$p \leq q \in P \text{ and } q \Vdash_P \text{“}\eta \in T(\mathbf{m})\text{”}.$$

Clearly this is enough. Let $\langle \nu_\rho : \rho \in {}^\omega \omega \rangle$ be the list of splitting nodes of p such that $\rho_1 \subsetneq \rho_2$ implies $\nu_{\rho_1} \subsetneq \nu_{\rho_2}$. Thus $\langle \nu_{\rho \wedge \langle n \rangle}(|\nu_\rho|) : n < \omega \rangle$ are distinct and w.l.o.g. are strictly increasing, so $(*) \ \nu_{\rho \wedge \langle n \rangle}(|\nu_\rho|) \geq n$.

For each $\rho \in {}^\omega > \omega$ let

$$A_\rho = \{n < \omega : \eta \upharpoonright (\text{Range } \nu_{\rho \wedge \langle n \rangle} \setminus \text{Range } \nu_\rho) \text{ is identically zero}\}.$$

11. *Fact.* For $\rho \in {}^\omega > \omega$, A_ρ is infinite.

[*Proof.* $p \in N$ and let $s \in 2^{<\omega}$ be a condition in Cohen forcing. Then there is n , by (*), such that

$$\text{dom}(s) \cap (\text{Range } \nu_{\rho \wedge \langle n \rangle} \setminus \text{Range } \nu_\rho) = \emptyset.$$

Thus we can extend s to $t \in 2^{<\omega}$ such that

$$t \upharpoonright (\text{Range } \nu_{\rho \wedge \langle n \rangle} \setminus \text{Range } \nu_\rho)$$

is identically zero. Thus, because η is Cohen over N , we have that A_ρ is infinite.

□

Now we define q by

$$q = \{\nu \in p : (\forall \ell \leq |\nu|)(\nu \upharpoonright \ell = \nu_{\rho \wedge \langle n \rangle} \rightarrow n \in A_\rho)\}$$

and $q \Vdash_P \text{“}\eta \in T(\mathbf{m})\text{”}$. □

12. Corollary. *If $X \in V$ is a Luzin set, then X is not a Luzin set in any extension of V containing a Miller real over V .* □

Now we are ready to show the main Theorem.

13. Theorem. *Cons(ZF) implies Cons(ZFC + $U(\mathcal{M}) = U(\mathcal{N}) = \aleph_1$ + there are neither Luzin nor Sierpinski sets).*

Proof. Let us start with $V = L$. Let P_{ω_2} be the countable support iteration of P , of length ω_2 . Then the following holds in $V^{P_{\omega_2}}$.

- (i) $U(\mathcal{M}) = \aleph_1$: In [Go] it is proved that the property of being non-meager is preserved by a countable support iteration. It is easy to see that P satisfies the covering properties established in [Go], §6.20. Therefore $V \cap 2^\omega$ is a non-meager set in $V^{P_{\omega_2}}$.

- (ii) $U(\mathcal{N}) = \aleph_1$: In [Go] it is proved that the property of being non-null is preserved by countable support iteration. In [BJS] it is proved that P satisfies the covering properties established in [Go] §6.8. Therefore $V \cap 2^\omega$ is a non-null set in $V^{P_{\omega_2}}$.
- (iii) There are no Luzin sets in $V^{P_{\omega_2}}$: by corollary 6.
- (iv) There are no Sierpinski sets in $V^{P_{\omega_2}}$: by corollary 12. \square

14. *Remark.* In the ω_2 -iteration of Laver reals we have that $U(\mathcal{N}) = \aleph_1$ and there are no Sierpinski sets. We don't know if in this model there are uncountable strongly meager sets. We know that Miller reals do not kill strong measure zero sets. This is a consequence of a Rothberger theorem. See [BJ2].

REFERENCES

- [[BGJS]] Bartoszyński, M. Goldstern, H. Judah and S. Shelah, *All meager filters may be null*, accepted by the Proc. of the A.M.S..
- [[BJ1]] Bartoszyński and H. Judah, *Measure and Category: The asymmetry*, in preparation.
- [[BJ2]] ———, *Strong measure zero sets*, to appear in Set Theory of Reals, ed. by H. Judah.
- [[BJS]] Bartoszyński, H. Judah and S. Shelah, *The Cichon diagram*, submitted to the Journal of Symbolic Logic.
- [[Go]] M. Goldstern, *Tools for your forcing constructions*, to appear in Set Theory of Reals, ed. by H. Judah.
- [[Ku]] K. Kunen, *Set Theory - An Introduction to Independence Proofs*, North-Holland, Amsterdam, 1980.
- [[Mi]] A. Miller, *Rational perfect set forcing*, Contemporary Mathematics, edited by J. Baumgartner, D. Martin, S. Shelah.
- [[Sh]] S. Shelah, *Independence results*, Journal of Symbolic Logic **45** (1980).