

# The Bounded Proper Forcing Axiom

Revised version, January 1994

**Martin GOLDSTERN**<sup>1</sup>

Free University of Berlin

**Saharon SHELAH**<sup>1,2</sup>

Hebrew University of Jerusalem

ABSTRACT. The bounded proper forcing axiom BPFA is the statement that for any family of  $\aleph_1$  many maximal antichains of a proper forcing notion, each of size  $\aleph_1$ , there is a directed set meeting all these antichains.

A regular cardinal  $\kappa$  is called  $\Sigma_1$ -reflecting, if for any regular cardinal  $\chi$ , for all formulas  $\varphi$ , “ $H(\chi) \models \varphi$ ” implies “ $\exists \delta < \kappa$ ,  $H(\delta) \models \varphi$ ”

We show that BPFA is equivalent to the statement that two nonisomorphic models of size  $\aleph_1$  cannot be made isomorphic by a proper forcing notion, and we show that the consistency strength of the bounded proper forcing axiom is exactly the existence of a  $\Sigma_1$ -reflecting cardinal (which is less than the existence of a Mahlo cardinal).

We also show that the question of the existence of isomorphisms between two structures can be reduced to the question of rigidity of a structure.

---

<sup>1</sup> The authors thank the DFG (grant Ko 490/7-1) and the Edmund Landau Center for research in Mathematical Analysis, supported by the Minerva Foundation (Germany)

<sup>2</sup> Publication 507

## Introduction

The proper forcing axiom has been successfully employed to decide many questions in set-theoretic topology and infinite combinatorics. See [Ba1] for some applications, and [Sh b] and [Sh f] for variants.

In the recent paper [Fu], Fuchino investigated the following two consequences of the proper forcing axiom:

- (a) If a structure  $\mathfrak{A}$  of size  $\aleph_1$  cannot be embedded into a structure  $\mathfrak{B}$ , then such an embedding cannot be produced by a proper forcing notion.
- (b) If two structures  $\mathfrak{A}$  and  $\mathfrak{B}$  are not isomorphic, then they cannot be made isomorphic by a proper forcing notion.

He showed that (a) is in fact equivalent to the proper forcing axiom, and asked if the same is true for (b).

In this paper we find a natural weakening of the proper forcing axiom, the “bounded” proper forcing axiom and show that it is equivalent to property (b) above.

We then investigate the consistency strength of this new axiom. While the exact consistency strength of the proper forcing axiom is still unknown (but large, see [To]), it turns out that the bounded proper forcing axiom is equiconsistent to a rather small large cardinal.

For notational simplicity we will, for the moment, only consider forcing notions which are complete Boolean algebras. See 0.4 and 4.6.

We begin by recalling the forcing axiom in its usual form: For a forcing notion  $P$ ,  $\text{FA}(P, \kappa)$  is the following statement:

Whenever  $\langle A_i : i < \kappa \rangle$  is a family of maximal antichains of  $P$ , then there is a filter  $G^* \subseteq P$  meeting all  $A_i$ .

If  $\underline{f}$  is a  $P$ -name for a function from  $\kappa$  to the ordinals, we will say that  $G^* \subseteq P$  decides  $\underline{f}$  if for each  $i < \kappa$  there is a condition  $p \in G^*$  and an ordinal  $\alpha_i$  such that  $p \Vdash \underline{f}(i) = \alpha_i$ . (If  $G^*$  is directed, then this ordinal must be unique, and we will write  $\underline{f}[G^*]$  for the function  $i \mapsto \alpha_i$ .) Now it is easy to see that the  $\text{FA}(P, \kappa)$  is equivalent to the following statement:

Whenever  $\underline{f}$  is a  $P$ -name for a function from  $\kappa$  to the ordinals, then there is a filter  $G^* \subseteq P$  which decides  $\underline{f}$ .

This characterization suggests the following weakening of the forcing axiom:

**0.1 Definition:** Let  $P$  be a forcing notion, and let  $\kappa$  and  $\lambda$  be infinite cardinals.

$\text{BFA}(P, \kappa, \lambda)$  is the following statement: Whenever  $\underline{f}$  is a  $P$ -name for a function from  $\kappa$  to  $\lambda$  then there is a filter  $G^* \subseteq P$  which decides  $\underline{f}$ , or equivalently:

Whenever  $\langle A_i : i < \kappa \rangle$  is a family of maximal antichains of  $P$ , each of size  $\leq \lambda$ , then there is a filter  $G \subseteq P$  which meets all  $A_i$ .

**0.2 Notation:**

- (1)  $\text{BFA}(P, \lambda)$  is  $\text{BFA}(P, \lambda, \lambda)$ , and  $\text{BFA}(P)$  is  $\text{BFA}(P, \omega_1)$ .
- (2) If  $\mathcal{E}$  is a class or property of forcing notions, we write  $\text{BFA}(\mathcal{E})$  for  $\forall P \in \mathcal{E} \text{ BFA}(P)$ , etc.
- (3)  $\text{BPFA}$  = the bounded proper forcing axiom =  $\text{BFA}(\text{proper})$ .

**0.3 Remark:** For the class of ccc forcing notions we get nothing new:  $\text{BFA}(\text{ccc}, \lambda)$  is equivalent to Martin's axiom  $\text{MA}(\lambda)$ , i.e.,  $\text{FA}(\text{ccc}, \lambda)$ . ☺ 0.3

**0.4 Remark:** If the forcing notion  $P$  is not a complete Boolean algebra but an arbitrary

poset, then it is possible that  $P$  does not have any small antichains, so it could satisfy the second version of  $\text{BFA}(P)$  vacuously. The problem with the first definition, when applied to an arbitrary poset, is that a filter on  $\text{ro}(P)$  which interprets the  $P$ -name ( $=\text{ro}(P)$ -name)  $\underline{f}$  does not necessarily generate a filter on  $P$ . So for the moment our official definition of  $\text{BFA}(P)$  for arbitrary posets  $P$  will be

$$\text{BFA}(P) : \iff \text{BFA}(\text{ro}(P))$$

In 4.4 and 4.5 we will find a equivalent (and more natural?) definition  $\text{BFA}'(P)$  which does not explicitly refer to  $\text{ro}(P)$ .

Contents of the paper: in section 1 we show that the “bounded forcing axiom” for any forcing notion  $P$  is equivalent to Fuchino’s “potential isomorphism” axiom for  $P$ . In section 2 we define the concept of a  $\Sigma_1$ -reflecting cardinal, and we show that from a model with such a cardinal we can produce a model for the bounded proper forcing axiom. In section 3 we describe a (known) forcing notion which we will use in section 4, where we complement our consistency result by showing that a  $\Sigma_1$ -reflecting cardinal is necessary: If BPFA holds, then  $\aleph_2$  must be  $\Sigma_1$ -reflecting in  $L$ .

Notation: We use  $\odot$  to denote the end of a proof, and we write  $\odot$  when we leave a proof to the reader.

We will use gothic letters  $\mathfrak{A}, \mathfrak{B}, \mathfrak{M}, \dots$  for structures ( $=$ models of a first order language), and the corresponding latin letters  $A, B, M, \dots$  for the underlying universes. Thus, a model  $\mathfrak{A}$  will have the universe  $A$ , and if  $A' \subseteq A$  then we let  $\mathfrak{A}'$  be the submodel (possibly with partial functions) with universe  $A'$ , etc.

## 1. Fuchino’s problem and other applications

Let  $\mathcal{E}$  be a class of forcing notions.

**1.1 Definition:** Let  $\mathfrak{A}$  and  $\mathfrak{B}$  be two structures for the same first order language, and let  $\mathcal{E}$  be a class (or property) of forcing notions. We say that  $\mathfrak{A}$  and  $\mathfrak{B}$  are  $\mathcal{E}$ -potentially isomorphic ( $\mathfrak{A} \simeq_{\mathcal{E}} \mathfrak{B}$ ) iff there is a forcing  $P \in \mathcal{E}$  such that  $\Vdash_P \text{“}\mathfrak{A} \simeq \mathfrak{B} \text{”}$ .  $\mathfrak{A} \simeq_P \mathfrak{B}$  means  $\mathfrak{A} \simeq_{\{P\}} \mathfrak{B}$ .

**1.2 Definition:** We say that a structure  $\mathfrak{A}$  is nonrigid, if it admits a nontrivial automorphism. We say that  $\mathfrak{A}$  is  $\mathcal{E}$ -potentially nonrigid, if there is a forcing notion  $P \in \mathcal{E}$ ,  $\Vdash_P \text{“}\mathfrak{A} \text{ is nonrigid”}$ .

### 1.3 Definition:

- (1)  $\text{PI}(\mathcal{E}, \lambda)$  is the statement: Any two  $\mathcal{E}$ -potentially isomorphic structures of size  $\lambda$  are isomorphic.
- (2)  $\text{PA}(\mathcal{E}, \lambda)$  is the statement: Any  $\mathcal{E}$ -potentially nonrigid structure of size  $\lambda$  is nonrigid.

$\text{PI}(\mathcal{E}, \lambda)$  was defined by Fuchino [Fu]. It is clear that

$$\text{FA}(\mathcal{E}, \lambda) \implies \text{BFA}(\mathcal{E}, \lambda) \implies \text{PI}(\mathcal{E}, \lambda) \& \text{PA}(\mathcal{E}, \lambda)$$

for all  $\mathcal{E}$ , and Fuchino asked if  $\text{PI}(\mathcal{E}, \lambda)$  implies  $\text{FA}(\mathcal{E}, \lambda)$ , in particular for the cases  $\mathcal{E}=\text{ccc}$ ,  $\mathcal{E}=\text{proper}$  and  $\mathcal{E}=\text{stationary-preserving}$ .

We will show in this section that the the three statments BFA, PA and PI are in fact equivalent. Hence in particular  $\text{PI}(\text{ccc}, \lambda)$  is equivalent to  $\text{MA}(\lambda)$ .

In the next sections we will show that for  $\mathcal{E}=\text{proper}$ , the first implication cannot be re-

versed, by computing the exact consistency strength of BPFA and comparing it to the known lower bounds for the consistency strength of PFA.

**1.4 Theorem:** For any forcing notion  $P$  and for any  $\lambda$ , the following are equivalent:

PI( $P, \lambda$ )

PA( $P, \lambda$ )

BFA( $P, \lambda$ )

Proof of PI  $\Rightarrow$  PA: This follows from theorem 1.13 below. Here we will give a shorter proof under the additional assumption that we have not only PI( $P$ ) but also PI( $P_p$ ) for all  $p \in P$ , where  $P_p$  is the set of all elements of  $P$  which are stronger than  $p$ :

Let  $\mathfrak{M}$  be a potentially nonrigid structure. So there is a  $P$ -name  $\underline{f}$  such that

$$\Vdash_P \text{“}\underline{f} \text{ is a nontrivial automorphism of } \mathfrak{M}\text{”}$$

We can find a condition  $p \in P$  and two elements  $a \neq b$  of  $\mathfrak{M}$  such that

$$p \Vdash_P \text{“}\underline{f}(a) = b\text{”}$$

Since we can replace  $P$  by  $P_p$ , we may assume that  $p$  is the weakest condition of  $P$ . So we have that  $(\mathfrak{M}, a)$  and  $(\mathfrak{M}, b)$  are potentially isomorphic. Any isomorphism from  $(\mathfrak{M}, a)$  to  $(\mathfrak{M}, b)$  is an automorphism of  $\mathfrak{M}$  mapping  $a$  to  $b$ , so we are done. ☺  $PI \Rightarrow PA$

We will now describe the framework of the proof of the second part of our theorem: PA  $\Rightarrow$  BFA. We start with a forcing notion  $P$ . Recall that (for the moment) all our forcing notions are complete Boolean algebras. Fix a small family of small antichains. Our structure will consist of a disjoint union of the free groups generated by the antichains. On each of the free groups the translation by an element of the corresponding antichain will be a nontrivial

automorphism, and if all these translating elements are selected from the antichains by a directed set, then the union of these automorphisms will be an automorphism of the whole structure. We will also ensure that “essentially” these are the only automorphisms, so every automorphism will define a sufficiently generic set.

**1.5 Definition:** For any set  $X$  let  $F(X)$  be the free group on the generators  $X$ , and for  $w \in F(X)$  define  $\text{supp}(w) = \bigcap \{Y \subseteq X : w \in \langle Y \rangle\}$ , i.e.  $\text{supp}(w)$  is the set of elements  $x$  of  $X$  which occur (as  $x$  or as  $x^{-1}$ ) in the reduced representation of  $w$ . (If you prefer, you can change the proof below by using the free abelian group generated by  $X$  instead of the free group, or the free abelian group of order 2, ...).

**1.6 Setup:** Let  $P$  be a complete Boolean algebra, and let  $(A_i : i \in I)$  be a system of  $\lambda$  many maximal antichains of size  $\lambda$ . We may assume that this is a directed system, i.e., for any  $i, j \in I$  there is a  $k \in I$  such that  $A_k$  refines both  $A_i$  and  $A_j$ . So if we write  $i < j$  for “ $A_j$  refines  $A_i$ ”,  $(I, <)$  becomes a partially ordered upwards directed set. (We say that  $A$  refines  $B$  if each element of  $A$  is stronger than some unique element of  $B$ , or in the Boolean sense if there is a partition  $A = \dot{\bigcup}_{b \in B} A_b$  of the set  $A$  satisfying  $\forall b \in B \sum_{a \in A_b} a = b$ .)

Assuming  $\text{PA}(P, \lambda)$ , we will find a filter(base) meeting all the sets  $A_i$ .

**1.7 Definition:**

- (a) Let  $(F_i, *)$  be the free group generated by  $A_i$ , and let  $M$  be the disjoint union of the sets  $F_i$ .
- (b) For  $i \in I, z \in F_i$  let

$$R_{i,z} = \{(y, z * y) : y \in F_i\}$$

- (c) If  $i < j$ , then there is a “projection” function  $h_i^j$  from  $A_j$  to  $A_i$ : For  $p \in A_j$ ,

$h_i^j(p)$  is the unique element of  $A_i$  which is compatible with (and in fact weaker than)  $p$ .  $h_i^j$  extends to a unique homomorphism from  $F_j$  to  $F_i$ , which we will also call  $h_i^j$ .

**1.8 Fact:** (1) The functions  $h_i^j$  commute, i.e., if  $i < j < k$  then  $h_i^k = h_i^j \circ h_j^k$ .

(2) If  $i < j$ ,  $p \in A_j$ , then  $p$  is stronger than  $h_i^j(p)$ . ☺ 1.8

Now let  $\mathfrak{M} = (M, (F_i)_{i \in I}, (R_{i,z})_{i \in I, z \in F_i}, (h_i^j)_{i \in I, j \in I, i < j})$ , where we treat all sets  $F_i$ ,  $R_{i,z}$ ,  $h_i^j$  as relations on  $M$ .

**1.9 Definition:** Let  $G \subseteq P$  be a filter which meets all the sets  $A_i$ , say  $G \cap A_i = \{y_i(G)\}$ . Define  $f_G : M \rightarrow M$  as follows: If  $x \in F_i$ , then  $f_G(x) = x * y_i(G)$  (here  $*$  =  $*_i$  is the group operation on  $F_i$ ).

**1.10 Fact:** If  $G$  is a filter which meets all sets  $A_i$ , then  $f_G$  is an automorphism of  $\mathfrak{M}$  without fixpoints.

Proof: It is clear that the sets  $F_i$  and the relations  $R_{i,z}$  are preserved. Note that for  $i < j$  we have  $h_i^j(y_j) = y_i$ , since  $y_i$  and  $y_j$  are compatible. Since the functions  $h_i^j$  are homomorphisms, we have  $h_i^j(f_G(x)) = h_i^j(x * y_j) = h_i^j(x) * y_i = f_G(h_i^j(x))$ , so also  $h_i^j$  is preserved. ☺ 1.10

So  $\mathfrak{M}$  is potentially nonrigid. So by PA( $P, \lambda$ ) we know that  $\mathfrak{M}$  is really nonrigid.

Finally we will show how a nontrivial automorphism of  $\mathfrak{M}$  defines a filter  $G^*$  meeting all the sets  $A_i$ .

So let  $F$  be an automorphism. Let  $1_i$  be the neutral element of  $F_i$ , and assume  $F(1_i) = w_i$ .

Since the sets  $F_i$  are predicates in our structure, we must have  $w_i \in F_i$ . Using the predicates



$h_i^j$  we can show: If  $i < j$ , then  $h_i^j(w_j) = w_i$ , and using the predicates  $R_{i,z}$  we can show that for all  $z \in F_i$  we must have  $F(z) = z * w_i$ .

Therefore, as  $F$  is not the identity, we can find  $i^* \in I$  such that  $w_{i^*} \neq 1_{i^*}$ . From now on we will work only with  $I^* = \{i \in I : i^* \leq i\}$ . Since every antichain  $A_i$  is refined by some antichain  $A_j$  with  $j \in I^*$  it is enough to find a directed set which meets all antichains  $A_j$  for  $j \in I^*$ .

Let  $u_i = \text{supp}(w_i)$ . So for all  $i \in I^*$  the set  $u_i$  is finite and nonempty (since  $h_{i^*}^i[u_i] \supseteq u_{i^*} \neq \emptyset$ ).

### 1.11 Fact:

- (1) If  $J \subseteq I^*$  is a finite set, then there is a family  $\{p_i : i \in J\}$  such that
  - (a) for all  $i \in J$ :  $p_i \in u_i$
  - (b) for all  $i, j \in J$ : If  $i < j$ , then  $h_i^j(p_j) = p_i$ .
- (2) There is a family  $\{p_i : i \in I^*\}$  such that
  - (a) for all  $i \in I^*$ :  $p_i \in u_i$
  - (b) for all  $i, j \in I^*$ : If  $i < j$ , then  $h_i^j(p_j) = p_i$ .
- (3) If  $\{p_i : i \in I^*\}$  is as in (2), then this set generates a filter which will meet all sets  $A_i$ .

Proof of (1): As  $I^*$  is directed, we can find an upper bound  $j$  for  $J$ . Let  $p$  be an element of  $w_j$  such that  $p_i := h_i^j(p_j) \in w_i$  for all  $i \in J$ .

(2) follows from (1), by the compactness theorem of propositional calculus. (Recall that all sets  $u_i$  are finite.)

(3): We have to show that for any  $i_1, i_2 \in I^*$  the conditions  $p_{i_1}$  and  $p_{i_2}$  are compatible,

i.e., have a common extension. Let  $j$  be an upper bound of  $i_1$  and  $i_2$ . Then  $p_j$  witnesses that  $p_{i_1}$  and  $p_{i_2}$  are compatible, as  $h_{i_1}^j(p_j) = p_{i_1}$  and  $h_{i_2}^j(p_j) = p_{i_2}$ . ☺ 1.11 ☺ 1.4.

For the theorem 1.13 below we need the following definitions.

**1.12 Definition:** A tree on a set  $X$  is a nonempty set  $T$  of finite sequences of elements of  $X$  which is closed under restrictions, i.e., if  $\eta : k \rightarrow X$  is in  $T$  and  $i < k$ , then also  $\eta \upharpoonright i \in T$ .

The tree ordering  $\leq_T$  is given by the subset (or extension) relation:  $\eta \leq \nu$  iff  $\eta \subseteq \nu$  iff  $\exists i : \eta = \nu \upharpoonright i$ .

For  $\eta \in T$  let  $\text{Suc}_T(\eta) := \{x \in X : \eta \frown x \in T\}$ .

For  $A \subseteq T$ ,  $\eta \in T$  we let  $\text{rk}(\eta, A)$  be the rank of  $\eta$  with respect to  $A$ , i.e., the rank of the (inverse) tree ordering on the set

$$\{\nu : \eta \leq \nu \in T, \forall \nu' : \eta \leq \nu' < \nu \Rightarrow \nu' \notin A\}$$

In other words,  $\text{rk}(\eta, A) = 0$  iff  $\eta \in A$ ,  $\text{rk}(\eta, A) = \infty$  iff there is an infinite branch of  $T$  starting at  $\eta$  which avoids  $A$ , and  $\text{rk}(\eta, A) = \sup\{\text{rk}(\nu, A) + 1 : \nu \text{ a direct successor of } \eta\}$  otherwise.

**1.13 Theorem:** For any two structures  $\mathfrak{A}$  and  $\mathfrak{B}$  there is a structure  $\mathfrak{C} = \mathfrak{C}(\mathfrak{A}, \mathfrak{B})$  such that in any extension  $V' \supseteq V$  of the universe,  $V' \models \text{“}\mathfrak{A} \simeq \mathfrak{B} \leftrightarrow \mathfrak{C} \text{ is not rigid.}”$

Proof: Wlog  $|A| \leq |B|$ . Also wlog  $\mathfrak{A}$  and  $\mathfrak{B}$  are structures in a purely relational language  $\mathcal{L}$ , and we may also assume that  $A \cap B = \emptyset$ .

We will say that a tree  $T$  on  $A \cup B$  “codes  $A$ ” iff

- (1)  $\text{Suc}_T(\eta) \in \{A, B\}$  for all  $\eta \in T$ .
- (2) Letting  $T^A := \{\eta \in T : \text{Suc}_T(\eta) = A\}$ , the ranks  $\text{rk}(\eta, T^A)$  are  $< \infty$  for all

$\eta \in T$ .

(3) The function  $\eta \mapsto \text{rk}(\eta, T^A \setminus \{\eta\})$  is 1-1 on  $T^A$ .

Such a tree can be constructed inductively as  $T = \bigcup_n T_n$ , where the  $T_n$  are well-founded trees, each  $T_{n+1}$  end-extends  $T_n$ , and all nodes in  $T_{n+1} - T_n$  are from  $B$  except those at the top (i.e., those whose immediate successors will be in  $T_{n+2} - T_{n+1}$ ). Because we have complete freedom in what the rank of the tree ordering for each connected component of  $T_{n+1} - T_n$  should be (and because all the  $T_n$  have size  $= |B|$ ), we can arrange to satisfy (1), (2) and (3).

Moreover, we can find trees  $T_0$  and  $T_1$ , both coding  $A$ , such that

(4)  $\text{Suc}_{T_0}(\emptyset) = A$ ,  $\text{Suc}_{T_1}(\emptyset) = B$ .

We will replace the roots ( $\emptyset$ ) of the trees  $T_0$  and  $T_1$  by some new and distinct objects  $\emptyset_0$  and  $\emptyset_1$ . So the trees  $T_0$  and  $T_1$  will be disjoint (by (4)).

Now define the structure  $\mathfrak{C}$  as follows: We let  $C = T_0 \cup T_1$ .

The underlying language of  $\mathfrak{C}$  will be the language  $\mathcal{L}$  plus an additional binary relation symbol  $\leq$ , which is to be interpreted as the tree order. Whenever  $R$  is an  $n$ -ary relation in the language  $\mathcal{L}$ , we interpret  $R$  in  $\mathfrak{C}$  by

$$R^{\mathfrak{C}} := \{(\eta \hat{\ } a_1, \dots, \eta \hat{\ } a_n) : \eta \in T_0 \cup T_1, \text{ and} \\ \text{Suc}(\eta) = A \Rightarrow (a_1, \dots, a_n) \in R^{\mathfrak{A}}, \\ \text{Suc}(\eta) = B \Rightarrow (a_1, \dots, a_n) \in R^{\mathfrak{B}} \}$$

Now work in any extension  $V' \supseteq V$ . First assume that  $f : \mathfrak{A} \rightarrow \mathfrak{B}$  is an isomorphism. We will define a map  $g : T_0 \rightarrow T_1$  such that the map  $g \cup g^{-1}$  is a (nontrivial) automorphism of  $\mathfrak{C}$ .

$g$  is defined inductively as follows:

- (a)  $g(\emptyset_0) = \emptyset_1$ .
- (b) If  $\text{Suc}_{T_0}(\eta) = \text{Suc}_{T_1}(g(\eta))$ , then  $g(\eta \frown a) = g(\eta) \frown a$ .
- (c) Otherwise,  $g(\eta \frown a) = g(\eta) \frown f(a)$  or  $g(\eta \frown a) = g(\eta) \frown f^{-1}(a)$ , as appropriate.

It is easy to see that  $g \cup g^{-1}$  will then be a nontrivial automorphism.

Now assume conversely that  $g : \mathfrak{C} \rightarrow \mathfrak{C}$  is a nontrivial automorphism. Recall that the tree ordering is a relation on the structure  $\mathfrak{C}$ , so it must be respected by  $g$ .

First assume that there are  $i, j \in \{0, 1\}$  and an  $\eta$  such that

$$(*) \quad \eta \in T_i \quad g(\eta) \in T_j \quad \text{Suc}_{T_i}(\eta) \neq \text{Suc}_{T_j}(g(\eta))$$

So without loss of generality  $\text{Suc}_{T_i}(\eta) = A$  and  $\text{Suc}_{T_j}(g(\eta)) = B$ . Now define a map  $f : A \rightarrow B$  by requiring

$$g(\eta \frown a) = g(\eta) \frown f(a)$$

and check that  $f$  must be an isomorphism.

Now we show that we can always find  $i, j, \eta$  as in  $(*)$ . If not, then we can first see that  $g$  respects  $T_0$  and  $T_1$ , i.e.,  $g(\eta) \in T_0$  iff  $\eta \in T_0$ . Next, our assumption implies that the functions  $g \upharpoonright T_0$  respect the sets  $T_0^A$ , i.e.,  $\eta \in T_0^A$  iff  $g(\eta) \in T_0^A$ . Hence for all  $\eta \in T_0^A$ ,  $\text{rk}(\eta, T_0^A) = \text{rk}(g(\eta), T_0^A)$ , so (by condition (3) above)  $g(\eta) = \eta$  for all  $\eta \in T_0^A$ . Since every  $\nu \in T_0$  can be extended to some  $\eta \in T_0^A$  and  $g$  respects  $<$ , we must have  $g(\nu) = \nu$  for all  $\nu \in T_0$ . The same argument shows that also  $g \upharpoonright T_1$  is the identity. ☺ 1.13

**1.14 Remarks on other applications:** Which other consequences of PFA (see, e.g., [Ba1]) are already implied by BPFA? On the one hand it is clear that if PFA is only

needed to produce a sufficiently generic function from  $\omega_1$  to  $\omega_1$ , then the same proof should show that BPFA is a sufficient assumption. For example:

BPFA implies “all  $\aleph_1$ -dense sets of reals are isomorphic.”

On the other hand, as we will see in the next section, the consistency strength of BPFA is quite weak. So BPFA cannot imply any statement which needs large cardinals, such as “there is an Aronszajn tree on  $\aleph_2$ .” In particular, BPFA does not imply PFA.

We do not know if BPFA already decides the size of the continuum, but Woodin has remarked that the bounded **semiproper** forcing axiom implies  $2^{\aleph_0} = \aleph_2$ .

## 2. The consistency of BPFA

**2.1 Definition:** For any cardinal  $\chi$ ,  $H(\chi)$  is the collection of sets which are hereditarily of cardinality  $< \chi$ : Letting  $trcl(x)$  be the transitive closure of  $x$ ,  $trcl(x) = \{x\} \cup \bigcup x \cup \bigcup \bigcup x \cup \dots$ , we have

$$H(\chi) = \{x : |trcl(x)| < \chi\}$$

(Usually we require  $\chi$  to be regular)

**2.2 Definition:** Let  $\kappa$  be an regular cardinal. We say that  $\kappa$  is “reflecting” or more precisely,  $\Sigma_1$ -reflecting, if:

For any first order formula  $\varphi$  in the language of set theory, for any  $a \in H(\kappa)$ :

IF there exists a regular cardinal  $\chi \geq \kappa$  such that  $H(\chi) \models \varphi(a)$

THEN there is a cardinal  $\delta < \kappa$  such that  $a \in H(\delta)$  and  $H(\delta) \models \varphi(a)$ .

**2.3 Remark:** (1) We may require  $\delta$  to be regular without changing the concept of “ $\Sigma_1$ -reflecting”.

(2) We can replace “for all  $\chi$ ” by “for unboundedly many  $\chi$ ”

Proof: (1) Assume that  $H(\chi) \models \varphi(a)$ ,  $\chi$  regular. Choose some large enough  $\chi_1$  such that  $H(\chi) \in H(\chi_1)$ ,  $\chi_1$  a successor cardinal. So  $H(\chi_1) \models “\exists \chi, \chi \text{ regular, } H(\chi) \text{ exists and } H(\chi) \models ‘\varphi(a)’”$ . We can find a (successor)  $\delta_1 < \kappa$  such that  $H(\delta_1) \models “\exists \delta, \delta \text{ regular, } H(\delta) \models ‘\varphi(a)’”$  So  $\delta$  is really regular.

(2): If  $\chi < \chi_1$  then  $H(\chi) \models “\varphi”$  iff  $H(\chi_1) \models “H(\chi) \models ‘\varphi’”$ . ☺ 2.3

**2.4 Remark:** It is easy to see that if  $\kappa$  is reflecting, then  $\kappa$  is a strong limit, hence inaccessible. Applying  $\Sigma_1$  reflection, we get that  $\kappa$  is hyperinaccessible, etc. ☺ 2.4

**2.5 Remark:** (1) There is a closed unbounded class  $C$  of cardinals such that every regular  $\kappa \in C$  (if there are any) is  $\Sigma_1$  reflecting. So if “ $\infty$  is Mahlo”, then there are many  $\Sigma_1$ -reflecting cardinals.

(2) If  $\kappa$  is reflecting, then  $L \models “\kappa \text{ is reflecting}”$ .

Proof: (1) For any set  $a$  and any formula  $\varphi$  let  $f'(a, \varphi) = \min\{\chi \in RCard : H(\chi) \models \varphi(a)\}$  (where  $RCard$  is the class of regular cardinals, and we define  $\min \emptyset = 0$ ). Now let  $f : RCard \rightarrow RCard$  be defined by  $f(\alpha) = \sup\{f'(a, \varphi) : \varphi \text{ a formula, } a \in H(\alpha)\}$ , and let  $C = \{\delta \in Card : \forall \alpha \in RCard \cap \delta \ f(\alpha) < \delta\}$ .

(2) is also easy. ☺ 2.5

Our main interest in this concept is its relativization to  $L$ . In this context we recall the following fact:

**2.6 Fact:** Assume  $V = L$ . Then for all (regular) cardinals  $\chi$ ,  $H(\chi) = L_\chi$ . ☺ 2.6

**2.7 Fact:** Assume  $P \in H(\lambda)$  is a forcing notion,  $\chi > 2^{2^\lambda}$  is regular. Then

- (1) For any  $P$ -name  $\underline{x}$  there is a  $P$ -name  $\underline{y} \in H(\chi)$  such that  $\Vdash_P \text{“}\underline{x} \in H(\chi) \Rightarrow \underline{x} = \underline{y}\text{”}$ . (And conversely, if  $\underline{x} \in H(\chi)$ , then  $\Vdash_P \text{“}\underline{x} \in H(\chi)\text{”}$ .)
- (2) If  $\underline{x} \in H(\chi)$ ,  $\varphi(\cdot)$  a formula, then

$$\Vdash \text{“}H(\chi) \models \varphi(\underline{x})\text{”} \Leftrightarrow \text{“}H(\chi) \models \text{‘}\Vdash \varphi(\underline{x})\text{’”}$$

Proof: (1) is by induction on the rank of  $\underline{x}$  in  $V^P$ , and (2) uses (1). ☺ 2.7

**2.8 Fact:** Let  $P$  be a forcing notion,  $P \in H(\lambda)$ ,  $\chi > 2^{2^\lambda}$  regular. Then  $P$  is proper iff  $H(\chi) \models \text{“}P \text{ is proper”}$ . ☺ 2.8

**2.9 Lemma:** Assume that  $\kappa$  is reflecting,  $\lambda < \kappa$  is a regular cardinal,  $\mathfrak{A}$  and  $\mathfrak{B}$  are structures in  $H(\lambda)$ .

If there is a proper forcing notion  $P$  such that  $\Vdash_P \text{“}\mathfrak{A} \simeq \mathfrak{B}\text{”}$ , then there is such a (proper) forcing notion in  $H(\kappa)$ .

Proof: Fix  $P$ , and let  $\chi$  be a large enough regular cardinal. So  $H(\chi) \models \text{“}P \text{ proper, } P \in H(\mu), (2^{2^\mu}) \text{ exists”}$ . Also, there is a  $P$ -name  $\underline{f} \in H(\chi)$  such that  $\Vdash_P \text{“}H(\chi) \models \text{‘}\underline{f} : \mathfrak{A} \rightarrow \mathfrak{B} \text{ is an isomorphism’”}$ , so by 2.7(2),  $H(\chi) \models \text{“}\Vdash_P \text{‘}\underline{f} : \mathfrak{A} \rightarrow \mathfrak{B} \text{ is an isomorphism’”}$ .

Now we use the fact that  $\kappa$  is reflecting. We can find  $\delta < \kappa$ ,  $\delta > \lambda$  and  $\chi' \in H(\delta)$  such that  $H(\delta) \models \text{“}\exists \nu \exists Q \in H(\nu), Q \text{ proper, } \exists \underline{g} \Vdash_Q \text{‘}\underline{g} : \mathfrak{A} \rightarrow \mathfrak{B} \text{ is an isomorphism’}$ , and  $(2^{2^\nu})$  exists.” So this  $Q$  is really proper, and  $Q$  forces that  $\mathfrak{A}$  and  $\mathfrak{B}$  are isomorphic. ☺ 2.9

**2.10 Fact:** If  $\kappa$  is reflecting,  $P \in H(\kappa)$  is a forcing notion, then  $\Vdash_P \text{“}\kappa \text{ is reflecting”}$ .

Proof: Let  $P \in H(\lambda)$ ,  $\lambda < \kappa$ . Assume that  $p \Vdash \text{“}H(\chi) \models \text{‘}\varphi(\underline{a})\text{’, } \underline{a} \in H(\kappa)\text{”}$ . We may assume that  $\underline{a} \in H(\kappa)$ . By 2.7 we have  $H(\chi) \models \text{“}p \Vdash \text{‘}\varphi(\underline{a})\text{’”}$ , so there is a  $\delta < \kappa$ ,  $\delta > \lambda$ , such that  $H(\delta) \models \text{“}p \Vdash \text{‘}\varphi(\underline{a})\text{’”}$ , hence  $p \Vdash \text{“}H(\delta) \models \text{‘}\varphi(\underline{a})\text{’”}$ .  $\delta$  is a cardinal in  $V^P$ ,

because  $|P| < \lambda < \delta$ .

☺ 2.10

**2.11 Theorem:** If “there is a reflecting cardinal” is consistent with ZFC, then also PI(proper) (and hence BPFA, by 1.4) is consistent with ZFC.

Proof: (Short version) We will use an CS iteration of length  $\kappa$ , where  $\kappa$  reflects. All intermediate forcing notions will have hereditary size  $< \kappa$ . By a bookkeeping argument we can take care of all possible pairs of structures on  $\omega_1$ . If in the intermediate model there is a proper forcing notion making two structures isomorphic, then there is such a forcing notion of size  $< \kappa$ , so we continue. Note that once two structures have been made isomorphic, they continue to stay isomorphic.

Proof: (More detailed version) Assume that  $\kappa$  reflects. We define a countable support iteration  $(P_i, Q_i : i < \kappa)$  of proper forcing notions and a sequence  $\langle \mathfrak{M}_i, \mathfrak{N}_i : i < \kappa \rangle$  with the following properties for all  $i < \kappa$ :

- (1)  $P_i \in H(\kappa)$
- (2)  $Q_i$  is a  $P_i$ -name,  $\Vdash_{P_i}$  “ $Q_i$  is proper,  $Q_i \in H(\kappa)$ ”.
- (3)  $\Vdash_{P_i} 2^{\aleph_1} < \kappa$ . (This follows from (1) and (2))
- (4)  $\mathfrak{M}_i$  and  $\mathfrak{N}_i$  are names for structures on  $\omega_1$ .
- (5)  $\Vdash_{P_i}$  “If  $\mathfrak{M}_i \underset{\text{proper, } < \kappa}{\simeq} \mathfrak{N}_i$ , then  $\Vdash_{Q_i}$  ‘ $\mathfrak{M}_i \simeq \mathfrak{N}_i$ ’”.

With the usual bookkeeping argument we can also ensure that

- (6) Whenever  $\mathfrak{M}$  and  $\mathfrak{N}$  are  $P_i$ -names for structures on  $\omega_1$  for some  $i$ , then there are unboundedly (or even stationarily) many  $j > i$  with  $\Vdash_j$  “ $\mathfrak{M}_j = \mathfrak{M}$ ,  $\mathfrak{N}_j = \mathfrak{N}$ ”

From (1) we also get the following two properties:

- (7)  $P_\kappa \models \kappa\text{-cc}$



- (8) Whenever  $\mathfrak{M}$  is a  $P_\kappa$ -name for a structure on  $\omega_1$ , then there are  $i < \kappa$  and a  $P_i$ -name  $\mathfrak{M}'$  such that  $\Vdash_{-\kappa} \mathfrak{M} = \mathfrak{M}'$ .

From these properties we can now show  $\Vdash_{-\kappa}$  BPFA.  $P_\kappa$  is proper, so  $\omega_1$  is not collapsed. Let  $p$  be a condition, and let  $\mathfrak{M}$  and  $\mathfrak{N}$  be  $P_\kappa$ -names for structures on  $\omega_1$ , and assume that

$$p \Vdash_{-\kappa} \text{“}\underline{Q} \text{ proper, } \Vdash_{-\underline{Q}} \mathfrak{M} \simeq \mathfrak{N}\text{”}$$

where  $\underline{Q}$  is a  $P_\kappa$ -name. So by (8) we may assume that for some large enough  $i < \kappa$   $\mathfrak{M}$  and  $\mathfrak{N}$  are  $P_i$ -names. By (6) wlog we may assume that  $\mathfrak{M} = \mathfrak{M}_i$ ,  $\mathfrak{N} = \mathfrak{N}_i$ . Now letting  $R$  be the  $P_i$ -name  $(P_\kappa/G_i) * \underline{Q}$ , we get

$$p \Vdash_{-i} \text{“}\Vdash_{-R} \mathfrak{M} \simeq \mathfrak{N}\text{”}$$

But by 2.10,  $\Vdash_{-i}$  “ $\kappa$  is reflecting”, so by the definition of  $\underline{Q}_i$  and by 2.9 we get that  $p \Vdash_{-i+1} \mathfrak{M} \simeq \mathfrak{N}$ . ☺ 2.11

**2.12 Remark:** Since 2.8 is also true with “proper” replaced by “semiproper”, we similarly get that the consistency of a  $\Sigma_1$ -reflecting cardinal implies the consistency of the bounded semiproper forcing axiom. ☺ 2.12

### 3. Sealing the $\omega_1$ -branches of a tree

In this section we will define a forcing notion which makes the set of branches of an  $\omega_1$ -tree absolute.

**3.1 Definition:** Let  $T$  be a tree of height  $\omega_1$ . We say that  $B \subseteq T$  is an  $\omega_1$ -branch if  $B$  is a maximal linearly ordered subset of  $T$  and has order type  $\omega_1$ .

**3.2 Lemma:** Let  $T$  be a tree of height  $\omega_1$ . Assume that every node of  $T$  is on some  $\omega_1$ -branch, and that there are at uncountably many  $\omega_1$ -branches. (These assumptions are just to simplify the notation). Then there is a proper forcing notion  $P'_T$  (in fact,  $P'_T$  is a composition of finitely many  $\sigma$ -closed and ccc forcing notions) forcing the following:

- (1)  $T$  has  $\aleph_1$  many  $\omega_1$ -branches, i.e., there is a function  $b : \omega_1 \times \omega_1 \rightarrow T$  such that each set  $B_\alpha = \{b(\alpha, \beta) : \beta < \omega_1\}$  is an end segment of a branch of  $T$  (enumerated in its natural order), and every  $\omega_1$ -branch is (modulo a countable set) equal to one of the  $B_\alpha$ s, and the sets  $B_\alpha$  are pairwise disjoint.
- (2) There is a function  $g : T \rightarrow \omega$  such that for all  $s < t$  in  $T$ , if  $g(s) = g(t)$  then there is some (unique)  $\alpha < \omega_1$  such that  $\{s, t\} \subseteq B_\alpha$ .

The proof consists of two parts. In the first part (3.3) we show that we may wlog assume that  $T$  has actly  $\aleph_1$  many branches. This observation is a special case of a theorem of Mitchell [Mi, 3.1].

In the second part we describe the forcing notion which works under the additional assumption that  $T$  has only  $\aleph_1$  many branches. This forcing notion is essentially the same as the one used by Baumgartner in [Ba2, section 8].

**3.3 Fact:** Let  $T$  be a tree of height  $\omega_1$ ,  $\kappa > |T|$ , and let  $R_1$  be the forcing notion adding

$\kappa$  many Cohen reals. In  $V^{R_1}$ , let  $R_2$  be a  $\sigma$ -closed forcing notion. Then every branch of  $T$  in  $V^{R_1 * R_2}$  is already in  $V^{R_1}$  (and in fact already in  $V$ ).

Hence, taking  $R_2$  to be the Levy collapse of the number of branches of  $T$  to  $\aleph_1$  (with countable conditions),  $T$  will have at most  $\aleph_1$  many branches in  $V^{R_1 * R_2}$ .

Proof: Assume that  $\underline{b}$  is a name of a new branch. So the set

$$T_{\underline{b}} := \{t \in T : \exists p \in R_2 p \Vdash t \in \underline{b}\}$$

is (in  $V^{R_1}$ ) a perfect subtree of  $T$ . In particular, there is an order-preserving function  $f : 2^{<\omega} \rightarrow T_{\underline{b}}$ . Since  $\kappa$  was chosen big enough, we can find a real  $c \in 2^\omega \cap V^{R_1}$  which is not in  $V[f]$ . Now note that  $T'$  is  $\sigma$ -closed, so there is  $t^* \in T$  such that  $\forall n f(c|n) \leq t^*$ . But this implies that

$$c = \bigcup \{s \in 2^{<\omega} : f(s) \leq t^*\}$$

can be computed in  $V[f]$ , a contradiction. ☺ 3.3

Now we describe a forcing notion  $P'_T$  which works under the assumption that  $T$  has not more than  $\aleph_1$  branches. In the general case we can then use the forcing  $P_T = R_1 * R_2 * P'_T$ .

**3.4 Definition:** Let  $T$  be a tree of height  $\omega_1$  with  $\aleph_1$  many  $\omega_1$ -branches  $\{B_i : i < \omega_1\}$  and assume that each node of  $T$  is on some  $\omega_1$ -branch. Let  $B'_j = B_j \setminus \bigcup_{i < j} B_i$ ,  $x_j = \min(B'_j)$  so that the sets  $B'_j$  are disjoint end segments of the branches  $B_j$ , and they form a partition of  $T$ . Let  $A = \{x_i : i < \omega_1\}$ .

The forcing “sealing the branches of  $T$ ” is defined as

$$P'_T = \{f : f \text{ a finite function from } A \text{ to } \omega, \text{ and if } x < y \text{ are in } \text{dom}(f), \text{ then } f(x) \neq f(y)\}$$

**3.5 Lemma:**  $P'_T$  satisfies the countable chain condition. (In fact, much more is true: If

$\langle p_i : i < \omega_1 \rangle$  are conditions in  $P$ , then there are uncountable sets  $S_1, S_2 \subseteq \omega_1$  such that whenever  $i \in S_1, j \in S_2$ , then  $p_i$  and  $p_j$  are compatible. See [Sh f, XI])

Proof: Essentially the same as in [Ba2, 8.2].

☺ 3.5

To conclude the proof of 3.2, note that any generic filter  $G$  on  $P'_T$  induces a generic  $f_G : A \rightarrow \omega$ . Let  $g_G : T \rightarrow \omega$  be defined by  $g_G(y) = f_G(x_i)$  for all  $y \in B_i$ . This function  $g_G$  fulfills the requirement 3.2(2).

☺ 3.2

#### 4. BPFA and reflecting cardinals are equiconsistent

In this section we will show that

**4.1 Theorem:** If BPFA holds, then the cardinal  $\aleph_2$  (computed in  $V$ ) is  $\Sigma_1$ -reflecting in  $L$ .

Before we start the proof of this theorem, we show some general properties of “sufficiently generic” filters.

First a remark on terminology: When we consider  $\text{BFA}(P, \lambda)$ , then by “for all sufficiently generic  $G^* \subseteq P$ ,  $\varphi(G^*)$  holds” we mean: “there is a  $P$ -name  $\underline{f} : \lambda \rightarrow \lambda$  such that: whenever a filter  $G^*$  interprets  $\underline{f}$ , then  $\varphi(G^*)$  will hold”. A description of the name  $\underline{f}$  can always be deduced from the context. Instead of a single name  $\underline{f}$  we usually have a family of  $\lambda$  many names.

The first lemma shows that from any sufficiently generic filter we can correctly compute the first order theory (that is, the part of it which is forced), or equivalently, the first order diagram, of any small structure in the extension.

**4.2 Lemma:** Let  $P$  be a forcing notion,  $\Vdash_P$  “ $\mathfrak{M}$  is a structure with universe  $\lambda$  with  $\lambda$

many relations  $(\underline{R}_i : i < \lambda)$ ". Assume  $\text{BFA}(P, \lambda)$ . Then for every sufficiently generic filter  $G^* \subseteq P$ , letting  $\mathfrak{M}^* = (\lambda, \underline{R}_i[G^*])_{i < \lambda}$ , (where  $\underline{R}_i[G^*] := \{(x_1, \dots, x_k) \in \lambda^n : \exists p \in G^* p \Vdash \mathfrak{M} \models \underline{R}_i(x_1, \dots, x_k)\}$ ) we have:

Whenever  $\varphi$  is a closed formula such that  $\Vdash_P \mathfrak{M} \models \varphi$ ,  
then  $\mathfrak{M}^* \models \varphi$ .

Proof: Let  $\chi$  be a large enough cardinal, and let  $N$  be an elementary submodel of  $H(\chi)$  of size  $\lambda$  containing all the necessary information (i.e.,  $\lambda \subseteq N$ ,  $(P, \leq) \in N$ ,  $(\underline{R}_i : i < \lambda) \in N$ ).

By  $\text{BFA}(P, \lambda)$  we can find a filter  $G^* \subseteq P$  which decides all  $P$ -names of elements of  $\mathfrak{M}$  which are in  $N$  and all first order statements about  $\mathfrak{M}$ , i.e.,

- (1) For all  $\alpha \in N$ , if  $\Vdash_P \text{"}\alpha \in \lambda\text{"}$  then there is  $\beta \in \lambda$  and  $p \in G^*$  such that  $p \Vdash_P \text{"}\alpha = \check{\beta}\text{"}$ .
- (2) For all  $\alpha_1, \dots, \alpha_k \in \lambda$  and all formulas  $\varphi(x_1, \dots, x_k)$  there is  $p \in G^*$  such that either  $p \Vdash \text{"}\mathfrak{M} \models \varphi(\alpha_1, \dots, \alpha_k)\text{"}$  or  $p \Vdash \text{"}\mathfrak{M} \models \neg\varphi(\alpha_1, \dots, \alpha_k)\text{"}$ .

We now claim that for every formula  $\varphi(x_1, \dots, x_k)$ , for every  $\underline{a}_1, \dots, \underline{a}_k \in N$ : If  $\Vdash_P \text{"}\mathfrak{M} \models \varphi(\underline{a}_1, \dots, \underline{a}_k)\text{"}$ , then  $\mathfrak{M}^* \models \varphi(\underline{a}_1[G^*], \dots, \underline{a}_k[G^*])$ . We assume that  $\varphi$  is in prefix form, so in particular negation signs appear only before atomic formulas. The proof is by induction on the complexity of  $\varphi$ , starting from atomic and negated atomic formulas. We will only treat the case  $\varphi = \exists x \varphi_1$ . So assume that  $\Vdash_P \mathfrak{M} \models \exists x \varphi_1(x, \underline{a}_1, \dots, \underline{a}_k)$ . We can find a name  $\underline{b} \in N$  such that  $\Vdash_P \mathfrak{M} \models \varphi_1(\underline{b}, \underline{a}_1, \dots, \underline{a}_k)$ , so by induction hypothesis we get  $\mathfrak{M}^* \models \varphi_1(\underline{b}[G^*], \underline{a}_1[G^*], \dots, \underline{a}_k[G^*])$ . ☺ 4.2

**4.3 Remark:** In a sense the previous lemma characterizes “sufficiently generic” filters. More precisely, the following is (trivially) true: Let  $P$  be a complete Boolean algebra, let  $\Vdash_P \underline{f} : \lambda \rightarrow \lambda$ , and let  $\mathfrak{M} = (\lambda, \underline{f})$ , where we treat  $\underline{f}$  as a relation. For any ultrafilter

$G^* \subseteq P$  the model  $\mathfrak{M}^* = (\lambda, \underline{f}[G^*])$  is well-defined. Since  $\underline{f}$  is forced to be a function, we have  $\Vdash_P \text{“}\mathfrak{M} \models \forall\alpha \exists\beta (\alpha, \beta) \in \underline{f}\text{”}$ . Clearly  $G^*$  “decides”  $\underline{f}$  (as a function) iff  $\mathfrak{M}^*$  satisfies the same  $\forall\exists$  statement. ☺ 4.3

This last remark suggests the following easy characterization of  $\text{BFA}(P)$ :

**4.4 Definition:** Let  $P$  be an arbitrary forcing notion, not necessarily a complete Boolean algebra. If  $\underline{f}$  is a  $P$ -name of a function from  $\lambda$  to  $\lambda$ , then let the “(forced) diagram” of  $\mathfrak{M} = (\lambda, \underline{f})$  be defined by

$$D^{\parallel-}(\underline{M}) = D^{\parallel-}(\underline{f}) = \{(\varphi, \alpha_1, \dots, \alpha_n) : \varphi(x_1, \dots, x_n) \text{ a first order formula, } \alpha_1, \dots, \alpha_n \in \lambda, \Vdash_P \varphi(\alpha_1, \dots, \alpha_n)\}$$

The “open (forced) diagram”  $D_{qf}^{\parallel-}(\underline{f})$  is defined similarly, but  $\varphi$  ranges only over quantifier-free formulas.

**4.5 Definition:** For any forcing notion  $P$  let  $\text{BFA}'(P, \lambda)$  be the statement

$$\text{BFA}'(P, \lambda) = \text{Whenever } \underline{f} : \lambda \rightarrow \lambda \text{ is a } P\text{-name of a function, then there is a function } f^* \text{ such that } (\lambda, f^*) \models D_{qf}^{\parallel-}(\underline{f}).$$

**4.6 Fact:** For any forcing notion  $P$ ,  $\text{BFA}(P, \lambda)$  iff  $\text{BFA}'(P, \lambda)$ .

Proof:  $\text{BFA}'(P, \lambda)$  is clearly equivalent to  $\text{BFA}'(\text{ro}(P), \lambda)$ . The same is true (by definition) for  $\text{BFA}$ . So we may wlog assume that  $P$  is a complete Boolean Algebra. It is clear that  $\text{BFA}(P, \lambda) \Rightarrow \text{BFA}'(P, \lambda)$ .

Conversely, if  $f^*$  is a function as in  $\text{BFA}'$ , then we claim that the set  $\{\llbracket \underline{f}(\alpha) = f^*(\alpha) \rrbracket : \alpha \in \lambda\}$  generates a filter on  $P$  (where  $\llbracket \varphi \rrbracket$  denotes the Boolean value of a closed statement

$\varphi$ ). Proof of this claim: If not, then there are ordinals  $\alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_n$  such that

$$f^*(\alpha_1) = \beta_1 \ \& \ \dots \ \& \ f^*(\alpha_n) = \beta_n$$

but the Boolean value

$$\llbracket f(\alpha_1) = \beta_1 \ \& \ \dots \ \& \ f(\alpha_n) = \beta_n \rrbracket$$

is 0. This is a contradiction to the fact that  $f^*$  witnesses  $\text{BFA}'(P, \lambda)$ . ☺ 4.6

After this digression we now continue our preparatory work for the proof of theorem 4.1. Our next lemma shows that a generic filter will not only reflect first order statements about small structures, but will also preserve their wellfoundedness.

**4.7 Lemma:** Assume that  $\Vdash_P$  “ $\mathfrak{M} = (\lambda, \underline{E})$  is a well-founded structure,  $\lambda$  is a cardinal”. Assume that  $cf(\lambda) > \omega$ , and assume that  $\text{BFA}(P, \lambda)$  holds. Then for every sufficiently generic filter  $G^* \subseteq P$  we have that  $\mathfrak{M}^* := (\lambda, \underline{E}[G^*])$  is well-founded.

(We will use this lemma only for the case where  $P$  is proper and  $\lambda = \omega_1$ .)

Proof: For each  $\alpha < \lambda$  let  $\underline{r}_\alpha$  be the name of the canonical rank function on  $(\alpha, \underline{E})$ , i.e.,

$$\Vdash_P \text{ “dom}(\underline{r}_\alpha) = \alpha, \ \forall \beta < \alpha \ \underline{r}_\alpha(\beta) = \sup\{\underline{r}_\alpha(\gamma) + 1 : \gamma \underline{E} \beta\}\text{”}$$

As  $\Vdash_P$  “ $\lambda$  is a cardinal”, we have  $\Vdash_P$  “ $\text{rng}(\underline{r}_\alpha) \subseteq \lambda$ ”, so any sufficiently generic filter  $G^*$  will interpret all the functions  $\underline{r}_\alpha$ . Applying lemma 4.2 to the structure  $(\alpha, \underline{E}[G^*], \underline{r}_\alpha[G^*])$  we see that  $\underline{r}_\alpha[G^*]$  is indeed a rank function witnessing that  $(\alpha, \underline{E}[G^*])$  is well-founded. Since  $cf(\lambda) > \omega$  this now implies that also  $(\lambda, \underline{E}[G^*])$  is well-founded. ☺ 4.7

We now start the proof of 4.1. The definitions in the following paragraphs will be valid throughout this section.

Assume BPFA. Let  $\kappa := \aleph_2$ . We will show that  $\kappa$  is reflecting in  $L$ . It is clear that  $\kappa$  is

regular in  $L$ .

**4.8 Claim:** Without loss of generality we may assume:

- (1)  $0^\#$  does not exist, i.e., the covering lemma holds for  $L$ .
- (2)  $\aleph_2^{\aleph_1} = \aleph_2$ .
- (3) There is  $A \subseteq \aleph_2$  such that whenever  $X \subseteq Ord$  is of size  $\leq \aleph_1$ , then  $X \in L[A]$ .

Proof: (1) If  $0^\#$  exists, then  $L_\kappa \prec L$ , and it is easy to see that this implies that  $\kappa$  is a reflecting cardinal in  $L$ .

(2) Let  $P = Levy(\aleph_2, \aleph_2^{\aleph_1})$ , i.e., members of  $P$  are partial functions from  $\aleph_2$  to  $\aleph_2^{\aleph_1}$  with bounded domain. Since  $P$  does not add new subsets of  $\aleph_1$  and  $P$  is proper, also  $V^P$  will satisfy  $PI(\text{proper}, \aleph_1)$ . Also  $\aleph_2^V = \aleph_2^{V^P}$  and  $V^P \models \aleph_2^{\aleph_1} = \aleph_2$ , so we can wlog work in  $V^P$  instead of  $V$ .

(3) By (2) we can find a set  $A \subseteq \aleph_2$  such that  $\aleph_2^{L[A]} = \aleph_2$  and every function from  $\aleph_1$  to  $\aleph_2$  is already in  $L[A]$ . By (1), every set  $X$  of ordinals of size  $\leq \aleph_1$  can be covered by a set  $Y \in L$ ,  $|Y| = \aleph_1$ . Let  $j : Y \rightarrow otp(Y)$  be order preseving, then  $j[X] \in L[A]$ ,  $j \in L$ , so  $X \in L[A]$ . ☺ 4.8

Proof of 4.1: Let  $\varphi(x)$  be a formula,  $a \in L_\kappa$ , and assume that  $\chi > \kappa$ ,  $L_\chi \models \varphi(a)$ ,  $\chi$  a regular cardinal in  $L$ . We have to find an  $L$ -cardinal  $\chi' < \kappa$  such that  $a \in L_{\chi'}$  and  $L_{\chi'} \models \varphi(a)$ .

By 2.3, we may assume that  $\chi$  is a cardinal in  $L[A]$  or even in  $V$ .

Informal outline of the proof: We will define a forcing notion  $P$ . In  $V^P$  we will construct a model  $\mathfrak{M} = (M, \in, \chi, x, \dots) \prec V^P$  of size  $\aleph_1$  containing all necessary information. This model has an isomorphic copy  $\bar{\mathfrak{M}}$  with underlying set  $\omega_1$ . We will find a “sufficiently



generic” filter  $G^*$  which will “interpret”  $\bar{\mathfrak{M}}$  as  $\mathfrak{M}^*$ . By 4.7 we may assume that  $\mathfrak{M}^* = (\omega_1, E^*, \chi^* \dots)$  will be well-founded, so we can form its transitive collapse  $\mathfrak{M}' = (M', \in, \chi', \dots)$ . By 4.2 we have that  $\mathfrak{M}' \models \text{“}\chi' \text{ is a cardinal in } L\text{”}$ , i.e.,  $\chi'$  is a cardinal in  $L_{M' \cap Ord}$ . The main point will be to show that any filter on our forcing notion  $P$  will code enough information to enable us to conclude that  $\chi'$  is really a cardinal of  $L$ .

**4.9 Definition of the forcing notions  $Q_0$  and  $Q_1$ :** Let  $Q_0$  be the Levy-collapse of  $L_\chi[A]$  to  $\aleph_1$ , i.e. the set of countable partial functions from  $\omega_1$  to  $L_\chi[A]$  ordered by extension.

In  $V^{Q_0}$  let  $T$  be the following tree: Elements of  $T$  are of the form

$$\langle \langle \mu_i : i < \alpha \rangle, \langle f_{ij} : i \leq j < \alpha \rangle \rangle$$

(we will usually write them as  $\langle \mu_i, f_{ij} : i \leq j < \alpha \rangle$ ), where the  $\mu_i$  are ordinals  $< \chi$ , the  $f_{ij}$  are a system of commuting order-preserving embeddings, and  $\alpha < \omega_1$ .  $T$  is ordered by the relation “is an initial segment of”.

If  $B$  is a branch of  $T$  (in  $V^{Q_0}$ , or in any bigger universe) of length  $\delta$  then  $B$  defines a directed system  $\langle \mu_i, f_{ij} : i \leq j < \delta \rangle$  of well-orders. We will call the direct limit of this system  $(\gamma_B, <_B)$ . In general this may not be a well-order, but it is clear that if the length of  $B$  is  $\omega_1$ , then  $(\gamma_B, <_B)$  will be a well-order.

Let  $Q_1 = P_T$  be the forcing “sealing the  $\omega_1$ -branches of  $T$ ” described in 3.2. We let  $P = Q_0 * Q_1$ . So  $P$  is a proper order, in fact it is a finite iteration of  $\sigma$ -closed and ccc partial orderings.

**4.10 Definition:** In  $V^P$  we define a model  $\mathfrak{M}$  as follows: Let  $\Omega$  a large enough regular cardinal of  $V$ . Let  $(M, \in)$  be an elementary submodel of  $(H(\Omega)^{V^P}, \in)$  of size  $\aleph_1$  containing all necessary information, in particular  $M \supseteq L_\chi[A]$ . We now expand  $(M, \in)$  to a model

$\mathfrak{M} = (M, \in, \chi, A, \dots)$  by adding the following functions, relations and constants:

- a constant for each element of  $L_\xi$  (where  $\xi$  is chosen such that  $a \in L_\xi$ )
- relations  $M_0$  and  $M_1$  which are interpreted as  $M \cap H(\Omega)^V$  and  $M \cap H(\Omega)^{V^{Q_0}}$ , respectively.
- constants  $\chi, A, \kappa, T, g, b$  ( $b$  is the function enumerating the branches of  $T$  from 3.2, and  $g$  is the specializing function  $g : T \rightarrow \omega$  also from 3.2)
- a function  $c : \chi \times \omega_1 \rightarrow \chi$  such that for all  $\delta < \chi$ : If  $cf(\delta) = \aleph_1$ , then  $c(\delta, \cdot) : \omega_1 \rightarrow \delta$  is increasing and cofinal in  $\delta$ .

Since  $M$ , the underlying set of  $\mathfrak{M}$ , is of cardinality  $\aleph_1$ , we can find an isomorphic model

$$\bar{\mathfrak{M}} = (\omega_1, \bar{E}, \bar{\chi}, \dots)$$

In  $V$  we have names for all the above:  $\bar{\mathfrak{M}}, \bar{E}$ , etc. Now let  $G^*$  be a sufficiently generic filter, i.e.,  $G^*$  will interpret all these names. Writing  $E^*$  for  $\bar{E}[G^*]$ , etc., and letting  $\mathfrak{M}^* = (\omega_1, E^*, \chi^*, \dots)$ , we may by 4.7 and 4.2 assume that the following holds:

**4.11 Fact:**

- (1)  $(\omega_1, E^*)$  is well-founded.
- (2) If  $\psi$  is a closed formula such that  $\Vdash_P \text{“}\mathfrak{M} \models \psi\text{”}$ , then  $\mathfrak{M}^* \models \psi$ .

**4.12 Main definition:** We let

$$\mathfrak{M}' = (M', \in, \chi', \dots)$$

be the Mostowski collapse of  $\mathfrak{M}^*$ . This is possible by 4.11(1).  $\mathfrak{M}'_0 = (M'_0, \in)$  and  $\mathfrak{M}'_1 = (M'_1, \in)$  will be “inner models” of  $\mathfrak{M}'$ .

**Note:** We will now do several computations and absoluteness arguments involving the

universes  $V$ ,  $L[A']$ ,  $\mathfrak{M}'$ ,  $L[A']^{\mathfrak{M}'} = L_{M' \cap Ord}[A']$ , etc. By default, all set-theoretic functions, quantifiers, etc., are to be interpreted in  $V$ , but we will often also have to consider relativized notions, like  $\mathfrak{M}' \models "L[A'] \models \dots"$  (which is of course equivalent to  $L_{M' \cap Ord}[A] \models \dots$ ), or  $cf^{L[A']}$ , etc.

Note that  $\mathfrak{M}' \models "L[A'] \models \kappa' = \aleph_2"$ , so we get  $\aleph_1^{\mathfrak{M}'} = \aleph_1^V$ .

We will finish the proof of 4.1 with the following two lemmas:

**4.13 Lemma:**  $a \in L_{\chi'}$ ,  $L_{\chi'} \subseteq M'$  and  $L_{\chi'} \models \varphi(a)$ .

**4.14 Lemma:**  $L \models \chi'$  is a cardinal.

Proof of 4.13: Since  $\chi' + 1 \subseteq M'$  and  $\mathfrak{M}'$  satisfies a large fragment of ZFC, we have  $L_{\chi'} \subseteq M'$  and  $L_{\chi'} \in M'$ . For each  $y \in L_{\xi}$  let  $c_y$  be the associated constant symbol, then by induction (using 4.11(2)) it is easy to show that  $y = c_y^{\mathfrak{M}'}$  for all  $y \in L_{\xi}$ . Since  $\Vdash_P \mathfrak{M} \models [L_{\chi} \models \varphi(a)]$ , we thus have  $\mathfrak{M}' \models "L_{\chi'} \models \varphi(a)"$ . But  $L_{\chi'} \subseteq M'$ , so  $L_{\chi'} \models \varphi(a)$ . ☺ 4.13

So we are left with proving 4.14. In  $L[A']$  let  $\mu$  be the cardinality of  $\chi'$ , and (again in  $L[A']$ ) let  $\nu$  be the successor of  $\mu$ . We will prove 4.14 by showing the following fact:

**4.15 Lemma:**  $\nu \subseteq M'$ .

Proof of 4.14(using 4.15): In fact we show that 4.15 implies that  $\chi'$  is a cardinal even in  $L[A']$ : If not, then  $\mu < \chi'$ , and since  $\nu$  is a cardinal in  $L[A']$  we can find a  $\gamma < \nu$  such that  $L_{\gamma}[A'] \models$ “there is a function from  $\mu$  onto  $\chi'$ ”. By 4.15,  $\gamma \in M'$ , so by the well-known absoluteness properties of  $L$  we have  $L_{\gamma}[A'] \subseteq M'$ , so  $\mathfrak{M}' \models "L[A'] \models \chi'$  is not a cardinal.” But we also have  $\Vdash_P \mathfrak{M} \models "L[A] \models \chi$  IS a cardinal”, so we get a contradiction to 4.11(2). ☺ 4.14

Proof of 4.15: We will distinguish two cases, according to what the cofinality of  $\mu$  is.

**Case 1:**  $cf(\mu) = \aleph_0$ . (This is the “easy” case, for which we do not need to know anything about the forcing  $Q_1$  other than that it is proper, so the class  $\{\delta : cf(\delta) = \aleph_0\}$  is the same in  $V, V^{Q_0}, V^P, L[A]$ ). We start our investigation of case 1 with the following remark:

**4.16 Fact:**

- (1) For all  $\delta$ : If  $cf^{L[A]}(\delta) > \aleph_0$ , then  $cf(\delta) > \aleph_0$ .
- (2)  $\Vdash_P$  “For all  $\delta < \chi$ : If  $cf^{L[A]}(\delta) > \aleph_0$ , then  $cf(\delta) = \aleph_1$ ”.
- (3) If  $\mathfrak{M}' \models cf^{L[A]}(\mu) > \aleph_0$ , then  $\mathfrak{M}' \models cf(\mu) = \aleph_1$ .
- (4) If  $\mathfrak{M}' \models “cf(\mu) = \aleph_1”$ , then  $cf(\mu) = \aleph_1$ .

Proof: (1): By the choice of  $A$ . (4.8(3)).

(2): Use (1) and the fact that  $P$  is proper, hence does not cover old uncountable sets by new countable sets.

(3): Use (2) and 4.2.

(4): If  $\mathfrak{M}' \models “cf(\mu) = \aleph_1”$ , then the function  $c'(\mu, \cdot)$  is increasing and cofinal in  $\mu$ . (Recall that  $\omega_1^V = \omega_1^{\mathfrak{M}'}$ ) ☺ 4.16

**4.17 Conclusion:** Since  $cf(\mu) = \aleph_0$ , we get from (3) and (4):  $\mathfrak{M}' \models “L[A'] \models ‘cf(\mu) = \aleph_0’”$ .

Let  $\mathfrak{M}' \models “\nu_1$  is the  $L[A']$ -successor of  $\mu.”$  We will show that  $\nu_1 = \nu$ . This suffices, because  $M'$  is transitive.

So assume that  $\nu_1 < \nu$ . Working in  $L[A']$  we have  $|\mu^{\aleph_0}| = \nu$  and  $|L_{\nu_1}[A']| < \nu$ , so we can find a  $y \in [\mu]^{\aleph_0}$ ,  $y \in L_\gamma[A'] \setminus L_{\nu_1}[A']$  for some  $\gamma < \nu$ . Working in  $V$ , let  $L_\gamma[A'] = \bigcup_{i < \omega_1} X_i$ , where  $\langle X_i : i < \omega_1 \rangle$  is a continuous increasing chain of elementary countable submodels of

$L_\gamma[A']$ , with  $y, A' \in X_0$ . In  $\mathfrak{M}'_1 = (M'_1, \in)$  we can find a continuous increasing sequence  $\langle Y_i : i < \omega_1 \rangle$  of countable elementary submodels of  $L_\mu[A']$  with  $\bigcup_{i < \omega_1} Y_i = L_\mu[A']$  and  $A' \in Y_0$ . We can find an  $i$  such that  $X_i \cap L_\mu[A'] = Y_i$ .

Let  $j : (X_i, \in, A, Y_i) \rightarrow (L_{\hat{\gamma}}[\hat{A}], \in, \hat{A}, L_{\hat{\mu}}[\hat{A}])$  be the collapsing isomorphism.

Now note that  $Y_i = X_i \cap L_\mu[A']$  is a transitive subset of  $X_i$ , so  $j \upharpoonright Y_i$  is exactly the Mostowski collapse of  $(Y_i, \in)$ , so  $j \upharpoonright Y_i \in M'_1$  and  $\hat{A} \in M'_1$ . Hence also  $j(y) \in L_{\hat{\gamma}}[\hat{A}] \subseteq M'_1$ , so we can compute

$$y = \{\alpha : (j \upharpoonright Y_i)(\alpha) \in j(y)\}$$

in  $\mathfrak{M}'_1$ . Hence  $y \in M'_1$ . But  $\mathfrak{M}' \models "[\mu]^{\aleph_0} \cap M'_1 = [\mu]^{\aleph_0} \cap M'_0 = [\mu]^{\aleph_0} \cap L[A']"$  (the first equality holds because  $Q_0$  is a  $\sigma$ -closed forcing notion, the second because of our assumption 4.8(3))

Hence  $\mathfrak{M}' \models y \in L[A']$ , so  $\mathfrak{M}' \models y \in L_{\nu_1}[A']$ , a contradiction to our choice of  $\nu$ .

☺ 4.15 Case 1

**Case 2:**  $cf(\mu) = \aleph_1$ . Let  $\gamma < \nu$ . We have to show that  $\gamma \in M'$ . Since  $L[A'] \models |\gamma| = \mu$ , we can in  $L[A']$  find an increasing sequence  $\langle A_\xi : \xi < \mu \rangle$ ,  $\gamma = \bigcup_{\xi < \mu} A_\xi$ , where each  $A_\xi$  has (in  $L[A']$ ) cardinality  $< \mu$ . Let  $\alpha_\xi$  be the order type of  $A_\xi$ , then the inclusion map from  $A_\xi$  into  $A_\zeta$  naturally induces an order preserving function  $f_{\xi\zeta} : \alpha_\xi \rightarrow \alpha_\zeta$ . Let  $B = \langle \alpha_\xi, f_{\xi\zeta} : \xi \leq \zeta < \mu \rangle$ , and write  $B \upharpoonright \beta$  for  $\langle \alpha_\xi, f_{\xi\zeta} : \xi \leq \zeta < \beta \rangle$ . Clearly the direct limit of this system is a well-ordered set of order type  $\gamma$ .

So  $B$  is in  $L[A']$ , but we can moreover show that each initial segment  $B \upharpoonright \beta$  is already in  $L_\mu[A']$ . This follows from the fact that each such initial segment can be canonically coded by a bounded subset of  $\mu$ .

Since  $L_\mu[A'] \subseteq L_{\chi'}[A'] \subseteq M'_1$ , we know that  $B \restriction \beta$  is in  $M'_1$  for all  $\beta < \mu$ . In  $M'_1$  let  $\langle \xi_i : i < \omega_1 \rangle$  be an increasing cofinal subsequence of  $\mu$ . Let  $\beta_i = \alpha_{\xi_i}$ ,  $h_{ij} = f_{\xi_i, \xi_j}$ . Note that the direct limit of the system  $\langle \beta_i, h_{ij}; i \leq j < \omega_1 \rangle$  is still a well-ordered set of order type  $\gamma$ .

So for each  $\delta < \omega_1$  we know that the sequence  $b_\delta := \langle \beta_i, f_{ij} : i \leq j \leq \delta \rangle$  is in  $M'_1$ , and  $\mathfrak{M}'_1 \models b_\delta \in T'$ .

Now we can (in  $V$ ) find an uncountable set  $C \subseteq \omega_1$  and a natural number  $n$  such that for all  $\delta \in C$  we have  $g'(b_\delta) = n$ . Now recall the characteristic property of  $g$  (see 3.2) and hence of  $g'$  (by 4.11): for each  $\delta_1 < \delta_2$  in  $C$  we have a unique branch  $B'_\alpha = \{b'(\alpha, \beta) : \beta < \omega_1\}$  with  $\{b_{\delta_1}, b_{\delta_2}\} \subseteq B_\alpha$ . A priori this  $\alpha$  depends on  $\delta_1$  and  $\delta_2$ , but since  $B_\alpha \cap B_\beta = \emptyset$  for  $\alpha \neq \beta$  we must have the same  $\alpha$  for all  $\delta \in C$ .

So the sequence  $\langle b_\delta : \delta \in C \rangle$  is cofinal on some branch  $B'_\alpha$  which is in  $\mathfrak{M}'$ . So we get that  $\gamma$ , the order type of the limit of this system, is also in  $M'$ . ☺ 4.15 Case 2 ☺ 4.1 ☺ [GoSh 507]

References.

- [Ba1] J. Baumgartner, *Applications of the Proper forcing axiom*, in: Handbook of set-theoretic topology, 915–959.
- [Ba2] J. Baumgartner, *Iterated forcing*, in: Surveys in set theory (A. R. D. Mathias, editor), London Mathematical Society Lecture Note Series, No. 8, Cambridge University Press, Cambridge, 1983.
- [Fu] Sakae Fuchino, *On potential embedding and versions of Martin’s Axiom*, Notre Dame Journal of Formal Logic, vol 33, 1992.
- [Mi] Bill Mitchell, *Aronszajn Trees and the independence of the transfer property*, Annals of Mathematical Logic, 5 (1971), pp.21–46.
- [Sh 56] Saharon Shelah, *Refuting Ehrenfeucht Conjecture on rigid models*, Proc. of the Symp. in memory of A. Robinson, Yale, 1975, A special volume in the *Israel J. of Math.*, 25 (1976) 273-286.
- [Sh 73] Saharon Shelah, *Models with second order properties II. On trees with no undefinable branches*, Annals of pure and applied logic, 14 (1978), pp.73–87.
- [Sh b] S. Shelah, *Proper Forcing*, Lecture Notes in Mathematics Vol. 942, Springer Verlag.
- [Sh f] S. Shelah, *Proper and Improper Forcing*, to appear.
- [To] S. Todorčević, *A Note on The Proper Forcing Axiom*, in: Axiomatic set theory, J. Baumgartner, D.A. Martin, (eds.), Contemporary Mathematics 31 (1984).

Martin Goldstern

Institut für Algebra und Diskrete Mathematik,

TU Wien

Wiedner Hauptstraße 8–10/118.2

A-1040 Wien, Austria

goldstrn@email.tuwien.ac.at

Saharon Shelah

Department of Mathematics

Hebrew University of Jerusalem

91904 Jerusalem, Givat Ram

Israel

shelah@math.huji.ac.il