

Randomness and Semigenericity*

John T. Baldwin[†]

Department of Mathematics, Statistics and Computer Science
University of Illinois at Chicago

Saharon Shelah[‡]

Department of Mathematics
Hebrew University of Jerusalem
Rutgers University

October 6, 2003

Abstract

Let L contain only the equality symbol and let L^+ be an arbitrary finite symmetric relational language containing L . Suppose probabilities are defined on finite L^+ structures with ‘edge probability’ $n^{-\alpha}$. By T^α , the almost sure theory of random L^+ -structures we mean the collection of L^+ -sentences which have limit probability 1. T_α denotes the theory of the generic structures for \mathbf{K}_α , (the collection of finite graphs G with $\delta_\alpha(G) = |G| - \alpha \cdot |\text{edges of } G|$ hereditarily nonnegative.)

0.1 Theorem. T^α , the almost sure theory of random L^+ -structures is the same as the theory T_α of the \mathbf{K}_α -generic model. This theory is complete, stable, and nearly model complete. Moreover, it has the

*Mathematics Review Numbers: 03C10, 05C80, keywords: random graphs, 0 – 1-laws, stability

[†]Partially supported by NSF grant 9308768 and a visit to Simon Fraser University

[‡]This is paper 528. Both authors thank Rutgers University and the Binational Science Foundation for partial support of this research.

finite model property and has only infinite models so is not finitely axiomatizable.

This paper unites two apparently disparate lines of research. In [8], Shelah and Spencer proved a 0–1-law for first order sentences about random graphs with edge probability $n^{-\alpha}$ where α is an irrational number between 0 and 1. Answering a question raised by Lynch [5], we extend this result from graphs to hypergraphs (i.e. to arbitrary finite symmetric relational languages). Let T^α denote the set of sentences with limit probability 1. The Spencer-Shelah proof proceeded by a process of quantifier elimination which implicitly showed the theories T^α were *nearly model complete* (see below) and complete.

Hrushovski in [3] refuted a conjecture of Lachlan by constructing an \aleph_0 -categorical strictly stable pseudoplane. Baldwin and Shi [1] considered a variant on his methods to construct strictly stable (but not \aleph_0 -categorical) theories T_α indexed by irrational α . In this paper we show that for each irrational α , $T^\alpha = T_\alpha$ and thus deduce that T_α is not finitely axiomatizable and that T^α is stable.

Each T_α is the theory of a ‘generic’ model M_α of an amalgamation class \mathbf{K}_α of finite structures. Although the Hrushovski examples are easily seen to be nearly model complete this is less clear for the T_α since they are not \aleph_0 -categorical. We show that each T_α is nearly model complete.

In the first, purely model theoretic, section of the paper we describe our basic framework and prove a sufficient condition for certain theories, including the T_α , to be nearly model complete. These conditions depend upon a generalization of the notion of *genericity* of a structure: *semigenericity*, which is introduced in this paper. In the second section we consider the addition of random relations and deduce the main results for this case: The almost-sure theory and the theory of the generic model are equal, complete, stable, nearly model complete, and not finitely axiomatizable. From the model theoretic standpoint the extension from graphs to an arbitrary finite relational language is not a big step; it was

spelled out in [9]. The distance is larger from the probability standpoint and the problem of making such an extension had been raised by Lynch [5].

The first author greatly benefited from discussions on this paper with M. Albert, G. Cherlin, M. Itai, A.H. Lachlan, C. Laskowski, D. Kueker and D. Marker. We want to thank Shmuel Lifsches for a careful reading of Section 1.

1 Near model completeness

After Hrushovski's construction of counterexamples to the conjectures of Lachlan and Zil'ber a number of authors explored generalizations of the variation he had introduced on the Fraissé-Jonsson construction. Hrushovski had noted that in his situation, where the generic model was ω -saturated, the theory of the generic admitted the level of quantifier elimination which we christen "nearly model complete" in

this paper. We reprise one general setting for this study here and in the next section connect it with certain random models. Baldwin and Shi [1] studied a situation where the homogeneous-universal model, renamed *generic* by Kueker and Laskowski [4], is not ω -saturated. Kueker and Laskowski investigated the conditions in which the theory constructed from a generic admitted various levels of quantifier elimination. After the first author noticed the connection between [3] and [8], we began to consider the quantifier complexity of the theory T^α . There is no explicit elimination of quantifiers result in [8] but a lemma similar to our Lemma 1.30 which is the crucial technical step. The second author had already begun notes generalizing [8]; the 0 – 1-law in Section 2 contains a more concrete version of his approach. Shelah has continued this approach to the probability aspect in more generality in [6]. A close look at the quantifier elimination results in [3, 8], suggests the following definition.

1.1 Definition. A theory T is said to be *nearly model complete* if every formula is equivalent in T to a Boolean combination of Σ_1 -formulas.

Thus, T is nearly model complete if the type of any finite sequence is determined by exactly the family of Σ_1 -formulas it satisfies. Near model completeness lies strictly in strength between model completeness and 1-model completeness (every formula is equivalent to a Σ_2 -formula).

1.2 Notation. Fix a finite relational language L . For any class \mathbf{K} of structures, $S(\mathbf{K})$ denotes the class of all substructures of members of \mathbf{K} . Let \mathbf{K}_0 be a collection of finite L -structures and \mathbf{K} be a class of models whose finite substructures are in $S(\mathbf{K}_0)$. We always assume that the empty structure is in \mathbf{K}_0 . We will consider several different choices for \mathbf{K}_0 in this paper. In the following, A, B, C vary over \mathbf{K}_0 ; M, N over \mathbf{K} . If A, B are subsets of N , we write AB for the L -structure contained in N with universe $A \cup B$.

If $B \cap C = A$ we write $B \otimes_A C$ for the structure with universe $B \cup C$ and no relations other than those on B or C . If A, B, C are substructures of N such that the structure imposed by N on BC is isomorphic to $B \otimes_A C$ we say B and C are *freely joined over A in N* . In general we do not assume \mathbf{K}_0 is closed under \otimes but this assertion will turn out to be an important property of some classes we consider. We write $X \subseteq_\omega Y$ to indicate X is a finite subset of Y .

We will first discuss a class of finite structures equipped with a dimension function satisfying certain natural properties. Then we define from this dimension function a notion of strong submodel. The main quantifier elimination result is proved in terms of the strong submodel concept. But, the connection with random models is obtained by exploiting an appropriate dimension function. The fact that this dimension function (in Example 1.6) is the same as that employed by Hrushovski to construct a strictly stable \aleph_0 -categorical pseudoplane is the key to the argument for the stability of the almost sure theory of random graphs with edge probability $n^{-\alpha}$ when α is irrational.

1.3 Definition. Let δ be an arbitrary function assigning a real number to each isomorphism type of finite L -structure with $\delta(\emptyset) = 0$. $\delta(A/B)$ equals by definition $\delta(AB) - \delta(B)$. This yields immediately:

$$\delta(AB/C) = \delta(A/BC) + \delta(B/C).$$

Note that the structure with universe AB (and thus $\delta(A/B)$) is not determined by the separate structures on A and B but by some embedding of both into an element of \mathbf{K}_0 .

1.4 Notation. We deal only with structures on which the relations of L are symmetric (i.e. $R(\bar{a})$ holds just if it holds for any permutation of \bar{a}) and ir-reflexive (i.e. hold only for sequences of distinct elements). Thus the relations are on sets rather than sequences.

We require the following conditions on δ .

1.5 Axiom. \mathbf{K}_0 and δ satisfy for $A, B, C \dots \in S(\mathbf{K}_0)$ and $N, M \in \mathbf{K}$:

1. $\delta : S(\mathbf{K}_0) \mapsto \mathfrak{R}^+$ (the nonnegative reals) and $\delta(\emptyset) = 0$.

2. If $A, B,$ and C are disjoint subsets of N then $\delta(A/B) \geq \delta(A/BC)$.
3. For every $n \in \omega$ there is an $\epsilon_n > 0$ such that if $|C| < n$ and A, C are disjoint subsets of M with $\delta(CA/A) < 0$ then $\delta(CA/A) \leq -\epsilon_n$.
4. There is a real number $\epsilon > 0$ such that if A, B, B' are disjoint subsets of a model N and $\delta(A/B) - \delta(A/BB') < \epsilon$ then $R(A, B, B') = \emptyset$ and $\delta(A/B) = \delta(A/BB')$.
5. If f is a 1 – 1 homomorphism from A to B then for every $X \subseteq Y \subseteq A$, $\delta(Y/X) \geq \delta(f(Y)/f(X))$.

Axioms iii) and iv) play no explicit role in the argument presented here. But they are important in establishing the stability of T_α in [1] so are used in the proof of Theorem 1.34. Note that Axiom 1.5 iv) is stronger than the assertion that if f is a 1 – 1-homomorphism, $\delta(X) \geq \delta(f(X))$.

Axiom 1.5 i) requires that the range of δ be the nonnegative reals. This allows us to obtain an important monotonicity property by modifying δ to $d : \mathbf{K} \times S(\mathbf{K}_0) \mapsto \mathfrak{R}^+$ by defining for each $N \in \mathbf{K}$,

$$d(N, A) = \inf\{\delta(B) : A \subseteq B \subseteq_\omega N\}.$$

We usually write $d(N, A)$ as $d_N(A)$. We will omit the subscript N if it is clear from context. This operator serves only as a notational convenience within this paper but plays an essential role in establishing the stability of T_α in [1]. The nonnegativity requirement on $\delta|_{\mathbf{K}_0}$ not only justifies the definition of $d_N(A)$ but is necessary for the important Lemma 1.17.

The classes $(\mathbf{K}_\alpha, \delta_\alpha)$, which are defined as follows, are important examples of this situation.

1.6 Example. Let the relation symbols of L be $\langle R_i : i < p \rangle$. Let $w_i(A)$ be the realizations of R_i in A . Fix a sequence $\bar{\alpha}$ with $0 < \alpha_i \leq 1$ for $i < p$. Then for each A , let $e(A) = \sum w_i \alpha_i$. Let \mathbf{K}_α denote the class of all finite L -structures A such that for all substructures A' of A ,

$$\delta_\alpha(A') = |A'| - e(A') \geq 0.$$

See [1] for the straightforward verification of the axioms in this example.

From the dimension function we define certain special notions of submodel which make it easier to formulate our argument.

1.7 Definition. For finite A, B contained in N , define the relative dimension of A over B , $d_N(A/B)$ as $d_N(A/B) = d_N(AB) - d_N(B)$. If $B \subseteq A \subseteq_\omega N$ this simplifies to $d_N(A/B) = d_N(A) - d_N(B)$

1.8 Definition. For $M \subseteq N \in S(\mathbf{K})$, define $M \leq_s N$ if for each finite $X \subseteq M$, $d_M(X) = d_N(X)$. We say M is a *strong* submodel of N . We say $f : M \mapsto N$ is a *strong embedding* if $fM \leq_s N$. We write $M <_s N$ if $M \leq_s N$ but $M \neq N$.

We introduce a second kind of distinguished substructure by defining \leq_i from \leq_s as follows. Note that the definition yields that $A \leq_i A$.

1.9 Definition. For $A, B \in S(\mathbf{K}_0)$, $A \leq_i B$ if $A \subseteq B$ but there is no A' with $A \subseteq A' <_s B$. If $A \leq_i B$, we say B is an *intrinsic* extension of A .

In terms of the dimension function $A \leq_i B$ means $A = B$ or $\delta(B/A) < 0$ and $\delta(B/A) < \delta(B'/A)$ for any intermediate B' .

1.10 Lemma. Consider the situation described in Definition 1.3. If δ is a dimension function satisfying the properties of Axiom 1.5 and \leq_s is defined as in Definition 1.8 then (\mathbf{K}, \leq_s) satisfies the following conditions for $M, N, N' \in S(\mathbf{K})$.

- A1.** $M \leq_s M$.
- A2.** If $M \leq_s N$ then $M \subseteq N$.
- A3.** If $M \leq_s N \leq_s N'$ then $M \leq_s N'$.
- A4.** If $M \leq_s N$, $N' \subseteq N$ then $M \cap N' \leq_s N'$.
- A5.** \leq_i is preserved under 1 – 1 homomorphism.
- A6.** For all $M \in S(\mathbf{K})$, $\emptyset \leq_s M$.

1.11 Remark. Note that **A4** implies that if $M, N, N' \in S(\mathbf{K})$ and $M \leq_s N$, $N' \subseteq N$ and $M \subseteq N'$ then $M \leq_s N'$.

The quantifier elimination results of this section could be obtained by taking as primitive a class \mathbf{K}_0 equipped with a notion of strong submodel, and regarding the results of Lemma 1.10 and Lemma 1.17 below as axioms. Naturally, we would then require that \mathbf{K}_0 and \leq_s be closed under isomorphism. The dimension function is needed for the calculations in Section 2.

1.12 Remark. In earlier formulations, the relation \leq_s was defined just on \mathbf{K}_0 rather than on $S(\mathbf{K}_0)$. This leads to difficulties in phrasing Axiom **A4**.

Our current formulation extends the ideas of [1] to encompass the Baudisch construction of a new \aleph_1 -categorical group [2]. For our purposes in this paper, we could have identified \mathbf{K}_0 with $S(\mathbf{K}_0)$ and we make that restriction in Section 2.

1.13 Remark. Axiom **A6** holds in the examples at hand because the range of δ is nonnegative as specified in Axiom 1.5. In Section 2 we will begin with a δ mapping all finite L - structures into the reals. The requirement that δ is nonnegative requires revising the choice of \mathbf{K}_0 (and thus \mathbf{K}) to guarantee that if $M \in \mathbf{K}$, then for every $A \subseteq_\omega M$, $\delta(A) \geq 0$. We show it is harmless to make this assumption in Lemma 2.19.

1.14 Lemma. 1. For any $A \subseteq C \in \mathbf{K}_0$, we can choose B with $A \leq_i B \leq_s C$.
2. \leq_i is transitive.

Proof. For i), let B have minimal cardinality among the subsets X of C that contain A with $X \leq_s C$. Use **A4** for ii).

1.15 Definition. For any L -structure M , let A, B be finite substructures of M with $A \subseteq B$. Then

1. By a *copy* of B over A in M we mean the image of an extension \hat{f} to B of an embedding f from A into M .
2. $\chi_M(B/A)$ is the number of distinct copies of B over A in M .
3. $\chi_M^*(B/A)$ is the supremum of the cardinalities of maximal families of disjoint (over A) copies of B over A in M .

1.16 Lemma. If \mathcal{B} and \mathcal{B}' are maximal families of disjoint over A copies of B over A then $|\mathcal{B}| \leq |B - A||\mathcal{B}'|$.

Proof. Define a map from \mathcal{B} to \mathcal{B}' by mapping each element of \mathcal{B} to an element of \mathcal{B}' that it intersects off A . This map is at most $|B - A|$ -to-one since the members of \mathcal{B} are disjoint over A .

In particular, this shows that the supremum in the definition of χ_M^* is achieved. As one varies over the entire family of examples of structures

constructed in this manner (e.g. in [3], [1], etc.) the dimension function produces an important trichotomy concerning pairs $A \subseteq B$. Consider an infinite (\mathbf{K}_0, \leq_s) -generic (Definition 1.23) model M . $\chi_M^*(B/A)$ will be bounded if $A \leq_i B$, infinite if $\delta(B/A) > 0$, and will vary with the choice of (\mathbf{K}_0, \leq_s)

if $\delta(B/A) = 0$. The key to the 0–1 law in Section 2 is that when α denotes a sequence, which is linearly independent with 1, the third case cannot occur. The uniform bound on $\chi_M^*(B/A)$ follows from our restricting \mathbf{K}_0 so δ is nonnegative. In [7], Shelah considers a different probability measure which does not permit the nonnegativity restriction; in that situation $\chi_M^*(B/A)$ is a slow growing function. In our situation we have the following.

1.17 Lemma. *There is a binary function $t : \omega \times \omega \mapsto \omega$ which is monotone increasing in both arguments such that if $A \leq_i B$ then for any $M \in \mathbf{K}$ with $A \subseteq M$, $\chi_M(B/A) \leq t(|A|, |B|)$.*

1.18 Remark. This follows easily from Lemma 3.19 of [1]. One must note that $A \leq_i B$ if and only if there is a sequence $A = A_0, A_1, \dots, A_n = B$ such that (A_i, A_{i+1}) is a minimal pair in the sense of [1].

1.19 Definition. For any $M \in \mathbf{K}$, any $m \in \omega$, and any $A \subseteq M$,

$$\text{cl}_M^m(A) = \cup \{B : A \leq_i B \subseteq M \text{ \& } |B - A| < m\}.$$

The following are immediate from Lemma 1.17 and the definitions.

1.20 Lemma. *There is a function f mapping $\omega \times \omega$ into ω such that for any $A \subseteq_\omega M \in \mathbf{K}$ and m , $\text{cl}_M^m(A)$ is finite and its cardinality is uniformly bounded by $f(|A|, m)$.*

1.21 Lemma. *For any M, A, m, n there exists a p depending on $|A|, m, n$ but not on the embedding of A into M with $\text{cl}_M^m(\text{cl}_M^n(A)) \subseteq \text{cl}_M^p(A)$.*

Proof. Let $p = m + f(|A|, n)$ and check.

The next result is immediate noting that $A \leq_i X$ does not depend on any ambient model containing X .

1.22 Lemma. *If $A \subseteq B \subseteq C$ and $\text{cl}_C^m(A) \subseteq B$, then $\text{cl}_C^m(A) = \text{cl}_B^m(A)$.*

1.23 Definition. The countable model $M \in \mathbf{K}$ is (\mathbf{K}_0, \leq_s) -generic if

1. If $A \leq_s M, A \leq_s B \in \mathbf{K}_0$, then there exists $B' \leq_s M$ with $B \cong_A B'$; and
2. M is the union of $\langle A_i : i < \omega \rangle$ where each $A_i \in \mathbf{K}_0$ and $A_i \leq_s A_{i+1}$.

1.24 Definition. A class (\mathbf{K}, \leq_s) has the *amalgamation property* if for any three structures $A, B, C \in \mathbf{K}$ with strong embeddings f, g from A into B, C there exists $D \in \mathbf{K}$ and strong embeddings $f' : B \mapsto D, g' : C \mapsto D$ with $f'f = g'g$.

Following the Fraissé-Jonsson construction, it is easy to show the following result.

1.25 Fact. If (\mathbf{K}_0, \leq_s) satisfies **A0** through **A6** and has the amalgamation property then there is a unique countable (\mathbf{K}_0, \leq_s) -generic model.

We need a more local notion. This is the key new idea of this paper; it arose from the notion of a full model in [1] and from considering the role of $\text{cl}_M^m(A)$ in [8].

1.26 Definition. The countable model M is (\mathbf{K}_0, \leq_s) -semigeneric, or just semigeneric, if

1. $M \in \mathbf{K}$
2. If $A \leq_s B \in \mathbf{K}_0$ and $g : A \mapsto M$, then for each finite m there exists an embedding \hat{g} of B into M which extends g such that
 - (a) $\text{cl}_M^m(\hat{g}B) = \hat{g}B \cup \text{cl}_M^m(gA)$
 - (b) $M \upharpoonright \text{cl}_M^m(gA)\hat{g}B$ is the free join over gA of $\text{cl}_M^m(gA)$ and $\hat{g}B$.

In our applications any generic model is semigeneric (Lemma 1.35), so Fact 1.25 provides us with a semigeneric model. But while generic models are unique there are many semigeneric models in the situations that we deal with here.

We describe below an infinite set of first order formulas $\phi_{A,B,C}^m$ which allow us to axiomatize the class of semigeneric models by the following lemma, which is immediate once we have made the definitions. Note that these are Π_3 -formulas as there is a universal quantifier hidden in the last clause.

1.27 Lemma. *The structure $N \in \mathbf{K}$ is semigeneric, if and only if for each $A \leq_s B$, each $m < \omega$, and $C \in \mathcal{D}_A^m$, $N \models \phi_{A,B,C}^m$*

In establishing the following notation we are suppressing a fixed correspondence between enumerations of the structures A, B, C, D and the variables $\bar{x}, \bar{z}, \bar{y}, \bar{w}$. This correspondence is chosen to preserve natural inclusions among the structures and the variables. Intuitively, \mathcal{D}_A^m is the set of possible isomorphism types for $\text{cl}_M^m(A)$. The structure M satisfies $\phi_{A,B,C}^m$ just if the definition of semigenericity holds for the finite structures $A \subseteq C$ and $A \leq_s B$ when $C \approx \text{cl}_M^m(A)$.

1.28 Notation. 1. Write $A \leq_i^m D$ if for each $d \in D - A$, there is a B with $Ad \subseteq B$, $A \leq_i B$, and $|B - A| < m$.

2. Let $A \in \mathbf{K}_0$.

$$\mathcal{D}_A^m = \{D \in \mathbf{K}_0 : A \leq_i^m D\}.$$

Note that by Lemma 1.20 if $D \in \mathcal{D}_A^m$, $|D| < f(|A|, m)$.

3. For $C \in \mathcal{D}_A^m$ let $\mathcal{D}_{A,C}^m$ be the set of $D \in \mathcal{D}_A^m$ which cannot be embedded in C .

4. For any finite A , $\delta_A(\bar{x})$ denotes the atomic diagram of A .

5. For $C \in \mathcal{D}_A^m$, $\theta_{A,C}^m(\bar{x}, \bar{y})$ is the formula

$$\delta_A(\bar{x}) \wedge \delta_C(\bar{x}, \bar{y}) \wedge (\forall \bar{w}_D) \bigwedge_{D \in \mathcal{D}_{A,C}^m} \neg \delta_D(\bar{x}, \bar{w}_D).$$

Then for $\bar{a}\bar{c}$ an enumeration of C , $A \subseteq C \subseteq N$, $N \models \theta_{A,C}^m(\bar{a}, \bar{c})$ if and only if $C = \text{cl}_N^m(A)$.

6. For $A \leq_s B$ and $C \in \mathcal{D}_A^m$, let $\phi_{A,B,C}^m$ be the sentence

$$(\forall \bar{x})(\forall \bar{y})(\exists \bar{z})[\delta_A(\bar{x}) \wedge \theta_{A,C}^m(\bar{x}, \bar{y}) \rightarrow (\delta_{C \otimes_A B}(\bar{x}, \bar{y}, \bar{z}) \wedge \theta_{B, C \otimes_A B}^m(\bar{x}, \bar{y}, \bar{z}))].$$

1.29 Theorem. *If (\mathbf{K}_0, \leq_s) satisfies **A1-A6** of Lemma 1.10 and Lemma 1.17, then for every formula $\phi(\bar{x})$ there is a Boolean combination of existential formulas $\psi_\phi(\bar{x})$ such that if M is (\mathbf{K}_0, \leq_s) -semigeneric then $\psi_\phi(\bar{x})$ is equivalent to $\phi(\bar{x})$ on M .*

Proof. We first show:

1.30 Lemma. *For any formula $\phi(x_1 \dots x_r)$ there is an integer $\ell = \ell_\phi$, such that for any pair of semigenetics $M, M' \in \mathbf{K}$ and any r -tuples $\bar{a} \in M$ and $\bar{a}' \in M'$ if $\text{cl}_M^{\ell_\phi}(\bar{a}) \approx \text{cl}_{M'}^{\ell_\phi}(\bar{a}')$ by an isomorphism taking \bar{a} to \bar{a}' , then $M \models \phi(\bar{a})$ if and only if $M' \models \phi(\bar{a}')$.*

Proof. The proof is by induction on formula complexity; the cases involving Boolean connectives are easy. So suppose $\phi(\bar{x})$ is of the form $(\exists y)\psi(\bar{x}, y)$. Suppose $M \models \phi(\bar{a})$, so there is a b such that $M \models \psi(\bar{a}, b)$.

Choose p_1 large enough so that for any $N \in \mathbf{K}$, any r -tuple \bar{c} from N and any $d \in N$, $|\text{cl}_N^{\ell_\psi}(\bar{c}, d)| < p_1$. Set $p = \max(p_1, \ell_\psi)$. For $i \leq p$, for any $N \in \mathbf{K}$, for any $\bar{a} \in N$ define by induction $A_0^N = A_0^N(\bar{a}) = \bar{a}$ and $A_{i+1}^N = A_{i+1}^N(\bar{a}) = \text{cl}_N^p(A_i)$. Now applying Lemma 1.21, choose ℓ_ϕ so that for every \bar{a} of length r , and every semigenetic N , $A_p^N(\bar{a}) \subseteq \text{cl}_N^{\ell_\phi}(\bar{a})$.

We want to show that for any semigenetics M and M' , for any $\bar{a} \in M^r$, $\bar{a}' \in M'^r$, and $b \in M$ if $\text{cl}_M^{\ell_\phi}(\bar{a}) \approx \text{cl}_{M'}^{\ell_\phi}(\bar{a}')$ then there is a $b' \in M'$ with $\text{cl}_M^{\ell_\psi}(\bar{a}, b) \approx \text{cl}_{M'}^{\ell_\psi}(\bar{a}', b')$ by an isomorphism taking \bar{a} to \bar{a}' . Let H_0 be the substructure of M with universe (\bar{a}, b) and $H_1 = \text{cl}_M^{\ell_\psi}(H_0)$.

Fix g which maps \bar{a} to \bar{a}' and $\text{cl}_M^{\ell_\phi}(\bar{a})$ isomorphically onto $\text{cl}_{M'}^{\ell_\phi}(\bar{a}')$. By the choice of ℓ_ϕ , for each $i \leq p$, g maps $A_i^M(\bar{a})$ isomorphically onto $A_i^{M'}(\bar{a}')$. (Use Lemma 1.22 and induct.) To avoid superscripts, for each i , let A'_i denote the image of $A_i = A_i^M$ under g . Notice that for some $j \leq p$,

$$(A_{j+1}^M - A_j^M) \cap (H_1 - H_0) = \emptyset.$$

Since $p > |H_1 - A_j^M|$ this implies $A_j^M \leq_s A_j^M H_1$.

Since M' is semigenetic, $M' \models \phi_{A_j, H_1, A_{j+1}}^p$. Thus, there is an isomorphism \hat{g} extending g and mapping H_1 into M' with $\text{cl}_{M'}^p(A'_j \hat{g} H_1) = \text{cl}_{M'}^p(A'_j) \cup \hat{g} H_1$ and so that $M' \upharpoonright (\text{cl}_{M'}^p(A'_j) \hat{g} H_1)$ is a free join of $\text{cl}_{M'}^p(A'_j)$ and $\hat{g} H_1$ over A'_j . Let $H'_1 = \hat{g} H_1$ and $b' = \hat{g}(b)$. We need to show $\text{cl}_M^{\ell_\psi}(\bar{a}, b) \approx \text{cl}_{M'}^{\ell_\psi}(\bar{a}', b')$.

By the choice of \hat{g} and H'_1 , $A'_j H'_1 \cong A_j H_1$ which contains $\text{cl}_M^{\ell_\psi}(\bar{a}, b)$, so it suffices (by Lemma 1.22) to show $A'_j H'_1$ contains $\text{cl}_{M'}^{\ell_\psi}(\bar{a}', b')$. Note $\text{cl}_{M'}^{\ell_\psi}(\bar{a}', b') \subseteq \text{cl}_{M'}^p(\bar{a}', b') \subseteq \text{cl}_{M'}^p(A'_j \hat{g} H_1) = A'_{j+1} H'_1$. By Lemma 1.22, $\text{cl}_{M'}^{\ell_\psi}(\bar{a}', b') = \text{cl}_{A'_{j+1} H'_1}^{\ell_\psi}(\bar{a}', b')$. Since A'_{j+1} and H'_1 are freely joined over A'_j , \hat{g}^{-1} is a 1-1

homomorphism from $A'_{j+1}H'_1$ onto $A_{j+1}H_1$. Applying **A5** from Lemma 1.10, we see $\text{cl}_{A'_{j+1}H'_1}^{\ell_\psi}(\bar{a}', b') \subseteq A'_jH'_1$ whence $\text{cl}_{A'_{j+1}H'_1}^{\ell_\psi}(\bar{a}', b') = \text{cl}_{A'_jH'_1}^{\ell_\psi}(\bar{a}', b')$.

The proof of the following corollary encompasses the derivation of Theorem 1.29 from Lemma 1.30.

1.31 Corollary. *Suppose there is a (\mathbf{K}_0, \leq_s) -semigeneric L -structure. The theory of the class of (\mathbf{K}_0, \leq_s) -semigeneric L -structures is nearly model complete.*

Proof. We have shown that in each semigeneric model the truth of $\phi(\bar{a})$ is determined by the isomorphism type of $\text{cl}_M^{\ell_\phi}(\bar{a})$ and does not depend on the particular embedding of $\text{cl}_M^{\ell_\phi}(\bar{a})$ in M . There are only finitely many possibilities for this closure and each is determined by a conjunction of existential and universal sentences (specifying which B with $|B| < \ell_\phi$ and with \bar{a} enumerating an intrinsic substructure of B occur).

1.32 Corollary. *The theory of the semigeneric models is complete.*

Proof. If N is semigeneric, $N \in \mathbf{K}$ so, by A6, N does not contain any substructure A , with $\delta(A) < 0$. Thus, $\text{cl}_N(\emptyset) = \emptyset$; completeness follows from Lemma 1.30.

Recall from [1]:

1.33 Definition. \mathbf{K}_0 has the *full amalgamation property* if $B \cap C = A$ and $A \leq_s B$ implies $D = B \otimes_A C \in \mathbf{K}_0$ and $C \leq_s D$.

The following result is proved in [1]

1.34 Theorem. $(\mathbf{K}_\alpha, \leq_s)$ has the *full amalgamation property*. There is a generic model M_α and the theory T_α of this generic model is stable.

Using the full amalgamation property, it is easy to see

1.35 Lemma. *The generic model M_α for \mathbf{K}_α is semigeneric.*

Combining the above results we have

1.36 Theorem. T_α is nearly model complete.

The strength of this remark is emphasized by the following observation.

1.37 Theorem. *The theory T_α is not model complete.*

Proof. If T is model complete with generic M , the type of any finite subset X is determined by positive assertions of the substructures that contain X . Fix $A <_i B <_i C \in \mathbf{K}_0$. Suppose $A_1, A_2 \subset M$ with $f : A \approx A_1$ and $f : A \approx A_2$ and suppose $A_1 <_i B_1 \leq_s M$, $A_2 <_i C_2 \leq_s M$ with f and g extending to \hat{f} , \hat{g} such that $\hat{g} : B \approx B_1$ and $\hat{g} : C \approx C_2$. Then every existential formula true of A_1 is true of A_2 but the converse is obviously false. To see the nonobvious assertion, let D be arbitrary with $A_1 \subseteq D \subset M$ and D not contained in B_1 . Then, $B_1 \leq_s B_1 D$. Pulling back to A, B, C , by full amalgamation, there is a $D' \supset D$ such that $BD' \otimes_B C \in \mathbf{K}_0$ and $C \leq_s CD'$. Extending \hat{g} from C to CD' provides the required witness.

1.38 Remark. Let L contain a single binary relation and restrict to the class of graphs. Baldwin and Shi noted [1] that full amalgamation holds for the class \mathbf{K}'_α consisting of those graphs in \mathbf{K}_α which omit squares. Laskowski observed that this argument applies as well to the class K_α^n of graphs which omit cliques of size n . Thus the theory of the generic model associated with each of these classes is stable and nearly model complete.

2 Adding Random relations

In this section we begin with the collection of finite models for a language L with only the equality symbol (i.e. n -element sets for arbitrary n) and add additional ‘random’ relations with respect to probability measures described below.

We show that a 0 – 1 law holds for the set of first order sentences in the expanded language and that the almost sure theory (the sentences with limit probability 1) is stable. Adding a single symmetric irreflexive binary relation gives the family of theories investigated independently by Shelah-Spencer and Baldwin-Shi. Viewing this situation as an expansion of the language of equality may seem eccentric but we expect to exploit this viewpoint for more interesting base languages in the future. This project is well-advanced in [6].

2.1 Context. Let L contain only the equality symbol. The L -structure M_n is a set with n elements. \mathbf{K}_0 is the class of all finite sets and \mathbf{K} the class of all sets. On \mathbf{K} , \leq_s is just \subseteq and $A \leq_i B$ just if $A = B$.

2.2 Remark. The properties **A1-A6** and the conclusion of Lemma 1.17 hold for \mathbf{K} in Context 2.1. Moreover, \mathbf{K}_0 has the full amalgamation property.

2.3 Definition. We say that B is a *primitive* extension of A if $A \leq_s B$ and for every B' with B' properly contained between A and B , B' is not a strong submodel of B .

Now, we show how to define the notion of independent random relations (with edge probability ' $n^{-\alpha}$ ') for an arbitrary finite relational language L^+ . Then we define the notions of dimension and strong submodel in the extended language L^+ and show that the properties **A1-A5** hold for the extended language and **A6** holds with probability 1.

2.4 Notation. We write $[X]^m$ for the collection of m -element subsets of a set X . We will write either $C \in [X]^m$ or (surreptitiously fixing an enumeration of C) $\bar{c} \in [X]^m$ to indicate a member of this set.

2.5 Adding Random relations. Fix an enumeration $\langle R_i : i < p \rangle$ of the relation symbols in $L^+ - L$ and let k_i denote the arity of R_i . Let L_i contain only R_i . Let t denote the largest arity of the R_i . Fix also a sequence of numbers α_i with $0 < \alpha_i \leq 1$ and γ_i with $0 \leq \gamma_i \leq 1$ for $i < p$. (We will require later that the α_i and 1 be linearly independent over the rationals.)

We will define for each isomorphism type of an L^+ structure of size n , the probability of a random structure of size n , having that isomorphism type.

We assume that each new relation in the expanded structure is symmetric and irreflexive in the sense of Paragraph 1.4. Note that this formalism does not describe what one should mean by a random directed graph.

Let N be an L^+ structure of cardinality n . Let C , enumerated as \bar{c} , be a subset of N with size k_i . Let

$$q_{i,n}(C|L_i) = \begin{cases} \gamma_i n^{-\alpha_i} & \text{if } N \models R_i(\bar{c}) \\ 1 - \gamma_i n^{-\alpha_i} & \text{if } N \models \neg R_i(\bar{c}) \end{cases}$$

and,

$$P_n(N) = \prod_{i < p} \prod \{q_{i,n}(C) : C \in [N]^{k_i}\}.$$

2.6 Remark. Let $w_j(N) = |\{C \in [N]^{k_j} : N \models R_j(\bar{c})\}|$ and $\bar{w}_j(N) = |\{C \in [N]^{k_j} : N \models \neg R_j(\bar{c})\}|$. Then

$$P_n(N) = \prod_{j < p} (\gamma_j n^{-\alpha_j})^{w_j(N)} (1 - \gamma_j n^{-\alpha_j})^{\bar{w}_j(N)}.$$

If L^+ has a single binary edge relation and the probability of a two element structure is $n^{-\alpha}$ when the points are related and $1 - n^{-\alpha}$ if not, we return to the situation of [8].

Recall from Lemma 1.27 the sentences axiomatizing the class of semi-generic models. We want to show that the almost sure theory exists and is exactly the theory of the semigenetics. To this end, we will show

$$\lim_{n \rightarrow \infty} P_n(\phi_{A,B,C}^m) = 1$$

for each m, A, B, C .

2.7 Notation. Henceforth, $A, B, \dots, M, N \dots$ range over L -structures. A^+, B^+ etc. denote an expansion of A , respectively B to L^+ . We refer to the universe of A^+ or A by either of these terms rather than the more accurate $|A^+|$ or $|A|$ and reserve $||$ for cardinality. Thus $A^+|L = A$ and we use these notations interchangeably.

We now translate our probability assignment into a class (\mathbf{K}, δ) as in Example 1.6.

2.8 Notation. Let \mathbf{K}_0^* be the collection of all finite L^+ structures.

1. For $A^+ \in \mathbf{K}_0^*$, define $\delta(A^+) = \delta_\alpha(A^+)$ as in Example 1.6, using only the relation symbols in $L^+ - L$ and using the parameters α_i from Paragraph 2.5.
2. $e(B^+/A^+)$ denotes $e(A^+B^+) - e(A^+)$.
3. \mathbf{K}_0^+ denotes the collection of $A^+ \in \mathbf{K}_0^*$ such that for each $A' \subseteq A$, $\delta(A') \geq 0$.
4. $\gamma(A^+, B^+) = \prod_{i < p} \{\gamma_i : |C| = k_i, C \subseteq B, C \not\subseteq A \& B \models R_i(\bar{c})\}$.

2.9 Remark. The link between the function δ and the probabilistic situation is provided in Remark 2.18 where we show that the expectation of the existence of a copy over A^+ of a structure B^+ is determined by $\delta(B^+/A^+)$. Lemma 2.19 implies that replacing \mathbf{K}_0^* by \mathbf{K}_0^+ does not change the almost sure theory. But it does make the model theory conform with the framework of Section 1. The major calculations of this section were carried out in K_0^* in [8].

2.10 Assumption. [Irrationality Hypothesis] The coefficients α_i and 1 are linearly independent over the rationals. This generalizes for an arbitrary finite language the assumption in the case of random graphs with $p = n^{-\alpha}$ that α is irrational. This hypothesis easily implies that for any L -structures $A \leq_s B$ and expansions $A^+ \subseteq B^+$, $\delta(B^+/A^+) \neq 0$. This is a key property (see Remark 2.27).

We have the notions of \leq_s and \leq_i as a relation on members of \mathbf{K}_0 . The definition of δ on \mathbf{K}_0^+ induces corresponding

relations on \mathbf{K}_0^+ . Since we work directly with δ it is worthwhile recording the translation.

2.11 Definition. For $A^+ \subseteq B^+ \in \mathbf{K}_0^+$,

1. $A^+ \leq_s B^+$ if $\delta(B_1^+/A^+) > 0$ for every B_1^+ with $A^+ \subset B_1^+ \subseteq B^+$.
2. $A^+ \leq_i B^+$ if $\delta(B^+/B_1^+) < 0$ for every B_1^+ with $A^+ \subseteq B_1^+ \subset B^+$.
3. B^+ is a *primitive extension* of A^+ , if $\delta(B^+/A^+) > 0$ and $\delta(B^+/A_1^+) \leq 0$ for each A_1^+ with $A^+ \subset A_1^+ \subseteq B^+$.

Note that:

2.12 Remark. \mathbf{K}_0^+ satisfies axioms **A0-A6**. We will be using the following monotonicity properties which follow formally as in Section 1.

- 2.13 Lemma.**
1. If $A^+ \leq_i B^+$ and $A^+ \subseteq B_1^+ \subseteq B^+$ then $B_1^+ \leq_i B^+$.
 2. If $A^+ \leq_s B^+$ and $A^+ \subseteq B_1^+ \subseteq B^+$ then $A^+ \leq_s B_1^+$.

2.14 Remark. The exact phrasing of the following notions is extremely delicate. We consider a fixed pair of finite L^+ -structures, $A^+ \leq_s B^+$. The L -structures $\langle M_n : n < \omega \rangle$ naturally form a chain so an embedding f of A into M_n can naturally be regarded as a map of A into M_m for $m > n$. We are concerned with the properties of *extensions* of f . Thus, the immediately following definition of an L^+ -homomorphism extending f is agnostic concerning the preservation of relations on A .

2.15 Definition. Let $A^+ \subseteq B^+$. Let f be a 1 – 1 map from A into M_n , and let G^+ be an L^+ structure expanding M_n . Let T denote the range of f .

1. We say an injective map $g : B^+ \mapsto G^+$ which extends f is an L^+ -homomorphism relative to A is an if for any L^+ -relation R , and any $\bar{b} \in B$ but not in A , $B^+ \models R(\bar{b})$ implies $G^+ \models R(g\bar{b})$.
2. For any G^+ expanding M_n , and $W \subseteq M_n$ with $|W| \leq n$ we say $N(f, A^+, B^+, W) = k$ in G^+ , if

$$k = |\{g : B^+ \mapsto W \subseteq M_n \text{ is an } L^+\text{-homomorphism relative to } A \text{ and } g \supseteq f\}|.$$

If $W = M_n$ we omit it.

3. We say G^+ is in the event Y_f , which depends on a constant c_1 , if

$$n^{\delta(B^+/A^+)}(\log n)^{-(v(B/A)+1)} < N(f, A^+, B^+) < c_1 n^{\delta(B^+/A^+)}.$$
4. Let U denote the range of f . For each $S \subseteq M_n$ with $S \cap U = \emptyset$ and $|S| = |B - A|$, fix (if possible) an L -isomorphism g_S between B and US which extends f . (Since L -isomorphism just means 1 – 1 map, such S and g_S exist whenever $n \geq |B|$).
5. For each such S with fixed L -isomorphism g_S of B into M_n , let $X_{f,S}$ be a random variable such that $X_{f,S}(G^+)$ is

$$\begin{cases} 1 & \text{if } g_S \text{ maps } B \text{ onto } US \text{ and is an } L^+\text{-homomorphism relative to } A \text{ into } G^+ \\ 0 & \text{otherwise.} \end{cases}$$

For $W \subseteq M_n$ with $|W| \leq n$, let

$$X_{f,W} = \sum \{X_{f,S} : g_S : B \mapsto M_n \text{ and } g_S \supseteq f \text{ and } S \subseteq W\}.$$

If $W = M_n$, we write X_f for $X_{f,W}$.

- 2.16 Notation.** 1. For any property P of structures, in particular a first order property, the assertion, ‘for almost all sufficiently large M , $M \models P$ ’ (abbreviated a.a.) means ‘for every $\epsilon > 0$ there is an N such that if $n > N$, $P_n(\{G^+ | L = M_n : G^+ \models P\}) > 1 - \epsilon$ ’.
2. By an *indicator random variable* we mean one which takes values 0 or 1 and thus indicates a set.
3. We write $f \approx g$ if $f = O(g)$ and $g = O(f)$.

The next lemma expresses the key observation linking the probability with the dimension function δ .

2.17 Lemma. *For all sufficiently large n and all $f : A \mapsto M_n$, and any $W \subseteq M_n$ with $|W| \leq n$, the expectation*

$$E(X_{f,W}) \approx |W|^{v(B/A)} n^{-e(B^+/A^+)}.$$

Proof. The probability of an L -embedding of $B^+ | L$ into M^+ actually being an L^+ homomorphism is $\gamma(A^+, B^+) n^{-e(B^+/A^+)}$. The number of such embeddings has order of magnitude $|W|^{v(B/A)}$. Since expectation is additive this yields

$$E(X_{f,W}) \approx |W|^{v(B/A)} \gamma(A^+, B^+) n^{-e(B^+/A^+)}$$

The constant is absorbed by the approximation \approx . In particular, we have:

2.18 Remark. If $W = M_n$ this simplifies to

$$E(X_f) \approx n^{\delta(B^+/A^+)}.$$

In Theorem 2.30 we guarantee that our extensions are L^+ -isomorphisms (no new relations) rather than just L^+ -homomorphisms. Now we justify the restriction from \mathbf{K}_0^* to \mathbf{K}_0^+ .

2.19 Lemma. *If $\delta(B^+) < 0$ then a.a. there is no embedding of B^+ into G^+ .*

Proof. The expected number of copies of B^+ is $n^{\delta(B^+)}$. If $\delta(B^+) < 0$, this tends to 0.

2.20 Theorem. Fix $A^+ \subseteq B^+$ with $A^+ \leq_s B^+$.

Let V be the event: for all $f : A \mapsto M_n$, the event Y_f holds.
 Then, for some choice of c_1 (recall Y_f depends on c_1 .),

$$\lim_{n \rightarrow \infty} P_n(V) = 1.$$

Proof. By a straightforward induction, we can reduce to the case that B^+ is a primitive extension of A^+ . The proof of this case proceeds through several definitions and lemmas. Considering the definition of Y_f , one can see that we need to establish both lower and upper bounds. The lower bound argument proceeds as follows. Roughly speaking, for $f : A \mapsto M_n$ and $W \subseteq M_n$, we say (f, W) is bad if there is no extension of f to an L^+ -homomorphism (in the sense of Definition 2.15) of B^+ into W . In Lemma 2.22 we show that if W meets a cardinality requirement specified in Definition 2.21 then the probability that (f, W) is bad is less than $1/2$. By strengthening the requirements on W as in Definition 2.23 we improve the upper bound on the probability that (f, W) is bad in Lemma 2.24. Finally, taking into account the number of possible W 's, we complete the proof of the lower bound in Paragraph 2.25. After several preliminary definitions and lemmas we complete the proof of the upper bound in Paragraph 2.29.

2.21 Definition. We have fixed $A^+ \subseteq B^+$ with B^+ a primitive extension of A^+ . For G^+ an L^+ -expansion of M_n and $W \subseteq M_n$ and f an L -isomorphism of A into M_n , (f, W) is bad in G^+ if there is no g defined on $B - A$ into W such that $f \cup g$ defines an L^+ -homomorphism from B^+ into G^+ .

2.22 Lemma. There is a constant s such that for all sufficiently large n and any L -isomorphism $f : A \mapsto M_n$, if $|W|$ is the integer $m = m_s = \lceil sn^{e(B^+/A^+)/v(B/A)} \rceil$ then

$$P_n((f, W) \text{ is bad}) < 1/2.$$

Proof. Without serious loss of precision, $W \cap \text{rng } f = \emptyset$. We use the notation from Definition 2.15.

Now

$$P_n((f, W) \text{ bad}) = P_n(X_{f,W} = 0)$$

so we want to show that for all sufficiently large n ,

$$P_n(X_{f,W} = 0) < 1/2.$$

By Chebyshev's inequality,

$$P_n(X = 0) \leq \frac{\text{Var}(X)}{E(X)^2}.$$

By Lemma 2.17,

$$E(X) = E(X_{f,W}) \approx |W|^{v(B/A)} n^{-e(B^+/A^+)}.$$

Using the fact that $|W| = m_s$ this shows $E(X)$ is a polynomial of degree v in s , as the powers of n cancel. We will obtain the required result by showing $\text{Var}(X)$ is a polynomial of degree $2v - 1$ in s which implies that for sufficiently large s , $\frac{\text{Var}(X)}{E(X)^2} < 1/2$.

Now,

$$\text{Var}(X) = \sum_S \text{Var}(X_S) + \sum_{S \neq T} \text{Cov}(X_S, X_T)$$

where S, T range over subsets of M_n disjoint from the image of f .

An easy calculation shows that for any set of indicator random variables

$$E(\sum_S X_S) \geq \sum_S \text{Var}(X_S);$$

so we have

$$\text{Var}(X) \leq E(X) + \sum_{j=0}^v \sum_{|(S \cap T) - A| = j} \text{Cov}(X_S, X_T).$$

If $|(S \cap T) - A| = 0$ then $S \cap T = \emptyset$ and $\text{Cov}(X_S, X_T)$ is zero. Always, $\text{Cov}(X_S, X_T) \leq E(X_S X_T)$ which, since these are indicator random variables, is just $P_n(X_S X_T)$. Recall the definition of the probability measure from Definition 2.5; t is largest arity in the language.

$$\begin{aligned} P_n(X_S X_T) &= \prod_{m \leq t} \prod_{A \in [ST]^m} q_m((M|L^{\leq m})|A) \\ &\leq \prod_{m \leq t} \left(\frac{\prod_{A \in [S]^m} q_m((M|L^{\leq m})|A) \prod_{A \in [T]^m} q_m((M|L^{\leq m})|A)}{\prod_{A \in [S \cap T]^m} q_m((M|L^{\leq m})|A)} \right). \end{aligned}$$

Let B' be $g_s^{-1}(S \cap T)$. So $|B'| = j$. Abbreviating the notations from 2.8, let $c_S = c_T = \gamma(A^+, B^+)$ and $c_{B'} = \gamma(A^+, B')$. Similarly, let $e = e_S = e_T = e(B^+/A^+)$ and $u' = e(B'/A^+)$. With this notation we can rewrite the last inequality as

$$P_n(X_S X_T) \leq \frac{c_S c_T}{c_{B'}} n^{-(2e-u')}.$$

(The key to the inequality is that $2e - u'$ may undercount the number of relations on ST but this undercount can only overestimate the probability). $|(S \cap T) - A| = j$, $|ST - A| = 2v - j$ so $\chi_W(ST/A) \approx m^{2v-j}$.

If $\frac{u'}{j} > \frac{e}{v}$ then $\frac{e}{v} > \frac{e-u'}{v-j}$ which contradicts the fact that B^+ is a primitive extension of A^+ . So $u' \leq \frac{j e}{v}$. Thus,

$$\sum_{|(S \cap T) - A| = j} \text{Cov}(X_S, X_T) \leq m^{2v-j} n^{u'-2e} \leq (sn^{e/v})^{2v-j} n^{je/v-2e} = (s^v)^{2-j/v}.$$

(We can drop the constants in the last computation as $\frac{c_S c_T}{c_B} < 1$.) So,

$$\text{Var}(X) \leq E(X) + \sum_{j=1}^v (s^v)^{2-j/v} \leq E(X) + v s^{2v-1}.$$

Since $E(X)$ has degree v in s , this implies $\text{Var}(X) \leq E^2(X)/2$ for sufficiently large s and so

$$P_n((f, W) \text{ is bad}) < 1/2.$$

Now we want to modify the choice of W to get a better upper bound on the probability that (f, W) is bad.

2.23 Definition. Choose s by Lemma 2.22. As before, let $m_s = \lfloor sn^{e(B^+/A^+)/v(B/A)} \rfloor$. We say that $W \subseteq M_n$ is k -appropriate if $|W| = \lfloor 1 + km_s \ln n \rfloor$.

2.24 Lemma. For all sufficiently large n and any L -isomorphism $f : A \mapsto M_n$,

for sufficiently large k , if $W \subseteq n$ is k -appropriate,

$$P_n((f, W) \text{ is bad}) < \frac{n^{-|A|-1}}{2}.$$

Proof. Again, assume without loss of generality that $W \cap A = \emptyset$. Suppose W contains $k \ln n$ disjoint subsets W_i each with cardinality m_s . For (f, W) to be bad, each of the $k \ln n$ independent events that (f, W_i) is bad must occur and by Lemma 2.22 $P_n((f, W) \text{ is bad}) < 1/2$. Thus,

$$P_n((f, W) \text{ is bad}) < 2^{-k \ln n}.$$

But for all sufficiently large n and k ,

$$2^{-k \ln n} < \frac{n^{-|A|-1}}{2}$$

so we have the result.

We have shown that for each f , a.a. there is a W such that (f, W) is not bad. The next paragraph strengthens this assertion.

2.25 Proof of Lower Bound in Theorem 2.20. Fix k satisfying the conclusion of Lemma 2.24. For an L -isomorphism f of A into M_n , let the random variable $Z_f(G^+)$ be the number of k -appropriate $W \subseteq M_n$ (i.e. $|W| = w = [1 + km_s \ln n]$ with m from Definition 2.23) such that (f, W) is bad. Let γ denote the number of possible k -appropriate W . (The value of γ is not used in the first stage of the argument.) Then, $E(Z_f) < \gamma n^{-|A|-1}$. So, by Markov's inequality,

$$P_n(Z_f \geq \gamma/2) \leq 2E(Z_f)/\gamma < 2n^{-|A|-1}.$$

But then, since there are only $n^{|A|}$ choices for A , a.a. for each f at most half of the W are bad for f .

Let v denote $|B - A| = v(B/A)$. Each extension g of f to B is contained in at most $\binom{n-v}{w-v}$ k -appropriate W , since there are approximately $\binom{n-v}{w-v}$ choices for the elements which comprise $W - \text{rng } g$. So at most $|N(f, A^+, B^+)| \binom{n-v}{w-v}$ k -appropriate W contain an extension of f but at least $\gamma/2$ do. Now note that $\gamma = \binom{n}{w}$

Thus, a.a. for all f ,

$$|N(f, A^+, B^+)| \binom{n-v}{w-v} \geq \frac{1}{2} \binom{n}{w}.$$

Noting that $\binom{n-v}{w-v}$ is approximately $\binom{n}{w}(\frac{w}{n})^v$, we have

$$|N(f, A^+, B^+)| \geq n^{-e(B^+/A^+)}(n/w)^v/2.$$

Recalling that $w = [1 + km_s \ln n]$, this implies for every f , a.a.

$$|N(f, A^+, B^+)| > n^{\delta(B^+/A^+)}(2ks \ln n)^{-v}$$

which establishes the lower bound $n^{\delta(B^+/A^+)}(\ln n)^{-c}$ by taking $c = v + 1$.

2.26 Remark. The statement and proofs of of the probability analysis are based on the argument in [8]. The first author acknowledges discussions with Albert, Cherlin, Lachlan, and Laskowski on the details of the current argument, and supplemental remarks to the original paper by Spencer.

2.27 Remark. The irrationality hypothesis is necessary to make fruitful application of this result. If there exist $A \leq_s B$ with $\delta(B/A) = 0$ then the lower bound we have established is less than one rather than tending to infinity as n does. This destroys the argument of Theorem 2.30.

2.28 Remark. From Lemma 1.17 we have: Let $A^+ \leq_i C^+$. There exists a K such that a.a. for every embedding f of A^+ into an expansion G^+ of M_n , there are fewer than K L^+ -homomorphisms extending f from C^+ into G^+ .

2.29 Proof of upper bound in Theorem 2.20. Since B^+ is a primitive extension of A^+ , $(Ab)^+ \leq_i B^+$ for any $b \in B - A$. Thus, by Remark 2.28 there are fewer than K extensions g_i with any fixed image of b . The range of each extension g_i can intersect at most $K|B|^2$ other extensions so if $N(f, A^+, B^+) = s'$, there is a set of $s = s'/(K|B|^2 + 1)$ disjoint extensions. Let $p = |A|$, $v = v(B/A) = |B - A|$ and $e = e(B^+/A^+)$. For an appropriate constant $c < 1$, there are less than $cn^p(n^{vs}/s!)$ pairs of a function f taking A into M_n and a set of s extensions (disjoint over $\text{rng } f$) $\langle g_1, \dots, g_s \rangle$. The probability that each of the g_i is an L^+ -homomorphism is n^{-e} so the probability of such a pair of a function and s homomorphisms is at most $n^p(n^{vs}n^{-es})/s!$ By Stirling's formula, this is less than $\frac{n^p(2.72n^v)^s n^{-es}}{(2\pi)^{1/2} s^{s+1/2} (2.72)^{1/(12s+1)}}$ which is much less than 1 if $s \geq 3n^{\delta(B^+/A^+)}$. (Observe that $n^p(\frac{2.72}{3})^s$ tends to 0 as n tends to infinity.) Thus, a.a. $s \leq 3n^{\delta(B^+/A^+)}$.

a.a. for each f ,

$$N(f, A^+, B^+) = s' \leq 3(K|B|^2 + 1)n^{\delta(B^+/A^+)}$$

proving Theorem 2.20.

We now want to show that each of the axioms for semigenericity has limit probability 1. Roughly, the program is to show that for $A^+ \leq_s B^+$ and $f : A \mapsto M_n$, the number of extensions of f to 1 – 1-homomorphisms of B^+ is much greater than the number of such extensions which fail to witness the definition of semigenericity. Since there are a bounded number of types of failure, it suffices to check each type separately as we do in the following argument.

In general, embeddings $f : A^+ \mapsto M_n$ and $\hat{f} : B^+ \mapsto M_n$ fail to witness semigenericity of G^+ if

1. $\text{cl}_{G^+}^m(\hat{f}B) \neq \hat{f}B \cup \text{cl}_{G^+}^m(fA)$ or
2. $\text{cl}_{G^+}^m(fA)$ and $\hat{f}B$ are not freely joined over fA in G^+ .

In considering ϕ_{A^+, B^+, C^+}^m , we are fixing on C^+ as a specific candidate for the isomorphism type of $\text{cl}_{G^+}^m(fA)$.

2.30 Theorem. *If $A^+ \leq_s B^+$ and $A^+ \leq_i C^+$ with $|C^+| < m$ then*

$$\lim_{n \rightarrow \infty} P_n(\phi_{A^+, B^+, C^+}^m) = 1.$$

Proof. For any f mapping C into M_n , and a 1 – 1 homomorphism f' extending f to E , (G^+, f') fails as a witness for C and f if

1. f' is not an L^+ -isomorphism or
2. $\text{cl}_{G^+}^m(f'B) \neq f'B \cup fC$ or
3. fC and $f'B$ are not freely joined over fA in G^+ .

Note that $C^+ \leq_s C^+ \otimes_{A^+} B^+$ and, letting E^+ denote $C^+ \otimes_{A^+} B^+$, $\delta(B^+/A^+) = \delta(E^+/C^+)$. By Theorem 2.20, more specifically Paragraph 2.25, a.a. for each f ,

$$|N(f, C^+, E^+)| > n^{\delta(B^+/A^+)}(\log n)^{-c}$$

where $c = v(B^+/A^+) + 1$. For conditions i) and iii) consider any F^+ which is an expansion of E^+ by adding additional relations. Then $\delta(F^+/A^+) = v(E/A) - e(F^+/A^+)$ and $e(F^+/A^+) > e(E^+/A^+)$. By Theorem 2.20

$$N(f, C^+, F^+) < c_1 n^{\delta(F^+/A^+)} < n^{\delta(E^+/A^+)} (\log n)^{-c}.$$

For condition ii) for any $D^+ \in \mathcal{D}_{B^+, C^+}^m$,

$$\delta(D^+B^+/C^+) < \delta(B^+C^+/C^+) = \delta(B^+/A^+).$$

If C^+ is not strong in D^+B^+ then by Lemma 2.28, $N(f', C^+, D^+B^+) < K$. If $C^+ \leq_s D^+B^+$ then by Theorem 2.20, more specifically Paragraph 2.29, a.a.

$$N(f, C^+, D^+B^+) < c_1 n^{\delta(D^+B^+/C^+)} < n^{\delta(B^+C^+/C^+)} (\log n)^{-c}.$$

Now the number of isomorphism types of extensions C that have failures f' is bounded in terms of the cardinality of A^+ , B^+ , and m ; it does not depend on n . If this number is L , the total number of failures of any sort is less than $Ln^{\delta(E^+/A^+)} (\log n)^{-c}$. Thus, the probability that for each f , one of the extensions of f witnesses ϕ_{A^+, B^+, C^+}^m tends to one as required.

2.31 Lemma. *For every m , a.a. $\text{cl}_{L^+}^m(\emptyset) = \emptyset$.*

Proof. $A^+ \subseteq \text{cl}_{L^+}^m(\emptyset)$ just if $\delta(A^+) < 0$. But in passing from \mathbf{K}_0^* to \mathbf{K}_0^+ (cf. Lemma 2.19, we have forbidden such A .

We collect our results in the following theorem which requires the definition of two theories.

2.32 Notation. Let L contain only the equality symbol and let L^+ be an arbitrary finite relational language containing L . Suppose probabilities are defined on finite L^+ structures as in Definition 2.5 with the α_i and 1 linearly independent over the rationals. By T^α , the almost sure theory of random L^+ -structures we mean the collection of L^+ -sentences which have limit probability 1. Recall that T_α is the theory of the generic structures for \mathbf{K}_α (Definition 1.6) whose existence is guaranteed by Theorem 1.34.

A theory T has the finite model property if every theorem of T has a finite model.

2.33 Theorem. *Under the hypotheses in Notation 2.32, T^α , the almost sure theory of random L^+ -structures is the same as the theory T_α of the \mathbf{K}_α -generic model. This theory is complete, stable, and nearly model complete. Moreover, it has the finite model property and has only infinite models so is not finitely axiomatizable.*

Proof. By Theorem 2.30 and the choice of \mathbf{K}_0^+ , every model of T^α is $(\mathbf{K}_\alpha, \leq_\alpha)$ -semigeneric. By Corollary 1.31, T^α is nearly model complete. By Corollary 1.32 and Lemma 2.31, T^α is complete. Since the generic model for \mathbf{K}_α is semigeneric, $T^\alpha = T_\alpha$. [1] shows that T_α is stable.

Since each theorem of T^α has limit probability 1, for arbitrarily large n , there is nonzero probability that there is a model of size n . Thus, T^α has the finite model property.

2.34 Remark. The major novelty of this result is the identification of the two theories, thereby obtaining the stability of T^α and the non-finite axiomatizability of T_α . The notion of near model completeness specifies the precise degree of quantifier elimination in T^α . In addition, we have extended the 0 – 1 law from a language with a single binary relation to an arbitrary finite relational language.

References

- [1] J.T. Baldwin and Niandong Shi. Stable generic structures. *Annals of Pure and Applied Logic*, 199x. to appear.
- [2] A. Baudisch. A new \aleph_1 -categorical pure group. 1992.
- [3] E. Hrushovski. A stable \aleph_0 -categorical pseudoplane. preprint, 1988.
- [4] D.W. Kueker and C. Laskowski. On generic structures. *Notre Dame Journal of Formal Logic*, 33:175–183, 1992.
- [5] J. Lynch. Probabilities of sentences about very sparse random graphs. *Random Structures and Algorithms*, 3:33–53, 1992.
- [6] S. Shelah. 0-1 laws. preprint 550, 199?

- [7] S. Shelah. Zero-one laws with probability varying with decaying distance. Shelah 467, 199x.
- [8] S. Shelah and J. Spencer. Zero-one laws for sparse random graphs. *Journal of A.M.S.*, 1:97–115, 1988.
- [9] F. Wagner. Relational structures and dimensions. In *Automorphisms of first order structures*, pages 153–181. Clarendon Press, Oxford, 1994.