

APPLICATIONS OF PCF THEORY  
SH589

SAHARON SHELAH

Institute of Mathematics  
The Hebrew University  
Jerusalem, Israel

Rutgers University  
Department of Mathematics  
New Brunswick, NJ USA

ABSTRACT. We deal with several pcf problems; we characterize another version of exponentiation: number of  $\kappa$ -branches in a tree with  $\lambda$  nodes, deal with existence of independent sets in stable theories, possible cardinalities of ultraproducts and the depth of ultraproducts of Boolean Algebras<sup>1</sup>. Also we give cardinal invariants for each  $\lambda$  with a pcf restriction and investigate further  $T_D(f)$ . The sections can be read independently, although there are some minor dependencies.

modified:2005-07-06

---

Partially supported by the basic research fund, Israeli Academy  
This version came from sections of Sh580  
I thank Alice Leonhardt for the excellent typing  
First revision after the Journal [?] - 04/Oct/29  
Latest Revision - 05/July/4  
Pub. No. 589  
Saharon references to 513:p.32,34, see §2 4.8

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

## ANNOTATED CONTENT

§1  $T_D$  via true cofinality

[Assume  $D$  is a filter on  $\kappa$ ,  $\mu = \text{cf}(\mu) > 2^\kappa$ ,  $f \in {}^\kappa\text{Ord}$ , and:  $D$  is  $\aleph_1$ -complete or  $(\forall \sigma < \mu)(\sigma^{\aleph_0} < \mu)$ . We prove that if  $T_D(f) \geq \mu$  (i.e. there are  $f_\alpha <_D f$  for  $\alpha < \mu$  such that  $f_\alpha \neq_D f_\beta$  for  $\alpha < \beta < \mu$ ) then for some  $A \in D^+$  and regular  $\lambda_i \in (2^\kappa, f(i)]$  we have:  $\mu$  is the true cofinality of  $\prod_{i < \kappa} \lambda_i / (D + A)$ . We end summing up conditions equivalent to  $T_{D+A}(f) \geq \mu$  for some  $A \in A^+$ .]

## §2 The tree revised power

[We characterize more natural cardinal functions using pcf. The main one is  $\lambda^{\kappa, \text{tr}}$ , the supremum on the number of  $\kappa$ -branches of trees with  $\lambda$  nodes, where  $\kappa$  is regular uncountable. If  $\lambda > \kappa^{\kappa, \text{tr}}$  it is the supremum on  $\max \text{pcf}\{\theta_\zeta : \zeta < \kappa\}$  for an increasing sequence  $\langle \theta_\zeta : \zeta < \kappa \rangle$  of regular cardinals with  $\zeta < \kappa \Rightarrow \lambda \geq \max \text{pcf}\{\theta_\varepsilon : \varepsilon < \zeta\}$ .]

## §3 On the depth behaviour of ultraproducts

[We deal with a problem of Monk on the depth of ultraproducts of Boolean algebras; this continues [Sh 506, §3]. We try to characterize for a filter  $D$  on  $\kappa$  and  $\lambda_i = \text{cf}(\lambda_i) > 2^\kappa$ , and  $\mu = \text{cf}(\mu)$ , when does  $(\forall i < \kappa)[\lambda_i \leq \text{Depth}^+(B_i)] \Rightarrow \mu < \text{Depth}^+(\prod_{i < \kappa} B_i / D)$  (where  $\text{Depth}^+(B) = \cup\{\mu^+ : \text{in } B \text{ there is an increasing sequence of length } \mu\}$ ). When  $D$  is  $\aleph_1$ -complete or  $(\forall \sigma < \mu)[\sigma^{\aleph_0} < \mu]$  the characterization is reasonable: for some  $A \in D^+$  and  $\lambda'_i = \text{cf}(\lambda'_i) < \lambda_i$  we have  $\mu = \text{tcf} \prod_{i < \kappa} \lambda'_i / (D + A)$ . We then proceed to look at  $\text{Depth}_h^{(+)}$  (closing under homomorphic images), and with more work succeed. We use results from §1.]

## §4 On the existence of independent sets for stable theories

[Bays [Bays Ph.D.] has continued work in [Sh:c] on existence of independent sets (in the sense of non-forking) for stable theories. We connect those problems to pcf and shed some light. Note that the combinatorial Claim 4.1 continues [Sh 430, §3].]

## §5 Cardinal invariants for general cardinals: restriction on the depth

[We show that some (natural) cardinal invariants defined for any regular  $\lambda (> \aleph_0)$ , as functions of  $\lambda$  satisfies inequalities coming from pcf (more accurately

norms for  $\aleph_1$ -complete filters). They are variants of depth, supremum of length of sequences from  ${}^\lambda\lambda$  (increasing in a suitable sense) and also the supremum of sizes of  $\lambda$ -MAD families. Contrast this with Cummings Shelah [CuSh 541]. Also we connect pcf and the ideal  $I[\lambda]$ ; see 5.20.]

§6 The class of cardinal ultraproducts mod  $D$

[Let  $D$  be an ultrafilter on  $\kappa$  and let  $\text{reg}(D) = \text{Min}\{\theta : \text{the filter } D \text{ is not } \theta\text{-regular}\}$ , so  $\text{reg}(D)$  is regular itself. We prove that if  $\mu = \mu^{\text{reg}(D)} + 2^\kappa$  then  $\mu$  can be represented as  $|\prod_{i < \kappa} \lambda_i / D|$ , and for suitable  $\mu$ 's get  $\mu$ -like such ultraproducts.]

We thank Todd Eisworth for doing much in corrections and improving presentation, and Andres Villaveces similarly for §4.

§1  $T_D$  VIA TRUE COFINALITY

We improve here results of [Sh 506, §3] but do not depend on it. See more related things in §6. Our main result is 1.6, which we will use in §3 in our analysis of ultraproducts of Boolean Algebras<sup>2</sup>.

**1.1 Claim.** 1) *Assume*

- (a)  $J$  is an  $\aleph_1$ -complete ideal on  $\kappa$
- (b)  $f \in {}^\kappa\text{Ord}$ , each  $f(i)$  an infinite ordinal  $\geq \mu$
- (c)  $T_J^2(f) \geq \lambda = \text{cf}(\lambda) > \mu \geq \kappa$  (see 1.2(1) below)
- (d)  $\mu = 2^\kappa$ , or at least
  - (d)<sup>-</sup>(i) if  $\mathfrak{a} \subseteq \text{Reg}$ , and
 
$$(\forall \theta \in \mathfrak{a})(\mu \leq \theta < \lambda \ \& \ \mu \leq \theta < \sup_{i < \kappa} f(i))$$
 and  $|\mathfrak{a}| \leq \kappa$ , then  $|\text{pcf}(\mathfrak{a})| \leq \mu$
  - (ii)  $|\mu^\kappa/J| < \lambda$
  - (iii)  $2^\kappa < \lambda$ .

Then for some  $A \in J^+$  and  $\bar{\lambda} = \langle \lambda_i : i \in A \rangle$  such that  $\mu \leq \lambda_i = \text{cf}(\lambda_i) \leq f(i)$  we have  $\prod_{i \in A} \lambda_i / (J \upharpoonright A)$  has true cofinality  $\lambda$ .

2) If we can clause (e) below that we can add  $i \in A \Rightarrow \lambda_i = f(i)$

- (e) if  $g \in {}^\kappa\text{Ord}$  and  $g <_J f$  then  $T_J^2(g) < \lambda$  or just
- (e)' if  $A \in J^+$  then  $\prod_{i \in A} f(i) / (J \upharpoonright A)$  is not  $\lambda$ -directed.

1.2 Remark. 1) Remember  $T_J^2(f) = \text{Min}\{|F| : F \subseteq \prod_{i < \kappa} f(i)$  and for every

$g \in \prod_{i < \kappa} f(i)$  for some  $g' \in F$  we have  $\neg(g \neq_J g')$ \}. See [Sh 506, §3] on the relationship of relatives of this definition; they agree when  $> 2^\kappa$ . The inverse of the claim is immediate, i.e., the conclusion implies that  $\lambda \leq T_J^2(f)$ .

2) If  $A_1 = \{i < \kappa : f(i) \geq \lambda\} \in J^+$  then the conclusion is immediate, with  $\lambda_i = \lambda$ .

3) Note if  $A_2 = \{i < \kappa : f(i) < (2^\kappa)^+\} \in J^+$  then  $T_J^2(f) \leq 2^\kappa$ . If in addition

<sup>2</sup>Claim 1.1 was revised June 2004

$\kappa \setminus A_2 \in J$  then any  $\lambda$  satisfying the conclusion satisfies  $\lambda \leq 2^\kappa$ .

4) We can omit the assumption clause  $(d)^-(iii)$  and weaken (here and in 2.6) the assumption “ $|\mu^\kappa/J| < \lambda$ ” (in clause  $(d)^-$ ) and just ask:

$\bigoplus_{J,\mu,\lambda}$  there is  $F \subseteq {}^\kappa\mu$  of cardinality  $< \lambda$  such that for every  $g \in {}^\kappa\mu$  we can find  $F' \subseteq F$  of cardinality  $\leq \mu$  such that for every  $A \in J^+$  for some  $f \in F'$  we have  $\{i \in A : g(i) = f(i)\} \in J^+$ , or even

$\bigoplus_{J,\mu,\lambda}^-$  we require the above only for all  $g \in G$ , where  $G \subseteq {}^\kappa\mu$  has cardinality  $< \lambda$  and: if  $\langle \theta_i : i < \kappa \rangle$  is a sequence of regulars in  $[\aleph_0, \mu]$  and  $g' \in \prod_{i < \kappa} \theta_i$  then for some  $g'' \in G$  we have  $g' <_J g'' <_J \langle \theta_i : i < \kappa \rangle$ .

Considering  $(d)^-(iii)$  in the proof we weaken  $g_n \upharpoonright A \in N$  for some  $g', A' \subseteq \kappa$  from  $g_n \upharpoonright A =_J g' \upharpoonright A'$ .

5) Also in 1.6 and 1.7 we can replace the assumption  $\lambda > 2^\kappa$  by the existence of a  $\mu$  satisfying  $\lambda > \mu \geq \kappa$  such that  $(d)^-$  as weakened above holds.

6) Note that we do not ask  $(\forall \alpha < \lambda)[|\alpha|^{< \text{reg}(J)} < \lambda]$ .

7) Of course, we can apply the claim to  $J \upharpoonright A$  for every  $A \in J^+$  hence  $\{A/J : A \in J^+, \text{ and for some } \bar{\lambda} = \langle \lambda_i : i \in A \rangle \text{ such that } \mu \leq \lambda_i = \text{cf}(\lambda_i) \leq f(i) \text{ we have } \prod_{i \in A} \lambda_i / (J \upharpoonright A) \text{ has true cofinality } \lambda\}$  is dense in the Boolean Algebra  $\mathcal{P}(\kappa)/J$ .

*1.3 Remark.* The changes in the proof of 1.1 below required for weakening in 1.1 the clause  $|\mu^\kappa/J| < \lambda$  to  $\bigoplus_{J,\mu,\lambda}^-$  from 1.2(4) are as follows.

As  $J, \mu, \lambda \in N$  there are  $F \subseteq {}^\kappa\mu, G \subseteq {}^\kappa\mu$  as required in  $\bigoplus_{J,\mu,\lambda}^-$  belonging to  $N$  (hence  $\subseteq N$ ). After choosing  $g^{n,1}$  and  $B_n$  apply the assumption on  $G$  to  $g^{n,3} \in {}^\kappa\mu$  when  $g^{n,3} \upharpoonright B_n = (g^{n,2} \upharpoonright B_n)$  and  $g^{n,3} \upharpoonright (\kappa \setminus B_n)$  is constantly zero and  $\bar{\theta} = \langle \theta_i : i < \kappa \rangle$  where  $\theta_i = \text{cf}(g_n(i))$  if  $i \in B_n$  and  $\theta_i = \aleph_0$  if  $i \in \kappa \setminus B_n$ .

So we get some  $g^{n,4} \in G$  such that  $g^{n,3} <_J g^{n,4} <_J \langle \theta_i : i < \kappa \rangle$ . As  $G \in N$ ,  $|G| < \lambda$  clearly  $G \subseteq N$  hence  $g^{n,4} \in G$ . Let  $F'_n$  be a subset of  $F$  of cardinality  $\leq \mu$  such that: for every  $A \in J^+$  for some  $f \in F'_n$  we have  $\{i \in A : g^{n,4}(i) = f(i)\} \in J^+$ .

Now continue as there but defining  $g_{n+1}$  use  $g^{n,4}$  instead  $g^{n,3}$  and choose  $\mathcal{P}_{n+1}^1$  as

$$\left\{ \{i < \kappa : g^{n,4}(i) = f(i)\} : f \in F'_n \right\}.$$

The rest is straight.

Remember

1.4 *Fact.* Assume

- (a)  $N \prec (\mathcal{H}(\chi), \in, <_\chi^*)$  and  $\mu < \lambda < \chi$  and  $\{\mu, \lambda\} \in N$ ,
- (b)  $N \cap \lambda$  is an ordinal,
- (c)  $i^* \leq \mu$ , and for  $i < i^*$  we have  $\mathfrak{a}_i \subseteq \text{Reg} \setminus \mu^+$ ,  $|\mathfrak{a}_i| \leq \mu$ ,  $\theta_i \in \text{pcf}(\mathfrak{a}_i) \cap \lambda$  and  $(\mathfrak{a}_i, \theta_i) \in N$ , and let  $\mathfrak{a} = \bigcup_{i < i^*} \mathfrak{a}_i$ .

Then

- (\*) for every  $g \in \Pi \mathfrak{a}$  there is  $f$  such that:
  - ( $\alpha$ )  $g < f \in \Pi \mathfrak{a}$
  - ( $\beta$ )  $f \upharpoonright \mathfrak{b}_{\theta_i}[\mathfrak{a}_i] \in N$ , and if  $\theta_i = \max \text{pcf}(\mathfrak{a}_i)$  we have  $f \upharpoonright \mathfrak{a}_i \in N$ .

*Proof.* By [Sh:g, Ch.II,3.4] or [Sh:g, VIII,§1].

*Proof of 1.1.* 1) We can find  $f' \leq_J f$  such that  $f' \in \prod_{i < \kappa} f(i) + 1$  which satisfies the

requirements on  $f$  and clause (e), so it is enough to prove part (2).

2) Note that assuming  $2^\kappa < \lambda$  slightly simplifies the proof, as then we can demand  $g_{A,n} = g_n \upharpoonright A$ . Assume toward contradiction that the conclusion fails. Let  $\chi$  be large enough, and let  $N$  be an elementary submodel of  $(\mathcal{H}(\chi), \in, <_\chi^*)$  of cardinality  $< \lambda$  such that  $\{f, \lambda, \mu\}$  belongs to  $N$  and  $N \cap \lambda$  is an ordinal and if we assume only clause (d)<sup>3</sup> then<sup>3</sup>

- ☒ for every  $f \in {}^\kappa \mu$  there is  $g \in N \cap {}^\kappa \mu$  such that  $f = g \text{ mod } J$  (if  $J \in N$  this is immediate).

Let  $\mathcal{F} = \left( \prod_{i < \kappa} f(i) \right) \cap N$  so  $\mathcal{F}$  cannot exemplify that  $T_J^2(f) \leq |\mathcal{F}| (< \lambda)$ , thus giving a contradiction.

As  $\mathcal{F} \subseteq \prod_{i < \kappa} f(i)$ , there is  $g$  witnessing “ $\mathcal{F}$  does not exemplify  $T_J^2(f) \leq |\mathcal{F}|$ ” hence

- (\*)  $g \in \prod_{i < \kappa} f(i)$  is such that for every  $g' \in \mathcal{F}$  we have  $(g \neq_J g')$  i.e.
 
$$\{i < \kappa : g'(i) = g(i)\} \in J.$$

---

<sup>3</sup>note we did not forget to ask  $J \in N$ , we just want to help reading this as a proof of 1.5, too for the case  $2^{|J|} \geq \lambda$ , so there  $J'$  does not necessarily belong to  $N$ .

We now define by induction on  $n < \omega$  the function  $g_n$  and the family  $\mathcal{P}_n$  and ideal  $J_n$  on  $\kappa$  such that:

- ⊠ (i)  $g_0 = f, g_n \in {}^\kappa\text{Ord}$ , and  $g_{n+1} \leq g_n$
- (ii)  $g_{n+1} < g_n \text{ mod } J_{n+1}, J_0 = J, J_n \subseteq J_{n+1}$
- (iii)(α)  $\mathcal{P}_n$  is a family of  $\leq \mu$  members of  $J^+$
- (β)  $A \in \mathcal{P}_n \wedge B \in J_n \Rightarrow A \cap B \in J$
- (γ) if  $n = m + 1$  and  $A \in \mathcal{P}_n$  then for some  $A' \in \mathcal{P}_m$  we have  $A \subseteq A'$
- (iv) if  $A \in \mathcal{P}_n$  then  $g_{A,n} =: g_n \upharpoonright A \in N$  hence  $A \in N$  but if  $2^\kappa \geq \lambda$  we just assume that for some  $g_{A,n} \in \prod_{i \in A} f(i)$  we have  $g_{A,n} = g_n \upharpoonright A \text{ mod } J$  and  $g_{A,n} \in N$  hence  $A \in N$
- (v)  $\mathcal{P}_0 = \{\kappa\}$  and  $J_0 = J$
- (vi) if  $A \in \mathcal{P}_n$  and  $B \subseteq A$  and  $B \in J_{n+1}^+$  and  $(\forall B' \in J_n)(B \cap B' \in J)$  then for some  $A' \in \mathcal{P}_{n+1}$  we have  $A' \subseteq A$  &  $A' \cap B \in J_{n+1}^+$
- (vii)  $g < g_n \text{ mod } J_n$
- (viii)(α)  $g(i) \leq g_n(i)$
- (β)  $g(i) < g_n(i) \Rightarrow g_{n+1}(i) < g_n(i)$
- (γ)  $g_{n+1}(i) \leq g_n(i)$
- (δ)  $\{i : g_{n+1}(i) = g_n(i)\} \in J_n$
- (ε) if  $i \in A \in \mathcal{P}_{n+1}$  then  $g_{n+1}$  then  $g(i) < g_n(i)$ .
- (ix)  $J_{n+1} \equiv \{A_1 \cup A_2 : A_1 \in J_n, A_2 \subseteq \kappa \text{ and if } A_2 \notin J_n \text{ then } \prod_{i \in A_2} g_{n+1}(i) / (J_n \upharpoonright A_2) \text{ is } \lambda\text{-directed}\}$ , note that:  $J_1 = J$  by the assumption toward contradiction.

Why is carrying the definition enough?

Let

- (\*)<sub>1</sub>  $J_n^* := \{A : A \in J_n \text{ or } A \in J_n^+ \text{ and } \lambda > \text{cf}(\prod_{i \in A} g_n \upharpoonright (J \upharpoonright A))\}$  so
- (\*)<sub>2</sub>  $J_{n+1} = \{A \subseteq \kappa : \text{if } B \in (J_{n+1}^*)^+ \text{ then } A \cap B \in J_n\}$
- (\*)<sub>3</sub> for  $n > 0$ , let  $g_\alpha^n \in \prod_{i < \kappa} (g_n(i) + 1)$  for  $\alpha < \lambda$  be  $<_{J_n^*}$ -increasing.

[Why it exists? By  $(*)_1 + (*)_2$  and the pcf theorem.]

For every  $i < \kappa$  let  $n(i)$  be the minimal  $n$  such that  $g_n(i) \leq g(i)$  (equivalently  $g_n(i) = g(i)$ ) and let  $g_\alpha^*(i) \in g_1(i)$  code  $\langle g_\alpha^n(i) : n \in [1, n(\alpha)] \rangle$ .

$$(*)_4 \quad g_\alpha^* \in \prod_{i < \kappa} g_1(i) \text{ for } \alpha < \lambda$$

$$(*)_5 \quad \text{if } \alpha < \beta < \lambda \text{ then } B =: \{i < \kappa : g_\alpha^*(i) = g_\beta^*(i)\} \in J.$$

[Why? If not, then for some  $n(*)$  we have  $B_* = \{i \in B : n(i) = n(*)\} \in J^+$ . Now we try to choose  $A_m \in \mathcal{P}_m$  by induction on  $m \leq n(*) + 2$  such that  $B_* \cap A_m \notin J$  and  $m = k + 1 \Rightarrow A_m \subseteq A_k$ . For  $m = 0$  this is possible as we can choose  $A_m = \kappa$  (see  $\boxtimes(v)$ ) so  $B_* \cap A_m = B_* \notin J$ .

Assume  $m < n(*)$  and  $A_0, \dots, A_m$  has been defined and we cannot choose  $A_{m+1}$  by the choice of  $J_{m+1}$ , see  $\boxtimes(ix)$  our inability to choose  $A_{m+1}$  it follows that  $B_* \cap A_m \in J_{n+1}$ . However, by  $\boxtimes$  we know that  $\mathcal{P}(A_m) \cap J_m = \mathcal{P}(A_m) \cap J$  hence  $\mathcal{P}(A_m) \cap J_m^* = \mathcal{P}(A_m) \cap J_m^*$  hence  $\langle g_\gamma^* \upharpoonright A_m : \gamma < \lambda \rangle$  to  $\langle J_m^* \upharpoonright A_m \rangle$ -increasing hence  $\langle J_m \upharpoonright A_m \rangle$ -increasing hence by the choice of  $B^*$  we have  $B^* \cap A_m \in J_m$ , contradictin. So we can choose  $A_{m+1}$ . So we have chosen  $A_0, \dots, A_{n(*)+2}$  as required.

But  $i \in A \in \mathcal{P}_m \wedge i \in A \Rightarrow n(i) \geq m$ , by  $\boxtimes(viii)(\varepsilon)$  but  $B_* \cap A_{n(*)+2} \neq \emptyset$ , so there is  $i \in B_* \cap A_{n(*)+2}$ , contradiction.]

Together

$$(*)_6 \quad \langle g_\alpha^* : \alpha < \lambda \rangle \text{ witness } T_J(g_1) \geq \lambda.$$

But  $f = g_0$  by  $\boxtimes(i)$ ,  $g_0 < g_1 \text{ mod } J_1$  by  $\boxtimes(ii)$  and  $J_1 = J_0, \boxtimes(ix)$ , so

$$(*)_7 \quad g_1 < f \text{ mod } J.$$

However  $(*)_6 + (*)_7$  contradict clause (e) of the assumption.

Carrying the induction  $\boxtimes$ : Also the case  $n = 0$  is easy by (i) + (v). So assume we have  $g_n, \mathcal{P}_n, J_n$  and we shall define  $g_{n+1}, \mathcal{P}_{n+1}, J_{n+1}$ . In  $N$  there is a two-place function  $\mathbf{e}$ , written  $\mathbf{e}_\delta(i)$  such that  $\mathbf{e}_\delta(i)$  is defined iff  $\delta \in \{\alpha : \alpha \text{ a non-zero ordinal } \leq \sup f(i)\}$ , and  $i < \text{cf}(\delta)$ , and if  $\delta$  is a limit ordinal, then  $\langle \mathbf{e}_\delta(i) : i < \text{cf}(\delta) \rangle$  is strictly increasing with limit  $\delta$  and  $\mathbf{e}_{\alpha+1}(0) = \alpha$ ; of course,  $\text{Dom}(\mathbf{e}_{\alpha+1}) = \{0\}$ .

We also know by assumption (d) or (d)<sup>-</sup>(i) that

$$\otimes \quad \text{for every } A \in \mathcal{P}_n \text{ letting } \mathbf{a}_A^n =: \{\text{cf}(g_{A,n}(i)) : i \in A\} \setminus \mu^+, \text{ the set } \text{pcf}(\mathbf{a}_A^n) \text{ has at most } \mu \text{ members.}$$

So  $\mathcal{Y} =: \{(A, \mathbf{a}_A^n, \theta) : A \in \mathcal{P}_n \text{ and } \theta \in \lambda \cap \text{pcf}(\mathbf{a}_A^n)\}$  has at most  $|\mathcal{P}_n| \times \mu \leq \mu \times \mu = \mu$  members (as  $|\mathcal{P}_n| \leq \mu$  and  $|\text{pcf}(\mathbf{a}_A^n)| \leq \mu$  by  $\otimes$  above) so let



$\{(A_\varepsilon^n, \mathfrak{a}_\varepsilon^n, \theta_\varepsilon^n) : \varepsilon < \varepsilon_n^*\}$  list them with  $\varepsilon_n^* \leq \mu$ . Clearly  $\mathfrak{a}_\varepsilon^n \in N$  (as  $g_{A,n} \upharpoonright A_\varepsilon^n \in N$ ), and since  $\mu + 1 \subseteq N$  and  $|\text{pcf}(\mathfrak{a}_\varepsilon^n)| \leq \mu$ , we have  $\mathcal{Y} \subseteq N$ . For each  $\varepsilon < \varepsilon_n^*$  we define  $h_\varepsilon^n \in \Pi \mathfrak{a}_\varepsilon^n$  by:

$$h_\varepsilon^n(\theta) = \text{Min} \left\{ \zeta < \theta : \text{if } i \in A_\varepsilon^n, g(i) < g_n(i), \text{ and} \right. \\ \left. \theta = \text{cf}(g_n(i)) \text{ then } g(i) < \mathbf{e}_{g_n(i)}(\zeta) \right\}.$$

[Why is  $h_\varepsilon^n$  well defined? The number of possible  $i$ 's is  $\leq |A_\varepsilon^n| \leq \kappa \leq \mu$ , for each relevant  $i$ , every  $\zeta < \theta$  large enough is OK as  $\langle \mathbf{e}_{g_n(i)}(\zeta) : \zeta < \theta \rangle$  is increasing continuous with limit  $g_n(i)$ . Lastly,  $\theta = \text{cf}(\theta) > \mu$  (by the choice of  $\mathfrak{a}_\varepsilon^n$ ) so all the demands together hold for every large enough  $\zeta < \theta$ .]

Let  $\mathfrak{a}_n = \bigcup_{\varepsilon < \varepsilon_n^*} \mathfrak{a}_\varepsilon^n$  and let  $h_n \in \Pi \mathfrak{a}_n$  be defined by  $h_n(\theta) = \sup\{h_\varepsilon^n(\theta) : \varepsilon < \varepsilon_n^* \text{ and } \theta \in \mathfrak{a}_\varepsilon^n\}$ , it is well defined by the argument above. So by 1.4 there is a function  $g^{n,1} \in \Pi \mathfrak{a}_n$  such that:

- ( $\alpha$ )  $h_n < g^{n,1}$
- ( $\beta$ )  $g^{n,1} \upharpoonright \mathfrak{b}_{\theta_\varepsilon^n}[\mathfrak{a}_\varepsilon^n] \in N$  (and  $\theta_\varepsilon^n = \max \text{pcf}(\mathfrak{a}_\varepsilon^n) \Rightarrow \mathfrak{b}_{\theta_\varepsilon^n}[\mathfrak{a}_\varepsilon^n] = \mathfrak{a}_\varepsilon^n$ ).

Also we can define  $g^{n,2} \in {}^\kappa \text{Ord}$  by:

$$g^{n,2}(i) = \text{Min}\{\zeta < \text{cf}(g_n(i)) : \mathbf{e}_{g_n(i)}(\zeta) \geq g(i)\}.$$

So letting  $B_n = \{i : 1 \leq \text{cf}(g_n(i)) \leq \mu\}$  clearly  $g^{n,2} \upharpoonright B_n \in {}^{B_n} \mu$ . Now if assumption (d) holds, then  $\mu^\kappa / J < \lambda$ , hence  $\mu^\kappa \subseteq N$  so we can find  $g^{n,3} \in N$  such that  $g^{n,2} = g^{n,3} \text{ mod } (J + (\kappa \setminus B_n))$ ; if assumption (d) fails we still can get such  $g^{n,3}$  by  $\boxtimes$  above. Lastly, we define  $g_{n+1} \in {}^\kappa \text{Ord}$ :

$$g_{n+1}(i) = \begin{cases} \mathbf{e}_{g_n(i)}(g^{n,1}(\text{cf}(g_n(i)))) & \underline{\text{if}} \quad \text{cf}(g_n(i)) \in \mathfrak{a}_n \text{ and } g_n(i) > g(i) \\ \mathbf{e}_{g_n(i)}(g^{n,3}(\text{cf}(g_n(i)))) & \underline{\text{if}} \quad \text{cf}(g_n(i)) \in [1, \mu] \text{ and } g_n(i) > g(i) \\ g_n(i) & \underline{\text{if}} \quad g(i) = g_n(i) \text{ or } \mu < \text{cf}(g_n(i)) \notin \mathfrak{a}_n \end{cases}$$

and  $\mathcal{P}_{n+1} = (\mathcal{P}_{n+1}^0 \cup \mathcal{P}_{n+1}^1) \setminus J$  where

$$\mathcal{P}_{n+1}^0 = \left\{ \{i \in A_\varepsilon^n : \text{cf}(g_{A_\varepsilon^n, n}(i)) \in \mathfrak{b}_{\theta_\varepsilon^n}[\mathfrak{a}_\varepsilon^n]\} : \varepsilon < \varepsilon_n^* \right\}$$

and

$$\mathcal{P}_{n+1}^1 = \left\{ \{i \in A^* : \text{cf}(g_{A^*,n}(i)) \leq \mu\} : A^* \in \mathcal{P}_n \right\}.$$

$J_{n+1}$  is defined as in clause (ix).

(Note: possibly  $(\mathcal{P}_{n+1}^0 \cup \mathcal{P}_{n+1}^1) \cap J \neq \emptyset$  but this does not cause problems).

So let us check clauses (i) – (ix).

Clause (i): Trivial by the choice of  $\mathbf{e}$  and  $g_{n+1}$ .

Clause (ii): By the definition of  $g_{n+1}(i)$  above it is  $< g_n(i)$  except when  $g_n(i) = g(i)$ , but by clause (vii) we know that  $g < g_n \bmod J_n$  hence necessarily  $\{i < \kappa : g_n(i) = g(i)\} \in J_n$ , so really  $g_{n+1} < g_n \bmod J_n$ .

Clause (iii): Clearly if  $A \in \mathcal{P}_{n+1}$  then  $A \subseteq \kappa$  and  $A \in (J + \mathcal{U}_{n+1})^+$  by the choice of  $\mathcal{P}_{n+1}$ ,  $|\mathcal{P}_{n+1}| \leq |\mathcal{P}_n| + |\varepsilon_n^*| + \aleph_0$  and  $|\mathcal{P}_n| \leq \mu$  by clause (iii) for  $n$  (i.e. the induction hypothesis) and during the construction we have shown that  $|\varepsilon_n^*| = |\mathcal{Y}| \leq \mu$ . The last phrase of clause (iii) holds by the choice of  $J_{n+1}$ .

Clause (iv): let  $A \in \mathcal{P}_{n+1}$  so we have two cases.

Case 1:  $A \in \mathcal{P}_{n+1}^0$ .

So for some  $\varepsilon < \varepsilon_n^*$  we have  $(\theta_\varepsilon^n \in \lambda \cap \text{pcf}(\mathfrak{a}_\varepsilon^n)$  and)  $A =: \{i \in A_\varepsilon^n : \text{cf}(g_{A_\varepsilon^n,n}(i)) \in \mathfrak{b}_{\theta_\varepsilon^n}[\mathfrak{a}_\varepsilon^n]\}$ . Let  $g_{A,n+1} \in \prod_{i \in A} f(i)$  be defined by  $g_{A,n+1}(i) = \mathbf{e}_{g_{A_\varepsilon^n,n}(i)(\varepsilon)}(g^{n,1}(\text{cf}(g_{A_\varepsilon^n,n}(i))))$ .

By the choice of  $g^{n,1} \in \Pi \mathfrak{a}_n$  we have:

$$g^{n,1} \upharpoonright \mathfrak{b}_{\theta_\varepsilon^n}[\mathfrak{a}_\varepsilon^n] \in N.$$

Now the set  $A$  is definable from  $A_\varepsilon^n, g_{A_\varepsilon^n,n}$  and  $\mathfrak{b}_{\theta_\varepsilon^n}[\mathfrak{a}_\varepsilon^n]$ , all of which belong to  $N$  hence  $A \in N$ . Also  $A_\varepsilon^n \in N$  and clearly  $g_{A,n+1}$  is definable from the functions  $g^{n,1} \upharpoonright \mathfrak{b}_{\theta_\varepsilon^n}[\mathfrak{a}_\varepsilon^n], g^{n,2}, g_{A_\varepsilon^n,n}, A_\varepsilon^n$  and the function  $\mathbf{e}$  (see the definition of  $g_{n+1}$  by cases), but all four are from  $N$  so  $g_{A,n+1} \in N$ . Lastly,  $g_{n+1} \upharpoonright A \equiv_J g_{A,n+1}$  as  $i \in A$  &  $g_{A_\varepsilon^n,n}(i) = g_n(i)$  &  $g_n(i) > g(i) \Rightarrow g_{n+1}(i) = g_{A,n+1}(i)$  and each of the three assumptions fail only for a set of  $i \in A$  that belongs to  $J$ .

Case 2:  $A \in \mathcal{P}_{n+1}^1$ .

So for some  $A^* \in \mathcal{P}_n$  we have

$$A = \{i < \kappa : i \in A^* \text{ and } \text{cf}(g_{A^*,n}(i)) \leq \mu\}.$$

Let  $g_{A,n+1}(i) \equiv \mathbf{e}_{g_{A,n}}(g^{n,3}(\text{cf}(g_{A^*,n}(i))))$ . Again,  $g_{A,n+1} \in N, g_{A,n+1} \equiv_J g_{n+1} \upharpoonright A$ . Looking at the definition of  $g_{A,n+1}$ , clearly  $g_{A,n}$  is definable from  $g^{n,2} \in N, g_{A^*,n}$  and the function  $\mathbf{e}$ , all of which belong to  $N$ .

Clause (v): Holds trivially.

Clause (vi): Assume  $A \in \mathcal{P}_n$  and  $B \subseteq A$  satisfies  $B \in J^+$  (so also  $A \in J^+$ ) and  $(\forall B' \in J_{n+1})(B' \cap B \in J)$  we have to find  $A' \in \mathcal{P}_{n+1}$ , such that  $A' \subseteq A$  &  $A' \cap B \in J^+$ .

Case 1:  $B_1 = \{i \in B : \text{cf}(g_{A,n}(i)) \leq \mu\} \in J^+$ .

In this case  $A' =: \{i \in A : \text{cf}(g_{A,n}(i)) \leq \mu\} \in \mathcal{P}_{n+1}^1 \subseteq \mathcal{P}_{n+1}$  and  $A' \cap B \in J^+$  by the assumption of the case.

Case 2: For some  $\varepsilon < \varepsilon_n^*$  we have  $A = A_\varepsilon^n$  and

$$B_2 = \{i \in B : \text{cf}(g_{A,n}(i)) \in \mathfrak{b}_{\theta_\varepsilon^n}[\mathfrak{a}_\varepsilon^n]\} \in J^+.$$

In this case  $A' =: \{i \in A : \text{cf}(g_{A,n}(i)) \in \mathfrak{b}_{\theta_\varepsilon^n}[\mathfrak{a}_\varepsilon^n]\} \in J^+$  belongs to  $\mathcal{P}_{n+1}^1 \subseteq \mathcal{P}_{n+1}$ , is  $\subseteq A$  and  $B \cap A' \in J^+$  by the assumption of the case (remember  $B \cap \mathcal{U}_{n+1} = \emptyset$ ).

Case 3: Neither Case 1 nor Case 2.

So  $B_3 = B \setminus B_1 \in J^+$  and let  $\lambda_i = \text{cf}(g_{A,n}(i))$ .

We shall show that  $\prod_{i \in B_3} \text{cf}(g_{A,n}(i))/J$  is  $\lambda$ -directed. This suffices as letting

$\lambda_i =: \text{cf}(g_{A,n}(i)) \in (\mu, f(i)]$ , by [Sh:g, II,1.4](1),p.46,50 for some  $\lambda'_i = \text{cf}(\lambda'_i) \leq \lambda_i$ , we have

$\liminf_{J \upharpoonright B_3} \langle \lambda'_i : i \in B_3 \rangle = \liminf_{J \upharpoonright B_3} \langle \lambda_i : i \in B_2 \rangle$  and

$\lambda = \text{tcf} \prod_{i \subseteq B_3} \lambda'_i / (J \upharpoonright B_3)$  and this shows that the conclusion of 1.1 holds, contra-

dicting our initial assumption, so the  $\lambda$ -directedness really suffices.

Now  $i \in B \setminus B_1 \Rightarrow \lambda_i = \text{cf}(g_n(i)) > \mu$ ; and if  $\prod_{i \in B_3} \lambda_i / J$  is not  $\lambda$ -directed, by

[Sh:g],I,§1 for some  $B_4 \subseteq B_3$  and  $\theta = \text{cf}(\theta) < \lambda$  we have:  $B_4 \in J^+$  and  $\prod_{i \in B_4} \lambda_i / J$

has true cofinality  $\theta$ . Hence  $\theta \in \text{pcf}\{\text{cf}(g_{A,n}(i)) : i \in A \text{ and } \text{cf}(g_n(i)) > \mu\}$ , and as  $\theta > \mu$ , for some  $\varepsilon < \varepsilon_n^*$  we have  $A = A_\varepsilon^n$  and  $\theta = \theta_\varepsilon^n$  so  $A' = \{i \in A : \text{cf}(g_{A,n}(i)) \in \mathfrak{b}_{\theta_\varepsilon^n}[\mathfrak{a}_\varepsilon^n]\}$  is as required in case 2 on  $B_2$  (note: we could have restricted ourselves to  $\theta$ 's like that).

Clause (vii): By the choice of  $g^{n,1}, g^{n,2}$  and  $g^n$  clearly  $i < \kappa$  &  $g(i) < g_n(i) \Rightarrow g(i) \leq g_{n+1}(i)$ . As  $g < g_n \text{ mod } D$  it suffices to prove  $B =: \{i : g(i) = g_{n+1}(i)\} \in J$ .

modified:2005-07-06

(589) revision:2005-07-05

If not, we choose by induction on  $\ell \leq n+1$  a member  $B_\ell$  of  $\mathcal{P}_\ell$  such that  $B_\ell \cap B \in J^+$ . For  $\ell = 0$  let  $B_\ell = \kappa \in \mathcal{P}_0$ , for  $\ell + 1$  apply clause (vi) for  $\ell$  (even when  $\ell = n$  we have just proved it). So  $B_{n+1} \cap B \in J^+$  and  $g_{n+1} \upharpoonright (B_{n+1} \cap B) = g \upharpoonright (B_{n+1} \cap B)$  hence  $\neg(g_{n+1} \upharpoonright B_{n+1} \neq_J g \upharpoonright B_{n+1})$  but  $g_{n+1} \upharpoonright B_{n+1} \in N$  so we have contradicted the choice of  $g$  as contradicting (\*).

Clause (viii): Easy.

Clause (ix): By the choice of  $J_{n+1}$ . □<sub>1.1</sub>

**1.5 Claim.** *Assume*

- (a)  $J$  is an ideal<sup>4</sup> on  $\kappa$
- (b)  $f \in {}^\kappa \text{Ord}$ , each  $f(i)$  an infinite ordinal
- (c)  $T_J^2(f) \geq \lambda = \text{cf}(\lambda) > \mu > \kappa$
- (d)  $\mu = (2^\kappa)^+$  or at least
  - (d)<sup>-</sup> (i) if  $\mathfrak{a} \subseteq \text{Reg}$ , and
    - $(\forall \theta \in \mathfrak{a})(\mu \leq \theta < \lambda \ \& \ \mu \leq \theta < f(i))$
    - and  $|\mathfrak{a}| \leq \kappa$  then  $|\text{pcf}(\mathfrak{a})| \leq \mu$
  - (ii)  $|\mu^\kappa/J| < \lambda \vee (\forall g \in {}^\kappa \mu)[|\Pi g/J| < \lambda]$  and  $\mu$  is regular
- (e)  $\alpha < \lambda \Rightarrow |\alpha|^{\aleph_0} < \lambda$ .

Then for some  $A \in J^+$  and  $\bar{\lambda} = \langle \lambda_i : i \in A \rangle$  such that  $\mu \leq \text{cf}(\lambda_i) = \lambda_i \leq f(i)$  we have  $\prod_{i \in A} \lambda_i/J$  has true cofinality  $\lambda$ .

*Proof.* We repeat the proof of 1.1 but we choose  $N$  such that  ${}^\omega N \subseteq N$ , (possible by assumption (e) as  $\lambda$  is regular), and let  $F =: (\prod_{i < \kappa} f(i)) \cap N$ . If  $2^\kappa < \lambda$  then clearly

$$F = \left\{ g \in \prod_{i < \kappa} f(i) : \text{for some partition } \langle A_n : n < \omega \rangle \text{ of } \kappa \text{ and} \right.$$

$$g_n \in N \cap \prod_{i < \kappa} f(i) \text{ we have}$$

$$g = \bigcup_{n < \omega} (g_n \upharpoonright A_n) \left. \right\}.$$

---

<sup>4</sup>compared to 1.1 we are omitting “ $J$  is  $\aleph_1$ -complete”.

Then assume (\*) (from the proof of 1.1) fails and  $g \in \prod_{i < \kappa} f(i)$  exemplifies it and we let  $J'$  be the ideal  $J' = \{A \subseteq \kappa : g \upharpoonright A = g' \upharpoonright A \text{ for some } g' \in F\}$ .

Clearly  $J'$  is  $\aleph_1$ -complete,  $J' \subseteq J$  (as  $g$  is a counterexample to (\*)) and the representation of  $F$  above) and we continue as there getting the conclusion for  $J'$  hence for  $J$ .

If  $2^\kappa \geq \lambda$ , let  $F' = N \cap \prod_{i < \kappa} f(i)$ , then

$\otimes$  for  $g \in \prod_{i < \kappa} f(i)$  and  $A \in J^+$  we have (i)  $\Leftrightarrow$  (ii) where:

- (i) there are  $g'_n \in F'$  for  $n < \omega$  such that  $\{i < \kappa : \bigvee_{n < \omega} g(i) = g'_n(i)\} \supseteq A \text{ mod } J$
- (ii) for some  $g' \in F'$  we have  $\{i < \kappa : g(i) = g'(i)\} \supseteq A \text{ mod } J$ .

[Why?  $\Leftarrow$  is trivial; now  $\Rightarrow$  holds as  $g_n \in N$  also  $\langle g_n : n < \omega \rangle \in N$  hence  $\langle \{g_n(i) : n < \omega\} : i < \kappa \rangle \in N$  and use  $\omega^\kappa/J \leq \mu^\kappa/J < \lambda$  (or just  $\bigoplus_{J, \mu, \lambda}$  from 1.2(4).]

Let  $g \in \prod_{i < \kappa} f(i)$  be such that  $f \in N \cap \prod_{i < \kappa} f(i) \Rightarrow g \neq_J f$ . Now we repeat the proof of 1.1 with our  $\kappa, f, \lambda, N, F, g$  this time using the demands in clause (viii) (i.e.  $g(i) \leq g_n(i)$ ). The proof does not change except that we do not get a contradiction from  $n < \omega \Rightarrow g_{n+1} <_J g_n$ . However, for each  $i < \kappa, \langle g_n(i) : n < \omega \rangle$  is non-increasing (by clause (viii)) hence eventually constant and by that clause eventually equal to  $g(i)$ . So clause (i) of  $\otimes$  above holds hence clause (ii) so we are done.  $\square_{1.5}$

**1.6 Conclusion.** Assume  $J$  is an ideal on  $\kappa, f \in {}^\kappa \text{Ord}, i < \kappa \Rightarrow f(i) > 2^\kappa, \lambda = \text{cf}(\lambda) > 2^\kappa$ , and

(\*)  $J$  is  $\aleph_1$ -complete or  $(\forall \alpha < \lambda)(|\alpha|^{\aleph_0} < \lambda)$ .

Then (a)  $\Leftrightarrow$  (b)  $\Leftrightarrow$  (b)<sup>+</sup>  $\Leftrightarrow$  (c)  $\Leftrightarrow$  (c)<sup>+</sup> where

- (a) for some  $A \in J^+$  we have  $T_{J \upharpoonright A}^2(f \upharpoonright A) \geq \lambda$
- (b) for some  $A \in J^+$  and  $\lambda_i = \text{cf}(\lambda_i) \in (2^\kappa, f(i)]$  (for  $i \in A$ ) we have  $\prod_{i \in A} \lambda_i / (J \upharpoonright A)$  is  $\lambda$ -directed

modified:2005-07-06

(589) revision:2005-07-05

- (b)<sup>+</sup> like (b) but  $\prod_{i \in A} \lambda_i / (J \upharpoonright A)$  has true cofinality  $\lambda$
- (c) for some  $A \in J^+$ , and  $\bar{n} = \langle n_i : i < \kappa \rangle \in {}^\kappa \omega$  and ideal  $J^*$  on  $A^* = \bigcup_{i \in A} (\{i\} \times n_i)$  satisfying
- $$(\forall B \subseteq A)[B \in J \Leftrightarrow \bigcup_{i \in B} (\{i\} \times n_i) \in J^*]$$

and regular cardinals  $\lambda_{(i,n)} \in (2^\kappa, f(i)]$  we have  $\prod_{(i,n) \in A^*} \lambda_{(i,n)} / J^*$  is

$\lambda$ -directed

- (c)<sup>+</sup> as in (c) but  $\prod_{(i,n) \in A^*} \lambda_{(i,n)} / J^*$  has true cofinality  $\lambda$ .

*Proof.* Clearly  $(b)^+ \Rightarrow (b)$ ,  $(b) \Rightarrow (c)$ ,  $(b)^+ \Rightarrow (c)^+$  and  $(c)^+ \Rightarrow (c)$ . Also  $(b) \Rightarrow (b)^+$  by [Sh:g, Ch.II,1.4](1), and similarly  $(c) \Rightarrow (c)^+$ . Now we prove  $(c) \Rightarrow (a)$ ; let  $\lambda_i = \max\{\lambda_{(i,n)} : n < n_i\}$  and let  $g_i$  be a one-to-one function from

$\prod_{n < n_i} \lambda_{(i,n)}$  into  $\lambda_i$  and let  $\langle f_\alpha : \alpha < \lambda \rangle$  be a  $<_{J^*}$ -increasing sequence in

$\prod_{(i,n) \in A^*} \lambda_{(i,n)}$ . Define  $f_\alpha^* \in \prod_{i \in A} \lambda_i$  by  $f_\alpha^*(i) = g_i(f_\alpha \upharpoonright (\{i\} \times n_i))$ . So if  $\alpha < \beta$ , then

$$\left\{ i \in A : f_\alpha^*(i) = f_\beta^*(i) \right\} = \left\{ i : \bigwedge_{n < n_i} f_\alpha((i,n)) = f_\beta((i,n)) \right\}$$

so by the assumption on  $J^*$  and the choice of  $\langle f_\alpha : \alpha < \lambda \rangle$ , for  $\alpha < \beta < \lambda$  we get  $f_\alpha^* \neq_J f_\beta^*$  hence  $\{f_\alpha^* : \alpha < \lambda\}$  is as required in clause (a).

Lastly  $(a) \Rightarrow (b)$  by 1.1 (in the case  $J$  is  $\aleph_1$ -complete) or 1.5 (in the case  $(\forall \alpha < \lambda)(|\alpha|^{\aleph_0} < \lambda)$ ). We have gotten enough implications to prove the conclusions.  $\square_{1.6}$

**1.7 Conclusion.** Let  $D$  be an ultrafilter on  $\kappa$ . If  $\left| \prod_{i < \kappa} f(i) / D \right| \geq \lambda = \text{cf}(\lambda) > 2^\kappa$  and  $(\forall \alpha < \lambda)[|\alpha|^{\aleph_0} < \lambda]$ , then for some regular  $\lambda_i \leq f(i)$  (for  $i < \kappa$ ) we have  $\lambda = \text{tcf}(\prod_{i < \kappa} \lambda_i / D)$ .

*Remark.* On  $|\prod_{i < \kappa} \lambda_i / D|$ , see [Sh 506, 3.9B].

§2 THE TREE REVISED POWER

**2.1 Definition.** For  $\kappa$  regular and  $\lambda \geq \kappa$  let

$$\lambda^{\kappa, \text{tr}} = \sup\{|\text{lim}_\kappa(T)| : T \text{ a tree with } \leq \lambda \text{ nodes and } \kappa \text{ levels}\}$$

where  $\text{lim}_\kappa(T)$  is the set of  $\kappa$ -branches of  $T$ ; and let when  $\lambda \geq \mu \geq \kappa$  and  $\theta \geq 1$

$$\lambda^{\langle \kappa, \theta \rangle} = \text{Min} \left\{ \mu : \text{if } T \text{ is a tree with } \lambda \text{ nodes and } \kappa \text{ levels,} \right. \\ \left. \text{then there is } \mathcal{P} \in [[T]^\theta]^\mu \text{ such that} \right. \\ \left. \eta \in \text{lim}_\kappa(T) \Rightarrow (\exists A \in \mathcal{P})(\eta \subseteq A) \right\}.$$

$$\lambda^{\langle \kappa \rangle} = \lambda^{\langle \kappa, \kappa \rangle}.$$

Recall  $[A]^\kappa =: \{B : B \subseteq A \text{ and } |B| = \kappa\}$ .

*2.2 Remark.* 1) Clearly  $\lambda^{\langle \kappa, \theta \rangle} \leq \lambda^{\kappa, \text{tr}} \leq \lambda^{\langle \kappa, \theta \rangle} + \theta^\kappa$ .

2) If  $\kappa = \aleph_0$  then obviously  $\lambda^{\kappa, \text{tr}} = \lambda^\kappa$ .

3) Of course,  $\lambda^{\langle \kappa, \theta \rangle} \leq \text{cov}(\lambda, \theta^+, \kappa^+, \kappa)$  and  $\kappa \leq \theta \leq \sigma \leq \lambda \Rightarrow \lambda^{\langle \kappa, \theta \rangle} \leq \lambda^{\langle \kappa, \sigma \rangle} + \text{cov}(\lambda, \theta^+, \kappa^+, \kappa)$ . (See [Sh:g] if these concepts are unfamiliar.)

**2.3 Theorem.** *Let  $\kappa$  be regular uncountable  $\leq \lambda$ . Then the following cardinals are equal:*

- (i)  $\lambda^{\langle \kappa \rangle}$
- (ii)  $\lambda + \sup\{\max \text{pcf}(\mathbf{a}) : \mathbf{a} \subseteq \text{Reg} \cap \lambda \setminus \kappa, \mathbf{a} = \{\theta_\zeta : \zeta < \kappa\} \text{ strictly increasing, and if } \xi < \kappa \text{ then } \max \text{pcf}(\{\theta_\zeta : \zeta < \xi\}) \leq \theta_\xi \leq \lambda\}$ .

*Remark.* We can add

- (ii)<sup>-</sup> like (ii) but we demand only  $\max \text{pcf}(\{\theta_\zeta : \zeta < \xi\}) \leq \lambda$ .

*Proof. First inequality.* Cardinal of (i) (i.e.  $\lambda^{(\kappa)}$ ) is  $\leq$  cardinal of (ii).

Assume not and let  $\mu$  be the cardinal from clause (ii) so  $\mu \geq \lambda$ .

Let  $T$ , a tree with  $\kappa$  levels and  $\lambda$  nodes, exemplify  $\lambda^{(\kappa)} > \mu$ . Without loss of generality  $T \subseteq {}^{\kappa}>\lambda$  and  $\langle T = \triangleleft \upharpoonright T$ .

Let  $\{T, \kappa, \lambda, \mu\} \in \mathfrak{B}_n \prec (\mathcal{H}(\chi), \in <_{\chi}^*)$ ,  $\mu + 1 \subseteq \mathfrak{B}_n$ ,  $\|\mathfrak{B}_n\| = \mu$ , for  $n < \omega$ ,

$\mathfrak{B}_n \in \mathfrak{B}_{n+1}$ ,  $\mathfrak{B}_n \prec \mathfrak{B}_{n+1}$  and let  $\mathfrak{B} =: \bigcup_{n < \omega} \mathfrak{B}_n$ . So  $\mathcal{P} =: \mathfrak{B} \cap [T]^{\leq \kappa}$  cannot

exemplify (i). So there is  $\eta \in \lim_{\kappa}(T)$  such that  $(\forall A \in \mathcal{P})[\{\eta \upharpoonright \zeta : \zeta < \kappa\}] \not\subseteq A$ .

We choose by induction on  $n$ ,  $N_n^0, N_n^1$  such that:

- (a)  $N_n^0 \prec N_n^1 \prec \mathfrak{B}_n$
- (b)  $N_0^1 = \text{Sk}_{\mathfrak{B}_0}(\{\zeta : \zeta < \kappa\} \cup \{\eta \upharpoonright \zeta : \zeta < \kappa\} \cup \{\kappa, \mu, \lambda, T\})$  and  
 $N_0^0 = \text{Sk}_{\mathfrak{B}_0}(\{\zeta : \zeta < \kappa\} \cup \{\kappa, \mu, \lambda, T\})$
- (c)  $\|N_n^{\ell}\| = \kappa$
- (d)  $N_n^0 \in \mathfrak{B}_{n+1}$
- (e)  $N_n^1 = \text{Sk}_{\mathfrak{B}_n}(N_n^0 \cup \{\eta \upharpoonright \zeta : \zeta < \kappa\})$
- (f)  $\theta \in \lambda^+ \cap \text{Reg} \cap N_n^0 \setminus \kappa^+ \Rightarrow \sup(N_{n+1}^0 \cap \theta) > \sup(N_n^1 \cap \theta)$ .

(Here ‘‘Sk’’ denotes the Skolem hull.)

Let us carry the induction.

For  $n = 0$ : No problem.

For  $n + 1$ : Let  $\mathfrak{a}^n =: N_n^0 \cap \text{Reg} \cap \lambda^+ \setminus \kappa^+$ , so  $\mathfrak{a}^n \in \mathfrak{B}_{n+1}$  and  $\mathfrak{a}^n$  is a set of cardinality  $\leq \kappa$  of regular cardinals  $\in (\kappa, \lambda^+)$ .

Let  $g^n \in \Pi \mathfrak{a}^n$  be defined by  $g^n(\theta) =: \sup(N_n^1 \cap \theta)$ . Let

$$(*)_1 \quad I^n = \{\mathfrak{b} \subseteq \mathfrak{a}^n : \text{for some } f \in (\Pi \mathfrak{a}^n) \cap \mathfrak{B}_{n+1} \text{ we have } g^n \upharpoonright \mathfrak{b} < f\},$$

so we need to show  $\mathfrak{a}^n \in I^n$ .

An easy induction on  $\text{pcf}(\mathfrak{a}^n)$  tells us that

$$(*)_2 \quad J_{\leq \mu}[\mathfrak{a}^n] \subseteq I^n \text{ (in particular all singletons are in } I^n).$$

Fact: There is  $f^* \in \mathfrak{B}_{n+1} \cap \Pi \mathfrak{a}^n$  such that:

$$\mathfrak{b}^n =: \{\theta \in \mathfrak{a}^n : f^*(\theta) < g^n(\theta)\}$$

satisfies

$$[\mathfrak{b}^n]^{< \kappa} \subseteq J_{\leq \lambda}[\mathfrak{a}^n]$$



(yes! not  $J_{\leq \mu}[\mathfrak{a}^n]$ ).

*Proof.* In  $\mathfrak{B}_{n+1}$  there is a list  $\{a_{n,\varepsilon} : \varepsilon < \kappa\}$  of  $N_n^0$ . For each  $\nu \in T$  let  $\nu$  be of level  $\zeta$  and let  $N_{n,\nu}^1 = \text{Sk}_{\mathfrak{B}_n}(\{(a_{n,\varepsilon}, \nu \upharpoonright \varepsilon) : \varepsilon < \zeta\})$ . So the function  $\nu \mapsto N_{n,\nu}^1$  (i.e. the set of pairs  $\langle (\nu, N_{n,\nu}^1) : \nu \in T \rangle$ ) belongs to  $\mathfrak{B}_{n+1}$ . Clearly  $\langle N_{n,\eta \upharpoonright \zeta}^1 : \zeta < \kappa \rangle$  is increasing continuous with union  $N_n^1$ . Let  $g_{n,\nu}^1 \in \Pi(\mathfrak{a}^n \cap N_{n,\nu}^1)$  be defined by  $g_{n,\nu}^1(\theta) = \sup(\theta \cap N_{n,\nu}^1)$ , so  $\{(\mathfrak{a}^n \cap N_{n,\nu}^1, g_{n,\nu}^1) : \nu \in T\} \in \mathfrak{B}_{n+1}$ . Now  $\Pi \mathfrak{a}^n / J_{\leq \lambda}[\mathfrak{a}^n]$  is  $\lambda^+$ -directed, hence as  $|T| \leq \lambda$  there is  $f^* \in \Pi \mathfrak{a}^n$  such that:

$$(*)_3 \quad \nu \in T \Rightarrow g_{n,\nu}^1 <_{J_{\leq \lambda}[\mathfrak{a}^n]} f^*,$$

and by the previous sentence without loss of generality  $f^* \in \mathfrak{B}_{n+1}$ . Note that for  $\theta \in \mathfrak{a}^n$  the sequence  $\langle g_{n,\eta \upharpoonright \zeta}^1(\theta) : \zeta < \kappa \rangle$  is non-decreasing with limit  $g^n(\theta)$ . Let  $\mathfrak{c} = \{\theta \in \mathfrak{a}^n : f^*(\theta) < g^n(\theta)\}$ , now note

$$(*)_4 \quad \text{if } \theta \in \mathfrak{c} \text{ then for every } \zeta < \kappa \text{ large enough, } f^*(\theta) < g_{n,\eta \upharpoonright \zeta}^1(\theta).$$

Hence  $\mathfrak{c}' \in [\mathfrak{c}]^{< \kappa} \Rightarrow \mathfrak{c}' \in J_{\leq \lambda}[\mathfrak{a}^n]$  as required in the fact.

(Why the implication? Because if  $\mathfrak{c}' \subseteq \mathfrak{c}, |\mathfrak{c}'| < \kappa$  then by  $(*)_4$  for some  $\zeta < \kappa$  we have  $f^* \upharpoonright \mathfrak{c}' < g'_{n,\eta \upharpoonright \zeta} \upharpoonright \mathfrak{c}'$  which by  $(*)_3$  gives  $\mathfrak{c}' \in J_{\leq \lambda}[\mathfrak{a}^n]$ ) so let  $\mathfrak{b}^n = \mathfrak{c}$ .  $\square_{\text{Fact}}$

Now if  $\mathfrak{b}^n$  is in  $J_{\leq \mu}[\mathfrak{a}^n]$ , by  $(*)_1 + (*)_2$  above we can finish the induction step. If not, some  $\tau^* \in \text{Reg} \setminus \mu^+$  satisfies  $\tau^* \in \text{pcf}(\mathfrak{b}^n)$ ; let  $\langle \mathfrak{c}_\zeta : \zeta < \kappa \rangle$  be an increasing continuous sequence of subsets of  $\mathfrak{a}^n$  each of cardinality  $< \kappa$  such that  $\mathfrak{b}^n = \bigcup_{\zeta < \kappa} \mathfrak{c}_\zeta$  and so (by the fact above)  $\zeta < \kappa \Rightarrow \tau^* > \lambda \geq \max \text{pcf}(\mathfrak{c}_\zeta)$ . We

know that this implies that for some club  $E$  of  $\kappa$  and  $\theta_\zeta \in \text{pcf}(\mathfrak{c}_\zeta)$ , for  $\zeta \in E$ ,  $\tau^* \in \text{pcf}_{\kappa\text{-complete}}(\{\theta_\zeta : \zeta \in E\})$  and  $\langle \theta_\zeta : \zeta \in E \rangle$  is strictly increasing and  $\max \text{pcf}\{\theta_\zeta : \zeta \in E \cap \xi\} \leq \theta_\xi$  for  $\xi \in E$ , by [Sh:g, Ch.VIII,1.5](2),(3),p.317.

Now  $\max \text{pcf}\{\theta_\varepsilon : \varepsilon \in \zeta \cap E\} \leq \max \text{pcf}(\mathfrak{c}_\zeta) \leq \lambda$  so  $\mu < \tau^* \leq$  the cardinal from clause (i) of 2.3, against an assumption. So we have carried out the inductive step in defining  $N_n^0, N_n^1$ .

So  $N_n^0, N_n^1$  are well defined for every  $n$ , clearly  $\bigcup_{n < \omega} N_n^0 \cap \lambda = \bigcup_{n < \omega} N_n^1 \cap \lambda$  (see [Sh:g, Ch.IX,3.3A,p.379]) hence  $\bigcup_{n < \omega} N_n^0 \cap T = \bigcup_{n < \omega} N_n^1 \cap T$ , hence for some  $n, N_n^0 \cap \{\eta \upharpoonright \zeta : \zeta < \kappa\}$  has cardinality  $\kappa$ . Now

$$A = \{\nu \in T : \text{for some } \rho \text{ we have } \nu \triangleleft \rho \in N_n^0\}$$

belongs to  $\mathfrak{B}_{n+1} \cap [T]^\kappa$  and  $\{\eta \upharpoonright \zeta : \zeta < \kappa\} \subseteq A$ , contradicting the choice of  $\eta$ .

modified:2005-07-06

revision:2005-07-05

(589)

Second inequality Cardinal of  $(ii) \leq$  cardinal of  $(i)$ .

By the proof of [Sh:g, II,3.5].

□<sub>2.3</sub>

**2.4 Definition.** 1) Assume  $I \subseteq J \subseteq \mathcal{P}(\kappa)$ ,  $I$  an ideal on  $\kappa$ ,  $J$  an ideal or the complement of a filter on  $\kappa$ , e.g.,  $J = \mathcal{P}^-(\kappa) = \mathcal{P}(\kappa) \setminus \{\kappa\}$  stipulating  $f \neq_J g \Leftrightarrow \{i < \kappa : f(i) = g(i)\} \in J$ . We let

$$T_{I,J}^+(f, \lambda) = \sup\{|F|^+ : F \in \mathcal{F}_{I,J}(f, \lambda)\}$$

and

$$T_{I,J}(f, \lambda) = \sup\{|F| : F \in \mathcal{F}_{I,J}(f, \lambda)\},$$

where

$$\mathcal{F}_{I,J}(f, \lambda) = \{F \subseteq \prod_{i < \kappa} f(i) : f \neq g \in F \Rightarrow f \neq_J g$$

$$\text{and } A \in I \Rightarrow \lambda \geq |\{f \upharpoonright A : f \in F\}|\}.$$

2) For  $J$  an ideal on  $\kappa$ ,  $\theta \geq \kappa$  and  $f \in {}^\kappa(\text{Ord} \setminus \{0\})$ , we let

$$\mathbf{U}_J(f, \theta) = \text{Min}\{|\mathcal{P}| : \mathcal{P} \subseteq [\sup \text{Rang}(f)]^\theta \text{ and for every } g \in \prod_{i < \kappa} f(i)$$

$$\text{for some } a \in \mathcal{P} \text{ we have } \{i < \kappa : g(i) \in a\} \in J^+\}.$$

If  $\theta = \kappa$  ( $= \text{Dom}(J)$ ), then we may omit  $\theta$ . If  $f$  is constantly  $\lambda$  we may write  $\lambda$  instead of  $f$ .

3) For  $I \subseteq J$ ,  $I$  ideal on  $\kappa$ ,  $J$  an ideal or complement of a filter on  $\kappa$ ,  $\mu \geq \theta \geq \kappa$  and  $f \in {}^\kappa(\text{Ord} \setminus \{0\})$  let

$$\mathbf{U}_{I,J}(f, \theta, \mu) = \sup\{\mathbf{U}_J(F, \theta) : F \in \mathcal{F}_I^-(f, \mu)\}$$

where

$$\mathcal{F}_I^-(f, \mu) = \{F : F \subseteq \prod_{i < \kappa} f(i) \text{ and}$$

$$A \in I \Rightarrow \mu \geq |\{f \upharpoonright A : f \in F\}|\}$$

and

$$\mathbf{U}_J(F, \theta) = \text{Min}\{|\mathcal{P}| : \mathcal{P} \subseteq [\text{sup Rang}(f)]^\theta \text{ and for every } f \in F \\ \text{for some } a \in \mathcal{P} \text{ we have } \{i < \kappa : f(i) \in a\} \in J^+\}.$$

2.5 *Fact.* Let  $\lambda \geq \kappa = \text{cf}(\kappa) > \aleph_0$ .

1)  $\lambda^{\kappa, \text{tr}} = T_{J_\kappa^{\text{bd}}, \mathcal{P}^-(\kappa)}(\lambda, \lambda)$  and  $\lambda^{\langle \kappa, \theta \rangle} \leq \mathbf{U}_{J_\kappa^{\text{bd}}}(\lambda, \theta)$ .

2) If  $\lambda \geq \mu$ , then  $\lambda^{\kappa, \text{tr}} \geq \mu^{\kappa, \text{tr}}$  and  $\lambda^{\langle \kappa \rangle} \geq \mu^{\langle \kappa \rangle}$ .

3)  $\lambda^{\kappa, \text{tr}} = \lambda^{\langle \kappa \rangle} + \kappa^{\kappa, \text{tr}}$ .

4) Assume  $I \subseteq J$  are ideals on  $\kappa$ . Then  $T_I^+(f, \lambda) > \mu$  if:

(i) each  $f(i)$  is a regular cardinal  $\lambda_i \in (\kappa, \lambda)$

(ii)  $\prod_{i < \kappa} f(i)/J$  is  $\mu$ -directed

(iii) for some  $A_\zeta \subseteq \kappa$  for  $\zeta < \zeta^* < \text{Min}f(j)$  we have:

$$\max \text{pcf}\{f(i) : i \in A_\zeta\} \leq \lambda \text{ (hence } \text{cf}\left(\prod_{i \in A_\zeta} f(i)\right) \leq \lambda) \text{ and } \{A_\zeta : \zeta < \zeta^*\}$$

generates an ideal on  $\kappa$  extending  $I$  but included in  $J$ .

5)  $\mathbf{U}_J(\lambda) \leq \mathbf{U}_J(\lambda, \theta) \leq \mathbf{U}_J(\lambda) + \text{cf}([\theta]^\kappa, \subseteq) \leq \mathbf{U}_J(\lambda) + \theta^\kappa$  and  $T_I(f) \leq \mathbf{U}_I(f) + 2^\kappa$  and  $\mathbf{U}_{I,J}(f, \lambda) \leq T_{I,J}(f, \lambda) \leq \mathbf{U}_{I,J}(f, \lambda) + 2^\kappa$  where  $I \subseteq J$  are ideals on  $\kappa$ .

Also obvious monotonicity properties (in  $I, J, \lambda, \theta, f$ ) hold.

*Proof.* 1) Easy. Let us prove the first equation. First assume  $F \in \mathcal{F}_{J_\kappa^{\text{bd}}, \mathcal{P}^-(\kappa)}(\lambda, \lambda)$ , and we define a tree as follows: for  $i < \kappa$  the  $i$ th level is

$$T_i = \{f \upharpoonright i : f \in F\}$$

and

$$T = \bigcup_{i < \kappa} T_i, \text{ with the natural order } \subseteq .$$

Clearly  $T$  is a tree with  $\kappa$  levels, the  $i$ -th level being  $T_i$ .

By the definition of  $\mathcal{F}_{J_\kappa^{\text{bd}}, \mathcal{P}^-(\kappa)}(\lambda, \lambda)$  as  $i < \kappa \Rightarrow \{j : j < i\} \in J_\kappa^{\text{bd}}$ , clearly  $|T_i| \leq \lambda$ .

Now for each  $f \in F$ , clearly  $t_f =: \langle (f \upharpoonright i) : i < \kappa \rangle$  is a  $\kappa$ -branch of  $T$ , and  $f_1 \neq f_2 \in F \Rightarrow t_{f_1} \neq t_{f_2}$  so  $T$  has at least  $|F|$   $\kappa$ -branches.

The other direction is easy, too. Note that the proof gives  $=^+$ ; i.e., the supremum is obtained in one side iff it is obtained in the other side.

2) If  $T$  is a tree with  $\mu$  nodes and  $\kappa$  levels then we can add  $\lambda$  nodes adding  $\lambda$  branches. Also the other inequality is trivial.

3) First  $\lambda^{\kappa, \text{tr}} \geq \lambda^{<\kappa>}$  because if  $T$  is a tree with  $\lambda$  nodes and  $\kappa$  levels, then we know  $|\lim_{\kappa}(T)| \leq \lambda^{\kappa, \text{tr}}$ , hence  $\mathcal{P} = \{t : t \text{ is a } \kappa\text{-branch of } T\}$  has cardinality  $\leq \lambda^{\kappa, \text{tr}}$  and satisfies the requirement in the definition of  $\lambda^{<\kappa>}$ .

Second  $\lambda^{\kappa, \text{tr}} \geq \kappa^{\kappa, \text{tr}}$  by part (2) of 2.5.

Lastly,  $\lambda^{\kappa, \text{tr}} \leq \lambda^{<\kappa>} + \kappa^{\kappa, \text{tr}}$  because if  $T$  is a tree with  $\lambda$  nodes and  $\kappa$  levels, we know by Definition 2.1 that there is  $\mathcal{P} \subseteq [T]^{\kappa}$  of cardinality  $\leq \lambda^{<\kappa>}$  such that every  $\kappa$ -branch of  $T$  is included in some  $A \in \mathcal{P}$ , without loss of generality  $x <_T y \in A \in \mathcal{P} \Rightarrow x \in A$ ; so

$$\begin{aligned} |\lim_{\kappa}(T)| &= |\{t : t \text{ a } \kappa\text{-branch of } T\}| \\ &= \left| \bigcup_{A \in \mathcal{P}} \{t \subseteq A : t \text{ a } \kappa\text{-branch of } T\} \right| \\ &\leq \sum_{A \in \mathcal{P}} |\lim_{\kappa}(T \upharpoonright A)| \\ &\leq |\mathcal{P}| + \kappa^{\kappa, \text{tr}} \leq \lambda^{<\kappa>} + \kappa^{\kappa, \text{tr}}. \end{aligned}$$

4) Like the proof of [Sh:g, Ch.II,3.5].

5) Left to the reader. □<sub>2.5</sub>

## 2.6 Lemma. Assume

- (a)  $I \subseteq J$  are ideals on  $\kappa$
- (b)  $I$  is generated by  $\leq \mu^*$  sets,  $\mu^* \geq \kappa$
- (c)  $T_{I,J}^+(f, \lambda) > \mu = \text{cf}(\mu) > \mu^* \geq T_{I,J}(\mu^*, \kappa)$
- (d)  $\kappa$  is not the union of countably many members of  $I$ .

Then We can find  $A_0 \subseteq A_1 \subseteq \dots \subseteq A_n \subseteq \dots$  from  $I^+$  with union  $\kappa$ , such that for each  $n$  there is  $\langle \lambda_i^n : i \in A_n \rangle$ ,  $\mu^* < \lambda_i^n = \text{cf}(\lambda_i^n) \leq f(i)$  such that:

$$\prod_{i \in A_n} \lambda_i^n / J \text{ is } \mu\text{-directed}$$

$$A \subseteq A_n, A \in I \Rightarrow \text{cf}\left(\prod_{i \in A} \lambda_i^n\right) \leq \lambda.$$

*2.7 Remark.* The point in the proof is that if  $I$  is generated by  $\{B_\gamma : \gamma < \gamma^* \leq \mu^*\}$ , and  $\{\eta_\alpha : \alpha < \mu^+\}$  are distinct branches and  $f \in {}^A(\lambda + 1 \setminus \{0\})$ ,  $A \subseteq \kappa$  and  $i \in A \Rightarrow \text{cf}(f(i)) > \mu^*$ , then for some  $g < f$  for every  $\gamma < \gamma^*$  and  $\alpha < \mu^+$ ,  $\{i < \gamma : \text{if } \eta_\alpha(i) < f(i) \text{ then } \eta_\alpha(i) < g(i)\} = \gamma \bmod J_{<\lambda^+}(f \upharpoonright \gamma)$ .

*Proof.* Similar to the proof of 1.1 adding the main point of the proof of 2.3, the “fact” there.

We can further generalize

**2.8 Definition.** For  $I \subseteq J \subseteq \mathcal{P}(\kappa)$ , function  $f^* \in {}^\kappa\text{Reg}$  and  $\lambda$ , we let

$$\mathcal{F}_{(I,J,\lambda)}^1(f^*) = \left\{ F \subseteq \prod_{i < \kappa} f^*(i) : \text{if } A \in J \text{ then } \lambda \geq |\{(f \upharpoonright A)/I : f \in F\}| \right\}$$

(so  $I$  is without loss of generality an ideal on  $\kappa$  and this is just  $\mathcal{F}_I^-(f^*, \lambda)$ )

$$\mathcal{F}_{(I,J,\lambda)}^2(f^*) = \left\{ F \subseteq \prod_{i < \kappa} f^*(i) : \text{if } A \in J, \text{ and } f, g \in F \text{ are distinct then } \{i \in A : f(i) = g(i)\} \in I \right\}$$

$$\mathcal{F}_{(I,J,\lambda,\bar{\theta})}^3(f^*) = \left\{ F \subseteq \prod_{i < \kappa} f^*(i) : \text{if } A \in J, \text{ then for some } G \subseteq \prod_{i \in A} [f^*(i)]^{\theta_i} \text{ of cardinality } \leq \lambda \text{ we have } (\forall f \in F)(\exists g \in G)\{i \in A : f(i) \notin g(i)\} \in I \right\}.$$

If  $\Xi$  is a set of such tuples, then we let  $\mathcal{F}_\Xi^\ell(f^*) = \bigcap_{\Upsilon \in \Xi} \mathcal{F}_\Upsilon^\ell(f^*)$

If in all the tuples  $\lambda$  is the third element, we write triples and  $f, \lambda$  instead of  $f$ .

For any  $\mathcal{F}_\Upsilon^\ell$  we let  $T_\Upsilon^\ell(f^*) = \sup\{|F| : F \in \mathcal{F}_\Upsilon^\ell(f^*)\}$  but: instead of  $T$  we have  $F \in \mathcal{F}_I(f)$  exemplifying  $\mathbf{U}_{I,J}(f, \lambda) > \mu$ ; i.e.  $\mathbf{U}_{I,J}(F, \lambda) > \mu$ . Then  $\eta \in F$  satisfies  $(\forall A \in \mathcal{P})[\{i : \eta(i) \in A\} \in J]$ . We choose  $N_n^0, N_n^1$  satisfying (a)-(f) with  $\gamma_n = 1$ .

## §3 ON THE DEPTH BEHAVIOUR OF ULTRAPRODUCTS

The problem originates from Monk [M] and see on it Roslanowski Shelah [RoSh 534] and then [Sh 506, §3] but the presentation is self-contained. We would like to have (letting  $B_i$  denote Boolean algebra), for  $D$  an ultrafilter on  $\kappa$ :

$$\text{Depth}\left(\prod_{i<\kappa} B_i/D\right) \geq \left| \prod_{i<\kappa} \text{Depth}(B_i)/D \right|.$$

(If  $D$  is just a filter, we should use  $T_D$  instead of product in the right side). Because of the problem of attainment (serious by Magidor Shelah [MgSh 433]), we rephrase the question:

⊗ for  $D$  an ultrafilter on  $\kappa$ , does  $\lambda_i < \text{Depth}^+(B_i)$  for  $i < \kappa$  imply

$$\left| \prod_{i<\kappa} \lambda_i/D \right| < \text{Depth}^+\left(\prod_{i<\kappa} B_i/D\right)$$

at least when  $\lambda_i > 2^\kappa$ ;

⊗' for  $D$  a filter on  $\kappa$  does  $\lambda_i < \text{Depth}^+(B_i)$  for  $i < \kappa$  imply (assuming  $\lambda_i > 2^\kappa$  for simplicity):

$$\begin{aligned} \mu = \text{cf}(\mu) < T_{D+A}^+(\langle \lambda_i : i < \kappa \rangle) \text{ for some } A \in D^+ &\Rightarrow \\ \mu < \text{Depth}^+\left(\prod_{i<\kappa} B_i/(D+A)\right) \text{ for some } A \in D^+. & \end{aligned}$$

As found in [Sh 506], this actually is connected to a pcf problem, whose answer under reasonable restrictions is 1.6. So now we can clarify the connections.

Also, by changing the invariant (closing under homomorphisms, see [M]) we get a nicer result; this shall be dealt with here.

The results here (mainly 3.5) supercede [Sh 506, 3.26].

---

Done 24/Feb/95

**3.1 Definition.** 1) For a partial order  $P$  (e.g. a Boolean algebra) let

$$\text{Depth}^+(P) = \min\{\lambda : \text{we cannot find } a_\alpha \in P \text{ for } \alpha < \lambda \text{ such that } \alpha < \beta \Rightarrow a_\alpha <_P a_\beta\}.$$

2) For a Boolean algebra  $B$  let

$$D_h^+(B) = \text{Depth}_h^+(B) = \sup\{\text{Depth}^+(B') : B' \text{ is a homomorphic image of } B\}.$$

3)  $\text{Depth}(P) = \sup\{\mu : \text{there are } a_\alpha \in P \text{ for } \alpha < \mu \text{ such that } \alpha < \beta < \mu \Rightarrow a_\alpha <_P a_\beta\}.$

4)  $\text{Depth}_h(P) = D_h(P) = \sup\{\text{Depth}(B') : B' \text{ is a homomorphic image of } B\}.$

5) We write  $D_r$  or  $D_{h,r}$  or  $\text{Depth}_r$  if we restrict ourselves to regular cardinals. Of course we could have looked at the ordinals.

**3.2 Definition.** 1) For a linear order  $\mathcal{I}$ , let the interval Boolean algebra,  $BA[\mathcal{I}]$  be the Boolean algebra of subsets of  $\mathcal{I}$  generated by  $\{[s, t)_{\mathcal{I}} : s < t \text{ are from } \{-\infty\} \cup \mathcal{I} \cup \{+\infty\}\}$ .

2) For a Boolean algebra  $B$  and regular  $\theta$ , let  $\text{com}_{<\theta}(B)$  be the  $(< \theta)$ -completion of  $B$ , that is the closure of  $B$  under the operations  $-x$  and  $\bigvee_{i < \alpha} x_i$  for  $\alpha < \theta$  inside the completion of  $B$ .

*3.3 Fact.* 1) If  $B$  is the interval Boolean algebra of the ordinal  $\gamma \geq \omega$  then

- (a)  $D_h^+(B) = |\gamma|^+$
- (b)  $\text{Depth}^+(B) = |\gamma|^+.$

2) If  $B'$  is a subalgebra of a homomorphic image of  $B$ , then  $D_h^+(B) \geq D_h^+(B')$ .

3) If  $D' \supseteq D$  are filters on  $\kappa$  and for  $i < \kappa$ ,  $B'_i$  is a subalgebra of a homomorphic image of  $B_i$  then:

( $\alpha$ )  $\prod_{i < \kappa} B'_i/D'$  is a subalgebra of a homomorphic image of  $\prod_{i < \kappa} B_i/D$ , hence

( $\beta$ )  $D_h^+(\prod_{i < \kappa} B_i/D) \geq D_h^+(\prod_{i < \kappa} B'_i/D').$

4) In parts (2), (3) we can replace  $D_h$  by  $D$  if we omit “homomorphic image”.

*Proof.* Straightforward.

**3.4 Claim.** 1) If  $D$  is a filter on  $\kappa$  and for  $i < \kappa$ ,  $B_i$  a Boolean algebra,  $\lambda_i < \text{Depth}_h^+(B_i)$  then

- (a)  $\text{Depth}_h^+(\prod_{i < \kappa} B_i/D) \geq \sup_{D_1 \supseteq D} \left( \text{tcf}(\prod_{i < \kappa} \lambda_i/D_1) \right)^+$   
*(i.e. sup on the cases tcf is well defined)*
- (b)  $\text{Depth}_h^+(\prod_{i < \kappa} B_i/D)$  is  $\geq \text{Depth}_h^+(\mathcal{P}(\kappa)/D)$  and is at least  $\sup\{[\text{tcf}(\prod_{i < \kappa} \lambda'_i/D_1)]^+ : \lambda'_i < \text{Depth}^+(B_i), D_1 \supseteq D\}$ .

2)  $\mu < \text{Depth}_h^+(B)$  iff for some  $a_i \in B$  for  $i < \mu$  we have that:  $\alpha < \beta < \mu, n < \omega$ , and  $\alpha_\ell < \beta_\ell < \mu$  for  $\ell < n$  together imply that  $B \models "(a_\beta - a_\alpha) - \bigcup_{\ell < n} (a_{\alpha_\ell} - a_{\beta_\ell}) > 0"$ .

3) Let  $A \in D^+$  ( $D$  a filter on  $\kappa$ ). In  $\prod_{i < \kappa} B_i/D$  there is a chain of order type  $\Upsilon$  if in  $\prod_{i < \kappa} B_i/(D + A)$  there is such a chain. If  $\Upsilon = \lambda; \text{cf}(\lambda) > 2^\kappa$  also the inverse is true.

4) If  $\mu < \text{Depth}_h^+(\prod_{i < \kappa} B_i/D)$  and  $\text{cf}(\mu) > 2^\kappa$ , then we can find  $A \in D^+$  and

$f_\alpha \in \prod_{i < \kappa} B_i$  for  $\alpha < \mu$  such that letting  $D^* = D + A$ :

$\alpha < \beta < \mu \Rightarrow (\prod_{i < \kappa} B_i/D^*) \models f_\alpha/D^* < f_\beta/D^*$  moreover  $f_\alpha <_{D^*} f_\beta$ .

5) Like (1) replacing  $\text{Depth}_h^+$  by  $\text{Depth}^+$ ,  $D_1 \supseteq D$  by  $\{D + A : A \in D^+\}$ .

*Proof.* Check, e.g.:

2) The "if" direction:

Let  $I$  be the ideal of  $B$  generated by  $\{a_\alpha - a_\beta : \alpha < \beta < \mu\}$ ,  $h : B \rightarrow B/I$  the canonical homomorphism, so  $\langle a_\alpha/I : \alpha < \mu \rangle$  is strictly increasing in  $B/I$ .

The "only if" direction:

Let  $h$  be a homomorphism from  $B$  onto  $B_1$  and  $\langle b_\alpha : \alpha < \mu \rangle$  be a (strictly) increasing sequence of elements of  $B_1$ . Choose  $a_\alpha \in B$  such that  $h(a_\alpha) = b_\alpha$ , so  $\alpha < \beta \Rightarrow a_\alpha \setminus a_\beta \in \text{Ker}(h)$  but  $a_\alpha \notin \text{Ker}(h)$ , moreover  $\beta < \alpha \Rightarrow a_\alpha - a_\beta \notin \text{Ker}(h)$ .

3) The first implication is trivial, the second follows from part (4).

4) First, assume  $\mu$  is regular. Let  $\langle f_\alpha/D : \alpha < \mu \rangle$  exemplify  $\mu < \text{Depth}_h^+(\prod_{i < \kappa} B_i/D)$ .



Then  $\alpha < \beta < \mu \Rightarrow f_\alpha \leq_D f_\beta$  &  $\neg(f_\alpha =_D f_\beta)$ , so for each  $\alpha, \langle \{i < \kappa : f_\alpha(i) = f_\beta(i)\} / D : \beta < \mu \rangle$  is decreasing and  $|2^\kappa / D| < \mu = \text{cf}(\mu)$  hence for some  $\beta_\alpha \in (\alpha, \mu)$  we have  $(\forall \beta)(\beta_\alpha \leq \beta < \mu \Rightarrow \{i < \kappa : f_\alpha(i) \neq f_{\beta_\alpha}(i)\} = \{i < \kappa : f_\alpha(i) \neq f_\beta(i)\} \text{ mod } D$  (as  $f_\gamma / D$  is increasing). So  $\langle \{i : f_\alpha(i) = f_{\beta_\alpha}(i)\} / D : \alpha < \mu \rangle$  is decreasing and  $|2^\kappa / D| \leq 2^\kappa < \mu$ , hence for some  $A^* \subseteq \kappa$  the set  $E = \{\alpha < \mu : \{i < \kappa : f_\alpha(i) < f_{\beta_\alpha}(i)\} = A^* \text{ mod } D\}$  unbounded and even stationary in  $\mu$ . Let  $D^* = D + A^*$ , so for  $\alpha < \beta < \mu$  we have  $f_\alpha \leq_D f_\beta$  hence  $f_\alpha \leq_{D^*} f_\beta$ , but  $\alpha \in E$  &  $\beta \geq \beta_\alpha \Rightarrow f_\alpha \neq_{D^*} f_\beta$ . Hence some  $E' \subseteq \{\delta \in E : (\forall \alpha < \delta \cap E)(\beta_\alpha < \delta)\}$  is unbounded in  $\mu$  and clearly  $(\forall \alpha, \beta)(\alpha < \beta \text{ \& } \alpha \in E' \text{ \& } \beta \in E' \Rightarrow f_\alpha <_{D^*} f_\beta)$ .

So  $\{f_\alpha : \alpha \in E'\}$  exemplifies the conclusion.

Second, if  $\mu$  is singular, let  $\mu = \sum_{\zeta < \text{cf}(\mu)} \mu_\zeta, \mu_\zeta > 2^\kappa; \mu_\zeta$  strictly increasing and each

$\mu_\zeta$  is regular. So given  $\langle f_\alpha : \alpha < \mu \rangle$ , for each  $\zeta < \text{cf}(\mu)$  we can find  $E_\zeta \subseteq \mu_\zeta^+$  of cardinality  $\mu_\zeta^+$  and  $A_\zeta \in D^+$  such that  $\alpha \in E_\zeta$  &  $\beta \in E_\zeta$  &  $\alpha < \beta \Rightarrow f_\alpha <_{D+A_\zeta} f_\beta$ . For some  $A, \text{cf}(\mu) = \sup\{\zeta : A_\zeta = A\}$ ; so  $A$  and the  $f_\alpha$ 's for  $\alpha \in \{E_\zeta \setminus \{\text{Min}(E_\zeta)\} : \zeta < \text{cf}(\mu)\}$  is such that  $A_\zeta = A$  are as required.  $\square_{3.4}$

We now give lower bound of depth of reduced products of Boolean algebras  $B_i$  from the depths of the  $B_i$ 's.

**3.5 First Main Lemma.** *Let  $D$  be a filter on  $\kappa$  and  $\langle \lambda_i : i < \kappa \rangle$  a sequence of cardinals ( $> 2^\kappa$ ) and  $2^\kappa < \mu = \text{cf}(\mu)$ . Then:*

1)  $(\alpha) \Leftrightarrow (\alpha)^+ \Leftrightarrow (\beta) \Leftrightarrow (\beta)^- \Leftrightarrow (\gamma)^+ \Rightarrow (\gamma) \Rightarrow (\delta)$ .

2) *If in addition  $(\forall \sigma < \mu)(\sigma^{\aleph_0} < \mu) \vee (D \text{ is } \aleph_1\text{-complete})$  we also have  $(\gamma) \Leftrightarrow (\gamma)^+ \Leftrightarrow (\delta)$  so all clauses are equivalent*

where:

( $\alpha$ ) *if  $B_i$  is a Boolean algebra,  $\lambda_i \leq \text{Depth}^+(B_i)$  then  $\mu < \text{Depth}^+(\prod_{i < \kappa} B_i / D)$*

( $\beta$ ) *there are cardinals  $\gamma_i < \lambda_i$  for  $i < \kappa$  such that, letting  $B_i$  be  $BA[\gamma_i]$  = the interval Boolean algebra of (the linear order)  $\gamma_i$ , we have  $\mu < \text{Depth}^+(\prod_{i < \kappa} B_i / D)$*

( $\gamma$ ) *there are  $\langle \langle \lambda_{i,n} : n < n_i \rangle : i < \kappa \rangle$  where  $\lambda_{i,n} = \text{cf}(\lambda_{i,n}) < \lambda_i$  and a non-trivial filter  $D^*$  on  $\bigcup_{i < \kappa} (\{i\} \times n_i)$  such that:*

(i)  $\mu = \text{tcf}(\prod_{(i,n)} \lambda_{i,n} / D^*)$

(ii) *for some  $A^* \in D^+$  we have*

$$D + A^* = \{A \subseteq \kappa : \text{the set } \bigcup_{i \in A} (\{i\} \times n_i) \text{ belongs to } D^*\}$$

- ( $\delta$ ) for some filter  $D' = D + A, A \in D^+$  and cardinals  $\lambda'_i < \lambda_i$  we have  $\mu \leq T_{D'}(\langle \lambda_i : i < \kappa \rangle)$
- ( $\beta$ )' like ( $\beta$ ) we allow  $\gamma_i$  to be an ordinal
- ( $\beta$ )<sup>-</sup> letting  $B_i$  be the disjoint sum of  $\{BA[\gamma] : \gamma < \lambda_i\}$  we have:  
 $\mu < \text{Depth}^+(\prod_{i < \kappa} B_i/D)$ .
- ( $\gamma$ )<sup>+</sup> for some filter  $D^*$  of the form  $D + A$  and  $\lambda'_i = \text{cf}(\lambda'_i) < \lambda_i$  we have  $\mu = \text{tcf}\left(\prod_{i < \kappa} \lambda'_i/D^*\right)$
- ( $\alpha$ )<sup>+</sup> if  $B_i$  is a Boolean algebra,  $\lambda_i \leq \text{Depth}^+(B_i)$  then for some  $A \in D^+$  we have, setting  $D^* = D + A$ , that  $\mu < \text{Depth}^+\left(\prod_{i < \kappa} B_i, <_{D^*}\right)$ ; moreover for some  $f_\alpha \in \prod_{i < \kappa} B_i$  for  $\alpha < \mu$  we have  $\alpha < \beta \Rightarrow \{i : B_i \models f_\alpha(i) < f_\beta(i)\} = \kappa \bmod D^*$ .

*Proof.* 1) We shall prove  $(\alpha) \Leftrightarrow (\beta) \Rightarrow (\beta)' \Rightarrow (\beta)^- \Rightarrow (\beta)' \Rightarrow (\gamma)^+ \Rightarrow (\beta)$  and  $(\alpha)^+ \Leftrightarrow (\alpha)$  and  $(\gamma)^+ \Rightarrow (\gamma) \Rightarrow (\delta)$ .

This suffices.

Now for  $(\alpha)^+ \Rightarrow (\alpha)$  note that if  $(\lambda_i, B_i$  for  $i < \kappa$  are given and)  $A \in D^+$ ,  $\langle f_\alpha : \alpha < \lambda \rangle$  exemplify  $(\alpha)^+$  then letting  $f'_\alpha = (f_\alpha \upharpoonright A) \cup 0_{(\kappa \setminus A)}$ ; i.e.,  $f'_\alpha(i)$  is  $f_\alpha(i)$  when  $i \in A$  and  $0_{B_i}$  if  $i \in \kappa \setminus A$ , easily  $\langle f'_\alpha : \alpha < \lambda \rangle$  exemplifies  $(\alpha)$ . Next  $(\alpha) \Rightarrow (\alpha)^+$  by 3.4(4).

Now  $(\beta) \Rightarrow (\beta)' \Rightarrow (\beta)^-$  holds trivially and for  $(\beta)' \Rightarrow (\gamma)^+$  repeat the proof of [Sh 506, 3.24,p.35] or the relevant part of the proof of 3.6 below (with appropriate changes, the case there is more complicated). Also  $(\beta)^- \Rightarrow (\beta)'$  as in the proof of 3.6 below. Easily  $(\gamma)^+ \Rightarrow (\beta)$ ; also  $(\beta) \Rightarrow (\alpha)$  because

- (i) if  $\gamma_i$  a cardinal  $< \text{Depth}^+(B_i)$ , the Boolean Algebra  $BA[\gamma_i]$  can be embedded into  $B_i$ , and
- (ii) if  $B'_i$  is embeddable into  $B_i$  for  $i < \kappa$  then  $B' = \prod_{i < \kappa} B'_i/D$  can be embedded into  $\prod_{i < \kappa} B_i/D$
- (iii) if  $B'$  is embeddable into  $B$  then  $\text{Depth}^+(B') \leq \text{Depth}^+(B)$ .

Now  $(\alpha) \Rightarrow (\beta)$  trivially. Also  $(\gamma)^+ \Rightarrow (\gamma)$  trivially and  $(\gamma) \Rightarrow (\delta)$  as in the proof of 1.6. Next we note  $(\beta) \Rightarrow (\delta)$ , as if  $B_i = BA[\gamma_i]$  and  $\gamma_i < \lambda_i$  and  $\mu < \text{Depth}^+(\prod B_i/D)$ , then by 3.4(4) there is a sequence  $\langle f_\alpha : \alpha < \mu \rangle$  satisfying  $f_\alpha \in \prod_{i < \kappa} B_i$  and  $A^* \in D^+$  such that  $\alpha < \beta < \mu \Rightarrow f_\alpha <_{D+A} f_\beta$ . So  $\{f_\alpha : \alpha < \mu\}$  exemplifies that  $T_{D+A}(\langle |B_i| : i < \kappa \rangle) \geq \mu$ , as required in clause  $(\delta)$ .  
 2) Assume  $(\forall \sigma < \mu)(\sigma^{\aleph_0} < \mu) \vee (D \text{ is } \aleph_1\text{-complete})$ .  
 Now 1.6 gives  $(\delta) \Rightarrow (\gamma)^+$  hence  $(\gamma) \Leftrightarrow (\gamma)^+ \Leftrightarrow (\delta)$ . □<sub>3.5</sub>

Now we turn to the other variant,  $D_h^+$ .

**3.6 Second Main Lemma.** *Let  $D$  be a filter on  $\kappa$  and  $\langle \lambda_i : i < \kappa \rangle$  be a sequence of cardinals  $(> 2^\kappa)$  and  $2^\kappa < \mu = \text{cf}(\mu)$ . Then (see below on  $(\alpha), \dots$ ):*

- 1)  $(\alpha) \Leftrightarrow (\alpha)^+ \Leftrightarrow (\beta)' \Leftrightarrow (\beta)^- \Leftrightarrow (\gamma)$  and  $(\gamma)^+ \Rightarrow (\gamma) \Leftrightarrow (\beta) \Rightarrow (\delta)$ .
- 2) If  $(\forall \sigma < \mu)(\sigma^{\aleph_0} < \mu) \vee (D \text{ is } \aleph_1\text{-complete})$  we also have  $(\beta) \Leftrightarrow (\gamma) \Leftrightarrow (\gamma)^+ \Leftrightarrow (\delta)$  (so all clauses are equivalent) where:

( $\alpha$ ) if  $B_i$  is a Boolean algebra,  $\lambda_i \leq \text{Depth}_h^+(B_i)$  then  $\mu < \text{Depth}_h^+(\prod_{i < \kappa} B_i/D)$

( $\beta$ ) there are cardinals  $\gamma_i < \lambda_i$  for  $i < \kappa$  such that, letting  $B_i$  be  $BA[\gamma_i]$  = the interval Boolean algebra of (the linear order)  $\gamma_i$ , we have  $\mu < \text{Depth}_h^+(\prod_{i < \kappa} B_i/D)$

( $\gamma$ ) there are  $\langle \langle \lambda_{i,n} : n < n_i \rangle : i < \kappa \rangle$  where  $\lambda_{i,n} = \text{cf}(\lambda_{i,n}) < \lambda_i$  and a non-trivial filter  $D^*$  on  $\bigcup_{i < \kappa} \{i\} \times n_i$  such that:

$$\mu = \text{tcf}(\prod_{(i,n)} \lambda_{i,n}/D^*) \text{ and } D \subseteq \{A \subseteq \kappa : \text{the set } \bigcup_{i \in A} \{i\} \times n_i \text{ belongs to } D^*\}$$

( $\delta$ ) for some filter  $D^* \supseteq D$  and cardinals  $\lambda'_i < \lambda_i$  we have  $\mu \leq T_{D^*}(\langle \lambda_i : i < \kappa \rangle)$

- ( $\beta$ )' like ( $\beta$ ) but allowing  $\gamma_i$  to be any ordinal  $< \lambda_i$
- ( $\beta$ )<sup>-</sup> letting  $B_i$  be the disjoint sum of  $\{BA[\gamma] : \gamma < \lambda_i\}$  (so  $\text{Depth}_h^+(B_i) = \lambda_i$ ) we have:  
 $\mu < \text{Depth}_h^+(\prod_{i < \kappa} B_i/D)$

( $\gamma$ )<sup>+</sup> there are  $\lambda'_i = \text{cf}(\lambda'_i) \in (2^\kappa, \lambda_i)$  for  $i < \kappa$  and filter  $D_1^* \supseteq D$  such that  $\prod_{i \in A} \lambda'_i/D^*$  has true cofinality  $\mu$

modified:2005-07-06

(589) revision:2005-07-05

$(\alpha)^+$  if  $B_i$  is a Boolean algebra,  $\lambda_i \leq \text{Depth}_h^+(B_i)$  then for some filter  $D^* \supseteq D$  we have  $\mu < \text{Depth}_h^+\left(\prod_{i < \kappa} B_i/D^*\right)$ .

*Proof.* Now  $(\beta) \Rightarrow (\beta)'$  trivially and  $(\beta)' \Rightarrow (\beta)^-$  by 3.3(3) as  $BA[\gamma_i]$  can be embedded into  $B_i$ , and similarly  $(\beta) \Rightarrow (\alpha)$  by 3.3(3), and  $(\alpha) \Rightarrow (\beta)$  trivially. Also  $(\alpha) \Rightarrow (\alpha)^+$  trivially and  $(\alpha)^+ \Rightarrow (\alpha)$  easily (e.g. by 3.3(3)).

Also  $(\gamma)^+ \Rightarrow (\beta)$  trivially and  $(\beta) \Rightarrow (\delta)$  easily (as in the proof of 3.5).

We shall prove below  $(\gamma) \Rightarrow (\beta)$ ,  $(\beta)' \Rightarrow (\gamma)$  and  $(\beta)^- \Rightarrow (\beta)'$ .

Together we have  $(\alpha) \Rightarrow (\alpha)^+ \Rightarrow (\alpha) \Rightarrow (\beta) \Rightarrow (\beta)' \Rightarrow (\beta)^- \Rightarrow (\beta)' \Rightarrow (\gamma) \Rightarrow (\beta) \Rightarrow (\alpha)$  and  $(\gamma)^+ \Rightarrow (\gamma) \Rightarrow (\delta)$ ; this is enough for part (1).

Lastly, to prove part (2) of 3.6, by part (1) it is enough to prove  $(\delta) \Rightarrow (\gamma)^+$  as in the proof of 3.5.

$(\gamma) \Rightarrow (\beta)$

So we have  $\lambda_{i,n}$  (for  $n < n_i, i < \kappa$ ),  $D^*$  as in clause  $(\gamma)$  and let  $\langle g_\varepsilon : \varepsilon < \mu \rangle$  be

$<_{D^*}$ -increasing cofinal in  $\prod_{(i,n)} \lambda_{i,n}$  but abusing notation we may write  $g_\varepsilon(i, n)$  for

$g_\varepsilon((i, n))$ . Let  $\gamma_i =: \max\{\lambda_{i,n} : n < n_i\}$  and  $B_i =: BA[\gamma_i]$ , clearly  $\gamma_i < \lambda_i$ , a (regular) cardinal as by assumption  $\lambda_{i,n} < \lambda_i \leq \text{Depth}^+(B_i)$  is regular for  $n < n_i$ . In  $B_i$  we have a strictly increasing sequence of length  $\gamma_i$ . Without loss of generality  $\{\lambda_{i,n} : n < n_i\}$  is with no repetition (see [Sh:g, I,1.3](8)) and  $\lambda_{i,0} > \lambda_{i,1} > \dots > \lambda_{i,n_i-1}$ .

So for each  $i$  we can find  $a_{i,n} \in B_i$  (for  $n < n_i$ ) pairwise disjoint and  $\langle a_{i,n,\zeta} : \zeta < \lambda_{i,n} \rangle$  (again in  $B_i$ ) strictly increasing and  $< a_{i,n}$ .

Let  $b_{i,\varepsilon} \in B_i$  be  $\bigcup_{n < n_i} a_{i,n,g_\varepsilon(i,n)}$  (it is a finite union of members of  $B_i$  hence a

member of  $B_i$ ). Let  $b_\varepsilon \in \prod_{i < \kappa} B_i/D$  be  $b_\varepsilon = \langle b_{i,\varepsilon} : i < \kappa \rangle/D$ . Let  $J$  be the ideal of

$B =: \prod_{i < \kappa} B_i/D$  generated by  $\{b_\varepsilon - b_\zeta : \varepsilon < \zeta < \mu\}$ . Clearly  $\varepsilon < \zeta < \mu \Rightarrow b_\varepsilon \leq$

$b_\zeta \text{ mod } J$ , so by 3.4(2) what we have to prove is: assuming  $\varepsilon < \zeta < \mu, k < \omega$  and  $\varepsilon_m < \zeta_m < \mu$  for  $m < k$ , then  $B \models "b_\zeta - b_\varepsilon - \bigcup_{m < k} (b_{\varepsilon_m} - b_{\zeta_m}) \neq 0"$ .

Now

$$Y =: \left\{ (i, n) : g_\varepsilon(i, n) < g_\zeta(i, n) \text{ and } g_{\varepsilon_m}(i, n) < g_{\zeta_m}(i, n) \text{ for } m = 0, 1, \dots, k-1 \right\}$$

is known to belong to  $D^*$ , hence it is not empty so let  $(i^*, n^*) \in Y$ . Now  $B_{i^*} \models b_{i^*, \xi} \cap a_{i^*, n^*} = a_{i^*, n^*, g_\xi(i^*, n^*)}$ , for every  $\xi < \mu$ , in particular for  $\xi$  among  $\varepsilon, \zeta, \varepsilon_m, \zeta_m$  (for  $m < k$ ). As  $(i^*, n^*) \in Y$  we have

$$\begin{aligned} B_{i^*} \models (b_{i^*, \zeta} - b_{i^*, \varepsilon}) \cap a_{i^*, n^*} &\geq b_{i^*, \zeta} \cap a_{i^*, n^*} - b_{i^*, \varepsilon} \cap a_{i^*, n^*} \\ &= a_{i^*, n^*, g_\zeta(i^*, n^*)} - a_{i^*, n^*, g_\varepsilon(i^*, n^*)} > 0 \end{aligned}$$

(as  $g_\zeta(i^*, n^*) > g_\varepsilon(i^*, n^*)$  since  $(i^*, n^*) \in Y$ ) and similarly

$$B_{i^*} \models (b_{i^*, \varepsilon_m} - b_{i^*, \zeta_m}) \cap a_{i^*, n^*} = 0.$$

Hence

$$B_{i^*} \models "b_{i^*, \zeta} - b_{i^*, \varepsilon} - \bigcup_{m < k} (b_{i^*, \varepsilon_m} - b_{i^*, \zeta_m}) \neq 0"$$

As this holds for every  $(i^*, n^*) \in Y$  and  $Y \in D^*$ , by the assumptions on  $D^*$  we have

$$\{i^* < \kappa : B_{i^*} \models "b_{i^*, \zeta} - b_{i^*, \varepsilon} - \bigcup_{m < k} (b_{i^*, \varepsilon_m} - b_{i^*, \zeta_m}) \neq 0"\} \in D^+$$

hence in  $B, b_\zeta - b_\varepsilon \notin J$  as required.

( $\beta$ )'  $\Rightarrow$  ( $\gamma$ )

Let  $B_i$  be the interval Boolean algebra for  $\gamma_i$ , an ordinal  $< \lambda_i$ .

To prove clause ( $\gamma$ ) we assume that our regular  $\mu$  is  $< \text{Depth}_h^+(\prod_{i < \kappa} B_i/D)$ , and we have to find  $n_i < \omega, \lambda_{i, n} < \lambda_i$  for  $i < \kappa, n < n_i$  and  $D^*$  as in the conclusion of clause ( $\gamma$ ). So there are  $f_\alpha \in \prod_{i < \kappa} B_i$  for  $\alpha < \mu$  and an ideal  $J$  of the Boolean algebra  $B =: \prod_{i < \kappa} B_i/D$  such that  $f_\alpha/D < f_\beta/D \text{ mod } J$  for  $\alpha < \beta$ .

Remember  $\mu > 2^\kappa$ . Let  $f_\alpha(i) = \bigcup_{\ell < n(\alpha, i)} [j_{\alpha, i, 2\ell}, j_{\alpha, i, 2\ell+1})$  where  $j_{\alpha, i, \ell} < j_{\alpha, i, \ell+1} \leq \gamma_i$  for  $\ell < 2n(\alpha, i)$ . As  $\mu = \text{cf}(\mu) > 2^\kappa$ , without loss of generality  $n(\alpha, i) = n_i$  for all  $\alpha < \mu$ . By [Sh 430, 6.6D] (better yet, see [Sh 513, 6.1] or [Sh 620, 7.0]) we can find  $A \subseteq A^* =: \{(i, \ell) : i < \kappa, \ell < 2n_i\}$  and  $\langle \gamma_{i, \ell}^* : i < \kappa, \ell < 2n_i \rangle$  such that  $(i, \ell) \in A \Rightarrow \gamma_{i, \ell}^*$  is a limit ordinal of cofinality  $> 2^\kappa$  and

(\*) for every  $f \in \prod_{(i, \ell) \in A} \gamma_{i, \ell}^*$  and  $\alpha < \mu$  there is  $\beta \in (\alpha, \mu)$  such that:

$$\begin{aligned} (i, \ell) \in A^* \setminus A &\Rightarrow j_{\beta, i, \ell} = \gamma_{i, \ell}^* \\ (i, \ell) \in A &\Rightarrow f(i, \ell) < j_{\beta, i, \ell} < \gamma_{i, \ell}^* \end{aligned}$$

For  $(i, \ell) \in A^*$  define  $\beta_{i, \ell}^*$  by  
 $\beta_{i, \ell}^* =: \sup\{\gamma_{i, m}^* : (i, m) \in A^* \text{ and } \gamma_{i, m}^* < \gamma_{i, \ell}^* \text{ and } m < 2n_i$   
 (actually  $m < \ell$  suffices) $\}$ .

Now  $\beta_{i, \ell}^* < \gamma_{i, \ell}^*$  as the supremum is on a finite set, and the case  $0 = \beta_{i, \ell}^* = \gamma_{i, \ell}^*$  does not occur if  $(i, \ell) \in A$ . Let

$$Y = \left\{ \alpha < \mu : \begin{aligned} &\text{if } (i, \ell) \in A^* \setminus A \text{ then } j_{\alpha, i, \ell} = \gamma_{i, \ell}^* \\ &\text{and if } (i, \ell) \in A \text{ then } \beta_{i, \ell}^* < j_{\alpha, i, \ell} < \gamma_{i, \ell}^* \end{aligned} \right\}.$$

Clearly  $\{f_\alpha : \alpha \in Y\}$  satisfies (\*), so without loss of generality  $Y = \mu$ .

Clearly

(\*)<sub>1</sub>  $\langle \gamma_{i, \ell}^* : \ell < 2n_i \rangle$  is non-decreasing (for each  $i$ ).

Let  $u_i = \{\ell < 2n_i : (\forall m < \ell)[\gamma_{i, m}^* < \gamma_{i, \ell}^*]\}$ .

For  $i < \kappa, \ell < 2n_i$  define  $b_{i, \ell} =: f_\alpha(i) \cap [\beta_{i, \ell}^*, \gamma_{i, \ell}^*) \in B_i$ .

Let  $w_i =: \{\ell \in u_i : \text{for every (equivalently some) } \alpha < \mu \text{ we have}$   
 $B_i \models "[\beta_{i, \ell}^*, \gamma_{i, \ell}^*) \cap f_\alpha(i) \text{ is } \neq 0 \text{ and } \neq [\beta_{i, \ell}^*, \gamma_{i, \ell}^*]" \}$ .

So

(\*)<sub>2</sub>  $f_\alpha(i) \setminus \bigcup_{\ell \in w_i} b_{i, \ell}$  does not depend on  $\alpha$ , call it  $c_i (\in B_i)$ .

Let for  $\ell \in w_i$

$$u_{i, \ell} =: \left\{ n < n_i : [j_{\alpha, i, 2n}, j_{\alpha, i, 2n+1}) \text{ is not disjoint to } [\beta_{i, \ell}^*, \gamma_{i, \ell}^*) \right.$$

$$\left. \text{for some (equivalently every) } \alpha < \mu \right\}.$$

$$A_0 = \left\{ (i, \ell) : i < \kappa, \ell \in w_i \text{ and for some } n \in u_{i,\ell} \text{ we have, for some} \right. \\ \left. (\equiv \text{ every}) \alpha < \mu \text{ that } j_{\alpha,i,2n} \leq \beta_{i,\ell}^* < j_{\alpha,i,2n+1} < \gamma_{i,\ell}^* \right\}.$$

$$A_1 = \left\{ (i, \ell) : i < \kappa, \ell \in w_i \text{ and for some } n \in u_{i,\ell} \text{ we have, for some} \right. \\ \left. (\equiv \text{ every}) \alpha < \mu \text{ that } \beta_{i,\ell}^* < j_{\alpha,i,2n} < \gamma_{i,\ell}^* \leq j_{\alpha,i,2n+1} \right\}.$$

Let

$$b_i^0 =: \bigcup \left\{ [\beta_{i,\ell}^*, \gamma_{i,\ell}^*] : \ell \in w_i \text{ and } (i, \ell) \in A_0 \right\} \in B_i$$

$$b_i^1 =: \bigcup \left\{ [\beta_{i,\ell}^*, \gamma_{i,\ell}^*] : \ell \in w_i \text{ and } (i, \ell) \in A_1 \right\} \in B_i$$

$$c_i^1 = b_i^0 \cap b_i^1, c_i^2 = b_i^0 \cap (1 - b_i^1), c_i^3 = (1 - b_i^0) \cap b_i^1, c_i^4 = (1 - b_i^0) \cap (1 - b_i^1)$$

$$b_0 =: \langle b_i^0 : i < \kappa \rangle / D \in B$$

$$b_1 =: \langle b_i^1 : i < \kappa \rangle / D \in B$$

$$c_t = \langle c_i^t : i < \kappa \rangle / D \in B$$

$$c = \langle c_i : i < \kappa \rangle / D \in B.$$

Let  $J_1 = \{b \in B : \langle (f_\alpha/D) \cap b : \alpha < \mu \rangle \text{ is eventually constant modulo } J, \text{ i.e., } (\exists \alpha < \mu)(\forall \beta)[\alpha \leq \beta < \mu \rightarrow (f_\alpha/D) \cap b - (f_\beta/D) \cap b \in J]\}$ . Also  $B \models c \leq f_\alpha/D$ .

Clearly  $J_1$  is an ideal of  $B$  extending  $J$  and  $1_B \notin J_1$ . Also if  $x \in J_1^+$  then for some closed unbounded  $E \subseteq \mu$  we have:  $\langle (f_\alpha/D) \cap x : \alpha \in E \rangle$  is strictly increasing modulo  $J$ .

Hence by easy manipulations without loss of generality:

- (\*)<sub>3</sub>(a) if  $c_t \in J_1^+$  then  $\langle (f_\alpha/D) \cap c_t : \alpha < \mu \rangle$  is strictly increasing modulo  $J$
- (b) for at least one  $t, c_t \in J_1^+$ .

By (\*) we can find  $0 < \alpha_0 < \alpha_1 < \alpha_2 < \mu$  such that:

$$(*)_4 \text{ if } i < \kappa, \ell < 2n_i, \bigwedge_{\alpha < \mu} \gamma_{i,\ell}^* > j_{\alpha,i,\ell} \text{ and } k < 2 \text{ then} \\ \sup\{j_{\alpha_k,i,\ell_1} : j_{\alpha_k,i,\ell_1} < \gamma_{i,\ell}^* \text{ and } \ell_1 < 2n_i\} < j_{\alpha_{k+1},i,\ell}.$$

Now if in  $(*)_3, c_4 \in J_1^+$  occurs then

$$B_i \models "f_{\alpha_0}(i) \cap f_{\alpha_1}(i) \cap c_i^4 - c_i = \bigcup\{(f_{\alpha_0}(i) \cap f_{\alpha_1}(i)) \\ \cap [\beta_{i,\ell}^*, \gamma_{i,\ell}^*] : \ell \in w_i \text{ and } (i, \ell) \notin A_0, (i, \ell) \notin A_1\} \\ = \bigcup_{\ell \in w_i} 0_{B_i} = 0_{B_i}"$$

(as for each  $\ell \in w_i$  such that  $(i, \ell) \notin A_0 \cup A_1$ , the intersection is the intersection of two unions of intervals which are pairwise disjoint) whereas we know  $(f_{\alpha_0}/D) \cap (f_{\alpha_1}/D) \cap c_4 - c = {}_J f_{\alpha_0}/D \cap c_4 - c \notin J$ ; contradiction.

Next if in  $(*)_3, c_3 \in J_1^+$  holds then

$$B_i \models "(f_{\alpha_1}(i) \cap c_i^3 - c_i) - (f_{\alpha_0}(i) \cap c_i^3 - c_i) = \bigcup_i \{(f_{\alpha_1}(i) \cap [\beta_{i,\ell}^*, \gamma_{i,\ell}^*] \\ - f_{\alpha_0}(i) \cap [\beta_{i,\ell}^*, \gamma_{i,j}^*]) : \ell \in w_i \text{ and } (i, \ell) \in A_1 \setminus A_0\} = \bigcup_{\ell \in w_i} 0_{B_i} = 0_{B_i}"$$

(as for each  $\ell \in w_i$  such that  $(\ell, i) \in A_1 \setminus A_0$  the term is the difference of two unions of intervals but the first is included in the right most interval of the second) and we have a contradiction.

Now if in  $(*)_3, c_1 \in J^+$  holds then

$$B_i \models "(f_{\alpha_2}(i) \cap c_i^1 - c_i) - (f_{\alpha_1}(i) \cap c_i^1 - c_i) \cup (f_{\alpha_0}(i) \cap c_i^1 - c_i) \\ = \bigcup_i \{((f_{\alpha_2}(i) - f_{\alpha_1}(i) \cup f_{\alpha_0}(i)) \cap [\beta_{i,\ell}^*, \gamma_{i,\ell}^*]) : \ell \in w_i \text{ and } (i, \ell) \in A_0 \cap A_1\} \\ = \bigcup_{\ell \in w_i} 0_{B_i} = 0_{B_i}"$$

and we get a similar contradiction.

So

$$(*)_5 \text{ in } (*)_3, c_2 \in J_1^+.$$



Without loss of generality

- (\*)<sub>6</sub> for  $\alpha < \mu, i < \kappa$  and  $\ell < 2n_i$  such that  $(i, \ell) \in A$  we have  
 $\sup\{j_{2\alpha, i, \ell_1} : \ell_1 < 2n_i \text{ and } j_{2\alpha, i, \ell_1} < \gamma_{i, \ell}^* < j_{2\alpha+1, i, \ell}\}.$

Let  $v_i = \{\ell \in w_i : (i, \ell) \in A_0, (i, \ell) \notin A_1\}$ , so  $c_i^2 = \cup\{[\beta_{i, \ell}^*, \gamma_{i, \ell}^*) : \ell \in v_i\}$ . As  $\ell \in v_i \Rightarrow (i, \ell) \in A_0$  necessarily

- (\*)<sub>7</sub> if  $\ell \in v_i$  then  $\ell$  is odd and  $j_{\alpha, i, \ell-1} = \beta_{i, \ell}^* < j_{\alpha, i, 2\ell+1} < \gamma_{i, \ell}^*.$

Now for every  $\alpha < \mu$  define  $f'_\alpha \in \prod_{i < \kappa} B_i$  by

$$f'_\alpha(i) = \bigcup_{\ell \in v_i} [\beta_{i, \ell}^*, \max\{j_{2\alpha, i, 2n+1} : n \in u_i, \ell\}).$$

Clearly

$$B_i \models "f_{2\alpha}(i) \cap c_i^2 - c_i \leq f'_\alpha(i) \leq f_{2\alpha+1}(i) \cap c_i^2 - c_i".$$

Let  $Y^* =: \bigcup_{i < \kappa} (\{i\} \times v_i)$  and we shall define now a family  $D_0$  of subsets of  $Y^*$ .

For  $Y \subseteq Y^*$ , and for  $\alpha < \mu$  define  $f_{\alpha, Y} \in \prod_{i < \kappa} B_i$  by

$$f_{\alpha, Y}(i) = \cup\{[j_{\alpha, i, 2\ell}, j_{\alpha, i, 2\ell+1}) : \ell \in v_i \text{ and } (i, \ell) \notin Y\}.$$

For  $g \in G =: \prod_{(i, \ell) \in Y^*} [\beta_{i, \ell}^*, \gamma_{i, \ell}^*)$  define  $f_g \in \prod_{i < \kappa} B_i$  by  $f_g(i) = \bigcup_{\ell \in v_i} [\beta_{i, \ell}^*, g((i, \ell))]$ , now

- (\*)<sub>8</sub> for every  $\alpha < \mu$  for some  $g = g_\alpha^* \in G$  we have  $f'_\alpha = f_g$ .  
 [Why? By the previous analysis; in particular (\*)<sub>7</sub>].

Let

$$D_0 = \{Y \subseteq Y^* : \text{for some } g_1 \in G \text{ for every } g \in G \text{ satisfying} \\ [(i, \ell) \in Y^* \setminus Y \Rightarrow g(i, 0) = \beta_{i, \ell}^*] \text{ we have} \\ f_g/D - f_{g_1}/D \text{ belongs to } J_1\}$$

it is a filter on  $Y^*$ .

- (\*)<sub>9</sub> if  $g_1, g_2 \in G$  then  
 (a)  $g_1 \leq_{D_0} g_2 \Leftrightarrow B \models (f_{g_1}/D) \cap c_2 \leq (f_{g_2}/D) \cap c_2$   
 (b)  $g_1 <_{D_0} g_2 \Leftrightarrow B \models (f_{g_1}/D) \cap c_2 < (f_{g_2}/D) \cap c_2$

modified:2005-07-06

(589) revision:2005-07-05

(\*)<sub>10</sub> for every  $g' \in G$  for some  $\alpha(g') < \mu$  we have  $g' < g_{\alpha(g')}^*$  (see (\*)<sub>8</sub>).  
[Why? By (\*).]

Now

$$\otimes \text{cf}\left(\prod_{(i,\ell) \in Y^*} \gamma_{i,\ell}^*/D_0\right) \geq \mu.$$

[Why? If not, we can find  $G^* \subseteq G = \prod_{(i,\ell) \in Y^*} [\beta_{i,\ell}^*, \gamma_{i,\ell}^*]$  of cardinality  $< \mu$ ,

cofinal in  $\prod_{(i,\ell) \in Y^*} \gamma_{i,\ell}^*/D_0$ . For each  $g \in G^*$  for some  $\alpha(g) < \mu$  we have  $g < g_{\alpha(g)}^*$ , hence  $\alpha \in [\alpha(g), \mu) \Rightarrow g <_{D_0} g_{\alpha}^*$ , let  $\alpha(*) = \sup\{\alpha(g) : g \in G\}$  so  $\alpha(*) < \mu$  so  $\bigwedge_{g \in G} g <_{D_0} g_{\alpha(*)}^*$ ; contradiction, so  $\otimes$  holds].

So for some ultrafilter  $D^*$  on  $Y^*$  extending  $D_0$ ,  $\mu \leq \text{tcf}\left(\prod_{(i,\ell) \in Y^*} \gamma_{i,\ell}^*/D^*\right)$ , hence

$\mu \leq \text{tcf}\left(\prod_{(i,\ell) \in Y^*} \text{cf}(\gamma_{i,\ell}^*/D^*)\right)$  and by [Sh:g, II,1.3] for some

$\lambda'_{i,\ell} = \text{cf}(\lambda'_{i,\ell}) \leq \text{cf}(\gamma_{i,\ell}^*) \leq \gamma_i < \lambda_i$  we have  $\mu = \text{tcf}\left(\prod_{(i,\ell) \in Y^*} \lambda'_{i,\ell}/D^*\right)$  as required

(we could, instead of relying on this quotation, analyze more).

( $\beta$ )<sup>-</sup>  $\Rightarrow$  ( $\beta$ )'

Let  $B_{i,\gamma}$  be the interval Boolean algebra on  $\gamma$  for  $\gamma < \lambda_i, i < \kappa$ , and we let  $B_{i,\gamma}^*$  be generated by  $\{a_j^{i,\gamma} : j < \gamma\}$  freely except  $a_{j_1}^{i,\gamma} \leq a_{j_2}^{i,\gamma}$  for  $j_1 < j_2 < \gamma$ .

So without loss of generality  $B_i$  is the disjoint sum of  $\{B_{i,\gamma}^* : \gamma < \lambda_i\}$ . Let  $e_{i,\gamma} = 1_{B_{i,\gamma}^*}$  (so  $\langle e_{i,\gamma} : \gamma < \lambda_i \rangle$  is a maximal antichain of  $B_i$ ,  $B_i \upharpoonright \{x \in B_i : x \leq e_{i,\gamma}\}$  is isomorphic to  $B_{i,\gamma}^*$  and  $B_i$  is generated by  $\{x : (\exists \gamma < \lambda_i)(x \leq e_{i,\gamma})\}$ ). Let  $\langle f_\alpha : \alpha < \mu \rangle$  and an ideal  $J$  of  $B$  exemplify clause ( $\beta$ )<sup>-</sup>.

Let  $I_i$  be the ideal of  $B_i$  generated by  $\{e_{i,\gamma} : \gamma < \lambda_i\}$ , so it is a maximal ideal; let  $I$  be such that  $(B, I) = \prod_{i < \kappa} (B_i, I_i)/D$  so clearly  $|B/I| = |2^\kappa/D| \leq 2^\kappa < \text{cf}(\mu)$

(actually  $|B/I| = 2$  if  $D$  is an ultrafilter on  $\kappa$ ), so without loss of generality  $\alpha < \beta \leq \mu \Rightarrow f_\alpha/D = f_\beta/D \text{ mod } I$ . We can use  $\langle f_{1+\alpha}/D - f_0/D : \alpha < \mu \rangle$ , so without loss of generality  $f_\alpha/D \in I$ , hence without loss of generality  $f_\alpha(i) \in I_i$  for  $\alpha < \mu, i < \kappa$ .

Let  $f_\alpha(i) = \tau_{\alpha,i}(\dots, e_{i,\gamma(\alpha,i,\varepsilon)}, a_{j(\alpha,i,\varepsilon)}^{i,\gamma(\alpha,i,\varepsilon)}, \dots)_{\varepsilon < n_{\alpha,i}}$  where  $n_{\alpha,i} < \omega$  and  $\tau_{\alpha,i}$  is a Boolean term. As  $\mu$  is regular  $> 2^\kappa$ , without loss of generality  $\tau_{\alpha,i} = \tau_i$  and  $n_{\alpha,i} = n_i$ . Let  $\gamma_{\alpha,i,\varepsilon}^0 = \gamma(\alpha,i,\varepsilon)$  and  $\gamma_{\alpha,i,\varepsilon}^1 = j(\alpha,i,\varepsilon)$ .

By [Sh 430, 6.6D] (or better [Sh 620, 7.0]) we can find a subset  $A$  of  $A^* = \{(i, n, \ell) : i < \kappa \text{ and } n < n_i \text{ and } \ell < 2\}$  and  $\langle \gamma_{i,n,\ell}^* : i < \kappa \text{ and } n < n_i \text{ and } \ell < 2 \rangle$  such that:

$$(*) (A) \quad (i, n, \ell) \in A \Rightarrow \text{cf}(\gamma_{i,n,\ell}^*) > 2^\kappa$$

$$(B) \quad \text{for every } g \in \prod_{(i,n,\ell) \in A} \gamma_{i,n,\ell}^* \text{ for arbitrarily large } \alpha < \mu \text{ we have}$$

$$(i, n, \ell) \in A^* \setminus A \Rightarrow \gamma_{\alpha,i,n}^\ell = \gamma_{i,n,\ell}^*$$

$$(i, n, \ell) \in A \Rightarrow g(i, n, \ell) < \gamma_{\alpha,i,n}^\ell < \gamma_{i,n,\ell}^*.$$

Let

$$\beta_{i,n,\ell}^* = \sup\{\gamma_{i,n',\ell'}^* : n' < n_i, \ell' < 2 \text{ and } \gamma_{i,n',\ell'}^* < \gamma_{i,n,\ell}^*\}$$

Without loss of generality

$$(i, n, \ell) \in A \ \& \ \alpha < \mu \Rightarrow \gamma_{\alpha,i,n}^\ell \in (\beta_{i,n,\ell}^*, \gamma_{i,n,\ell}^*)$$

$$(i, n, \ell) \in A^* \setminus A \ \& \ \alpha < \mu \Rightarrow \gamma_{\alpha,i,n}^\ell = \gamma_{i,n,\ell}^*.$$

Also without loss of generality

$$(*) \quad \text{for } \alpha < \mu \text{ and } (i, n, \ell) \in A \text{ we have}$$

$$\gamma_{2\alpha+1,i,n}^\ell > \sup \left\{ \gamma_{2\alpha,i,n'}^{\ell'} : i < \kappa, \ell' < 2, n' < n_i, \right. \\ \left. \text{and } \gamma_{2\alpha,i,n'}^{\ell'} < \gamma_{i,n,\ell}^* \right\}.$$

Let  $\Delta_i = \{\gamma_{i,n,0}^* : n < n_i \text{ and } (i, n, 0) \in A^* \setminus A\}$  and

$$B'_i = B_i \upharpoonright \sum \{e_{i,\gamma} : \gamma \in \Delta_i\}.$$

We define  $f'_\alpha \in \prod_{i < \kappa} B'_i$  by  $f'_\alpha(i) = f_{2\alpha+1}(i) \cap (\bigcup_{\gamma \in \Delta_i} e_{i,\gamma}) \in B'_i \subseteq B_i$ .

Now easily  $f'_\alpha/D \leq f_{2\alpha+1}/D$  and (in B)  $f_{2\alpha}/D - f'_\alpha/D \leq f_{2\alpha}/D - f'_{2\alpha+1}/D \in J$ , hence  $\langle f'_\alpha : \alpha < \lambda \rangle$  is increasing modulo  $J$ , even strictly. So  $\langle B'_i : i < \kappa \rangle, \langle f'_\alpha : \alpha < \mu \rangle$  form a witness, too. But  $B'_i$  is isomorphic to the interval Boolean algebra of the ordinal  $\gamma_i = \sum_{\gamma \in \Delta_i} \gamma < \lambda_i$ , so we are almost done. Well,  $\gamma_i$  is an ordinal, not necessarily a cardinal, but we are proving  $(\beta)'$  not  $(\beta)$ . □<sub>3.6</sub>

§4 ON THE EXISTENCE OF INDEPENDENT SETS FOR STABLE THEORIES

The following is motivated by questions of Bays [Bay] which continues some investigations of [Sh:a] (better see [Sh:c]) dealing with questions on  $\text{Pr}_T(\mu), \text{Pr}_T^*$  for stable  $T$  (see Definition 4.2 below). We connect this to pcf, using [Sh 430, 3.17] and also [Sh 513, 6.12]). We assume basic knowledge on non-forking (see [Sh:c, Ch.III,I]) and we say some things on the combinatorics but the rest of the paper does not depend on this section. For simplicity, we concentrate on the regular case.

**4.1 Claim.** *Assume  $\lambda > \theta \geq \kappa$  are regular uncountable. Then the following are equivalent:*

- (A) *If  $\mu < \lambda$  and  $a_\alpha \in [\mu]^{<\kappa}$  for  $\alpha < \lambda$  then for some  $A \in [\lambda]^\lambda$  we have  $\bigcup_{\alpha \in A} a_\alpha$  has cardinality  $< \theta$*
- (B) *if  $\delta = \text{cf}(\delta) < \kappa$  and  $\eta_\alpha \in {}^\delta \lambda$  for  $\alpha < \lambda$  and  $|\{\eta_\alpha \upharpoonright i : \alpha < \lambda, i < \delta\}| < \lambda$  then for some  $A \in [\lambda]^\lambda$  the set  $\{\eta_\alpha \upharpoonright i : \alpha \in A, i < \delta\}$  has cardinality  $< \theta$ .*

*Remark.* Of course, if  $a_\alpha$  is just a set of cardinality  $< \kappa$ , by renaming  $a_\alpha \in [\lambda]^{<\kappa}$  and for some stationary  $S \subseteq \lambda$  and  $\alpha^* < \mu, \langle a_\alpha \setminus \alpha^* : \alpha \in S \rangle$  are pairwise disjoint, renaming  $\alpha^* = \mu < \lambda$ , etc., see more in [Sh 430, §2].

*Proof.* (A)  $\Rightarrow$  (B). Immediate.

$\neg(A) \Rightarrow \neg(B)$

Case 1: For some  $\mu \in (\theta, \lambda)$  we have  $\text{cf}(\mu) < \kappa$  and  $\text{pp}(\mu) \geq \lambda$ . Without loss of generality  $\mu$  is minimal. So

$$(*) \mathfrak{a} \subseteq \text{Reg} \cap \mu \setminus \theta, |\mathfrak{a}| < \kappa, \text{sup}(\mathfrak{a}) < \mu \Rightarrow \max \text{pcf}(\mathfrak{a}) < \mu.$$

Subcase 1a:  $\lambda < \text{pp}^+(\mu)$ .

So by [Sh:g, Ch.VIII,1.6](2),p.321, (if  $\text{cf}(\mu) > \aleph_0$ ) and [Sh 430, 6.5] (if  $\text{cf}(\mu) = \aleph_0$ ) we can find  $\langle \lambda_\alpha : \alpha < \text{cf}(\mu) \rangle$ , a strictly increasing sequence of regulars from  $(\theta, \mu)$  with limit  $\mu$  and an ideal  $J$  on  $\text{cf}(\mu)$  satisfying  $J_{\text{cf}(\mu)}^{\text{bd}} \subseteq J$  such that  $\lambda =$

$$\text{tcf} \left( \prod_{\alpha < \text{cf}(\mu)} \lambda_\alpha / J \right) \text{ and } \max \text{pcf}\{\lambda_\beta : \beta < \alpha\} < \lambda_\alpha. \text{ By [Sh:g, II,3.5], there is}$$

modified:2005-07-06

revision:2005-07-05

(589)

$\langle f_\zeta : \zeta < \lambda \rangle$  which is  $<_J$ -increasing cofinal in  $\prod_{\alpha < \text{cf}(\mu)} \lambda_\alpha / J$  with  $|\{f_\zeta \upharpoonright \alpha : \zeta < \lambda\}| < \lambda_\alpha$ .

Easily  $\langle f_\zeta : \zeta < \lambda \rangle$  exemplifies  $\neg(B)$  : if  $A \in [\lambda]^\lambda$  and  $B =: \bigcup_{\zeta \in A} \text{Range}(f_\zeta)$  has

cardinality  $< \mu$  let  $g \in \prod \lambda_\alpha$  be:  $g(\alpha) = \sup(\lambda_\alpha \cap B)$  if  $< \lambda_\alpha$ , zero otherwise and let  $\alpha_0 = \text{Min}\{\alpha < \text{cf}(\mu) : \lambda_\alpha > |B|\}$ . So  $\alpha_0 < \text{cf}(\mu)$  and  $\zeta \in A \Rightarrow f_\zeta \upharpoonright [\alpha_0, \text{cf}(\mu)) < g$ , contradiction to “ $<_J$ -cofinal”.

Subcase 1b:  $\text{cf}(\mu) > \aleph_0$  and  $\text{pp}^+(\mu) = \text{pp}(\mu) = \lambda$ .

Use [Sh 513, §6] and finish as above.

Subcase 1c:  $\text{cf}(\mu) = \aleph_0$  and  $\lambda = \text{pp}^+(\mu) = \text{pp}(\mu) = \lambda$ .

Let  $\mathfrak{a}, \langle \mathfrak{b}_\tau : \tau \in R \rangle, \langle f_\tau : \tau \in R \rangle$  be as in [Sh 513, §6], so  $|\mathfrak{b}_\tau| = \aleph_0$ . Let  $\eta_\tau$  be an  $\omega$ -sequence of ordinals enumerating  $\text{Rang}(f_\tau)$  for  $\tau \in R$ , now  $\{\eta_\tau : \tau \in R\}$  is as required.

Case 2: Not Case 1.

So by [Sh:g, Ch.II,5.4], we have  $\theta \leq \mu < \lambda \Rightarrow \text{cov}(\mu, \theta, \kappa, \aleph_1) < \lambda$ .

As we are assuming  $\neg(A)$ , we can find  $\mu_0 < \lambda, a_\alpha \in [\mu_0]^{<\kappa}$  for  $\alpha < \lambda$  such that  $A \in [\lambda]^\lambda \Rightarrow |\bigcup_{\alpha \in A} a_\alpha| \geq \theta$ , but by the previous sentence we can find  $\mu_1 < \lambda$  and  $\{b_\beta : \beta < \mu_1\} \subseteq [\mu_0]^{<\theta}$  such that: every  $a \in [\mu_0]^{<\kappa}$  is included in the union of  $\leq \aleph_0$  sets from  $\{b_\beta : \beta < \mu_1\}$ . So we can find  $c_\alpha \in [\mu_1]^{\aleph_0}$  for  $\alpha < \lambda$  such that  $a_\alpha \subseteq \bigcup_{\beta \in c_\alpha} b_\beta$ . Now for  $A \in [\lambda]^\lambda$ , if  $|\bigcup_{\alpha \in A} c_\alpha| < \theta$  then

$$\begin{aligned} |\bigcup \{a_\alpha : \alpha \in A\}| &\leq |\bigcup_{\beta \in c_\alpha} b_\beta : \alpha \in A| \\ &= |\bigcup \{b_\beta : \beta \in \bigcup_{\alpha \in A} c_\alpha\}| < \min\{\sigma : \sigma = \text{cf}(\sigma) > |b_\beta| \text{ for } \beta < \mu_1\} \\ &\quad + |\bigcup_{\alpha \in A} c_\alpha|^+ \leq \theta + \theta = \theta \end{aligned}$$

contradicting the choice of  $\langle a_\alpha : \alpha < \lambda \rangle$ .

So

$$\begin{aligned} (*) \quad c_\alpha &\in [\mu_1]^{\leq \aleph_0}, \text{ for } \alpha < \lambda, \mu_1 < \lambda \text{ and} \\ A \in [\lambda]^\lambda &\Rightarrow |\bigcup_{\alpha \in A} c_\alpha| \geq \theta. \end{aligned}$$

Let  $\eta_\alpha$  be an  $\omega$ -sequence enumerating  $c_\alpha$ , so  $\langle \eta_\alpha : \alpha < \lambda \rangle$  is a counterexample to clause (B). □<sub>4.1</sub>

We concentrate below on  $\lambda, \theta, \kappa$  regular (others can be reduced to it).

**4.2 Definition.** Let  $\mathbf{T}$  be a complete first order theory; which is stable ( $\mathfrak{C}$  the monster model of  $\mathbf{T}$  and  $A, B, \dots$  denote subsets of  $\mathfrak{C}^{\text{eq}}$  of cardinality  $< \|\mathfrak{C}^{\text{eq}}\|$ ).

1)  $\text{Pr}_{\mathbf{T}}(\lambda, \chi, \theta)$  means:

- (\*) if  $A \subseteq \mathfrak{C}^{\text{eq}}, |A| = \lambda$  then we can find  $A' \subseteq A, |A'| = \chi$  and  $B', |B'| < \theta$  such that  $A'$  is independent over  $B'$   
(i.e.  $a \in A' \Rightarrow \text{tp}(a, B' \cup (A' \setminus \{a\}))$  does not fork over  $B'$ ).

2)  $\text{Pr}_{\mathbf{T}}^*(\lambda, \mu, \chi, \theta)$  means:

- (\*\*) if  $A \subseteq \mathfrak{C}^{\text{eq}}$  is independent over  $B$  where  $|A| = \lambda$  and  $|B| < \mu, B \subseteq \mathfrak{C}^{\text{eq}}$  then there are  $A' \subseteq A, |A'| = \chi$  and  $B' \subseteq B$  satisfying  $|B'| < \theta$  such that  $\text{tp}(A', B)$  does not fork over  $B'$  (hence  $A'$  is independent over  $B'$ ).

3)  $\text{Pr}_{\mathbf{T}}^*(\lambda, \chi, \theta)$  means  $\text{Pr}_{\mathbf{T}}^*(\lambda, \lambda, \chi, \theta)$ .

*4.3 Fact.* Assume  $\lambda$  is regular  $> \theta \geq \kappa_r(\mathbf{T})$  then

- (1) if  $\chi = \lambda$  then  $\text{Pr}_{\mathbf{T}}(\lambda, \chi, \theta) \Leftrightarrow \text{Pr}_{\mathbf{T}}^*(\lambda, \lambda, \chi, \theta)$
- (2) if  $\lambda \geq \chi \geq \mu \geq \theta$  then  $\text{Pr}_{\mathbf{T}}(\lambda, \chi, \theta) \Rightarrow \text{Pr}_{\mathbf{T}}^*(\lambda, \mu, \chi, \theta)$ .

*Proof.* 1) The direction  $\Leftarrow$  is by the proof in [Sh:a, III].

[In detail, let  $A, B$  be given (the  $B$  is not really necessary), such that  $\lambda = |A| > |B| + \kappa_r(\mathbf{T})$  so let  $A = \{a_i : i < \lambda\}$ ; define

$A_i =: \{a_j : j < i\}, S = \{i < \lambda : \text{cf}(i) \geq \kappa_r(\mathbf{T})\}$ , so by the definition of  $\kappa_r(\mathbf{T})$  for  $\alpha \in S$  there is  $j_\alpha < \alpha$  such that  $\text{tp}(a_\alpha, A_\alpha \cup B)$  does not fork over  $A_{j_\alpha} \cup B$  so for some  $j^*$  the set  $S' = \{\delta \in S : j_\delta = j^*\}$  is stationary, now apply the right side with  $\{a_\delta : \delta \in S'\}, A_{j^*} \cup B$ , here standing for  $A, B$  there].

The other direction  $\Rightarrow$  follows by part (2).

2) This is easy, too, by the non-forking calculus [Sh:a, III,Th.0.1 + ](0)-(4),pgs.82-84 but we give details. So we are given a set  $A \subseteq \mathfrak{C}^{\text{eq}}$  independent over  $B$ , where  $|A| = \lambda$  and  $|B| < \mu$ . As we are assuming  $\text{Pr}_{\mathbf{T}}(\lambda, \chi, \theta)$  there is  $A' \subseteq A, |A'| = \chi$  and  $B', |B'| < \theta$  such that  $A'$  is independent over  $B'$ . So for every finite  $\bar{c} \subseteq B$  for some  $A_{\bar{c}} \subseteq A'$  of cardinality  $< \kappa(\mathbf{T})$  ( $\leq \kappa_r(\mathbf{T})$ ) we have:  $A' \setminus A_{\bar{c}}$  is independent over  $B' \cup \bar{c}$ . So  $A^* = \bigcup \{A_{\bar{c}} : \bar{c} \subseteq B \text{ finite}\}$  has cardinality  $< \kappa_r(\mathbf{T}) + |B|^+ \leq \chi$  so necessarily

$A' \setminus A^*$  has cardinality  $\chi$  and it is independent over  $\cup\{\bar{c} : \bar{c} \subseteq B \text{ finite}\} \cup B' = B \cup B'$ .  $\square_{4.3}$

*4.4 Discussion.* So in order to understand the model theoretic property it suffices to prove the equivalence

$\text{Pr}_{\mathbf{T}}^*(\lambda, \mu, \chi, \theta) \Leftrightarrow \text{Pr}(\lambda, \mu, \chi, \theta, \kappa)$  with  $\kappa = \kappa_r(\mathbf{T})$ , where

**4.5 Definition.** Assume

(\*)  $\lambda \geq \max\{\mu, \chi\} \geq \min\{\mu, \chi\} \geq \theta \geq \kappa > \aleph_0$  and  $\mu > \theta$  and for simplicity  $\lambda, \theta, \kappa$  are regular if not said otherwise (as the general case can be reduced to this case).

1)  $\text{Pr}(\lambda, \mu, \chi, \theta, \kappa)$  is defined as follows: if  $u_\alpha \in [\mu]^{<\kappa}$  for  $\alpha < \lambda$  and  $|\bigcup_{\alpha < \lambda} u_\alpha| < \mu$

then there is  $Y \in [\lambda]^\chi$  such that  $|\bigcup_{\alpha \in Y} u_\alpha| < \theta$ ;

2)  $\text{Pr}^{\text{tr}}(\lambda, \mu, \chi, \theta, \kappa)$  is defined similarly but for some tree  $T$  each  $u_\alpha$  is a branch of  $T$ .

3) We write  $\text{Pr}(\lambda, \leq \mu, \chi, \theta, \kappa)$  for  $\text{Pr}(\lambda, \mu^+, \chi, \theta, \kappa)$  and similarly for  $\text{Pr}^{\text{tr}}$  and  $\text{Pr}_{\mathbf{T}}^*$ .

*4.6 Fact.* Assume  $\lambda, \mu, \chi, \theta, \kappa = \kappa_r(\mathbf{T})$  satisfies (\*) of Definition 4.5. Then

1)  $\text{Pr}(\lambda, \mu, \chi, \theta, \kappa_r(\mathbf{T})) \Rightarrow \text{Pr}_{\mathbf{T}}^*(\lambda, \mu, \chi, \theta) \Rightarrow \text{Pr}^{\text{tr}}(\lambda, \mu, \chi, \theta, \kappa_r(\mathbf{T}))$ .

2)  $\text{Pr}(\lambda, \chi, \chi, \theta, \kappa_r(\mathbf{T})) \Rightarrow \text{Pr}_{\mathbf{T}}(\lambda, \chi, \theta) \Rightarrow \text{Pr}^{\text{tr}}(\lambda, \chi, \chi, \theta, \kappa_r(\mathbf{T}))$ .

3) We have obvious monotonicity properties.

*Proof.* Straight.

1) First we prove the first implication so assume  $\text{Pr}(\lambda, \mu, \chi, \theta, \kappa_r(\mathbf{T}))$ , let  $\kappa = \kappa_r(\mathbf{T})$ , hence (\*) of 4.5 holds and we shall prove  $\text{Pr}_{\mathbf{T}}^*(\lambda, \mu, \chi, \theta)$ . So (see Definition 4.2(2)) we have  $A \subseteq \mathfrak{C}^{\text{eq}}$  is independent over  $B \subseteq \mathfrak{C}^{\text{eq}}$ ,  $|A| = \lambda$  and  $|B| < \mu$ . Let  $A = \{a_\alpha : \alpha < \lambda\}$  with no repetitions and  $B = \{b_j : j < j(*)\}$  so  $j(*) < \mu$ . For each  $\alpha < \lambda$ , there is a subset  $u_\alpha$  of  $j(*)$  of cardinality  $< \kappa_r(\mathbf{T}) = \kappa$  such that  $\text{tp}(a_\alpha, B)$  does not fork over  $\{b_j : j \in u_\alpha\}$ . So  $u_\alpha \in [\mu]^{<\kappa}$  and  $|\bigcup_{\alpha < \lambda} u_\alpha| \leq |j(*)| < \mu$  hence

as we are assuming  $\text{Pr}(\lambda, \mu, \chi, \theta, \kappa)$ , there is  $Y \in [\lambda]^\chi$  such that  $|\bigcup_{\alpha \in Y} u_\alpha| < \theta$ . Let

$B' = \{b_j : j \in \bigcup_{\alpha \in Y} u_\alpha\}$ ,  $A' = \{a_\alpha : \alpha \in Y\}$  so  $B' \subseteq B$ ,  $|B'| < \theta$  and  $A' \subseteq A$ ,  $|A'| = \chi$



and by the nonforking calculus,  $\text{tp}(A', B)$  does not fork over  $B'$  (even  $\{a_\alpha : \alpha \in Y\}$  is independent over  $(B, B')$ ).

Second, we prove the second implication, so we assume  $\text{Pr}_{\mathbf{T}}^*(\lambda, \mu, \chi, \theta)$  and we shall prove  $\text{Pr}^{\text{tr}}(\lambda, \mu, \chi, \theta, \kappa_r(\mathbf{T}))$ . Let  $\kappa = \kappa_r(\mathbf{T})$ .

Let  $T$  be a tree and for  $\alpha < \lambda, u_\alpha$  a branch,  $|u_\alpha| < \kappa, |\bigcup_{\alpha < \lambda} u_\alpha| < \mu$ . Without loss

of generality  $T = \bigcup_{\alpha < \lambda} u_\alpha, \lambda = \bigcup_{\zeta < \kappa} A_\zeta$ , where  $A_\zeta = \{\alpha : \text{otp}(u_\alpha) = \zeta\}$ . Without loss

of generality  $T \subseteq {}^{\kappa >} \mu, T = \bigcup_{\zeta < \kappa} T_\zeta$  where  $T_\zeta = \bigcup \{u_\alpha : \alpha \in A_\zeta\}$  and

$$\eta \in T_\zeta \setminus \{<>\} \Rightarrow \eta(0) = \zeta.$$

Now  $T_\zeta$  can be replaced by  $\{\eta \upharpoonright C_\zeta : \eta \in T_\zeta\}$  where  $0 \in C_\zeta, \text{otp}(C_\zeta) = 1 + \text{cf}(\zeta), \text{sup}(C) = \zeta$ . So without loss of generality

$$T = \bigcup \{T_\sigma : \sigma \in \text{Reg} \cap \kappa\}$$

$$<> \neq \eta \in T_\sigma \Rightarrow \eta(0) = \sigma.$$

Without loss of generality  $\lambda = \bigcup \{A_\sigma : \sigma \in \text{Reg} \cap \kappa\}$  and  $\bigcup_{\alpha \in A_\sigma} u_\alpha = T_\sigma$ . It

is enough to take care of one  $\sigma$  (otherwise a little more work is required). So without loss of generality:

$$\alpha < \lambda \Rightarrow \text{otp}(u_\alpha) = \sigma.$$

As  $\sigma = \text{cf}(\sigma) < \kappa$  there are  $A_i \subseteq \mathfrak{C}^{\text{eq}}$  such that  $\langle A_i : i \leq \sigma \rangle$  increases continuously and  $p \in S(A_\sigma)$  and for each  $i < \sigma$  the type  $p \upharpoonright A_{i+1}$  forks over  $A_i$  say  $\varphi(x, c_i) \in p \upharpoonright A_{i+1}$  forks over  $A_i$  and  $A_i = \{c_j : j < i\}$ , (recall we work in  $\mathfrak{C}^{\text{eq}}$ ).

By the nonforking calculus we can find  $\langle f_\eta : \eta \in T \rangle, f_\eta$  elementary mapping

$$\text{Dom}(f_\eta) = A_{\ell g(\eta)}$$

$\langle f_\eta : \eta \in T \rangle$  nonforking tree, that is

$$\nu \triangleleft \eta \Rightarrow \text{tp}(\text{Rang}(f_\eta), \bigcup \{\text{Rang}(f_\rho) : \rho \in T, \rho \upharpoonright (\ell g(\nu) + 1) \not\triangleleft \eta\})$$

does not fork over  $A_\nu$ .

For  $\alpha < \lambda$ , let  $g_\alpha = \bigcup \{f_\nu : \nu \in a_\alpha\}, A_\alpha = \bigcup_{\nu \in a_\alpha} \text{Rang}(f_\nu) = g_\alpha(A_\sigma)$  and  $p_\alpha = g_\alpha(p)$ .

Let  $b_\alpha \in \mathfrak{C}$  realize  $p_\alpha$  for  $\alpha < \lambda$  be such that:

$\text{tp}(b_\alpha, \bigcup_{\eta \in T} \text{Rang}(f_\eta) \cup \{b_\beta : \beta \neq \alpha\})$  does not fork over  $A_\alpha$ .

Now we apply  $\text{Pr}_T^*(\lambda, \mu, \chi, \theta)$  on

$$A = \{b_\alpha : \alpha < \lambda\}$$

$$B = \bigcup_{\eta \in T} \text{Rang}(f_\eta).$$

So there are  $A' \subseteq A, |A'| = \chi$  and  $B' \subseteq B, |B'| < \theta$ ,  $\text{tp}(A', B)$  does not fork over  $B'$ , hence (for some  $Y \in [\lambda]^\chi$ ) we have  $A' = \{a_\alpha : \alpha \in Y\}$  independent over  $B'$ . So there is  $T' \subseteq T_\alpha$  subtree such that  $|T'| = |B'| + \sigma < \theta$  and such that  $B' \subseteq \bigcup_{\rho \in T'} A_\rho$ .

Throwing “few” ( $< |B'| + \kappa_r(\mathbf{T})$ ) members of  $A'$  that is of  $Y$  we get  $A'$  independent over  $B'$  as by the nonforking calculus, if  $\alpha \in Y$  then  $\text{tp}(b_\alpha, \bigcup \text{Rang}(f_\eta))$  does not fork over  $\bigcup_{\eta \in T'} \text{Rang}(f_\eta)$  hence  $u_\alpha \subseteq T'$ . So clearly  $Y$  is as required.

2) By part (1) and 4.3.

3) Left to the reader. □<sub>4.6</sub>

**4.7 Discussion** So by 4.6(1) if  $\text{Pr}$  and  $\text{Pr}^{\text{tr}}$  are equivalent,  $\kappa = \kappa_r(\mathbf{T})$  then  $\text{Pr}_T^*$  is equivalent to them (for the suitable cardinal parameter, so we would like to prove such equivalence). Now Claim 4.1 gives the equivalence when  $\theta = \kappa_r(\mathbf{T}), \lambda = \chi = \text{cf}(\lambda)$  and “for every  $\mu < \lambda$ ”. We give below more general cases; e.g. if  $\lambda$  is a successor of regular or  $\{\delta < \lambda : \text{cf}(\delta) = \theta^*\} \in I(\lambda)$  or ...

**4.8 Fact.** Assume  $\lambda, \mu, \chi, \theta, \kappa$  are as in (\*) of Definition 4.5 and  $\mu^* \in [\theta, \mu)$  and  $\text{cf}(\mu^*) < \kappa$ .

0)  $\text{Pr}(\lambda, \mu, \chi, \theta, \kappa) \Rightarrow \text{Pr}^{\text{tr}}(\lambda, \mu, \chi, \theta, \kappa)$ .

[Why? Straight].

1) If  $\kappa < \lambda$  and  $\mu < \lambda$  and  $\text{cf}(\mu) \geq \kappa$ , then  $\text{Pr}(\lambda, \leq \mu, \chi, \theta, \kappa) \Leftrightarrow (\forall \mu_1 < \mu) \text{Pr}(\lambda, \leq \mu_1, \chi, \theta, \kappa)$ ; similarly for  $\text{Pr}^{\text{tr}}$ .

2) If  $\text{pp}(\mu^*) > \lambda$  then  $\neg \text{Pr}^{\text{tr}}(\lambda, \mu, \chi, \theta, \kappa)$  (by [Sh 355, 1.5A], see [Sh 513, 6.10]).

3) If  $\text{pp}(\mu^*) \geq \lambda$  and

(a)  $\{\delta < \lambda : \text{cf}(\delta) = \theta\} \in I[\lambda]$  or just

(a)<sup>-</sup> for some  $S \in I[\lambda], (\forall \delta \in S), \text{cf}(\delta) = \theta$  and

(a)<sub>S</sub> for every closed  $e \subseteq \lambda$  of order type  $\chi, e \cap S \neq \emptyset$ .

Then  $\neg \text{Pr}^{\text{tr}}(\lambda, \mu, \chi, \theta, \kappa)$ .

[Why? As in [Sh:g, Ch.VIII,6.4] based on [Sh:g, Ch.II,5.4] better still [Sh:g, Ch.II,3.5]].

- 4) If  $\lambda$  is a successor of regular and  $\theta^+ < \lambda$ , then the assumption (b) of part (3) holds (see [Sh:g, Ch.VIII,6.1] based on [Sh 351, §4]).
- 5) If  $\mu < \lambda$  and  $\text{cov}(\mu, \theta, \kappa, \aleph_1) < \lambda$  (equivalently  $(\forall \tau)[\theta < \tau \leq \mu \ \& \ \text{cf}(\tau) < \kappa \rightarrow \text{pp}_{\aleph_1\text{-complete}}(\tau) < \lambda]$ ), then  $\neg \text{Pr}(\lambda, \mu^+, \chi, \theta, \kappa)$  implies that for some  $\mu_1 \in (\mu, \lambda)$  we have  $\neg \text{Pr}(\lambda, \mu_1, \chi, \theta, \aleph_1)$  (as in Case 2 in the proof of 4.1).
- 6)  $\text{Pr}(\lambda, \mu, \chi, \theta, \aleph_1) \Leftrightarrow \text{Pr}^{\text{tr}}(\lambda, \mu, \chi, \theta, \aleph_1)$ .
- 7)  $\text{Pr}(\lambda, \mu, \lambda, \theta, \kappa)$  iff for every  $\tau \in [\theta, \mu)$  we have:  $\text{Pr}(\lambda, \leq \tau, \lambda, \tau, \kappa)$ ; similarly for  $\text{Pr}^{\text{tr}}$ .
- 8)  $\text{Pr}(\lambda, \leq \mu, \lambda, \theta, \kappa)$  iff  $\text{Pr}^{\text{tr}}(\lambda, \leq \mu, \lambda, \theta, \kappa)$  (by 4.1).

**4.9 Claim.** *Under GCH we get equivalence:*  $\text{Pr}(\lambda, \mu, \chi, \theta, \kappa) \Leftrightarrow \text{Pr}^{\text{tr}}(\lambda, \mu, \chi, \theta, \kappa)$ .

*Proof.*  $\text{Pr} \Rightarrow \text{Pr}^{\text{tr}}$  is trivial; so let us prove  $\neg \text{Pr} \Rightarrow \neg \text{Pr}^{\text{tr}}$ , so assume  $\{a_\alpha : \alpha < \lambda\} \subseteq [\mu]^{<\kappa}$  exemplifies  $\neg \text{Pr}(\lambda, \mu, \chi, \theta, \kappa)$ . Without loss of generality  $|a_\alpha| = \kappa^* < \kappa$ . By 4.8(1) without loss of generality  $\lambda > \mu$ , so necessarily

- (c)  $\lambda = \mu^+, \mu > \kappa^* \geq \text{cf}(\mu)$  or
- (d)  $\lambda = \mu^+, \kappa = \lambda$ .

In Case (a) let  $T$  be the set of sequences of bounded subsets of  $\mu$  each of cardinality  $\leq \kappa^*$  of length  $< \text{Min}\{\text{cf}(\mu), \kappa^*\}$ . For each  $\alpha < \lambda$  let  $\bar{b}^\alpha = \langle b_{\alpha, \varepsilon} : \varepsilon < \text{cf}(\mu) \rangle$  be a sequence, every initial segment is in  $T$  and  $a_\alpha = \bigcup_{\varepsilon < \text{cf}(\mu)} b_{\alpha, \varepsilon}$ , so

$t_\alpha = \{\bar{b}^\alpha \upharpoonright \zeta : \zeta < \text{cf}(\mu)\}$  is a  $\text{cf}(\mu)$ -branch of  $T$ , and it should be clear.

*4.10 Remark.* We can get an independence result by instances of Chang's Conjecture (so the consistency strength seems somewhat more than huge cardinals, see Foreman [For], Levinski Magidor Shelah [LMSH 198]).

modified:2005-07-06

(589) revision:2005-07-05

§5 CARDINAL INVARIANTS FOR GENERAL REGULAR  
CARDINALS: RESTRICTIONS ON THE DEPTH

Cummings and Shelah [CuSh 541] prove that there are no non-trivial restrictions on some cardinal invariants like  $\mathfrak{b}_\lambda$  and  $\mathfrak{d}_\lambda$ , even for all regular cardinals simultaneously; i.e., on functions like  $\langle \mathfrak{b}_\lambda : \lambda \in \text{Reg} \rangle$ . But not everything is independent of ZFC. Consider the cardinal invariants  $\mathfrak{dp}_\lambda^{\ell+}$ , defined below.

**5.1 Definition.** 1) We are given an ideal  $J$  on a regular cardinal  $\lambda$ .

If  $\lambda > \aleph_0$  let

$$\mathfrak{dp}_\lambda^{1+} = \text{Min} \left\{ \mu : \text{there is no sequence } \langle C_\alpha : \alpha < \mu \rangle \text{ such that:} \right. \\ \left. \begin{array}{l} (a) \ C_\alpha \text{ is a club of } \lambda, \\ (b) \ \beta < \alpha \Rightarrow |C_\alpha \setminus C_\beta| < \lambda, \\ (c) \ C_{\alpha+1} \subseteq \text{acc}(C_\alpha) \end{array} \right\},$$

where  $\text{acc}(C)$  is the set of accumulation points of  $C$ .

If  $\lambda \geq \aleph_0$  let

$$\mathfrak{dp}_{\lambda, J}^{2+} = \text{Min} \left\{ \mu : \text{there are no } f_\alpha \in {}^\lambda \lambda \text{ for} \right. \\ \left. \alpha < \mu \text{ such that } \alpha < \beta < \mu \Rightarrow f_\alpha <_J f_\beta \right\}.$$

If  $\lambda \geq \aleph_0$  let

$$\mathfrak{dp}_{\lambda, J}^{3+} = \text{Min} \left\{ \mu : \text{there is no sequence } \langle A_\alpha : \alpha < \mu \rangle \text{ such that:} \right. \\ \left. \begin{array}{l} A_\alpha \in J^+ \text{ and} \\ \alpha < \beta < \mu \Rightarrow [A_\beta \setminus A_\alpha \in J^+ \ \& \ A_\alpha \setminus A_\beta \in J] \end{array} \right\}.$$

If  $J = J_\lambda^{\text{bd}}$ , we may omit it. We can replace  $J$  by its dual filter.

2) For  $\ell \in \{1, 2, 3\}$  let  $\mathfrak{dp}_\lambda^\ell = \sup\{\mu : \mu < \mathfrak{dp}_\lambda^{\ell+}\}$ .

3) For a regular cardinal  $\lambda$  let

$$\mathfrak{d}_\lambda = \text{Min} \left\{ |F| : F \subseteq {}^\lambda \lambda \text{ and } (\forall g \in {}^\lambda \lambda)(\exists f \in F)(g <_{J_\lambda^{\text{bd}}} f) \right\}$$

(equivalently  $g < f$ )

$$\mathfrak{b}_\lambda = \text{Min} \left\{ |F| : F \subseteq {}^\lambda \lambda \text{ and } \neg(\exists g \in {}^\lambda \lambda)(\forall f \in F)[f <_{J_\lambda^{\text{bd}}} g] \right\}.$$

We shall prove here that in the “neighborhood” of singular cardinals there are some connections between the  $\mathfrak{d}\mathfrak{p}_\lambda^{\ell+}$ ’s (hence by monotonicity, also with the  $\mathfrak{b}_\lambda$ ’s).

We first note connections for “one  $\lambda$ ”.

5.2 *Fact.* 1) If  $\lambda = \text{cf}(\lambda) > \aleph_0$  then

$$\mathfrak{b}_\lambda < \mathfrak{d}\mathfrak{p}_\lambda^{1+} \leq \mathfrak{d}\mathfrak{p}_\lambda^{2+} \leq \mathfrak{d}\mathfrak{p}_\lambda^{3+}.$$

2)  $\mathfrak{b}_{\aleph_0} < \mathfrak{d}\mathfrak{p}_{\aleph_0}^{2+} = \mathfrak{d}\mathfrak{p}_{\aleph_0}^{3+}$ .

3) In the definition of  $\mathfrak{d}\mathfrak{p}_\lambda^{1+}$ ,  $C_{\alpha+1} \subseteq \text{acc}(C_\alpha) \text{ mod } J_\lambda^{\text{bd}}$  suffices.

*Proof.* 1) First inequality:  $\mathfrak{b}_\lambda < \mathfrak{d}\mathfrak{p}_\lambda^{1+}$ .

We choose by induction on  $\alpha < \mathfrak{b}_\lambda$ , a club  $C_\alpha$  of  $\lambda$  such that  $\beta < \alpha \Rightarrow |C_\alpha \setminus C_\beta| < \lambda$  and  $C_{\beta+1} \subseteq \text{acc}(C_\beta)$ .

For  $\alpha = 0$  let  $C_\alpha = \lambda$ , for  $\alpha = \beta+1$  let  $C_\alpha = \text{acc}(C_\beta)$ , and for  $\alpha$  limit let, for each  $\beta < \alpha$ ,  $f_\beta \in {}^\lambda \lambda$  be defined by  $f_\beta(i) = \text{Min}(C_\alpha \setminus (i+1))$ . So  $\{f_\beta : \beta < \alpha\}$  is a subset of  ${}^\lambda \lambda$  of cardinality  $\leq |\alpha| < \mathfrak{b}_\lambda$ , so there is  $g_\alpha \in {}^\lambda \lambda$  such that  $\beta < \alpha \Rightarrow f_\beta <_{J_\lambda^{\text{bd}}} g_\alpha$ .

Lastly, let  $C_\alpha = \{\delta < \lambda : \delta \text{ a limit ordinal such that } (\forall \zeta < \delta)[g_\alpha(\zeta) < \delta]\}$ , now  $C_\alpha$  is as required.

So  $\langle C_\alpha : \alpha < \mathfrak{b}_\lambda \rangle$  exemplifies  $\mathfrak{b}_\lambda < \mathfrak{d}\mathfrak{p}_\lambda^{1+}$ .

Second inequality:  $\mathfrak{d}\mathfrak{p}_\lambda^{1+} \leq \mathfrak{d}\mathfrak{p}_\lambda^{2+}$

Assume  $\mu < \mathfrak{d}\mathfrak{p}_\lambda^{1+}$ . Let  $\langle C_\alpha : \alpha < \mu \rangle$  exemplify it, and let us define for  $\alpha < \mu$  the function  $f_\alpha \in {}^\lambda \lambda$  by:  $f_\alpha(\zeta)$  is the  $(\zeta + 1)$ -th member of  $C_\alpha$ ; clearly  $f_\alpha \in {}^\lambda \lambda$  and  $f_\alpha$  is strictly increasing. Also, if  $\beta < \alpha$  then  $C_\alpha \setminus C_\beta$  is a bounded subset of  $\lambda$ , say by  $\delta_1$ , and there is  $\delta_2 \in (\delta_1, \lambda)$  such that  $\text{otp}(\delta_2 \cap C_\beta) = \delta_2$ . So for every  $\zeta \in [\delta_2, \lambda)$  clearly  $f_\beta(\zeta) =$  the  $(\zeta + 1)$ -th member of  $C_\beta =$  the  $(\zeta + 1)$ -th member of  $C_\beta \setminus \delta_1 \leq$  the  $(\zeta + 1)$ -th member of  $C_\alpha$ . So  $\beta < \alpha \Rightarrow f_\beta \leq_{J_\lambda^{\text{bd}}} f_\alpha$ . Lastly, for  $\alpha < \mu$ ,  $C_{\alpha+1} \subseteq \text{acc}(C_\alpha)$  hence  $f_\alpha(\zeta) =$  the  $(\zeta + 1)$ -th member of  $C_\alpha <$  the  $(\zeta + \omega)$ -th member of  $C_\alpha \leq$  the  $(\zeta + 1)$ -th member of  $\text{acc}(C_\alpha) \leq$  the  $(\zeta + 1)$ -th member of  $C_{\alpha+1}$ . So  $\beta < \alpha \Rightarrow f_\beta <_{J_\lambda^{\text{bd}}} f_{\beta+1} \leq_{J_\lambda^{\text{bd}}} f_\alpha$ , so  $\langle f_\alpha : \alpha < \lambda \rangle$  exemplifies  $\mu < \mathfrak{d}\mathfrak{p}_\lambda^{2+}$ .

Third inequality:  $\mathfrak{dp}_\lambda^{2+} \leq \mathfrak{dp}_\lambda^{3+}$

Assume  $\mu < \mathfrak{dp}_\lambda^{2+}$  and let  $\langle f_\alpha : \alpha < \mu \rangle$  exemplify this.

Let  $c : \lambda \times \lambda \rightarrow \lambda$  be one to one and let

$$A_\alpha = \{c(\zeta, \xi) : \zeta < \lambda \text{ and } \xi < f_\alpha(\zeta)\}.$$

Now  $\langle A_\alpha : \alpha < \mu \rangle$  exemplifies  $\mu < \mathfrak{dp}_\lambda^{3+}$ .

2), 3) Easy. □<sub>5.2</sub>

**5.3 Observation.** Suppose  $\lambda = \text{cf}(\lambda) > \aleph_0$ .

1) If  $\langle f_\alpha : \alpha \leq \gamma^* \rangle$  is  $<_{J_\lambda^{\text{bd}}}$ -increasing then we can find a sequence  $\langle C_\alpha : \alpha < \gamma^* \rangle$  of clubs of  $\lambda$ , such that  $\alpha < \beta \Rightarrow |C_\alpha \setminus C_\beta| < \lambda$  and  $C_{\alpha+1} \subseteq \text{acc}(C_\alpha) \text{ mod } J_\lambda^{\text{bd}}$ .

2)  $\mathfrak{dp}_\lambda^{1+} = \mathfrak{dp}_\lambda^{+2}$  or for some  $\mu$ ,  $\mathfrak{dp}_\lambda^{1+} = \mu^+$ ,  $\mathfrak{dp}_\lambda^{2+} = \mu^{++}$  (moreover though there is in  $({}^\lambda \lambda, <_{J_\lambda^{\text{bd}}})$  an increasing sequence of length  $\mu^+$ , there is none of length  $\mu^+ + 1$ ).

*Proof.* 1) Let

$$C^* = \left\{ \delta < \lambda : \delta \text{ a limit ordinal and } (\forall \beta < \delta) f_{\gamma^*}(\beta) < \delta \right. \\ \left. \text{and } \omega^\delta = \delta \text{ (ordinal exponentiation)} \right\};$$

this is a club of  $\lambda$ .

For each  $\alpha < \gamma^*$  let

$$C_\alpha = \left\{ \delta + \omega^{f_\alpha(\delta)} \cdot \beta : \delta \in C^* \text{ and } \beta < f_\alpha(\delta) \right. \\ \left. \text{and } f_\alpha(\delta) < f_{\gamma^*}(\delta) \right\}.$$

2) Follows. □<sub>5.3</sub>

Now we come to our main concern.

**5.4 Theorem.** *Assume*

- (a)  $\kappa$  is regular uncountable,  $\ell \in \{1, 2, 3\}$
- (b)  $\langle \mu_i : i < \kappa \rangle$  is (strictly) increasing continuous with limit  $\mu$ ,  
 $\lambda_i = \mu_i^+, \lambda = \mu^+$
- (c)  $2^\kappa < \mu$  and  $\mu_i^\kappa < \mu$
- (d)  $D$  a normal filter on  $\kappa$
- (e)  $\theta_i < \mathfrak{dp}_{\lambda_i}^{\ell+}$  and  $\theta = \text{tcf}(\prod_{i < \kappa} \theta_i / D)$  or just  
 $\theta < \text{Depth}^+(\prod_{i < \kappa} \theta_i / D)$ .

Then  $\theta < \mathfrak{dp}_\lambda^{\ell+}$ .

*Proof.* By 5.15, 5.16, 5.6 below for  $\ell = 1, 2, 3$  respectively (the conditions there are easily checked). □<sub>5.4</sub>

*5.5 Remark.* 1) Concerning assumption (e), e.g. if  $2^{\mu_i} = \mu_i^{+5}$  and  $2^\mu = \mu^{+5}$ , then necessarily  $\mu^{+\ell} = \text{tcf}(\prod_{i < \kappa} \mu_i^{+\ell} / D)$  for  $\ell = 1, \dots, 5$  and so  $\bigwedge_{i < \kappa} \mathfrak{dp}_{\lambda_i}^{\ell+} = 2^{\mu_i} \Rightarrow \mathfrak{dp}_\lambda = 2^\mu$  and we can use  $\mu_i = (2^\kappa)^{+i}, \lambda_i = \mu_i^+, \theta_i = \mu_i^{+5}, \theta = \mu^{+5}$ .

So this theorem really says that the function  $\lambda \mapsto \mathfrak{dp}_\lambda$  has more than the cardinality exponentiation restrictions.

2) Note that Theorem 5.4 is trivial if  $\prod_{i < \kappa} \lambda_i = 2^\mu = \lambda$ , so (see [Sh:g, V]) it is natural

to assume  $E =: \{D' : D' \text{ a normal filter on } \kappa\}$  is nice, but this will not be used.

3) Note that the proof of 5.16 (i.e. the case  $\ell = 2$ ) does not depend on the longer proof of 5.6, whereas the proof of 5.15 does.

4) Recall that for an  $\aleph_1$ -complete filter  $D$ , say on  $\kappa$ , and  $f \in {}^\kappa \text{Ord}$  we define  $\|f\|_D$  by  $\|f\|_D = \cup\{\|g\|_D + 1 : g \in {}^\kappa \text{Ord} \text{ and } g <_D f\}$ .

5) Below we shall use the assumption

(\*)  $\|\lambda\|_{D+A} = \lambda$  for every  $A \in D^+$ .

This is not a strong assumption as

(a) if SCH holds, then the only case of interest is if  $\langle \chi_i : i < \kappa \rangle$  is increasing continuous with limit  $\chi$  and  $\|\langle \chi_i^+ : i < \kappa \rangle\|_D = \chi^+$  for any normal filter  $D$  on  $\kappa$ ; so our statements degenerate and say nothing,

(b) if SCH fails, there are nice filters for which this phenomenon is “popular” see [Sh:g, V,1.13,3.10] (see more in **5.18**).

→ scite{5.11A} ambiguous

modified:2005-07-06

revision:2005-07-05

(589)

**5.6 Theorem.** *Assume*

- (a)  $D$  is an  $\aleph_1$ -complete filter on  $\kappa$
- (b)  $\langle \lambda_i : i < \kappa \rangle$  is a sequence of regular cardinals  $> (2^\kappa)^+$
- (c)  $\|\langle \lambda_i : i < \kappa \rangle\|_{D+A} = \lambda$  for  $A \in D^+$
- (d)  $\mu_i < \mathfrak{dp}_{\lambda_i}^{3+}$
- (e)  $\mu = \text{tcf}(\prod \mu_i / D)$  or at least
- (e<sup>-</sup>)  $\mu < \text{Depth}^+(\prod \mu_i, <_D)$  and  $\mu > 2^\kappa$ .

Then  $\mu < \mathfrak{dp}_\lambda^{3+}$ .

*Remark.* Why not assume just  $\|f\|_D = \lambda$  for  $f =: \langle \lambda_i : i < \kappa \rangle$ ? Note that  $\text{cla}_f^\alpha(f, A)$ , see below, does not make much sense.

We delay the proof of 5.6 until we complete some preliminary work.

*5.7 Fact.* Assuming 5.6(a), for any  $f \in {}^\kappa(\text{Ord} \setminus (2^\kappa)^+)$  we have:  $T_D(f)$  is smaller or equal to the cardinality of  $\|f\|_D$  remembering (5.5(4) above and)

$$T_D(f) = \sup \left\{ |F| : F \subseteq \prod_{i < \kappa} f(i) \text{ and } f \neq g \in F \Rightarrow f \neq_D g \right\}.$$

*Proof.* Why? Let  $F$  be as in the definition of  $T_D(f)$ , note:  $f_i \neq_D f_j$  &  $f_i \leq_D f_j \Rightarrow f_i <_D f_j$ . Note that as  $i < \kappa \Rightarrow f(i) \geq (2^\kappa)^+$ , necessarily  $|F| > 2^\kappa$ . Now for each ordinal  $\alpha$  let  $F^{[\alpha]} =: \{f \in F : \|f\|_D = \alpha\}$ . Clearly  $F^{[\alpha]}$  has at most  $2^\kappa$  members, as otherwise some  $f_i \in F^{[\alpha]}$  for  $i < (2^\kappa)^+$  are pairwise distinct so for some  $i < j$ ,  $f_i <_D f_j$  (by [Sh 111, §2]) or simply use Erdős-Rado on  $c(i, j) = \min\{\zeta < \kappa : f_i(\zeta) > f_j(\zeta)\}$ .

So  $\|f\|_D \geq \sup\{\|g\|_D : g \in F\} \geq \text{otp}\{\alpha : F^{[\alpha]} \neq \emptyset\} \geq |\{\alpha : F^{[\alpha]} \neq \emptyset\}| \geq |F|/2^\kappa = |F|$ . So  $\|f\|_D \geq T_D(f)$ . □<sub>5.7</sub>



**5.8 Definition.** For  $f \in {}^\kappa\text{Ord}$  (natural to be mainly interested in the case  $0 \notin \text{Rang}(f)$ ) and  $D$  an  $\aleph_1$ -complete filter on  $\kappa$  let  $\prod_{i < \kappa}^* f(i) = \{g : \text{Dom}(g) = \kappa, f(i) > 0 \Rightarrow g(i) < f(i) \text{ and } f(i) = 0 \Rightarrow g(i) = 0\}$  and

$$1) \text{ cla}(f, D) = \left\{ (g, A) : g \in \prod_{i < \kappa}^* f(i) \text{ and } A \in D^+ \right\}$$

$$\text{cla}^\alpha(f, D) = \{(g, A) \in \text{cla}(f, D) : \|g\|_{D+A} = \alpha\}.$$

Here “cla” abbreviates “class”.

2) For  $(g, A) \in \text{cla}(f, D)$  let

$$J_D(g, A) = \{B \subseteq \kappa : \text{if } B \in (D + A)^+ \text{ then } \|g\|_{(D+A)+B} > \|g\|_{D+A}\}.$$

3) We say  $(g', A') \approx (g'', A'')$  if (both are in  $\text{cla}(f, D)$  and)  $A' = A'' \text{ mod } D$  and  $J_D(g', A') = J_D(g'', A'')$  and  $g' = g'' \text{ mod } J_D(g', A')$ .

4) For  $I$  an ideal on  $\kappa$  disjoint to  $D$  we let

$$I * D = \{A \subseteq \kappa : \text{for some } X \in D \text{ we have } A \cap X \in I\},$$

(usually we have  $\{\kappa \setminus A : A \in D\} \subseteq I$  so  $I * D = I$ ) and let

$$\text{cla}_I(f, D) = \{(g, A) : g \in \prod_{i < \kappa}^* f(i) \text{ and } A \in (I * D)^+\}.$$

5) On  $\text{cla}_I(f, D)$  we define a relation  $\approx_I$

$(g_1, A_1) \approx_I (g_2, A_2)$  if:

(a)  $A_1 = A_2 \text{ mod } D$  and

(b) there is  $B_0 \in I$  such that: if  $B_0 \subseteq B \in I$  then

$$\|g_1\|_{(D+A_1)+(\kappa \setminus B)} = \|g_2\|_{(D+A_2)+(\kappa \setminus B)} \text{ and } J_{(D+A_1)+(\kappa \setminus B)}(g_1, A_1) = J_{(D+A_2)+(\kappa \setminus B)}(g_2, A_2).$$

$$6) \quad J_{D,I}(g_1, A_1) = \{A \subseteq \kappa : \text{for some } B_0 \in I \text{ if } B_0 \subseteq B \in I \text{ we have } A \in J_{(D+A_1)+(\kappa \setminus B)}(g_1, A_1)\}.$$

7) Let  $\text{com}(D)$  be the maximal  $\theta$  such that  $D$  is  $\theta$ -complete.

5.9 *Fact.* For  $f \in {}^\kappa\text{Ord}$  and  $D$  an  $\aleph_1$ -complete filter on  $\kappa$  and  $A \in D^+$ :

- 0) If  $f_1 \leq f_2$  then  $\text{cla}(f_1, D) \subseteq \text{cla}(f_2, D)$  and for  $g', g'' \in \prod_{i < \kappa}^* f_1(i), A \subseteq \kappa$  we have  $(g', A) \approx (g'', A)$  in  $\text{cla}(f_1, D)$  iff  $(g', A) \approx (g'', A)$  in  $\text{cla}(f_2, D)$  (so we shall be careless about this).
- 1)  $J_D(g, A)$  is an ideal on  $\kappa$ ,  $\text{com}(D)$ -complete, and normal if  $D$  is normal.
- 2)  $A$  does not belong to  $J_D(g, A)$ , and it includes  $\{B \subseteq \kappa : B = \emptyset \text{ mod } (D + A)\}$ . If  $B \in J_D^+(g, A)$  then  $A \cap B \in D^+$  and  $\|g\|_{D+(A \cap B)} = \|g\|_{D+A}$ .
- 3)  $\approx$  is an equivalence relation on  $\text{cla}(f, D)$ , similarly  $\approx_I$  on  $\text{cla}_I(f, D)$ .
- 4) Assume

(i)  $(g, A) \in \text{cla}^\alpha(f, D), g' \in \prod_{i < \kappa}^* f(i)$  and

- (ii) (a)  $g' = g \text{ mod } (D + A)$  or  
 (b) for some  $B \in J_D(g, A)$  we have  $\alpha \in B \Rightarrow g'(\alpha) > \|g\|_D$   
 (or just  $\|g\|_{D+A} < \|g'\|_{D+B}$ ) and  
 $g' \upharpoonright (\kappa \setminus B) = g \upharpoonright (\kappa \setminus B) \text{ mod } D$ .

Then  $(g', A) \approx (g, A)$ .

- 5) For each  $\alpha$ , in  $\text{cla}^\alpha(f, D)/\approx$  there are at most  $2^\kappa$  classes.
- 6) For  $f \in {}^\kappa(\text{Ord})$ , in  $\text{cla}(f, D)/\approx$  there are at most  $2^\kappa + \sup_{A \in D^+} \|f\|_{D+A}$  classes.

*Proof.* 0) Easy.

- 1) Straight (e.g., it is an ideal as for  $B \subseteq \kappa$  we have  $\|g\|_D = \text{Min}\{\|g\|_{D+A}, \|g\|_{D+(\kappa-A)}\}$ , where we stipulate  $\|g\|_{\mathcal{P}(\kappa)} = \infty$  see [Sh 71]).
- 2) Check.
- 3) Check.
- 4) Check.
- 5) We can work also in  $\text{cla}^\alpha(f + 1, D)$  (this change gives more elements and by (0) it preserves  $\approx$ ). Assume  $\alpha$  is a counterexample (note that " $\leq 2^{2^\kappa}$ " is totally immediate). Let  $\chi$  be large enough; choose  $N \prec (\mathcal{H}(\chi), \in, <_\chi^*)$  of cardinality  $2^\kappa$  such that  $\{f, D, \kappa, \alpha\} \in N$  and  ${}^\kappa N \subseteq N$ . So necessarily there is  $(g, A) \in \text{cla}^\alpha(f, D)$  such that the equivalence class  $(g, A)/\approx$  does not belong to  $N$ , by the definition of  $\text{cla}^\alpha$ , clearly  $\|g\|_{D+A} = \alpha$ . Let  $B =: \{i < \kappa : g(i) \notin N\}$ .

Case 1:  $B \in J_D(g, A)$ .

Let  $g' \in \prod_{i < \kappa} (f(i) + 1)$  be defined by:  $g'(i) = g(i)$  if  $i \in \kappa \setminus B$  and  $g'(i) = f(i)$  if  $i \in B$ . By part (4) we have  $(g', A) \approx (g, A)$  and by the choice of  $N$  we have

$(g', A) \in N$  as  $A \in \mathcal{P}(\kappa) \subseteq N, g' \in N$  (as  $\text{Rang}(g') \subseteq N$  &  ${}^\kappa N \subseteq N$ ). Thus, there is  $(g', A) \in N$  such that  $(g', A) \approx (g, A)$  as required.

Case 2:  $B \notin J_D(g, A)$ .

Let  $g' \in {}^\kappa \text{Ord}$  be:  $g'(i) = \text{Min}(N \cap (f(i) + 1) \setminus g(i)) \leq f(i)$  if  $i \in B, g'(i) = g(i)$  if  $i \notin B$  (note:  $f(i) \in N, g(i) < f(i)$  so  $g'$  is well defined).

Clearly  $g' \in N$ , (as  $\text{Rang}(g') \subseteq N$  and  ${}^\kappa N \subseteq N$ ), and

$$(\mathcal{H}(\chi), \in, <^*_\chi) \models (\exists x)(x \in \prod_{i < \kappa}^* f(i) \wedge (\forall i \in \kappa \setminus B)(x(i) = g'(i)) \wedge (\forall i \in B)(x(i) < g'(i)) \wedge \|x\|_{D+(A \cap B)} = \alpha).$$

(Why? Because  $x = g$  is like that, last equality as  $B \notin J_D(g, A)$ ). So there is such  $x$  in  $N$ , call it  $g''$ . So  $g'' \in \prod_{i < \kappa} (f(i) + 1)$  and  $\|g''\|_{D+(A \cap B)} = \alpha$  and for

$i \in B, g''(i) \in g'(i) \cap N$  hence  $g''(i) < g(i)$  by the definition of  $g'(i)$ .

So  $g'' < g \text{ mod } D+(A \cap B)$ , but this contradicts  $\|g''\|_{D+(A \cap B)} = \alpha = \|g\|_{D+(A \cap B)}$ , the last equality as  $B \notin J_D(g, A)$ .

6) Immediate from (5). □<sub>5.9</sub>

*5.10 Fact.* Assume  $f \in {}^\kappa \text{Ord}$  and  $D$  an  $\aleph_1$ -complete filter on  $\kappa$  and  $I$  an  $\text{com}(D)$ -complete ideal on  $\kappa$ .

1) If  $(g, A) \in \text{cla}_I(f, D)$  then  $J_{D,I}(g, A)$  is an ideal on  $\kappa$ , which is  $\text{com}(D)$ -complete and normal if  $D, I$  are normal.

If  $B \in (J_{D,I}(g, A))^+$  then  $\|g\|_{D+(A \cap B)} = \|g\|_{D+A}$ , and  $(D + (A \cap B)) \cap I = \emptyset$ .

2)  $\approx_I$  is an equivalence relation on  $\text{cla}(f, D)$ .

3) If  $(g, A) \in \text{cla}(f, D)$  and  $g' \in \prod_{i < \kappa}^* f(i)$  and  $g' = g \text{ mod } J_{D,I}(g, A)$  then for some  $A'$  we have  $(g', A') \approx_I (g, A)$  so  $(g', A') \in \text{cla}(f, D)$  and  $\|g'\|_{D+A'} = \|g\|_{D+A'}$  (in fact  $A' = \{i \in A : g'(i) = g(i)\}$  is O.K.).

*Proof.* Easy.

*5.11 Fact.* Let  $\kappa, f, D$  be as in 5.10.

1) If  $f_\zeta \in {}^\kappa \text{Ord}$ , for  $\zeta \leq \delta, \text{cf}(\delta) > \kappa$  and for each  $i$  the sequence  $\langle f_\zeta(i) : \zeta \leq \delta \rangle$  is increasing continuous then  $\|f_\delta\|_D = \sup_{\zeta < \delta} \|f_\zeta\|_D$ .

2) If  $\delta = \|f\|_D, \text{cf}(\delta) > 2^\kappa$  then  $\{i : \text{cf}(f(i)) \leq 2^\kappa\} \in J_D(f, \kappa)$ .

- 3) If  $\|f\|_D = \delta$ ,  $A \in J_D^+(f, \kappa)$  then  $\prod_{i < \kappa}^* f(i)/(D + A)$  is not  $(\text{cf}(\delta))^+$ -directed.
- 4) If  $\|f\|_D = \delta$  and  $A \in J_D^+(f, \kappa)$  then  $\text{cf}(\delta) \leq \text{cf}(\prod_{i < \kappa}^* f(i)/(D + A))$ .
- 5) If  $\|f\|_D = \delta$  and  $A \subseteq \kappa$ ,  $(\forall i \in A) \text{cf}(f(i)) > \kappa$  and  $\max \text{pcf}\{f(i) : i \in A\} < \text{cf}(\delta)$   
 (or just  $\text{cf}(\delta) > \max\{\text{cf} \prod_{i < \kappa}^* f(i)/D' : D' \text{ an ultrafilter extending } D + A\}$ ) then  
 $A \in J_D(f, \kappa)$ .
- 6) If  $\|f\|_D = \delta$ ,  $\text{cf}(\delta) > 2^\kappa$ , then  $\prod_{i < \kappa}^* f(i)/J_D(f, \kappa)$  is  $\text{cf}(\delta)$ -directed.
- 7) If  $\|f\|_D = \delta$ ,  $\text{cf}(\delta) > 2^\kappa$ , then for some  $A \in J_D^+(f, \kappa)$  we have  
 $\prod_{i < \kappa}^* f(i)/J_D(f, \kappa) + (\kappa \setminus A)$  has true cofinality  $\text{cf}(\delta)$ .
- 8) Assume  $\|f\|_D = \lambda = \text{cf}(\lambda) > 2^\kappa$ .  
 Then  $(\forall A \in D^+)(\|f\|_{D+A} = \lambda)$  implies  $\text{tcf}(\prod_{i < \kappa}^* f(i)/D) = \lambda$ .
- 9) If  $\|f\|_D = \delta$ ,  $\text{cf}(\delta) > 2^\kappa$  then  $\text{tcf} \prod_{i < \kappa}^* f(i)/J_D(f, \kappa) = \text{cf}(\delta)$ .

*Proof.* 1) Let  $g <_D f_\delta$ , so  $A = \{i < \kappa : g(i) < f_\delta(i)\} \in D$ , now for each  $i \in A$  we have  $g(i) < f_\delta(i) \Rightarrow (\exists \alpha < \delta)(g(i) < f_\alpha(i)) \Rightarrow$  there is  $\alpha_i < \delta$  such that  $(\forall \alpha)[\alpha_i \leq \alpha \leq \delta \Rightarrow g(i) < f_\alpha(i)]$ . Hence  $\alpha(*) =: \sup\{\alpha_i : i \in A\} < \delta$  as  $\text{cf}(\delta) > \kappa$ , so  $g <_D f_{\alpha(*)}$  hence  $\|g\|_D < \|f_{\alpha(*)}\|_D$ ; this suffices for one inequality, the other is trivial.

2) Let  $A = \{i : \text{cf}(i) \leq 2^\kappa\}$ , and assume toward contradiction that  $A \in J_D^+(f, \kappa)$ . For each  $i \in A$  let  $C_i \subseteq f(i)$  be unbounded of order type  $\text{cf}(f(i)) \leq 2^\kappa$ .

Let  $F = \{g \in \prod_{i < \kappa}^* (f(i) + 1) : \text{if } i \in A \text{ then } g(i) \in C_i, \text{ if } i \in \kappa \setminus A \text{ then } g(i) = f(i)\}$ .

So  $|F| \leq 2^\kappa$  and:

(\*) if  $g <_{D+A} f$  then for some  $g' \in F$ ,  $g <_{D+A} g'$ ,

hence  $\delta = \|f\|_{D+A} = \sup\{\|g\|_{D+A} : g \in F\}$  but the supremum is on  $\leq |F| < \text{cf}(\delta)$  ordinals each  $< \delta$  because  $g' \in F \Rightarrow g' <_{D+A} f$  as  $\|f\|_D = \delta \Rightarrow f \neq_D 0_\kappa$ , contradiction to  $\text{cf}(\delta) > 2^\kappa$ .

- 3) Assume this fails, so  $\|f\|_D = \delta$ ,  $A \in J_D^+(f, \kappa)$  and  $\prod_{i < \kappa}^* f(i)/(D + A)$  is  $(\text{cf}(\delta))^+$ -directed. Let  $C \subseteq \delta$  be unbounded of order type  $\text{cf}(\delta)$ ; as  $\|f\|_{D+A} = \delta$  (because  $A \in J_D^+(f, A)$ ) for each  $\alpha \in C$  there is  $f_\alpha <_{D+A} f$  such that  $\|f_\alpha\|_{D+A} \geq \alpha$  (even  $= \alpha$  by the definition of  $\| - \|_{D+A}$ ). As  $\prod_{i < \kappa}^* f(i)/(D + A)$  is  $(\text{cf}(\delta))^+$ -directed there is  $f' <_{D+A} f$  such that  $\alpha \in C \Rightarrow f_\alpha <_{D+A} f'$ . By the first inequality  $\|f'_{D+A}\| < \|f\|_{D+A} = \delta$ , and by the second inequality  $\alpha \in C \Rightarrow \alpha \leq \|f_\alpha\|_{D+A} \leq \|f'\|_{D+A}$  hence  $\delta = \sup(C) \leq \|f'\|_{D+A}$ , a contradiction.
- 4) Same proof as part (2).
- 5) By part (4) and [Sh:g, Ch.II,3.1].
- 6) Follows.
- 7) Toward contradiction assume that not; by part (2) without loss of generality  $\forall i[\text{cf}(f(i)) > 2^\kappa]$ ; let  $C \subseteq \delta$  be unbounded,  $\text{otp}(C) = \text{cf}(\delta)$ . For each  $\alpha \in C$  and  $A \in J_D^+(f, \kappa)$  choose  $f_{\alpha,A} <_D f$  such that  $\|f_{\alpha,A}\|_{D+A} = \alpha$ . Let  $f_\alpha$  be  $f_\alpha(i) = \sup\{f_{\alpha,A}(i) : A \in J_D^+(f, \kappa)\}$ . As  $(\prod_{i < \kappa}^* f_\alpha(i), <_{J_D(f, \kappa)})$  is  $\text{cf}(\delta)$ -directed (see part (6)), by the assumption toward contradiction and the pcf theorem we have  $\prod_{i < \kappa}^* f(i)/J_D(f, \kappa)$  is  $(\text{cf}(\delta))^+$ -directed. Hence we can find  $f^* < f$  such that  $\alpha \in C \Rightarrow f_\alpha <_{J_D(f, \kappa)} f^*$ . Let  $\beta = \sup\{\|f^*\|_{D+B} : B \in J_D^+(f, A)\}$ , it is  $< \delta$  as  $\text{cf}(\delta) > 2^\kappa$ ; hence there is  $\alpha, \beta < \alpha \in C$ , so by the choice of  $f^*$  we have  $f_\alpha <_{J_D(f, \kappa)} f^*$ , and let  $A =: \{i < \kappa : f_\alpha(i) < f^*(i)\}$  so  $A \in J_D^+(f, \kappa)$ , so  $f_{\alpha,A} \leq f_\alpha <_{D+A} f^*$  hence  $\alpha \leq \|f_{\alpha,A}\|_{D+A} \leq \|f_\alpha\|_{D+A} \leq \|f^*\|_{D+A} \leq \beta$  contradicting the choice of  $\alpha$ .
- 8) For every  $\alpha < \lambda$  we can choose  $f_\alpha <_D f$  such that  $\|f_\alpha\|_D = \alpha$ . Let  $a_\alpha = \{\|f_\alpha\|_{D+A} : A \in D^+\}$ , as  $A \in D^+ \Rightarrow \alpha \leq \|f_\alpha\|_D \leq \|f_\alpha\|_{D+A} < \|f\|_{D+A} = \lambda$ , clearly  $a_\alpha$  is a subset of  $\lambda \setminus \alpha$ , and its cardinality is  $\leq 2^\kappa < \lambda$ . So we can find an unbounded  $E \subseteq \lambda$  such that  $\alpha < \beta \in E \Rightarrow \sup(a_\alpha) < \beta$ . So if  $\alpha < \beta, \alpha \in E, \beta \in E$ , let  $A = \{i < \kappa : f_\alpha(i) \geq f_\beta(i)\}$ , and if  $A \in D^+$ , then  $\|f_\beta\|_{D+A} \leq \|f_\alpha\|_{D+A} \leq \sup(a_\alpha) < \beta$ , contradiction. Hence  $A = \emptyset \text{ mod } D$ , that is  $f_\alpha <_D f_\beta$ . Also if  $g <_D f$ , then  $a =: \{\|g\|_{D+A} : A \in D^+\}$  is again a subset of  $\lambda$  of cardinality  $\leq 2^\kappa$  hence for some  $\beta < \lambda$ ,  $\sup(a) < \beta$ , so as above  $g <_D f_\beta$ . Together  $\langle f_\alpha : \alpha \in E \rangle$  exemplify  $\lambda = \text{tcf}(\prod f(i), <_D)$ .
- 9) Similar proof (to part (8)), using parts (6), (7). □<sub>5.11</sub>

5.12 Remark. We think Claims 5.9, 5.10, 5.11 (and Definition 5.8) can be applied to the problems from [Sh 497] probably saving some uses of niceness so weakening

some assumptions; but we have not checked.

*Proof of 5.6.* Fix  $f \in {}^\kappa\text{Ord}$  as  $f(i) = \lambda_i$  and let  $\approx, \approx_I$  be as in Definition 5.8. For each  $i < \kappa$  let  $\bar{X}^i = \langle X_\alpha^i : \alpha < \mu_i \rangle$  be a sequence of members of  $[\lambda_i]^{\lambda_i}$  such that

$$\alpha < \beta < \mu_i \Rightarrow X_\alpha^i \setminus X_\beta^i \in J_{\lambda_i}^{\text{bd}} \ \& \ X_\beta^i \setminus X_\alpha^i \notin J_{\lambda_i}^{\text{bd}}.$$

(it exists by assumption (d)).

Let  $\bar{g}^* = \langle g_\zeta^* : \zeta < \mu \rangle$  be a  $<_D$ -increasing sequence of members of  $\prod_{i < \kappa} \mu_i$ , it exists

by assumption (e) or (e)<sup>-</sup>.

Let  $I =: \{B \subseteq \kappa : \text{if } B \in D^+ \text{ then } \|f\|_{D+B} > \lambda\}$ , it is a  $\text{com}(D)$ -complete ideal on  $\kappa$  disjoint to  $D$ , i.e.,  $I = J_D(\bar{\lambda}, \kappa) \supseteq \{\kappa \setminus A : A \in D\}$ , and  $\approx_I, \approx$  are equal because  $I$  is the ideal on  $\kappa$  dual to  $D$  which holds by assumption (c). For any sequence  $\bar{X} = \langle X_i : i < \kappa \rangle \in \prod_{i < \kappa} [\lambda_i]^{\lambda_i}$ , let

$$Y[\bar{X}] =: \left\{ \|h\|_{D+A} : h \in \prod_{i < \kappa} X_i \text{ and } A \in I^+ \right\}$$

and

$$\mathscr{Y}[\bar{X}] =: \left\{ (h, A) / \approx : h \in \prod_{i < \kappa} X_i \text{ and } (h, A) \in \text{cla}_I^\alpha(\bar{\lambda}, D) \right. \\ \left. \text{for some } \alpha < \lambda \right\}$$

Note:  $Y[\bar{X}] \subseteq \lambda$  and  $\mathscr{Y}[\bar{X}] \subseteq \mathscr{Y}^* =: \bigcup_{\alpha < \lambda} \text{cla}^\alpha(\bar{\lambda}, D) / \approx$ .

Note that by 5.9(6)

$$\boxtimes \bigcup_{\alpha < \lambda} \text{cla}^\alpha(f, D) / \approx \text{ has cardinality } \leq \lambda.$$

(\*)<sub>0</sub> for  $\bar{X} \in \prod_{i < \kappa} [\lambda_i]^{\lambda_i}$ , the mapping  $(g, A) / \approx_I \mapsto \|g\|_{D+A}$  is from  $\mathscr{Y}[\bar{X}]$  onto  $Y[\bar{X}]$  with every  $\alpha \in Y[\bar{X}]$  having at most  $2^\kappa$  preimages [why? by 5.9(5)]

(\*)<sub>1</sub> if  $\bar{X} \in \prod_{i < \kappa} [\lambda_i]^{\lambda_i}$  then  $\mathscr{Y}[\bar{X}]$  has cardinality  $\lambda$ .

[why? by the definition of  $\| - \|_D$  for every  $\alpha < \lambda$  for some  $g \in \prod_{i < \kappa} \lambda_i/D$  we

have  $\|g\|_D = \alpha$ ; as  $\sup(X_i) = \lambda_i > g(i)$  we can find  $g' \in \prod_{i < \kappa} (X_i \setminus g(i))$  such

that  $g \leq g' < \langle \lambda_i : i < \kappa \rangle$ , so  $\alpha = \|g\|_D \leq \|g'\|_D < \|\langle \lambda_i : i < \kappa \rangle\|_D = \lambda$ .

Clearly for some  $\alpha'$  and  $A$ ,  $(g', A) \in \text{cla}^{\alpha'}(f, A)$ , so  $A \in I^+ \subseteq D^+$ , and

$\alpha \leq \alpha' = \|g'\|_{D+A} < \|f\|_{D+A} = \lambda$  (as  $A \in I^+$ ). So  $\alpha' \in Y[\bar{X}]$  hence

$Y[\bar{X}] \not\subseteq \alpha$ ; as  $\alpha < \lambda$  was arbitrary,  $Y[\bar{X}]$  has cardinality  $\geq \lambda$ , by  $\boxtimes$  equality holds hence (by (\*)<sub>0</sub>) also  $\mathscr{Y}[\bar{X}]$  has cardinality  $\lambda$ .]

(\*)<sub>2</sub> if  $\bar{X}', \bar{X}'' \in \prod_{i < \kappa} [\lambda_i]^{\lambda_i}$ , and  $\{i < \kappa : X'_i \subseteq X''_i \text{ mod } J_{\lambda_i}^{\text{bd}}\} \in D$  then

(a)  $Y[\bar{X}'] \subseteq Y[\bar{X}''] \text{ mod } J_{\lambda}^{\text{bd}}$

(b)  $\mathscr{Y}[\bar{X}'] \setminus \mathscr{Y}[\bar{X}'']$  has cardinality  $< \lambda$ .

[Why? Define  $g \in \prod_{i < \kappa} \lambda_i$  by  $g(i) = \sup(X'_i \setminus X''_i)$  if

$i \in A^* =: \{i < \kappa : X'_i \subseteq X''_i \text{ mod } J_{\lambda_i}^{\text{bd}}\}$  and  $g(i) = 0$  other-

wise. Let  $\alpha(*) = \sup\{\|g\|_{D+A} + 1 : A \in I^+\}$ , as  $\lambda$  is regular

$> 2^\kappa$  clearly  $\alpha(*) < \lambda$  (see assumption (c) or definition of  $I$ ). As-

sume  $\beta \in Y[\bar{X}'] \setminus \alpha(*)$  and we shall prove that  $\beta \in Y[\bar{X}'']$ , moreover,

$\mathscr{Y}[\bar{X}'] \cap (\text{cla}^\beta(\bar{X}, D)/ \approx_I) \subseteq \mathscr{Y}[\bar{X}'']$ , this clearly suffices for both

clauses. We can find  $f^* \in \prod_{i < \kappa} ((X'_i \cap X''_i) \cup \{0\})$  such that  $\|f^*\|_D > \beta$ .

So let a member of  $\mathscr{Y}[\bar{X}'] \cap (\text{cla}^\beta(\bar{\lambda}, D)/ \approx)$  have the form  $(h, A)/ \approx_I$ ,

where  $A \in I^+$ ,  $h \in \prod_{i < \kappa} X'_i$  and  $\beta = \|h\|_{D+A}$  and let

$A_1 =: \{i < \kappa : h(i) \leq g(i)\}$ . We know

$\beta = \|h\|_{D+A} = \text{Min}\{\|h\|_{D+(A \cap A_1)}, \|h\|_{D+(A \setminus A_1)}\}$  (if  $A \cap A_1 = \emptyset \text{ mod } D$ , then  $\|h\|_{D+A \cap A_1}$  can be considered  $\infty$ ).

If  $\beta = \|h\|_{D+(A \cap A_1)}$  then note  $h \leq_{D+(A \cap A_1)} g$  hence

$\beta = \|h\|_{D+(A \cap A_1)} \leq \|g\|_{D+(A \cap A_1)} < \alpha(*)$ , contradicting an assumption on  $\beta$ . So  $\beta = \|h\|_{D+(A \setminus A_1)}$  and  $A \cap A_1 \in J_{D,I}(h, A)$ . Now

define  $h' \in \prod_{i < \kappa} f(i)$  by:  $h'(i)$  is  $h(i)$  if  $i \in A \setminus A_1$  and  $h'(i)$  is  $f^*(i)$  if  $i \in$

$\kappa \setminus (A \setminus A_1)$ . So  $h' \in \prod_{i < \kappa} f(i)$  and  $h' =_{D+(A \setminus A_1)} h$  hence  $\|h'\|_{D+(A \setminus A_1)} =$

$\|h\|_{D+(A \setminus A_1)} = \beta$ , and clearly  $\beta = \|h'\|_{D+(A \setminus A_1)} \in Y[\bar{X}'']$ , as required

for clause (a), moreover  $(h, A) \approx (h', A)$  so  $((h', A)/ \approx) \in \mathscr{Y}[\bar{X}'']$  as required for clause (b).]

modified:2005-07-06

revision:2005-07-05

(589)

(\*)<sub>3</sub> If  $\bar{X}', \bar{X}'' \in \prod_{i < \lambda} [\lambda_i]^{\lambda_i}$  and  $\{i < \kappa : X_i'' \not\subseteq X_i' \text{ mod } J_\lambda^{\text{bd}}\} \in D$  then

$\mathcal{Y}[\bar{X}''] \setminus \mathcal{Y}[\bar{X}']$  has cardinality  $\lambda$ .

[Why? Let  $\alpha < \lambda$ , it is enough to find  $\beta \in [\alpha, \lambda)$  such that

$$(\mathcal{Y}[\bar{X}''] \setminus \mathcal{Y}[\bar{X}']) \cap (\text{cla}^\beta(f, D) / \approx) \neq \emptyset.$$

We can find  $g \in \prod_{i < \kappa} \lambda_i$  such that  $\|g\|_D = \alpha$ . Define  $g' \in \prod_{i < \kappa} X_i''$  by:  $g'(i)$  is  $\text{Min}(X_i'' \setminus X_i' \setminus g(i))$  when well defined,  $\text{Min}(X_i'')$  otherwise. By assumption  $g \leq_D g'$  and, of course,  $g' \in \prod_{i < \kappa} X_i'' \subseteq \prod_{i < \kappa} \lambda_i$ , so  $\|g'\|_D \geq \alpha$ . So

$((g', \kappa) / \approx) \in \mathcal{Y}[\bar{X}'']$  but trivially  $((g', \kappa) / \approx) \notin \mathcal{Y}[\bar{X}']$ , so we are done.]

Together  $(*)_0 - (*)_3$  give that  $\langle \mathcal{Y}[\langle X_{g_\zeta^*}^i : i < \kappa \rangle] : \zeta < \mu \rangle$  is a sequence of subsets of  $\mathcal{Y}^*$  of length  $\mu$  (see  $(*)_1$ ),  $|\mathcal{Y}^*| = \lambda$ , which is increasing modulo  $[\mathcal{Y}^*]^{< \lambda}$  (by  $(*)_2$ ), and in fact, strictly increasing (by  $(*)_3$ , see choice of  $\langle g_\zeta^* : \zeta < \mu \rangle$  in the beginning of the proof). So modulo changing names we have finished. (In fact, also  $\langle Y[\langle X_{g_\zeta^*}^i : i < \kappa \rangle] : \zeta < \mu \rangle$  is as required.)  $\square_{5.6}$

A related theorem

### 5.13 Definition.

$$\alpha_\lambda = \text{Min} \left\{ \mu : \text{there is no } \mathcal{P} \subseteq [\lambda]^\lambda \text{ of cardinality } \mu \text{ such that } A \neq B \in \mathcal{P} \Rightarrow |A \cap B| < \lambda \right\}.$$

### 5.14 Theorem. Assume

- (a)  $D$  is an  $\aleph_1$ -complete filter on  $\kappa$
- (b)  $\langle \lambda_i : i < \kappa \rangle$  is a sequence of regular cardinals  $> (2^\kappa)^+$
- (c)  $\|\langle \lambda_i : i < \kappa \rangle\|_{D+A} = \lambda$  for  $A \in D^+$
- (d)  $\mu_i < \mathfrak{a}_{\lambda_i}$
- (e)  $\mu = \text{tcf}(\prod \mu_i / D)$  or at least
- (e<sup>-</sup>)  $\mu < \text{Depth}^+(\prod \mu_i, <_D)$  and  $\mu > 2^\kappa$ .



Then  $\mu < \mathfrak{a}_\lambda$ .

*Proof of 5.14.* Similar to the proof of 5.6

**5.15 Theorem.** *Assume*

- (a)  $D$  an  $\aleph_1$ -complete filter on  $\kappa$
- (b)  $\bar{\lambda} = \langle \lambda_i : i < \kappa \rangle$  is a sequence of regular cardinals  $> 2^\kappa$
- (c)  $\lambda = \|\bar{\lambda}\|_{D+A}$  for  $A \in D^+$
- (d)  $\mu_i < \mathfrak{dp}_{\lambda_i}^{1+}$
- (e)  $\mu < \text{Depth}^+(\prod_{i < \kappa} \mu_i, <_D)$ .

Then  $\mu < \mathfrak{dp}_\lambda^{1+}$ .

*Proof.* Let  $\text{Club}(\lambda) = \{C : C \text{ a club of } \lambda\}$  so  $\text{Club}(\lambda) \subseteq [\lambda]^\lambda$  for  $\lambda = \text{cf}(\lambda) > \aleph_0$ .

For any sequence  $\bar{C} \in \prod_{i < \kappa} \text{Club}(\lambda_i)$  let  $\mathcal{C}(\bar{C})$  be the set  $\text{acc}(\text{cl}(Y(\bar{C})))$  where  $Y[\bar{C}] =: \{\|g\|_D : g \in \prod_{i < \kappa} C_i\} (\subseteq \lambda)$ ; i.e.  $\mathcal{C}(\bar{C}) = \{\delta < \lambda : \delta = \sup(\delta \cap Y[\bar{C}])\}$ .

Clearly

- (\*)<sub>1</sub> for  $\bar{C} \in \prod_{i < \kappa} \text{Club}(\lambda_i)$  we have  $\mathcal{C}(\bar{C}) \in \text{Club}(\lambda)$

[the question is why it is unbounded, and this holds as  $\|\bar{\lambda}\|_D = \lambda$  by its definition]

- (\*)<sub>2</sub> if  $\bar{C}', \bar{C}'' \in \prod_{i < \kappa} \text{Club}(\lambda_i)$ ,  $g^* \in \prod_{i < \kappa} \lambda_i$ , and  $C'_i = C''_i \setminus g^*(i)$  then

$$\mathcal{C}(\bar{C}') = \mathcal{C}(\bar{C}'') \text{ mod } J_\lambda^{\text{bd}}.$$

[Why? Let  $\alpha(*) = \sup\{\|g^*\|_{D+A} : A \in D^+ \text{ and } \|g^*\|_{D+A} < \lambda\} + 1$ , so as  $2^\kappa < \lambda = \text{cf}(\lambda)$  clearly  $\alpha(*) < \lambda$ . We shall show  $\mathcal{C}(\bar{C}') \setminus \alpha(*) = \mathcal{C}(\bar{C}'') \setminus \alpha(*)$ ; for this it suffices to prove  $Y(\bar{C}') \setminus \alpha(*) = Y(\bar{C}'') \setminus \alpha(*)$ . If  $\alpha \in Y(\bar{C}') \setminus \alpha(*)$  let  $\alpha = \|h\|_D$  where  $h \in \prod_i C'_i$ , and let  $A = \{i <$

$\kappa : h(i) < g^*(i)\}$ , so if  $A \in (J_D(\bar{\lambda}, \kappa))^+$  then  $\alpha \leq \|h\|_{D+A} < \lambda$  and  $\|h\|_{D+A} \leq \|g^*\|_{D+A} < \alpha(*)$  but  $\alpha \geq \alpha(*)$ , a contradiction. So  $A \in J_D(\bar{\lambda}, \kappa)$  hence  $A \notin D^+$  by clause (c) of the assumption, so  $g^* \leq_D h$ . Now clearly there is  $h' =_D h$  with  $h' \in \prod_{i < \kappa} C''_i$ , so  $\alpha = \|h\|_D = \|h'\|_D \in \mathcal{C}(\bar{C}'')$ . The

other inclusion is easier.]

(\*)<sub>3</sub> if  $\bar{C}', \bar{C}'' \in \prod_{i < \kappa} \text{Club}(\lambda_i)$  and  $\{i < \kappa : C_i'' \subseteq \text{acc}(C_i')\} \in D$  then

$$\mathcal{C}(\bar{C}'') \subseteq \text{acc}(\mathcal{C}(\bar{C}')).$$

[Why? Let  $\beta \in \mathcal{C}[\bar{C}'']$  but  $\beta \notin \text{acc}(\mathcal{C}(\bar{C}'))$  and we shall get a contradiction. Clearly  $\beta > \sup(\mathcal{C}(\bar{C}') \cap \beta)$  (as  $\beta \notin \text{acc}(\mathcal{C}(\bar{C}'))$ ). As  $\mathcal{C}[\bar{C}'']$  is  $\text{acc}(clY[\bar{C}''])$ , clearly there is  $\alpha \in Y[\bar{C}'']$  such that  $\beta > \alpha > \sup(\mathcal{C}(\bar{C}') \cap \beta)$ , but  $Y[\bar{C}''] = \{\|g\|_D : g \in \prod_{i < \kappa} C_i''\}$ , so there is  $g \in \prod_{i < \kappa} C_i''$  such that  $\|g\|_D = \alpha$ . As  $\{i : C_i'' \subseteq \text{acc}(C_i')\} \in D$ , clearly

$$B =: \{i < \kappa : g(i) \in \text{acc}(C_i')\} \in D.$$

So if  $h \in \prod_{i < \lambda} \lambda_i, h <_D g$  then we can find  $h' \in \prod_{i < \kappa} C_i'$  such that  $h <_D h' <_D g$

(just  $h'(i) = \text{Min}(C_i' \setminus (h(i) + 1))$  noting  $B \in D$ ) hence

$\alpha = \|g\|_D = \sup\{\|h\|_D : h(i) \in g(i) \cap C_i' \text{ when } i \in B, h(i) = \text{Min}(C_i') \text{ otherwise}\}$   
and in this set there is no last element and it is included in  $Y[\bar{C}']$ , so necessarily  $\alpha \in \mathcal{C}(\bar{C}')$ , contradicting the choice of  $\alpha : \beta > \alpha > \sup(\mathcal{C}(\bar{C}') \cap \beta)$ .]

(\*)<sub>4</sub> if  $\bar{C}', \bar{C}'' \in \prod_{i < \kappa} \text{Club}(\lambda_i)$  and  $\{i : C_i'' \subseteq \text{acc}(C_i') \text{ mod } J_{\lambda_i}^{\text{bd}}\} \in D$  then

$$\mathcal{C}(\bar{C}'') \subseteq \text{acc}(\mathcal{C}(\bar{C}')) \text{ mod } J_{\lambda}^{\text{bd}}.$$

[Why? By (\*)<sub>2</sub> + (\*)<sub>3</sub>, i.e., define  $C_i'''$  to be  $C_i'' \setminus g(i)$  where  $g(i) =: \sup(C_i'' \setminus \text{acc}(C_i')) + 1$  when  $C_i'' \subseteq \text{acc}(C_i')$  and the empty set otherwise. Now by (\*)<sub>2</sub> we know  $\mathcal{C}(\bar{C}'') = \mathcal{C}(\bar{C}''') \text{ mod } J_{\lambda}^{\text{bd}}$  and by (\*)<sub>3</sub> we know  $\mathcal{C}(\bar{C}''') \subseteq \text{acc}(\mathcal{C}(\bar{C}'))$ .]

Now we can prove the conclusion of 5.15. Let  $\langle C_{\alpha}^i : \alpha < \mu_i \rangle$  witness  $\mu_i < \mathfrak{dp}_{\lambda_i}^{1+}$  and  $\langle g_{\alpha} : \alpha < \mu \rangle$  witness  $\mu < \text{Depth}^+(\prod_{i < \kappa} \lambda_i, <_D)$ . Let  $C_{\alpha} =: \mathcal{C}(\langle C_{g_{\alpha}(i)}^i : i < \kappa \rangle)$  for  $\alpha < \mu$ . So  $\langle C_{\alpha} : \alpha < \mu \rangle$  witnesses  $\mu < \mathfrak{dp}_{\lambda}^{1+}$ . □<sub>5.15</sub>

**5.16 Theorem.** *Assume*

- (a)  $\kappa$  is regular uncountable
- (b)  $\bar{\lambda} = \langle \lambda_i : i < \kappa \rangle$  is a sequence of regular cardinals  $> \kappa$
- (c)  $D$  is a normal filter on  $\kappa$  (or just  $\aleph_1$ -complete)
- (d)  $\lambda = \|\bar{\lambda}\|_D = \text{tcf}(\prod_{i < \kappa} \lambda_i / D), \lambda$  regular
- (e)  $\mu_i < \mathfrak{dp}_{\lambda_i}^{2+}$

$$(f) \mu < \text{Depth}^+(\prod_{i < \kappa} \mu_i, <_D).$$

Then  $\mu < \mathfrak{dp}_\lambda^{2+}$ .

*Proof.* Let  $\langle f_\alpha^i : \alpha < \mu_i \rangle$  exemplify  $\mu_i < \mathfrak{dp}_{\lambda_i}^{+2}$ , let  $\langle g_\alpha : \alpha < \mu \rangle$  exemplify  $\mu < \text{Depth}^+(\prod_{i < \kappa} \mu_i, <_D)$ , and let  $\langle h_\zeta : \zeta < \lambda \rangle$  exemplify  $\lambda = \text{tcf}(\prod_{i < \kappa} \lambda_i, <_D)$ .

Now for each  $\alpha < \mu$  we define  $f_\alpha \in {}^\lambda \lambda$  as follows:

$$f_\alpha(\zeta) = \|\langle f_{g_\alpha(i)}^i(h_\zeta(i)) : i < \kappa \rangle\|_D.$$

Clearly  $f_\alpha(\zeta)$  is an ordinal and as  $f_{g_\alpha(i)}^i \in {}^{(\lambda_i)} \lambda_i$  clearly  $\langle f_{g_\alpha(i)}^i(h_\zeta(i)) : i < \kappa \rangle <_D \langle \lambda_i : i < \kappa \rangle$  hence  $f_\alpha(\zeta) < \|\bar{\lambda}\|_D = \lambda$ , so

$$(*)_1 f_\alpha \in {}^\lambda \lambda.$$

The main point is to prove  $\beta < \alpha < \mu \Rightarrow f_\beta <_{J_\lambda^{\text{bd}}} f_\alpha$ .

Suppose  $\beta < \alpha < \mu$ , then  $g_\beta <_D g_\alpha$  hence  $A =: \{i < \kappa : g_\beta(i) < g_\alpha(i)\} \in D$  so  $i \in A \Rightarrow f_{g_\beta(i)}^i <_{J_{\lambda_i}^{\text{bd}}} f_{g_\alpha(i)}^i$ . We can define  $h \in \prod_{i < \kappa} \lambda_i$  by:

$h(i)$  is  $\sup\{\zeta + 1 : f_{g_\beta(i)}^i(\zeta) \geq f_{g_\alpha(i)}^i(\zeta)\}$  if  $i \in A$ , and  $h(i)$  is zero otherwise.

But  $\langle h_\zeta : \zeta < \lambda \rangle$  is  $<_D$ -increasing and cofinal in  $(\prod_{i < \kappa} \lambda_i, <_D)$  hence there is

$\zeta(*) < \lambda$  such that  $h <_D h_{\zeta(*)}$ .

So it suffices to prove:

$$\zeta(*) \leq \zeta < \lambda \Rightarrow f_\beta(\zeta) < f_\alpha(\zeta).$$

So let  $\zeta \in [\zeta(*), \lambda)$ , so

$$B =: \{i < \kappa : h(i) < h_{\zeta(*)}(i) \leq h_\zeta(i) \text{ and } i \in A\}$$

belongs to  $D$  and by the definition of  $A$  and  $B$  and  $h$  we have

$$i \in B \Rightarrow f_{g_\beta(i)}^i(h_\zeta(i)) < f_{g_\alpha(i)}^i(h_\zeta(i)).$$

So

$$\langle f_{g_\beta(i)}^i(h_\zeta(i)) : i < \kappa \rangle <_D \langle f_{g_\alpha(i)}^i(h_\zeta(i)) : i < \kappa \rangle$$

hence (by the definition of  $\| - \|_D$ )

$$\|\langle f_{g_\beta}^i(h_\zeta(i)) : i < \kappa \rangle\|_D < \|\langle f_{g_\alpha}^i(h_\zeta(i)) : i < \kappa \rangle\|_D$$

which means

$$f_\beta(\zeta) < f_\alpha(\zeta).$$

As this holds for every  $\zeta \in [\zeta(*), \lambda)$  clearly

$$f_\beta <_{J_\lambda^{\text{bd}}} f_\alpha.$$

So  $\langle f_\alpha : \alpha < \mu \rangle$  is  $<_{J_\lambda^{\text{bd}}}$ -increasing, so we have finished.  $\square_{5.16}$

**5.17 Discussion:** Now assumption (c) in 5.15 (and in 5.6) is not so serious once we quote [Sh:g, V] (to satisfy the assumption in the usual case we are given  $\lambda = \text{cf}(\lambda)$ ,  $\mu < \lambda \leq \mu^\kappa$ ,  $\text{cf}(\mu) = \kappa$ ,  $(\forall \alpha < \mu)(|\alpha|^\kappa < \mu)$  and we like to find  $\langle \lambda_i : i < \kappa \rangle$ , and normal  $D$  such that  $\|\langle \lambda_i : i < \kappa \rangle\|_{D+A} = \lambda$ ). E.g., ([Sh:g, Ch.V]) if SCH fails above  $2^{2^\theta}$ ,  $\theta$  regular uncountable,  $D$  a normal filter on  $\theta$ ,  $\|f\|_D \geq \lambda = \text{cf}(\lambda) > 2^{2^\theta}$ , (so if  $\mathcal{E}$  = family of normal filters on  $\theta$ , so  $\mathcal{E}$  is nice and  $\text{rk}_E^3(f) \geq \|f\|_D \geq \lambda$ ), so  $g_\kappa$  from [Sh:g, Ch.V,3.10,p.244] is as required.

Still we may note

**5.18 Fact.** Assume

- (a)  $D$  is an  $\aleph_1$ -complete filter on  $\kappa$
- (b)  $f^* \in {}^\kappa\text{Ord}$  and  $\text{cf}(f^*(i)) > 2^\kappa$  for  $i < \kappa$ .

Then for any  $\bar{C} = \langle C_i : i < \kappa \rangle$ ,  $C_i$  a club of  $f^*(i)$  and  $\alpha < \|f^*\|_D$  we can find  $f \in \prod_{i < \kappa} C_i$  such that:

- ( $\alpha$ )  $A \in (J_D(f^*, \kappa))^+ \Rightarrow \alpha < \|f\|_{D+A} = \|f\|_D < \|f^*\|_D$
- ( $\beta$ )  $A \in J_D(f^*, \kappa) \cap D^+ \Rightarrow \|f\|_{D+A} \geq \|f^*\|_D$

*Proof.* We choose by induction on  $\zeta \leq \kappa^+$  a function  $f_\zeta$  and  $\langle f_{\zeta,A} : A \in (J_D(f^*, \kappa))^+ \rangle$  such that:

- (a)  $f_\zeta \in \prod_{i < \kappa} C_i$
- (b)  $\varepsilon < \zeta \Rightarrow \bigwedge_i f_\varepsilon(i) < f_\zeta(i)$

- (c) for  $\zeta$  limit  $f_\zeta(i) = \sup_{\varepsilon < \zeta} f_\varepsilon(i)$
- (d) for  $A \in (J_D(f^*, A))^+$ , letting  $\alpha_{\zeta, A} =: \|f_\zeta\|_{D+A}$  we have  
 $f_{\zeta, A} \in \prod_{i < \kappa} f^*(i), \|f_{\zeta, A}\|_D > \alpha_{\zeta, A}$  and  
 $f_{\zeta, A}(i) \geq f_\zeta(i)$  for  $i < \kappa$
- (e)  $f_{\zeta, A}(i) < f_{\zeta+1}(i)$  for  $i < \kappa, A \in (J_D(f^*, A))^+$
- (f)  $\|f_0\|_D \geq \alpha$  and  
 $A \in J_D(f^*, \kappa) \Rightarrow \|f_0\|_{D+A} \geq \|f^*\|_D$ .

There is no problem to carry out the definition: for defining  $f_0$  for each  $A \in J_D(f^*, \kappa)$  choose  $g_A <_{D+A} f^*$  such that  $\|g_A\|_{D+A} \geq \|f^*\|_D$  (possible as  $\|f^*\|_{D+A} > \|f^*\|_D$  by the assumption on  $A$ ). Let  $g^* < f^*$  be such that  $\|g^*\|_D \geq \alpha$ , (possible as  $\alpha < \|f^*\|_D$ ) and let  $f_0 \in \prod_{i < \kappa} f^*(i)$  be defined by  $f_0(i) = \text{Min}(C_i \setminus \sup\{g^*(i), g_A(i) : A \in J_D(f^*, \kappa)\})$ . For  $\zeta$  limit there is no problem to define  $f_\zeta$ ; and also for  $\zeta$  successor. If  $f_\zeta$  is defined, we should choose  $f_{\zeta, A}$ . For clause (d) note that  $\|f^*\|_{D+A} = \|f^*\|_D$  as  $A \in (J_D(f^*, A))^+$  and use the definition of  $\|f\|_D$ . We use, of course,  $\bigwedge_i \text{cf}(f^*(i)) > 2^\kappa$ .

Now  $f_{\kappa^+}$  is as required. Note:  $f <_D f_{\kappa^+} \Rightarrow \bigvee_{\zeta < \kappa^+} f <_D f_\zeta$ , and for

$$A \in (J_D(f^*, \kappa))^+, \|f_{\kappa^+}\|_{D+A} = \sup_{\zeta < \kappa^+} \|f_\zeta\|_{D+A} = \sup_{\zeta < \kappa^+} \alpha_{\zeta, A} \leq \sup_{\zeta < \kappa^+} \|f_{\zeta+1}\|_D = \|f_{\kappa^+}\|_D.$$

□ **5.18**

→ scite{5.11A} ambiguous

*5.19 Conclusion.* 1) In 5.15 we can weaken assumption (c) to

(1)  $(c)^-$   $\|\langle \lambda_i : i < \kappa \rangle\|_D = \lambda$ .

2) In 5.6 we can weaken assumption (c) to  $(c)^-$ .

*Proof.* 1) In the proof of 5.15, choose  $g^{**} \in \prod_{i < \kappa} \lambda_i$  satisfying (exists by **5.18**):

→ scite{5.11A} ambiguous

$(*)_0$   $A \in J_D(\bar{\lambda}, \kappa) \cap D^+ \Rightarrow \|g^{**}\|_{D+A} \geq \lambda$  (which is  $\|\bar{\lambda}\|_D$ ).

We redefine  $Y[\bar{C}]$  as  $\{\|g\|_D : g \in \prod_{i < \kappa} C_i \text{ but } g(i) \geq g^{**}(i) \text{ for } i < \kappa\}$ . The only change is during the proof of  $(*)_2$  there. Now if  $\alpha \in Y[\bar{C}'] \setminus \alpha(*)$  then there is  $h \in \prod_{i < \kappa} \lambda_i$  such that  $[i < \kappa \Rightarrow h(i) \geq g^{**}(i)]$  and  $\|h\|_D = \alpha$  and let  $A = \{i < \kappa : h(i) < g^*(i)\}$ . Now if  $A \in (J_D(\bar{\lambda}, \kappa))^+$  we get a contradiction as there and if  $A = \emptyset \text{ mod } D$  we finish as there. So we are left with the case  $A \in J_D(\bar{\lambda}, \kappa) \cap D^+$ ,  $\|\bar{\lambda}\|_{D+A} > \|\bar{\lambda}\|_D \leq \lambda$  hence  $\|g^{**}\|_{D+A} \leq \lambda$  hence  $\|h\|_{D+A} \leq \lambda > \alpha$  hence necessarily  $\|h\|_{D+(\kappa \setminus A)} = \alpha$  (as  $\|h\|_D = \text{Min}\{\|h\|_{D+A}, \|h\|_{D+(\kappa \setminus A)}\}$ ). Now choose  $h' \in \prod_{i < \kappa} \lambda_i$  by  $h' \upharpoonright (\kappa \setminus A) = h \upharpoonright (\kappa \setminus A)$  and  $[i \in A \Rightarrow h'(i) = \text{Min}(C''_i \setminus h(i))]$  so  $h' \in \prod_{i < \kappa} C''_i$ ,  $h \leq h' < \bar{\lambda}$ ,  $\lambda \leq \|h\|_{D+A} \leq \|h'\|_{D+A} \leq \|h'\|_{D+A}$  and so  $\|h'\|_D = \text{Min}\{\|h'\|_{D+A}, \|h'\|_{D+(\kappa \setminus A)}\} = \alpha$ . So we are done.

2) Let  $g^{**}$  be as in the proof of part (1). In the proof of 5.6 we let

$$Y[\bar{X}] =: \left\{ \|h\|_{D+A} : h \in \prod_{i < \kappa} (X_i \setminus g^{**}(i)) \text{ and } A \in I^+ \right\},$$

remembering  $I = J_D(\bar{\lambda}, \kappa)$ .

$$\mathcal{Y}[\bar{X}] =: \left\{ (h, A) / \approx_I : h \in \prod_{i < \kappa} (X_i \setminus g^{**}(i)) \text{ and } (h, A) \in \text{cl}_I^\alpha(\lambda, D) \text{ for some } \alpha < \lambda \right\}$$

and we can restrict ourselves to sequences  $\bar{X}$  such that  $X_i \cap g^{**}(i) = \emptyset$ . In the proof of  $(*)_2$  make  $g \geq g^{**}$ . □<sub>5.19</sub>

**5.20 Claim.** *Assume*

- (a)  $J$  is a filter on  $\kappa$
- (b)  $\lambda$  a regular cardinal,  $\lambda_i > 2^\kappa, \theta > 2^\kappa$
- (c)  $\prod_{i < \kappa} \lambda_i / J$  is  $\lambda$ -like, i.e.,
  - (i)  $\lambda = \text{tcf} \prod \lambda_i / J$
  - (ii)  $T_J(\langle \lambda_i : i < \kappa \rangle) = \lambda$  (follows from (i) + (iii) actually) and
  - (iii) if  $\mu_i < \lambda_i$  then  $T_J(\langle \mu_i : i < \kappa \rangle) < \lambda$

- (d)  $\kappa < \theta = \text{cf}(\theta) < \lambda_i$  for  $i < \kappa$
- (e)  $i < \kappa \Rightarrow S_\theta^{\lambda_i} = \{\delta < \lambda_i : \text{cf}(\delta) = \theta\} \in I[\lambda_i]$  (see below)
- (f)  $(\forall \alpha < \theta)[|\alpha|^\kappa < \theta]$ .

Then  $S_\theta^\lambda = \{\delta < \lambda : \text{cf}(\delta) = \theta\} \in I[\lambda]$ .

*Remark.* Remember that for  $\lambda$  regular uncountable

$$I[\lambda] = \left\{ A \subseteq \lambda : \text{for some club } E \text{ of } \lambda \text{ and } \bar{\mathcal{P}} = \langle \mathcal{P}_\alpha : \alpha < \lambda \rangle \text{ with} \right.$$

$$\mathcal{P}_\alpha \subseteq \mathcal{P}(\alpha), |\mathcal{P}| < \lambda,$$

$$\text{for every } \delta \in A \cap E, \text{cf}(\delta) < \delta \text{ and for some closed}$$

$$\text{unbounded subset } a \text{ of } \delta \text{ of order type } < \delta,$$

$$\left. (\forall \alpha < \delta)(\exists \beta < \delta)(a \cap \alpha \in \mathcal{P}_\beta) \right\}.$$

*Proof.* Clearly each  $\lambda_i$  is a regular cardinal and  $\lambda = \text{tcf}(\prod_{i < \kappa} \lambda_i / J)$ , so let

$\bar{f} = \langle f_\alpha : \alpha < \lambda \rangle$  be a  $<_J$ -increasing sequence of members of  $\prod_{i < \kappa} \lambda_i$ , which is cofinal

in  $\prod_{i < \kappa} \lambda_i / J$ . So without loss of generality if  $\bar{f} \upharpoonright \delta$  has a  $<_J$ -eub  $f'$  then  $f_\delta =_J f'$ .

For each  $i < \kappa$  (see the references above) we can find  $\bar{e}^i = \langle e_\alpha^i : \alpha < \lambda_i \rangle$  and  $E_i$  such that:

- (i)  $E_i$  is a club of  $\lambda_i$
- (ii)  $e_\alpha^i \subseteq \alpha$  and  $\text{otp}(e_\alpha^i) \leq \theta$
- (iii) if  $\beta \in e_\alpha^i$  then  $e_\beta^i = e_\alpha^i \cap \beta$
- (iv) if  $\delta \in E_i$  and  $\text{cf}(\delta) = \theta$ , then  $\delta = \sup(e_\delta^i)$ .

Choose  $\bar{N} = \langle N_i : i < \lambda \rangle$  such that  $N_i \prec (\mathcal{H}(\chi), \in, <_\chi^*)$  where, e.g.,  $\chi = \beth_8(\lambda)^+$ ,  $\|N_i\| < \lambda$ ,  $N_i$  is increasing continuous,  $\bar{N} \upharpoonright (i+1) \in N_{i+1}$ ,  $N_i \cap \lambda$  is an ordinal, and  $\{\bar{f}, J, \lambda, \langle \lambda_i : i < \kappa \rangle, \langle \bar{e}^i : i < \kappa \rangle\} \in N_0$ . Let  $E = \{\delta < \lambda : N_\delta \cap \lambda = \delta\}$ , so it suffices to prove

(\*) if  $\delta \in E \cap S_\theta^\lambda$  then there is  $a$  such that:

- (i)  $a \subseteq \delta$
- (ii)  $\delta = \sup(a)$
- (iii)  $|a| < \lambda$
- (iv)  $\alpha < \delta \Rightarrow a \cap N_\alpha \in N_\delta$ .

By clause (b) in the assumption necessarily  $\bar{f} \upharpoonright \delta$  has a  $<_J$ -eub ([Sh:g, Ch.II,§1]) so necessarily  $f_\delta$  is a  $<_J$ -eub of  $\bar{f} \upharpoonright \delta$ . Moreover,  $A^* = \{i < \kappa : \text{cf}(f_\delta(i)) = \theta \text{ and } f_\delta(i) \in E_i\} = \kappa \text{ mod } J$  by clause (f) of the assumption. So for each  $i \in A^*$ ,  $e_{f_\delta(i)}^i$  is well-defined, and let  $e_{f_\delta(i)}^i = \{\alpha_\zeta^i : \zeta < \theta\}$  with  $\alpha_\zeta^i$  increasing with  $\zeta$ . For each  $\zeta < \theta$  we have  $\langle \alpha_\zeta^i : i < \kappa \rangle <_J f_\delta$  hence for some  $\gamma(\zeta) < \delta$  we have  $\langle \alpha_\zeta^i : i < \kappa \rangle <_J f_{\gamma(\zeta)}$ , but  $T_D(f_{\gamma(\zeta)}) < \lambda$  and  $\gamma(\zeta) \in N_{\gamma(\zeta)+1}$  hence  $f_{\gamma(\zeta)} \in N_{\gamma(\zeta)+1}$  hence for some  $g_\zeta <_J f_{\gamma(\zeta)}$  we have:  $g_\zeta \in N_{\gamma(\zeta)+1}$  and  $A_\zeta = \{i < \kappa : g_\zeta(i) = \alpha_\zeta^i\} \neq \emptyset \text{ mod } J$ . As  $\theta = \text{cf}(\theta) > 2^\kappa$  for some  $A \subseteq \kappa$  we have  $B =: \{\zeta < \theta : A_\zeta = A\}$  is unbounded in  $\theta$ .

Now for  $\zeta < \theta$  let

$$a_\zeta = \left\{ \text{Min}\{\gamma < \lambda : \neg(f_\gamma \leq_{J+(\kappa \setminus A)} g)\} : \right. \\ \left. g \in \prod_{i < \kappa} \{\alpha_\varepsilon^i : \varepsilon < \zeta\} = \prod_{i < \kappa} e_{(\alpha_\zeta^i)}^i \right\}.$$

Clearly  $\zeta < \xi < \theta \Rightarrow a_\zeta \subseteq a_\xi$ . Also for  $\zeta < \theta$ ,  $a_\zeta$  is definable from  $\bar{f}$  and  $g_\zeta \upharpoonright A$ , hence belongs to  $N_{\gamma(\zeta)+1}$ , but its cardinality is  $\leq \theta + 2^\kappa < \lambda$  hence it is a subset of  $N_{\gamma(\zeta)+1}$ . Moreover, also  $\langle a_\xi : \xi < \zeta \rangle$  is definable from  $\bar{f}$  and  $\langle \langle \{\alpha_\varepsilon^i : \varepsilon < \xi\} : i < A \rangle : \xi \leq \zeta \rangle$  hence from  $\bar{f}$  and  $g_\zeta \upharpoonright A$  and  $\langle \bar{e}^i : i < \kappa \rangle$ , all of which belong to  $N_0 \prec N_{\gamma(\zeta)+1}$ , hence  $\zeta \in B \Rightarrow \langle a_\xi : \xi \leq \zeta \rangle \in N_{\gamma(\zeta)+1}$  &  $a_\zeta$  is a bounded subset of  $\delta$ . Now

(\*)  $\bigcup_{\xi < \theta} a_\xi$  is unbounded in  $\delta$ .

[Why? Let  $\beta < \delta$ , so for some  $\zeta < \theta$  we have:

$$f_\beta(i) < f_\delta(i) \Rightarrow f_\beta(i) < \alpha_\zeta^i < f_\delta(i)$$

so

$$\text{Min}\{\gamma : \neg(f_\gamma \leq_{J+(\kappa \setminus A)} \langle \alpha_\zeta^i : i < \kappa \rangle)\} \in (\beta, \delta) \cap a_{\zeta+1}.$$



Let  $w = \{\zeta < \theta : a_\zeta \text{ is bounded in } a_{\zeta+1}\}$

$$a'_\zeta = \{\text{Min}\{\gamma \in a_{\xi+1} : \gamma \text{ is an upper bound of } a_\xi\} : \xi < \zeta\}.$$

So  $\cup\{a'_\zeta : \zeta < \theta\}$  is as required. □<sub>5.20</sub>

*5.21 Remark.* 1) If we want to weaken clause (c) in claim 5.20 retaining only (i) there (and omitting (ii) + (iii)), it is enough if we add:

(g) for each  $i < \kappa$  and  $\delta \in S_\theta^{\lambda_i}$ ,  $\{\gamma < \delta : \text{cf}(\gamma) > \kappa \text{ and } \gamma \in e_\delta^i\}$  is a stationary subset of  $\delta$ .

2) In part (1) of this remark, we can replace  $\text{cf}(\gamma) > \kappa$  by  $\text{cf}(\gamma) = \sigma$ , if  $D$  is  $\sigma^+$ -complete or at least not  $\sigma$ -incomplete.

3) This is particularly interesting if  $\lambda = \mu^+ = \text{pp}(\mu)$ .

modified:2005-07-06

**(589) revision:2005-07-05**

§6 THE CLASS OF CARDINAL ULTRAPRODUCTS MODULO  $D$ 

We presently concentrate on ultrafilters (for filters: two versions). This continues [Sh 506, §3], see history there and in [CK], [Sh:g].

Recall

**6.1 Definition.** 1) A filter  $D$  is  $\theta$ -regular if there are  $A_\varepsilon \in D$  for  $\varepsilon < \theta$  such that the intersection of any infinitely many  $A_\varepsilon$ 's is empty.

2) For a filter  $D$ , let  $\text{reg}(D) = \min\{\theta : D \text{ is not } \theta\text{-regular}\}$ .

Note that  $\text{reg}(D)$  is a regular cardinal.

*6.2 Fact.* Assume

(a)  $D$  is an ultrafilter on  $\kappa$  and  $\theta = \text{reg}(D)$

(b)  $\mu = \text{cf}(\mu)$  and  $\alpha < \mu \Rightarrow |\alpha|^{< \text{reg}(D)} < \mu$

(c)  $\bar{n} = \langle n_i : i < \kappa \rangle, 0 < n_i < \omega, A^* = \bigcup_{i < \kappa} (\{i\} \times n_i)$

(d) for each  $i < \kappa, n < n_i$  we have  $\lambda_{(i,n)}$  is regular  $> \kappa$  strictly increasing with  $n$ , stipulating  $\lambda_{(i,n_i)} = \mu$ .

Then for some  $\langle m_i : i < \kappa \rangle \in \prod_{i < \kappa} (n_i + 1)$  and  $B \in D$  we have:

( $\alpha$ )  $\mu \leq \text{tcf}(\prod_{i < \kappa} \lambda_{(i,m_i)} / D)$

( $\beta$ )  $\mu > \max \text{pcf}\{\lambda_{(i,n)} : i \in B \text{ and } n < m_i\}$ .

*Proof.* We try to choose by induction on  $\zeta < \text{reg}(D), B_\zeta$  and  $\langle n_i^\zeta : i < \kappa \rangle$  such that:

(i)  $B_\zeta \in D$

(ii)  $n_i^\zeta < n_i$  non-decreasing in  $\zeta$

(iii)  $B_\zeta = \{i : n_i^\zeta < n_i^{\zeta+1}\}$  and

(iv)  $\max \text{pcf}\{\lambda_{(i,n)} : i < \kappa \text{ and } n \leq n_i^\zeta\} < \mu$ .

If we succeed, then  $\{B_\zeta : \zeta < \text{reg}(D)\}$  exemplifies  $D$  is  $\text{reg}(D)$ -regular, contradiction. During the induction we choose  $B_\zeta$  in step  $\zeta + 1$ . For  $\zeta = 0$  try  $n_i^\zeta = 0$ , if this fails then  $m_i = 0$  (for  $i < \kappa$ ) is as required. For  $\zeta$  limit let  $n_i^\zeta = n_i^\xi$  for every  $\xi < \zeta$  large enough, this is O.K. as

$$\max \text{pcf}\{\lambda_{(i,n)} : i < \kappa \text{ and } n < n_i^\zeta\} \leq \prod_{\xi < \zeta} \max \text{pcf}\{\lambda_{(i,n)} : i < \kappa \text{ and } n \leq n_i^\xi\} < \mu$$

by assumption (b). Lastly, for  $\zeta = \xi + 1$ ,  $\{i < \kappa : n_i^\xi < n_i\} \in D$  (otherwise contradiction as  $\lambda_{(i,n_i)} = \mu$  and clause (iv) contradict assumption (d)), and if  $\mu \leq \text{tcf}(\prod_{i < \kappa} \lambda_{n_i^\xi+1}/D)$  we are done with  $m_i = n_i^\xi + 1$ , if not there is  $B_\xi \in D$  such that  $\max \text{pcf}\{\lambda_{n_i^\xi+1} : i \in B\} < \mu$  and let

$$n_i^\zeta = \begin{cases} n_i^\xi + 1 & \text{if } i \in B_\xi, n_i^\xi < n_i \\ n_i^\xi & \text{if otherwise.} \end{cases}$$

□<sub>6.2</sub>

**6.3 Lemma.** *Assume*

- (i)  $D$  is an ultrafilter on  $\kappa$
- (ii)  $\mu = \text{cf}(\mu)$  and  $\alpha < \mu \Rightarrow |\alpha|^{< \text{reg}(D)} < \mu$
- (iii) at least one of the following occurs:
  - (α)  $\alpha < \mu \Rightarrow |\alpha|^{\text{reg}(D)} < \mu$
  - (β)  $D$  is closed under decreasing sequences of length  $\text{reg}(D)$ .

Then there is a minimal  $g/D$  such that:

$$\mu = \text{tcf}\left(\prod_{i < \kappa} g(i)/D\right) \text{ and } \bigwedge_{i < \kappa} \text{cf}(g(i)) > \kappa.$$

We shall prove it somewhat later.

**6.4 Remark.** 1) Note that necessarily (in 6.3)

$$\{i < \kappa : g(i) \text{ a regular cardinal}\} \in D.$$

2)  $g$  is also  $<_D$ -minimal under:  $\mu \leq \text{tcf}\left(\prod_{i < \kappa} g(i)/D\right) \ \& \ \{i : \text{cf}(g(i)) > \kappa\} \in D.$

[Why? assume  $g' <_D g, \mu \leq \text{tcf}\left(\prod_{i < \kappa} g'(i)/D\right)$ , and  $X = \{i : \text{cf}(g(i)) \leq \kappa\} =$

modified:2005-07-06

(589) revision:2005-07-05

$\emptyset \bmod D$ ; clearly  $\mu \leq \text{tcf}\left(\prod_{i<\kappa} \text{cf}(g'(i))/D\right)$ . If  $\text{Lim}_D \text{cf}(g'(i))$  is singular, by [Sh:g, II,1.4](1),p.50 for some  $\langle \lambda_i : i < \kappa \rangle$ , we have  $\mu = \text{tcf}(\prod \lambda_i/D)$  and  $\text{Lim}_D \lambda_i = \text{Lim}_D \text{cf}(g(i))$  and  $(\forall i)[\text{cf}(g(i)) > \kappa \rightarrow \lambda_i \geq \kappa]$ , so again without loss of generality  $\bigwedge_{i<\kappa} \lambda_i > \kappa$ . Now  $\langle \lambda_i : i < \kappa \rangle$  contradicts the choice of  $g$ . If  $\text{Lim}_D \text{cf}(g(i))$  is regular, it is  $\mu$  and all is easier.]

3) If  $|\kappa^\kappa/D| < \mu$  then we can omit (in the conclusion of 6.3 and of 6.4(2)) the clause “ $\{i : \text{cf}(g(i)) > \kappa\} \in D$ ”.

**6.5 Conclusion.** If assumptions (i)-(iii) of 6.3 hold and

$$(iv) \quad \mu > 2^\kappa$$

then without loss of generality each  $g(i)$  is a regular cardinal and  $\left(\prod_{i<\kappa} g(i)/D, <_D\right)$  is  $\mu$ -like (i.e. of cardinality  $\mu$  but every proper initial segment has smaller cardinality).

**6.6 Remark.** We use  $\mu > 2^\kappa$  in 6.5 rather than  $\mu > |\kappa^\kappa/D|$  as in 6.4(3) (which concerns 6.3, 6.4(3)) as the proof of 6.5 uses 1.4.

*Proof of 6.5.* If  $D$  is  $\aleph_1$ -complete this is trivial, so assume not hence  $\text{reg}(D) > \aleph_0$ .

Let  $g \in {}^\kappa(\mu+1)$  be as in 6.3, so without loss of generality as in 6.4(2), and remember 6.4(1) so without loss of generality each  $g(i)$  is a regular cardinal. Clearly  $\prod_{i<\kappa} g(i)$  has cardinality  $\geq \mu$ . Assume first  $\mu = \chi^+$ .

Let  $g' \in \prod_{i<\kappa} g(i)$ , then by 6.4(3) and choice of  $g$

$$\sup\{\text{tcf} \prod \lambda_i/D : \lambda_i \leq g'(i) \text{ for } i < \kappa\} \leq \chi.$$

But as  $\text{reg}(D) > \aleph_0$  by clause (ii) of the assumption we have  $\alpha < \mu \Rightarrow |\alpha|^{\aleph_0} < \mu$  so 1.5 applies (say for  $J = \{\kappa \setminus A : A \in D\}$ , as  $D$  is an ultrafilter clearly  $T_J^2(f) = \left(\prod_{i<\kappa} f(i)/D\right)$  and by assumption (ii), clause (e) of 1.5 holds. So we get

$$\left|\prod_{i<\kappa} g'(i)/D\right| \leq \chi, \text{ so really } \prod_{i<\kappa} g(i)/D \text{ is } \mu\text{-like.}$$

If  $\mu$  is not a successor, then it is weakly inaccessible and  $\mu = \sup(Z)$ , where  $Z = \{\chi^+ : |\kappa^\kappa/D| < \chi^{\aleph_0} = \chi < \mu\}$ , so for each  $\chi \in Z$  we can find  $g_\chi \in {}^\kappa(\mu + 1)$  such that  $\prod_{i < \kappa} g_\chi(i)/D$  is  $\chi$ -like so necessarily for  $\chi_1 < \chi_2$  in  $Z$  we have  $g_{\chi_1} <_D g_{\chi_2}$ .

It is enough to find a  $<_D$ -lub for  $\langle f_\chi : \chi \in Z \rangle$ , and as  $\mu > 2^\kappa$  this is immediate.

□<sub>6.5</sub>

*Proof of 6.3.* First try to choose, by induction on  $\alpha$ ,  $f_\alpha$  such that:

(A)  $f_\alpha \in {}^\kappa(\mu + 1)$

(B)  $\mu = \text{tcf}\left(\prod_{i < \kappa} f_\alpha(i)/D\right)$

(C)  $\beta < \alpha \Rightarrow f_\alpha <_D f_\beta$

(D) each  $f_\alpha(i)$  is a regular cardinal  $> \kappa$ .

Necessarily for some  $\alpha^*$  we have:  $f_\alpha$  is well-defined iff  $\alpha < \alpha^*$ . Now  $\alpha^*$  cannot be zero as the constant function with value  $\mu$  can serve as  $f_0$ . Also if  $\alpha^*$  is a successor ordinal, say  $\alpha^* = \beta + 1$ , then  $f_\beta$  is as required in the desired conclusion (by 6.4(2)'s proof).

So  $\alpha^*$  is a limit ordinal, and by passing to a subsequence, without loss of generality  $\alpha^* = \text{cf}(\alpha^*)$  and call it  $\theta$ .

Without loss of generality

(E)  $\mu = \max \text{pcf}\{f_\alpha(i) : i < \kappa\}$ .

We now try to choose by induction on  $\zeta < \text{reg}(D)$  the objects  $\alpha_\zeta, A_\zeta, \mathfrak{b}_\zeta$  such that:

(a)  $\alpha_\zeta < \theta$  is strictly increasing with  $\zeta$

(b)  $A_\zeta \in D$

(c)  $\mathfrak{b}_\zeta \subseteq \{f_{\alpha_\xi}(i) : \xi \leq \zeta, \text{ and } i \in A_\xi\}$

(d)  $\mathfrak{b}_\zeta$  is increasing with  $\zeta$

(e)  $\max \text{pcf}(\mathfrak{b}_\zeta) < \mu$

(f) for each  $i$  the sequence

$\langle f_{\alpha_\xi}(i) : \xi \leq \zeta \text{ and } i \in A_\xi \text{ and } f_{\alpha_\xi}(i) \notin \mathfrak{b}_\zeta \rangle$  is strictly decreasing

(g)  $\alpha_0 = 0, A_0 = \kappa, \mathfrak{b}_\zeta = \emptyset$

(h)  $\alpha_{\zeta+1} = \alpha_\zeta + 1$  and

$A_{\zeta+1} = \{i \in A_\zeta : f_{\alpha_{\zeta+1}}(i) < f_{\alpha_\zeta}(i) \text{ and } f_{\alpha_\zeta}(i) \in \mathfrak{b}_\zeta\}$

modified:2005-07-06

(589) revision:2005-07-05

- (i) for  $\zeta$  limit,  $\alpha_\zeta$  is the first  $\alpha < \theta$  which is  $\geq \bigcup_{\varepsilon < \zeta} \alpha_\varepsilon$  such that for some  $B \in D$  we have:  
 $\mu > \max \text{pcf}\{f_{\alpha_\xi}(i) : \xi < \zeta, i \in A_\xi \text{ and } i \in B \text{ and } f_{\alpha_\xi}(i) \leq f_\alpha(i)\}$
- (j)  $\mathfrak{b}_{\zeta+1} = \mathfrak{b}_\zeta$
- (k) for  $\zeta$  limit  $A_\zeta$  satisfies the requirements on  $B$  in clause (i) and  
 $\mathfrak{b}_\zeta = \bigcup_{\varepsilon < \zeta} \mathfrak{b}_\varepsilon \cup \bigcup \{f_\xi(i) : \xi < \zeta \text{ and } i \in A_\zeta, A_\xi \cap A_\zeta \text{ and } f_{\alpha_\xi}(i) \leq f_{\alpha_\zeta}(i)\}$
- (l) for  $\xi \leq \zeta$  we have  $\{i \in A_\xi : f_{\alpha_\xi}(i) \notin \mathfrak{b}_\zeta\} \in D$ .

So for some  $\zeta^* \leq \text{reg}(D)$  we have  $(\alpha_\zeta, A_\zeta, \mathfrak{b}_\zeta)$  is well defined iff  $\zeta < \zeta^*$ . We check the different cases and get a contradiction in each (so  $\alpha^*$  must have been a successor ordinal giving the desired conclusion).

CASE 1:  $\zeta^* = 0$ .

We choose  $\alpha_0 = 0, A_0 = \kappa, \mathfrak{b}_0 = \emptyset$ ; so clause (g) holds, first part of clause (a) (i.e.  $\alpha_\zeta < \theta$ ) holds, clause (b) and clause (c) are totally trivial, clause (e) holds as  $\max \text{pcf}(\emptyset) = 0$  (formally we should have written  $\sup \text{pcf}(\mathfrak{b}_\zeta)$ ), clause (f) speaks on the empty sequence, and the other clauses are empty in this case.

CASE 2:  $\zeta^* = \zeta + 1$ .

We choose  $\alpha_{\zeta^*} = \alpha_{\zeta+1} = \alpha_\zeta + 1, A_{\zeta^*} = \{i \in A_\zeta : f_{\alpha_{\zeta+1}}(i) < f_{\alpha_\zeta}(i) \text{ and } f_{\alpha_\zeta}(i) \notin \mathfrak{b}_\zeta\}$  and  $\mathfrak{b}_{\zeta+1} \supseteq \mathfrak{b}_\zeta$  is defined by clause (j). Clearly  $\alpha_\zeta < \alpha_{\zeta+1} < \theta$  and  $A_{\zeta+1} \in D$  as  $A_\zeta \in D$  and  $f_{\alpha_{\zeta+1}} <_D f_{\alpha_\zeta}$  and  $\{i : f_{\alpha_\zeta}(i) \notin \mathfrak{b}_\zeta\} \in D$  by clause (l); so clause (b) holds. Now clause (a) holds trivially and clauses (g) and (i) are irrelevant. Clause (h) holds by our choice.

For clause (f), the new cases are when  $f_{\alpha_{\zeta+1}}(i)$  appears in the sequence, i.e.,  $i \in A_{\zeta+1}$  such that  $f_{\alpha_{\zeta+1}}(i) \notin \bigcup_{\xi \leq \zeta+1} \mathfrak{b}_\xi = \mathfrak{b}_{\zeta+1} = \mathfrak{b}_\zeta$  but  $i \in A_{\zeta+1} \Rightarrow i \in A_\zeta$  &  $f_{\alpha_\zeta}(i) \notin \mathfrak{b}_\zeta$  so also  $f_{\alpha_\zeta}(i)$  appears in the sequence and as  $i \in A_{\zeta+1} \Rightarrow f_{\alpha_\zeta}(i) > f_{\alpha_{\zeta+1}}(i) = f_{\alpha_{\zeta+1}}(i)$  plus the induction hypothesis; we are done.

As for clause (l) for  $\xi \leq \zeta + 1$ , if  $\xi \leq \zeta$  this holds by the induction hypothesis (as  $\mathfrak{b}_{\zeta+1} = \mathfrak{b}_\zeta$ ) so assume  $\xi = \zeta + 1$ . Clearly  $\{i \in A_\xi : f_{\alpha_\xi}(i) \notin \mathfrak{b}_{\zeta+1}\} = A_\xi \cap \{i < \kappa : f_{\alpha_\xi}(i) \notin \mathfrak{b}_{\zeta+1}\}$ . Now the first belongs to  $D$  by clause (b) proved above and the second belongs to  $D$  as  $\max \text{pcf}(\mathfrak{b}_{\zeta+1}) < \mu$  by clause (e) proved below as

$$\text{tcf}\left(\prod_{i < \kappa} f_{\alpha_\xi}(i)/D\right) = \mu \text{ by clause (B).}$$

We have chosen  $\mathfrak{b}_{\zeta+1} = \mathfrak{b}_\zeta$ , so (using the induction hypothesis) clauses (c), (d), (e) trivially hold and also clause (j) holds by the choice of  $\mathfrak{b}_{\zeta^*}$ , and clause (k) is irrelevant so we are done.

CASE 3:  $\zeta^* = \zeta$  is a limit ordinal  $< \text{reg}(D)$ .

Let  $\mathfrak{b}_\zeta^* = \bigcup_{\xi < \zeta} \mathfrak{b}_\xi$ , so by basic pcf:

$$\boxtimes \quad \max \text{pcf}(\mathfrak{b}_\zeta^*) \leq \prod_{\xi < \zeta} \max \text{pcf}(\mathfrak{b}_\xi) < \mu$$

as

$$\mu = \text{cf}(\mu) \ \& \ (\forall \alpha < \mu)[|\alpha|^{< \text{reg}(D)} < \mu] \ \& \ \zeta < \text{reg}(D).$$

Now we try to define  $\alpha_\zeta$  by clause (i).

SUBCASE 3A:  $\alpha_\zeta$  is not well defined.

Let  $w_i = \{\xi < \zeta : i \in A_\xi \text{ and } f_{\alpha_\xi}(i) \notin \mathfrak{b}_\zeta^*\}$ . Note that by the induction hypothesis (clause (f)) for each  $\varepsilon < \zeta$  and  $i < \kappa$  we have the sequence  $\langle f_{\alpha_\xi}(i) : \xi < \varepsilon \text{ and } i \in A_\xi \text{ and } f_{\alpha_\xi}(i) \notin \mathfrak{b}_\varepsilon \rangle$  is strictly decreasing, so as  $\mathfrak{b}_\varepsilon \subseteq \mathfrak{b}_\zeta^*$  clearly  $\langle f_{\alpha_\xi}(i) : \xi < \varepsilon \text{ and } \xi \in w_i \rangle$  is strictly decreasing. As this holds for each  $\varepsilon < \zeta$  and  $\zeta$  is a limit ordinal, clearly  $\langle f_{\alpha_\xi}(i) : \xi \in w_i \rangle$  is strictly decreasing hence  $w_i$  is finite.

Now for each  $B \in D$  we have (first inequality by clause (E) and clause (b) on the induction hypothesis on  $\zeta$ , second by the definition of the  $w_i$ 's)

$$\begin{aligned} \mu &\leq \max \text{pcf} \left\{ f_\xi(i) : \xi < \zeta, i \in A_\xi \text{ and } i \in B \right\} \\ &\leq \max \left\{ \max \text{pcf}(\mathfrak{b}_\zeta), \max \text{pcf} \{ f_\xi(i) : \xi \in w_i \text{ and } i \in B \} \right\}, \end{aligned}$$

and  $\max \text{pcf}(\mathfrak{b}_\zeta^*) < \mu$  as said above, hence necessarily

$$(*) \quad B \in D \Rightarrow \mu \leq \max \text{pcf} \{ f_{\alpha_\xi}(i) : \xi \in w_i \text{ and } i \in B \}.$$

As  $w_i$  is finite and each  $f_\alpha(i)$  is a regular cardinal  $> \kappa$  we have  $\{i : w_i \neq \emptyset\} \in D$ .

By Claim 6.2 (the case there of  $\{i : m_i = n_i\} \in D$  is impossible by (\*) above) we can find  $g \in \prod_{i < \kappa} w_i/D$ , more exactly  $g \in {}^\kappa \text{Ord}, w_i \neq \emptyset \Rightarrow g(i) \in w_i$  and  $B \in D$

such that:

$$\begin{aligned} (\alpha) \quad \mu &\leq \text{tcf} \left( \prod_{i < \kappa} g(i)/D \right) \\ (\beta) \quad \mu &> \max \text{pcf} \{ f_{\alpha_\xi}(i) : \xi \in w_i \text{ and } i \in B \text{ and } f_{\alpha_\xi}(i) < g(i) \}. \end{aligned}$$

Now by the choice of  $\langle f_\alpha : \alpha < \theta \rangle$  and clause  $(\alpha)$  necessarily (and [Sh:g, Ch.II,1.4](1),p.50) for some  $\alpha < \theta$  we have  $f_\alpha <_D g$ . Now for  $\xi < \zeta$ , let  $B_\alpha^\xi = \{i < \kappa : f_\alpha(i) \geq f_{\alpha_\xi}(i)\}$ , if  $B_\alpha^\xi \in D$  then  $B^* = \{i < \kappa : \xi \in w_i \text{ and } i \in B \text{ and } g_\alpha(i) > f_{\alpha_\xi}(i)\} \supseteq \{i < \kappa : i \in A_\xi\} \cap \{i < \kappa : f_\alpha(i) \notin \mathfrak{b}_\zeta^*\} \cap \{i < \kappa : f_\alpha(i) \geq f_{\alpha_\xi}(i)\}$  which is the intersection of three members of  $D$  hence belongs to  $D$ , but  $\{f_{\alpha_\xi}(i) : i \in B^*\}$  is included in the set in the right side of clause  $(\beta)$  hence  $\mu > \max \text{pcf}\{f_{\alpha_\xi}(i) : i \in B^*\}$  contradicting  $B^* \in D$ ,  $\text{tcf}(\prod_{i < \kappa} f_{\alpha_\xi}(i)/D) = \mu$ . So necessarily  $B_\alpha^\xi \notin D$ , hence  $f_\alpha <_D f_{\alpha_\xi}$  hence  $\alpha < \alpha_\xi$ . So  $\bigcup_{\xi < \zeta} \alpha_\xi \leq \alpha < \theta$ . Let  $B' = B \cap \{i < \kappa : f_\alpha(i) < g(i)\}$  so  $B' \in D$  and [first inclusion by the choice of  $B'$ , second inclusion by the choice of  $\mathfrak{b}_\zeta^*$ ]

$$\begin{aligned} & \left\{ f_{\alpha_\xi}(i) : \xi < \zeta, i \in A_\xi \text{ and } i \in B' \text{ and } f_{\alpha_\xi}(i) \leq f_\alpha(i) \right\} \subseteq \\ & \left\{ f_{\alpha_\xi}(i) : \xi < \zeta, i \in A_\xi \text{ and } i \in B \text{ and } f_{\alpha_\xi}(i) < g(i) \right\} \subseteq \\ & \mathfrak{b}_\zeta^* \cup \left\{ f_{\alpha_\xi}(i) : \xi \in w_i \text{ and } f_{\alpha_\xi}(i) < g(i) \right\} \end{aligned}$$

hence

$$\begin{aligned} & \max \text{pcf} \left\{ f_{\alpha_\xi}(i) : \xi < \zeta, i \in A_\xi \text{ and } i \in B' \text{ and } f_{\alpha_\xi}(i) \leq f_\alpha(i) \right\} \leq \\ & \max \left\{ \max \text{pcf}(\mathfrak{b}_\zeta), \max \text{pcf} \left\{ f_{\alpha_\xi}(i) : \xi \in w_i \text{ and } i \in B \text{ and } \right. \right. \\ & \left. \left. f_{\alpha_\xi}(i) < g(i) \right\} \right\} < \mu \end{aligned}$$

(the first term is  $< \mu$  as the statement  $\boxtimes$  was proved in the beginning of Case 3, the second term is  $< \mu$  by clause  $(\beta)$ ). So  $\alpha$  is as required in clause (i) so  $\alpha_\zeta$  is well defined; contradiction.

CASE 3B:  $\alpha_\zeta$  is well defined.

Let  $B \in D$  exemplify it. We choose  $A_\zeta$  as  $B$  and we define  $\mathfrak{b}_\zeta$  by clause (k).

Now clause (a) follows from clause (i) (which holds by the assumption of the subcase), clause (b) holds by the choice of  $B$  (and of  $A_\zeta$ ), clause (c) by the choice of  $\mathfrak{b}_\zeta$ , clause (d) by the choice of  $\mathfrak{b}_\zeta$ , clause (e) by the choice of  $\mathfrak{b}_\zeta$ . Now for clause (f) by the induction hypothesis and clause (d) we should consider only  $f_{\alpha_\xi}(i) > f_{\alpha_\zeta}(i)$



when  $\xi < \zeta, i \in A_\xi \cap A_\zeta$  and  $f_{\alpha_\xi}(i), f_{\alpha_\zeta}(i) \notin \mathfrak{b}_\zeta$ , but clauses (i) + (k) (i.e. the choice of  $\mathfrak{b}_\zeta$ ) take care of this, clauses (g), (h), (j) are irrelevant, and clause ( $\ell$ ) follows from clause (e).

So we are done.

CASE 4:  $\zeta^* = \text{reg}(D)$ .

The proof is split according to the two cases in the assumption (iii).

SUBCASE 4A:  $\alpha < \mu \Rightarrow |\alpha|^{\text{reg}(D)} < \mu$ .

Let  $\mathfrak{b} = \cup\{\mathfrak{b}_\xi : \xi < \zeta^*\}$  so  $\max \text{pcf}(\mathfrak{b}) < \mu$ , hence for each  $\xi < \zeta^*$  we have  $A'_\xi =: \{i \in A_\xi : f_{\alpha_\xi}(i) \notin \mathfrak{b}\} \in D$ . Let  $w_i = \{\xi < \zeta^* : i \in A'_\xi \text{ and } f_{\alpha_\xi}(i) \notin \mathfrak{b}\}$ . Now for any  $\zeta < \zeta^*$  and  $i < \kappa$  the sequence  $\langle f_{\alpha_\xi}(i) : \xi < \zeta \text{ and } \xi \in w_i \rangle$  is strictly decreasing (by clause (f)) hence  $\langle f_{\alpha_\xi}(i) : \xi < \zeta^* \text{ and } \xi \in w_i \rangle$  is strictly decreasing hence  $w_i$  is finite. Also for each  $\xi < \zeta^*$  the set  $A'_\xi$  belongs to  $D$ , so  $\{A'_\xi : \xi < \zeta^*\}$  exemplifies  $D$  is  $|\zeta^*|$ -regular, but  $\zeta^* = \text{reg}(D)$ , contradiction.

SUBCASE 4B:  $D$  is closed under decreasing sequences of length  $\text{reg}(D)$ .

Let  $\mathfrak{b} = \bigcup_{\zeta < \zeta^*} \mathfrak{b}_\zeta$ .

In this case, for each  $\xi < \zeta^*$ , the sequence  $\langle \{i \in A_\xi : f_{\alpha_\xi}(i) \notin \mathfrak{b}_\zeta\} : \zeta \in [\xi, \zeta^*] \rangle$  is a decreasing sequence of length  $\zeta^* = \text{reg}(D)$  of members of  $D$  so the intersection,  $A'_\xi = \{i \in A_\xi : f_{\alpha_\xi}(i) \notin \mathfrak{b}\} \in D$ , and we continue as in the first subcase.  $\square_{6.3}$

**6.7 Definition.** 1) For an ultrafilter  $D$  on  $\kappa$  let  $\text{reg}'(D)$  be:  $\text{reg}(D)$  if  $D$  is closed under intersection of decreasing sequences of length  $\text{reg}(D)$  and  $(\text{reg}(D))^+$  otherwise.  
2)  $\text{reg}''(D)$  is:  $\text{reg}(D)$  iff  $(a)^-$  below holds and  $(\text{reg}(D))^+$  otherwise

(a)  $\text{reg}'(D) = \text{reg}(D)$  or just

(a)<sup>-</sup> letting  $\theta = \text{reg}(D)$ , in  $\theta^\kappa/D$  there is a  $<_D$ -first function above the constant functions.

**6.8 Theorem.** *If  $D$  is an ultrafilter on  $\kappa$  and  $\theta = \text{reg}'(D)$  then  $\mu = \mu^{<\theta} \geq |2^\kappa/D| \Rightarrow \mu \in \{\prod \lambda_i/D : \lambda_i \in \text{Card}\}$ .*

*Proof.* Apply Lemma 6.5 with  $D, \kappa, \mu^+$  here standing for  $D, \kappa, \mu$  there; note that assumption (iii) there holds as the definition of  $\text{reg}'(D)(= \theta)$  was chosen appropriately.

Let  $g^*/D = \langle \lambda_i^* : i < \kappa \rangle$  be as there, so as  $\left( \prod_{i < \kappa} \lambda_i^*/D \right)$  is  $\mu^+$ -like, for some  $f \in \prod_{i < \kappa} \lambda_i$ , we have  $|\prod_{i < \kappa} f(i)/D| = \mu$  as required.  $\square_{6.8}$

*Remark.* Can  $\text{reg}'(D) \neq \text{reg}(D)$ ? This is equivalent to:  $D$  is not closed under intersections of decreasing sequences of length  $\theta = \text{reg}(D)$ . So if  $\text{reg}'(D) \neq \text{reg}(D) = \theta$  then  $\theta$  is regular and for some function  $\mathbf{i} : \kappa \rightarrow \theta$  the ultrafilter  $D' = \{A \subseteq \theta : \mathbf{i}^{-1}(A) \in D\}$  is an ultrafilter on  $\theta$ , with  $\text{reg}(D') = \theta$  so  $D'$  is not regular.

This leads to the well known problem (Kanamori [Kn]): if  $D$  is a uniform ultrafilter on  $\kappa$  with  $\text{reg}(D) = \kappa$  does  $\kappa^\kappa/D$  have a first function above the constant ones?

6.9 *Fact.* If  $\mu = \theta = \text{reg}(D) < \text{reg}'(D)$ ,  $\mu = \sum_{i < \theta} \mu_i$ ,  $\mu_i^\kappa = \mu_i < \mu_{i+1}$  and

$$|\prod_{i < \kappa} f(i)/D| \geq \mu \text{ then } |\prod_{i < \kappa} f(i)/D| \geq \mu^\theta = \mu^\kappa.$$

## REFERENCES.

- [Bay] Timothy James Bays. *Multi-cardinal phenomena in stable theories*. PhD thesis, UCLA, 1994.
- [CuSh 541] James Cummings and Saharon Shelah. Cardinal invariants above the continuum. *Annals of Pure and Applied Logic*, **75**:251–268, 1995. arxiv:math.LO/9509228.
- [For] Matthew Foreman. ??? PhD thesis.
- [Kn] Akihiro Kanamori. Weakly normal filters and irregular ultra-filters. *Trans. of A.M.S.*, **220**:393–396, 1976.
- [LMSH 198] Jean Pierre Levinski, Menachem Magidor, and Saharon Shelah. Chang’s conjecture for  $\aleph_\omega$ . *Israel Journal of Mathematics*, **69**:161–172, 1990.
- [MgSh 433] Menachem Magidor and Saharon Shelah. Length of Boolean algebras and ultraproducts. *Mathematica Japonica*, **48**(2):301–307, 1998. arxiv:math.LO/9805145.
- [M] J. Donald Monk. Cardinal functions of Boolean algebras. circulated notes.
- [RoSh 534] Andrzej Rosłanowski and Saharon Shelah. Cardinal invariants of ultraproducts of Boolean algebras. *Fundamenta Mathematicae*, **155**:101–151, 1998. arxiv:math.LO/9703218.
- [Sh:a] Saharon Shelah. *Classification theory and the number of nonisomorphic models*, volume 92 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam-New York, xvi+544 pp, \$62.25, 1978.
- [Sh 71] Saharon Shelah. A note on cardinal exponentiation. *The Journal of Symbolic Logic*, **45**:56–66, 1980.
- [Sh 111] Saharon Shelah. On power of singular cardinals. *Notre Dame Journal of Formal Logic*, **27**:263–299, 1986.
- [Sh:c] Saharon Shelah. *Classification theory and the number of nonisomorphic models*, volume 92 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, xxxiv+705 pp, 1990.
- [Sh 351] Saharon Shelah. Reflecting stationary sets and successors of singular cardinals. *Archive for Mathematical Logic*, **31**:25–53, 1991.

- [Sh 355] Saharon Shelah.  $\aleph_{\omega+1}$  has a Jonsson Algebra. In *Cardinal Arithmetic*, volume 29 of *Oxford Logic Guides*, chapter II. Oxford University Press, 1994.
- [Sh:g] Saharon Shelah. *Cardinal Arithmetic*, volume 29 of *Oxford Logic Guides*. Oxford University Press, 1994.
- [Sh 430] Saharon Shelah. Further cardinal arithmetic. *Israel Journal of Mathematics*, **95**:61–114, 1996. arxiv:math.LO/9610226.
- [Sh 497] Saharon Shelah. Set Theory without choice: not everything on cofinality is possible. *Archive for Mathematical Logic*, **36**:81–125, 1997. A special volume dedicated to Prof. Azriel Levy. arxiv:math.LO/9512227.
- [Sh 506] Saharon Shelah. The pcf-theorem revisited. In *The Mathematics of Paul Erdős, II*, volume 14 of *Algorithms and Combinatorics*, pages 420–459. Springer, 1997. Graham, Nešetřil, eds.. arxiv:math.LO/9502233.
- [Sh 620] Saharon Shelah. Special Subsets of  ${}^{\text{cf}(\mu)}\mu$ , Boolean Algebras and Maharam measure Algebras. *Topology and its Applications*, **99**:135–235, 1999. 8th Prague Topological Symposium on General Topology and its Relations to Modern Analysis and Algebra, Part II (1996). arxiv:math.LO/9804156.
- [Sh 513] Saharon Shelah. PCF and infinite free subsets in an algebra. *Archive for Mathematical Logic*, **41**:321–359, 2002. arxiv:math.LO/9807177.