

Embedding Cohen algebras using pcf theory

Saharon Shelah*

Institute of Mathematics
The Hebrew University of Jerusalem
91904 Jerusalem, Israel
and
Department of Mathematics
Rutgers University
New Brunswick, NJ 08854, USA

done: July 1995

printed: October 6, 2003

Abstract

Using a theorem from pcf theory, we show that for any singular cardinal ν , the product of the Cohen forcing notions on κ , $\kappa < \nu$ adds a generic for the Cohen forcing notion on ν^+ .

The following question (problem 5.1 in Miller's list [Mi91]) is attributed to Rene David and Sy Friedman:

Does the product of the forcing notions $\aleph_n^{>2}$ add a generic for the forcing $\aleph_{\omega+1}^{>2}$?

We show here that the answer is yes in ZFC. Previously Zapletal [Za] has shown this result under the assumption $\square_{\aleph_{\omega+1}}$.

In fact, a similar theorem can be shown about other singular cardinals as well. The reader who is interested only in the original problem should read $\aleph_{\omega+1}$ for λ , \aleph_ω for μ and $\{\aleph_n : n \in (1, \omega)\}$ for \mathfrak{a} .

Definition 1 1. Let \mathfrak{a} be a set of regular cardinals. $\prod \mathfrak{a}$ is the set of all functions f with domain \mathfrak{a} satisfying $f(\kappa) \in \kappa$ for all $\kappa \in \mathfrak{a}$.

2. A set $\mathfrak{b} \subseteq \mathfrak{a}$ is bounded if $\sup \mathfrak{b} < \sup \mathfrak{a}$. \mathfrak{b} is cobounded if $\mathfrak{a} \setminus \mathfrak{b}$ is bounded.

*The research partially supported by "The Israel Science Foundation" administered by of The Israel Academy of Sciences and Humanities. Publication 595.

3. If J is an ideal on \mathfrak{a} , $f, g \in \prod \mathfrak{a}$, then $f <_J g$ means $\{\kappa \in \mathfrak{a} : f(\kappa) \not\leq g(\kappa)\} \in J$. We write $\prod \mathfrak{a}/J$ for the partial (quasi)order $(\prod \mathfrak{a}, <_J)$.
4. $\lambda = \text{pcf}(\prod \mathfrak{a}/J)$ (λ is the true cofinality of $\prod \mathfrak{a}/J$) means that there is a strictly increasing cofinal sequence of functions in the partial order $(\prod \mathfrak{a}, <_J)$.
5. $\text{pcf}(\mathfrak{a}) = \{\lambda : \exists J \lambda = \text{pcf}(\prod \mathfrak{a}/J)\}$.

We will use the following theorem from pcf theory:

Lemma 2 *Let μ be a singular cardinal. Then there is a set \mathfrak{a} of regular cardinals below μ , $|\mathfrak{a}| = \text{cf}(\mu) < \min \mathfrak{a}$ and $\mu^+ \in \text{pcf}(\mathfrak{a})$.*

Moreover, we can even have $\text{pcf}(\prod \mathfrak{a}/J^{\text{bd}}) = \mu^+$, where J^{bd} is the ideal of all bounded subset of \mathfrak{a} .

PROOF See [Sh 355, theorem 1.5].

Theorem 3 *Let \mathfrak{a} be a set of regular cardinals, $\mu = \sup \mathfrak{a} \notin \mathfrak{a}$, $2^{<\lambda} = 2^\mu$, $\lambda > \mu$, $\lambda \in \text{pcf}(\mathfrak{a})$, and moreover:*

- (*) *There is an ideal J on \mathfrak{a} containing all bounded sets such that $\lambda = \text{pcf}(\prod \mathfrak{a}/J)$.*

Then the forcing notion $\prod_{\kappa \in \mathfrak{a}} \kappa^{>2}$ adds a generic for $\lambda^{>2}$.

Corollary 4 *If ν is a singular cardinal, and P is the product of the forcing notions $\kappa^{>2}$ for $\kappa < \nu$, then P adds a generic for $\nu^{+>2}$.*

PROOF By lemma 2 and theorem 3

Remark 5 1. *The condition (*) in the theorem is equivalent to:*

(**) *For all bounded sets $\mathfrak{b} \subset \mathfrak{a}$ we have $\lambda \in \text{pcf}(\mathfrak{a} \setminus \mathfrak{b})$.*

2. *Clearly the assumption $2^{<\lambda} = 2^\mu$ is necessary, because otherwise the forcing notion $\prod_{\kappa \in \mathfrak{a}} \kappa^{>2}$ would be too small to add a generic for $\lambda^{>2}$.*

Proof of the theorem: By our assumption we have some ideal J containing all bounded sets such that $\text{pcf}(\prod \mathfrak{a}/J) = \lambda$.

We will write $\forall^J \kappa \in \mathfrak{a} \varphi(\kappa)$ for

$$\{\kappa \in \mathfrak{a} : \neg \varphi(\kappa)\} \in J$$

So we have a sequence $\langle f_\alpha : \alpha < \lambda \rangle$ such that

- (a) $f_\alpha \in \prod \mathfrak{a}$
- (b) If $\alpha < \beta$, then $\forall^J \kappa \in \mathfrak{a} f_\alpha(\kappa) < f_\beta(\kappa)$
- (c) $\forall f \in \prod \mathfrak{a} \exists \alpha \forall^J \kappa \in \mathfrak{a} f(\kappa) < f_\alpha(\kappa)$.

The next lemma shows that if we allow these functions to be defined only almost everywhere, then we can additionally assume that in each block of length μ these functions have disjoint graphs:

Lemma 6 *Assume that \mathfrak{a} , λ , μ are as above. Then there is a sequence $\langle g_\alpha : \alpha < \lambda \rangle$ such that*

- (a) $\text{dom}(g_\alpha) \subseteq \mathfrak{a}$ *cobounded (so in particular $\forall^J \kappa \in \mathfrak{a} : \kappa \in \text{dom}(g_\alpha(\kappa))$).*
- (b) *If $\alpha < \beta$, then $\forall^J \kappa \in \mathfrak{a} g_\alpha(\kappa) < g_\beta(\kappa)$*
- (c) $\forall f \in \prod \mathfrak{a} \exists \alpha \forall^J \kappa \in \mathfrak{a} f(\kappa) < g_\alpha(\kappa)$. *Moreover, we may choose α to be divisible by μ .*
- (d) *If $\alpha < \beta < \alpha + \mu$, then $\forall \kappa \in \text{dom}(g_\alpha) \cap \text{dom}(g_\beta) : g_\alpha(\kappa) < g_\beta(\kappa)$.*

PROOF Let $\langle f_\alpha : \alpha < \lambda \rangle$ be as above. Now define $\langle g_\alpha : \alpha < \lambda \rangle$ by induction as follows:

If $\alpha = \mu \cdot \zeta$, then let g_α be any function that satisfies $g_\beta <_J g_\alpha$ for all $\beta < \alpha$, and also $f_\alpha <_J g_\alpha$. Such a function can be found because set of functions of size $< \lambda$ can be $<_J$ -bounded by some f_β .

If $\alpha = \mu \cdot \zeta + i$, $0 < i < \mu$, then let

$$g_\alpha(\kappa) = \begin{cases} g_{\mu \cdot \zeta}(\kappa) + i & \text{if } i < \kappa \\ \text{undefined} & \text{otherwise} \end{cases}$$

It is easy to see that (a)–(d) are satisfied.

Definition 7 1. Let P_κ be the set ${}^{\kappa > 2}$, partially ordered by inclusion (= sequence extension). Let $P = \prod_{\kappa \in \mathfrak{a}} P_\kappa$. [We will show that P adds a generic for ${}^{\lambda > 2}$]

- 2. Assume that $\langle g_\alpha : \alpha < \lambda \rangle$ is as in lemma 6.
- 3. Let $H : {}^\mu 2 \rightarrow {}^{\lambda > 2}$ be onto.
- 4. For $\kappa \in \mathfrak{a}$, let η_κ be the P_κ -name for the generic function from κ to 2. Define a P -name of a function $h : \lambda \rightarrow 2$ by

$$\dot{h}(\alpha) = \begin{cases} 0 & \text{if } \forall^J \kappa \in \mathfrak{a} \eta_\kappa(g_\alpha(\kappa)) = 0 \\ 1 & \text{otherwise} \end{cases}$$

5. For $\xi < \lambda$ let ρ_ξ be a P -name for the element of ${}^\mu 2$ that satisfies $\rho_\xi \simeq \underline{h} \upharpoonright [\mu \cdot \xi, \mu \cdot (\xi + 1))$, i.e.,

$$i < \mu \Rightarrow \Vdash_P \rho_\xi(i) = \underline{h}(\mu \cdot \xi + i).$$

Define $\rho \in {}^\lambda 2$ by

$$\rho = H(\rho_0) \frown H(\rho_1) \frown \cdots \frown H(\rho_\xi) \frown \cdots$$

Main Claim 8 ρ is generic for $\lambda > 2$.

Definition 9 For $\alpha < \lambda$ let $P^{(\alpha)}$ be the set of all conditions p satisfying $\forall^J \kappa : \text{dom}(p_\kappa) = g_\alpha(\kappa)$.

Remark 10 $\bigcup_{\zeta < \lambda} P^{(\mu \cdot \zeta)}$ is dense in P .

PROOF By lemma 6(c).

Fact 11 Let $\alpha = \mu \cdot \zeta$, $p \in P^{(\alpha)}$, $\sigma \in {}^\mu 2$. Then there is a condition $q \in P^{(\alpha + \mu)}$, $q \geq p$ and

$$\forall j < \mu \forall^J \kappa q_\kappa(g_{\alpha+j}(\kappa)) = \sigma(j)$$

PROOF Let $p = (p_\kappa : \kappa \in \mathfrak{a})$. There is a set $\mathfrak{b} \in J$ such that: For all $\kappa \in \mathfrak{a} \setminus \mathfrak{b}$ we have $\text{dom}(p_\kappa) = g_\alpha(\kappa)$. Define $q \in P^{(\alpha + \mu)}$, $q = (q_\kappa : \kappa \in \mathfrak{a})$ as follows:

$$q_\kappa(\gamma) = \begin{cases} p_\kappa(\gamma) & \text{if } \gamma \in \text{dom}(p_\kappa) \\ \sigma(j) & \text{if } \gamma = g_{\alpha+j}(\kappa), \kappa \in \mathfrak{a} \setminus \mathfrak{b} \\ 0 & \text{otherwise} \end{cases}$$

We have to explain why q is well-defined: First note that the first and the second case are mutually exclusive. Indeed, if $\gamma = g_{\alpha+j}(\kappa)$, then $\gamma > g_\alpha(\kappa)$, whereas $\kappa \notin \mathfrak{b}$ implies that $\text{dom}(p_\kappa) = g_\alpha(\kappa)$, so $\gamma \notin \text{dom}(p_\kappa)$.

Next, by the property (d) from lemma 6 there is no contradiction between various instances of the second case.

Hence we get that for all $j < \mu$, whenever $\kappa \in \mathfrak{a} \setminus \mathfrak{b}$, and $\kappa > j$, then $q_\kappa(g_{\alpha+j}(\kappa)) = \sigma(j)$. Since J contains all bounded sets, this means that $\forall^J \kappa : q_\kappa(g_{\alpha+j}(\kappa)) = \sigma(j)$.

Remark 12 Assume that $\alpha = \mu \cdot \zeta$, and p, q, σ are as above. Then $q \Vdash \rho_\zeta = \sigma$.

Proof of the main claim: Let $p \in P$, and $D \subseteq {}^{\lambda}2$ be a dense open set. We may assume that for some $\alpha^* < \lambda$, $\zeta^* < \lambda$ we have $\alpha^* = \mu \cdot \zeta^*$ and $p \in P^{(\alpha^*)}$, i.e., for some $\mathfrak{b} \in J$: $\forall \kappa \notin \mathfrak{b} : \text{dom}(p_\kappa) = g_{\alpha^*}(\kappa)$

So p decides the values of $h \upharpoonright \alpha^*$, and hence also the values of ρ_ζ for $\zeta < \zeta^*$. Specifically, for each $\zeta < \zeta^*$ we can define $\sigma_\zeta \in {}^{\mu}2$ by

$$\sigma_\zeta(i) = \begin{cases} 0 & \text{if } \forall^J \kappa p_\kappa(g_{\mu \cdot \zeta + i}(\kappa)) = 0 \\ 1 & \text{otherwise} \end{cases}$$

(Note that for all $\zeta < \zeta^*$, for all $i < \mu$, for almost all κ the value of $p_\kappa(g_{\mu \cdot \zeta + i}(\kappa))$ is defined.)

Clearly $p \Vdash \rho_\zeta = \sigma_\zeta$. Since D is dense and H is onto, we can now find $\sigma_{\zeta^*} \in {}^{\mu}2$ such that

$$H(\sigma_0) \frown \dots \frown H(\sigma_{\zeta^*}) \in D$$

Using 11 and 12, we can now find $q \geq p$ such that $q \Vdash \rho_{\zeta^*} = \sigma_{\zeta^*}$.

Hence $q \Vdash \rho \in D$, and we are done.

References

- [Mi91] Arnold Miller. Arnie Miller's problem list. In Haim Judah, editor, *Set Theory of the Reals*, volume 6 of *Israel Mathematical Conference Proceedings*, pages 645–654. Proceedings of the Winter Institute held at Bar–Ilan University, Ramat Gan, January 1991.
- [Za] Jindrich Zapletal. *Some Results in Set Theory and Boolean Algebras*. PhD thesis, Penn State University, 1995.
- [Sh 355] Saharon Shelah. $\aleph_{\omega+1}$ has a Jonsson Algebra. In *Cardinal Arithmetic*, volume 29 of *Oxford Logic Guides*, chapter II. Oxford University Press, 1994.