

**NNR REVISITED**  
**SH656**

SAHARON SHELAH

ABSTRACT. We are interested in proving that if we use CS iterations of forcing notions not adding reals that satisfies additional conditions then the limit forcing does not add reals. As a result we prove that we can amalgamate two earlier methods and prove the consistency with ZFC + G.C.H. of two statements gotten separately earlier: SH and non-club guessing. We also prove the consistency of further cases of “strong failure of club guessing” solving a problem of Justin Moore.

---

*Date:* January 26, 2015.

*2010 Mathematics Subject Classification.* Primary; Secondary:

*Key words and phrases.* set theory, forcing, set theory of the reals, iterated forcing, preservation theorem, no new reals.

The author thanks Alice Leonhardt for the beautiful typing. First typed 1997/June/17.

## Anotated Content

## §0 Introduction, pg. 4

[We give lengthy explanations of the problems and proofs for No-New-Reals iterations (for proper forcing, CS iterations).]

## §1 Preservation of not adding reals, pg.9

{a24} [We present sufficient conditions for CS iteration of proper forcing not to add reals. For this we define “reasonable parameters  $\mathfrak{p}$ ” and we have two main demands. One (clause (c) of Definition 1.8) is a weakening of “ $\alpha$ -proper for every  $\alpha < \omega_1$ ”. This time it has the form (on  $\mathbb{Q}_i$ ),  $\mathfrak{p}$ -proper which informally says that: if  $\mathfrak{p} \in N, Y \subseteq \{M \in N : M \text{ appropriate}\}$  is  $\alpha$ -large then for some  $(N, \mathbb{Q}_i)$ -generic condition  $q \geq p, q$  forces that  $\{M \in Y : M[\mathbf{G}_{\mathbb{Q}}] \cap \mathbf{V} = M\}$  is  $\alpha$ -large (the meaning of  $\alpha$ -large depends on  $\mathfrak{p}$ ), hence without loss of generality,  $\mathfrak{p}$  has length  $\omega_1$ . The other main demand

{a24} (clause (d) of Definition 1.8) is a “weak diamond preventive”.

{a48} We then show that  $\alpha$ -properness for  $\alpha < \omega_1$  is sufficient for the first main demand (in 1.19(3)). The demand on the games for  $\mathfrak{p}$  helps to prove the preservation of  $\mathfrak{p}$ -properness.]

## §2 Delayed properness, pg. 19

[The preservation theorem in the first section does not, for standard  $\mathfrak{p}$ , cover shooting a club  $C \subseteq \omega_1$  running away for  $C_\delta \subseteq \delta = \sup(C_\delta), C_\delta$  small (see §3). For this we will use  $(\mathfrak{p}, \alpha, \beta)$ -proper for enough pairs  $\alpha \leq \beta < \ell g(\mathfrak{p})$  (so starting from  $\beta$ -large we get  $\alpha$ -large; for many  $\alpha$  we can choose  $\beta = \alpha$  but during the inductive proof we pass through cases of  $\alpha < \beta$ ). Here we introduce various definitions and basic facts needed. We discuss axioms, version of the properties preserved by CS iterations and strengthening of the iteration Lemmas of §1.]

## §3 Example: shooting a thin club, pg.25

[We present the natural forcing showing  $\kappa = 2$  is interesting (not only  $\kappa = \aleph_0$ ) (from [Sh:b, Ch.VIII,§4]). We show that the natural forcing (see above) for running away from  $C_\delta \subseteq \delta$ , of small order type (see [Sh:f, Ch.XVIII,§2]) falls under our framework for delayed properness. We give examples: running away from  $\langle C_{\delta,0}, C_{\delta,1} : \delta < \omega_1 \text{ limit} \rangle, C_{\delta,0}, C_{\delta,1}$  are disjoint closed subsets of  $\delta$  with no restrictions on their order type so we ask for  $C, C \cap C_{\delta,0}$  or  $C \cap C_{\delta,1}$  is bounded in  $\delta$  and more.]

## §4 Second preservation of not adding reals, pg.32

[We give a sufficient condition for the limit forcing not to add reals. We here are weakening the demand “ $\mathfrak{p}$ -proper”, using  $(\mathfrak{p}, \alpha, f(\alpha))$ -proper instead  $(\mathfrak{p}, \alpha, \alpha)$ -proper, what we called delayed properness. The price is that here  $\mathfrak{p}$  has length of large cofinality, so essentially we catch our tails on a club of it. Also the Lemma here covers the examples.]

- §5 Problematic Forcing, pg.40  
[We discuss further generalizations.]
- §6 Spelling out the axioms, pg.41

## § 0. INTRODUCTION

{intro}

We try to explain our problems and results. If the explanations look opaque, try to return to them after reading at least part of the proof. Sections §0, §1 are based on lectures in the logic seminar in the Hebrew University, Spring 1997, whose participants I thank. On the history see in [Sh:f, Ch.V,§7,Ch.VIII,§4,Ch.XVIII,§1,§2], [Sh:666, §3].

Lately, Justin Moore makes a great advance: solving a problem from [Sh:666, §3], prove in ZFC + CH that, e.g. some proper forcing notion not adding and satisfying a “strong form of the medicine against weak diamond” has no generic, in fact, is a tree with no branch (earlier we know that if, e.g.  $\mathbf{V} = \mathbf{L}$  preservation of NNR fail). He raises a question and we deduce here a solution to it.

{y3}

This work was done and circulated in the late nineties and even submitted but for various reasons work was resumed only after listening to a lecture of Moore.

**Definition 0.1.** 1) Let  $K_0$  be the family of CS iterations,  $\bar{\mathbb{Q}} = \langle \mathbb{P}_i, \mathbb{Q}_i : i < \alpha \rangle$ , we denote  $\mathbb{P}_\alpha = \text{Lim}(\bar{\mathbb{Q}})$ .

2) We say  $\bar{\mathbb{Q}}$  is proper when each  $\mathbb{Q}_i$  is, hence  $\mathbb{P}_j/\mathbb{P}_i$  is proper for  $i < j \leq \alpha$  (see [Sh:b] or [Sh:f]).

3) We say  $\bar{\mathbb{Q}}$  is  $\omega$ -bounding or bounded when each  $\mathbb{Q}_i$  is (which means: every  $f \in (\omega\omega)^{\mathbf{V}^{\mathbb{P}_{i+1}}} = (\omega\omega)^{(\mathbf{V}^{\mathbb{P}_i})^{\mathbb{Q}_i}}$  is bounded<sup>1</sup> by some  $g \in (\omega\omega)^{\mathbf{V}^{\mathbb{P}_i}}$ , hence  $\mathbb{P}_j/\mathbb{P}_i$  is  $\omega$ -bounding for  $i < j \leq \alpha$ ).

4) We say  $\bar{\mathbb{Q}}$  is NNR if  $i < \alpha \Rightarrow \mathbb{P}_{i+1}$  adds no reals.

[Equivalently:  $i < \alpha \Rightarrow \mathbb{Q}_i$  adds no reals and  $\beta < \alpha \Rightarrow \mathbb{P}_\beta$  adds not reals.]

**Discussion 0.2.** It would be nice if also this (NNR) would be preserved in limit. But this is wrong for two known reasons, obstacles, explained below:

- ⊗<sub>1</sub> weak diamond
- ⊗<sub>2</sub> existence of clubs.

Our aim here is to weaken the medicine one uses against ⊗<sub>2</sub> for CS iterations of proper forcing notions. Let us explain the “obstacles”.

Concerning the weak diamond:

Let  $\bar{\eta} = \langle \eta_\delta : \delta < \omega_1, \delta \text{ limit} \rangle$ ,  $\eta_\delta = \langle \eta_\delta(n) : n < \omega \rangle$  where  $\eta_\delta$  is an  $\omega$ -sequence of ordinals (strictly) increasing with limit  $\delta$ . Let  $D$  be a non-principal ultrafilter on  $\omega$ .

{y6}

For  $f \in {}^{\omega}2$ ,  $\delta < \omega_1$  limit, let  $\text{Av}_D(f, \eta_\delta) = \ell$  iff  $\{n : f(\eta_\delta(n)) = \ell\} \in D$ .

*Question 0.3.* [CH] Given  $\bar{e} = \langle e_\delta : \delta < \omega_1 \text{ limit} \rangle$ ,  $e_\delta \in \{0, 1\}$  is there  $f \in {}^{\omega}2$  such that for a club of  $\delta < \omega_1$  we have  $e_\delta = \text{Av}_D(f, \eta_D)$ ?

Naturally, trying to prove consistency we should use a CS iteration  $\bar{\mathbb{Q}}$ , for simplicity we assume

{y6}

- (\*) (a)  $\mathbf{V} \models \text{GCH}$ ,
- (b)  $\bar{\mathbb{Q}} = \langle \mathbb{P}_i, \mathbb{Q}_i : i < \omega_2 \rangle$
- (c)  $\mathbb{Q}_i = \mathbb{Q}_{\bar{e}}$  (i.e.  $\mathbb{Q}_i = \mathbb{Q}_{\bar{e}}$  for some  $\bar{e}$  as in 0.3) where
  - $\mathbb{Q}_{\bar{e}} = \{f : \text{for some } \zeta < \omega_1, f \in {}^\zeta 2 \text{ and for every limit } \delta \leq \zeta \text{ we have } \text{Av}_D(f, \eta_\delta) = e_\delta\}$ .

<sup>1</sup>where  $\mathbb{Q}$  is bounded (in the universe  $\mathbf{V}$ ) if every  $f \in (\omega\omega)^{\mathbf{V}^{\mathbb{Q}}}$  is bounded by some  $g \in (\omega\omega)^{\mathbf{V}}$

This is a very nice forcing notion - it is proper (even  $< \omega_1$ -proper, see below) and NNR and for every  $\alpha < \omega_1$ ,  $\mathcal{I}_\alpha = \{f \in \mathbb{Q}_{\bar{e}} : \alpha \subseteq \text{Dom}(f)\}$  is a dense open subset of it.

But the weak diamond ([DvSh:65] or see [Sh:b, Ch.XII,§1] = [Sh:f, AP,§1]) tells us for this case that the answer is no, that is:  $\exists \bar{e} \forall f \in \omega_1 2 \exists^{\text{stat}} \delta (e_\delta \neq \text{Av}_D(f, \eta_\delta))$ .

In fact this holds for any function  $\text{Av}' : \bigcup_{\delta < \omega_1} \delta 2 \rightarrow \{0, 1\}$ .

Why is  $\mathbb{Q}_{\bar{e}}$  NNR?

Let  $\langle N_i : i \leq \omega^2 \rangle$  be increasing continuously with  $i, \mathbb{Q}_{\bar{e}} \in N_0$  where  $N_i \prec (\mathcal{H}(\chi), \in)$  is countable,  $\bar{N} \upharpoonright (i+1) \in N_{i+1}$ ; let  $\delta_i = \delta(i) = N_i \cap \omega_1$ . So  $\langle \delta_i : i \leq \omega^2 \rangle$  is strictly increasing continuously. But  $\{\eta_{\delta(\omega^2)}(n) : n < \omega\} \subset \delta(\omega^2)$  has order type  $\omega$ , so  $W = \{i < \omega^2 : \exists n (\delta_i \leq \eta_{\delta(\omega^2)}(n) < \delta_{i+1})\}$  has order type  $\omega^2$ .

So we can find  $\ell_n < \omega$  such that  $\bigwedge_{n < \omega} \bigwedge_{m < \omega} \omega \times n + \ell_n + m \notin W$ . Let  $p \in \mathbb{Q}_{\bar{e}} \cap N_0$ , let  $\tau : \omega \rightarrow \text{Ord}, \tau \in N_0$ . We choose by induction on  $n, p_n \in \mathbb{Q}_{\bar{e}}$  such that  $p \leq p_n, p_{n-1} \leq p_n, p_n \in N_{\omega n + \ell_n + 1}, p_n$  force some value to  $\tau(n)$  and on  $[\delta_{\omega m + \ell_m}, \delta_{\omega(m+1) + \ell_{m+1}}) \cap \text{Rang}(\eta_\delta) \setminus \text{Dom}(p)$  which agrees with  $e_{\delta(\omega^2)}$ .

So

- (\*) the desired demand on  $\bar{\mathbb{Q}}_i$  (guaranteeing  $\mathbb{P}_\alpha$  in NNR) should exclude the  $\mathbb{Q}_{\bar{e}}$ 's.

{y9}

**Definition 0.4.** Let  $K_1$  be the class of proper  $\omega$ -bounding iteration  $\bar{\mathbb{Q}} \in K_0$ , let  $K_2$  be the class of NNR iterations  $\bar{\mathbb{Q}} \in K_1$ , and let  $K_3$  be the class of  $\bar{\mathbb{Q}} \in K_2$  such that

(\*) if (A) then (B) where

- (A) (a)  $\chi$  is large enough (so  $\bar{a} \in \mathcal{H}(\chi)$ ),
- (b)  $N \prec (\mathcal{H}(\chi), \in)$  countable,
- (c)  $\bar{\mathbb{Q}} \in N$ ,
- (d)  $i \in \ell g(\bar{\mathbb{Q}}) \cap N$ ,
- (e)  $p \in \mathbb{P}_{i+1} \cap N$ ,
- (f)  $q_0, q_1 \in \mathbb{P}_i$  are  $(N, \mathbb{P}_i)$ -generic (i.e.  $q_\ell \Vdash "N[\mathbb{G}_{\mathbb{P}_i}] \cap \mathbf{V} = N"$ ) and moreover
- (g)  $q_\ell \Vdash "\mathbb{G}_{\mathbb{P}_i} \cap N = \mathbf{G}^*" \text{ and}$
- (h)  $p \upharpoonright i \leq q_\ell$

(B) we can find  $q'_0, q'_1, \mathbf{G}^{**}$  such that for  $\ell = 1, 2$  we have

- (a)  $q_\ell \leq q'_\ell \in \mathbb{P}_{i+1}$ ,
- (b)  $p \upharpoonright (i+1) \leq q'_\ell$
- (c)  $q'_\ell \Vdash "\mathbb{G}_P \cap N = \mathbf{G}^{**}"$ ,
- (d)  $q'_\ell$  is  $(N, \mathbb{P}_{i+1})$ -generic, so  $\mathbf{G}^{**} \subseteq \mathbb{P}_{i+1} \cap N$  is generic over  $N$ .

This tries to say:

We know  $\mathbb{G}_{\mathbb{P}_i} \cap N$  (as being  $\mathbf{G}^*$ ) and we are looking at  $N[\mathbf{G}^*]$  (formally, only its isomorphism type). So we know  $\mathbb{Q}_i^N[\mathbf{G}_i^*]$ .

We would like to find  $\mathbf{G}' \subseteq \mathbb{Q}_i^N[\mathbf{G}^*]$  generic over  $N[\mathbf{G}^*]$ , so that  $\mathbf{G}^*, \mathbf{G}'$  will determine  $\mathbf{G}^{**}$ . But we need a guarantee that  $\mathbf{G}'$  will have an upper bound in

$\mathbb{Q}_i[\mathbf{G}_{\mathbb{P}_i}]$ . If we know  $\mathbf{G}_{\mathbb{P}_i}$ , fine; but in a sense, we are given 2 candidates (by  $q_0, q_1$  and can increase then to  $q'_0 \upharpoonright i, q'_1 \upharpoonright i$ ) and have to find  $\mathbf{G}'$  “accepted” by both.

This (weak diamond) obstacle was overcome, with a price, i.e. more than being in  $K_3$ , [Sh:f, Ch.V,§7] using  $\aleph_1$ -completeness systems and [Sh:f, Ch.XVIII,§4] for 2-completeness systems (phrased in [Sh:f, Ch.XVIII,§2] without them).

Unfortunately, this does not suffice: there is an extreme example where e.g. if for  $q_0, q_1$  incompatible in  $\mathbb{Q}_i, \bar{E}_i$  a  $\mathbb{Q}_i$ -name of a club, for some  $\alpha(q_0, q_1)$  we have:

$$q_\ell \leq q'_\ell$$

$$q_\ell \Vdash “\bar{E}_i \cap \delta = E_i^\delta \Rightarrow E_i^{\delta_0} \cap E_1^{\delta_1} \setminus \alpha(q_0, q_1) \text{ is finite}”.$$

This represents the reason, the obstacle which we shall call  $\otimes_2$ , it was overcome (with a price) either with  $(< \omega_1)$ -properness or by a kind of “finite powers are proper” (see below).

{y12}

**Definition 0.5.** 1)  $\mathbb{Q}$  is  $\alpha$ -proper when:

if  $\bar{N} = \langle N_i : i \leq \alpha \rangle$  is  $\prec$ - increasing continuously,  $\alpha \in N_0$

$N_i \prec (\mathcal{H}(X), \in)$  countable,  $\bar{N} \upharpoonright (i+1) \in N_{i+1}$ ,

$\mathbb{Q} \in N_0$  and  $p \in \mathbb{Q} \cap N_0$

then there is  $q, p \leq q \in \mathbb{Q}_i$  such that  $q$  is  $(N_i, q)$ -generic for  $i \leq \alpha$ .

2) A forcing notion  $\mathbb{Q}$  is  $(<^+ \omega_1)$ -proper if the above holds for any  $\alpha < \omega_1$  even omitting “ $\alpha \in N_0$ ”. We say  $\mathbb{Q}$  is  $(< \omega_1)$ -proper if  $\mathbb{Q}$  is  $\alpha$ -proper for any  $\alpha < \omega_1$ .

{y14}

**Discussion 0.6.** So  $(< \omega_1)$ -proper is an antidote to such problems, i.e. against “reason  $\otimes_2$ ”. Okay for specializing a Aronszajn tree and many others, but it seems to me since [Sh:177] too strong: it kills the following:

{y15}

*Question 0.7.* Let  $\bar{C} = \langle C_\delta : \delta < \omega_1, \delta \text{ limit} \rangle, C_\delta \subseteq \delta = \sup(C_\delta), \text{otp}(C_\delta) = \omega$  or at least  $< \delta$ , is there a club  $E$  of  $\omega_1$  such that  $\delta < \omega_1 \Rightarrow \delta > \sup(C_\delta \cap E)$ ? (i.e. is this consistent with CH).

We consider

$$\mathbb{Q}_{\bar{C}}^1 = \{ \bar{f} : \text{for some non-limit } \alpha < \omega_1 \text{ we have } f \in {}^\alpha 2, f^{-1}(\{1\}) \text{ closed} \\ \text{and } \delta < \alpha \text{ limit } \Rightarrow \sup(f^{-1}(\{1\}) \cap C_\delta) < \delta \}.$$

This is the natural forcing for adding a club such that  $\bigwedge_{\delta} [C_\delta \cap E \text{ bounded in } \delta]$ . So

$E$  “runs away” from each  $C_\delta$ . This forcing notion is NOT  $\omega$ -proper: if  $\langle N_i : i \leq \omega \rangle$  satisfies  $C_{N_\omega \cap \omega_1} = \{N_i \cap \omega_1 : i < \omega\}$ , then no  $f \in \mathbb{Q}$  is  $(N_i, \mathbf{G})$ -generic, for infinitely many  $i$ 's.

A solution ([Sh:f, Ch.XVIII,§2]) was to demand “essentially” that, e.g.  $P_i \times P_i$  is proper for  $i < \ell g(\mathbb{Q})$ . While this is fine for  $\mathbb{Q}_i^1$ , this seems to exclude specializing an Aronszajn tree without adding reals. We will deal with a condition implied by both  $(< \omega_1)$ -proper and (essentially) “the square of the forcing notion is proper”.

Continuing explanation:

So for CS iteration  $\mathbb{Q}$  of proper forcing the “reasons”, “dangers” for adding reals may come from:

- $\otimes_0$  (0-reason, danger) some  $\mathbb{Q}_i$  adds reals and
- $\otimes_1$  weak diamond.

Against this, we will assume something like (Definition 0.4): many times in some sense  $q_0, q_1 \in \mathbb{P}_i$  are  $(N, \mathbb{P}_i)$ -generic,  $p \in \mathbb{Q}_i \cap N, q_\ell \Vdash_{\mathbb{P}_i} \text{“}\mathbf{G}_{\mathbb{P}_i} \cap N = \mathbf{G}^*\text{”}$  and for some  $\mathbf{G}', q'_0 \geq q_0, q'_1 \geq q_1$  in  $\mathbb{P}_{i+1}$  we have  $\mathbf{G}' \subseteq (\mathbb{Q}_i \cap N)[\mathbf{G}^*]$  and  $q'_\ell \Vdash_{\mathbb{P}_i} \text{“}\mathbf{G}_{\mathbb{Q}_i} \cap N[\mathbf{G}^*] = \mathbf{G}'\text{”}$  and  $p \in \mathbf{G}'$ . {y9}

It is simpler in the proof to allow  $q_\ell (\ell < n)$  for some  $n < \omega$ ; anyhow in addition we have the obstacle:

- $\otimes_2$  adding almost disjoint clubs ([Sh:f, Ch.XVIII,§1]).

There were two medicines:

- ( $\alpha$ )  $\alpha$ -proper for every  $\alpha < \omega_1$
- ( $\beta$ ) something like  $\mathbb{P}_i \times \mathbb{P}_i$  is proper.

In the proofs we have a situation:

- (\*) (a)  $\bar{Q} \in N_0 \in N$
- (b)  $N_0 \prec (\mathcal{H}(\chi), \in)$  and  $N \prec (\mathcal{H}(\chi), \in)$  are countable
- (c)  $q_\ell$  is  $(N, \mathbb{P}_i)$ -generic and  $(N_0, \mathbb{P}_i)$ -generic
- (d)  $q_\ell$  forces that  $\mathbf{G}_{\mathbb{P}_i} \cap N = \mathbf{G}_\ell$ , (for  $\ell < 2$ )
- (e)  $\mathbf{G}^* = \mathbf{G}_1 \cap N_0 = \mathbf{G}_2 \cap N_0$
- (f)  $i, j, p \in N_0[\mathbf{G}^*], i \leq j \leq \ell g(\bar{Q})$
- (g)  $p \in P_j, p \upharpoonright i \in \mathbf{G}^*$  (possibly more).

We would like to find  $\mathbf{G}' \subseteq \mathbb{P}_j^N / \mathbf{G}^*$  generic over  $N_0$  such that  $q_0$  and  $q_1$  both forces that it has an upper bound in  $P_j / \mathbf{G}_{\mathbb{P}_i}$ . If  $j = i + 1$  this means  $\mathbf{G}' \subseteq \mathbb{Q}_i[\mathbf{G}^*]$  is generic over  $N_0$  such that  $q_0, q_1$  both force that  $\mathbf{G}'$  has an upper bound in  $\mathbb{Q}_i[\mathbf{G}_{\mathbb{P}_i}]$ .

It is natural to demand  $\mathbf{G}' \in N$ , otherwise the two possible generic extensions (for  $q_0$  and  $q_1$ ) become not related. For the case  $j = i + 1$ , a “weak diamond medicine” should help us. But we need it for every  $j$ , naturally we prove it by induction on  $j$ , and the successor case can be reduced to the case  $j = i + 1$ .

But to continue in a limit we need  $\mathbf{G}' \in N$  and more: for some intermediate  $N_1, N_0 \in N_1 \in N$  also  $\bigwedge_{\ell} [q_\ell \Vdash N_1[\mathbf{G}_{\mathbb{P}_i}] \cap \mathbf{V} = N_1]$ . So the clubs of elementary submodels which  $q_0, q_1$  induce on  $\{M \prec N : M \in N\}$  should have non-trivial intersection. This is a major point and it has always appeared in some form. Here the medicine against  $\otimes_2$  should help, in some way there will be many possible  $N_1$ 's; but its help has a price: we have to carry it during the induction. On the other hand the models playing the role of  $N_1$  may change, we may “consume it and discard it”.

Note that the discussion is on two levels. Necessary limitations of universes with CH on the one hand, and how we try to carry the inductive proof on appropriate iterations on the other hand; the connection though is quite tight.

So we shall try for  $j \in \ell g(\bar{Q}) \cap N_0$  to extend the situation with  $i$  being replaced by  $j$  while  $\mathbf{G}^*$  is being increased to  $G^{**}$ . We shall prove by induction suitable facts, with  $\mathbf{G}^{**}$  the object we are really interested in. We are given  $q_1, q_2 \in \mathbb{P}_i$  and would like to find suitable  $q'_1, q'_2 \in \mathbb{P}_j$  such that  $q'_\ell \upharpoonright i = q_\ell$  (otherwise in limit why is there an upper bound?)

So the real action occur for  $j$  limit, hence we choose  $\zeta_n \in N \cap [i, j)$  such that  $\zeta_0 = i, \zeta_n < \zeta_{n+1}$  (sometimes better to have  $i$  and each  $\zeta_n$  non-limit) and  $\bigcup_{n < \omega} \zeta_n = \sup(j \cap N)$ .

You can think of:

in each case of limit  $j$ , proving the inductive statement, we choose a “surrogate” for  $N$  called  $N_1$ , during the induction it serves like  $N$ , in the limit dealing with  $\zeta_0, \zeta_1, \dots$  using the induction hypothesis on  $N_1$  we get  $\mathbf{G}^{**}$  which may not be in  $N_1$  but is in  $N$ .

So we try to choose by induction on  $n, q_{0,n}, q_{1,n}, \mathbf{G}_n^*$  such that:  $q_{\ell,n} \in \mathbb{P}_{\zeta_n}$  is  $(N, \mathbb{P}_{\zeta_n})$ -generic,  $q_{\ell,0} = q_{\ell}, q_{\ell,n+1} \upharpoonright \zeta_n = q_{\ell,n}, \mathbf{G}_n^* \in N_1, \mathbf{G}_n^* \subseteq P_{\zeta_n} \cap N$  is generic over  $N$  and  $q_{\ell,n} \Vdash \mathbf{G}_{\mathbb{P}_{\zeta_n}} \cap N = \mathbf{G}_n^*$ . The construction of the  $\mathbf{G}_n^*$  should use little information on the actual  $q_{\ell,n}$  so that the choices of the  $\mathbf{G}_n^*$  can be carried say inside  $N_1$  so that  $\langle \mathbf{G}_n^* : n < \omega \rangle \in N$ . In fact several models will play a role like  $N_1$ .

By the proof of the preservation of  ${}^\omega\omega$ -bounding we can choose some  $N_1$  and demand “ $q_{\ell,n}$  gives to each  $\mathbb{P}_{\zeta_n}$ -name of an ordinal  $\tau_n \in N_1$ , only finitely many possibilities”.

Now how does  $(< \omega_1)$ -proper help?

We can assume in the beginning that  $\langle N_{1,\gamma} : \gamma \in A \rangle \in N$  is  $\prec$ -increasing continuously,  $N_0 \prec N_{1,\gamma} \prec N, \langle N_{1,\gamma} : \gamma \leq \beta \rangle \in N_{\beta+1}$  with  $A =: (j+1) \cap N \setminus i$  and assume  $q_{\ell}$  is  $(N_{1,\gamma}, \mathbb{P}_i)$ -generic for  $\gamma \in A$  (similarly for  $q'_0, q'_1, j$  in the conclusion) and demand  $q_{\ell,n}$  is  $(N_{1,\gamma}, \mathbb{P}_{\zeta_n})$ -generic for  $n < \omega$  and  $\gamma \in A \setminus \zeta_n$ .

We are ignoring several points including how the induction change and having  $\ell < 2$  rather than  $\ell < n (< \omega)$  which complicates life.

How does “ $\mathbb{Q} \times \mathbb{Q}$  is proper” help?

We demand things like “ $(q_0, q_1)$  is  $(N_1, \mathbb{P}_i \times \mathbb{P}_i)$ -generic” so this gives many common  $N_1$ ’s, but to preserve this we need more complicated situations. Instead of a “tower” of models of countable length, we have a finite tower of models (say of length 5) where on the bottom we are computing  $G^{**} \cap \mathbb{P}_{\zeta_n}$  and as we go up less and less is demanded.

The medicine in the present work is  $\mathfrak{p}$ -properness where “ $\mathbb{Q}$  is  $\mathfrak{p}$ -proper” say that if  $Y$  is a large family of  $M \prec N$  and  $p \in \mathbb{Q} \cap N, \mathbb{Q} \in N$  then for some  $q$  we have  $p \leq q$  and  $q$  is  $(N, \mathbb{Q})$ -generic and  $q \Vdash \{M \in Y : M[\mathbf{G}_{\mathbb{Q}}] \cap \mathbf{V} = M\}$  is large”. (The idea of the finite tower is retained in the proof). This is quite obvious in hindsight.

Why is it important to be inside  $N$ ? Otherwise, we could forget about  $N$  and we have  $q_0, q_1$ . We know they have a common candidate but we need to increase them to know it and in limit by the knowledge that  $P_i$  is bounded.

“We need a real not a name of a real.”

Note that a sufficient condition for  $\mathfrak{p}$ -properness for  $\mathbb{Q}$ , if  $\mathfrak{p}$  is standard, is homogeneity.

{y18}

*Notation* 0.8.  $\mathfrak{p}$  denote a reasonable parameter.

We thank Todd Eisworth for many corrections; he has continued this work.



§ 1. PRESERVATION OF NOT ADDING REALS

On comparing the results of this section with those of [Sh:f], see 1.20.

{preservation}  
{a51}  
{a3}

**Definition 1.1.** We say  $\mathbf{p} = (\bar{\chi}, \bar{R}, \bar{\mathcal{E}}, \bar{D}) = (\bar{\chi}^{\mathbf{p}}, \bar{R}^{\mathbf{p}}, \bar{\mathcal{E}}^{\mathbf{p}}, \bar{D}^{\mathbf{p}})$  is a reasonable parameter when for some ordinal  $\alpha^*$  called  $\ell g(\mathbf{p})$  we have<sup>2</sup>:

- (a)  $\bar{\chi} = \langle \chi_\alpha : \alpha < \alpha^* \rangle, \chi_\alpha$  a regular cardinal,  $\mathcal{H}((\bigcup_{\beta < \alpha} \chi_\beta)^+) \in \mathcal{H}(\chi_\alpha)$   
[this is just technical]
- (b)  $\bar{R} = \langle R_\alpha : \alpha < \alpha^* \rangle, R_\alpha \in \mathcal{H}(\chi_\alpha)$ ;  
[we could have asked “ $R_\alpha$  a relation on  $\mathcal{H}(\chi_\alpha)$ ”, no real difference for our purpose; in a sense it codes a club of  $[\mathcal{H}(\chi_\alpha)]^{\leq \aleph_0}$ .]
- (c)  $\bar{\mathcal{E}} = \langle \mathcal{E}_\alpha : \alpha < \alpha^* \rangle$  where  $\mathcal{E}_\alpha \subseteq [\mathcal{H}(\chi_\alpha)]^{\leq \aleph_0}$  is stationary
- (d)  $\bar{D} = \langle D_\alpha : \alpha < \alpha^* \rangle, D_\alpha$  is a function with domain  $\mathcal{E}_\alpha, a \in \mathcal{E}_\alpha \Rightarrow D_\alpha(a)$  is a pseudo-filter on  $a$ , i.e.  $D_\alpha(a)$  is a family of subsets of  $a$  closed under supersets, non-empty if  $\alpha > 0$   
[and let  $D_\alpha^-(a) = (D_\alpha(a))^- = \mathcal{P}(a) \setminus D_\alpha(a)$ ]
- (e) for  $\alpha < \alpha^*$  we let  $\mathbf{p}^{[\alpha]} =: \langle \bar{\chi} \upharpoonright \alpha, \bar{R} \upharpoonright (\alpha + 1), \bar{\mathcal{E}} \upharpoonright \alpha, \bar{D} \upharpoonright \alpha \rangle$ , so it belongs to  $\mathcal{H}(\chi_\alpha)$ .  
[Why  $\bar{R} \upharpoonright (\alpha + 1)$ ? This makes it an easy demand on  $\mathcal{E}_\alpha : N \in \mathcal{E}_\alpha \Rightarrow R_\alpha \in N$ ]
- (f) if  $a \in \mathcal{E}_\alpha$ , then for some countable  $N \prec (\mathcal{H}(\chi_\alpha), \in)$  we have:  
 $a$  is the universe of  $N$ , so we may write  $D_\alpha(N)$  instead of  $D_\alpha(a)$  and  $N \in \mathcal{E}_\alpha$  instead of  $|N| \in \mathcal{E}_\alpha$
- (g) if  $\alpha < \alpha^*$  and  $N \in \mathcal{E}_\alpha$ , then  $\mathbf{p}^{[\alpha]} \in N$  so  $\alpha \in N$
- (h) for  $N \in \mathcal{E}_\alpha$  and  $X \subseteq N$  we have:  
 $X \in D_\alpha(N)$  iff  $(\bigcup_{\beta < \alpha} \mathcal{E}_\beta) \cap X \cap N \in D_\alpha(a)$
- (i) if  $N \in \mathcal{E}_\alpha, X \in D_\alpha(N), \beta \in \alpha \cap N$  and  $y \in N \cap \mathcal{H}(\chi_\beta^{\mathbf{p}})$ , then for some  $M \in \mathcal{E}_\beta \cap X$  we have  $X \cap M \in D_\beta(M)$  and  $y \in M$
- (j) we may add  $\lambda$  such that  $\alpha < \lambda \Rightarrow |\alpha|^{\aleph_1} < \lambda$  and usually  $\lambda = \lambda^{< \lambda}$  (do? see §6)

{a6}

*Remark 1.2.* 1) Note that  $\langle \mathcal{E}_\alpha : \alpha < \ell g(\mathbf{p}) \rangle$  are pairwise disjoint by clause (g) (and clause (e)) so  $D(N)$  can be well defined as  $D_\alpha(N)$  for the unique  $\alpha$  such that  $N \in \mathcal{E}_\alpha$ .

2) Clearly only  $D_\alpha(N) \cap \mathcal{P}(\bigcup_{\beta < \alpha} \mathcal{E}_\beta)$  matters.

3) Note that the most natural case is “ $D(N)$  is  $\{X \subseteq N : X \neq \emptyset \text{ mod } D^{\text{fil}}\}$ ” for some filter  $D^{\text{fil}}$  on  $N$ .

4) Natural cases are:

- (a)  $D_\alpha(a)$  is a filter on  $a$ , and
- (b)  $D_\alpha(a) = \{b \subseteq a : a \setminus b \notin D'_\alpha(a)\}$  for a filter  $D'_\alpha(a)$  on  $a$ ; we say  $D_\alpha$  is dual to  $D'_\alpha$  or  $D_\alpha = (D'_\alpha)^+$ . Later we shall use them for  $\mathbb{Q}_{\bar{c}}^2$ . [FILL!]

5) We may add in clause (i) of 1.1 that some  $X' \in X \cap M$  belongs to  $N \cap D_\beta(M)$ .  
See no harm but not necessary at present.

{a3}

<sup>2</sup>Clause a gives a nice framework, see 1.8(a).

{a24}

{a9}  
{a3}**Definition 1.3.** In the context of 1.1.1) We say  $\bar{D}$  is standard when for every  $\alpha < \alpha^* (= \ell g(\mathfrak{p}))$  and  $N \in \mathcal{E}_\alpha$  we have

$$D_\alpha(N) = \{X \subseteq N : \begin{array}{l} \text{for every } \gamma \in N \cap \alpha \text{ and} \\ y \in N \cap \cup \{\mathcal{H}(\chi_\beta) : \beta \in \alpha \cap N\} \text{ for some} \\ \beta \in N \cap \alpha \setminus \gamma \text{ and} \\ M \in X \cap \mathcal{E}_\beta \text{ we have } y \in M, \\ X \cap M \in D_\beta(M)\}. \end{array}$$

2) We say  $\mathfrak{p}$  is simple if  $\alpha \leq \beta < \alpha^* \Rightarrow \alpha \leq_{\mathfrak{p}} \beta$ , see below.2A) We define partial order  $\leq_{\mathfrak{p}}$  on  $\alpha^* = \ell g(\mathfrak{p})$  as follows:  $\alpha \leq_{\mathfrak{p}} \beta$  iff

- (a)  $\alpha \leq \beta < \alpha^* = \ell g(\mathfrak{p})$
- (b)  $N \in \mathcal{E}_\beta \wedge \alpha \in N \Rightarrow M =: N \cap \mathcal{H}(\chi_\alpha^{\mathfrak{p}}) \in \mathcal{E}_\alpha^{\mathfrak{p}}$
- (c)  $N \in \mathcal{E}_\beta \wedge \alpha \in N \wedge Y \in D_\beta(N) \Rightarrow Y \cap \bigcup_{\gamma < \alpha} \mathcal{E}_\gamma^{\mathfrak{p}} \in D_\alpha(M)$ .

3) We say  $\mathfrak{p}$  is standard if ( $\mathfrak{p}$  is a reasonable parameter such that)  $\bar{D}^{\mathfrak{p}}$  is standard.4) If  $N \prec (\mathcal{H}(\chi), \in)$  and  $N \cap \mathcal{H}(\chi_\alpha^{\mathfrak{p}}) \in \mathcal{E}_\alpha$ , (hence  $\alpha \in N, \mathfrak{p} \upharpoonright \alpha \in N, R_\alpha^{\mathfrak{p}} \in N$ ), then we let  $D_\alpha(N) = D_\alpha^{\mathfrak{p}}(N)$  be  $D_\alpha^{\mathfrak{p}}(N \cap \mathcal{H}(\chi_\alpha^{\mathfrak{p}}))$ .

{a12}

**Convention 1.4.** If  $\bar{D} = \bar{D}^{\mathfrak{p}}$  is standard, we may omit it. If  $\mathfrak{p}$  clear from the context, we may write  $\mathcal{E}_\alpha$  instead  $\mathcal{E}_\alpha^{\mathfrak{p}}$ .

{a15}

**Definition 1.5.** 1) We say that  $\mathfrak{p}$  is a winner or a  $\mathfrak{D}$ -winner if:for every  $\alpha < \ell g(\mathfrak{p}), \alpha > 0$  and  $N \in \mathcal{E}_\alpha^{\mathfrak{p}}$ , in the game  $\mathfrak{D}_\alpha(N) = \mathfrak{D}_\alpha(N, \mathfrak{p})$  (defined below)

the chooser player has a winning strategy, where:

2) For  $N$  and  $\alpha$  as above,  $\mathfrak{D}_\alpha(N, \mathfrak{p})$  is the following game:a play lasts  $\omega$  moves, in the  $n$ -th movethe challenger choose  $X_n \in D_\alpha(N)$  such that  $m < n \Rightarrow X_m \subseteq X_n$ the chooser chooses  $M_n \in X_n$  and  $Y_n \subseteq M_n \cap X_n$  satisfying  $Y_n \in D(M_n) \cap N$ the challenger chooses  $Z_n \subseteq Y_n$  such that  $Z_n \in D(M_n)$ .In the end the chooser wins if  $\cup\{\{M_n\} \cup Z_n : n < \omega\} \in D_\alpha(N)$ .3) Assume  $N \in N' \prec (\mathcal{H}(\chi), \in)$  and  $\mathfrak{p} \upharpoonright \alpha \in N'$ , and, of course,  $N \prec N'$  are countable. The game  $\mathfrak{D}'_\alpha(N, N', \mathfrak{p})$  is defined similarly to  $\mathfrak{D}_\alpha(N, \mathfrak{p})$  but during the  $n$ -th move, we demand that all the chosen objects belong to  $N'$ , (this means only then  $X_n \in N'$ ) and in the end (of the  $n$ -th move) the chooser also chooses  $X'_n \subseteq X_n, X'_n \in D(N) \cap N'$  and the challenger in the next move has to satisfy  $X_{n+1} \subseteq X'_n$ .3A) Omitting  $N'$ , i.e. writing  $\mathfrak{D}'_\alpha(N, \mathfrak{p})$  we mean: for any such  $N'$  the demand holds.4) We say that  $\mathfrak{p}$  is a non- $\mathfrak{D}$ -loser if for  $\alpha < \ell g(\mathfrak{p}), \alpha > 0$  and  $N \in \mathcal{E}_\alpha$  the challenger has no winning strategy in  $\mathfrak{D}_\alpha(N, \mathfrak{p})$ .5) “ $\mathfrak{D}'_\alpha$ -winner” or “non- $\mathfrak{D}'_\alpha$ -loser” means we (in part (1) or part (4)) use  $\mathfrak{D}'_\alpha(N, \mathfrak{p})$ . We say that “the chooser/challenger wins the game  $\mathfrak{D}_\alpha(N)$ ” if he has a winning

strategy and so “the chooser/challenger does not win the game  $\mathcal{D}_\alpha(N)$ ” says the negation. (Similarly for the other games in this paper).

6) Omitting  $\alpha$  means for every  $\alpha < \ell g(\mathfrak{p})$ .

**Observation 1.6.** 1) If  $\mathfrak{p}$  is a reasonable parameter with the standard  $\bar{D}$ , then  $\mathfrak{p}$  is a winner. {a18}

2) If  $\mathfrak{p}$  is a  $\mathcal{D}_\alpha$ -winner then  $\mathfrak{p}$  is a  $\mathcal{D}'_\alpha$ -winner; if  $\mathfrak{p}$  is a  $\mathcal{D}$ -winner, then  $\mathfrak{p}$  is a  $\mathcal{D}'$ -winner; similarly for a non-loser.

*Proof.* Straightforward.  $\square_{1.6}$

**Definition 1.7.** Assume  $\mathfrak{p}$  is a reasonable parameter,  $N \in \mathcal{E}_\alpha^{\mathfrak{p}}, y \in N$  and  $\mathbb{P} \in N$  is a forcing notion. We let  $\mathcal{M}_{\mathbb{P}}[\mathbf{G}_{\mathbb{P}}, N, y] =: \{M \in N : \mathbb{P}, y \in M \text{ and } \mathbf{G}_{\mathbb{P}} \cap M \text{ is a subset of } \mathbb{P} \cap M \text{ generic over } M\}$ , so this is a  $\mathbb{P}$ -name and  $\mathcal{M}_{\mathbb{P}}[\mathbf{G}, N, y]$  is well defined for any  $\mathbf{G}$ . If  $\mathbb{P}$  is clear from the context, we may omit it; note that  $\mathcal{M}_{\mathbb{P}}[\mathbf{G}, N, y] = \mathcal{M}_{\mathbb{P}}[\mathbf{G} \cap N, N, y]$  so we may write  $\mathbf{G} \cap N$  instead of  $\mathbf{G}$ . If  $y = \emptyset$  we may omit it. {a21}

**Definition 1.8.** We say  $\bar{\mathbb{Q}}$  is a  $\mathfrak{p} - \text{NNR}_{\aleph_0}^0$  iteration when: {a24}

(a)  $\bar{\mathbb{Q}} = \langle \mathbb{P}_i, \mathbb{Q}_i : i < j(*) \rangle$  is a CS iteration of proper forcing notions which belongs to  $\mathcal{H}(\chi_0^{\mathfrak{p}})$ , even  $\mathcal{P}(\mathbb{P}_{j(*)}) \in \mathcal{H}(\chi_0^{\mathfrak{p}})$

(b) forcing with  $\mathbb{P}_{j(*)} = \text{Lim}(\bar{\mathbb{Q}})$  does not add reals

(c) [long properness] if  $(*)_1$  then  $(*)_2$  when

$(*)_1$  (a)  $i \leq j \leq j(*), \alpha < \ell g(\mathfrak{p})$ ,

(b)  $N \in \mathcal{E}_\alpha, \{i, j, \bar{\mathbb{Q}}\} \in N$ ,

(c) the condition  $q \in \mathbb{P}_i$  is  $(N, \mathbb{P}_i)$ -generic

(d)  $q$  forces  $\mathbf{G}_{\mathbb{P}_i} \cap N = \mathbf{G}$

(e)  $p \in \mathbb{P}_j \cap N$  and  $p \upharpoonright i \in \mathbf{G}$

(f)  $Y \subseteq \mathcal{M}_{\mathbb{P}_i}[\mathbf{G}, N, y]$  where  $y = \langle \bar{\mathbb{Q}}, i, j \rangle$  and  $Y \in D_\alpha(N)$  (note that from  $\bar{\mathbb{Q}}, \mathbf{G}$  the ordinal  $i$  is reconstructible);

$(*)_2$  there are  $\mathbf{G}'', q'$  such that:

(g)  $q' \in \mathbb{P}_j, p \leq q'$  and  $q \leq q' \upharpoonright i$

(h)  $q'$  is  $(N, \mathbb{P}_j)$ -generic

(i)  $q'$  forces  $\mathbf{G}_{\mathbb{P}_j} \cap N = \mathbf{G}''$

(j)  $Y \cap \mathcal{M}_{\mathbb{P}_j}[\mathbf{G}'', N, y] \in D_\alpha(N)$

(in §2 this is called  $\mathfrak{p}$ -proper)

(d) [anti-w.d.] if  $\bullet_1$  then  $\bullet_2$  where

$\bullet_1$   $i \leq j \leq j(*), \alpha < \ell g(\mathfrak{p}), N_0 \in N_1 \in \mathcal{E}_\alpha, N_0 \in \bigcup_{\beta < \alpha} \mathcal{E}_\beta, \text{otp}(N_0 \cap [i, j]) <$

$\alpha$  and<sup>3</sup>  $n < \omega$ , for  $\ell < n$  we have  $q_\ell \in \mathbb{P}_i$  is  $(N_1, \mathbb{P}_i)$ -generic and  $q_\ell$  forces  $\mathbf{G}_{\mathbb{P}_i} \cap N_1 = \mathbf{G}^\ell, \bigwedge_{\ell < n} [\mathbf{G}^\ell \cap N_0 = \mathbf{G}^*]$  where  $\mathbf{G}^* \subseteq \mathbb{P}_i \cap N_0$

is generic over  $N_0$  and  $Y =: \bigcap_{\ell < n} \mathcal{M}[\mathbf{G}^\ell, N_1] \in D_\alpha(N_1)$  and

$p \in \mathbb{P}_j \cap N_0, p \upharpoonright i \in \mathbf{G}^*$

<sup>3</sup>so naturally  $\ell g(\mathfrak{p}) = \omega_1$ ; we use the parallel of “ $\aleph_0$ -completeness system” rather than “2-completeness system” of [Sh:f] as things are complicated enough anyhow; see later in 1.14, 1.15 {a32}

- <sub>2</sub> for some  $\mathbf{G}^{**} \subseteq \mathbb{P}_j \cap N_0$  generic over  $N_0$  we have  $p \in \mathbf{G}^{**} \in N_1$  and
 
$$\bigwedge_{\ell < n} \bigvee_{q \in G^\ell} [q_\ell \Vdash \text{“}\mathbf{G}^{**} \text{ has an upper bound in } \mathbb{P}_j/\mathbf{G}_{\mathbb{P}_i}\text{”}].$$

{a25}

{b3g}

*Remark 1.9.* We may like to phrase clause (c) as a condition on each  $\mathbb{Q}_i$ , for this see Definitions 2.6, 2.8, 2.12; this is a slight loss if we deal with the case  $i < j$ ,  $i$  non limit  $\Rightarrow \mathbb{P}_j/\mathbb{P}_i$  is proper. As no need arise here we ignore this.

{a27}

The following is the section's main claim.

**Main Claim 1.10.** *Assume  $\bar{\mathbb{Q}}$  is a CS iteration,  $\bar{\mathbb{Q}} \in \mathcal{H}(\chi_0^{\mathbb{p}})$  and  $\mathcal{P}(\text{Lim}\bar{\mathbb{Q}}) \in \mathcal{H}(\chi_0^{\mathbb{p}})$ ,  $\mathbb{p}$  a reasonable parameter of length  $\omega_1$ ,  $\delta = \text{lg}(\bar{\mathbb{Q}})$  is a limit ordinal and for every  $\alpha < \delta$ ,  $\bar{\mathbb{Q}} \upharpoonright \alpha$  is a  $\mathbb{p}$ -NNR $_{\aleph_0}^0$  iteration and  $\mathbb{p}$  is a  $\mathfrak{D}$ -winner.*

*Then  $\bar{\mathbb{Q}}$  is a  $\mathbb{p}$ -NNR $_{\aleph_0}^0$  iteration.*

{a24}

*Proof.* Note that  $\delta$  here stands for  $j(*)$  in Definition 1.8.

Proof of clause (a) of Definition 1.8:

{a27}

Holds trivially by an assumption of 1.10.

Proof of clause (b) of Definition 1.8:

{a24}

Follows from clause (d) of Definition 1.8 proved below.

We shall prove clause (c) later, clause (d) has a not-so-short proof.

Proof of clause (d):

Let  $i, j, \alpha, N_0, N_1, n, q_0, \dots, q_{n-1}, G^\ell, G^*, p$  be as in the assumptions of clause (d). Let  $\alpha' = \text{otp}(N_0 \cap [i, j])$ , hence  $\alpha' < \omega_1$  so  $\alpha' \in N_1$ . If  $j < j(*) := \delta$  use “ $\bar{\mathbb{Q}} \upharpoonright j$  is a  $\mathbb{p}$ -NNR $_{\aleph_0}^0$  iteration”, so assume  $j = j(*)$ ; if  $i = j$  the conclusion is trivial so assume  $i < j$ . Let  $i_m \in N_0 \cap j$  be such that  $i_0 = i, i_m < i_{m+1}$  and  $\langle i_m : m < \omega \rangle \in N_1$  and  $\bigcup_{m < \omega} i_m = \text{sup}(N_0 \cap j)$ . Choose  $M_\ell$  for  $\ell < 5$  such

that  $y^* =: \{i, j, \alpha, \alpha', \bar{\mathbb{Q}}, N_0, \langle i_m : m < \omega \rangle\} \in M_\ell \in \mathcal{E}_{\alpha'} \cap N_1 \cap \bigcap_{\ell < n} \mathcal{M}[G^\ell, N_1]$

and  $M_0 \in M_1 \in M_2 \in M_3 \in M_4$  and  $\bigcap_{\ell < n} \mathcal{M}[G^\ell, M_0, y^*] \in D_\alpha(M_0)$ ; note that

$N_0 \prec M_\ell \prec N_1$  and  $M_\ell \in N_1$  follows as well as  $\mathbf{G}^\ell \cap M_\ell$  a generic subset of  $\mathbb{P}_i \cap M_\ell$ .

Now for  $\ell < n$  we can choose  $q'_\ell \in \mathbf{G}^\ell \cap M_4$  which forces (for  $\mathbb{P}_{i_0} = \mathbb{P}_i$ ) a value to  $\mathbf{G}_{\mathbb{P}_{i_0}} \cap M_3$  which necessarily is  $\mathbf{G}^\ell \cap M_3$  and necessarily  $q'_\ell \leq q_\ell$  and require, of course,  $q'_\ell$  is  $(M_k, \mathbb{P}_{i_0})$ -generic forcing  $\mathbf{G}_{\mathbb{P}_{i_0}} \cap M_k = \mathbf{G}^\ell \cap M_k$  for  $k = 0, 1, 2, 3$  and  $\mathbf{G}_{\mathbb{P}_{i_0}} \cap N_0 = \mathbf{G}^*$ .

Let  $\langle \mathcal{I}_m^* : m < \omega \rangle \in M_0$  list the maximal antichains of  $\mathbb{P}_j$  that belongs to  $N_0$ . Now we choose by induction on  $m < \omega$ , the objects  $r_m, \mathbf{G}_m^*, p_m, n_m, \mathbf{G}_m^\ell$  (for  $\ell < n_m$ ) and  $y_m$  such that:

- (\*)<sub>1</sub> (a)  $r_m \in \mathbb{P}_{i_m} \cap M_4$
- (b)  $\text{Dom}(r_m) \subseteq [i, i_m)$
- (c)  $r_{m+1} \upharpoonright i_m = r_m$
- (d)  $q'_\ell \cup r_m (\in \mathbb{P}_{i_m})$  is  $(M_k, \mathbb{P}_{i_m})$ -generic for  $k = 0, 1, 2, 3$  and is  $(N_0, \mathbb{P}_{i_m})$ -generic (note that  $q'_\ell, r_m$  have disjoint domains)
- (e) for every predense subset  $\mathcal{I}$  of  $\mathbb{P}_{i_m}$  which belongs to  $M_2$ , for some finite  $\mathcal{J} \subseteq \mathcal{I} \cap M_2$  the set  $\mathcal{I}$  is predense above  $q'_\ell \cup r_m$  for each  $\ell < n$

- (f)  $n_m < \omega$  and for  $\ell < n_m$  we have:  $\mathbf{G}_m^\ell$  is a subset of  $\mathbb{P}_{i_m} \cap M_0$  generic over  $M_0$ ,  $\mathbf{G}_m^\ell \in M_1$
- (g) if  $\ell < n_{m+1}$  then  $\mathbf{G}_{m+1}^\ell \cap \mathbb{P}_{i_m} \in \{\mathbf{G}_m^k : k < n_m\}$
- (h)  $n_0 = n$ ,  $\mathbf{G}_0^\ell = \mathbf{G}^\ell \cap M_0$
- (i)  $q'_\ell \cup r_m \Vdash_{\mathbb{P}_{i_m}} \text{“}\mathbf{G}_{\mathbb{P}_{i_m}} \cap M_0 \in \{\mathbf{G}_m^\ell : \ell < n_m\}$ ”
- (j)  $\mathbf{G}_m^*$  is a subset of  $\mathbb{P}_{i_m} \cap N_0$  generic over  $N_0$
- (k)  $\mathbf{G}_m^* \subseteq \mathbf{G}_m^\ell$  for  $\ell < n_m$
- (l)  $p_m \in \mathbb{P}_j \cap N_0, p_m \upharpoonright i_m \in \mathbf{G}_m^*, p_{m+1} \in \mathcal{I}_m^*, p_0 = p, p_m \leq p_{m+1}$
- (m)  $Y_m =: \bigcap_{\ell < n_m} \mathcal{M}[G_m^\ell, M_0, y^*] \in D_{\alpha'}(M_0)$  where  $y^* =: \{N_0, \langle i_k : k < \omega \rangle, \bar{Q}, i, j\}$ .

Why is this sufficient?

During the construction above we choose inductively members of  $M_4$  and all the parameters used are from  $M_4$ , so if we choose a well ordering  $<^*$  of  $M_4$  and always choose the  $<^*$ -first object the construction is determined. Clearly there is such  $<^* \in N_1$ . Now

- ( $\alpha$ )  $r = \bigcup_m r_m$  (i.e. the unique  $r \in \mathbb{P}_j$  satisfying  $m < \omega \Rightarrow r \upharpoonright i_m = r_m$ ) belongs to  $\mathbb{P}_j$  and to  $N_1$ .

[Why? Recall clause  $(*)_1(c)$ .]

- ( $\beta$ )  $\mathbf{G}^{**} = \{p' \in \mathbb{P}_j \cap N_0 : \bigvee_{m < \omega} [p' \leq p_m]\}$  belongs to  $M_4$  and is a subset of  $\mathbb{P}_j \cap N_0$  generic over  $N_0$ .

[Why? By the choice of  $\langle \mathcal{I}_m^* : m < \omega \rangle$  and clause  $(*)_1(k)$ .]

- ( $\gamma$ )  $q'_\ell \cup r$  is above  $\mathbf{G}^{**}$  (in  $\mathbb{P}_j$ ).

[Why? By the second statement in  $(*)_1(\ell)$ .]

So we are done. □<sub>1.10</sub>

Why can we carry out the construction?

For  $m = 0$  there is no problem. So assume we have it for  $m$  and we shall choose for  $m + 1$ .

Stage A: Choose  $p_{m+1} \in N_0 \cap \mathcal{I}_m^*$  such that  $p_m \leq p_{m+1}$  and  $p_{m+1} \upharpoonright i_m \in \mathbf{G}_m^*$ . No problem.

Stage B: Choose  $\mathbf{G}_{m+1}^* \subseteq \mathbb{P}_{i_{m+1}} \cap N_0$  generic over  $N_0$  such that  $\mathbf{G}_m^* \subseteq \mathbf{G}_{m+1}^* \in M_0, p_{m+1} \upharpoonright i_{m+1} \in \mathbf{G}_{m+1}^*$  and  $\bigwedge_{\ell < n_m} \bigvee_{r \in \mathbf{G}_m^\ell} [r \Vdash_{\mathbb{P}_{i_m}} \text{“}\mathbf{G}_{m+1}^*$  has an upper bound in  $\mathbb{P}_{i_{m+1}}/\mathbf{G}_{\mathbb{P}_{i_m}}]$ .

This is easy by “ $\bar{Q} \upharpoonright i_{m+1}$  is a  $p$ - $NNR_{\aleph_0}^0$  iteration” applied with  $i_m, i_{m+1}, \alpha', p_{m+1} \upharpoonright i_m, \mathbf{G}_m^*, \langle \mathbf{G}_m^\ell : \ell < n_m \rangle, N_0, M_0$  here standing for  $i, j, \alpha, p, \mathbf{G}^*, \langle \mathbf{G}^\ell : \ell < n \rangle, N_0, N_1$  there, (i.e. we use clause (d) of the Definition 1.8); we are using  $\text{otp}(N_0 \cap [i_m, i_{m+1})) < \{\mathbf{a24}\}$   $\text{otp}(N_0 \cap [i_m, j)) = \alpha'$ .

Stage C:

Now  $\mathbf{G}_{\mathbb{P}_{i_m}} \cap M_1$  is a  $\mathbb{P}_{i_m}$ -name of an object from  $\mathbf{V}$  (as  $\mathbb{P}_{i_m}$  is proper not adding reals), so  $\mathcal{I} =: \{p \in \mathbb{P}_{i_m} : p \text{ forces a value to } \mathbf{G}_{\mathbb{P}_{i_m}} \cap M_1 \in \mathbf{V}\}$  is a dense open subset of  $\mathbb{P}_{i_m}$  and  $\mathcal{I} \in M_2$ . By clause  $(*)_1(e)$  in the induction hypothesis there is a finite  $\mathcal{J} \subseteq \mathcal{I} \cap M_2$  such that:  $\ell < n \Rightarrow \mathcal{J}$  is predense above  $q'_\ell \cup r_m$ . Without loss of generality  $\mathcal{J}$  is minimal. Let  $n_{m+1} = |\mathcal{J}|$ .

Let  $\mathcal{J} = \{p_m^\ell : \ell < n_{m+1}\}$ , let  $p_m^\ell \Vdash \mathbf{G}_{\mathbb{P}_{i_m}} \cap M_1 = H_m^\ell$ . So  $H_m^\ell \in M_2$ , and as  $\mathcal{J}$  is minimal,  $H_m^\ell \cap M_0 \in \{\mathbf{G}_m^\ell : \ell < n_m\}$  so let  $H_m^\ell \cap M_0 = \mathbf{G}_m^{h(\ell)}$  where  $h : n_{m+1} \rightarrow n_m$ .

Let  $Y =: \bigcap_{\ell < m_n} \mathcal{M}[\mathbf{G}_m^\ell, M_0, y^*] \in D_{\alpha'}(M_0)$ . Now we choose by induction on  $\ell \leq n_{m+1}$  a condition  $r_m^\ell \in M_1$  such that:

- (\*)<sub>2</sub> (α)  $r_m^\ell \in \mathbb{P}_{i_{m+1}} \cap M_1$  and  $r_m^\ell \restriction i_m \in H_m^\ell$
- (β)  $r_m^\ell$  is  $(M_0, \mathbb{P}_{i_m})$ -generic and force a value to  $\mathbf{G}_{\mathbb{P}_{i_m}} \cap M_0$  called  $\mathbf{G}_{m+1}^\ell$
- (γ)  $r_m^\ell$  is above  $\mathbf{G}_{m+1}^*$  (chosen in the previous stage), moreover above  $p_m^{h(\ell)}$
- (δ)  $Y \cap \bigcap_{k < \ell} \mathcal{M}[\mathbf{G}_{m+1}^k, M_0, y^*] \in D_{\alpha'}(M_0)$ .

{a24} For the induction step, apply clause (c) in Definition 1.8 (as  $\bar{\mathbb{Q}} \restriction i_{m+1}$  is a  $\mathfrak{p}$ - $NNR_{\aleph_0}^0$ -iteration) with  $i_m, i_{m+1}, \alpha', M_0$ , large enough member of  $H_m^\ell, p_m^{h(\ell)}, Y_m^\ell = Y \cap \bigcap_{k < \ell} \mathcal{M}[\mathbf{G}_{m+1}^k, M_0, y^*]$  here standing for  $i, j, \alpha, N, q, p, Y$  there, noting  $Y_m^\ell \in M_1$ .

Stage D:

We can choose  $r_{m+1}$  as required such that  $\{r_m^\ell : \ell < n_{m+1}\}$  is predense over it by [Sh:f, Ch.XVIII,2.6] (can first do it for each  $r_m^\ell$  separately and then put them together<sup>4</sup>).

So we have finished proving clause (d).

Proof of clause (c):

We prove this by induction on  $\alpha$ .

Let  $i, j, \alpha, N, p, q, Y$  be as there. If  $j < j(*)$  we can apply “ $\bar{\mathbb{Q}} \restriction j$  is a  $\mathfrak{p}$ - $NNR_{\aleph_0}^0$  iteration”, so without loss of generality  $j = j(*)$ . If  $i = j$  the statement is trivial so assume  $i < j$ . Choose  $i_n$  for  $n < \omega$  such that  $i_0 = i, i_n \in N \cap j, i_n < i_{n+1}$  and  $\bigcup_{n < \omega} i_n = \sup(j \cap N)$ . Let  $\langle (y_n, \beta_n) : n < \omega \rangle$  list the pairs  $(y, \beta) \in N \times (\alpha \cap N)$  such that  $y \in \mathcal{H}(\chi_\beta^p)$ . Let  $\mathbf{St}$  be a winning strategy for the chooser in the game  $\partial_\alpha(N)$ .

Let  $\langle \mathcal{I}_n : n < \omega \rangle$  list the dense open subsets of  $P_j$  which belongs to  $N$ .

Now we choose by induction on  $n < \omega$ , the objects  $q_n, p_n, M_n, Y_n$  such that:

- (\*)<sub>3</sub> (a)  $q_n \in \mathbb{P}_{i_n}, q_0 = q$
- (b)  $q_n$  is  $(N, \mathbb{P}_{i_n})$ -generic
- (c)  $q_{n+1} \restriction i_n = q_n$
- (d)  $p_n$  is a  $\mathbb{P}_{i_n}$ -name of a member of  $(\mathbb{P}_j / \mathbf{G}_{\mathbb{P}_{i_n}}) \cap N$
- (e)  $p_n$  is forced to belong to  $\mathcal{I}_n$

<sup>4</sup>If we work in free iterations instead CS (see [Sh:f, Ch.IX,§1], which is equivalent in our context, we just use disjunction of the relevant possibilities.

- (f)  $\underline{M}_n$  is a  $\mathbb{P}_{i_n}$ -name of a member of  $\mathcal{E}_{\beta_n} \cap N$
  - (g) if  $\mathbf{G}_j \subseteq \mathbb{P}_j$  is generic over  $\mathbf{V}$ ,  $q_n \in \mathbf{G}_j$ ,  $p_n[\mathbf{G}_j \cap \mathbb{P}_{i_n}] \in \mathbf{G}_j$ ,  
 $M = \underline{M}_n[\mathbf{G}_j]$ , then
    - ( $\alpha$ )  $\mathbf{G}_j \cap M$  is a subset of  $\mathbb{Q} \cap M$  generic over  $M$
    - ( $\beta$ )  $\mathcal{M}[\mathbf{G}_j \cap M, M, y^*] \cap Y \in D_{\beta_n}[M]$
    - ( $\gamma$ )  $p_n[\mathbf{G}_j]$  belongs to  $M$
- $\langle Y_m \cap \mathcal{M}[\mathbf{G}_{i_m}, \underline{M}_m, y^*], \underline{Y}_m, P_{i_m}, \underline{M}_m : m \leq n \rangle$  is forced by  $q_n$  to be an initial segment of a play of the game  $\mathcal{D}_\alpha(N)$  in which the chooser uses the fixed winning strategy **St**.

The proof is straight by the induction hypothesis on  $\beta$  and “ $\bar{\mathbb{Q}} \upharpoonright i_n$  is a  $\mathfrak{p} - NNR_{\aleph_0}^0$ -iteration” remembering that  $\underline{M}_n, \underline{Y}_n$  are  $\mathbb{P}_{i_n}$ -names but of objects in  $\mathbf{V}$ .

Alternatively, see more in the proof in §4 or see below.

{a30}

*Remark 1.11.* 1) We could have used the “adding no reals” and clause (d) in the proof of clause (c) in order to weaken “winner” to “not loser”; also we can use  $\mathcal{D}'_\alpha(N, N', \mathbb{P})$ , see §4. Also the “ $N_0 \in \bigcup_{\beta < \alpha} \mathcal{E}_\beta$ ” can be replaced by “ $N_0 \in \mathcal{E}'_0$ ” with

$\mathcal{E}'_0 \subseteq [\mathcal{H}(\chi_0^{\mathfrak{p}})]^{\aleph_0}$  stationary; also we can put extra restrictions on  $\mathbf{G}^*$  (and  $\mathbf{G}^{**}$ ), e.g.  $\mathcal{M}[\mathbf{G}^*, N_0, y^*]$  large.

2) Of course, the use of  $\langle \chi_\alpha : \alpha < \ell g(\mathfrak{p}) \rangle$  is not really necessary, we could have used subsets of  $\mathcal{P}(\mathbb{P}_{\ell g(\bar{\mathbb{Q}})})$ , (so  $\mathcal{E}_\alpha$  is changed accordingly) as all the properties depend just on  $N_\ell \cap \mathcal{P}(\mathbb{P}_{\ell g(\bar{\mathbb{Q}})})$ . We feel the present way is more transparent.

3) The proof of clause (c) being preserved can be applied to  $\bar{\mathbb{Q}}$  satisfying (a) + (c) of Definition 1.8 (so possibly adding reals), but then we have to replace  $D_\alpha(N)$  by a definition of such pseudo filters with the winning strategy being absolute enough, e.g. for standard  $\bar{D}$ .

{a24}

4) We can also replace  $D_\alpha(a)$  by a partial ordered set  $L_\alpha(a)$  and a function  $v_a : L_\alpha(a) \rightarrow \mathcal{P}(a)$  (i.e. by the pair  $(L_\alpha, v_a)$ ).

5) In the case we adopt remark (3), then remark (1), (non-losing) becomes less clear, as the universe and even the set of reals changes during the proof. We may consider to weaken “the chooser has a winning strategy” in  $\mathcal{D}'_\alpha(N)$  (game depends on  $\bar{D}$ ), e.g. to not losing in a game with finitely many boards, possibly splitting in each move (no real novelty in the proof), but it is not clear how interesting this is. But if  $D_\alpha(N)$  is inductively defined as sums over an ultrafilter (which is preserved), it just seems that not losing is enough.

{a33}

**Definition 1.12.**  $\bar{\mathbb{Q}}$  a CS iteration will be called  $\mathfrak{p}$ -proper if clauses (a) + (c) of Definition 1.8 hold.

{a24}  
{a36}

*Question 1.13.* Is this notion of interest in proper iterations adding reals; in bounding iterations adding reals?

We still like to consider the parallel of having 2-completeness systems. Also we like to demand only non-losing rather than winning in the assumption of 1.10.

{a27}  
{a39}

**Definition 1.14.** Let  $\kappa \in [2, \omega]$ . We say that  $\bar{\mathbb{Q}}$  is a  $\mathfrak{p} - NNR_\kappa^0$ -iteration if: from Definition 1.8 we have:

{a24}

- (a) as there
- (b) as there

(c) [long properness] as there

(d) [ $\kappa$ -anti w.d.] as in (d) there but  $n < 1 + \kappa$  and  $N_0 \in \mathcal{E}_0$ .

{a42}

**Main Claim 1.15.** Assume  $2 \leq \kappa < \aleph_1$ ,  $\bar{\mathbb{Q}}$  is a CS iteration,  $\mathcal{P}(\text{Lim}(\bar{\mathbb{Q}})) \subseteq \mathcal{H}(\chi_0^{\mathfrak{p}})$ ,  $\mathfrak{p}$  a reasonable parameter of length  $\omega_1$  which is a non- $\mathfrak{D}'$ -loser,  $\delta = \text{lg}(\bar{\mathbb{Q}})$  a limit ordinal and  $i < \delta \Rightarrow \bar{\mathbb{Q}} \upharpoonright i$  is a  $\mathfrak{p} - \text{NNR}_\kappa^0$ -iteration. Then  $\bar{\mathbb{Q}}$  is a  $\mathfrak{p} - \text{NNR}_\kappa^0$ -iteration.

{a43}

{a27} *Remark 1.16.* 1) As it is just combining the proof of 1.10 and [Sh:f, Ch.XVIII,§2] (or [Sh:b, Ch.VIII,4.10] - [Sh:f, Ch.VIII,4.10]) we elaborate less.

2) See more in §4.

{a27}

*Proof.* Similar to the proof of 1.10, with some changes (as in [Sh:b, Ch.VIII,§4], [Sh:f, Ch.VIII,§4], [Sh:f, Ch.XVIII,2.10C]). During the proof of clause (d) we add to clauses (a)-(m):

(n)  $n_m$  is a power of 2, say  $2^{n_m^*}$  and so we can rename  $\{\mathbf{G}_m^\ell : \ell < n_m\}$  as  $\{\mathbf{G}_m^\eta : \eta \in {}^{n_m^*}2\}$

(o) ( $\alpha$ )  $M_\eta \in M_1 \cap \mathcal{E}_{j_\eta}$  for  $\eta \in ({}^{n_m^*}2)$  where  $j_0 = \text{otp}([i, j] \cap N_0)$  and if  $\eta = \nu^{\hat{< i >}}$  then  $j_\eta = \text{otp}([i, j] \cap M_\nu)$

( $\beta$ )  $M_{<>} = N_0$

( $\gamma$ )  $M_\eta \in M_{\eta^{\hat{< 0 >}}} \cap M_{\eta^{\hat{< 1 >}}}$

( $\delta$ )  $\eta \triangleleft \nu_1 \in {}^{n_m^*}2 \wedge \eta \triangleleft \nu_2 \in {}^{n_m^*}2 \Rightarrow \mathbf{G}_m^{\nu_1} \cap M_\eta = \mathbf{G}_m^{\nu_2} \cap M_\eta$  so we call it  $K_m^\eta$

( $\varepsilon$ )  $M_{\eta^{\hat{< 0 >}}} = M_{\eta^{\hat{< 1 >}}}$  when  $\eta \in {}^{n_m^*}2$  call it  $N_\eta$

( $\zeta$ )  $N_\eta \in \mathcal{E}_{j_{\eta^{\hat{< \ell >}}}}$  for  $\eta \in ({}^{m_m^*}2)$ ,  $\ell < 2$

( $\eta$ )  $Y_m^\eta = \mathcal{M}[K_m^{\eta^{\hat{< 0 >}}}, N_\eta] \cap \mathcal{M}[K_m^{\eta^{\hat{< 1 >}}}, N_\eta] \in D_{j_{\eta^{\hat{< 0 >}}}}(N_\eta)$ .

{a45}

**Conclusion 1.17.** Let  $\bar{\mathbb{Q}}$  be a CS iteration and  $\mathfrak{p}$  a non- $\mathfrak{D}'$ -loser reasonable parameter,  $2 \leq n(*) \leq \aleph_0$ ,  $\text{lg}(\mathfrak{p}) = \omega_1$ .

Then  $\bar{\mathbb{Q}}$  is a  $\mathfrak{p} - \text{NNR}_{n(*)}^0$ -iteration iff for each  $i < \text{lg}(\bar{\mathbb{Q}})$

(\*) $_i$   $\mathbb{Q}_i$  is a proper forcing and  $\mathbb{P}_i, \mathbb{Q}_i$  satisfies clauses (d) + (c) of the Definition “ $\mathfrak{p} - \text{NNR}_{n(*)}^0$ -iteration” with  $i, i + 1$  here standing for  $i, j$  there.

*Remark 1.18.* Oct. 2012 - was not checked.

{a27}

*Proof.* By induction on  $j = \text{lg}(\bar{\mathbb{Q}})$ . For  $j = 0$  there is nothing, for  $j$  limit use 1.10 (or 1.11) and if  $j$  successor just read the definitions.  $\square_{1.17}$

{a30}

{a27}

We point out here that clause (c) of Definitions 1.8 and 1.10 really follows from earlier properties which play parallel roles.

{a48}

**Claim 1.19.** 1) Assume that  $\mathfrak{p}$  is a reasonable parameter, and is standard and is non-zero  $\alpha < \text{lg}(\mathfrak{p})$ ,  $N \in \mathcal{E}_\alpha^{\mathfrak{p}}$ ,  $Y \in D_\alpha(N)$  and  $\delta \leq \omega_1 \cap N$  a limit ordinal.

Then we can find sequences  $\bar{N} = \langle N_i : i < \delta \rangle, \langle \gamma_i : i < \delta \rangle$  such that:

(a)  $N_i \in N$  (for  $i < \delta$ ) is countable,  $N \cap \alpha \subseteq N_i$  and  $N_i \in Y$

(b)  $N_i \subseteq \bigcup_{\beta \in \alpha \cap N} (\mathcal{H}(\chi_\beta^{\mathfrak{p}}), \in)$ , and  $\beta \in \alpha \cap N_i \Rightarrow N_i \upharpoonright \mathcal{H}(\chi_\beta^{\mathfrak{p}}) \prec (\mathcal{H}(\chi_\beta^{\mathfrak{p}}), \in)$



- (c)  $i < j \Rightarrow N_i \subseteq N_j$
- (d) for  $i$  limit  $N_i = \bigcup_{j < i} N_j$  and  $N \cap \bigcup \{ \mathcal{H}(\chi_\beta^p) : \beta \in \alpha \cap N \} = \bigcup_{j < \delta} N_j$  so we can stipulate  $N_\delta = N$
- (e)  $\beta \in \alpha \cap N \Rightarrow \langle \mathcal{H}(\chi_\beta^p) \cap N_j : j \leq i \rangle \in N_{i+1}$
- (f)  $\gamma_i \in N_i \cap \alpha, N_i \cap \mathcal{H}(\chi_{\gamma_i}^p) \in \mathcal{E}_{\gamma_i} \cap Y$  and  $\bar{\gamma} \upharpoonright (i+1) \in N_{i+1}$
- (g) if  $i \leq \delta$  is a limit ordinal, ( $i = \delta$  and  $\beta \in \alpha \cap N_i$ )  $\vee$  ( $i < \delta$  and  $\beta \in \gamma_i \cap N_i$ ) (so  $\beta \in \mathcal{H}(\chi_\beta^p)$ ) and  $y \in \mathcal{H}(\chi_\beta^p)$ , then for some  $j < i, \gamma_j = \beta, y \in N_j$ .

Moreover

- (g)<sup>+</sup> if  $i \leq \delta$  is a limit ordinal, then  $\{N_j \cap \mathcal{H}(\chi_{\gamma_j}^p) : j < i \text{ and } \gamma_j < \gamma_i\} \in D_{\gamma_i}(N) = D_{\gamma_i}(N \cap \mathcal{H}(\chi_{\gamma_i}^p))$
- (h) if  $\delta < N \cap \omega_1$  then  $\delta \in N_0$ , if  $\delta = N \cap \omega_1$  then  $i < \delta \Rightarrow i \in N_i$ .

2) If  $\mathfrak{p}$  is a standard, reasonable parameter,  $\bar{\mathbb{Q}}$  is a CS iteration,  $lg(\bar{\mathbb{Q}}) = \alpha + 1, \bar{\mathbb{Q}} \upharpoonright \alpha$  is  $\mathfrak{p} - \text{NNR}_{k(*)}^0$ -iteration and  $\Vdash_{\mathbb{P}_\alpha}$  “ $\bar{\mathbb{Q}}_\alpha$  is proper and  $(<^+ \omega_1)$ -proper”, then trying to apply 1.11 for  $\bar{\mathbb{Q}}$ , clauses (a), (c) (called  $\mathfrak{p}$ -proper in Definition ?? below) holds. {a30}

3) If  $lg(\mathfrak{p}) = \omega_1$ , then in part (2) it suffices to ask  $\Vdash_{\mathbb{P}_i}$  “ $\alpha < \omega_1 \Rightarrow \mathbb{Q}_i$  is  $\alpha$ -proper”, that is  $\Vdash_{\mathbb{P}_i}$  “ $\mathbb{Q}_i$  is  $(< \omega_1)$ -proper”.

*Proof.* 1) By induction on  $\beta \in (\alpha \cap N) \cup \{\alpha\}$  we prove that there is  $\langle N_j : j < \beta \cap N \rangle \in N$  satisfying the relevant requirements.

2) By Definition of  $(< \omega_1)$ -proper, if  $\mathbf{G}_\alpha \subseteq \mathbb{P}_\alpha$  is generic over  $\alpha, Y =: \mathcal{M}[G_\alpha, N, y^*] \in D_\alpha(N)$ , let  $\langle N_i : i < \delta \rangle$  be as in (1) for  $\delta = N \cap \omega_1$ , without loss of generality  $p \in N_0 \cap \mathbb{Q}_\alpha[\mathbf{G}_\alpha]$ , let  $q \geq p$  be  $(N_i, \mathbb{Q}_\alpha[\mathbf{G}_\alpha])$ -generic for every  $i < \delta$  (formally look at  $\bar{N}' = \langle N'_i : i < \delta \rangle, N'_i = N_i \upharpoonright \mathcal{H}(\chi_0^p)$  and apply to it the  $(< \omega_1)$ -proper).

3) Follows as  $\alpha \in N \Rightarrow \delta = \omega\alpha \in N \cap \omega_1$ . □<sub>1.19</sub>

**Discussion 1.20.** 1) This includes as special cases [Sh:b, Ch.V,§5,§7]. There is no direct comparison with [Sh:b, Ch.VIII,§4], [Sh:f, Ch.VIII,§4], but we can make the notion somewhat more complicated, to include the theorem there in our context, i.e., what is not included is a generalization there not really needed for the examples discussed there (see here in §3 below). The condition there involves having many sequences  $\langle N_\alpha : \alpha \leq \delta \rangle$  such that if  $p_0, p_1 \in P, p_\ell$  is  $(N_i, \mathbb{P}_\alpha)$ -generic for  $i, p_\ell \Vdash \mathbf{G}_\mathbb{P} \cap N_0 = \mathbf{G}^*$ , then there is  $\mathbf{G}' \subseteq \text{Gen}(\mathbb{Q}_i[\mathbf{G}^*], N_0[\mathbf{G}^*]), \mathbf{G}' \in N_0, \mathbb{P}_i \not\Vdash_{\mathbb{P}_i} \mathbf{G}_0$  has no bound in  $\mathbf{Q}_\alpha$ . This speaks on a family of sequences from  $[\mathcal{H}(\chi)]^{\aleph_0}$  rather than members. {a51}

2) For [Sh:f, Ch.XVIII,§2], the comparison is not so easy. Our problem is to “carry” good  $(N, \mathbb{P}_i, \langle \mathbf{G}_\ell : \ell < n \rangle), \mathbf{G}_\ell \in \text{Gen}(N, \mathbb{P}_i)$  with a bound, such that we can “increase  $i$ ” and we can find  $N', y \in N' \in N, N' \prec N$  such that  $(N', \mathbb{P}_i, \langle \mathbf{G}_i \cap N : \ell < n \rangle)$  is good enough. In [Sh:f, Ch.XVIII] we are carrying genericity in some  $\mathbb{P}_{\bar{\alpha}}, \bar{\alpha} \in \text{trind}(i)$ , here much less. But what we need is the implication “if  $(N, \mathbb{P}_i, \bar{G})$  is good we can extend it to good  $(N, \mathbb{P}_{i+1}, \bar{G}')$ ”, so making good weaker generates an incomparable notion and clearly there are other variants. We can consider other such notions (see §5).

2A) We can give alternative proofs of consistency of the questions in [Sh:f, Ch.XVIII,§1] by the present iteration theorem (see §3 below).

3) We can in 1.13(2) weaken the assumption “ $(< \omega_1)$ -proper” to things of the form: if  $\bar{N} = \langle N_i \leq \delta \rangle, p \in \mathbb{Q} \cap N_0, \mathbb{Q} \in N_0$  then there is  $q \geq p$  such that {a36}

$q \Vdash$  “for many  $i < \delta$ ,  $\mathfrak{G}_Q \cap N_i \in \text{Gen}(N_i, \mathbb{Q})$ ”. In particular the condition applies to the forcing notions considered in [Sh:f, Ch.XVIII,§1].

## § 2. DELAYED PROPERNESS

{delayed}

In this section we prove little, but the notions introduced are used in §3, §4. We concentrate here on simple parameters so the reader may assume it all the time. We give two versions, the simpler one is version 2 for which simplicity is a very natural demand.

{b3}

**Observation 2.1.** 1) Assume

- (a)  $\bar{\chi} = \langle \chi_\alpha : \alpha < \alpha^* \rangle$  increases fast enough, so  $\bigcup_{\beta < \alpha} \mathcal{H}(\chi_\beta) \in \mathcal{H}(\chi_\alpha)$
- (b)  $\mathcal{E}_\alpha \subseteq \{N : N \text{ a countable elementary submodel of } (\mathcal{H}(\chi_\alpha), \in)\}$  is stationary
- (c)  $R_\alpha \in \mathcal{H}(\chi_\alpha)$  and  $N \in \mathcal{E}_\alpha$  implies  $\langle \chi_\beta : \beta < \alpha \rangle \in N, \langle R_\beta : \beta \leq \alpha \rangle \in N$  and  $\langle \mathcal{E}_\beta : \beta < \alpha \rangle \in N$ .

Then there is a standard reasonable parameter  $\mathfrak{p}$  with  $\ell g(\mathfrak{p}) = \alpha^*$ ,  $\mathcal{E}_\alpha^{\mathfrak{p}} = \mathcal{E}_\alpha$ ,  $R_\alpha^{\mathfrak{p}} = R_\alpha$  (and  $\bar{D}^{\mathfrak{p}}$  standard).

2) If in addition clause (d) below holds, then  $\mathfrak{p}$  is a simple standard reasonable parameter (recall Definition 1.3(2)) where

{a9}

- (d)  $\beta \in N \in \mathcal{E}_\alpha, \beta < \alpha \Rightarrow N \cap \mathcal{H}(\chi_\beta) \in \mathcal{E}_\beta$ .

3) If  $\chi_\alpha = (\beth_{2\alpha+1})^+$  for  $\alpha < \alpha^*$ ,  $R_\alpha \in \mathcal{H}(\chi_\alpha)$ , then  $\chi_\alpha$  increases fast enough. If  $\langle \chi_\alpha : \alpha < \alpha^* \rangle, \langle R_\alpha : \alpha < \alpha^* \rangle$  are as in part (1),  $\chi \leq \chi_0, \mathcal{E} \subseteq [\mathcal{H}(\chi)]^{\leq \aleph_0}$  stationary and we let  $\mathcal{E}_\alpha = \{N : N \text{ a countable elementary submodel of } (\mathcal{H}(\chi_\alpha), \in)$  and  $\langle \chi_\beta : \beta < \alpha \rangle, \langle R_\beta : \beta \leq \alpha \rangle, \mathcal{E}$  belong to  $N$  and  $N \cap \mathcal{H}(\chi) \in \mathcal{E}\}$ , then the assumptions of parts (1) and (2) above holds (hence their conclusions).

*Proof.* Straightforward.

□<sub>2.1</sub>

{b6}

**Definition 2.2.** Let  $\mathfrak{p}$  be a reasonable parameter and  $\alpha \leq \beta < \ell g(\mathfrak{p})$ .

- 1) For  $N \in \mathcal{E}_\beta$  such that  $\alpha \in N$  we define a game  $\mathfrak{D}_{\alpha,\beta}(N) = \mathfrak{D}_{\alpha,\beta}(N, \mathfrak{p})$  as follows. A play lasts  $\omega$ -moves. In the  $n$ -th move:

- (a) the challenger chooses  $X_n \in D_\beta(N)$  such that  $m < n \Rightarrow X_n \subseteq X_m$
- (b) the chooser chooses  $\alpha_n \in \alpha \cap N$
- (c) the challenger chooses  $\beta'_n \in \beta \cap N$  and  $y'_n \in N \cap \mathcal{H}(\chi_{\alpha_n}^{\mathfrak{p}})$
- (d) the chooser chooses  $\beta_n \in \beta \cap N \setminus \beta'_n$  and  $M_n \in X_n \cap \mathcal{E}_{\beta_n}$  and  $y_n \in M_n \cap \mathcal{H}(\chi_{\alpha_n}^{\mathfrak{p}})$  satisfying  $\alpha_n \leq \beta_n, y'_n \in M_n, \alpha_n \in M_n$  and  $Y_n \in D_{\beta_n}(M_n)$  such that  $Y_n \subseteq X_n$  and  $Y_n \in N$
- (e) the challenger chooses  $M'_n \in Y_n \cap \mathcal{E}_{\alpha_n} \cap (M_n \cup \{M_n \cap \mathcal{H}(\chi_{\alpha_n}^{\mathfrak{p}})\})$  satisfying  $y_n, y'_n \in M'_n$  and chooses  $Z_n \in D_{\alpha_n}(M'_n) = D_{\alpha_n}(M'_n \cap \mathcal{H}(\chi_{\alpha_n}^{\mathfrak{p}}))$  such that  $Z_n \subseteq Y_n$ .

In the end the chooser wins the play if

$$\bigcup \{Z_n \cup \{M'_n\} : n < \omega\} \in D_\alpha^+(N) (= D_\alpha^+(N \cap \mathcal{H}(\chi_\alpha^{\mathfrak{p}})))$$

1A) We call  $\mathfrak{D}_{\alpha,\beta}(N) = \mathfrak{D}_{\alpha,\beta}(N, \mathfrak{p})$  version 1; version 2 means that in clause (e) we add the requirement  $M'_n = M_n \cap \mathcal{H}(\chi_{\alpha_n}^{\mathfrak{p}})$  and in clause (d) we require  $\alpha_n \leq_{\mathfrak{p}} \beta_n$ . If we do not mention the version it means that it holds for both.

2) Assume  $N \in N' \prec (\mathcal{H}(\chi), \in)$ . We define a game  $\mathcal{D}'_{\alpha, \beta}(N, N', \mathbf{p})$  similarly but replace a) - e) by

- (a)' the challenger chooses  $X_n \in D_\beta(N)$  such that  $m < n \Rightarrow X_n \subseteq X'_m$
- (b)' the chooser chooses  $\alpha_n \in \alpha \cap N$  and  $X'_n \subseteq X_n$  such that  $X'_n \in D_\beta(N)$
- (c)' like (c)
- (d)' like (d) but we replace “ $Y_n \in N$ ” by “ $Y_n \in N'$ ”
- (e)' like (e) but add  $Z_n \in N'$ .

Note: so every proper initial segment of a play belongs to  $N'$ .

{b9}

**Definition 2.3.** Let  $\mathbf{p}$  be a reasonable parameter.

1) For  $\alpha \leq \beta < \ell g(\mathbf{p})$  we say  $\mathbf{p}$  is an  $\mathcal{D}_{\alpha, \beta}$ -winner [non- $\mathcal{D}_{\alpha, \beta}$ -loser] when for some  $x \in \mathcal{H}(\chi_\beta^{\mathbf{p}})$  we have:

if  $\{x, \alpha\} \in N \in \mathcal{E}_\beta$ , then the chooser wins the game  $\mathcal{D}_{\alpha, \beta}(N, \mathbf{p})$  [the challenger does not win in the game  $\mathcal{D}_{\alpha, \beta}(N, \mathbf{p})$ ].

2) We can replace  $\mathcal{D}_{\alpha, \beta}$  by  $\mathcal{D}'_{\alpha, \beta}$ .

3) For any function  $f : \ell g(\mathbf{p}) \rightarrow \mathcal{P}(\ell g(\mathbf{p}))$  we can replace  $\alpha, \beta$  by  $f$  meaning: for every  $\alpha < \ell g(\mathbf{p})$  and  $\beta \in f(\alpha)$ , we have  $\mathbf{p}$  is a  $\mathcal{D}_{\alpha, \beta}$ -winner.

4) Let  $\mathcal{F}^{\mathbf{p}}$  be the family of functions from  $\ell g(\mathbf{p})$  to  $\mathcal{P}(\ell g(\mathbf{p}))$  such that for each  $\alpha < \ell g(\mathbf{p})$ ,  $f(\alpha)$  is a nonempty subset of  $[\alpha, \ell g(\mathbf{p}))$ . Let  $\mathcal{F}_{\text{club}}^{\mathbf{p}}$  be the set of  $f \in \mathcal{F}^{\mathbf{p}}$  such that for each  $\alpha < \ell g(\mathbf{p})$ ,  $f(\alpha)$  is a club of  $\ell g(\mathbf{p})$ . Let  $\mathcal{F}_{\text{club}}^{\mathbf{p}}$  be the set of  $f \in \mathcal{F}^{\mathbf{p}}$  such that for each  $\alpha < \ell g(\mathbf{p})$ ,  $f(\alpha)$  is an end segment of  $\ell g(\mathbf{p})$ , we then may identify  $f(\alpha)$  with  $\text{Min}(f(\alpha))$ .

5) Let  $F_{\text{ic}}^{\mathbf{p}}$  be the set of increasing continuous  $f \in \mathcal{F}_{\text{club}}^{\mathbf{p}}$ , i.e.

- (a)  $\alpha < \beta < \ell g(\mathbf{p}) \Rightarrow f(\alpha) \supseteq f(\beta)$
- (b) for limit  $\delta < \ell g(\mathbf{p})$  we have  $f(\delta) = \bigcap \{f(\alpha) : \alpha < \delta\}$ .

6) We call  $f$  decreasing continuous if  $\alpha_1 < \alpha_2 < \ell g(\mathbf{p}) \Rightarrow f(\alpha_2) \subseteq f(\alpha_1)$  and for limit  $\alpha < \ell g(\mathbf{p})$  we have  $f(\alpha) = \bigcap_{\gamma < \alpha} f(\gamma)$ . Let  $f \leq g$  mean  $(\forall \alpha < \ell g(\mathbf{p})) (g(\alpha) \subseteq f(\alpha))$ .

There are obvious monotonicity properties.

{b12}

**Claim 2.4.** Assume  $\mathbf{p}$  is a reasonable parameter.

1) If  $\alpha \leq_{\mathbf{p}} \alpha', \alpha' \leq \alpha \leq \beta = \beta' < \ell g(\mathbf{p})$ , and  $\mathbf{p}$  is a  $\mathcal{D}_{\alpha, \beta}$ -winner, then it is  $\mathcal{D}_{\alpha', \beta'}$ -winner. Similarly for  $\mathcal{D}'$ -winner, non- $\mathcal{D}$ -loser, non- $\mathcal{D}'$ -loser.

2) If  $\mathbf{p}$  is a  $\mathcal{D}_{\alpha, \beta}$ -winner, then  $\mathbf{p}$  is a  $\mathcal{D}'_{\alpha, \beta}$ -winner and a non- $\mathcal{D}_{\alpha, \beta}$ -loser. If  $\mathbf{p}$  is a  $\mathcal{D}'_{\alpha, \beta}$ -winner or non- $\mathcal{D}_{\alpha, \beta}$ -loser, then  $\mathbf{p}$  is non- $\mathcal{D}'_{\alpha, \beta}$ -loser.

3) Assume  $f, g \in \mathcal{F}^{\mathbf{p}}$  and  $f \leq g$ . If  $\mathbf{p}$  is a  $\mathcal{D}_f$ -winner [or  $\mathcal{D}'_f$ -winner] [or non- $\mathcal{D}_f$ -loser] [or non- $\mathcal{D}'_f$ -loser], then  $\mathbf{p}$  is a  $\mathcal{D}_g$ -winner [or  $\mathcal{D}'_g$ -winner] [or non- $\mathcal{D}_g$ -loser] [or non- $\mathcal{D}'_g$ -loser].

*Proof.* Straightforward (we are using the simplicity of  $\mathbf{p}$ ). □<sub>2.4</sub>

{b15}

**Claim 2.5.** 1) Assume  $\mathbf{p}$  is a standard reasonable parameter. Then  $\mathbf{p}$  is a winner (see Definition 1.4).

{a12}

2) If  $\mathbf{p}$  is a reasonable parameter and it is a winner, then  $\mathbf{p}$  is a  $\mathcal{D}_{\alpha, \alpha}$ -winner.

- {b9} 3) If  $\mathfrak{p}$  is a reasonable parameter and it is a winner and  $\alpha \leq \beta < \ell g(\mathfrak{p})$  then  $\mathfrak{p}$  is a  $\mathcal{D}_{\alpha,\beta}$ -winner (hence  $\mathcal{D}_f$ -winner for  $f$  as in 2.3(3)).  
 4) Similarly with  $\mathcal{D}'$ -winner,  $\mathcal{D}'_{\alpha,\beta}$  winner and/or with the “non-loser” cases.

*Proof.* Straight. □<sub>2.5</sub>

**Definition 2.6.** 1) Let  $\mathfrak{p}$  be a reasonable parameter and  $\mathbb{P}$  be a proper forcing notion not adding reals,  $\mathcal{P}(\mathbb{P}) \in \mathcal{H}(\chi_0^{\mathfrak{p}})$  and  $\mathbf{G}_{\mathbb{P}} \subseteq \mathbb{P}$  is generic over  $\mathbf{V}$ .

Then we interpret  $\mathfrak{p}$  in  $\mathbf{V}^{\mathbb{P}}$  as  $\mathfrak{p}' = \mathfrak{p}^{\mathbf{V}[\mathbf{G}_{\mathbb{P}}]}$ , or we may write  $\mathfrak{p}^{\mathbb{P}}$ , defined as follows:

- (a)  $\chi_{\alpha}^{\mathfrak{p}'} = \chi_{\alpha}^{\mathfrak{p}}$
- (b)  $R_{\alpha}^{\mathfrak{p}'} = \langle R_{\alpha}, P, \mathbf{G}_P \rangle$
- (c)  $\mathcal{E}_{\alpha}^{\mathfrak{p}'} = \{N[\mathbf{G}_{\mathbb{P}}] : N \in \mathcal{E}_{\alpha} \text{ and } \mathbb{P} \in N \text{ and } N[\mathbf{G}_{\mathbb{P}}] \cap V = N\}$
- (d)  $D_{\alpha}(N[\mathbf{G}_{\mathbb{P}}]) = \{\{M[\mathbf{G}_{\mathbb{P}}] \in \mathcal{E}_{\alpha} : M \in Y \cap \bigcup_{\beta < \alpha} \mathcal{E}_{\beta}^{\mathfrak{p}}\} : Y \in D_{\alpha}(N)\}$ .

**Claim 2.7.** Let  $\mathfrak{p}, \mathbb{P}$  be as in Definition 1.8. {b21}

- 1) In Definition 2.6,  $\mathfrak{p}^{\mathbf{V}[\mathbf{G}_{\mathbb{P}}]}$  is a reasonable parameter in  $\mathbf{V}[\mathbf{G}_{\mathbb{P}}]$ . {a24}
- 2) If  $\mathfrak{p}$  is, in  $\mathbf{V}$ , a  $\mathcal{D}$ -winner (or non- $\mathcal{D}$ -loser or  $\mathcal{D}'$ -winner or non- $\mathcal{D}'$ -loser), then  $\mathfrak{p}^{\mathbf{V}[\mathbf{G}_{\mathbb{P}}]}$  is so (in  $\mathbf{V}[\mathbf{G}_{\mathbb{P}}]$ ). {b18}
- 3) If  $\mathfrak{p}$  is, in  $\mathbf{V}$ , a  $\mathcal{D}_{\alpha,\beta}$ -winner (or a non- $\mathcal{D}_{\alpha,\beta}$ -lower or  $\mathcal{D}'_{\alpha,\beta}$ -winner or  $\mathcal{D}'_{\alpha,\beta}$ -non loser), then  $\mathfrak{p}^{\mathbf{V}[\mathbf{G}_{\mathbb{P}}]}$  is so in  $\mathbf{V}[\mathbf{G}_{\mathbb{P}}]$ .

*Proof.* Straightforward (we use  $\mathbb{P}$  is a proper forcing notion not adding reals). □<sub>2.7</sub>

\* \* \*

**Definition 2.8.** Let  $\mathfrak{p}$  be a reasonable parameter. {b24}

1) For  $\alpha \leq \beta < \ell g(\mathfrak{p})$ , we say a forcing notion  $\mathbb{Q}$  is  $(\mathfrak{p}, \alpha, \beta)$ -proper when: ( $\mathcal{P}(\mathbb{Q}) \in \mathcal{H}(\chi_0^{\mathfrak{p}})$  and):

- (\*) for some  $x \in \mathcal{H}(\chi_{\beta})$  if  $N \in \mathcal{E}_{\beta}$ ,  $\{x, \mathbb{Q}, \alpha\} \in N$ ,  $p \in N \cap \mathbb{Q}$  and  $Y \in D_{\alpha}(N)$ , then for some  $q$  we have:
  - (a)  $p \leq q \in \mathbb{Q}$
  - (b)  $q$  is  $(N, \mathbb{Q})$ -generic
  - (c) version (1): for some  $N' \in (\mathcal{E}_{\alpha} \cap N \cap Y) \cup \{N \cap \mathcal{H}(\chi_{\alpha}^{\mathfrak{p}})\}$  satisfying  $\alpha = \beta \Rightarrow N' = N$  we have  $q \Vdash_{\mathbb{Q}} \text{“}\mathcal{M}[\mathbf{G}_{\mathbb{Q}}, N', y^*] \cap Y \in D_{\alpha}(N')\text{”}$  where  $y^* = \langle x, p, \mathbb{Q} \rangle$  hence necessarily  $q$  is  $(N', \mathbb{Q})$ -generic.
- (Version 2: similarly but  $N' = N \upharpoonright \mathcal{H}(\chi_{\alpha}^{\mathfrak{p}})$  and naturally we demand  $\alpha \leq_{\mathfrak{p}} \beta$ ).

2) We say  $\mathbb{Q}$  is  $(\mathfrak{p}, f)$ -proper when  $(\mathfrak{p}, f)$  are as above and) for every  $\alpha < \ell g(\mathfrak{p})$  and  $\beta \in f(\alpha)$  the forcing notion  $\mathbb{Q}$  is  $(\mathfrak{p}, \alpha, \beta)$ -proper.

3) We say  $\mathbb{Q}$  is  $\mathfrak{p}$ -proper when  $\mathbb{Q}$  is  $(\mathfrak{p}, \alpha, \alpha)$ -proper for  $\alpha < \ell g(\mathfrak{p})$ . We say  $\mathbb{Q}$  is almost  $\mathfrak{p}$ -proper if  $\mathbb{Q}$  is  $(\mathfrak{p}, f)$ -proper for some  $f \in \mathcal{F}^{\mathfrak{p}}$  (see Definition 2.3(4)). {b9}

{b27}

**Claim 2.9.** Assume  $\mathfrak{p}$  is a simple reasonable parameter.

- 1) If  $\alpha' \leq \alpha \leq \beta \leq \beta' < \text{lg}(\mathfrak{p})$  (for version 2 we demand  $\alpha' \leq_{\mathfrak{p}} \beta'$  and  $\alpha \leq_{\mathfrak{p}} \beta$ ) and  $\mathbb{Q}$  is as  $(\mathfrak{p}, \alpha, \beta)$ -proper forcing notion, then  $\mathbb{Q}$  is a  $(\mathfrak{p}, \alpha', \beta')$ -proper forcing notion.
- 2) Assume  $f, f'$  are in  $\mathcal{F}^{\mathfrak{p}}$  and  $f \leq f'$ . If  $\mathbb{Q}$  is a  $(\mathfrak{p}, f)$ -proper forcing notion, then  $\mathbb{Q}$  is a  $(\mathfrak{p}, f')$ -proper forcing notion.

*Proof.* Straight.  $\square_{2.9}$

{b30}

**Discussion 2.10.** We may like to consider  $(\mathfrak{p}, f)$ -proper for iterations which may add reals. Then we have to replace  $D_{\alpha}(N)$  by a definition which is absolute enough, (and the non-loser versions have to be absolute enough).

It is natural to restrict ourselves to  $\mathfrak{p}$ -closed  $Y$ , see below.

{b33}

**Definition 2.11.** Let  $\mathfrak{p}$  be a simple reasonable parameter and  $N \in \mathcal{E}_{\alpha}$ . Now  $Y \subseteq N$  is called  $\mathfrak{p}$ -closed if (see 3.11):

{c27}

- (a)  $Y \subseteq N \cap \bigcup_{\beta < \alpha} \mathcal{E}_{\beta}$
- (b) if  $M \in N \cap \mathcal{E}_{\beta}$  and  $\beta < \alpha$  (hence  $\beta \in \alpha \cap M \subseteq \alpha \cap N$ ) and  $\gamma \in M \cap \beta$  and  $M \cap \mathcal{H}(\chi_{\gamma}^{\mathfrak{p}}) \in \mathcal{E}_{\gamma}$  (this requirement is redundant if  $\mathfrak{p}$  is simple or just  $\gamma \leq_{\mathfrak{p}} \beta$ ), then  $M \cap \mathcal{H}(\chi_{\gamma}^{\mathfrak{p}}) \in Y \Leftrightarrow M \in Y$
- (c) if  $\beta < \alpha, M_{\ell} \in N \cap \mathcal{E}_{\beta}$  (hence  $\beta \in \alpha \cap N$ ) and  $M_{\ell} \subseteq M_{\ell+1}$  for  $\ell < \omega$  and  $M = \bigcup_{\ell < \omega} M_{\ell} \in N \cap \mathcal{E}_{\beta}$ , and even  $\langle M_{\ell} : \ell < \omega \rangle \in N$ , then  $\bigwedge_{\ell} M_{\ell} \in Y \Rightarrow M \in Y$ .

{b37}

**Definition 2.12.** Assume  $\mathfrak{p}$  is a reasonable parameter,  $\mathbb{P}$  a forcing notion and  $\mathbb{Q}$  a  $\mathbb{P}$ -name of a forcing notion. We say  $\mathbb{Q}$  has  $(\kappa, \alpha, \beta)$ -anti w.d. above  $\mathbb{P}$  (or  $(\mathbb{P}, \mathbb{Q})$  has  $(\kappa, \alpha, \beta)$ -anti-w.d.  $(\kappa, f)$  means  $(\kappa, \alpha, \beta)$  when  $f \in \mathcal{F}^{\mathfrak{p}}, \beta \in f(\alpha)$ ) when  $\alpha \leq \beta < \text{lg}(\mathfrak{p})$  and (similarly to (d) of 1.8), clause (A) implies clause (B) when:

{a24}

- (A) (a)  $N_0 \in \mathcal{E}_{\alpha}$  and  $N_1 \in \mathcal{E}_{\beta}$
- (b)  $N_0 \in N_1$  and  $\{\mathbb{P}, \mathbb{Q}\} \in N_0$
- (c)  $n < 1 + \kappa$
- (d)  $q_{\ell} \in \mathbb{P}$  is  $(N_{\ell}, \mathbb{P})$ -generic for  $\ell < n$  and  $\iota = 0, 1$
- (e)  $q_{\ell} \Vdash \text{“}\mathbf{G} \cap N_1 = \mathbf{G}^{\ell}\text{”}$  for  $\ell < n$
- (f)  $\mathbf{G}^{\ell} \cap N_0 = \mathbf{G}^*$  for  $\ell < n$
- (g)  $Y = \bigcap_{\ell < n} \mathcal{M}[\mathbf{G}^{\ell}, N_1]$  belongs to  $D_{\beta}(N_1)$
- (h)  $\underline{q}$  is a  $\mathbb{P}$ -name of a member of  $\mathbb{Q}$
- (i)  $\underline{q} \in N_0$
- (B) there is a triple  $(p', q', \mathbf{G}^{**})$  such that:
  - (a)  $q'$  is a  $\mathbb{P}$ -name of a member of  $\mathbb{Q}$
  - (b)  $p'_{\ell}^{\xi} \Vdash \text{“}q \leq q'\text{”}$  for  $\ell < n$
  - (c)  $\mathbf{G}^{**} \in \text{Gen}(\mathbb{P} * \mathbb{Q}), N$  over  $N$
  - (d)  $(p'_{\ell}, q') \Vdash \text{“}\mathbf{G}_{\mathbb{P} * \mathbb{Q}} \cap N = \mathbf{G}^{**}\text{”}$  for  $\ell < n$ .

**Discussion 2.13.** As the sets  $\mathcal{H}(\chi_{\alpha})$  may change with forcing, we may prefer to use  $\mathcal{E}_{\alpha} \subseteq [\chi_{\alpha}]^{\leq \aleph_0}$ , for this we define:

(c)<sub>≡</sub> if (\*)<sub>1</sub> then (\*)<sub>2</sub> where (\*)<sub>1</sub>(α). FILL!

**Definition 2.14.** 1) We call **p** an o.b. (ordinal based) parameter if  $\mathbf{p} = (\bar{\chi}^{\mathbf{p}}, \bar{R}^{\mathbf{p}}, \bar{\mathcal{E}}^{\mathbf{p}}, \bar{D}^{\mathbf{p}})$ <sup>{b42}</sup> and for some ordinal  $\alpha^*$  called  $\ell g(\mathbf{p})$  we have:

- (a)  $\bar{\chi} = \langle \chi_\alpha : \alpha < \alpha^* \rangle$ ,  $\chi_\alpha$  is a regular cardinal and  $\mathcal{H}((\bigcup_{\beta < \alpha} \chi_\beta)^+) \in \mathcal{H}(\chi_\alpha)$
- (b)  $\bar{R} = \langle R_\alpha : \alpha < \alpha^* \rangle$ ,  $R_\alpha$  an  $n(R_\alpha)$ -place relation on some bounded subset of  $\chi_\alpha$  (we could have asked “on  $\chi_\alpha$ ”, no real difference)
- (c)  $\bar{\mathcal{E}} = \langle \mathcal{E}_\alpha : \alpha < \alpha^* \rangle$ ,  $\mathcal{E}_\alpha \subseteq [\chi_\alpha]^{\leq \aleph_0}$  is stationary
- (d)  $\bar{D} = \langle D_\alpha : \alpha < \alpha^* \rangle$  and  $D_\alpha$  is a function with domain  $\mathcal{E}_\alpha$  and for each  $a \in \mathcal{E}_\alpha$ ,  $D_\alpha(a)$  is a family of subsets of  $a$ , closed under supersets, non-empty if  $\alpha > 0$  (let  $D_\alpha^-(a) = (D_\alpha(a))^- = \mathcal{P}(a) \setminus D_\alpha(a)$ )
- (e) let  $\mathbf{p}^{[\alpha]} = \langle \bar{\chi} \upharpoonright \alpha, \bar{R} \upharpoonright (\alpha + 1), \bar{\mathcal{E}} \upharpoonright \alpha, \bar{D} \upharpoonright \alpha \rangle$
- (f) if  $a \in \mathcal{E}_\alpha$  and  $X \subseteq \mathcal{P}(a)$ , then:  $X \in D_\alpha(a) \Leftrightarrow X \cap \bigcup_{\beta < \alpha} \mathcal{E}_\beta \in D_\alpha(a)$ .

2) We say that an o.b. parameter **p** is simple when:

- (g) if  $a \in \mathcal{E}_\alpha$  and  $X \in D_\alpha(a)$  and  $\beta \in \alpha \cap a$ , then  $a \cap \chi_\beta \in \mathcal{E}_\beta$ .

3) For **p** as above let  $\mathbf{q} =: \mathbf{p}^V$  be defined by

$$\alpha^{*,\mathbf{q}} = \alpha^{*,\mathbf{p}}$$

and we define by induction on  $\alpha < \alpha^{*,\mathbf{p}}$ :

$$\chi_\alpha^{\mathbf{q}} = \chi_\alpha^{\mathbf{p}}$$

$$R_\alpha^{\mathbf{q}} = R_\alpha^{\mathbf{p}}$$

$$\mathcal{E}_\alpha^{\mathbf{q}} = \{N \prec (\mathcal{H}(\chi_\alpha^{\mathbf{q}}), \in) : N \text{ is countable, } N \cap \chi_\alpha^{\mathbf{p}} \in \mathcal{E}_\alpha^{\mathbf{p}}, \mathbf{q}^{[\alpha]} \in N\}$$

(note that  $\mathbf{q}^{[\alpha]}$  is well defined by the induction hypothesis)

$$D_\alpha^{\mathbf{q}}(N) = \{Y' : Y' \subseteq \bigcup_{\beta < \alpha} \mathcal{E}_\beta^{\mathbf{q}} \text{ and for some } y \in N \cap \bigcup_{\beta < \alpha} \mathcal{H}(\chi_\beta^{\mathbf{p}}) \text{ and } Y \in D_\alpha^{\mathbf{p}}(N \cap \chi_\alpha^{\mathbf{p}}) \text{ we have } Y' \supseteq \{M : M \in N \cap \bigcup_{\beta < \alpha} \mathcal{E}_\beta^{\mathbf{q}} \text{ and } y \in M \text{ and } M \cap \chi_\alpha^{\mathbf{p}} \in Y\}\}.$$

4) For such **p** we say  $\bar{Q}$  is a  $NNR_\kappa^0$ -iteration for **p** if it is an  $NNR_\kappa^0$ -iteration for  $\mathbf{p}^V$ . We say **p** is simple if  $\mathbf{p}^V$  is. Similarly for  $\mathcal{D}$ -winner, non- $\mathcal{D}$ -loser, etc.

{b45}

**Claim 2.15.** Assume **p** is an o.b. [simple] parameter in the universe **V**.

- 1) If  $\mathbb{P} \in \mathcal{H}(\chi_0^{\mathbf{p}})$  is a proper forcing notion (or at least preserve the stationarity of  $\mathcal{E}_\alpha^{\mathbf{p}}$  for each  $\alpha < \ell g(\mathbf{p})$ ), then  $\Vdash_{\mathbb{P}}$  “**p** is an o.b. [simple] parameter”.
- 2) If forcing with  $\mathbb{P}$  adds no reals, then also  $\mathcal{D}$ -winner, non- $\mathcal{D}$ -loser, etc., are preserved.
- 3)  $\mathbf{p}^V$  is a reasonable parameter.

{b48}

**Definition 2.16.** Let  $\mathfrak{p}$  be an o.b. parameter.

1) We say  $\mathbb{Q}$  is an  $NNR_\kappa^0$ -forcing for  $\mathfrak{p}$  or a  $\mathfrak{p} - NNR_\kappa^0$ -forcing notion when: ( $\mathbb{Q}$  is a forcing notion in a universe  $\mathbf{V}$  and) if for some transitive class  $\mathbf{V}_0, \mathfrak{p} \in \mathbf{V}_0$  and  $NNR_\kappa^0$ -iteration  $\bar{\mathbb{Q}} = \langle \mathbb{P}_i, \mathbb{Q}_j : i \leq \alpha, j < \alpha \rangle$ , we have  $\mathbf{V}_0^{\mathbb{P}_\alpha} = \mathbf{V}$ , then we can let  $\mathbb{Q}_\alpha = \mathbb{Q}$  and get  $\bar{\mathbb{Q}}'$ , an  $NNR_\kappa^0$ -iteration (i.e.  $\mathbf{V} = \mathbf{V}_0[\mathbf{G}_\alpha]$ ,  $\mathbf{G}_\alpha \subseteq \mathbb{P}_\alpha$  is generic over  $\mathbf{V}_0$  and there is a  $\mathbb{P}_\alpha$ -name  $\mathbb{Q}_\alpha$  such that  $\langle \mathbb{P}_i, \mathbb{Q}_i : i \leq \alpha \rangle$  is an  $NNR_\kappa^0$ -iteration and  $\mathbb{Q} = \mathbb{Q}_\alpha[\mathbf{G}_\alpha]$ ). In particular  $\mathbb{Q}$  is proper and does not add reals.

2) If we omit “for  $\mathfrak{p}$ ” we mean for any  $\mathfrak{p}$  which makes sense. Alternatively, we can put a family of  $\mathfrak{p}$ ’s.

3) We add “over  $x$ ” if this holds whenever  $x \in \mathbf{V}_0$ . We can use the same definition for other versions of  $NNR$ .

4)  $\text{Ax}_\lambda^\alpha(\mathfrak{p}, \kappa, 0)$  means: if  $\mathbb{Q}$  is an  $\aleph_2$ -e.c.c. (see [Sh:f, Ch.VII,§1]), ( $\aleph_2$ -pic if  $\lambda = \aleph_2$ ; see [Sh:f, Ch.VIII,§2]), an  $NNR_\kappa^0$ -forcing notion for  $\mathfrak{p}$ ,  $\mathcal{I}_\beta$  a dense open subset of  $\mathbb{Q}$  for  $\beta < \beta^* < \lambda$  and  $\mathcal{S}_i$  is a  $\mathbb{Q}_i$ -name of a stationary subset of  $\omega_1$  for  $i < i^* < \alpha$ , then for some directed  $\mathbf{G} \subseteq \mathbb{Q}$  we have:  $\beta < \beta^* \Rightarrow \mathbf{G} \cap \mathcal{I}_\beta \neq \emptyset, \mathcal{S}_i[\mathbf{G}] = \{\gamma < \omega_1 : (\exists r \in \mathbf{G})(r \Vdash_{\mathbb{Q}} “\gamma \in \mathcal{S}_i”)\}$  is a stationary subset of  $\omega_1$ .

Now we can deduce conclusions on preservation of being an  $NNR_\kappa^0$ -forcing notion for  $\mathfrak{p}$  and consistency of axioms.

{b51}

**Conclusion 2.17.** 1) If  $\mathfrak{p}$  is o.b. parameter, non- $\mathcal{D}'_{\text{id}}$ -loser,  $\ell g(\mathfrak{p}) = \omega_1$  and  $\bar{\mathbb{Q}}$  is an CS iteration such that  $\alpha < \ell g(\bar{\mathbb{Q}}) \Rightarrow \Vdash_{\mathbb{P}_\alpha} “\mathbb{Q}_\alpha \text{ is an } NNR_\kappa^0\text{-forcing notion for } \mathfrak{p}”$ , then  $\bar{\mathbb{Q}}$  is an  $NNR_\kappa^0$ -iteration for  $\mathfrak{p}$  (note 1.10 applies).

{a27}

2) Assume  $\text{CH} + \mu = \mu^{<\mu} \geq \lambda$ , if  $\mathfrak{p}$  is a non- $\mathcal{D}'_{\text{id}}$ -loser o.b. parameter,  $\chi_0^{\mathfrak{p}} > 2^\lambda$ , then for some  $\aleph_2$ -e.c.c., (if  $\lambda = \aleph_2, \aleph_2$ -pic)  $NNR_\kappa^0$ -forcing notion  $\mathbb{P}, |\mathbb{P}| = \mu$  we have  $\Vdash_{\mathbb{P}} “\text{Ax}_\lambda(\mathfrak{p}, \kappa, 0)”$ .

{a33}

3) The parallel of 1.12 holds.

*Proof.* Straight.

□<sub>2.17</sub>



§ 3. EXAMPLES: SHOOTING A THIN CLUB

{examples}

We would like here to see how some examples fit our framework. We already know that  $(< \omega_1)$ -proper forcings are  $\mathfrak{p}$ -proper (by 1.13). We first deal with a forcing notion for which the  $\kappa$  (in  $NNR_\kappa^x$ ) is  $< \aleph_0$  (in Definition 3.1). Second, we deal with shooting clubs of  $\omega_1$  running away from some  $C_\delta \subseteq \delta = \sup(C_\delta)$  which are small (see [Sh:f, Ch.XVIII,§1]). Those are the most natural non- $\omega$ -proper forcing notions not adding reals, (though this may depend on the set theory). The prototype for [Sh:b, Ch.VIII,§4], [Sh:f, Ch.VIII,§4] is 3.1(1) below.

{a36}  
{c3}

{c3}  
{c3}

**Definition 3.1.** 1) Let  $\bar{C} = \langle (C_\delta, n_\delta) : \delta < \omega_1 \text{ limit} \rangle$  where  $C_\delta$  is an unbounded subset of  $\delta$  of order type  $\omega$  and  $1 \leq n_\delta < \omega$  (e.g.  $n_\delta = 1$ ). Let  $\bar{u} = \langle u_\delta : \delta < \omega_1 \text{ limit} \rangle$ , where  $u_\delta \in [2n_\delta + 1]^{n_\delta}$ , (if  $n_\delta = 1$ , then  $u_\delta = \{k_\delta\}, k_\delta \in \{0, 1, 2\}$ ). Then we define

$$\mathbb{Q} = \mathbb{Q}_{\bar{C}, \bar{u}} = \{f : \text{for some } \alpha < \omega_1, f \text{ is a function from } \alpha \text{ to } \omega \text{ such that for every limit ordinal } \delta \leq \alpha, \text{ for some } k < 2n_\delta + 1, k \notin u_\delta \text{ and for every } i \in C_\delta \text{ large enough we have } f(i) = k\}$$

ordered by inclusion.

2) Assume  $\bar{C} = \langle C_\delta : \delta < \omega_1 \text{ limit} \rangle, C_\delta$  a closed subset of  $\delta$  of order type  $< \omega \times \delta$  and for  $\delta_1 < \delta_2$  limit,  $\sup(C_{\delta_1} \cap C_{\delta_2}) < \delta_1$ , and for limit  $\delta^*$  we have  $\{C_\delta \cap \delta^* : \delta < \omega_1\}$  is countable. Assume further  $\bar{\kappa} = \langle \kappa_\delta : \delta < \omega_1 \text{ limit} \rangle, \kappa_\delta \in \{2, 3, \dots, \aleph_0\}, \bar{D} = \langle D_\delta : \delta < \omega_1 \rangle, D_\delta$  is a family of subsets of  $\text{Dom}(D_\delta)$ , such that the intersection of any  $< \kappa_\delta$  of them is non-empty.

Let  $\bar{f} = \langle f_{\delta,x} : \delta < \omega_1, x \in \text{Dom}(D_\delta) \rangle$  satisfy  $f_{\delta,x} : C_\delta \rightarrow \omega$  and  $\bar{A} = \langle A_\delta : \delta < \omega_1 \rangle$  satisfy  $A_\delta \in D_\delta$ . Then we define  $\mathbb{Q} = \mathbb{Q}_{\bar{C}, \bar{D}, \bar{\kappa}, \bar{f}, \bar{A}}$  as

$$\{f : \text{for some } \alpha < \omega_1, f \text{ is a function from } \alpha \text{ to } \omega \text{ such that for every limit } \delta \leq \omega \text{ for some } x \in A_\delta \text{ we have } f_{\delta,x} \subseteq^* f \text{ i.e. for every large enough } i \in C_\delta \text{ we have } f_{\delta,x}(i) = f(i)\}$$

ordered by inclusion.

{c6}  
{c3}

**Claim 3.2.** 1) The forcing notion  $\mathbb{Q}$  from 3.1(1) is proper, does not add reals, and is  $\alpha$ -proper for  $\alpha < \omega_1$  and even is  $(< \omega_1)$ -proper, also  $\mathbb{D}$ -complete for some simple 2-completeness system, hence causes no problem to the demand (d) in Definition 1.10 which means:

{a27}

(\*) if  $\bar{Q}$  is a CS iteration,  $\ell g(\bar{Q}) = \alpha + 1, \bar{Q} \upharpoonright \alpha$  is  $NNR_2^0$ -iteration and  $\Vdash_{\mathbb{P}_\alpha} \text{“}\mathbb{Q}_\alpha = \mathbb{Q}_{\bar{c}, \bar{u}}\text{”}$ ,  $(\bar{c}, \bar{u})$  as above  $\in \mathbf{V}$  then  $\bar{Q}$  satisfies demand ((a),(b) and (d)) of Definition 3.11.

{c27}

2) Similarly for the forcing notion  $\mathbb{Q}$  from 3.1(2).

{c3}

*Proof.* 1) See [Sh:b, Ch.VIII,§4] or [Sh:f, Ch.VIII,§4].

2) Similarly. □<sub>3.2</sub>

*Remark 3.3.* May add uniformization of any function with the right domains, instead constant.

modified:2015-01-27

revision:2015-01-26

(656)

{c9}

**Definition 3.4.** Assume  $\bar{C} = \langle C_\delta : \delta < \omega_1 \rangle, C_\delta$  an unbounded subset of the limit ordinal  $\delta$  (think of the case  $C_\delta$  of order type  $< \delta$  but not necessarily). Let

$$\mathbb{Q}_{\bar{C}} = \{c : \text{for some } \alpha < \omega_1, c \text{ is a closed subset of } \alpha \\ \text{and for every limit ordinal } \delta \leq \alpha \text{ we have :} \\ \delta = \sup(c \cap \delta) \Rightarrow c \cap C_\delta \text{ bounded in } \delta\}$$

$c_1 < c_2$  iff  $c_1$  is an initial segment of  $c_2$ .

{c10}

*Remark 3.5.* 1) See [Sh:f, Ch.XVIII,§1], note  $\mathbb{Q}_{\bar{C}}$  may be not  $\omega$ -proper. We first deal with the simple case:  $\text{otp}(C_\delta) = \omega$ .

2) Note that  $c \in \mathbb{Q}_{\bar{C}}$  is a closed subset of  $\alpha$ , but not necessarily a closed subset of  $\omega_1$ .

{c12}

**Claim 3.6.** Assume  $\mathfrak{p}$  is a simple reasonable parameter,  $\bar{C}$  as in 3.4,  $\bigwedge_\delta \text{otp}(C_\delta) =$

{c9}

$\omega$ , let  $f \in \mathcal{F}^{\mathfrak{p}}$  be  $f(0) = 0, f(\beta) = 1 + \beta$ , then  $\mathbb{Q}_{\bar{C}}$  is  $(\mathfrak{p}, f)$ -proper.

{c15}

*Remark 3.7.* 1) Note that if  $\bigwedge_\delta \text{otp}(C_\delta) < \delta$ , we can split the analysis by, for each ordinal  $\gamma < \omega_1$  such that  $S_\gamma = \{\delta : \text{otp}(C_\delta) = \gamma\}$  is stationary, that is restricting ourselves to  $\{N : N \cap \omega_1 \in S_\gamma\}$ .

{c18}

**Claim 3.8.** 1) Assume

- (a)  $\mathfrak{p}$  is a simple reasonable parameter
- (b)  $\mathfrak{p}$  is non- $\neg_f$ -loser,  $f \in \mathcal{F}^{\mathfrak{p}}$
- (c)  $\gamma(*) \leq \omega_1$
- (d)  $\bar{C} = \langle C_\delta : \delta < \omega_1 \rangle, C_\delta \subseteq \delta = \sup(C_\delta)$
- (e)  $\text{otp}(C_\delta) \leq \omega^{\gamma(*)}$  for every  $\delta$ .

Define  $g \in \mathcal{F}^{\mathfrak{p}}$  by  $g(0) = 0, g(1) = f(1) + \gamma(*), g(\alpha + 1) = f(g(\alpha)) + \gamma(*) + 1$  (and necessarily, for limit  $\alpha, g(\alpha) = \sup_{\beta < \alpha} g(\beta)$ ).

Then for every  $\alpha < \ell g(\mathfrak{p})$ , the forcing notion  $\mathbb{Q} = \mathbb{Q}_{\bar{C}}$  is  $(\mathfrak{p}, \alpha, g(\alpha))$ -proper (version 1).

2) We can in part (1) get “version 2” when: if  $g(\delta) = \delta, N \in \mathcal{E}_\alpha, C \subseteq \omega_1 \cap N = \sup(C), \text{otp}(C) \leq \omega^{\gamma(*)}$  (for example  $C = C_\alpha \cap N$  for some  $\alpha$ ) and  $Y \in D_\alpha(N)$  then  $Y' = \{M \in Y : M \cap \omega_1 \notin C\} \in D_\alpha(N)$ , which is a mild condition.

3) If we weaken clause (b) to  $(b)_{f,g}$  below, then for  $\alpha < \ell g(\mathfrak{p})$ , the forcing notion  $\mathbb{Q}_{\bar{C}}$  is  $(\mathfrak{p}, \alpha, f(\gamma + \alpha))$ -proper, where  $(b)_{f,g} f \in \mathcal{F}^{\mathfrak{p}}$  and  $f(f(\alpha)) = f(\alpha)$  for every  $\alpha < \ell g(\mathfrak{p})$  and  $\mathfrak{p}$  is a non- $\neg'_f$ -loser.

*Proof.* First observe

- (\*) for each  $\alpha < \omega_1$  the set  $\mathcal{S}_\alpha^* = \{p \in \mathbb{Q}_{\bar{C}} : \text{there is } \beta \in p \text{ which is } \geq \alpha\}$  is a dense (and open) subset of  $\mathbb{Q}$ .

1) Similar to the proof of 2) noting version (2) is harder on the chooser, and using the extra freedom we can avoid the need for the extra assumption from (2).

2) We prove this by induction on  $\alpha$ , let  $\beta = g(\alpha)$ .

Let  $N \in \mathcal{E}_\beta$  be countable,  $p \in N \cap \mathbb{Q}_{\bar{C}}$  (and  $\mathbb{Q}_{\bar{C}}, \alpha, \beta \in N$ ) and  $Y \in D_\beta(N)$  be given. Let  $\delta = \delta_N =: N \cap \omega_1$ .

Case 1:  $\alpha = 0$ .

Comprehending the demand in Definition 2.8(1) it just means “ $\mathbb{Q}_{\bar{C}}$  is proper”. {b24}  
 Let  $\langle \mathcal{I}_n : n < \omega \rangle$  list the dense open subsets of  $\mathbb{Q}_{\bar{C}}$  that belongs to  $N$  and we shall choose by induction on  $n$ , a condition  $p_n$  such that:

- (i)  $p_0 = p, p_n \in N$
- (ii)  $p_n \leq p_{n+1} \in \mathcal{I}_n$
- (iii) the set  $p_{n+1} \cup \{\sup p_{n+1}\} \setminus (p_n \cup \{\sup p_n\})$  is disjoint to  $C_\delta$ .

If we succeed to carry out the induction clearly we are done as  $\bigcup p_n$  is as required, noting for each  $\alpha < N \cap \omega_1$ ,  $\mathcal{I}_\alpha^* \in \{\mathcal{I}_n : n < \omega\}$ , hence  $(\exists \beta \in (\bigcup_{n < \omega} p_n))[\alpha \leq \beta < \omega_1]$ .

Also there is no problem to choose  $p_0$ . So assume  $p_n$  has been chosen and we shall choose  $p_{n+1}$  as required. There is a function  $F_n \in N$  with domain  $\mathbb{Q}_{\bar{C}}$  such that:  $q \in \mathbb{Q}_{\bar{C}} \Rightarrow q \leq F_n(q) \in \mathcal{I}_n$ .

For  $\alpha < \omega_1$  let  $q^{[\alpha]} = q \cup \{\sup(q)\} \cup \{\sup(q) + 1 + \alpha\}$ , so clearly  $q \leq q^{[\alpha]} \in \mathbb{Q}_{\bar{C}}$  and the function  $(q, \alpha) \mapsto q^{[\alpha]}$  belongs to  $N$ . Define a function  $H : \omega_1 \rightarrow \omega_1$  by  $H(\alpha) = \sup(F_n(p_n^{[\alpha]}))$ , clearly it is well defined, belongs to  $N$  and let  $C =: \{\beta < \omega_1 : \beta \text{ a limit ordinal, moreover } \omega\beta = \beta \text{ and } (\forall \alpha < \beta)(H(\alpha) < \beta) \text{ and } \sup(p_n) < \beta\}$ , so  $C$  is a club of  $\omega_1$  which belongs to  $N$  and  $\gamma(*) \in N$  hence we can find  $\beta^* \in C$  such that  $\text{otp}(\beta^* \cap C)$  is divisible by  $\omega^{\gamma(*)}$ , but  $\text{otp}(C_\delta \cap \beta^*) < \omega^{\gamma(*)}$  hence for some  $\beta \in C$  we have  $\sup(C_\delta \cap \beta) < \beta$ . Let  $p_{n+1} = F_n(p_n^{\sup(C_\delta \cap \beta) + 1})$ ; it is as required.

Case 2:  $\alpha = 1$ .

Easily  $Y' =: \{M \in Y : M \cap \omega_1 \notin C_\delta\} \in D_{g(1)}(N)$  as  $\beta \geq g(1) = f(1) + \gamma(*)$ , just prove this by induction on  $\gamma(*)$ . Let  $\langle \mathcal{I}_n : n < \omega \rangle$  list the dense open subsets of  $\mathbb{Q}_{\bar{C}}$  which belong to  $N$  and let  $\delta = \bigcup_{n < \omega} \alpha_n$  and  $\alpha_n < \alpha_{n+1} < \delta$ . We now simulate

a strategy for the challenger in the game  $\partial_{\alpha, \beta}(N)$ , together with choosing in the  $n$ -th move, (in the end of it) also the challenger chooses  $p_{n+1} \in \mathbb{Q}_{\bar{C}} \cap N$  such that  $p_0 = p, p_n \leq p_{n+1} \in \mathcal{I}_n$  and during the play letting  $Z_n = \emptyset$  (in fact also the chooser has to use  $Y_n = \emptyset$ ) and  $p_{n+1}$  is  $(M_n, \mathbb{Q}_{\bar{C}})$ -generic and  $\sup(p_{n+1}) > \alpha_n$  and the set  $(p_{n+1} \cup \{\sup p_{n+1}\}) \setminus (p_n \cup \{\sup p_n\})$  is disjoint to  $C_\delta$ . This is possible by Case 1 and its proof because  $M_n \cap \omega_1 \notin C_\delta$  which holds as  $M_n \in Y'$ . As this is a legal strategy for the challenger, so it cannot be a winning strategy hence for some such play the chooser wins hence  $\{M_n : n < \omega\} \in D_1(N)$ , remember  $\alpha = 1$ . Now  $q = \bigcup_{n < \omega} p_n$  is well defined, and  $\sup(q) = \delta$  and  $q \cap C_\delta \subseteq p \cup \{\sup(p)\}$  and  $q \Vdash_{\mathbb{Q}_{\bar{C}}} \text{“}\{M_n : n < \omega\} \subseteq \mathcal{M}[\mathbf{G}_{\mathbb{Q}_{\bar{C}}}, N]\text{”}$ , so  $q$  is as required as the chooser has won the play.

Case 3:  $\alpha > 1, \alpha$  successor.

Similar to Case 2 only we use the induction hypothesis instead of using Case 1.

Case 4:  $\alpha$  a limit ordinal.

Similar to Case 2. Easy. □<sub>3.8</sub>

{c21}

**Definition 3.9.** 1) Assume

- (a)  $S \subseteq \omega_1$  is stationary
- (b)  $f \in {}^{\omega_1}(\omega_1)$

- (c)  $\bar{C} = \langle C_\delta : \delta < \omega_1 \rangle$   
 (d)  $C_\delta$  is an unbounded subset of  $\delta$ .

We define when “ $\bar{C}$  obeys  $f$  on  $S$ ” for  $f \in {}^{\omega_1}\omega_1$  a  $(\mathcal{D}_{\omega_1} + S, \gamma)$ -th function (see part (2)) by induction on  $\gamma$ . For  $f$  being  $(\mathcal{D}_{\omega_1} + S, \gamma)$ -function,  $\gamma < \omega_1$  this means  $\{\delta \in S : \text{otp}(C_\delta) \leq \omega^{1+f(\delta)}\} = S \pmod{\mathcal{D}_{\omega_1}}$ . In general it means that for some  $g : \omega_1 \rightarrow \omega_1$  and pressing down function on  $h$  on  $S$ , for every  $\zeta < \omega_1$  for which  $h^{-1}\{\zeta\}$  is stationary, for some  $\beta < \gamma$  and  $f_\beta$ , a  $(\mathcal{D} + h^{-1}\{\zeta\}, \beta)$ -th function, we have  $\langle C_{g(\delta)} \cap \delta : \delta \in h^{-1}\{\zeta\} \rangle$  obeys  $f_\beta$ .

2) We say  $f \in {}^{\omega_1}(\omega_1)$  is a  $(\mathcal{D}_{\omega_1} + S, \gamma)$ -th function when:  $S \Vdash_{(\mathcal{D}_{\omega_1}^+, \geq)}$  “in  $\mathbf{V}[\mathbf{G}]$ ,  $\{x \in \mathbf{V}^{\omega_1}/\mathbf{G} : \mathbf{V}^{\omega_1}/G \models x \text{ an ordinal } < f/\mathbf{G}\}$  has order type  $\gamma$ ”.

{c24}

**Claim 3.10.** *Assume*

- (a)  $\mathfrak{p}$  is a simple reasonable parameter,  $\ell g(\mathfrak{p})$  of uncountable cofinality  
 (b)  $S \in \mathcal{D}_{\omega_1}^+$  and  $N \in \bigcup_{\alpha} \mathcal{E}_\alpha^{\mathfrak{p}} \Rightarrow N \cap \omega_1 \in S$   
 (c)  $\mathfrak{p}$  is a non- $\bar{\mathcal{D}}_{\alpha, \alpha}$ -loser (or just non- $\bar{\mathcal{D}}'_{\alpha, \alpha}$ -loser) for all  $\alpha \in C^*$ ,  $C^*$  a club of  $\ell g(\mathfrak{p})$ ,  
 (d)  $\bar{C}$  obeys  $f$  on  $S$  which is a  $(\mathcal{D}_{\omega_1} + S, \gamma)$ -function  
 (e)  $0 < \text{Min}(C)$ ,  $g(\alpha) = \text{Min}(C^* \setminus \alpha)$ .

Then  $\mathbb{Q}_{\bar{C}}$  is  $(\mathfrak{p}, g)$ -proper.

*Proof.* Similar. □<sub>3.10</sub>

\* \* \*

Another example (which could have been done in [Sh:f, Ch.XVIII]) is:

{c27}

**Definition 3.11.** 1) We say  $\bar{\mathcal{D}} = \langle \mathcal{D}_\delta : \delta < \omega_1 \text{ limit} \rangle$  is an  $\omega_1$ -filter-sequence if:

- (a)  $\mathcal{D}_\delta$  is a filter on  $\delta$ , containing the co-bounded subsets  
 (b)  $\mathcal{D}_\delta$  is a  $P$ -filter and some  $C_\delta \in D_\delta$  has order type  $\omega$  ( $P$ -filter means that  $\mathcal{D}_\delta$  contains all co-finite sets and if  $A_n \in D_\delta$  for  $n < \omega$  then for some  $A \in D_\delta$  we have  $n < \omega \Rightarrow |A \setminus A_n| < \aleph_0$ ) (can generalize as in 3.1(2) but did not)  
 (c) for every club  $C \subseteq \omega_1$  and  $\alpha < \omega_1$ , the set  $A_C^\alpha[\bar{\mathcal{D}}]$  is stationary, where, by induction on  $\alpha$  we define:  
 $A_C^\alpha[\bar{\mathcal{D}}] = \{\delta < \omega_1 : \delta \text{ is a limit ordinal, } \delta \in C \text{ and for every } \beta < \alpha \text{ we have } \delta = \sup(\delta \cap A_C^\beta[\bar{\mathcal{D}}]), \text{ moreover } \delta \cap A_C^\beta[\bar{\mathcal{D}}] \in \mathcal{D}_\delta\}$ .

{c3}

2) We say a reasonable parameter  $\mathfrak{p}$  obeys  $\bar{\mathcal{D}}$  if for each  $\alpha < \ell g(\mathfrak{p})$  and  $N \in \mathcal{E}_\alpha$  we have:  $\bar{\mathcal{D}} \in N$  and we have

$\mathcal{D}_\alpha^p(N) = \{Y : Y \subseteq N \cap \bigcup_{\beta < \alpha} \mathcal{E}_\beta \text{ is closed (see 2.11) and if}$   
 $\alpha > 0$  then there are  
 $\bar{\beta} = \langle \beta_n : n < \omega \rangle, \bar{M} = \langle M_n : n < \omega \rangle$  satisfying  
 (a)  $\beta_n \in N \cap \alpha$  and  
 (b) either  $\bigwedge [\alpha = \beta_n + 1]$   
 or  $\beta_n < \beta_{n+1}, \sup_{n < \omega} \beta_n = \sup(\alpha \cap N)$   
 (c)  $M_n \in Y \cap \mathcal{E}_{\beta_n}, M_n \in M_{n+1},$   
 $\bigcup_{n < \omega} M_n = N \cap \bigcup_{\beta \in \alpha \cap N} \mathcal{H}(\chi_\beta)$   
 (d)  $\{M_n \cap \omega_1 : n < \omega\} \in \mathcal{D}_{N \cap \omega_1}\}.$

- 3) We say a forcing notion  $\mathbb{Q}$  is a  $\bar{\mathcal{D}} - NNR_\kappa^0$ -forcing if for every reasonable parameter  $\mathfrak{p}$  which obeys  $\bar{\mathcal{D}}$  we have:  $\mathbb{Q}$  is an  $NNR_\kappa^0$ -forcing over  $\mathfrak{p}$  (see 2.15). {b45}  
 4) For a  $\mathbb{P}$ -filter  $\mathcal{D}$ , we say a reasonable parameter  $\mathfrak{p}$  obeys  $\mathcal{D}$  if: for every  $N \in \mathcal{E}_\alpha$

$D_\alpha^p(N) = \{Y : Y \subseteq N \cap \bigcup_{\beta < \alpha} \mathcal{E}_\beta \text{ is closed (see 2.11) and if } \alpha > 0 \text{ then}$   
 for some  $\bar{\beta}, \bar{M}$  satisfying clauses (a), (b), (c)  
 of part (2) and  
 (d)'  $\{n : M_n \in Y\} \in \mathcal{D}$   
 (\*)  $\mathcal{D}$  is a  $P$ -filter on  $\omega\}$ .

- 5) In the parts 1) - 4) above we may replace the word filter by ultrafilter if the  $\mathcal{D}$ 's are ultrafilter (so we have " $\omega_1$ -ultrafilter sequence"). {c30}

**Claim 3.12.** 1) If  $\diamond_{\aleph_1}$  (or much less), then there is an  $\omega_1$ -ultrafilter sequence. {c30}  
 2) If  $\bar{\mathcal{D}}$  is an  $\omega_1$ -filter sequence and  $\langle (\chi_\alpha, \mathcal{E}_\alpha) : \alpha < \alpha^* \rangle$  is a in Definition 1.1, then there is a reasonable parameter  $\mathfrak{p}$  of length  $\omega_1$  obeying  $\bar{\mathcal{D}}$  which is a non- $\bar{\mathcal{D}}$ -loser  $\chi_\alpha^p = \chi_\alpha, \mathcal{E}_\alpha^p = \mathcal{E}_\alpha$ . {a3}  
 3) If  $\diamond_{\aleph_1}$  and  $\langle (\chi_\alpha, \mathcal{E}_\alpha) : \alpha < \alpha^* \rangle$  is as in Definition 1.1 and  $\mathcal{D}$  is a  $\mathbb{P}$ -filter on  $\omega$ , then some reasonable parameter  $\mathfrak{p}$  is  $\mathbb{P}$ -filter like, non- $\bar{\mathcal{D}}_{id}$ -loser with  $\chi_\alpha^p = \chi_\alpha, \mathcal{E}_\alpha^p = \mathcal{E}_\alpha, \alpha^{*,p} = \alpha^*$ . Similarly for ultrafilters. {a3}  
 4) Instead of  $\diamond_{\aleph_1}$  it is enough to assume CH and that for some  $\langle C_\delta : \delta < \omega_1 \text{ limit} \rangle$  and normal filter  $D$  on  $\omega_1$  and for every club  $C$  of  $\omega_1, \{\delta : \delta > \sup(C_\delta \setminus C)\} \in D$ .

*Proof.* Straightforward (in part (4), we define the  $C_\delta$ 's simultaneously). □<sub>3.12</sub> {c33}

**Claim 3.13.** 1) If  $\mathfrak{p}$  is a reasonable parameter obeying  $\mathcal{D}$ , a  $\mathbb{P}$ -filter on  $\omega$  [or  $P$ -ultrafilter on  $\omega$ ], then for some  $\bar{\mathcal{D}}, \bar{\mathcal{D}}$  is an  $\omega_1$ -filter-sequence [or  $\omega_1$ -ultrafilter-sequence] and  $\mathfrak{p}$  obeys  $\bar{\mathcal{D}}$ .  
 2) If  $\mathfrak{p}$  is a  $\mathbb{P}$ -point filter (or ultrafilter), then  $\mathfrak{p}$  is a non- $\bar{\mathcal{D}}$ -loser.

*Proof.* Should be clear. □<sub>3.13</sub> {c36}

*Remark 3.14.* We may consider higher order types than  $\omega$ . {c39}

**Claim 3.15.** 1) Assume  
 (a)  $\bar{C} = \langle C_{\delta,\ell} : \ell < k_\delta, \delta < \omega_1 \text{ limit} \rangle, 1 + \kappa \leq k_\delta \leq \omega, C_{\delta,\ell}$  is a closed unbounded subset of  $\delta$  and  $\ell < m < k_\delta \Rightarrow C_{\delta,\ell} \cap C_{\delta,m} = \emptyset$

(656) revision:2015-01-26 modified:2015-01-27

- (b)  $\mathbb{Q} = \mathbb{Q}_{\bar{C}} = \{C : C \text{ is a closed bounded subset of } \omega_1 \text{ such that for every limit } \delta < \sup(C) \text{ and for every } \ell < k_\delta \text{ except } < 1 + \kappa \text{ many, } \delta \leq \sup(C \cap C_{\delta, \ell})\}$
- (c)  $\mathfrak{p}$  obeys the  $\mathbb{P}$ -ultrafilter  $\mathcal{D}$  and is a reasonable parameter.

Then  $\mathbb{Q}$  is a  $\mathfrak{p} - \text{NNR}_\kappa^0$  forcing notion.

2) In part (1) if we add:  $N \in \mathcal{E}_\alpha$  and  $D_\alpha^{\mathfrak{p}}(N) = \{Y : \{n : M_n \in Y\} \in \mathcal{D}_N\}$ ,  $\mathcal{D}_N$  a  $P$ -point ultrafilter (see above) and  $\ell < k_\delta \Rightarrow \{n < \omega : M_n \cap \omega_1 \in C_{\delta, \ell}\} = \emptyset \text{ mod } \mathcal{D}_N$ , then we can allow  $C_{\delta, 0} = C_{\delta, 1}$ .

3) Assume

- (a)  $D_\delta$  is a family of subsets of  $\text{Dom}(D_\delta)$ , the intersection  $Y$  of any  $< 1 + \kappa$  of them satisfies

$$(*) \quad \exists n(\exists y_1, \dots, y_n \in Y)[\delta > \sup(\bigcap_{\ell=1}^n C_{\delta, y_\ell})]$$

- (b)  $\bar{C} = \langle C_{\delta, x} : x \in \text{Dom}(D_\delta) \text{ and } \delta \text{ is a limit ordinal } < \omega_1 \rangle$
- (c)  $\langle C_{\delta, x} : x \in \text{Dom}(D_\delta) \rangle$  is a sequence of pairwise disjoint subsets of  $\delta$
- (d)  $\bar{X} \in \prod_{\delta < \omega_1} \text{Dom}(D_\delta)$
- (e)  $\mathbb{Q}_{\bar{C}, \bar{X}, \bar{D}} = \{C : C \text{ is a closed bounded subset of } \omega_1 \text{ such that for every limit } \delta \leq \sup(C) \text{ we have } (\exists x \in X_\delta)(\delta > \sup(C \cap C_{\delta, x}))\}$  ordered by: end extension
- (f)  $\bar{\mathcal{D}}$  is a  $\mathbb{P}$ -ultrafilter sequence.

Then  $\mathbb{Q}$  is a  $\mathfrak{p} - \text{NNR}_\kappa^0$  forcing notion  $\mathfrak{p}$  which obeys  $\bar{\mathcal{D}}$ .

*Proof.* Straight.

{c39} 1) So let  $\mathbf{V} = \mathbf{V}_0^{\mathbb{P}_\alpha}, \Vdash_{\mathbb{P}_\alpha}$  “ $\mathbb{Q}_\alpha = \mathbb{Q}_{\bar{C}}$  is as above” and in  $\mathbf{V}_0, N_0 \in \mathcal{E}_0, N_0 \in N_1 \in \mathcal{E}_\alpha = \{M_n : n < \omega\}, D_{N_0}$  as in Definition 3.15(2),  $p \in \mathbb{P}_{\alpha+1} \cap N, G_m \subseteq N_1 \cap \mathbb{P}_\alpha$  is generic over  $N_1$  for  $m < k < 1 + \kappa$  (so  $k$  is now fixed). So for each  $\ell$  we have  $\mathcal{C}_{N_0 \cap \omega_1, \ell}[\mathbf{G}_m]$  is a closed subset of  $\delta$  and for  $\ell_1 < \ell_2 < k_{\delta_{N_0}}$  we have  $\mathcal{C}_{N_0 \cap \omega_1, \ell_1}[\mathbf{G}_m] \cap \mathcal{C}_{N_0 \cap \omega_1, \ell_2}[\mathbf{G}_m] = \emptyset$  so for some  $\ell(m) \in \{0, 1, \dots, k_{\delta_{N_0}} - 1\}$  we have  $\ell \neq \ell(m) \Rightarrow \mathcal{C}_{N_0 \cap \omega_1, \ell}[\mathbf{G}_m] = \emptyset \text{ mod } D_{N_0}$ . Now let  $B = \{n : \text{if } \ell < k_\delta, \ell \notin \{\ell(m) : m < k\} \text{ and } \ell < n \text{ then } M_n \cap \omega_1 \notin \mathcal{C}_{N_0 \cap \omega_1, \ell}[\mathbf{G}_m] \text{ for } m < k \text{ and } p \in M_n\}$  belongs to  $D_{N_0}$ . Clearly, without loss of generality  $\langle M_n : n < \omega \rangle \in N_1, D_{N_0} \in N_1, B \in N_1$ . Let  $B = \{n_i : i < \omega\}, n_i$  increasing with  $i$  and  $\langle \mathcal{I}_n : n < \omega \rangle$  list the dense open subsets of  $\mathbb{P}_{\alpha+1}$  which belongs to  $N_0$ . We can choose  $p_i$  by induction on  $i < \omega$  such that:

- (\*) (a)  $p_i \in N_{n_i} \cap P_{\alpha+1}$
- (b)  $p_i \upharpoonright \alpha \in \bigcap_{m < k} G_m$
- (c)  $p_i \in \cap \{\mathcal{I}_n : n < n_i, \mathcal{I}_n \in N_{n_i} \text{ and } i > 0\}$
- (d)  $p \leq p_i$
- (e)  $p_i \leq p_{i+1}$

$p_{i+1} \setminus p_i$  is disjoint to  $\cup \{\mathcal{C}_{N_0 \cap \omega_1, \ell}[\mathbf{G}_m] : \ell < k_0, \ell < n_i \text{ and } m < k \Rightarrow \ell \neq \ell(m)\}$ .

This is possible as

$$\otimes \quad (\bigcup_{\ell < k \text{ and } \ell \neq \ell(m)}^{m < k} \mathcal{C}_{N_0 \cap \omega_1, \ell}[\mathbf{G}_m]) \cap \omega_1 \cap M_n \text{ is a bounded subset of } M_n \cap \omega_1.$$

2), 3), 4) Similarly. □<sub>3.15</sub>

{c42}  
 {c39} **Observation 3.16.** 1) The forcing notion from 3.15(1) is  $NNR_{\aleph_0}^0$ -forcing notion for every  $\mathfrak{p}$ , non- $\mathcal{D}_{\text{id}}$ -loser.  
 2) If  $\mathfrak{p}$  is a reasonable parameter obeying  $\bar{\mathcal{D}}$ , an  $\omega_1$ -filter sequence, then any  $(< \omega_1)$ -proper forcing is  $\mathfrak{p}$ -proper.

*Proof.* Similar to earlier proofs. □<sub>3.16</sub>

## § 4. SECOND PRESERVATION OF NOT ADDING REALS

{second}

We shall concentrate on the simple case.

{d3}

**Definition 4.1.** Let  $\mathfrak{p}$  be a reasonable parameter. We say that  $\bar{\mathbb{Q}}$  is a  $\mathfrak{p} - NNR_\kappa^1$  iteration (where  $2 \leq \kappa \leq \aleph_0$  as we omit  $\kappa = \aleph_1$  for convenience) when for some  $f_i, g_i \in \mathbf{F}_{cc}^{\mathfrak{p}}$ , for  $i < \ell g(\bar{\mathbb{Q}})$  (see 2.3(4),(5) we have:

{b9}

- (a)  $\text{cf}(\ell g(\mathfrak{p})) > \ell g(\bar{\mathbb{Q}})$  (for simplicity)
- (b)  $\bar{\mathbb{Q}}$  is a countable support iteration of proper forcing notions such that  $i < \ell g(\bar{\mathbb{Q}}) \Rightarrow \mathbb{P}_i$  adds no reals (follows by other parts (close (d)) even for  $\mathbb{P}_{i+1}, i < \ell g(\bar{\mathbb{Q}})$ )
- (c) [long properness] for each  $i < \ell g(\bar{\mathbb{Q}})$ , we have  $\Vdash_{\mathbb{P}_i}$  “ $\mathbb{Q}_i$  is  $(\mathfrak{p}^{\mathbb{P}_i}, f_i)$ -proper, see Definition 2.8(2) + 2.8(4) (hence has club of fix points)”
- (d) [ $\kappa$ -anti w.d.] if  $\beta \in g_i(\alpha)$  and  $i < \ell g(\bar{\mathbb{Q}})$  then  $\mathbb{Q}_i$  has  $(\kappa, \alpha, \beta)$ -anti w.d. above  $\mathbb{P}_i$ .

{b24}

{d6}

*Remark 4.2.* 0) In [Sh:311] we intend to relax the requirement that each iterand is proper.

{c9}

1)  $\kappa$  is the amount of “ $\mathbb{D}$ -completeness”, in other words what versions of weak diamond we kill by our iteration. So  $\kappa = \aleph_0$  is easier and we first deal with it in 3.4.

2) Question: Why do we ask for  $f_i \in \mathbf{V}$  and not a  $\mathbb{P}_i$ -name  $\check{f}_i$  of such a function?

Answer: If  $i < \ell g(\bar{\mathbb{Q}}) \Rightarrow P_i \models \text{cf}(\ell g(\mathfrak{p}))\text{-c.c.}$ , then we can find  $f'_i \in \mathcal{F}^{\mathfrak{p}}, f_i \geq \check{f}_i$ ; so as we are assuming, for conveniency  $\text{cf}(\ell g(\mathfrak{p})) > |\mathbb{P}_\alpha|$  there is no point at present for  $f_i$  to be a  $\mathbb{P}_i$ -name.

2A) In clause (d) we have implicitly used:

- (\*) if  $\alpha \leq \beta' < \beta$  then clause (d) for  $(\alpha, \beta)$  and  $k$  implies clause (d) for  $(\alpha, \beta')$  and  $k$ .

{a3}

This holds by clause (i) of Definition 1.1.

We could replace  $f_i$  by a club  $E_i$  of  $\ell g(\mathfrak{p})$ , letting  $f_i(\alpha) = E_i \setminus \alpha$ .

3) No real reason to “use some  $\kappa$ ”, but also no real reason to use 2. For each  $\ell$  to find  $\mathbf{G}^{**}$  means essentially “forcing with  $\mathbb{Q}_i$  adds no reals”. The point is the common solution.

4) In clause (c) for a club  $C$  of  $\ell g(\mathfrak{p})$  we catch our tail, that is  $f_i(\alpha) \cap C = C \setminus \alpha$  for a club of  $\alpha < \ell g(\mathfrak{p})$ . We could use less, no real point now.

5) In clause (d) much of the freedom/variation will be due to the decision how “similar” are  $\langle G^\ell : \ell < k \rangle$  such that  $\mathbf{G}^{**}$  exists. Here we demand

- ( $\alpha$ )  $Y \in D_\beta(N_1)$ .

In [Sh:f, Ch.VIII,§2] it is essentially required that

- ( $\beta$ )  $\mathbf{G}^0 \times \mathbf{G}^1 \times \dots \times \mathbf{G}^{k-1} \subseteq (\mathbb{P}_i \times \dots \times \mathbb{P}_i) \cap N_1$  ( $k$  times) is generic over  $N_1$ .

In [Sh:f, Ch.V]

- ( $\gamma$ ) the common  $Y$  is a predetermined increasing sequence of models.



{d3} We have a trade-off; clause  $(\beta)$  makes demand  $(d)$  in 4.1 easier, but the parallel of  $(c)$  harder compared to clause  $(\alpha)$ . This explains why the present work doesn't supercede Ch.XVIII,§2.

6) It is harder to win with a "slower" function, so the assumption above is the strongest possible - although practically makes no difference, probably.

**Claim 4.3.** *Let  $\bar{\mathbb{Q}}$  be a  $\mathfrak{p} - \text{NNR}_\kappa^1$  iteration (see Definition 4.1),  $\kappa = \aleph_0$  and  $\mathfrak{p}$  is a reasonable parameter and  $\mathfrak{p}$  is a  $\mathcal{D}_f$ -winner for some  $f \in \mathcal{F}_{\text{club}}^{\mathfrak{p}}$  or at least  $\mathcal{D}'_f$ -non loser.*

1) *Forcing with  $\mathbb{P}_{\ell g(\bar{\mathbb{Q}})} = \text{Lim}(\bar{\mathbb{Q}})$  does not add reals (so consequently no  $\omega$ -sequences, as we are assuming properness).*

2) *If  $i \leq j \leq \ell g(\bar{\mathbb{Q}})$ , then*

(b)'  $\mathbb{P}_j/\mathbb{P}_i$  is proper

(c)'  $\mathbb{P}_j/\mathbb{P}_i$  is  $(\mathfrak{p}, f_{i,j})$ -proper, where  $f_{i,j} \in \mathcal{F}_{\text{club}}^{\mathfrak{p}}$  is increasing continuous and is computable from the  $f_\varepsilon \in \mathcal{F}^{\mathfrak{p}}$  for  $\varepsilon \in [i, j)$ ,

(d)' we have the parallel of clause  $(d)$ , this time without assuming that  $j = i + 1$ .

*Proof.* The proof is by induction on  $\ell g(\bar{\mathbb{Q}})$ . For notational simplicity we assume

☒ all  $f_i$  are even in  $\mathcal{F}_{\text{ndd}}^{\mathfrak{p}}$  so consider them as functions from  $\ell g(\mathfrak{p})$  to  $\ell g(\mathfrak{p})$  (increasing continuous), and we demand also the  $f_{i,j}$  are like that, increasing continuous and moreover  $f_{i,j}(f_{i,j}(\alpha)) = f_{i,j}(\alpha)$  and they are  $\geq f^*$ ,  $f^* \in \mathcal{F}_{\text{nd}}^{\mathfrak{p}}$  increasing continuous,  $\mathfrak{p}$  is  $\mathcal{D}_{f^*}$ -winner or at least  $\mathcal{D}'_{f^*}$ -non-loser.

Case 1:  $\ell g(\bar{\mathbb{Q}}) = 0$

Trivial.

Case 2:  $\ell g(\bar{\mathbb{Q}}) = i(*) + 1$

Part (1):  $\mathbb{P}_{i(*)}$  adds no reals by the induction hypothesis,  $\mathbb{Q}_{i(*)}$  adds no reals by clause  $(d)$  in Definition 4.1, hence  $\mathbb{P}_{i(*)+1} = \mathbb{P}_{i(*)} * \mathbb{Q}_{i(*)}$  adds no reals.

Part (2): Clause  $(b)'$  is known, (namely  $\mathbb{P}_j/\mathbb{P}_i$  is proper).

Clause  $(c)'$ :

Given  $i \leq j \leq \ell g(\bar{\mathbb{Q}})$  if  $j < i(*) + 1$  the conclusion follows by the induction hypothesis. So assume  $j = i(*) + 1$ . If  $i = j$ , the required demand is trivial, so assume  $i < j$ . If  $i = i(*)$ , use clause  $(c)$  of Definition 4.1 for  $i$ . So assume  $i < i(*)$ . Let  $f_{i,j,0} = f_{i(*)}$ ,  $f_{i,j,m+1} = f_{i(*)} \circ f_{i,j(*)} \circ f_{i,j,m}$  and  $f_{i,j}(\alpha) = \sup_{m < \omega} f_{i,j,m}(\alpha)$ . Check

that they are as required in ☒. For proving " $\mathbb{P}_j/\mathbb{P}_i$  is  $(\mathfrak{p}, f_{i,j})$ -proper" assume

- (\*)<sub>1</sub> (a)  $N \prec (\mathcal{H}(\chi), \in)$  is countable
- (b)  $\{\bar{\mathbb{Q}}, i, j, \alpha, \beta, f_{i,i(*)}, f_{i(*)}, f_{i,j}\} \in N$
- (c)  $\alpha \leq f_{i,j}(\alpha) \leq \beta < \ell g(\mathfrak{p})$
- (d)  $q \in \mathbb{P}_i$  is  $(N, \mathbb{P}_i)$ -generic
- (e)  $p \in N \cap \mathbb{P}_j$ ,  $p \upharpoonright i \leq q$
- (f)  $Y \in D_\beta(N)$
- (g)  $q \Vdash "Y \subseteq \mathcal{M}[\mathbf{G}_{\mathbb{P}_i}, N]"$ .

modified:2015-01-27

(656) revision:2015-01-26

{d9}  
{d3}

{d3}

{d3}

First we deal with version 2 and assume that  $\mathbf{p}$  is simple. Choose  $y^* \in N$  which codes enough; clearly  $\beta' =: f_{i(*)}(\alpha)$  belongs to  $N$ . So  $f_{i,i(*)}(\beta') \leq \beta$ , hence by the induction hypothesis there are  $q', Y'$  such that:  $p \upharpoonright i(*) \leq q, q \leq q' \in \mathbb{P}_{i(*)}, q'$  is  $(N, P_{i(*)})$ -generic,  $Y' \subseteq Y, Y' \in D_{\beta'}(N)$  and  $q' \Vdash "Y' \subseteq \mathcal{M}[\mathbf{G}_{\mathbb{P}_{i(*)}}, N, y^*]"$ .

{d3} Next, we apply clause (c) in the Definition 4.1 for  $i(*)$ , so there are  $q'', Y''$  such that  $p \leq q'', q' \leq q'' \in \mathbb{P}_{i(*)+1} = \mathbb{P}_j, q''$  is  $(N, \mathbb{P}_j)$ -generic,  $Y'' \subseteq Y', Y'' \in D_\alpha(N)$  and  $q'' \Vdash "Y \subseteq \mathcal{M}[\mathbf{G}_{\mathbb{P}_j}, N, y^*]"$ .

The proof for version 1 is similar.

Clause (d)':

Recall that we have demanded  $f_{i,j}(f_{i,j}(\alpha)) = f_{i,j}(\alpha)$  (see  $\boxtimes$  at the beginning of the proof).

{d3} Let  $N_0, N_1, \mathbf{G}^*, \alpha, \beta, i, j, p, k, q_\ell$  (for  $\ell < k$ ),  $\mathbf{G}^\ell$  (for  $\ell < k$ ) and  $\mathbf{G}^*$  be as in the assumption of clause (d) (from Definition 4.1) be given.

Without loss of generality  $i < j, i < i(*), j = i(*) + 1$  (as in the proof of clause (c) the other cases are trivial). First, choose  $\mathbf{G}^{**}$  for  $\mathbf{G}^*, \alpha, \beta, i, i(*), \mathbf{G}^{**} \in N_1$ . For each  $\ell < k$ , if for some  $s_\ell \in \mathbf{G}^\ell$  we have

(\*)<sub>2</sub>  $s_\ell \Vdash_{\mathbb{P}_i} " \in \mathbb{P}_{i(*)}/\mathbf{G}_{\mathbb{P}_i}$  there is an upper bound to  $\mathbf{G}^{**}"$ .

then (possibly increasing  $s_\ell$ , recalling  $\mathbf{G}_\ell$  is generic over  $N^*$ ) there are  $s_\ell \in \mathbf{G}^\ell, r_\ell \in P_{i(*)} \cap N_1$  such that  $s_\ell$  forces that  $r_\ell$  is an upper bound to  $\mathbf{G}^{**}$  and without loss of generality  $r_\ell \upharpoonright i \leq s_\ell$ . Now without loss of generality  $[\mathbf{G}_{\ell_1} = \mathbf{G}_{\ell_2} \Rightarrow s_{\ell_1} = s_{\ell_2}]$  and  $[\mathbf{G}_{\ell_1} \neq \mathbf{G}_{\ell_2} \Rightarrow s_{\ell_1}, s_{\ell_2}$  incompatible]. Now choose  $r \in \mathbb{P}_{i(*)} \cap N_1$  with domain  $\subseteq i(*) \setminus i$  as follows:  $\text{Dom}(r) = \bigcup_{\ell < k} \text{Dom}(r_\ell) \setminus i$  and  $r(\alpha)$  is  $r_\ell(\alpha)$  if  $s_\ell \in \mathbf{G}_{\mathbb{P}_i}, \ell < k$  and is  $\emptyset_{\mathbb{P}_\alpha}$  if this occurs for no  $\ell$ . Renaming  $r \in N_i \cap \mathbb{P}_{i(*)}, \text{Dom}(r) \subseteq i(*) \setminus i$  and  $s_\ell \in \mathbf{G}^\ell, r_\ell = s_\ell \cup r$  is above  $\mathbf{G}^{**}$  in  $\mathbb{P}_{i(*)}$ . Let  $\beta_\ell = f_{i,j,1+\ell}(\alpha)$  for  $\ell \leq k$ .

We choose by induction on  $\ell \leq k$  the objects  $Y_\ell, q'_\ell, M_\ell$  such that:

- (\*)<sub>3</sub> (a)  $Y_0 = Y$   
 (b)  $M_0 = N_1$   
 (c)  $N_0 \in M_{\ell+1}$   
 (d)  $M_{\ell+1} \in M_\ell \cap \mathcal{E}_{\beta_{k-\ell}}$   
 (e)  $Y_{\ell+1} \subseteq Y_\ell$   
 (f)  $Y_\ell \in D_{\beta_\ell}(M_\ell)$   
 (g)  $M_{\ell+1} \in Y_\ell$   
 (h)  $q_\ell \leq q'_\ell \in P_{i(*)}$   
 (i)  $q'_\ell$  is  $(M_{\ell+1}, \mathbb{P}_{i(*)})$ -generic  
 (j)  $q'_\ell$  forces a value to  $\mathbf{G}_{\mathbb{P}_{i(*)}} \cap M_{\ell+1}$   
 (k)  $q'_\ell$  is  $(N_0, \mathbb{P}_{i(*)})$ -generic  
 (l)  $q'_\ell \Vdash "Y_{\ell+1} \subseteq \mathcal{M}[\mathbf{G}_{\mathbb{P}_{i(*)}}, M_\ell]"$   
 (m)  $q'_\ell \upharpoonright i \in G^\ell$ .

(In the older version we have to increase  $\beta_\omega$  times if  $k$  not given, arriving to a fixed point.)

No problem, now apply clause (d) of the definition for  $i(*), N_0, M_k, \langle q'_\ell : \ell < k \rangle, Y_k, \mathbf{G}^{**}$  and get  $\mathbf{G}^{***}$  as required.

Comment: The  $Y$  transfers information between the various generics. In [Sh:f, Ch.V], in the first proof after  $\omega$  steps we are lost, but having the common tower of models help us.

Case 3:  $\delta = \ell g(\bar{\mathbb{Q}})$  is limit.

Part (1): Follows from clause (d) in part (2) and part (2) is proved below.

Part (2): Let  $f_{i,j}$  be fast enough (you can collect the demands used below).

Clause (b)' is obvious. We first prove clause (d)' and later (c)'.

Clause (d)': So again without loss of generality  $i < j = \delta$ . Let  $N_0, N_1, p, \mathbf{G}^*, \alpha, \beta, k < 1 + \kappa$  and  $\mathbf{G}_\ell, q_\ell$  for  $\ell < k$  be as in the assumption of clause (d) in Definition 4.1 {d3} but for  $i < j$ .

We for simplicity use  $\kappa = \aleph_0$ . We choose  $\gamma \in N_0, \alpha < \gamma < \beta$  such that  $\gamma$  is larger enough, in particular:  $i \leq i' < j' < j \Rightarrow f_{i',j'}(\gamma) = \gamma$ .

Choose  $\langle i_m : m < \omega \rangle \in N_1$  such that:

- (\*)<sub>4</sub> (a)  $i_0 = i$
- (b)  $i_m < i_{m+1}$
- (c)  $\sup_{m < \omega} i_m = \sup(j \cap N_0)$ .

Choose  $y^* \in N \cap \mathcal{H}(\chi_\gamma)$  coding enough. We choose  $M_0, M_1, M_2, M_3, M_4 \in N_1 \cap \mathcal{E}_\gamma \cap Y$  such that:

- (\*)<sub>5</sub> (a)  $N_0 \in M_0 \in M_1 \in M_2 \in M_3 \in M_4$
- (b)  $Y \cap M_m \in D_\gamma(M_m)$  for  $m < 5$ .

Choose  $q'_\ell \in \mathbf{G}^\ell \cap M_4$  above  $\mathbf{G}^\ell \cap M_3$  so  $q'_\ell$  is  $(M_0, \mathbb{P}_i)$ -generic,  $(M_1, \mathbb{P}_i)$ -generic,  $(M_2, \mathbb{P}_i)$ -generic,  $(M_3, \mathbb{P}_i)$ -generic. Let  $\langle \mathcal{S}_m : m < \omega \rangle \in M_0$  list the dense open subsets of  $\mathbb{P}_j$  from  $N_0$ . □<sub>4.3</sub>

**Explanation 4.4.** Now we shall use the diagonal argument, choose  $\mathbf{G}_\mathbb{P} \cap N_0, p_m \in \mathbb{P}_{i_m} \cap N_0, r_m \in \mathbb{P}_{i_m}$  as usual, using in the  $m$ -th step  $f_{i_m, i_{m+1}}$  + relevant (d)' things. We fulfill the above in  $M_4$ , so in the end can find a solution in  $N_1$ , by using a canonical construction, e.g. each time the  $\langle \cdot \rangle_\chi^*$ -choice. {d13}

But to carry this we need to have finitely many candidates for  $\mathbf{G}_{\mathbb{P}_{i_m}} \cap M_0$  with a common  $Y_m$ . (Note: if  $\mathbb{Q} \times \dots \times \mathbb{Q}$  is proper, like in Ch.XVIII, §2 we can get such a common  $Y_m$ , i.e. the present result is stronger for this stage.) To get this in the inductive step, we need in step  $m - 1$  that for  $M_1$  we just have finitely many candidates for  $\mathbf{G}_{\mathbb{P}_{i_m}} \cap M_1$ , and in turn to get this in the step  $m - 1$  we use that in step  $m - 2$  for  $M_2$  we have: from every maximal antichain we choose a finite subset. To get this we use that for  $M_3$  we just ask  $M_3[\mathbf{G}_{\mathbb{P}_{i_{m-3}}}] \cap \mathbf{V} = M_3$ . So along the way  $N_0, M_0, M_1, M_2, M_3$  our induction demands go down, but slowly, so that in each step  $m$ , advancing for say  $M_0$ , we have to preserve less than really knowing  $\mathbf{G}_{\mathbb{P}_{i_m}} \cap M_0$ , and are helped by our demand on  $M_1$ , just like in [Sh:f, Ch.XVIII]. So compared to [Sh:f, Ch.V], we have a finite tower.

Returning to the proof we choose by induction on  $m < \omega$  the objects  $r_m, \mathbf{G}_m^*, p_m, n_m, \langle \mathbf{G}_m^\ell : \ell < n_m \rangle, Y_m$  such that

- (\*)<sub>6</sub> (a)  $r_m \in \mathbb{P}_{i_m} \cap M_4$
- (b)  $\text{Dom}(r_m) \subseteq [i, i_m)$

modified:2015-01-27

(656) revision:2015-01-26

- (c)  $r_{m+1} \upharpoonright i_m = r_m$   
(comment:  $r_m$ 's act as a generic for  $N_0$ )
- (d)  $q'_\ell \cup r_m \in \mathbb{P}_{i_m}$  is  $(M_0, \mathbb{P}_{i_m})$ -generic,  $(M_1, \mathbb{P}_{i_m})$ -generic,  
 $(M_2, \mathbb{P}_{i_m})$ -generic,  $(M_3, \mathbb{P}_{i_m})$ -generic
- (e) if  $\ell < k$ ,  $\mathcal{J} \subseteq \mathbb{P}_{i_m}$  is dense open and  $\mathcal{J} \in M_2$ , then for some finite  
 $\mathcal{J} \subseteq \mathcal{J} \cap M_2$ ,  $\mathcal{J}$  is predense above  $q'_\ell \cup r_m$   
(like in the proof of  ${}^\omega\omega$ -bounding)
- (f)  $n_m < \omega$ , and  $\ell < n_m \Rightarrow \mathbf{G}_m^\ell$  is a subset of  $\mathbb{P}_{i_m} \cap M_0$  generic over  $M_0$ ,  
and  $\mathbf{G}_m^\ell \in M_1$
- (g)  $\mathbf{G}_{m+1}^\ell \cap \mathbb{P}_{i_m} \in \{\mathbf{G}_m^\ell : \ell < n_m\}$
- (h)  $n_0 = k$ ,  $\mathbf{G}_0^\ell = \mathbf{G}^\ell \cap M_1$
- (i)  $q_\ell \cup r_m \Vdash \text{“}\mathbf{G}_{\mathbb{P}_{i_m}} \cap M_1 \in \{\mathbf{G}_m^\ell : \ell < n_m\}$ ”
- (j)  $\mathbf{G}_m^*$  is a subset of  $\mathbb{P}_{i_m} \cap N_0$  generic over  $N_0$
- (k)  $\mathbf{G}_m^* \subseteq \mathbf{G}_m^\ell$ , so  $\mathbf{G}_m^* \subseteq \mathbf{G}_{m+1}^*$ ,  $\mathbf{G}_0^* = \mathbf{G}^*$
- (l)  $p_m \in \mathbb{P}_j \cap N$
- (m)  $p_0 = p$
- (n)  $p_m \upharpoonright i_m \in \mathbf{G}_m^*$
- (o)  $p_m \leq p_{m+1} \in \mathcal{I}_m$
- (p)  $Y_m \subseteq \mathcal{M}[\mathbf{G}_m^\ell, M_0, y^*]$
- (q)  $Y_m \in D_\gamma(M_0)$ .

Why is this sufficient? If we can carry out the induction, then without loss of generality the construction belongs to  $N_1$ . So  $\mathbf{G}^{**} = \{s \in \mathbb{P}_j \cap N_0 : \bigvee_{n < \omega} s \leq p_n\}$  is as required, as  $q'_\ell =: q_\ell \cup \bigcup_m r_m \in \mathbb{P}_j \cap N_1$ , and is above  $\mathbf{G}^{**}$  and  $p \leq q'_\ell$ .

### Induction:

$m = 0$ : Trivial.

$m + 1$ : Stage A: Choosing  $p_{m+1}$  is trivial, the demands are:  $p_{m+1} \geq p_m$ ,  $p_{m+1} \upharpoonright i_m \in \mathbf{G}_m^*$  and  $p_{m+1} \in \mathcal{I}_m$ .

Stage B: Choosing  $\mathbf{G}_{m+1}^*$ : apply the induction hypothesis using clause (d)' of what we are proving with  $i_m, i_{m+1}, \gamma, f_{i_m, i_{m+1}}(\gamma), N_0, M_1$  here standing for  $i, j, \alpha, \beta, N_0, N_1$  there.

Stage C: Let  $\{H_m^\ell : \ell < n_{m+1}\}$  list the possibilities of  $\mathbf{G}_{\mathbb{P}_{i_m}} \cap M_1$  (by clause (e) this exists).

Without loss of generality  $H_m^\ell \cap M_0 = G_m^{h(\ell)}$ , for some function  $h = h_m : n_{m+1} \rightarrow n_m$ . We choose  $s_m^\ell \in \mathbb{P}_{i_{m+1}} \cap M_1$ , above  $\mathbf{G}_{m+1}^*$ , such that  $s_m^\ell \upharpoonright i_m \in \mathbf{G}_m^{h(\ell)}$ . Now repeat the argument of the successor stage of shrinking  $Y$  (but now we have a fixed  $\gamma$ !).

So we can find  $t_m^\ell \in \mathbb{P}_{i_{m+1}} \cap M_1$ , above  $s_m^\ell$ ,  $t_m^\ell \upharpoonright i_m \in H_m^\ell$ ,  $t_n^\ell \Vdash \text{“}\mathbf{G}_{\mathbb{P}_{i_{m+1}}} \cap M_0 =: \mathbf{G}_{m+1}^\ell$ ” such that:

$$(*)_7 \ Y_{m+1} =: \bigcap_{i < n_{m+1}} \mathcal{M}[\mathbf{G}_{m+1}^\ell, M_0, y^*] \in D_\gamma(M_0).$$

The rest is as in §1 so we have finished proving clause (d)' in the case  $lg(\bar{\mathbb{Q}})$  is a limit ordinal (which is last).

Clause (c)': Again, without loss of generality  $i < j = \delta$ . So assume  $f_{i,j}(\alpha) \leq \beta, \{i, j, \alpha, \beta\} \in N^* \in \mathcal{E}_\beta, q$  is  $(N^*, \mathbb{P}_i)$ -generic,  $Y^* \in D_\beta(N^*), q \in \mathbb{P}_\alpha$  and  $q \Vdash "Y^* \subseteq \mathcal{M}[\mathbf{G}_{\mathbb{P}_i}, N^*, y^*]"$  are given. We prove the desired conclusion by induction on  $\alpha$ . For each  $\alpha$ , we would like to simulate a play of  $\mathcal{D}_{\alpha,\beta}(N^*)$ , supplying the challenger with a strategy. For this we apply the proof of clause (d)'. Choose  $N_0, N_1, M_0, \dots, M_4, q_0, \mathbf{G}_0, \mathbf{G}^*$  (and  $k = 1$ ) as there for some  $\alpha' < \gamma' < \beta'$  as there such that  $\beta < \alpha'$  such that  $N^*, q, Y^* \in N_0$  (easy to find).

{d17}

**Explanation 4.5.** Note: as  $\mathfrak{p}$  is simple, we can use  $N \prec (\mathcal{H}(\chi), \in), N \cap \mathcal{H}(\chi_\beta) \in \mathcal{E}_\beta$  with no mention - there were other places we could have said so. During the construction this time we demand  $p_m \in N^* \cap \mathbb{P}_j$ , so a generic for  $N_0$  is not necessarily created. But still  $p_m \leq p_{m+1}, p_m \upharpoonright i_m \in \mathbf{G}_m^*$ . Now  $p_m$  will be played by the chooser. Now  $g(1 + \alpha)$  will be a fixed point of  $f_{i_m, i_{m+1}}$ . So we can add the demand  $N^*[\mathbf{G}_m^* \cap N^*] \cap \mathbf{V} = N^*$ , i.e.  $\mathbf{G}_m^*$  is generic over  $N^*$  and

$$(*)_8 \mathcal{M}[\mathbf{G}_m^* \cap N^*, N^*, y^*] \in D_{\delta(1+\alpha)}(N^*).$$

The challenger chooses

$$(*)_9 X_{m+1} = \mathcal{M}[\mathbf{G}_m^* \cap N^*, N^*, y^*] \cap \bigcup_{\xi < \gamma(1+\alpha)} \mathcal{E}_\xi \cap \{M : p_n \in M\} \in N_0.$$

Now the chooser chooses  $\alpha_m, \beta'_m$  and the challenger chooses  $\beta_n \geq \beta'_m, f_{i_m, i_{m+1}}(\alpha)$  in  $N_0^* \cap j$  and the chooser chooses  $M_m^*$  such that  $p_m \in M_{m+1}$ .

Comment: The game was defined with this point in mind.

Now the chooser chooses  $Y_n \in D_{\beta_n}(M_0), Y_n \subseteq X_n \cap M_0, Y_n \in N_0^*$  (check definition of game!) Now we play  $Z_n$  for the challenger as follows:

- there is  $p_{m+1} \geq p_m, (M_{m+1}^*, \mathbb{P}_{i_{m+1}})$ -generic
- $p_{m+1} \upharpoonright i_m \in \mathbf{G}_m^*$
- and  $p'_{m+1}$  forces  $\mathbf{G}_{\mathbb{P}_{i_{m+1}}} \cap N^*$  and
- forces  $Z_n = Y_n \cap \mathcal{M}[\mathbf{G}_{\mathbb{P}_{i_{m+1}}}, M_m^*, y^*] \in D_{\alpha_n}(M_m^*)$ .

As the challenger does not have a winning strategy, there is a play he wins, giving  $\bigcup_{n < \omega} \mathbf{G}_n^* \cap N_0^*$  with a bound.

Continuation of the Proof: Choose also  $\langle i'_m : m < \omega \rangle \in N_0$  such that  $i_m \in N^*, i_0 = i, i_m < i_{m+1}, \sup\{i_m : m < \omega\} = \sup(N^* \cap j)$  and this time we let  $\langle \mathcal{I}'_m : m < \omega \rangle$  list the dense open subsets of  $\mathbb{P}_j$  from  $N^*$ . For  $\mathbf{m} < \omega$  let  $\mathcal{I}_{\mathbf{m}}$  be the set of the finite sequences  $\mathfrak{r}$  from  $M_4$  coding  $\langle r_{\mathfrak{r},m} : m \leq \mathbf{m} \rangle, \langle \mathbf{G}_{\mathfrak{r},m} : m \leq \mathbf{m} \rangle, \langle p_{\mathfrak{r},m} : m \leq \mathbf{m} \rangle, \langle n_{\mathfrak{r},m} : m \leq \mathbf{m} \rangle, \langle \mathbf{G}_{\mathfrak{r},m}^\ell : \ell \leq n_{\mathfrak{r},m}, m \leq \mathbf{m} \rangle, \langle Y_{\mathfrak{r},m} : m \leq \mathbf{m} \rangle$  and also  $(X_{\mathfrak{r},m}, \alpha_{\mathfrak{r},m}, \beta'_{\mathfrak{r},m}, \beta_{\mathfrak{r},m}, M_{\mathfrak{r},n}, y'_{\mathfrak{r},m}, M'_{\mathfrak{r},m}, y_{\mathfrak{r},m})$  for  $m \leq \mathbf{m}$  and  $Z_{\mathfrak{r},m}$  for  $m < \mathbf{m}$  satisfying: clauses (a)-(k),(m),(n),(p),(q) from the proof of (d)' above and:

- (\*)<sub>10</sub> (l)'  $p_m \in_j \cap N^*$
- (o)'  $p_m \leq p_{m+1} \in \mathcal{I}'_n$
- (r)'  $r_{\mathfrak{r},m}$  is  $(N^*, \mathbf{G}_{i_m})$ -generic for  $m \leq \mathbf{m}$

modified:2015-01-27

(656) revision:2015-01-26

- {b6} (s)'  $\langle (X_{\mathfrak{r},m}, \alpha_{\mathfrak{r},m}, \beta'_{\mathfrak{r},m}, \beta_{\mathfrak{r},m}, M_{\mathfrak{r},m}, Y_{\mathfrak{r},m}, M'_{\mathfrak{r},m}, Z_{\mathfrak{r},m'}) : m \leq \mathbf{m} \rangle$  belongs to  $N$  and is an initial segment of a play of the game  $\mathcal{D}'_{\alpha,\beta}(N^*, \mathbf{p})$  or just  $\mathcal{D}'_{\alpha,\beta}(N^*, N, \mathbf{p})$ , note that in the  $\mathbf{m}$ -th move the challenger has not yet chose  $Z_{\mathfrak{r},m}$ , (see clause (e) of Definition 2.2(1))
- (t)'  $Z_{\mathfrak{r},m} \subseteq Y_{m+1}$
- (u)'  $y_{\mathfrak{r},m}$  code  $p_m, \langle i_m : m < \omega \rangle$
- (v)'  $f_{i_m, i_{m+1}}(\alpha_m) \leq \beta'_m$  for  $m \leq \mathbf{m}$ .

Let  $\mathfrak{r} \triangleleft \mathfrak{q}$  has the natural meaning for  $\mathfrak{r} \in \mathcal{T}_{\mathbf{m}_1}, \mathfrak{q} \in \mathcal{T}_{\mathbf{m}_2}, \mathbf{m}_1 < \mathbf{m}_2$ .

Note

- ☒<sub>1</sub>  $\mathcal{T}_{\mathbf{m}} \subseteq N$  for  $\mathbf{m} < \omega$
- ☒<sub>2</sub>  $\mathcal{T}_0 \neq \emptyset$
- ☒<sub>3</sub> if  $\mathbf{x} \in \mathcal{T}_{\mathbf{m}}$ , then considering the game  $\mathcal{D}_{\alpha,\beta}(N^*, \mathbf{p})$ ,  $\mathbf{x}$  is an initial segment of a play of it (see clause (s)' above).

Assume

- (\*)<sub>11</sub>  $M'_{\mathbf{m}} \in Y_{\mathfrak{r},\mathbf{m}} \cap \mathcal{E}_{\alpha_{\mathbf{m}}} \cap (M_{\mathfrak{r},\mathbf{m}} \cup \{M_{\mathfrak{r},\mathbf{m}} \cap \mathcal{H}(\chi_{\mathbf{m}}^{\mathbf{p}})\})$  satisfying  $y_{\mathbf{m}}, y'_{\mathbf{m}} \in M'_{\mathbf{m}}$  and  $Z_{\mathbf{m}} \subseteq \mathcal{D}_{\alpha_{\mathfrak{r},\mathbf{m}}}(M'_{\mathbf{m}}), Z_{\mathbf{m}} \subseteq Y_{\mathbf{m}}$  (hence  $Z_{\mathbf{m}} \subseteq X_{\mathfrak{r},\mathbf{m}}$ ) and  $X_{\mathbf{m}+1} \in D_{\beta}(N^*) \cap X_{\mathfrak{r},\mathbf{m}}$  such that  $Z_{\mathbf{m}} \subseteq X_{\mathbf{m}+1}$  and  $\alpha_{\mathbf{m}+1} \in \alpha \cap N^*$  and any  $\beta'_{\mathbf{m}+1} \in \beta \cap N^* \setminus p_{i_{\mathbf{m}+1}, i_{\mathbf{m}+1}}(\alpha_{\mathbf{m}+1}), y'_{\mathbf{m}+1} \in N \cap \mathcal{H}(\chi_{\alpha_{\mathbf{m}+1}}^{\mathbf{p}})$  and any  $\beta_{\mathbf{m}} \in \beta \cap N \setminus \beta'_n \setminus \alpha_n$  and  $M_{\mathbf{m}+1} \in X_{\mathbf{m}+1} \cap \mathcal{E}_{\beta_{\mathbf{m}+1}}, y_{\mathbf{m}+1} \in M_{\mathbf{m}+1} \cap \mathcal{H}(\chi_{\alpha_{\mathbf{m}}}^{\mathbf{p}})$  satisfying  $y'_{\mathbf{m}+1} \in M_{\mathbf{m}+1}$  and  $Y_{\mathbf{m}+1} \in N \cap D_{X_{\mathbf{m}+1}}(M_{\mathbf{m}+1})$  and any  $M'_{\mathbf{m}+1} \in Y_{\mathbf{m}+1} \cap \mathcal{E}_{\alpha_n} \cap (M_{\mathbf{m}+1} \cup \{M_{\mathbf{m}+1} \cap \mathcal{H}(\chi_{\alpha_{\mathbf{m}}}^{\mathbf{p}})\})$  satisfying  $y_{\mathbf{m}+1}, y'_{\mathbf{m}+1} \in M'_{\mathbf{m}+1}$ .

Then

- (\*\*) there is  $\mathfrak{q} \in \mathcal{T}_{\mathbf{m}+1}$  such that  $\mathfrak{r} \triangleleft \mathfrak{q}$  and  $(Z_{\mathfrak{q},\mathbf{m}}, X_{\mathfrak{q},\mathbf{m}+1}, \alpha_{\mathfrak{q},\mathbf{m}+1}, \beta'_{\mathfrak{q},\mathbf{m}+1}, y'_{\mathfrak{q},\mathbf{m}+1}, \beta_{\mathfrak{q},\mathbf{m}+1}, y_{\mathfrak{q},\mathbf{m}+1}, M_{\mathfrak{q},\mathbf{m}+1}, Y_{\mathfrak{q},\mathbf{m}+1}, M'_{\mathfrak{q},\mathbf{m}+1})$  is equal to  $(Z_{\mathbf{m}}, X_{\mathbf{m}+1}, \alpha_{\mathbf{m}+1}, \beta'_{\mathbf{m}+1}, y'_{\mathbf{m}+1}, \beta_{\mathbf{m}+1}, y_{\mathbf{m}+1}, M_{\mathbf{m}+1}, Y_{\mathbf{m}+1}, M'_{\mathbf{m}+1})$ .

Why? Because  $f_{i_m, i_{m+1}}(\alpha_m) \leq \beta_{\mathbf{m}}$ , hence we know that  $\mathbb{P}_{i_{m+1}}/\mathbb{P}_{i_m}$  is  $(\mathbf{p}, \alpha_n, \beta_n)$ -proper and let  $\mathbf{G}_{i_m} \subseteq \mathbb{P}_{i_m}$  be generic over  $\mathbf{V}, r_{\mathbf{m}} \in \mathbf{G}_{i_m}$  to the model  $M'_{\mathfrak{r},\mathbf{m}}$  and the set  $Y_{\mathfrak{r},\mathbf{m}}$ . So now we can describe a strategy for the challenger in the game  $\mathcal{D}_{\alpha,\beta}(N^*, \mathbf{p})$  (or  $\mathcal{D}'_{\alpha,\beta}(N^*, N, \mathbf{p})$ ) delaying his choice of  $M'_m, Z_m$  to the  $(m+1)$ -th move, he just chose on the side  $\mathfrak{r}_m \in \mathcal{T}_m$  which “code” they play so far, and preserve  $\mathfrak{r}_m \triangleleft \mathfrak{r}_{m+1}$ .

By ☒<sub>3</sub> this is O.K. - all possible choices of the chooser are allowed, that is this gives a well defined strategy for the challenger (will he have some free choice, this does not hurt). But the chosen does not lose the game, so there is such a choice  $\langle \mathfrak{r}_m : m < \omega \rangle$  with  $\cup \{(M'_{\mathfrak{r}_{m+1}, \mathbf{m}}) \cup Y_{\mathfrak{r}_{m+1}, \mathbf{m}} : m < \omega\} \in D_{\alpha}(N)$  and  $\bigcup_{m < \omega} r_m$  is as required.

{c9} The adaptation for the proof of 3.4 when  $\kappa = 2$  should be clear.  
{d20}

{c9} **Claim 4.6.** Like 3.4 but  $\kappa = 2$ .

{d9} *Proof.* Similar to the proof of 4.3, with some changes. We choose  $\langle j_n : n < \omega \rangle \in M_0$  such that  $j < j_n < j_{n+1} < h^*(j, i), h^*(j_n, i_m) < j_{n+1}$ :

- (l)  $n_m$  is a power of 2, say  $2^{n_m^*}$  and so we can rename  $\{\mathbf{G}_m^\ell : \ell < n_m\}$  as  $\{\mathbf{G}_m^\eta : \eta \in 2^{n_m^*}\}$
- (m) (α)  $M_\eta \in M_1$  for  $\eta \in (n_m^* \geq) 2$
- (β)  $M_{<>} = N_0$
- (γ)  $M_\eta \in M_{\eta \hat{<} 0} \cap M_{\eta \hat{<} 1}$
- (δ)  $\eta \triangleleft \nu_1 \in n_m^* 2, \eta \triangleleft \nu_2 \in n_m^* 2 \Rightarrow \mathbf{G}_m^{\nu_1} \cap M_\eta = \mathbf{G}_m^{\nu_2} \cap M_\eta$  so we call it  $K_m^\eta$
- (ε)  $M_{\eta \hat{<} 0} = M_{\eta \hat{<} 1}$  when  $\eta \in n_m^* 2$  call it  $N_\eta$
- (ζ)  $N_\eta \in \mathcal{E}_{\ell g(\eta)}$  for  $\eta \in (m_m^* >) 2$
- (η)  $Y_\eta = \mathcal{M}[K_{\eta \hat{<} 0}, N_\eta] \cap \mathcal{M}[K_{\eta \hat{<} 1}, N_\eta] \in D_{j_{\ell g(\eta)}}(N_\eta)$ .

□<sub>3.6</sub>

{d23}

**Discussion 4.7.** We may be interested in non-proper forcing, say semi-proper and UP ones (see [Sh:f, Ch.X,XI,XV] and [Sh:311]). Here the change from (reasonable) parameter  $\mathfrak{p} = \mathfrak{p}^V$  to  $\mathfrak{p}^{V[G]}$  is more serious as  $\{N \cap \chi_\alpha : N \in \mathcal{E}^{\mathfrak{p}^{V[G]}}\}$  is in general not equal to  $\{N \cap \chi_\alpha : N \in \mathcal{E}^{\mathfrak{p}}\}$ . We intend to spell it out in [Sh:311].

We may be interested in combining this work with [Sh:587], [Sh:F259]. Intend to deal with it later (see also [Sh:669]).

## § 5. PROBLEMATIC FORCING

{problematic}

**Discussion 5.1.** 1) In the examples the “barely adding reals but not so” may occur on some stationary  $\mathcal{E} \subseteq [\chi^*]^{N_0}$  and otherwise just properness is asked. We do not bother to do this in the examples.

2) We may like to put the present lemmas and [Sh:f, Ch.XVIII,§2] and more together. The way is clear, we concentrate on  $\kappa \in N_0$ .

So instead of a “tower” with six levels we have one with  $n(*) + 1$  levels.

{e6}

**Definition 5.2.** 1) We say  $\bar{\mathbb{Q}}$  is an  $\mathfrak{p} - NNR_{N_0, \bar{\text{Pr}}, \bar{\Xi}}^0$ -iteration for  $\mathfrak{p}$  when:

- (a)  $\bar{\mathbb{Q}}$  is a CS iteration of proper notions forcing,  $\mathfrak{p}$ -proper
- (b) forcing with  $\text{Lim}(\bar{\mathbb{Q}}) = \mathbb{P}_{\ell g(\bar{\mathbb{Q}})}$  does not add reals
- (c)  $\bar{\text{Pr}} = \langle \text{Pr}_\ell : \ell < n(*) \rangle, \bar{\Xi} = \langle \Xi_\ell : \ell \leq n(*) \rangle, \Xi_\ell \subseteq \ell g(\mathfrak{p}) \times \ell g(\mathfrak{p})$
- (d)  $\text{Pr}_\ell(N, \bar{\mathbf{G}}, \mathbb{P})$  implies:  $\mathbb{P}$  a forcing notion,  $N$  is a countable elementary submodel of  $(\mathcal{H}(\chi), \in), \bar{\mathbf{G}} = \langle \mathbf{G}_m : m < k \rangle, k < \omega, \mathbf{G}_m \subseteq N \cap \mathbb{P}$  is generic over  $N$
- (e) if  $\ell = 0, \text{Pr}_\ell(N, \bar{\mathbf{G}}, \mathbb{P})$ , then  $\bigwedge_\ell \mathbf{G}_\ell = \mathbf{G}_0$
- (f) if  $\ell = n(*) - 1$ , then  $\text{Pr}_\ell(N, \bar{\mathbf{G}}, \mathbb{P})$ , iff the demand in clause (d) holds
- (g) if  $\text{Pr}_\ell(N, \bar{\mathbf{G}}, \mathbb{P})$  and  $\bar{\mathbf{G}} \triangleleft \bar{\mathbf{G}}', \ell g(\bar{\mathbf{G}}')$  finite and  $\text{Rang}(\bar{\mathbf{G}}) = \text{Rang}(\bar{\mathbf{G}}')$  then  $\text{Pr}_\ell(N, \bar{\mathbf{G}}', \mathbb{P})$
- (h)  $\text{Pr}_0(N, \bar{\mathbf{G}}, \mathbb{P})$  holds iff (the condition in clause (c) and)  $\bigwedge_\ell \mathbf{G}_\ell = \mathbf{G}_0$
- (i) if  $\ell < n(*), i < j \leq \ell g(\bar{\mathbb{Q}}), N_0 \prec N_1 \prec (\mathcal{H}(\chi), \in), \mathfrak{p} \in N_0, N_0, N_1$  are countable and for some  $(\alpha, \beta) \in \Xi_\ell \cap N_0$  we have  $N_0 \cap \mathcal{H}(\chi_\alpha^{\mathfrak{p}}) \in \mathcal{E}_\alpha, N_1 \cap \mathcal{H}(\chi_\beta^{\mathfrak{p}}) \in \mathcal{E}_\beta, \text{Pr}_\ell(N_0, \bar{\mathbf{G}}, \mathbb{P}_i), \text{Pr}_{\ell+1}(N_1, \bar{H}, \mathbb{P}_i), \ell g(\bar{\mathbf{G}}) = \ell g(\bar{H}), \mathbf{G}_k \subseteq H_k, \bar{p} = \langle p_k : k < \ell g(\bar{\mathbf{G}}) \rangle, p_\ell \in N_0 \cap \mathbb{P}_j, p_k \upharpoonright i \in \mathbf{G}_k, \ell = 0 \Rightarrow p_k = p_0$ , then we can find  $\bar{\mathbf{G}}^+ = \langle \mathbf{G}_k^+ : k < \ell g(\bar{\mathbf{G}}) \rangle \in N_1$  such that  $\text{Pr}(N_0, \bar{\mathbf{G}}^+, \mathbb{P}_j)$  and  $p_k \in \mathbf{G}_k^+$
- (j) assume  $\text{Pr}_\ell(N, \bar{\mathbf{G}}, \mathbb{P}_i), i_n < i_{n+1}, i_n \in N, i_0 = i, \sup_{n < \omega} i_n = \sup(N \cap j), (\alpha, \beta) \in N \cap \Xi_\ell, N' = N \cap \mathcal{H}(\chi_\alpha^{\mathfrak{p}}) \in \mathcal{E}_\alpha$ , then in the following game the  $\text{Pr}_\ell^+$ -player has a winning strategy (or just does not lose).

Before the  $n$ -th move  $\bar{\mathbf{G}}^n$  such that  $\text{Pr}_\ell(N, \bar{\mathbf{G}}^n, \mathbb{P}_{i_n})$  is chosen  $\ell g(\bar{\mathbf{G}}^n) = \ell g(\bar{\mathbb{Q}})$  with  $\bar{\mathbf{G}}^0 = \bar{\mathbf{G}}$ . In the  $n$ -th move the challenger chooses  $\bar{p}^n = \langle p_k^n : k < \ell g(\bar{\mathbf{G}}) \rangle, p_k^n \in N_0 \cap \mathbb{P}_{i_{n+1}}, p_k^n \upharpoonright i_n \in \mathbf{G}_k^n, \ell = 0 \Rightarrow p_k^n = p_0^n$  and then the chooser chooses  $\bar{\mathbf{G}}^{n+1}$  is above such that  $p_k^n \in \mathbf{G}_k^{n+1}$ . In the end of the play the chooser wins if  $\text{Pr}_\ell(N, \bar{\mathbf{G}}^*, \mathbb{P}_j)$  where  $\bar{\mathbf{G}}_k^* = \{p \in N \cap \mathbb{P}_j : \text{for every } n, p \upharpoonright i_n \in \mathbf{G}_k^n\}$ .



§ 6. SPELLING OUT THE AXIOMS

{spelling}

As is well known Iteration Theorems given consistency of axioms. So we should consider  $\kappa \in \{2, \aleph_0\}$ , (could also  $\kappa = \aleph_1$ ) and iterations as in §1 or as in §2 and §4.

§ 6(A). Axioms from the Results of §1.

{f2}

**Conclusion 6.1.** *If (A) then (B) where:*

- (A)  $(\mathbb{P}, \lambda)$  is an NNR-context meaning
  - (a) CH
  - (b)  $\lambda = \lambda^{<\lambda} \gg \aleph_1 \geq \kappa \geq 2$
  - (c)  $\mathfrak{p}$  is a reasonable parameter, see Definition 3.1 such that  $\lambda < \kappa_0^{\mathfrak{p}}$  and  $\lambda \leq \text{cf}(\ell g(\mathfrak{p}))$  {c3}
- (B) (a)  $\mathbb{P}_*$  is a proper  $\aleph_2$ -c.c. forcing notion of cardinality  $\aleph_2$  not adding reals
- (b)  $\mathbb{P}_* = \text{Lim}(\bar{\mathbb{Q}}, \bar{\mathbb{Q}}_*)$  a CS iteration  $\langle \mathbb{P}_\alpha, \mathbb{Q}_\alpha : \alpha < \lambda \rangle, \Vdash_{\mathbb{P}_\alpha} “|\mathbb{Q}_\alpha| < \lambda”$  such that  $\bar{\mathbb{Q}} \upharpoonright \alpha \in \mathcal{H}(\lambda)$  for  $\alpha < \lambda$ , so  $\mathbb{P} = \bigcup_{\alpha < \lambda} \mathbb{P}_\alpha$
- (c)  $\bar{\mathbb{Q}}_*$  is  $\mathfrak{p}$ -NNR $_*$ -iteration (see Definition 4.1) {d3}
- (d) if  $\mathbf{I}$  is a dense open subset of  $\mathbb{R} = \mathbb{R}_{\lambda, \mathfrak{p}} := (\{\bar{\mathbb{Q}}' : \bar{\mathbb{Q}}' \in H(\lambda) \text{ as in clause (b), } <_{\mathbb{R}}\} \text{ where } \bar{\mathbb{Q}}^1 \leq_{\mathbb{R}} \bar{\mathbb{Q}}^2 \Leftrightarrow \bar{\mathbb{Q}}^1 = \bar{\mathbb{Q}}^2 \upharpoonright \ell g(\bar{\mathbb{Q}}^1) \text{ and } \mathbf{I} \text{ is definable in } (\mathcal{H}(\lambda), \in) \text{ from a parameter, then } \lambda = \sup\{\alpha < \lambda : \bar{\mathbb{Q}} \upharpoonright \alpha \in \mathbf{I}\}$
- (e) if  $\mathbb{Q} \in \mathbf{V}^{\mathbb{P}_*}$  a forcing notion of cardinality  $\aleph_1$  so without loss of generality set of elements  $\subseteq \omega_1$  and  $\in \mathbf{V}^{\mathbb{P}_\alpha}$  for some  $\alpha < \lambda$  and  $\mathbb{Q}$  is absolute  $(\mathfrak{p} \upharpoonright \alpha, \lambda)$ -proper, see 6.6 below then for some  $\beta \in (\alpha, \lambda)$ ,  $\mathbb{Q}_\beta = \mathbb{Q}$  hence  $\mathbf{V}^{\mathbb{P}} \models \text{Ax}_\lambda(\mathbb{Q})$  {f13}
- (f) similarly for  $(\mathbb{Q}, \bar{\mathcal{I}})$ .

*Proof.* By 1.10 every  $\leq_*$ -increasing sequence in  $\mathbb{R}$  of length  $< \lambda$  has a  $\leq_*$ -lub. We also need Observation 6.2 below to note the proof is obvious. {a27} □ {f5} {f5}

**Observation 6.2.** *In 6.1,  $\mathbb{R}$  is non-empty and with no maximal member.* {f2}

*Proof.* The iteration of length zero belongs to  $\mathbb{R}$  and if  $\bar{\mathbb{Q}} = \langle \mathbb{P}_\beta, \mathbb{Q}_\beta : \beta < \alpha \rangle$  we can define  $\bar{\mathbb{Q}}' \in \mathbb{R}$  above it by letting  $\mathbb{Q}_\gamma = (\omega_1^{>2}, <)$ . □<sub>6.2</sub> {f7}

**Definition 6.3.** Assume  $\lambda = \lambda^{<\lambda}$  and  $\alpha < \lambda \Rightarrow |\alpha|^{\aleph_1} < \lambda$ . We define the  $(\lambda, 1)$ -standard  $\mathfrak{p} = \mathfrak{p}_\lambda$  by:

- $\ell g(\mathfrak{p}) = \lambda$
- $\chi_\alpha^{\mathfrak{p}} = \lambda$
- $R_\alpha^{\mathfrak{p}} = \emptyset$
- $\bar{D}$  is standard, see 1.3(1). {a9} {f9}

**Claim 6.4.** *For  $\lambda$  as in 6.3,  $\mathfrak{p}_\lambda$  is a standard parameter and  $(\mathfrak{p}_\lambda, \lambda)$  is an NNR $_\kappa$ -content, i.e.  $(\lambda, \mathfrak{p}_\lambda)$  satisfies clause (A) of conclusion 6.1.* {f7} {f2}

modified:2015-01-27 (656) revision:2015-01-26

**Definition 6.5.** Let  $(\lambda, \kappa, \mathfrak{p}, \mathbb{R})$  be as in 6.1(A).

{f11}

{f2}

1) We say  $\mathbb{Q}$  is absolutely  $(\lambda, \mathfrak{p}, \mathbb{R})$ -NNR $_{\kappa}^0$  forcing above  $\bar{\mathbb{Q}}$  when:

- (a)  $\bar{\mathbb{Q}} \in \mathbb{R}$
- (b)  $\mathbb{Q}$  is a  $\text{Lim}(\bar{\mathbb{Q}})$ -name of a forcing notion from  $\mathcal{H}(\lambda)^{\mathbf{V}[\text{Lim}(\bar{\mathbb{Q}})]}$  temporarily  $(\bar{\mathbb{Q}}', \mathfrak{p})$ -NNR $_{\kappa}^0$
- (c) if  $\bar{\mathbb{Q}} \leq_{\mathbb{R}} \bar{\mathbb{Q}}^1$  then  $\mathbb{Q}$  is (or  $(\bar{\mathbb{Q}}', \bar{\mathbb{Q}}, \mathfrak{p})$ -NNR $_{\kappa}^0$ ) which means that there is  $\bar{\mathbb{Q}}^2 \in \mathbb{R}, \bar{\mathbb{Q}}^1 \leq_{\mathbb{R}} \bar{\mathbb{Q}}^2$  and  $\bar{\mathbb{Q}}_{\ell g(\bar{\mathbb{Q}}^1)}^2 = \mathbb{Q}$  so  $\mathbb{Q}$  is a  $\text{Lim}(\bar{\mathbb{Q}}^1)$ -name  $\in \mathcal{H}(\lambda)$ .

2) We say  $(\mathbb{Q}, \bar{\mathcal{I}})$  is an absolute  $(\lambda, \mathfrak{p}, \mathbb{R})$ -NNR $_{\kappa}^0$ -problem above  $\bar{\mathbb{Q}}$  when:

- (a)  $\mathbb{Q}$  is a  $\text{Lim}(\bar{\mathbb{Q}})$ -name of a forcing notion from  $\mathcal{H}(\lambda)^{\mathbf{V}[\text{Lim}(\bar{\mathbb{Q}})]}$
- (b)  $\bar{\mathcal{I}}$  is a sequence of  $< \lambda$  subsets of  $\mathbb{Q}$
- (c) if  $\bar{\mathbb{Q}} \leq_{\mathbb{R}} \bar{\mathbb{Q}}^1$  then there is  $\bar{\mathbb{Q}}^2$  such that  $\bar{\mathbb{Q}}^1 \leq_{\mathbb{R}} \bar{\mathbb{Q}}^2$  and  $\Vdash_{\text{Lim}(\bar{\mathbb{Q}}^2)} \text{“}(\mathbb{Q}, \bar{\mathcal{I}}) \text{ is solved meaning there is a directed } \mathbf{G} \subseteq \mathbb{Q} \text{ meeting } I_{\varepsilon} \text{ for every } \varepsilon < \ell g(\bar{\mathcal{I}})\text{”}$ .

It is natural to restrict ourselves to the linear case, but this is not a real difference when we allow to change the  $f$  a little.

{f13}

**Definition 6.6.** Let  $\mathfrak{p}$  be a reasonable parameter.

1) We say  $\mathfrak{p}$  is linear when: if  $\alpha < \ell g(\mathfrak{p}), N \in \mathcal{E}_{\alpha}$  and  $y \in D_{\alpha}^{\mathfrak{p}}(N)$  then there is  $Z$  such that

- (a)  $Z \in D_{\alpha}^{\mathfrak{p}}(N)$
- (b)  $Z \subseteq Y$
- (c) if  $a \in Z \cap \mathcal{E}_{\gamma}$  then  $N \upharpoonright a \prec N \upharpoonright \mathcal{H}(\chi_{\partial})$
- (d)  $Z$  is linear meaning
  - ( $\alpha$ )  $Z$  is well ordered by  $\in$  (and by  $\subseteq$ )
  - ( $\beta$ ) if  $a \in Z$  then  $Z \cap a \in N$  and  $\langle N \upharpoonright a : a \in Z \rangle$  is  $\subseteq$ -increasing continuous.

{a9} 2) We say  $\bar{D} = \mathcal{D}_{\mathfrak{p}}$  is linearly standard when we combine part (1) with Definition 1.3(1).

3) Assume  $\mathfrak{p}$  is a reasonable parameter,  $f \in \mathcal{F}^{\mathfrak{p}}$  and  $g \in \mathcal{F}^{\mathfrak{p}}$  is defined in  $g(\alpha) = \cup \{f(\beta) : \beta \in f(\alpha)\}$ . Let  $gq := \mathfrak{p}^{[f]}$  be defined by  $D_{\alpha}^{\mathfrak{p}}(N) := \{Y \in D_{\alpha}^{\mathfrak{p}}(N) : \text{for some } \beta \in N \cap f(\alpha) \text{ there is } Z \subseteq Y, Z \text{ linear and } Z \in N \upharpoonright \mathcal{H}(\chi_{\beta})\}$ .

{f15}

{f13}

**Claim 6.7.** If  $\mathfrak{p}$  is a reasonable parameter and  $f, g, \mathfrak{q}$  is as in 6.6, then

- (a)  $\mathfrak{q}$  is a reasonable parameter
- (b) if  $f$  is increasing continuous then so is  $g$
- (c) if  $f(\alpha) = \alpha$  then  $g(\alpha) = \alpha$  and  $D_{\alpha}^{\mathfrak{q}}(N) = D_{\alpha}^{\mathfrak{p}}(N)$ .

{f17}

**Claim 6.8.** If (A) then (B) where:

{f2}

- (A) (a)  $\mathfrak{p}, \bar{f}, \lambda, \mathbb{R}$  is as in 6.1
- (b)  $\bar{\mathbb{Q}} \in \mathbb{R}, \mathbb{P} = \text{Lim}(\bar{\mathbb{Q}})$
- (c)  $\mathfrak{p}$  is linear
- (B) (a) if  $\mathbb{Q}$  is a  $\mathbb{P}$ -name of a  $(<^+, \omega_1)$ -proper forcing notion from  $\mathcal{H}(\lambda)$  and  $f \in \mathcal{F}_{cc}^{\mathfrak{p}}$  then  $\Vdash_{\mathbb{P}} \text{“}\mathbb{Q} \text{ is } (\mathfrak{p}^{\mathbb{P}}, \alpha, f)\text{-proper”}$

- {b37} (b) if  $\mathbb{Q}$  satisfies  $\kappa$ -completeness system  $\mathbb{D} \in \mathbf{V}$  over  $\mathbb{P}$  then for some  $f \in \mathcal{F}_{cc}^{\mathbb{P}}$ , we have  $(\mathbb{P}, \mathbb{Q})$  is  $(\kappa, f)$ -anti w.d., see Definition 2.12
- (c) so if the pair  $(\mathbb{P}, \mathbb{Q})$  is as in [Sh:f, Ch.V,x.x] for  $\kappa = \aleph_1$  or [Sh:f, Ch.V,y.y] for  $\kappa = \aleph_0$  or in [Sh:f, Ch.VIII,4.x] for  $\kappa = 2$  then  $\mathbb{Q}$  is temporarily  $(\mathfrak{p}, \mathbb{Q})$ -NNR $_{\kappa}^1$ , (i.e. can be chosen in the  $\mathbb{Q}_{\ell g(\bar{\theta})}$ ).

*Proof.* FILL. □<sub>6.8</sub>

**Claim 6.9.** 1) For  $\mathfrak{p}, \lambda, \mathbb{R}, \bar{Q} = \bar{Q}_* \in R_{\lambda}, \mathbb{P} = \mathbb{P}_* = \text{Lim}(\bar{Q}_*)$  and in 6.1 the following  $\mathbf{V}_1 = \mathbf{V}^{\mathbb{P}_*}, \text{Ax}_{\lambda}(Q)$  holds for  $Q \in \mathbf{V}_1$  of the following for ? {f22}

2) The following  $\mathbb{P}$ -names  $\mathbb{Q}(\mathbf{I})$  fall under clause (B)(d) there (moreover if just  $\bar{Q} \in \mathbb{R} = \mathbb{R}_{<\lambda}, \mathbb{P} = \text{Lim}(\bar{Q})$  this holds) {f2}

- (a)  $\mathbb{Q}$  is as in 3.1 so depends on  $\{(C_{\delta}, n_{\delta}) : \delta < \omega_1 \text{ limit}\}$  which is a subset of  $\mathcal{H}(\aleph_1)$  {c3}
- (b)  $\mathbb{Q}$  is as in 3.4 for  $\bar{C} = \langle C_{\delta} : \delta < \omega_1 \text{ limit} \rangle, \text{otp}(C_{\delta}) = 4$  {c9}
- (c)  $\mathbb{Q}$  is as in 3.4 for  $\bar{C} = \langle C_{\delta} : \delta < \omega_1 \text{ limit} \rangle$  and for some countable  $\gamma(*), \delta < \omega_1 \Rightarrow \text{otp}(C_{\delta}) \leq \omega^{\gamma(*)}, \gamma(*) < \omega_1$ . {c9}

*Proof.* The  $(\mathfrak{p}, f)$ -properness is proved in §3. The  $(\mathfrak{p}, g)$ -anti w.d. is straight (see §0). □<sub>6.9</sub>

**Claim 6.10.** In 6.9: 1) If  $\lambda$  is strongly inaccessible<sup>5</sup> then every Aron tree is special. {f25}

2) If  $\mathbb{T}$  is a  $\mathbb{P}$ -name of an Aron tree then the pair  $(\mathbb{Q}_{\mathbb{T}}, \bar{\mathcal{I}})$  falls under - FILL. {f22}

*Proof.* For a  $\aleph_1$ -tree  $T$  let

- (\*)<sub>1</sub> (a)  $\mathbb{Q}_T = \{f : \text{for some } \alpha < \omega_1, f \text{ is a function from } T_{\leq \alpha} \text{ to } \omega \text{ such that } f(s) = f(t) \Rightarrow s \not\prec_T t\}$
- (b)  $\leq_{\mathbb{Q}_T}$  is unclear?
- (c)  $\bar{\mathcal{I}}_T = \langle \mathcal{I}_{T,\alpha} : \alpha < \omega_1 \rangle$  where  $\mathcal{I}_T = \{f \in \mathbb{Q}_T : T_{\leq \alpha} \subseteq \text{Dom}(f)\}$ .

Clearly

- (\*)<sub>2</sub> (a)  $(\mathbb{Q}_T, \bar{\mathcal{I}})$  is a problem
- (b) this problem has a solution iff  $T$  is a special Aronszajn Tree.

The rest should be clear. □<sub>6.10</sub>

<sup>5</sup>We may avoid this if we use iterations as in [Sh:f, Ch.VIII], i.e.  $\bar{Q} \in \mathbb{R}$  is only a class of  $(\mathcal{H}(\lambda), \in)$ , satisfying a strong version of  $\lambda_2$ -c.c., so  $\lambda = \aleph_2$  is O.K.

§ 7. PRIVATE APPENDIX  
MOORE QUESTION

{f28} We turn to the first Moore test

**Definition 7.1.** 1) Let  $\mathbf{cd}: \mathcal{H}(\aleph_1) \rightarrow \omega_1$  be one-to-one.

We say  $E$  solves  $\mathbf{cd}$  when  $E$  is a club of  $\omega_1$  and for every  $\alpha \in E$  we have  $\mathbf{cd}(e \cap (\alpha + 1)) < \min(E \setminus (\alpha + 1))$ .

1A) The statement  $JM_1$  (or  $(D)$ ) say: for every one-to-one function  $\mathbf{cd}$  from  $\mathcal{H}(\aleph_1)$  to  $\omega_1$ , some  $E$  solves it.

2) For  $\mathbf{cd}$  as above we define  $\mathbb{Q} = \mathbb{Q}_{\mathbf{cd}}$  as the following forcing notions:

(a)  $p \in \mathbb{Q}$  iff  $p$  is a closed bounded subset of  $\omega_1$  satisfying  $(\forall \alpha \in p)[\alpha \neq \max(p) \Rightarrow \mathbf{cd}(p \cap (\alpha + 1)) < \min(p \setminus (\alpha + 1))]$

{f31} (b)  $p \leq_{\mathbb{Q}} q$  iff  $p, q \in \mathbb{Q}$  and  $p$  is an initial segment of  $q$ .

*Question 7.2.* Moore first test says: is it consistent that

(a) CH

(b)  $JM_1$

(c) if  $\bar{C} = \langle C_\delta : \delta < \omega_1 \text{ limit} \rangle, C_\delta \subseteq \delta = \sup(C_\delta), \text{otp}(C_\delta) = \omega$  then for some club  $E$  of  $\omega_1, (\forall \delta)(\delta > \sup(C_\delta \cap E))$ .

{f33}  
{f2}

**Claim 7.3.** In 6.1,  $\mathbf{V}_1 = \mathbf{V}^{\mathbb{P}}$  satisfies in  $\mathbf{V}_1$  the first Moore question.

{f22} *Proof.* In 7.1, Clause (a) holds as 6.1(A)(a) says CH holds in  $\mathbf{V}$  and the forcing  
{f22} notion adds no new real. Clause (c) of 7.1 holds by 6.9(b) recalling 3.4.

{f36} For clause (b) use 7.4 below. □<sub>7.3</sub>

Note that  $\mathbb{Q}_{\mathbf{cd}}$  was chosen as a very “low”, “simple”, forcing as in [Sh:f, Ch.V] and  $\bar{Q}_{\bar{C}}, \bigwedge_{\delta} \text{otp}(C_\delta) > \omega$  as very “low”, “simple” forcing as in [Sh:f, Ch.XVIII, §1].

{f36}  
{f22}

**Claim 7.4.** 1) In 6.9 we could add  $\mathbb{Q}_{\mathbf{cd}}$  for  $\mathbf{cd}$  as above.

2)  $\mathbb{Q}_{\mathbf{cd}}$  is  $(<^+, \omega_1)$ -proper.

{f40} *Proof.* Straightforward. □<sub>7.3</sub>

**Definition 7.5.** 1) Let  $\mathbf{C}_{\mathbf{cd}} = \{\bar{C} : \bar{C} = \langle C_\delta : \delta < \omega_1 \text{ limit} \rangle, C_\delta \text{ a closed unbounded subset of } \delta\}$ .

2) Let  $\mathbf{C}_\gamma = \{\bar{C} \in \mathbf{C} : \text{otp}(C_\delta) \leq \omega^{1+\gamma} \text{ for every } \delta\}$ .

3) For  $\bar{C} \in \mathbf{C}_{\mathbf{cd}}$  let  $\mathbb{Q}_{\bar{C}}^2$  be the following forcing notion:

(A)  $p \in \mathbb{Q}_{\bar{C}}^1$  iff for some  $\alpha < \omega_1$

(a)  $p$  is a closed subset of  $\alpha$

(b) if  $\delta \leq \alpha$  is a limit ordinal then  $\delta > \sup(p \cap C_\delta)$  or  $\delta > \sup(p \setminus C_\delta)$

(B)  $\mathbb{Q}_{\bar{C}}^1 \models “p \leq q”$  is an initial segment of  $q$ .

{f42}  
{f40}

**Claim 7.6.** 1) In 7.5,  $\mathbb{Q} = \mathbb{Q}_{\bar{C}}^1$  is  $(\mathfrak{p}, f)$ -proper when: FILL. (consider moving!)

*Proof.* We prove it by induction on  $\alpha$  that

⊕ if (a) then (b) where

(a)  $\beta \in f(\alpha), N \in \mathcal{E}_\beta, \{\bar{C}, \alpha\} \subseteq N, Y \in D_\beta^{\mathfrak{p}}(N)$  and  $p \in N \cap \mathbb{Q}_{\bar{C}}^2, \delta = N \cap \omega_1$

- (b) there are  $q, Z, \mathbf{G}$  such that
  - ( $\alpha$ )  $q$  is  $(N, Q)$ -generic and forces  $\mathbf{G}_Q \cap N = \mathbf{G}$
  - ( $\beta$ )  $Q \Vdash "p \leq q"$ , equivalently  $p \in G$
  - ( $\gamma$ )  $q \setminus p$  is included in  $C_\delta$  or is disjoint to  $C_\delta$
  - ( $\delta$ )  $Z \in D_\alpha(N)$  and  $Z \subseteq Y$
  - ( $\varepsilon$ ) if  $M \in Z$  then  $q$  is  $(M, Q)$ -generic.

□<sub>7.6</sub>

{f43}

**Claim 7.7.** [CH] *If  $\mathbb{P}$  is proper NNR forcing and  $\Vdash_{\mathbb{P}} "\bar{C} \in \mathbf{C}_{cd}"$  then the pair  $(\mathbb{P}, \mathbb{Q}_{\bar{C}}^2)$  satisfies a simple  $\aleph_1$ -completeness system (hence for every non-trivial  $\mathfrak{p}$ , see Definition xxx - FILL).*

*Proof.* We define an  $\aleph_2$ -completeness system  $\bar{\mathbb{D}}$ .

Let  $\bar{\mathbb{D}} = \langle \mathbb{D}_N : N \in \mathbf{S} \rangle$  where

- $\mathbf{S} = \{N : N = (M_N, \bar{C}_N, Q_N, P_N) : \text{for some } \bar{C}' \in \mathbf{C}_{cd}, p' \in Q_{\bar{C}}^2 \cap M' \text{ and } \chi, M \prec (\mathcal{H}(\chi), \varepsilon) \text{ is countable, } \bar{C}' \in M \text{ and } (N, \bar{C}', Q_{\bar{C}'}, P) \text{ is the Mostowski collapse of } (M', \bar{C}', Q_{\bar{C}'}, P')\}$
- for  $N = (M, \bar{C}, Q) \in \mathbf{S}$  and  $p \in Q$  we let  $\text{gen}(N) = \{\mathbf{G} : \mathbf{G} \text{ a subset of } Q_N \text{ generic over } M_N \text{ to which } \mathbb{D}_N = \{\mathcal{G} : \text{for some countable, } \mathcal{C} \subseteq \{C : C \text{ a closed unbounded subset of } \delta(N) = \omega_1^N, \mathcal{G} = \mathcal{G}_{\mathcal{C}} = \{\mathbf{G} \in \text{gen}(N) : \text{if } C \in \mathcal{C} \text{ then } (\cup \mathbf{G}) \cap C \text{ is bounded below } \delta(N) \text{ or } (\cup \mathbf{G}) \setminus C \text{ is bounded below } \delta\}\}$

So this is just a repetition of  $Q$  being  $(\mathfrak{p}, f)$ -proper, in our case  $(\mathfrak{p}, \alpha, \beta)$ -proper, except the additional clause (c).

We prove also that  $Q$  is  $(\mathfrak{p}, \text{id}_\alpha)$ -anti w.d. We are given  $N, \mathbb{P}, Q = Q_{\bar{C}}^2$ , so  $\bar{C}$  is a  $P$ -name,  $\mathbb{P}$ -proper adding no reals.

So assume  $q_\ell \in P$  is  $(N, Q)$ -generic for  $\ell < n$ , so  $n$  finite,  $p_\ell \Vdash "\mathbf{G} \cap N = \mathbf{G}_*"$  and  $C_\delta = C_{\delta, \ell}$  for  $\ell < n$  and  $q \in N, \Vdash_{\mathbb{P}}$  and we should find  $\langle q_\ell : \ell < n \rangle, \mathbf{G}_{**}$  as required in Definition ??; clearly this suffices. Also without loss of generality  $\alpha = 0 \Rightarrow n = 1$ . {fxxx}

Case 1: There is a  $\mathbb{P}$ -name  $\bar{E}, \Vdash_{\mathbb{P}} "\bar{E} \text{ is a club of } \omega_1"$  such that  $\delta > \text{cor}(Y) \cap \bar{E}[\mathbf{G}_*] \setminus C_\delta$  recalling  $\text{cor}(Y) = \{M \cap \omega_1 : M \in Y\}$ .

Let  $\alpha_* = \sup(\bar{E}[\mathbf{G}_*] \setminus C_\delta) + 1$ , so as we can restrict ourselves to  $M$ 's such that " $\{\bar{E}, \alpha_*\} \in M$ " there is no problem.

Case 2: Not case 1.

Let  $\langle \delta_k : k < \omega \rangle$  be increasing with limit  $\delta$  and if  $\alpha = 0$  let  $\bigwedge_k \alpha_k = 1$ , if  $\alpha$  successor let  $\bigwedge_k \alpha_k = \alpha - 1$  and if  $\alpha$  is a limit ordinal let  $\langle \alpha_k : k < \omega \rangle$  be an increasing sequence of members of  $N \cap \alpha$  unbounded in it. Old question: check  $Y$  □<sub>7.7</sub>

**Discussion 7.8.** Thought: on Case 1 above: if  $Y \in D_{\beta_1}(N), C \subseteq \delta_N$  closed,  $\beta_2 \in \beta_1 \cap N$  then there is  $Z \subseteq Y$  such that  $C \subseteq Z$  or  $C \cap Z = \emptyset$ .

*Question 7.9.* (2012.10.12)

- 1) Should we replace  $R_\alpha = (R_{0, \alpha}, R_{1, \alpha}), R_{0, \alpha} \in \mathcal{H}(\chi_\alpha), R_{1, \alpha} \subseteq \mathcal{H}(\chi_\alpha)$ .

Writing  $R_\alpha$  means  $R_{0,\alpha} = R_\alpha \wedge R_{1,\alpha} = \mathcal{H}(\chi_\alpha)$ . If  $\bar{\chi}$  increases fast enough,  $\bar{R}_\alpha$  can be translated;  $\chi'_\alpha = (2^{<\chi_\alpha})^+$ ,  $R'_\alpha = (r_{0,\alpha}, R_{1,\alpha}, \chi_\alpha) \in \mathcal{H}(\chi)$ . Define  $\mathfrak{p}$  is  $\gamma$ -wide when  $\bar{\mathfrak{I}}_\gamma(\chi_\alpha) < \chi_{\alpha+1}$  for every  $\alpha$ .

- {d6} 2) In ?? should not (c) we adopted to  $\Xi$ ? but by 4.2?
- 3) Is long properness interesting while adding reals?
- {d3} 4) In 4.1, §4, clause (c), change to  $\mathfrak{p}^{\mathbb{P}_i}$ , ADD:  $(\mathbb{P}, \mathbb{Q})$  satisfies.
- {e6} 5) What does §5, in particular 5.2 do?
- 6) Add explicitly: force by “ $\bar{\mathbb{Q}}$  a  $\mathfrak{p}$ -NNR $^c_\kappa$  of length  $< \lambda$ ”,  $\lambda = \lambda^{<\kappa} \wedge (\forall \alpha < \lambda)[|\alpha|^{\aleph_1} < \lambda]$  or exist?
- 7) The gain in §4 is:
  - (a) weak diamond has delayed properness incorporated
  - (b) delayed properness itself.

What about  $f_i$  depends in  $i < \text{lg}(\bar{\mathbb{Q}})$ ?

- {y3} 8) CS iteration of length  $\alpha$ :  $P_\alpha$  included: yes by 0.1.
- {d2} 9) In 1.2, 1.10 don't we mean chains of models? Saharon check.
- {d6} 10) Concerning (6), in 4.2 - not really  $f_i$  depends on  $i$ , as each gives a club of  $\text{lg}[\mathfrak{p}]$  which has cofinality  $> \text{lg}(\bar{\mathbb{Q}})$ .

#### REFERENCES

- [Sh:b] Saharon Shelah, *Proper forcing*, Lecture Notes in Mathematics, vol. 940, Springer-Verlag, Berlin-New York, xxix+496 pp, 1982.
- [Sh:f] ———, *Proper and improper forcing*, Perspectives in Mathematical Logic, Springer, 1998.
- [DvSh:65] Keith J. Devlin and Saharon Shelah, *A weak version of  $\diamond$  which follows from  $2^{\aleph_0} < 2^{\aleph_1}$* , Israel Journal of Mathematics **29** (1978), 239–247.
- [Sh:177] Saharon Shelah, *More on proper forcing*, The Journal of Symbolic Logic **49** (1984), 1034–1038.
- [Sh:F259] ———, *Not collapsing cardinals  $\leq \kappa$  in  $(< \kappa)$ -support iterations II*, moved to 667.
- [Sh:311] ———, *A more general iterable condition ensuring  $\aleph_1$  is not collapsed*, preprint, arxiv:math.LO/0404221.
- [Sh:587] ———, *Not collapsing cardinals  $\leq \kappa$  in  $(< \kappa)$ -support iterations*, Israel Journal of Mathematics **136** (2003), 29–115, arxiv:math.LO/9707225.
- [Sh:666] ———, *On what I do not understand (and have something to say:) Part I*, Fundamenta Mathematicae **166** (2000), 1–82, arxiv:math.LO/9906113.
- [Sh:669] ———, *Non-Cohen Oracle c.c.c.*, Journal of Applied Analysis **12** (2006), 1–17, arxiv:math.LO/0303294.

EINSTEIN INSTITUTE OF MATHEMATICS, EDMOND J. SAFRA CAMPUS, GIVAT RAM, THE HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM, 91904, ISRAEL, AND, DEPARTMENT OF MATHEMATICS, HILL CENTER - BUSCH CAMPUS, RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY, 110 FRELINGHUYSEN ROAD, PISCATAWAY, NJ 08854-8019 USA

*E-mail address:* shelah@math.huji.ac.il

*URL:* http://shelah.logic.at