

The failure of the uncountable non-commutative Specker Phenomenon

By

Saharon Shelah^{*} and Lutz Strüningmann[†]

Abstract

Higman proved in 1952 that every free group is non-commutatively slender, this means that, for a free group G and for a homomorphism h from the free complete product $\times_{\omega} \mathbb{Z}$ of countably many copies of \mathbb{Z} into G , there exists a finite subset $F \subseteq \omega$ and a homomorphism $\bar{h} : *_F \mathbb{Z} \rightarrow G$ such that $h = \bar{h} \rho_F$ where ρ_F is the natural map from $\times_{\omega} \mathbb{Z}$ into $*_F \mathbb{Z}$. Due to the corresponding phenomenon for abelian groups this is called the non-commutative Specker Phenomenon. In the present paper we shall show that Higman's result fails if one passes from countable to uncountable and with it answer a question posed by K. Eda. In particular, we will see that, for an uncountable cardinal λ and for non-trivial groups G_{α} ($\alpha \in \lambda$), there are $2^{2^{\lambda}}$ homomorphisms from the free complete product of the G_{α} 's into the integers.

Introduction

In 1952 Higman [H] proved that every free group G is non-commutatively slender where slenderness means that any homomorphism h from the free complete product $\times_{\omega} \mathbb{Z}$ of countably many copies of the integers into G depends on finitely many coordinates only. A similar result was proven by Specker in 1950 [S] for abelian groups. Specker showed that any homomorphism from the product $\prod_{\omega} \mathbb{Z}$ of countably many copies of \mathbb{Z} into the integers is determined by only finitely many entries. These two phenomena are called the commutative and

Mathematics Subject Classification (2000): 20E06.

^{*}Number 729 in list of publications.

Supported by GIF project No. G-0545-173.06/97 of the *German-Israeli Foundation for Scientific Research & Development*

[†]Supported by the Graduiertenkolleg *Theoretische und Experimentelle Methoden der Reinen Mathematik* of Essen University.

the non-commutative Specker Phenomenon, respectively. Eda [E1] extended Higman's result by showing that, for any non-commutatively slender group S , for any non-trivial groups G_α ($\alpha \in I$) and for any homomorphism h from the free σ -product of the G'_α 's into S , there exist a finite subset F of I and a homomorphism $\bar{h} : *_{i \in F} G_i \rightarrow S$ such that $h = \bar{h} \rho_F$ where ρ_F is the natural map from $\times_{i \in I}^\sigma G_i$ to $*_{i \in F} G_i$ (for the definition of σ -product see 1.2). Motivated by this result Eda [E1, Question 3.8] asked whether or not the non-commutative Specker Phenomenon still holds if one passes from countable to uncountable cardinals replacing $\times_\omega \mathbb{Z}$ by the free complete product $\times_\lambda \mathbb{Z}$ for some uncountable cardinal λ (see 1.2). Here we shall give a negative answer to Eda's question by constructing, for a given uncountable cardinal λ and for non-trivial groups G_α ($\alpha \in \lambda$), a homomorphism h from the free complete product of the G_α 's into \mathbb{Z} for which the non-commutative Specker Phenomenon fails. In fact, we will show that there are 2^{2^λ} of these homomorphisms and so, in particular, we have that the cardinality of the set of all homomorphisms from $\times_{\alpha \in \lambda} G_\alpha$ into the additive group of the ring of integers is the largest one possible. This contrasts the countable case and also the abelian case.

Basics and notations

Let I be an arbitrary set. For groups G_i ($i \in I$), the free product is denoted by $*_{i \in I} G_i$ (for details on free products see [M]).

Given arbitrary subsets $X \subset Y$ of I we put $\rho_{XY} : *_{i \in Y} G_i \rightarrow *_{i \in X} G_i$ to be the canonical homomorphism. Moreover, we use the notation $X \Subset I$ for finite subsets X of I . Then the set $\{*_{i \in X} G_i : X \Subset I\}$ together with the homomorphisms ρ_{XY} ($X \subset Y \Subset I$) form an inverse system; its inverse limit $\varprojlim (*_{i \in X} G_i, \rho_{XY} : X \subset Y \Subset I)$ is called the *unrestricted free product* of the G_i 's (see [H]).

Eda [E1] introduced an infinite version of free products and defined the *free complete product* $\times_{i \in I} G_i$ of the groups G_i (for the exact definition see 1.2); it is isomorphic to the subgroup $\bigcap_{F \in I} \{*_{i \in F} G_i * \varprojlim (*_{i \in X} G_i, \rho_{XY} : X \subset Y \Subset I)\}$ of the unrestricted free product.

For the convenience of the reader who is not familiar with the notion of a free complete product we recall the definition of words of infinite length and also the definition of and some basic facts about $\times_{i \in I} G_i$ as can be found in [E1].

Definition 1.1 Let G_i ($i \in I$) be non-trivial groups such that $G_i \cap G_j = \{e\}$ for $i \neq j \in I$. The elements of $\bigcup_{i \in I} G_i$ are called *letters*.

A *word* W is a function $W : \bar{W} \rightarrow \bigcup_{i \in I} G_i$ from a linearly ordered set \bar{W} into the set of all letters $\bigcup_{i \in I} G_i$ such that $W^{-1}(G_i)$ is finite for any $i \in I$. If the domain \bar{W} of the word W is countable then we say that W is a σ -word.

The class of all words is denoted by $\mathcal{W}(G_i : i \in I)$ (abbreviated by \mathcal{W}) and the class of all σ -words is denoted by $\mathcal{W}^\sigma(G_i : i \in I)$ (abbreviated by \mathcal{W}^σ).

Two words U and V are said to be *isomorphic* ($U \cong V$) if there exists an order-isomorphism $\varphi : \overline{U} \rightarrow \overline{V}$ between the linearly ordered sets \overline{U} and \overline{V} such that $U(\alpha) = V(\varphi(\alpha))$ for all $\alpha \in \overline{U}$. Identifying isomorphic words it is easily seen that \mathcal{W} is, in fact, a set. Moreover, for words of finite length (i.e. with finite domain) the above definition obviously coincides with the usual definition of words.

For a subset X of the set I the *restricted word* (or *subword*) W_X of W is given by the function $W_X : \overline{W}_X \rightarrow \bigcup_{i \in X} G_i$ with $\overline{W}_X = \{\alpha \in \overline{W} : W(\alpha) \in \bigcup_{i \in X} G_i\}$

and $W_X(\alpha) = W(\alpha)$ for all $\alpha \in \overline{W}_X$. Therefore $W_X \in \mathcal{W}$. Using restricted words with respect to finite subsets of I we define an equivalence relation on \mathcal{W} by saying that two words U and V are *equivalent* ($U \sim V$) if $U_F = V_F$ for all $F \Subset I$ where we may consider U_F and V_F as elements of the free product $*_{i \in F} G_i$. The equivalence class of a word W is denoted by $[W]$ and the composition of two words as well as the inverse of a word are defined naturally. Thus $\mathcal{W}/\sim = \{[W] : W \in \mathcal{W}\}$ together with the representative-wise defined composition form a group.

Definition 1.2 Given groups G_i ($i \in I$) the *free complete product* $\times_{i \in I} G_i$ is defined to be the group $\mathcal{W}(G_i : i \in I)/\sim$ as described above. Moreover, the *free σ -product* $\times_{i \in I}^\sigma G_i$ is the group $\mathcal{W}^\sigma(G_i : i \in I)/\sim$ which is a subgroup of $\times_{i \in I} G_i$.

If G_i is isomorphic to a fixed group G for all $i \in I$ then we write $\times_I G$ and $\times_I^\sigma G$ instead of $\times_{i \in I} G_i$ and $\times_{i \in I}^\sigma G_i$, respectively.

Note, for a finite set I , we obviously have that $\times_{i \in I} G_i$ and $\times_{i \in I}^\sigma G_i$ are isomorphic to $*_{i \in I} G_i$. In general we have, by [E1, Proposition 1.8], the free complete product $\times_{i \in I} G_i$ is isomorphic to the subgroup $\bigcap_{F \Subset I} \{ *_{i \in F} G_i * \lim_{\leftarrow} (*_{i \in X} G_i, \rho_{XY} : X \subset Y \Subset I) \}$ of the unrestricted free product. Moreover, Eda [E1] proved that each equivalence class $[W]$ is determined uniquely by a reduced word; a word $W \in \mathcal{W}(G_i : i \in I)$ is said to be *reduced* if $W \cong UXV$ implies $[X] \neq e$ for any non-empty word X , where e is the identity, and it never occurs that the letters $W(\alpha)$ and $W(\beta)$ belong to the same G_i for neighbouring elements α and β of \overline{W} .

Lemma 1.3 (Eda, [E1]) *For any word $W \in \mathcal{W}(G_i : i \in I)$ there exists a reduced word $V \in \mathcal{W}(G_i : i \in I)$ such that $[W] = [V]$ and V is unique up to isomorphism.*

Furthermore, Eda [E1] showed the following lemma where a word $W \in \mathcal{W}(G_i : i \in I)$ is called *quasi-reduced* if the reduced word of W can be obtained by

multiplying neighbouring elements without cancellation.

Lemma 1.4 (Eda, [E1]) *For any two reduced words $W, V \in \mathcal{W}(G_i : i \in I)$ there exist reduced words $V_1, W_1, M \in \mathcal{W}(G_i : i \in I)$ such that $W \cong W_1 M$, $V \cong M^{-1} V_1$ and $W_1 V_1$ is quasi-reduced.*

We would like to remark that the free σ -product $\times_I^\sigma \mathbb{Z}$ is isomorphic to the fundamental group (see [E1]) and the free complete product $\times_I \mathbb{Z}$ is isomorphic to the big fundamental group of the Hawaiian earring with I -many circles (see [CC]). Hence free complete products are also of topological interest.

The uncountable Specker Phenomenon

In 1950 E. Specker [S] proved that, for any homomorphism h from the direct product \mathbb{Z}^ω of countably many copies of \mathbb{Z} into the additive group of the ring of integers \mathbb{Z} , there exist a finite subset F of ω and a homomorphism $\bar{h} : \mathbb{Z}^F \rightarrow \mathbb{Z}$ satisfying $h = \bar{h} \rho_F$ where $\rho_F : \mathbb{Z}^\omega \rightarrow \mathbb{Z}^F$ is the canonical projection. This result is called the Specker Phenomenon. It can be easily seen that Specker's result still holds if one considers homomorphisms into any free abelian group G instead of homomorphisms into \mathbb{Z} , i.e. free abelian groups are slender. In general, an abelian group G is said to be *slender* if G satisfies the above property for any homomorphisms $h : \mathbb{Z}^\omega \rightarrow G$. For generalizations to products of uncountably many copies of \mathbb{Z} within the category of abelian groups we refer to [EM] or [F1].

In [E2] Eda introduced a non-commutative version of slenderness and that is exactly what we shall consider here.

Definition 2.1 A group G is *non-commutatively slender* if, for any homomorphism $h : \times_\omega \mathbb{Z} \rightarrow G$, there exists a natural number n such that $h(\times_{\omega \setminus \{1, \dots, n\}} \mathbb{Z}) = \{e\}$ where e denotes the identity element of G .

Eda proved that non-commutatively slender groups are torsion-free and that non-commutative slenderness for abelian groups is the same as the ordinary (commutative) slenderness (see [E1, Theorem 3.3. and Corollary 3.4.]). Moreover, he proved that non-commutatively slender groups have the following nice property:

Proposition 2.2 (Eda, [E1]) *Let G_i ($i \in I$) be non-trivial groups, let S be a non-commutatively slender group, and let $h : \times_{i \in I}^\sigma G_i \rightarrow S$ be a homomorphism. Then there exist a finite subset F of I and a homomorphism $\bar{h} : \times_{i \in F} G_i \rightarrow S$ such that $h = \bar{h} \rho_F$ where ρ_F is the canonical map from $\times_{i \in I}^\sigma G_i$ to $\times_{i \in F} G_i$.*

Another interesting result is that the restricted direct product and the free product of non-commutatively slender groups S_j ($j \in J$) are non-commutatively slender (see [E1, Theorem 3.6.]). However, the first fundamental result on the class of non-commutatively slender groups was already obtained by Higman [H] in 1952; it is stated below.

Theorem 2.3 (Higman, [H]) *Every free group is non-commutatively slender.*

Contrary to Higman's result above we will show that the non-commutative Specker Phenomenon fails if one replaces the product of countably many groups by products of uncountably many. To be more precise: we actually show that, for an uncountable cardinal λ , there are 2^{2^λ} homomorphisms from the free complete product $\times_{\alpha \in \lambda} G_\alpha$ of non-trivial groups G_α ($\alpha \in \lambda$) into the additive group of the ring of integers.

To make the proof more transparent we first construct one homomorphism for which the Specker Phenomenon fails and then modify the construction to obtain our main result.

Theorem 2.4 *Let λ be any uncountable cardinal and G_α ($\alpha \in \lambda$) non-trivial groups. Then there exists a homomorphism $\varphi : \times_{\alpha \in \lambda} G_\alpha \rightarrow \mathbb{Z}$ for which the Specker Phenomenon fails.*

Proof. Let G_α ($\alpha \in \lambda$) be a collection of non-trivial groups with identity elements e_α and choose elements $g_\alpha \neq e_\alpha$ of G_α for each $\alpha \in \lambda$. For any regular uncountable cardinal $\kappa \leq \lambda$ we define the word $M_\kappa \in \times_{\alpha \in \lambda} G_\alpha$ as follows:

$$M_\kappa : (\kappa, <) \longrightarrow \bigcup_{\alpha \in \lambda} G_\alpha \text{ via } \beta \mapsto g_\beta$$

where $<$ is the natural ordering of λ . Note that M_κ is a word of uncountable cofinality (i.e. the domain of M_κ has uncountable cofinality) since κ is regular and uncountable. For $\beta < \kappa$ we put $M_{\kappa, \beta}$ to be the subword $M_\kappa \upharpoonright_{[\beta, \kappa)}$ of M_κ . Now, let X be any reduced word in $\times_{\alpha \in \lambda} G_\alpha$ and recall that a subset $J \subseteq (\overline{X}, <)$ is called convex if $x < y < z$ and $x, z \in J$ imply $y \in J$. We define

$$Occ_\kappa^+(X) := \{J \subseteq (\overline{X}, <) : J \text{ is convex and } X \upharpoonright_J \cong M_{\kappa, \beta} \text{ for some } \beta < \kappa\}.$$

Thus $Occ_\kappa^+(X)$ counts the occurrences of end segments of M_κ in X . Similarly we let

$$Occ_\kappa^-(X) := \{J \subseteq (\overline{X}, <) : J \text{ is convex and } X \upharpoonright_J \cong M_{\kappa, \beta}^{-1} \text{ for some } \beta < \kappa\}.$$

In order to avoid counting subsets of $(\overline{X}, <)$ more often than necessary we define the following equivalence relation on $Occ_\kappa^+(X)$ and $Occ_\kappa^-(X)$:

Dealing with Occ^+ two convex subsets J_1, J_2 of $(\overline{X}, <)$ are said to be equivalent ($J_1 \sim_\kappa J_2$) if they have a common end segment; in other words $J_1 \sim_\kappa J_2$ if there exist $j_1 \in J_1, j_2 \in J_2$ such that $X \upharpoonright_{S_1} \cong X \upharpoonright_{S_2}$ where $S_i = \{j \in J_i : j \geq j_i\}$ ($i = 1, 2$). Similarly, if we deal with Occ^- , we define the equivalence relation substituting end segments by initial segments. For simplicity we denote both equivalence relations by \sim_κ but the reader should keep in mind that \sim_κ is defined differently for Occ^+ and Occ^- .

First we prove that two subsets $J_1, J_2 \in Occ_\kappa^+(X)$ are either disjoint or equivalent. To do so assume that $J_1, J_2 \in Occ_\kappa^+(X)$ are not disjoint and let $j^* \in J_1 \cap J_2$ ($\neq \emptyset$). Moreover, there are ordinals $\beta_1, \beta_2 < \kappa$ ($\leq \lambda$) and isomorphisms $h_i : M_{\kappa, \beta_i} \rightarrow X \upharpoonright_{J_i}$ ($i = 1, 2$) since J_1, J_2 are elements of $Occ_\kappa^+(X)$. Thus we can find $\gamma_i \geq \beta_i$ such that $h_i(\gamma_i) = j^*$ and therefore $X(j^*) = g_{\gamma_i}$ for $i = 1, 2$. Hence $\gamma_1 = \gamma_2$ and by transfinite induction we conclude $X \upharpoonright_{T_1} \cong X \upharpoonright_{T_2}$, where $T_i = \{j \in J_i : j \geq j^*\}$. Note that h_i is an isomorphism of linearly ordered sets and hence h_i commutes with limits and the successor-function.

Similarly, two subsets J_1, J_2 of $Occ_\kappa^-(X)$ are either disjoint or equivalent.

Next we show that the set $Occ_\kappa^+(X)/\sim_\kappa$ is finite; by similar arguments it then also follows that $Occ_\kappa^-(X)/\sim_\kappa$ is finite. Let us assume the contrary, that is, there exist infinitely many pairwise non-equivalent $J_n \in Occ_\kappa^+(X)$ ($n \in \omega$). Then J_n and J_m are disjoint for any $n \neq m$ by the above. For each $n \in \omega$ let $X \upharpoonright_{J_n} \cong M_{\kappa, \beta_n}$ for some $\beta_n < \kappa$. Thus $\beta = \bigcup_{n \in \omega} \beta_n$ is strictly less than κ as κ is regular uncountable and hence $cf(\kappa) > \aleph_0$. Since $\beta \in [\beta_n, \kappa)$ for all $n \in \omega$ we can find $j_n \in J_n$ such that

$$X(j_n) = M_{\kappa, \beta_n}(\beta) = M_{\kappa, \beta}(\beta)$$

for all $n \in \omega$. But all J_n are pairwise disjoint and therefore $X^{-1}(G_\beta)$ is infinite which contradicts the definition of a word (see 1.1). Thus $Occ_\kappa^+(X)/\sim_\kappa$ and also $Occ_\kappa^-(X)/\sim_\kappa$ are finite sets.

We now define $\varphi_\kappa : \mathfrak{x}_{\alpha \in \lambda} G_\alpha \rightarrow \mathbb{Z}$ as follows:

$$W \mapsto |Occ_\kappa^+(X)/\sim_\kappa| - |Occ_\kappa^-(X)/\sim_\kappa|$$

where X is the reduced word corresponding to W . Note that φ_κ is well defined by Lemma 1.3. Moreover, it follows immediately from the definition that $\varphi_\kappa(X^{-1}) = -\varphi_\kappa(X)$ and also the Specker Phenomenon obviously fails for φ_κ . Note that in general the sets $Occ_\kappa^+(X)$ and $Occ_\kappa^-(X)$ are not of the same size, e.g. $\varphi(M_{\kappa, \beta}) = 1$. It remains to show, however, that φ_κ is a homomorphism. Therefore let X and Y be reduced words. By Lemma 1.4 there exist reduced words X_1, Y_1 and M such that $X \cong X_1 M$ and $Y \cong M^{-1} Y_1$ and $X_1 Y_1$ is quasi-reduced. Now it is easy to check that $\varphi_\kappa(XY) = \varphi_\kappa(X_1 Y_1)$ by definition and the fact that $XY = X_1 M M^{-1} Y_1$. Hence

$$\varphi_\kappa(XY) = \varphi_\kappa(X_1 Y_1) = \varphi_\kappa(X_1) + \varphi_\kappa(Y_1) = \varphi_\kappa(X) + \varphi_\kappa(Y),$$

as X_1Y_1 is quasi-reduced and thus the reduced word of X_1Y_1 is obtained without cancellation. \square

We would like to remark that the uncountability of κ in Theorem 2.4 is essential for the definition of the homomorphism φ_κ and can not be omitted because of Higman's theorem. Modifying the proof of Theorem 2.4 we obtain:

Theorem 2.5 *Let λ be any uncountable cardinal and G_α ($\alpha \in \lambda$) be non-trivial groups. Then there are 2^{2^λ} homomorphisms from the free complete product of the G_α 's into the additive group of the ring of integers. In fact there is an epimorphism from $\times_{\alpha \in \lambda} G_\alpha$ onto the free abelian group of 2^λ copies of the integers.*

Proof. Let λ be uncountable and $\{G_\alpha : \alpha \in \lambda\}$ be given as stated. We choose the following family of reduced words M_α for $\alpha \in 2^\lambda$. First we choose non-trivial elements $e_\gamma \neq g_\gamma \in G_\gamma$ for $\gamma \in \lambda$. Let $\{I_\epsilon : \epsilon \in \omega_1\}$ be a family of pairwise disjoint subsets of λ each of which has cardinality λ . It is well-known (see e.g. [EK]) that for every $\epsilon \in \omega_1$ we can find a family $\{I_{\epsilon,\alpha} \subseteq I_\epsilon : \alpha \in 2^\lambda\}$ of subsets of I_ϵ such that any finite Boolean combination of them is of cardinality λ and moreover there is $\gamma_\epsilon \in I_\epsilon$ that belongs to each $I_{\epsilon,\alpha}$ for every $\alpha \in 2^\lambda$. For every $\epsilon \in \omega_1$ and for every $\alpha \in 2^\lambda$ we choose a word $M_{\epsilon,\alpha}$ such that its domain $\overline{M}_{\epsilon,\alpha}$ equals $I_{\epsilon,\alpha}$ and $M_{\epsilon,\alpha}(\sigma) = g_\sigma$ for $\sigma \in I_{\epsilon,\alpha}$. Then the composition $M_\alpha = \sum_{\epsilon \in \omega_1} M_{\epsilon,\alpha}$ is a well-defined reduced word in $\mathcal{W}(G_\gamma : \gamma \in \lambda)$ for every $\alpha \in 2^\lambda$. Before defining the claimed homomorphism let us first state the crucial condition satisfied by the M_α 's ($\alpha \in 2^\lambda$):

- (i) the domain \overline{M}_α of M_α is well-ordered of order type $\lambda\omega_1$ (the ordinal product) for each $\alpha \in 2^\lambda$ and therefore has uncountable cofinality.

Now we repeat the construction given in Theorem 2.4 replacing κ by \overline{M}_α and for a reduced word X and $\alpha \in 2^\lambda$ we define

$$Occ_\alpha^+(X) = \{J \subseteq (\overline{X}, <) : J \text{ convex, } X \upharpoonright_J \cong M_{\alpha,\sigma} \text{ for some } \sigma \in \overline{M}_\alpha\}$$

and

$$Occ_\alpha^-(X) := \{J \subseteq (\overline{X}, <) : J \text{ convex, } X \upharpoonright_J \cong M_{\alpha,\sigma}^{-1} \text{ for some } \sigma \in \overline{M}_\alpha\},$$

where $M_{\alpha,\sigma}$ is the end segment of M_α starting with the element σ , i.e. $M_{\alpha,\sigma} = M_\alpha \upharpoonright_{\{\rho \in \overline{M}_\alpha : \rho \geq \sigma\}}$. As in the proof of Theorem 2.4 two equivalence relations are defined on the sets $Occ_\alpha^+(X)$ and $Occ_\alpha^-(X)$ both denoted by \sim_α .

We are now able to define homomorphisms φ_α from the free complete product of the G_γ 's to the integers for each $\alpha \in 2^\lambda$. As in the proof of Theorem 2.4

we can see that the sets $Occ_{\alpha}^{+}(X)/\sim_{\alpha}$ and $Occ_{\alpha}^{-}(X)/\sim_{\alpha}$ are finite for any reduced word X and $\alpha \in 2^{\lambda}$. Moreover, the maps $\varphi_{\alpha} : \times_{\beta \in \lambda} G_{\beta} \rightarrow \mathbb{Z}$ defined by

$$V \mapsto |Occ_{\alpha}^{+}(X)/\sim_{\alpha}| - |Occ_{\alpha}^{-}(X)/\sim_{\alpha}|$$

where X is the reduced word corresponding to V , are well defined homomorphisms since $\gamma_{\epsilon} \in I_{\epsilon, \alpha}$ for every $\alpha \in 2^{\lambda}$.

To obtain $2^{2^{\lambda}}$ homomorphisms we will show that there is a surjection onto the free abelian group of 2^{λ} copies of the integers. Define

$$\Phi : \times_{\alpha \in \lambda} G_{\alpha} \rightarrow \prod_{\alpha \in 2^{\lambda}} \mathbb{Z}$$

via

$$\Phi(V)(\alpha) = \varphi_{\alpha}(X)$$

for a word V where X is the reduced word corresponding to V . As all the φ_{α} 's ($\alpha \in 2^{\lambda}$) are homomorphisms, so is Φ and we claim that Φ is actually a homomorphism from the free complete product of the G_{α} 's onto the direct sum of 2^{λ} copies of the integers $\bigoplus_{\alpha \in 2^{\lambda}} \mathbb{Z}$. First assume that this mapping is not into,

then there is a reduced word X and a sequence of pairwise distinct ordinals α_n ($n \in \omega$) such that $\Phi_{\alpha_n}(X) \neq 0$ for all $n \in \omega$. Thus for each $n \in \omega$ there is a convex subset $J_n \subseteq \overline{X}$ such that w.l.o.g.

$$X \upharpoonright_{J_n} \cong M_{\alpha_n, \sigma_n}$$

for some $\sigma_n \in \overline{M}_{\epsilon_n, \alpha_n} \subseteq \overline{M}_{\alpha_n}$ ($\epsilon_n \in \omega_1$). But now, if $\kappa \in \omega_1$ such that $\epsilon_n < \kappa$ for all $n \in \omega$, then the element g_{κ} (since γ_{κ} belongs to all sets I_{κ, α_n}) appears infinitely many times in X , i.e. $\overline{X}^{-1}(G_{\kappa})$ is infinite - a contradiction. Thus the image of Φ is contained in the direct sum of 2^{λ} copies of the integers.

On the other hand, by the choice of the sets $I_{\epsilon, \alpha}$ we certainly have that

$$Occ_{\alpha}^{+}(M_{\beta}) = 0 = Occ_{\alpha}^{-}(M_{\beta})$$

for distinct $\alpha, \beta \in 2^{\lambda}$. Moreover,

$$Occ_{\alpha}^{+}(M_{\alpha}) = 1 \text{ and } Occ_{\alpha}^{-}(M_{\alpha}) = 0$$

for any $\alpha \in 2^{\lambda}$. Thus we obtain

$$\Phi(M_{\alpha}) = (0, \dots, 0, 1_{\alpha}, 0, \dots) \in \bigoplus_{\beta \in 2^{\lambda}} \mathbb{Z}$$

and therefore Φ is obviously surjective. Since there are $2^{2^{\lambda}}$ homomorphisms from the direct sum of copies of \mathbb{Z} to \mathbb{Z} itself we are done. \square

We have a short remark.

Remark 2.6 Note that the above proof gives us that

- (i) the free complete product $G = \mathbf{x}_{\alpha \in \lambda} G_\alpha$ contains a free subgroup H (the group generated by the words M_α) and;
- (ii) there is a projection onto H .

The following theorem gives us a complete description of all 'interesting' homomorphisms from $G = \mathbf{x}_{\alpha \in \lambda} G_\alpha$ to the integers for uncountable λ and groups G_α ($\alpha \in \lambda$). By interesting we mean interesting with respect to the Specker Phenomenon, i.e. if W is a finite subset of λ , then all homomorphisms from the subproduct $G_W = \mathbf{x}_{\alpha \in W} G_\alpha$ to the integers extend naturally to a homomorphism from G to \mathbb{Z} but these homomorphisms are not of particular interest for us and well-understood, hence we will restrict ourselves to homomorphisms from G to \mathbb{Z} which are already zero on every finite subproduct of G .

First note that the definition of φ_{M_α} in the proof of Theorem 2.5 did not really depend on the particular word M_α but only on the fact that \overline{M}_α had uncountable cofinality. It is immediate to see that for any word M such that the domain \overline{M} has uncountable cofinality we can define such a homomorphism $\varphi_M : G \rightarrow \mathbb{Z}$. Hence we define the following set:

$$I_G = \{M \in G : cf(\overline{M}) \geq \aleph_1\}$$

and let

$$\Phi_G : G \rightarrow \prod_{M \in I_G} \mathbb{Z}_M$$

be defined by $\Phi_G(V) = (\varphi_M(X) : M \in I_G)$ where X is the reduced word corresponding to V . Now Φ_G is well-defined and we have the following theorem.

Theorem 2.7 *Let $G = \mathbf{x}_{\alpha \in \lambda} G_\alpha$ for some uncountable cardinal λ and groups G_α ($\alpha \in \lambda$). Then any homomorphism $\psi : G \rightarrow \mathbb{Z}$ which is zero on every finite subproduct of G factors through Φ_G .*

Proof. For simplicity we assume that all words are already in reduced form. First we will show that the kernel of Φ_G is exactly $\text{Ker}(\Phi_G) = \{M \in G : M \text{ contains no monotonic sequence of length } \omega_1\}$, where a monotonic sequence of length ω_1 is just a subset of \overline{M} which is isomorphic to ω_1 or its inverse. Clearly we have that $\text{Ker}(\Phi_G)$ is contained in $\{M \in G : M \text{ contains no monotonic sequence of length } \omega_1\}$. Conversely, if M is a reduced word that contains no monotonic sequence of length ω_1 and $\Phi_G(M) \neq 0$, then there exists a reduced word $N \in I_G$, i.e. \overline{N} has uncountable cofinality, such that $\varphi_N(M) \neq 0$. But then $Occ_N^+(M)$ or $Occ_N^-(M)$ is non-trivial and hence M must contain a monotonic sequence of length ω_1 - a contradiction. It is now enough to prove that any homomorphism $f : G \rightarrow \mathbb{Z}$ which is zero on any finite subproduct of

G acts trivial on $\text{Ker}(\Phi_G)$. For this assume that M is a reduced word which contains no monotonic sequence of length ω_1 such that $f(M) \neq 0$ for some homomorphism $f : G \rightarrow \mathbb{Z}$. We distinguish between three cases:

Case a: There exist subwords N_n of M ($n \in \omega$) such that

- (i) \overline{N}_n is a convex subset of \overline{M} ;
- (ii) the \overline{N}_n 's ($n \in \omega$) are pairwise (almost) disjoint;
- (iii) $f(N_n) \neq 0$ (without loss of generality $f(N_n) > 0$).

Hence the composition N of the words N_n ($n \in \omega$) is a well-defined word in G and applying Theorem 2.3 together with [E1, Proposition 1.9] leads to a contradiction.

Case b: There is an initial segment \overline{M}^* of \overline{M} such that

- (i) $f(M^*) \neq 0$, where $M^* = M \upharpoonright_{\overline{M}^*}$;
- (ii) for every proper initial segment \overline{N} of \overline{M}^* we have $f(N) = 0$, where $N = M \upharpoonright_{\overline{N}}$;
- (iii) \overline{M}^* has no largest element or for every $t \in \overline{M}^*$ there exists a convex subset $\overline{N}_t \subseteq \{m \in \overline{M}^* : m \geq t\}$ such that $f(N_t) \neq 0$, where $N_t = M \upharpoonright_{\overline{N}_t}$.

Then the cofinality of \overline{M}^* has to be \aleph_0 by the assumptions and we choose an increasing, unbounded sequence $\{t_n : n \in \omega\}$ in \overline{M}^* and put

$$N_n = \{m \in \overline{M}^* : m \geq t_n\} \text{ or } N_n = N_{t_n} \text{ (hence } f(N_n) \neq 0\text{)}.$$

In both cases we easily obtain a contradiction. Similarly we can use the same arguments for the inverse of M to get a contradiction.

Case c: Neither case a nor case b is satisfied. Then it is easy to see that the set

$$J = \{t \in \overline{M} : f(M_t) \neq 0\}$$

is finite, where $M_t = M \upharpoonright_{\{t\}}$ (e.g. use Ramsey's Theorem). So without loss of generality we may assume that J is the empty set. We let

$$I = \{A \subset \overline{M} : A \text{ convex and } \forall B \subseteq A \text{ convex } f(M \upharpoonright_B) = 0\}.$$

Then I contains all singletons $t \in \overline{M}$, the empty set and it is downwards closed. Moreover, if A and B are elements of I and $A \cup B$ is convex, then $A \cup B \in I$. Finally every initial segment of \overline{M} with no largest element has an end segment

in I and hence $M \in I$ - a contradiction. Similarly we obtain a contradiction using the inverse M^{-1} instead of M . This finishes the proof. \square

For completeness let us state the following remark.

Remark 2.8 If $h : G \rightarrow \mathbb{Z}$ is any homomorphism, then an application of Theorem 2.3 shows that the set $\{\alpha \in \lambda : h(G_\alpha) \neq \{0\}\} = F$ is finite. Hence, regarding $*_{\alpha \in F} G_\alpha$ as a subgroup of G we let $h_0 = (h \upharpoonright_{*_{\alpha \in F} G_\alpha}) \rho_F$. Then $h - h_0$ satisfies the assumptions of Theorem 2.7 and thus factors through Φ_G .

Acknowledgement: The authors would like to thank the referee for his helpful comments and suggestions and in particular for his patience.

References

- [CC] **J. W. Cannon and G. Conner**, The big fundamental group, big Hawaiian earrings, and the big free groups, *Topology Appl.* **106** (2000), 273–291.
- [CE] **G. Conner and K. Eda**, Free subgroups of complete free products, preprint (2000).
- [E1] **K. Eda**, Free σ -products and noncommutatively slender groups, *J. of Algebra* **148** (1992), 243–263.
- [E2] **K. Eda**, The non-commutative Specker Phenomenon, *J. of Algebra* **204** (1998), 95–107.
- [EK] **R. Engelking and M. Karłowicz**, Some theorems of set theory and their topological consequences, *Fund. Math.* **57** (1965), 275–285.
- [EM] **P.C. Eklof and A. Mekler**, Almost Free Modules, Set-Theoretic Methods, North-Holland Amsterdam - New York (1990).
- [F1] **L. Fuchs**, Infinite Abelian Groups - Volume I and II, Academic Press, New York - London (1970 and 1973).
- [F2] **L. Fuchs**, Abelian Groups, Hungarian Academy of Science, Budapest (1958).
- [H] **G. Higman**, Unrestricted free products and varieties of topological groups, *J. London Math. Soc.* **27** (1952), 73–81.
- [M] **W.S. Massey**, Algebraic Topology: An Introduction, *Graduate Texts in Mathematics*, Springer Verlag New York - Heidelberg - Berlin (1967).

- [Sh] **S. Shelah**, Classification Theory, *Studies in Logic and The Foundations of Mathematics*, North-Holland Amsterdam - New York - Oxford - Tokyo **92** (1990).
- [S] **E. Specker**, Additive Gruppen von Folgen ganzer Zahlen, *Portugal. Math.* **9** (1950), 131–140.

Saharon Shelah
Department of Mathematics
Hebrew University
Jerusalem, Israel
and Rutgers University
Newbrunswick, NJ U.S.A.
e-Mail: Shelah@math.huji.ac.il

Lutz Strüingmann
Fachbereich 6, Mathematik
University of Essen
45117 Essen
Germany
e-Mail: lutz.struengmann@uni-essen.de