

RADICALS AND PLOTKIN'S PROBLEM CONCERNING GEOMETRICALLY EQUIVALENT GROUPS

RÜDIGER GÖBEL (ESSEN) AND SAHARON SHELAH (JERUSALEM)

ABSTRACT. If G and X are groups and N is a normal subgroup of X , then the G -closure of N in X is the normal subgroup $\overline{X}^G = \bigcap \{ \ker \varphi \mid \varphi : X \rightarrow G, \text{ with } N \subseteq \ker \varphi \}$ of X . In particular, $\overline{1}^G = R_G X$ is the G -radical of X . Plotkin [2, 6, 3] calls two groups G and H geometrically equivalent, written $G \sim H$, if for any free group F of finite rank and any normal subgroup N of F the G -closure and the H -closure of N in F are the same. Quasiidentities are formulas of the form $(\bigwedge_{i \leq n} w_i = 1 \rightarrow w = 1)$ for any words w, w_i ($i \leq n$) in a free group. Generally geometrically equivalent groups satisfy the same quasiidentities. Plotkin showed that nilpotent groups G and H satisfy the same quasiidentities if and only if G and H are geometrically equivalent. Hence he conjectured that this might hold for any pair of groups; see the Kourovka Notebook [2]. We provide a counterexample.

In a series of paper, B. I. Plotkin and his collaborators [6, 3, 4, 5] investigated radicals of groups and their relation to quasiidentities. If G is a group, then the G -radical $R_G X$ of a group X is defined by

$$R_G X = \bigcap \{ \ker \varphi; \varphi : X \rightarrow G \text{ any homomorphism} \}.$$

Clearly, $R_G X$ is a characteristic, hence a normal subgroup of X . The radical R_G can also be used to define the G -closure $\overline{U}^G = \overline{U}$ of a normal subgroup U of X , by saying that $\overline{U}/U = R_G(X/U)$. This immediately leads to Plotkin's definition of geometrically equivalent groups, see [6, 3, 4, 5] and [2, p. 113].

Definition 0.1. *Let G and H be two groups. Then G and H are geometrically equivalent, written $G \sim H$, if for any free group F of finite rank and any normal subgroup U of F the G - and H -closure of U in F are the same, i.e. for any normal subgroup U we have $\overline{U}^G = \overline{U}^H$.*

The authors are supported by the project No. G 0545-173, 06/97 of the German-Israeli Foundation for Scientific Research & Development.

GbSh 741 in Shelah's list of publications.

It is easy to see that $G \sim H$ if and only if $R_G K = R_H K$ for all finitely generated groups K . Plotkin notes that geometrically equivalent groups satisfy the same quasi-identities. The well-known notion of quasiidentities relates to quasivarieties of groups. A *quasiidentity* is an expression of the form

$$w_1 = 1 \wedge \cdots \wedge w_n = 1 \rightarrow w = 1 \text{ where } w_i, w \in F \ (i \leq n) \text{ are words.}$$

Moreover the following was shown in [6], see [2, p.113].

Theorem 0.2. (a) *If $G \sim H$, and G is torsion-free, then H is torsion-free.*
 (b) *If G, H are nilpotent, then $G \sim H$ if and only if G and H satisfy the same quasiidentities.*

This lead Plotkin to conjecture that two groups might be geometrically equivalent if and only if they satisfy the same quasiidentities, see the Kourovka Notebook [2, p.113, problem 14.71]. In this note we refute this conjecture. Clearly there are only countably many finitely presented groups which we enumerate as the set $\mathfrak{K} = \{K_n : n \in \omega\}$ and let $G = \prod_{n \in \omega} K_n$ be the restricted direct product. Then G satisfies only those quasiidentities satisfied by all groups and so if H is any group with $G \leq H$, G satisfies the same quasiidentities as H .

R. Camm [1, p. 68, p. 75 Corollary] proved there are 2^{\aleph_0} non-isomorphic, two-generator, simple groups, see also Lyndon, Schupp [7, p. 188, Theorem 3.2]. So there exists a 2-generated simple group L which cannot be mapped nontrivially into G . We consider the pair $G, H = L \times G$ and show the following:

Theorem 0.3. *If G, H and L are as above, $R_G L = L$ and $R_H L = 1$. In particular G and H are not geometrically equivalent. Since $G \leq H$ satisfy the same quasiidentities, this is the required counterexample.*

Proof. Since L is a two-generated simple group, L is an epimorphic image of a free group of rank 2. So it is enough to prove that $R_G L = L$ and $R_H L = 1$. The first equality follows since there is no nontrivial homomorphism of L into G . On the other hand, there is a canonical embedding $L \rightarrow H = L \times G$, so $R_H L = 1$. \square

REFERENCES

- [1] R. Camm, Simple free products, Journ. London Math. Soc. **28** (1953) 66–76.
- [2] E.I. Khukhro and V.D. Mazurov, *Unsolved problems in group theory; the Kourovka Notebook*, Russian Academy of Science, Novosibirsk, 1999.

- [3] B. Plotkin, Radicals in groups, operations on classes of groups, and radical classes, Transl., II Ser. Amer. Math. Soc. **119**, (1983) 89–118.
- [4] B. Plotkin, Radicals and verbals, Radical theory, Colloqu. Math. Soc. Janos Bolyai **38**, (1985) 379–403.
- [5] B. Plotkin, *Universal Algebra, Algebraic Logic, and Databases*, Kluwer Acad. Publ. Dordrecht, Boston, London 1994.
- [6] B. Plotkin, E. Plotkin, A. Tsurkov, Geometrical equivalence of groups, Commun. Algebra **27**, (1999) 4015–4025.
- [7] R. C. Lyndon and P. E. Schupp, *Combinatorial Group Theory*, Springer Ergebnisse **89**, Berlin–Heidelberg–New York, 1977.

FACHBEREICH 6, MATHEMATIK UND INFORMATIK, UNIVERSITÄT ESSEN, 45117 ESSEN, GERMANY
E-mail address: R.Goebel@Uni-Essen.De

DEPARTMENT OF MATHEMATICS, HEBREW UNIVERSITY, JERUSALEM, ISRAEL, AND RUTGERS
UNIVERSITY, NEW BRUNSWICK, NJ, U.S.A
E-mail address: e-mail: Shelah@math.huji.ac.il