

THE COMBINATORICS OF τ -COVERS

HEIKE MILDENBERGER, SAHARON SHELAH, AND BOAZ TSABAN

ABSTRACT. We solve four out of the six open problems concerning critical cardinalities of topological diagonalization properties involving τ -covers, show that the remaining two cardinals are equal, and give a consistency result concerning this remaining cardinal. Consequently, 21 open problems concerning potential implications between these properties are settled. We also give structural results based on the combinatorial techniques.

1. INTRODUCTION

Topological properties defined by diagonalizations of open or Borel covers have a rich history in various areas of general topology and analysis, and they are closely related to infinite combinatorial notions, see [10, 19, 7, 20] for surveys on the topic and some of its applications and open problems.

Let X be an infinite set. By a *cover of X* we mean a family \mathcal{U} with $X \notin \mathcal{U}$ and $X = \bigcup \mathcal{U}$. A cover \mathcal{U} of X is said to be

- (1) a *large cover of X* if: $(\forall x \in X) \{U \in \mathcal{U} : x \in U\}$ is infinite.
- (2) an ω -*cover of X* if: $(\forall \text{finite } F \subseteq X)(\exists U \in \mathcal{U}) F \subseteq U$.
- (3) a τ -*cover of X* if: \mathcal{U} is a large cover of X , and $(\forall x, y \in X) \{U \in \mathcal{U} : x \in U \text{ and } y \notin U\}$ is finite, or $\{U \in \mathcal{U} : y \in U \text{ and } x \notin U\}$ is finite.
- (4) a γ -*cover of X* if: \mathcal{U} is infinite and $(\forall x \in X) \{U \in \mathcal{U} : x \notin U\}$ is finite.

Let X be an infinite, zero-dimensional, separable metrizable topological space (in other words, a set of reals). Let Ω , T and Γ denote the collections of all *open* ω -covers, τ -covers and γ -covers of X , respectively. Additionally, denote the collection of all open covers of X by \mathcal{O} . Our restrictions on X imply that each member of any of the above classes contains a countable member of the same class [18]. We therefore confine attention in the sequel to *countable* covers, and restrict the above four classes to contain only their countable members. Let \mathcal{A} and \mathcal{B} be any of these four classes. Scheepers [9] introduced the following *selection hypotheses* that X might satisfy:

1991 *Mathematics Subject Classification.* 03E05, 54D20, 54D80.

Key words and phrases. combinatorial cardinal characteristics of the continuum, γ -cover, ω -cover, τ -cover, selection principles, Borel covers, open covers.

Supported by the Koshland Fellowship.

The authors were partially supported by: The Austrian “Fonds zur wissenschaftlichen Förderung”, grant no. 16334 and the University of Helsinki (first author), the Edmund Landau Center for Research in Mathematical Analysis and Related Areas, sponsored by the Minerva Foundation, Germany (first and third author), the United States-Israel Binational Science Foundation Grant no. 2002323 (second author), and the Golda Meir Fund (third author). This is the second author’s publication 858.

- $S_1(\mathcal{A}, \mathcal{B})$: For each sequence $\langle U_n : n \in \mathbb{N} \rangle$ of members of \mathcal{A} , there exist members $U_n \in \mathcal{U}_n, n \in \mathbb{N}$, such that $\{U_n : n \in \mathbb{N}\} \in \mathcal{B}$.
- $S_{fin}(\mathcal{A}, \mathcal{B})$: For each sequence $\langle U_n : n \in \mathbb{N} \rangle$ of members of \mathcal{A} , there exist finite (possibly empty) subsets $\mathcal{F}_n \subseteq U_n, n \in \mathbb{N}$, such that $\bigcup_{n \in \mathbb{N}} \mathcal{F}_n \in \mathcal{B}$.
- $U_{fin}(\mathcal{A}, \mathcal{B})$: For each sequence $\langle U_n : n \in \mathbb{N} \rangle$ of members of \mathcal{A} which do not contain a finite subcover, there exist finite (possibly empty) subsets $\mathcal{F}_n \subseteq U_n, n \in \mathbb{N}$, such that $\{\bigcup \mathcal{F}_n : n \in \mathbb{N}\} \in \mathcal{B}$.

Some of the properties are never satisfied, and many equivalences hold among the meaningful ones. The surviving properties appear in Figure 1, where an arrow denotes implication [17]. It is not known whether any other implication can be added to this diagram.

Below each property P in Figure 1 appears a serial number (to be used later), and the *critical cardinality* of the property, $\text{non}(P)$, which is the minimal cardinality of a space X not satisfying that property. The definitions of the cardinals appearing in this figure can be found in [3, 2], and the results were established in [5, 17, 15].




FIGURE 1. The surviving properties

The six framed entries in Figure 1 are critical cardinalities which were not found prior to the current work. In this paper we find four of them (as can be seen in the figure), and show that the remaining two are equal. We denote this possibly new cardinal by \mathfrak{od} , and prove that consistently, $\mathfrak{od} < \min\{s, b\}$. This allows us to rule out 21 (previously) potential new implications in the diagram – see Section 6.

The definition and study of τ -covers were originally motivated by the Minimal Tower Problem concerning the consistency of $\mathfrak{p} < \mathfrak{t}$, a classical open problem in infinitary combinatorics (see [16, 17], and references therein). Interestingly, this study

modified:2006-05-08

858 revision:2006-04-23

leads in Section 4 to a problem of a similar flavor – see Theorem 5.12 and the comment before it.

Let \mathcal{B}_Γ , \mathcal{B}_T , and \mathcal{B}_Ω denote the collections of *countable Borel* γ -covers, τ -covers, and ω -covers of X , respectively. Similarly, let C_Γ , C_T , and C_Ω denote the collections of (countable) *clopen* γ -covers, τ -covers, and ω -covers of X , respectively. \mathcal{B} and C denote the collections of all countable Borel and clopen covers of X , respectively. Since we restrict attention to countable covers, we have the following, where an arrow denotes inclusion:

$$\begin{array}{ccccccc} \mathcal{B}_\Gamma & \rightarrow & \mathcal{B}_T & \rightarrow & \mathcal{B}_\Omega & \rightarrow & \mathcal{B} \\ \uparrow & & \uparrow & & \uparrow & & \uparrow \\ \Gamma & \rightarrow & T & \rightarrow & \Omega & \rightarrow & \mathcal{O} \\ \uparrow & & \uparrow & & \uparrow & & \uparrow \\ C_\Gamma & \rightarrow & C_T & \rightarrow & C_\Omega & \rightarrow & C \end{array}$$

As each of the properties $\Pi(\mathcal{A}, \mathcal{B})$, $\Pi \in \{S_1, S_{fin}, U_{fin}\}$, is anti-monotonic in its first variable, we have that for each $x, y \in \{\Gamma, T, \Omega, \mathcal{O}\}$,

$$\Pi(\mathcal{B}_x, \mathcal{B}_y) \rightarrow \Pi(x, y) \rightarrow \Pi(C_x, C_y)$$

(here $C_\mathcal{O} := C$ and $\mathcal{B}_\mathcal{O} := \mathcal{B}$). In all previously studied instances, the critical cardinalities of the corresponding properties in the Borel, open, and clopen case were the same [14, 17]. Here too, we will derive the critical cardinalities of each property in the case of open covers from combinatorial characterizations of the corresponding Borel and clopen cases, between which the property is sandwiched as above.

2. $S_1(\Gamma, T)$ AND $S_{fin}(\Gamma, T)$

Since $S_1(\mathcal{B}_\Gamma, \mathcal{B}_\Gamma)$ implies $S_1(\mathcal{B}_\Gamma, \mathcal{B}_T)$, we have that

$$\mathfrak{b} = \text{non}(S_1(\mathcal{B}_\Gamma, \mathcal{B}_\Gamma)) \leq \text{non}(S_1(\mathcal{B}_\Gamma, \mathcal{B}_T)).$$

We will show that $\text{non}(S_{fin}(C_\Gamma, C_T)) \leq \mathfrak{b}$, thus settling the critical cardinalities of $S_1(\Gamma, T)$ and $S_{fin}(\Gamma, T)$, as well as their Borel and clopen counterparts.

Definition 2.1. We use the short notation \forall^∞ for “for all but finitely many” and \exists^∞ for “there exist infinitely many”.

- (1) $A \in \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$ is a γ -array if $(\forall n)(\forall^\infty m) A(n, m) = 1$.
- (2) $\mathcal{A} \subseteq \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$ is a γ -family if each $A \in \mathcal{A}$ is a γ -array.
- (3) A family $\mathcal{A} \subseteq \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$ is *finitely τ -diagonalizable* if there exist finite (possibly empty) subsets $F_n \subseteq \mathbb{N}$, $n \in \mathbb{N}$, such that:
 - (a) For each $A \in \mathcal{A}$: $(\exists^\infty n)(\exists m \in F_n) A(n, m) = 1$;
 - (b) For each $A, B \in \mathcal{A}$:

Either $(\forall^\infty n)(\forall m \in F_n) A(n, m) \leq B(n, m)$,
or $(\forall^\infty n)(\forall m \in F_n) B(n, m) \leq A(n, m)$.

modified:2006-05-08

858 revision:2006-04-23

Definition 2.2. Assume that \mathcal{U} is a countable cover of X , bijectively enumerated as $\langle U_n : n \in \mathbb{N} \rangle$. Define the *Marczewski characteristic function* of \mathcal{U} [8], $h_{\mathcal{U}} : X \rightarrow \{0, 1\}^{\mathbb{N}}$, by

$$h_{\mathcal{U}}(x)(n) = 1 \Leftrightarrow x \in U_n.$$

(Actually, $h_{\mathcal{U}}$ depends on the chosen enumeration of \mathcal{U} , but the properties of $h_{\mathcal{U}}$ which we will use do not depend on the chosen enumeration.)

$h_{\mathcal{U}}$ is continuous if the sets U_n are clopen, and Borel if the sets U_n are Borel.

$\{0, 1\}^{\mathbb{N} \times \mathbb{N}}$ is topologically the same as the Cantor space $\{0, 1\}^{\mathbb{N}}$.

Theorem 2.3. *For a set of reals X , the following are equivalent:*

- (1) X satisfies $\mathcal{S}_{fin}(\mathcal{B}_{\Gamma}, \mathcal{B}_{\mathbb{T}})$; and
- (2) For each Borel function $\Psi : X \rightarrow \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$, if $\Psi[X]$ is a γ -family, then it is finitely τ -diagonalizable.

The corresponding assertion for $\mathcal{S}_{fin}(C_{\Gamma}, C_{\mathbb{T}})$ holds when “Borel” is replaced by “continuous”.

Proof. We will prove the clopen case; the proof for the Borel case being identical.

(2 \Rightarrow 1) Assume that $\mathcal{U}_n = \{U_m^n : m \in \mathbb{N}\}$, $n \in \mathbb{N}$, is a clopen γ -cover of X . Then for each n , we have that for all but finitely many m , $h_{\mathcal{U}_n}(x)(m) = 1$. Define $\Psi : X \rightarrow \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$ by $\Psi(x)(n, m) = h_{\mathcal{U}_n}(x)(m)$. Since each $h_{\mathcal{U}_n}$ is continuous, Ψ is continuous. Moreover, for each $x \in X$, $\Psi(x)$ is a γ -array. By (2), $\Psi[X]$ is finitely τ -diagonalizable; let $\langle F_n : n \in \mathbb{N} \rangle$ witness that. Then $\bigcup_n \{U_m^n : m \in F_n\}$ is a τ -cover of X .

(1 \Rightarrow 2) Let $\Psi : X \rightarrow \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$ be continuous and such that for each $x \in X$, $\Psi(x)$ is a γ -array. Let $Y = \Psi[X]$. Since $\mathcal{S}_{fin}(C_{\Gamma}, C_{\mathbb{T}})$ is preserved under taking continuous images, Y satisfies $\mathcal{S}_{fin}(C_{\Gamma}, C_{\mathbb{T}})$. For each n and m define

$$U_m^n = \{y \in \{0, 1\}^{\mathbb{N} \times \mathbb{N}} : y(n, m) = 1\}.$$

Each U_m^n is clopen. Define $\mathcal{U}_n = \{U_m^n : m \in \mathbb{N}\}$ for each n . There are several cases to consider.

Case 1 (the interesting case). For each n and m , $Y \not\subseteq U_m^n$. Then $\mathcal{U}_n = \{U_m^n : m \in \mathbb{N}\}$ is a γ -cover of Y for each n . By $\mathcal{S}_{fin}(C_{\Gamma}, C_{\mathbb{T}})$, there exist finite sets $\mathcal{F}_n \subseteq \mathcal{U}_n$, $n \in \mathbb{N}$, such that $\bigcup_n \mathcal{F}_n$ is a τ -cover of Y . Choose $F_n = \{m : U_m^n \in \mathcal{F}_n\}$, $n \in \mathbb{N}$. Then $\langle F_n : n \in \mathbb{N} \rangle$ shows that Y is finitely τ -diagonalizable.

Case 2. There are only finitely many n for which there exists m with $Y \subseteq U_m^n$. In this case we can ignore these n 's (taking $F_n = \emptyset$ there) and apply Case 1 for the remaining n 's.

Case 3. There are infinitely many n for which there exists m_n with $Y \subseteq U_{m_n}^n$. In this case we take $F_n = \{m_n\}$ for these n 's and $F_n = \emptyset$ otherwise. \square

Theorem 2.4. *The critical cardinalities of the properties $\mathcal{S}_{fin}(\mathcal{B}_{\Gamma}, \mathcal{B}_{\mathbb{T}})$, $\mathcal{S}_{fin}(\Gamma, \mathbb{T})$, and $\mathcal{S}_{fin}(C_{\Gamma}, C_{\mathbb{T}})$, are all equal to \mathfrak{b} .*

Theorem 2.4 follows from Theorem 2.3 and Lemma 2.6 below.

Definition 2.5. For each $f \in \mathbb{N}^{\mathbb{N}}$ define a γ -array A_f by

$$A_f(n, m) = 1 \Leftrightarrow f(n) \leq m.$$

for all n and m . For γ -arrays A, B , define the following γ -array:

$$\text{cmp}(A, B)(n, m) = \max\{A(n, m), 1 - B(n, m)\}$$

for all n and m .

Lemma 2.6. *The minimal cardinality of a γ -family which is not finitely τ -diagonalizable is \mathfrak{b} .*

Proof. Let κ be the minimal cardinality we are looking for. Obviously, $\mathfrak{b} \leq \kappa$, so it remains to show that $\kappa \leq \mathfrak{b}$. Let F be a subset of $\mathbb{N}^{\mathbb{N}}$ such that $|F| = \mathfrak{b}$, and F is unbounded on each infinite subset of \mathbb{N} . (Any unbounded set F with all elements increasing has this property.) We claim that

$$\mathcal{A} = \{A_f : f \in F\} \cup \{\text{cmp}(A_f, A_h) : f, h \in F\}$$

is not finitely diagonalizable (thus $\kappa \leq |\mathcal{A}| = \mathfrak{b}$).

Assume that $\langle F_n : n \in \mathbb{N} \rangle$ is as in 2.1(2). Define a partial function $g : \mathbb{N} \rightarrow \mathbb{N}$ by $g(n) = \max F_n$ for all n with $F_n \neq \emptyset$. By 2.1(2)(a), $\text{dom}(g)$ is infinite, thus there exists $f \in F$ such that the set $D = \{n \in \text{dom}(g) : g(n) < f(n)\}$ is infinite. Fix any $h \in F$, and take $B = \text{cmp}(A_h, A_f)$. For each $n \in D$, $A_f(n, g(n)) = 0$ and $B(n, g(n)) = 1$. Since $g(n) \in F_n$ for each n , we have by 2.1(2)(b) that

$$(\forall^\infty n \in \text{dom}(g)) A_f(n, g(n)) \leq B(n, g(n)).$$

Let $D' = \text{dom}(g) \setminus D$. By 2.1(2)(a) for A_f , D' is infinite (If $m \in F_n$ is such that $A_f(n, m) = 1$, then $f(n) \leq m \leq \max F_n = g(n)$, so $n \in D'$). Now, for all but finitely many $n \in D'$,

$$1 = A_f(n, g(n)) \leq B(n, g(n)) = A_h(n, g(n)) \leq 1,$$

thus $A_h(n, g(n)) = 1$, that is, $h(n) \leq g(n)$. Thus, $g \upharpoonright D'$ dominates all elements of F on D' , a contradiction. \square

We obtain the following interesting characterization of \mathfrak{b} .

Definition 2.7. Say that a family $\mathcal{A} \subseteq \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$ is *semi τ -diagonalizable* if there exists a *partial* function $g : \mathbb{N} \rightarrow \mathbb{N}$ such that:

- (1) For each $A \in \mathcal{A}$: $(\exists^\infty n \in \text{dom}(g)) A(n, g(n)) = 1$;
- (2) For each $A, B \in \mathcal{A}$:
 Either $(\forall^\infty n \in \text{dom}(g)) A(n, g(n)) \leq B(n, g(n))$,
 or $(\forall^\infty n \in \text{dom}(g)) B(n, g(n)) \leq A(n, g(n))$.

Corollary 2.8. *The minimal cardinality of a γ -family which is not semi τ -diagonalizable is \mathfrak{b} .*

Proof. If κ is the minimal cardinality we are looking for, then $\mathfrak{b} \leq \kappa$, and κ is not greater than the cardinal defined in Lemma 2.6. \square

We can exploit the argument in the proof of Lemma 2.6 to obtain the following rather surprising result.

Theorem 2.9. *If X^2 satisfies $\mathcal{S}_{fin}(\Gamma, \mathbb{T})$, then X satisfies $\mathcal{U}_{fin}(\Gamma, \Gamma)$. (The corresponding assertion in the Borel and clopen cases also hold.)*

Proof. Assume, towards a contradiction, that X^2 satisfies $\mathcal{S}_{fin}(\Gamma, \mathbb{T})$ but X does not satisfy $\mathcal{U}_{fin}(\Gamma, \Gamma)$.

By Hurewicz' Theorem [4], there exists a continuous image Y_1 of X in $\mathbb{N}^{\mathbb{N}}$, such that Y_1 is unbounded. Fix $f_0 \in Y_1$. The mapping from $\mathbb{N}^{\mathbb{N}}$ to $\mathbb{N}^{\mathbb{N}}$ defined by $y(n) \mapsto \max\{0, y(n) - f_0(n)\}$ is continuous. Let Y_2 be the image of Y_1 under this mapping. Note that Y_2 is unbounded, and the constant zero function $0 \in \mathbb{N}^{\mathbb{N}}$ is a member of Y_2 . The mapping from $\mathbb{N}^{\mathbb{N}}$ to $\mathbb{N}^{\mathbb{N}}$ defined by $y(n) \mapsto y(0) + \dots + y(n)$ is also continuous, let Y be the image of Y_2 under this mapping. $0 \in Y$, and since all elements of Y are increasing and Y is unbounded, Y is nowhere bounded (i.e., $\{y \upharpoonright A : y \in Y\}$ is unbounded for each infinite $A \subseteq \mathbb{N}$). Y is a continuous image of X , therefore Y^2 is a continuous image of X^2 , and since $\mathcal{S}_{fin}(\Gamma, \mathbb{T})$ is preserved under taking continuous images, Y^2 satisfies $\mathcal{S}_{fin}(\Gamma, \mathbb{T})$. By the proof of Lemma 2.6,

$$\mathcal{A} = \{A_f : f \in Y\} \cup \{\text{cmp}(A_f, A_h) : f, h \in Y\}$$

is not finitely diagonalizable. Note that for each f , $\text{cmp}(A_f, A_0) = A_f$, thus $\mathcal{A} = \{\text{cmp}(A_f, A_h) : f, h \in Y\}$. The mapping $f \mapsto A_f$ is continuous, and so is $A \mapsto 1 - A$, therefore, the mapping defined on Y^2 by $(A, B) \mapsto \text{cmp}(A, B)$ is continuous, so \mathcal{A} is a continuous image of Y^2 (thus it satisfies $\mathcal{S}_{fin}(\Gamma, \mathbb{T})$) which is not finitely τ -diagonalizable, contradicting Theorem 2.3. \square

According to Scheepers [19, Problem 9.5], one of the more interesting problems concerning Figure 1 is whether $\mathcal{S}_1(\Omega, \mathbb{T})$ implies $\mathcal{U}_{fin}(\Gamma, \Gamma)$. $\mathcal{S}_1(\Omega, \Gamma)$ is preserved under taking finite powers [5], but it is not known whether $\mathcal{S}_1(\Omega, \mathbb{T})$ is preserved under taking finite powers [17, 19].

Corollary 2.10. *If X^2 satisfies $\mathcal{S}_{fin}(\Gamma, \mathbb{T})$ whenever X satisfies $\mathcal{S}_1(\Omega, \mathbb{T})$, then $\mathcal{S}_1(\Omega, \mathbb{T})$ implies $\mathcal{U}_{fin}(\Gamma, \Gamma)$.*

3. $\mathcal{S}_1(\mathbb{T}, \mathbb{T})$ AND $\mathcal{S}_{fin}(\mathbb{T}, \mathbb{T})$

Definition 3.1. A family $\mathcal{A} \subseteq \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$ is a τ -family if:

- (1) For each $A \in \mathcal{A}$: $(\forall n)(\exists^\infty m) A(n, m) = 1$;
- (2) For each $A, B \in \mathcal{A}$ and each n :
 Either $(\forall^\infty m) A(n, m) \leq B(n, m)$,
 or $(\forall^\infty m) B(n, m) \leq A(n, m)$.

A family $\mathcal{A} \subseteq \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$ is τ -diagonalizable if there exists a function $g : \mathbb{N} \rightarrow \mathbb{N}$, such that:

- (1) For each $A \in \mathcal{A}$: $(\exists^\infty n) A(n, g(n)) = 1$;

- (2) For each $A, B \in \mathcal{A}$:
 Either $(\forall^\infty n) A(n, g(n)) \leq B(n, g(n))$,
 or $(\forall^\infty n) B(n, g(n)) \leq A(n, g(n))$.

As in the proof of Theorem 2.3, we have the following.

Theorem 3.2. *For a set of reals X :*

- (1) X satisfies $S_1(\mathcal{B}_T, \mathcal{B}_T)$ if, and only if, for each Borel function $\Psi : X \rightarrow \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$, if $\Psi[X]$ is a τ -family, then it is τ -diagonalizable.
 (2) X satisfies $S_{fin}(\mathcal{B}_T, \mathcal{B}_T)$ if, and only if, for each Borel function $\Psi : X \rightarrow \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$, if $\Psi[X]$ is a τ -family, then it is finitely τ -diagonalizable.

The corresponding assertions for $S_1(C_T, C_T)$ and $S_{fin}(C_T, C_T)$ hold when “Borel” is replaced by “continuous”. \square

Theorem 3.3.

- (1) The critical cardinalities of the properties $S_1(\mathcal{B}_T, \mathcal{B}_T)$, $S_1(T, T)$, and $S_1(C_T, C_T)$ are all equal to \mathfrak{t} .
 (2) The critical cardinalities of the properties $S_{fin}(\mathcal{B}_T, \mathcal{B}_T)$, $S_{fin}(T, T)$, and $S_{fin}(C_T, C_T)$ are all equal to $\min\{\mathfrak{s}, \mathfrak{b}\}$.

Theorem 3.3 follows from Theorem 3.2 and the following.

Lemma 3.4.

- (1) The minimal cardinality of a τ -family which is not τ -diagonalizable is \mathfrak{t} .
 (2) The minimal cardinality of a τ -family which is not finitely τ -diagonalizable is $\min\{\mathfrak{s}, \mathfrak{b}\}$.

Proof. (1) Let κ be the minimal cardinality of a τ -family which is not τ -diagonalizable. By Figure 1 and Theorem 3.2, $\mathfrak{t} \leq \text{non}(S_1(T, T)) = \kappa$, so it remains to show that there exists a τ -family \mathcal{A} such that $|\mathcal{A}| = \mathfrak{t}$ and \mathcal{A} is not τ -diagonalizable. Let $T \subseteq [\mathbb{N}]^{\aleph_0}$ be such that $|T| = \mathfrak{t}$, T is linearly ordered by \subseteq^* , and T has no pseudo-intersection. For each $t \in T$ define $A_t^0, A_t^1 \in \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$ by:

$$A_t^0(n, m) = \begin{cases} \chi_{t \setminus n}(m) & n \text{ is even} \\ 1 & n \text{ is odd} \end{cases} \quad A_t^1(n, m) = \begin{cases} \chi_{t \setminus n}(m) & n \text{ is odd} \\ 1 & n \text{ is even} \end{cases}$$

where $\chi_{t \setminus n}$ denotes the characteristic function of $t \setminus n$.

Clearly, $\mathcal{A} = \{A_t^\ell : t \in T, \ell \in \{0, 1\}\}$ is a τ -family. Assume that \mathcal{A} is τ -diagonalizable, and let $g : \mathbb{N} \rightarrow \mathbb{N}$ be a witness for that. If the image of g is finite, then for all but finitely many even n , $A_t^0(n, g(n)) = \chi_{t \setminus n}(g(n)) = 0 < 1 = A_t^1(n, g(n))$, and for all but finitely many odd n , $A_t^1(n, g(n)) = \chi_{t \setminus n}(g(n)) = 0 < 1 = A_t^0(n, g(n))$, contradicting the fact that g is a τ -diagonalization of \mathcal{A} . Thus, either $g[E]$ or $g[O]$, where E and O are the sets of even and odd natural numbers, respectively, is infinite.

Assume that $g[E]$ is infinite. Fix any $t \in T$ such that $g[E] \not\subseteq^* t$. Then $g[E] \setminus t$ is infinite, and for each element $g(n) \in g[E] \setminus t$, $A_t^0(n, g(n)) = \chi_{t \setminus n}(g(n)) = 0 < 1 = A_t^1(n, g(n))$. Thus, $A_t^0(n, g(n)) \leq A_t^1(n, g(n))$ for all but finitely many n . For n odd, $A_t^0(n, g(n)) = 1$, therefore $\chi_{t \setminus n}(g(n)) = A_t^1(n, g(n)) = 1$ for all but finitely many

modified:2006-05-08

858 revision:2006-04-23

$n \in O$, that is, $g[O] \subseteq^* t$. Since $\chi_{t \setminus n}(g(n)) = 1$ implies that $n \leq g(n)$, $g[O]$ is infinite, and therefore a pseudo-intersection of T , a contradiction.

The case that $g[O]$ is infinite is similar.

(2) Let κ be the minimal cardinality of a τ -family which is not finitely τ -diagonalizable. By Theorems 3.2 and 2.4, $\kappa = \text{non}(\mathbf{S}_{fin}(\mathbb{T}, \mathbb{T})) \leq \text{non}(\mathbf{S}_{fin}(\Gamma, \mathbb{T})) = \mathfrak{b}$. Thus, to show that $\kappa \leq \min\{\mathfrak{s}, \mathfrak{b}\}$, it suffices to construct a τ -family \mathcal{A} such that $|\mathcal{A}| = \mathfrak{s}$ and \mathcal{A} is not finitely τ -diagonalizable. Let $S \subseteq [\mathbb{N}]^{\aleph_0}$ be a splitting family of size \mathfrak{s} and $T \subseteq [\mathbb{N}]^{\aleph_0}$ be as in (1). For each $t \in T$ and $s \in S$ define $A_{t,s}^0, A_{t,s}^1 \in \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$ by:

$$A_{t,s}^0(n, m) = \begin{cases} \chi_{t \setminus n}(m) & n \in s \\ 1 & n \notin s \end{cases} \quad A_{t,s}^1(n, m) = \begin{cases} \chi_{t \setminus n}(m) & n \notin s \\ 1 & n \in s \end{cases}$$

Let $\mathcal{A} = \{A_{t,s}^\ell : t \in T, s \in S, \ell \in \{0, 1\}\}$. \mathcal{A} is a τ -family, and since $\mathfrak{t} \leq \mathfrak{s}$, $|\mathcal{A}| = \mathfrak{t} \cdot \mathfrak{s} = \mathfrak{s}$. Assume that \mathcal{A} is finitely τ -diagonalizable, and let $\langle F_n : n \in \mathbb{N} \rangle$ witness that. Choose any function g with domain $\{n : F_n \neq \emptyset\}$ and such that $g(n) \in F_n$ for each $n \in \text{dom}(g)$, and a set $s \in S$ which splits $\text{dom}(g)$. Then we can restrict attention to $\text{dom}(g)$ and apply the analysis carried in (1) to obtain a contradiction.

We now prove that $\min\{\mathfrak{s}, \mathfrak{b}\} \leq \kappa$. Assume that \mathcal{A} is a τ -family and $|\mathcal{A}| < \min\{\mathfrak{s}, \mathfrak{b}\}$. We will show that \mathcal{A} is finitely τ -diagonalizable.

For each $A, B \in \mathcal{A}$, define $s_{A,B} = \{n : (\forall^\infty m) A(n, m) \leq B(n, m)\}$. Since $|\mathcal{A}| < \mathfrak{s}$, there is $s \in [\mathbb{N}]^{\aleph_0}$ which is not split by any of the sets $s_{A,B}$, $A, B \in \mathcal{A}$. Since $s_{A,B} \cup s_{B,A} = \mathbb{N}$, we have that for each $A, B \in \mathcal{A}$, $s \subseteq^* s_{A,B}$ or $s \subseteq^* s_{B,A}$.

For each $A, B \in \mathcal{A}$ define $g_{A,B} \in \mathbb{N}^{\mathbb{N}}$ by:

$$g_{A,B}(n) = \begin{cases} \min\{k : (\forall m \geq k) A(n, m) \leq B(n, m)\} & n \in s_{A,B} \setminus s_{B,A} \\ \min\{k : (\forall m \geq k) B(n, m) \leq A(n, m)\} & n \in s_{B,A} \setminus s_{A,B} \\ \min\{k : (\forall m \geq k) A(n, m) = B(n, m)\} & n \in s_{A,B} \cap s_{B,A} \end{cases}$$

Since $|\mathcal{A}| < \mathfrak{b}$, there exists $g_0 \in \mathbb{N}^{\mathbb{N}}$ which dominates all of the functions $g_{A,B}$, $A, B \in \mathcal{A}$. For each $A \in \mathcal{A}$, define $g_A \in \mathbb{N}^{\mathbb{N}}$ by

$$g_A(n) = \min\{m : g_0(n) \leq m \text{ and } A(n, m) = 1\}.$$

Choose $g_1 \in \mathbb{N}^{\mathbb{N}}$ which dominates the functions g_A , $A \in \mathcal{A}$ (here too, this is possible since $|\mathcal{A}| < \mathfrak{b}$).

For each $n \in s$, define $F_n = [g_0(n), g_1(n)]$. For $n \notin s$ let $F_n = \emptyset$. For each $A \in \mathcal{A}$ and all but finitely many n , $A(n, g_A(n)) = 1$ and $g_0(n) \leq g_A(n) \leq g_1(n)$, so $g_A(n) \in F_n$.

We now verify the remaining requirement. Let $A, B \in \mathcal{A}$. Without loss of generality it is the case that $s \subseteq^* s_{A,B}$. For all but finitely many n : either $n \notin s$ and $F_n = \emptyset$ so there is nothing to prove, or else $n \in s$, thus $n \in s_{A,B}$, therefore for each $m \in F_n$, $g_{A,B}(n) \leq g_0(n) \leq m$, and consequently $A(n, m) \leq B(n, m)$. \square

Remark 3.5. $\min\{\mathfrak{s}, \mathfrak{b}\}$ is sometimes referred to as the *partition number* par , see [2].

Here too, we can use the combinatorial construction to obtain the following. It is an open problem whether $S_1(T, T)$ implies (and is therefore equivalent to) $S_1(T, \Gamma)$.

Theorem 3.6. *The following are equivalent:*

- (1) $S_1(T, T)$ is equivalent to $S_1(T, \Gamma)$,
- (2) $S_1(T, T)$ is preserved under taking finite unions; and
- (3) $S_1(T, T)$ is preserved under taking unions of size less than \mathfrak{t} .

The corresponding assertions for Borel and clopen covers also hold.

Proof. (1 \Rightarrow 3 \Rightarrow 2) $S_1(T, \Gamma)$ is preserved under taking unions of size less than \mathfrak{t} [17].

(2 \Rightarrow 1) We will prove the clopen case. The proof for the Borel case is the same, but the proof in the open case requires tracing down the methods of the proofs since we do not have an analogous characterization in this case.

Assume that X satisfies $S_1(C_T, C_T)$ but not $S_1(C_T, C_\Gamma)$. Let $\binom{C_T}{C_\Gamma}$ denote the property that each member of C_T contains a member of C_Γ . Then $S_1(C_T, C_\Gamma)$ is equivalent to the conjunction of $S_1(C_T, C_T)$ and $\binom{C_T}{C_\Gamma}$ [17], thus X does not satisfy $\binom{C_T}{C_\Gamma}$, so by [16] there is a continuous image T of X in $[\mathbb{N}]^{\aleph_0}$ that is linearly ordered by \subseteq^* but has no pseudo-intersection. For each $\ell \in \{0, 1\}$, the mapping $t \mapsto A_t^\ell$ defined in (1) of Theorem 3.4's proof is continuous, and that proof shows that the union of the images of these mappings does not satisfy $S_1(C_T, C_T)$. \square

4. $S_1(T, \Omega)$ AND $S_1(T, \mathcal{O})$

The critical cardinalities of $S_1(T, \Omega)$ and $S_1(T, \mathcal{O})$ are still unknown. We will show that they are equal, and give a consistency result concerning this joint cardinal.

Lemma 4.1. *If all finite powers of X satisfy $S_1(T, \mathcal{O})$, then X satisfies $S_1(T, \Omega)$. (The corresponding assertions for the Borel and clopen case also hold.)*

Proof. Observe that for each k , if \mathcal{U} is a τ -cover of X , then $\mathcal{U}^k = \{U^k : U \in \mathcal{U}\}$ is a τ -cover of X^k . Moreover, \mathcal{U}^k is a cover of X^k if, and only if, \mathcal{U} is a k -cover of X (that is, for each $F \subseteq X$ with $|F| = k$, there is $U \in \mathcal{U}$ such that $F \subseteq U$).

Assume that for each k , X^k satisfies $S_1(T, \mathcal{O})$, and let $\langle \mathcal{U}_n : n \in \mathbb{N} \rangle$ be a sequence of open τ -covers of X . Let B_0, B_1, \dots be a partition of \mathbb{N} into infinitely many infinite sets. For each k , $\langle \mathcal{U}_n^k : n \in B_k \rangle$ is a sequence of τ -covers of X^k , and consequently there exist elements $U_n^k \in \mathcal{U}_n^k$, $n \in B_k$, such that $\{U_n^k : n \in B_k\}$ is a cover of X^k , and therefore $\{U_n : n \in B_k\}$ is a k -cover of X . Thus, $\{U_n : n \in \mathbb{N}\}$ is a k -cover of X for all k , that is, an ω -cover of X . \square

Corollary 4.2. $\text{non}(S_1(T, \Omega)) = \text{non}(S_1(T, \mathcal{O}))$.

Definition 4.3. Define $\text{o}\mathfrak{d} = \text{non}(S_1(T, \mathcal{O}))$, and call it the *o-diagonalization number*.

By Figure 1, $\text{cov}(\mathcal{M}) = \text{non}(S_1(\mathcal{O}, \mathcal{O})) \leq \text{non}(S_1(T, \mathcal{O})) \leq \text{non}(S_1(\Gamma, \mathcal{O})) = \mathfrak{d}$, thus $\text{cov}(\mathcal{M}) \leq \text{o}\mathfrak{d} \leq \mathfrak{d}$.

Definition 4.4. A τ -family \mathcal{A} is *o-diagonalizable* if there exists a function $g : \mathbb{N} \rightarrow \mathbb{N}$, such that:

$$(\forall A \in \mathcal{A})(\exists n) A(n, g(n)) = 1.$$

As in the proof of Theorem 2.3, we have the following.

Theorem 4.5. *For a set of reals X :*

- (1) X satisfies $\mathfrak{S}_1(\mathcal{B}_T, \mathcal{B})$ if, and only if, for each Borel function $\Psi : X \rightarrow \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$, if $\Psi[X]$ is a τ -family, then it is o -diagonalizable.
- (2) X satisfies $\mathfrak{S}_1(C_T, C)$ if, and only if, for each continuous function $\Psi : X \rightarrow \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$, if $\Psi[X]$ is a τ -family, then it is o -diagonalizable. \square

Corollary 4.6. \mathfrak{od} is equal to the minimal cardinality of a τ -family which is not o -diagonalizable. \square

Remark 4.7. In Definition 4.4, it is equivalent to require that $(\forall A \in \mathcal{A})(\exists^\infty n) A(n, g(n)) = 1$, or even that the family consisting of the sets $\{n : A(n, g(n)) = 1\}$, $A \in \mathcal{A}$, is centered.

The relation between $\text{cov}(\mathcal{M})$ and \mathfrak{od} is similar to the relation between \mathfrak{p} and \mathfrak{t} . The remainder of this section is dedicated to this phenomenon.

Consider Definition 3.1. One can define analogously an ω -family to be a family $\mathcal{A} \subseteq \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$ such that:

- (1) For each $A \in \mathcal{A}$: $(\forall n)(\exists^\infty m) A(n, m) = 1$;
- (2) For each n , the family of all sets $\{m : A(n, m) = 1\}$, $A \in \mathcal{A}$, is centered.

In other words, we have replaced “linearly (quasi)ordered by \subseteq^* ” by “centered”. This is exactly the way to change the definition of \mathfrak{t} to that of \mathfrak{p} .

Using the standard arguments, one gets that a set of reals X satisfies $\mathfrak{S}_1(\mathcal{B}_\Omega, \mathcal{B})$ if, and only if, for each Borel function $\Psi : X \rightarrow \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$, if $\Psi[X]$ is an ω -family, then it is o -diagonalizable (and similarly in the clopen case). As $\text{non}(\mathfrak{S}_1(\mathcal{B}_\Omega, \mathcal{B})) = \text{non}(\mathfrak{S}_1(C_\Omega, C)) = \text{cov}(\mathcal{M})$ [5, 14], we have the following.

Proposition 4.8. *The minimal cardinality of an ω -family that is not o -diagonalizable is equal to $\text{cov}(\mathcal{M})$.* \square

A classical open problem asks whether $\mathfrak{p} < \mathfrak{t}$ is consistent.

Problem 4.9. *Is it consistent (relative to ZFC) that $\text{cov}(\mathcal{M}) < \mathfrak{od}$?*

The major difficulty in proving the consistency of $\mathfrak{p} < \mathfrak{t}$ is that $\mathfrak{p} = \aleph_1$ implies $\mathfrak{t} = \aleph_1$ [3, 2], so that in any model where the continuum is (at most) \aleph_2 , $\mathfrak{p} = \mathfrak{t}$. Problem 4.9 has the same feature.

Theorem 4.10. *If $\text{cov}(\mathcal{M}) = \aleph_1$, then $\mathfrak{od} = \aleph_1$.*

Proof. Assume that $\mathfrak{od} > \aleph_1$. We will show that for each family $\{f_\alpha : \alpha < \aleph_1\} \subseteq \mathbb{N}^{\mathbb{N}}$, there is $\hat{g} \in \mathbb{N}^{\mathbb{N}}$ such that for each $\alpha < \aleph_1$, there is m with $\hat{g}(m) = f_\alpha(m)$. It is well known that this implies $\text{cov}(\mathcal{M}) > \aleph_1$ [1].

Fix a family $\{f_\alpha : \alpha < \aleph_1\} \subseteq \mathbb{N}^{\mathbb{N}}$. Choose a partition $\mathbb{N} = \bigcup_n A_n$ with each A_n infinite, and an increasing sequence of natural numbers t_i , $i \in \mathbb{N}$, such that for each i and each $n \leq i$, $|A_n \cap [t_i, t_{i+1})|/t_i \geq 1$.

modified:2006-05-08

858 revision:2006-04-23

By induction on $\alpha < \aleph_1$, choose f'_α such that for each i and each $m < t_{i+1}$, $f_\alpha(m) \leq f'_\alpha(i)$, and such that for all $\alpha < \beta$, $f'_\alpha \leq^* f'_\beta$. Using $\aleph_1 < \mathfrak{od} \leq \mathfrak{d}$, choose an increasing $h \in \mathbb{N}^{\mathbb{N}}$ witnessing that $\{f'_\alpha : \alpha < \aleph_1\}$ is not dominating.

Fix a natural number n . We construct, by induction on $\alpha < \aleph_1$, partial function $g_\alpha^n : A_n \rightarrow \mathbb{N}$ with the following properties:

- (1) For each i and each $m < t_{i+1}$, if $m \in \text{dom}(g_\alpha^n)$ then $g_\alpha^n(m) \leq h(i)$.
- (2) $\lim_{i \rightarrow \infty} |\text{dom}(g_\alpha^n) \cap [t_i, t_{i+1})|/t_i = 0$.
- (3) for all $\beta \leq \alpha$, $g_\beta^n \subseteq^* g_\alpha^n$ (i.e., $\text{dom}(g_\beta^n) \subseteq^* \text{dom}(g_\alpha^n)$ and for all but finitely many $k \in \text{dom}(g_\beta^n)$, $g_\alpha^n(k) \neq g_\beta^n(k)$).
- (4) For all but finitely many i with $f'_\alpha(i) \leq h(i)$, there is $m \in [t_i, t_{i+1}) \cap \text{dom}(g_\alpha^n)$ such that $f_\alpha(m) = g_\alpha^n(m)$.

Step $\alpha = 0$. For each i with $f'_0(i) \leq h(i)$, pick $m \in [t_i, t_{i+1}) \cap A_n$, put it into $\text{dom}(g_0^n)$ and set $g_0^n(m) = f_0(m)$.

Successor step $\alpha + 1$. g_α^n is given. For each i with $f'_\alpha(i) \leq h(i)$ and $|\text{dom}(g_\alpha^n) \cap [t_i, t_{i+1})|/t_i < 1/2$, add a point to g_α^n as in Step $\alpha = 0$, to obtain $g_{\alpha+1}^n$.

Limit step. Assume that $\alpha = \sup\{\alpha_k : k \in \mathbb{N}\}$. Choose an increasing sequence m_i , $i \in \mathbb{N}$, such that for each i :

- For all $k < k' \leq i$, $g_{\alpha_k}^n \upharpoonright [t_{m_i}, \infty) \subseteq g_{\alpha_{k'}}^n$.
- For each $k \geq m_i$, $|\text{dom}(g_\alpha^n) \cap [t_k, t_{k+1})|/t_k < 1/i$.

Take $g_\alpha^n = \bigcup_k g_{\alpha_k}^n \upharpoonright [t_{m_k}, \infty)$, and add some more values of f_α as in the successor step, to make sure that (4) is satisfied. This completes the inductive construction.

For all n and i , let F_i^n denote the set of all functions from $[t_i, t_{i+1}) \cap A_n$ to $h(i)$. Let $F_n = \bigcup_{i \in \mathbb{N}} F_i^n$. For each $\alpha < \aleph_1$, let

$$I_\alpha^n = \{i : f'_\alpha(i) \leq h(i) \text{ and } (\exists m \in [t_i, t_{i+1}) \cap \text{dom}(g_\alpha^n)) f_\alpha(m) = g_\alpha^n(m)\}.$$

I_α^n differs from $\{i : f'_\alpha(i) \leq h(i)\}$ by at most finitely many points, and is therefore infinite. Define

$$X_\alpha^n = \bigcup_{i \in I_\alpha^n} \{f \in F_i^n : g_\alpha^n \upharpoonright [t_i, t_{i+1}) \subseteq f\}.$$

X_α^n is an infinite subset of F_n . Since the f'_α are \leq^* -increasing with α and the g_α^n are \subseteq^* increasing with α , we have that for all $\alpha < \beta$, $X_\alpha^n \supseteq^* X_\beta^n$.

Fix bijections $d_n : F_n \rightarrow \mathbb{N}$. For each $\alpha < \aleph_1$, define $A_\alpha \in \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$ by

$$A_\alpha(n, m) = 1 \iff (\exists f \in X_\alpha^n) m = d_n(f).$$

Then $\{A_\alpha : \alpha < \aleph_1\}$ is a τ -family. Let g be an o -diagonalization of this family, and define $\hat{g} = \bigcup_{n \in \mathbb{N}} d_n^{-1}(g(n))$, and extend it to any function with domain \mathbb{N} . We will show that \hat{g} is as promised in the beginning of this proof.

Let $\alpha < \aleph_1$ be given. Take n such that $A_\alpha(n, g(n)) = 1$. By the definition of A_α , there is $f \in X_\alpha^n$ such that $g(n) = d_n(f)$. By the definition of X_α^n , there is i such that: $g_\alpha^n \upharpoonright [t_i, t_{i+1}) \subseteq f$, and there is $m \in [t_i, t_{i+1}) \cap \text{dom}(g_\alpha^n)$ such that $f_\alpha(m) = g_\alpha^n(m)$. Consequently,

$$\hat{g}(m) = d_n^{-1}(g(n))(m) = d_n^{-1}(d_n(f))(m) = f(m) = g_\alpha^n(m) = f_\alpha(m). \quad \square$$

It follows that in all “standard” models of ZFC, $\text{cov}(\mathcal{M}) = \mathfrak{od}$, either because $\mathfrak{c} \leq \aleph_2$, or because $\text{cov}(\mathcal{M}) = \mathfrak{d}$. Even in the models of $\mathfrak{u} = \nu < \mathfrak{d} = \delta$ from [13], we have $\mathfrak{od} \leq \mathfrak{u} = \nu$ (build a τ -family that cannot be diagonalized from the descending Mathias reals s_ξ , $\xi < \nu$), and $\text{cov}(\mathcal{M}) \geq \nu$ since the second iteration there is a finite support iteration.

5. A PARTIAL CHARACTERIZATION OF \mathfrak{od}

Definition 5.1. Fix $f \in \mathbb{N}^{\mathbb{N}}$ such that for all n , $f(n) \geq 2$. An f -sequence is an element $\sigma \in \prod_n P(f(n))$ (that is, such that $\sigma(n) \subseteq f(n)$ for each n). A family \mathcal{F} of f -sequences is o -diagonalizable if there exists $g \in \mathbb{N}^{\mathbb{N}}$ such that for each $\sigma \in \mathcal{F}$ there is n such that $g(n) \in \sigma(n)$.

θ_f is the minimal cardinality of a family \mathcal{F} of f -sequences such that:

- (1) For each $\sigma \in \mathcal{F}$: $(\forall^\infty n) \sigma(n) \neq \emptyset$,
- (2) For each $\sigma, \eta \in \mathcal{F}$: Either $(\forall^\infty n) \sigma(n) \subseteq \eta(n)$, or $(\forall^\infty n) \eta(n) \subseteq \sigma(n)$.
- (3) \mathcal{F} is not o -diagonalizable.

If there is no such family, we define $\theta_f = \mathfrak{c}^+$.

Lemma 5.2. If $f_1 \leq^* f_2$, then $\theta_{f_2} \leq \theta_{f_1}$. □

Lemma 5.3. For each $f \in \mathbb{N}^{\mathbb{N}}$, $\mathfrak{od} \leq \theta_f$.

Proof. Let \mathcal{F} be a witness for θ_f , $f^* \in \mathbb{N}^{\mathbb{N}}$ be defined by $f^*(n) = \sum_{k < n} f(k)$, and $\bar{B} = \langle B_n : n \in \mathbb{N} \rangle$ be a partition of \mathbb{N} into infinite sets.

For each $\sigma \in \mathcal{F}$, define $A_\sigma \in \{0, 1\}^{\mathbb{N} \times \mathbb{N}}$ by

$$A_\sigma(n, m) = \begin{cases} 1 & (\exists k \in B_n) m \in [f^*(k), f^*(k+1)) \text{ and } m - f^*(k) \in \sigma(k) \\ 0 & \text{otherwise} \end{cases}$$

Since \mathcal{F} is a witness for θ_f , $\mathcal{A} = \{A_\sigma : \sigma \in \mathcal{F}\}$ is a τ -family. We claim that \mathcal{A} is not o -diagonalizable. Assume that $g \in \mathbb{N}^{\mathbb{N}}$ is an o -diagonalization of \mathcal{A} . For each n , let k_n be the unique k such that $g(n) \in [f^*(k), f^*(k+1))$, and let $i(k_n)$ be the unique i such that $k_n \in B_i$. Let $h \in \mathbb{N}^{\mathbb{N}}$ be any function such that for each n , $h(k_n) = \max\{0, g(i(k_n)) - f^*(k_n)\}$.

For each $A_\sigma \in \mathcal{A}$, let n be such that $A_\sigma(n, g(n)) = 1$. Then $k_n \in B_n$, $g(n) \in [f^*(k_n), f^*(k_n+1))$, and $g(n) - f^*(k_n) \in \sigma(k_n)$. Since $k_n \in B_n$, we have that $i(k_n) = n$ and therefore $h(k_n) = g(n) - f^*(k_n) \in \sigma(k_n)$. Consequently, h is an o -diagonalization of \mathcal{F} . □

Definition 5.4. $\theta_* = \min\{\theta_f : f \in \mathbb{N}^{\mathbb{N}}\}$.

Lemma 5.3 implies the following.

Corollary 5.5. $\mathfrak{od} \leq \theta_*$. □

Definition 5.6. A forcing notion \mathbb{P} has the *Laver property* if for each $f \in \mathbb{N}^{\mathbb{N}} \cap V$ (where V is the ground model), each $p \in \mathbb{P}$, and each \mathbb{P} -name \check{g} for an element of $\mathbb{N}^{\mathbb{N}}$ such that $p \Vdash_{\mathbb{P}} (\forall n) \check{g}(n) \leq f(n)$, there exist $q \in \mathbb{P}$ stronger than p and $S \in V$ such that $|S(n)| \leq 2^n$ for all n , and $q \Vdash_{\mathbb{P}} (\forall n) \check{g}(n) \in S(n)$.

The Laver property is preserved under countable support iterations of proper forcing notions [1, Theorem 6.3.34]. The best known forcing notion with the Laver property is the Laver forcing [1, Theorem 7.3.29], more forcing notions with the Laver property are Miller’s superperfect tree forcing [1, Theorem 7.3.45] and the Mathias forcing [1, Corollary 7.4.7].

Theorem 5.7.

- (1) Assume that V is a model of the Continuum Hypothesis, and \mathbb{P} is a forcing notion with the Laver property. Then in $V^{\mathbb{P}}$, $\theta_* = \aleph_1$.
- (2) In the Laver model,¹ $\aleph_1 = \mathfrak{s} = \theta_* < \mathfrak{b} = \aleph_2$.
- (3) In the Miller (superperfect forcing) model, $\aleph_1 = \theta_* = \mathfrak{b} = \mathfrak{s} < \mathfrak{g} = \mathfrak{d} = \aleph_2$.
- (4) In the Mathias model, $\aleph_1 = \theta_* < \mathfrak{h} = \mathfrak{s} = \mathfrak{b} = \aleph_2$.

Proof. (1) We need the following lemma.

Lemma 5.8. Assume that $\{S_\alpha : \alpha < \aleph_1\} \subseteq \prod_n [4^n]^{2^n}$. Then there exists a sequence $\langle \sigma_\alpha : \alpha < \aleph_1 \rangle$ such that:

- (1) For each α , $\sigma_\alpha \in \prod_n P(4^n)$
- (2) For each α and n , $\sigma_\alpha(n)$ is nonempty, and $\sigma_\alpha(n) \cap S_\alpha(n) = \emptyset$,
- (3) For each α , $\lim_n |\sigma_\alpha(n)|/2^n = \infty$; and
- (4) For each $\alpha < \beta$, $\sigma_\beta(n) \subseteq \sigma_\alpha(n)$ for all but finitely many n .

Proof. This is proved by induction on $\alpha < \aleph_1$. For $\alpha = 0$ take $\sigma_0(n) = 4^n \setminus S_0(n)$ for all n . Assume that the construction was carried out up to stage α . We will define σ_α as follows. Enumerate $\alpha = \{\beta_k : k \in \mathbb{N}\}$. Let $k_0 = 0$, and define k_ℓ by induction on $\ell \in \mathbb{N}$ as follows: Since $F = \{\beta_k : k \leq \ell\}$ is finite, there exists by the induction hypothesis $k_\ell > k_{\ell-1}$ such that for each $n \geq k_\ell$ and $\gamma < \delta$ in F ,

$$\sigma_\delta(n) \subseteq \sigma_\gamma(n).$$

Let $\delta = \max F$. By the induction hypothesis, $\lim_n |\sigma_\delta(n)|/2^n = \infty$, therefore we can increase k_ℓ so that for all $n \geq k_\ell$,

$$|\sigma_\delta(n)| \geq \ell \cdot 2^n.$$

After the sequence $\langle k_\ell : \ell \in \mathbb{N} \rangle$ was defined, we can define $\sigma_\alpha(n)$ for each n by letting ℓ be such that $k_\ell \leq n < k_{\ell+1}$, and

$$\sigma_\alpha(n) = \sigma_{\max F}(n) \setminus S_\alpha(n).$$

Then $|\sigma_\alpha(n)|/2^n \geq \ell - 1$, so the induction hypotheses continue to hold. □

Define $f(n) = 4^n$ for all n . By Lemma 5.3, it suffices to show that $\theta_f = \aleph_1$. Let

$$\mathcal{S} = V \cap \prod_n [4^n]^{2^n}.$$

¹The *Laver model* is the model obtained by a length \aleph_2 countable support iteration of the Laver forcing over a model of the Continuum Hypothesis. A similar comment applies for the other named models in this theorem.

Enumerate $\mathcal{S} = \{S_\alpha : \alpha < \aleph_1\}$, and apply Lemma 5.8 to \mathcal{S} to obtain family $\mathcal{F} = \{\sigma_\alpha : \alpha < \aleph_1\}$. By the Laver property of \mathbb{P} , for each $g \in V^{\mathbb{P}} \cap \prod_n f(n)$, there is $S_\alpha \in \mathcal{S}$ such that $g(n) \in S_\alpha(n)$ for all n . Since $\sigma_\alpha(n) \cap S_\alpha(n) = \emptyset$ for all n , \mathcal{F} is not o -diagonalizable.

(2), (3), and (4) follow from (1), since all values of the other cardinals in the corresponding models are known [2]. \square

Theorem 5.9. $\min\{\mathfrak{s}, \mathfrak{b}, \mathfrak{od}\} = \min\{\mathfrak{s}, \mathfrak{b}, \theta_*\}$. In other words, if $\mathfrak{od} < \min\{\mathfrak{s}, \mathfrak{b}\}$, then there is $f \in \mathbb{N}^{\mathbb{N}}$ such that $\mathfrak{od} = \theta_f$.

Proof. By Lemma 5.3, it suffices to show that $\mathfrak{od} \geq \min\{\mathfrak{s}, \mathfrak{b}, \theta_*\}$. Assume that $\kappa < \{\mathfrak{s}, \mathfrak{b}, \theta_*\}$, and let \mathcal{A} be a τ -family. We will show that \mathcal{A} is o -diagonalizable. (In fact, we show a little more than that.)

Since $\kappa < \{\mathfrak{s}, \mathfrak{b}\}$, \mathcal{A} is finitely τ -diagonalizable (Lemma 3.4(2)); let $\langle F_n : n \in \mathbb{N} \rangle$ witness that. Enumerate $\{n : F_n \neq \emptyset\}$ bijectively as $\{k_n : n \in \mathbb{N}\}$. Define $f \in \mathbb{N}^{\mathbb{N}}$ by $f(n) = |F_{k_n}|$ for each n , and for each n and $m < f(n)$ let $F_{k_n}(m)$ denote the m th element of F_{k_n} . For each $A \in \mathcal{A}$, define an f -sequence σ_A by:

$$\sigma_A(n) = \{m < f(n) : A(k_n, F_{k_n}(m)) = 1\}.$$

As $\kappa < \theta_f$, $\{\sigma_A : A \in \mathcal{A}\}$ is o -diagonalizable; let $g \in \mathbb{N}^{\mathbb{N}}$ be a witness for that. Choose $h \in \mathbb{N}^{\mathbb{N}}$ such that $h(k_n) = F_{k_n}(g(n))$ for all n . Then h is an o -diagonalization of \mathcal{A} . (Moreover, we have that for each $A, B \in \mathcal{A}$, either $(\forall^\infty n) A(k_n, h(k_n)) \leq B(k_n, h(k_n))$, or $(\forall^\infty n) B(k_n, h(k_n)) \leq A(k_n, h(k_n))$.) \square

Definition 5.10. For a function $f : \mathbb{N} \rightarrow \mathbb{N} \setminus \{0\}$, define

$$\text{cov}_f(\mathcal{M}) = \min\{|F| : F \subseteq \prod_n f(n) \text{ and } (\forall g \in \mathbb{N}^{\mathbb{N}})(\exists h \in F)(\forall n) h(n) \neq g(n)\};$$

$$\text{cov}_*(\mathcal{M}) = \min\{\text{cov}_f(\mathcal{M}) : f : \mathbb{N} \rightarrow \mathbb{N} \setminus \{0\}\}$$

Clearly, if $f_1 \leq^* f_2$ then $\text{cov}_{f_2}(\mathcal{M}) \leq \text{cov}_{f_1}(\mathcal{M})$.

Remark 5.11. In Definition 5.10:

- (1) We may replace “ $\forall n$ ” by “ $\forall^\infty n$ ” without changing $\text{cov}_f(\mathcal{M})$.
- (2) If we replace $F \subseteq \prod_n f(n)$ by $F \subseteq \mathbb{N}^{\mathbb{N}}$, then we get $\text{cov}(\mathcal{M})$ instead of $\text{cov}_f(\mathcal{M})$ [1, Theorem 2.4.1]. Thus, $\text{cov}(\mathcal{M}) \leq \text{cov}_*(\mathcal{M})$.
- (3) $\text{cov}_*(\mathcal{M})$ is usually referred to as the minimal cardinality of a set of reals which is not strong measure zero.
- (4) If $\text{cov}(\mathcal{M}) < \mathfrak{b}$, then $\text{cov}(\mathcal{M}) = \text{cov}_*(\mathcal{M})$ [1].

Theorem 5.12. If $\text{cov}_*(\mathcal{M}) = \aleph_1$, then $\theta_* = \aleph_1$.

Proof. Let $f : \mathbb{N} \rightarrow \mathbb{N} \setminus \{0\}$ be such that $\text{cov}_f(\mathcal{M}) = \aleph_1$. We may assume that $f(n) \geq n$ for each n .

Choose a strictly increasing sequence $\langle n_i : i \in \mathbb{N} \rangle$ such that $n_0 = 0$ and $\lim_i (n_{i+1} - n_i) = \infty$. For each i let $X_i = \prod_{n_i}^{n_{i+1}-1} f(n)$, and set $X = \bigcup_i X_i$. For $Y \subseteq X_i$ define

$$\text{nor}(Y) = \min\{|Z| : Z \subseteq X_i \text{ and } (\forall \nu \in Y)(\exists \rho \in Z)(\forall n \in [n_i, n_{i+1})) \nu(n) \neq \rho(n)\}.$$

It is easy to see that $\text{nor}(X_i) = n_{i+1} + 1 - n_i$ for each i .

Let $F = \{\eta_\alpha : \alpha < \aleph_1\}$ be a witness for $\text{cov}_f(\mathcal{M}) = \aleph_1$. We define, by induction on $\alpha < \aleph_1$, sets $U_\alpha \subseteq X$ such that:

- (1) For each $\alpha < \aleph_1$, $\lim_i \text{nor}(U_\alpha \cap X_i) = \infty$,
- (2) For each $\beta < \alpha < \aleph_1$, $U_\alpha \subseteq^* U_\beta$; and
- (3) For each $\alpha < \aleph_1$, $U_{\alpha+1} = \{\nu \in U_\alpha : (\exists n \in \text{dom}(\nu)) \nu(n) = \eta_\alpha(n)\}$.

For $\alpha = 0$, we take $U_0 = X$. For $\alpha = \beta + 1$, we take U_α as in (3). Then for each i , $\text{nor}(U_\alpha \cap X_i) \geq \text{nor}(U_\beta \cap X_i) - 1$, since if Z is a witnesses that $\text{nor}(U_\alpha \cap X_i) = k$, then $Z \cup \{\eta_\alpha \upharpoonright [n_i, n_{i+1}]\}$ witnesses that $\text{nor}(U_\beta \cap X_i) \leq k + 1$. For limit α , let β_m , $m \in \mathbb{N}$, be increasing with limit α . By induction on m , choose an increasing sequence k_m , $m \in \mathbb{N}$, such that for each $i \geq k_m$,

$$U_{\beta_0} \cap X_i \supseteq U_{\beta_1} \cap X_i \supseteq \dots \supseteq U_{\beta_m} \cap X_i$$

and $\text{nor}(U_{\beta_m} \cap X_i) \geq m$. Take $U_\alpha = \bigcup_m \{U_{\beta_m} \cap X_i : i \in [k_m, k_{m+1}]\}$. This completes the inductive construction.

Since the functions η_α are witnesses for $\text{cov}_f(\mathcal{M})$, for each sequence $\langle \nu_i : i \in \mathbb{N} \rangle \in \prod_i X_i$ there is $\alpha < \aleph_1$ such that $\nu_i \notin U_\alpha$ for all i . By the inductive hypothesis (1), we may (by adding finitely many elements to each U_α) assume that for each $\alpha < \aleph_1$ and each i , $U_\alpha \cap X_i \neq \emptyset$.

Define $\tilde{f} \in \mathbb{N}^{\mathbb{N}}$ by

$$\tilde{f}(i) = \prod_{n=n_i}^{n_{i+1}-1} f(n),$$

and for each i fix a bijection $b_i : X_i \rightarrow \tilde{f}(i)$. We will show that $\theta_{\tilde{f}} = \aleph_1$. For each $\alpha < \aleph_1$ define $\sigma_\alpha \in \prod_n P(\tilde{f}(n))$ by

$$\sigma_\alpha(n) = b_i[U_\alpha \cap X_i].$$

Then the family $\mathcal{F} = \{\sigma_\alpha : \alpha < \aleph_1\}$ witnesses that $\theta_{\tilde{f}} = \aleph_1$. Indeed, for each $g \in \mathbb{N}^{\mathbb{N}}$, there is $\alpha < \aleph_1$ such that $b_i^{-1}(g(i)) \notin U_\alpha$ (and consequently $g(i) \notin b_i[U_\alpha \cap X_i] = \sigma_\alpha(i)$) for all i . \square

Remark 5.13. Kada has pointed out to us that in the Cohen model, $\theta_* = \mathfrak{c}^+$ for all f . This is proved in [6], where he also gives an elegant extension of Theorem 5.7.

6. NEW NONIMPLICATIONS

In Table 1 of [19], all known implications and nonimplications among the properties in Figure 1 were indicated. Until now, 76 possible implications remained unsettled. In Project 9.4 of [19] we are asked to settle any of these 76 problems. Our new results imply the solution of 21 of these problems (so there remain 55 possible implications).

The situation is summarized in Table 1, which updates Table 1 of [19]. Each entry (i, j) (i th row, j th column) contains a symbol. \checkmark means that property (i) in Figure 1 implies property (j) in Figure 1. \times means that property (i) does not (provably) imply property (j) , and $?$ means that the corresponding implication is still unsettled. The reader can easily verify the new results, which are framed, by consulting Figure

modified:2006-05-08

858 revision:2006-04-23

1. The reasoning is as follows: If P and Q are properties with $\text{non}(P) < \text{non}(Q)$ consistent, then Q does not imply P .

	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
0	✓	✓	✓	✓	×	×	×	×	×	×	×	×	✓	✓	×	?	×	×	✓	✓	✓	✓
1	?	✓	✓	✓	×	×	×	×	×	×	×	×	✓	✓	×	?	×	×	?	✓	✓	✓
2	×	×	✓	✓	×	×	×	×	×	×	×	×	×	✓	×	?	×	×	×	×	×	✓
3	×	×	×	✓	×	×	×	×	×	×	×	×	×	×	×	×	×	×	×	×	×	✓
4	✓	✓	✓	✓	✓	✓	✓	✓	×	×	?	?	✓	✓	✓	✓	×	?	✓	✓	✓	✓
5	?	✓	✓	✓	?	✓	✓	✓	×	×	?	?	✓	✓	✓	✓	×	?	?	✓	✓	✓
6	×	×	✓	✓	×	×	✓	✓	×	×	?	?	×	✓	×	✓	×	?	×	×	✓	✓
7	×	×	×	✓	×	×	×	✓	×	×	×	?	×	×	×	×	×	×	×	×	×	✓
8	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
9	?	✓	✓	✓	?	✓	✓	✓	?	✓	✓	✓	✓	✓	✓	✓	✓	✓	?	✓	✓	✓
10	×	×	✓	✓	×	×	✓	✓	×	×	✓	✓	×	✓	×	✓	×	✓	×	×	✓	✓
11	×	×	×	✓	×	×	×	✓	×	×	×	✓	×	×	×	×	×	×	×	×	×	✓
12	?	?	?	?	×	×	×	×	×	×	×	×	✓	✓	×	?	×	×	?	✓	✓	✓
13	×	×	×	×	×	×	×	×	×	×	×	×	×	✓	×	?	×	×	×	×	×	✓
14	?	?	?	?	×	×	×	×	×	×	×	×	✓	✓	✓	✓	×	?	?	✓	✓	✓
15	×	×	×	×	×	×	×	×	×	×	×	×	×	✓	×	✓	×	?	×	×	✓	✓
16	?	?	?	?	?	?	?	?	?	?	?	?	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
17	×	×	×	×	×	×	×	×	×	×	×	×	×	×	×	✓	×	✓	×	×	✓	✓
18	×	×	×	×	×	×	×	×	×	×	×	×	×	?	×	?	×	×	✓	✓	✓	✓
19	×	×	×	×	×	×	×	×	×	×	×	×	×	?	×	?	×	×	×	✓	✓	✓
20	×	×	×	×	×	×	×	×	×	×	×	×	×	?	×	?	×	×	×	×	✓	✓
21	×	×	×	×	×	×	×	×	×	×	×	×	×	×	×	×	×	×	×	×	×	✓

TABLE 1. Known implications and nonimplications

REFERENCES

- [1] T. Bartoszyński and H. Judah, *Set Theory: On the structure of the real line*, A. K. Peters, Massachusetts: 1995.
- [2] A. R. Blass, *Combinatorial cardinal characteristics of the continuum*, in: **Handbook of Set Theory** (M. Foreman, A. Kanamori, and M. Magidor, eds.), Kluwer Academic Publishers, Dordrecht, to appear.
- [3] E. K. van Douwen, *The integers and topology*, in: **Handbook of Set Theoretic Topology** (eds. K. Kunen and J. Vaughan), North-Holland, Amsterdam: 1984, 111–167.
- [4] W. Hurewicz, *Über Folgen stetiger Funktionen*, *Fundamenta Mathematicae* **9** (1927), 193–204.
- [5] W. Just, A. W. Miller, M. Scheepers, and P. J. Szeptycki, *The combinatorics of open covers II*, *Topology and its Applications* **73** (1996), 241–266.
- [6] M. Kada, *Covering a bounded set of slaloms by an increasing chain of slaloms*, *Topology and its Applications*, to appear.
- [7] Lj. D. R. Kočinac, *Selected results on selection principles*, in: **Proceedings of the 3rd Seminar on Geometry and Topology** (Sh. Rezapour, ed.), July 15-17, Tabriz, Iran, 2004, 71–104.
- [8] E. Marczewski (Szpilrajn), *The characteristic function of a sequence of sets and some of its applications*, *Fundamenta Mathematicae* **31** (1938), 207–233.
- [9] M. Scheepers, *Combinatorics of open covers I: Ramsey theory*, *Topology and its Applications* **69** (1996), 31–62.
- [10] M. Scheepers, *Selection principles and covering properties in topology*, *Note di Matematica* **22** (2003), 3–41.

858 revision:2006-04-23 modified:2006-05-08

- [11] S. Shelah, *Vive la différence I: Nonisomorphism of ultrapowers of countable models*, in: **Set Theory of the Continuum**, Mathematical Sciences Research Institute Publications **26** (1992), Springer Verlag, 357–405.
- [12] A. R. Blass and S. Shelah, *There may be simple P_{\aleph_1} - and P_{\aleph_2} -points, and the Rudin-Keisler ordering may be downward directed*, *Annals of Pure and Applied Logic* **33** (1987), 213–243.
- [13] A. R. Blass and S. Shelah, *Ultrafilters with small generating sets*, *Israel Journal of Mathematics* **65** (1989), 259–271.
- [14] M. Scheepers and B. Tsaban, *The combinatorics of Borel covers*, *Topology and its Applications* **121** (2002), 357–382.
- [15] S. Shelah and B. Tsaban, *Critical cardinalities and additivity properties of combinatorial notions of smallness*, *Journal of Applied Analysis* **9** (2003), 149–162.
- [16] B. Tsaban, *A topological interpretation of \mathfrak{t}* , *Real Analysis Exchange* **25** (1999/2000), 391–404.
- [17] B. Tsaban, *Selection principles and the minimal tower problem*, *Note di Matematica* **22** (2003), 53–81.
- [18] B. Tsaban, *The combinatorics of splittability*, *Annals of Pure and Applied Logic* **129** (2004), 107–130.
- [19] B. Tsaban, *Selection principles in Mathematics: A milestone of open problems*, *Note di Matematica* **22** (2003), 179–208.
- [20] B. Tsaban, *Some new directions in infinite-combinatorial topology*, to appear in: **Set Theory** (eds. J. Bagaria and S. Todorcevic), Trends in Mathematics, Birkhäuser: 2006. <http://arxiv.org/abs/math.GN/0409069>

UNIVERSITÄT WIEN, INSTITUT FÜR FORMALE LOGIK, WÄHRINGER STR. 25, A-1090 VIENNA, AUSTRIA

E-mail address: `heike@logic.univie.ac.at`

EINSTEIN INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM, GIVAT RAM, 91904 JERUSALEM, ISRAEL, AND MATHEMATICS DEPARTMENT, RUTGERS UNIVERSITY, NEW BRUNSWICK, NJ, USA

E-mail address: `shelah@math.huji.ac.il`

DEPARTMENT OF MATHEMATICS, THE WEIZMANN INSTITUTE OF SCIENCE, REHOVOT 76100, ISRAEL

E-mail address: `boaz.tsaban@weizmann.ac.il`

URL: <http://www.cs.biu.ac.il/~tsaban>