

**$\aleph_N$ -FREE ABELIAN GROUP WITH NO  
NON-ZERO HOMOMORPHISM TO  $\mathbb{Z}$   
SH883**

SAHARON SHELAH

The Hebrew University of Jerusalem  
Einstein Institute of Mathematics  
Edmond J. Safra Campus, Givat Ram  
Jerusalem 91904, Israel

Department of Mathematics  
Hill Center-Busch Campus  
Rutgers, The State University of New Jersey  
110 Frelinghuysen Road  
Piscataway, NJ 08854-8019 USA

ABSTRACT. We, for any natural  $n$ , construct an  $\aleph_n$ -free abelian groups which have few homomorphisms to  $\mathbb{Z}$ . For this we use “ $\aleph_n$ -free  $(n+1)$ -dimensional black boxes”. The method is hopefully relevant to other constructions of  $\aleph_n$ -free abelian groups.

modified:2009-03-08

(883) revision:2009-03-05

---

I would like to thank Alice Leonhardt for the beautiful typing.  
This research was supported by German-Israeli Foundation for Scientific Research and Development  
First Typed - 06/Mar/6  
Latest Revision - 09/Mar/5

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{T}\mathcal{E}\mathcal{X}$

## ANNOTATED CONTENT

§1 Constructing  $\aleph_{k(*)+1}$ -free Abelian group

[We introduce “ $\mathbf{x}$  is a combinatorial  $k(*)$ -parameter”. We also give a short cut for getting only “there is a non-Whitehead  $\aleph_{k(*)+1}$ -free non-free abelian group” (this is from 1.6 on). This is similar to [Sh 771, §5], so proofs are put in an appendix, except 1.14, note that 1.14(3) really belongs to §3.]

## §2 Black boxes

[We prove that we have black boxes in this context, see 2.1; it is based on the simple black box. Now 2.3 belongs to the short cut.]

## §3 Constructing abelian groups from combinatorial parameter

[For  $\mathbf{x} \in K_{k(*)+1}^{\text{cb}}$  we define a class  $\mathcal{G}_{\mathbf{x}}$  of abelian groups constructed from it and a black box. We prove they are all  $\aleph_{k(*)+1}$ -free of cardinality  $|\Gamma|^{\mathbf{x}} + \aleph_0$  and some  $G \in \mathcal{G}_{\mathbf{x}}$  satisfies  $\text{Hom}(G, \mathbb{Z}) = \{0\}$ .]

## §4 Appendix 1

[We give adaptation of the proofs from [Sh 771, §5] with the relevant changes.]

§0 INTRODUCTION

For regular  $\theta = \aleph_n$  we look for a  $\theta$ -free abelian group  $G$  with  $\text{Hom}(G, \mathbb{Z}) = \{0\}$ . We first construct  $G$  and a pure subgroup  $\mathbb{Z}z \subseteq G$  which is not a direct summand. If instead “not direct product” we ask “not free” so naturally of cardinality  $\theta$ , we know much: see [EM02].

We can ask further questions on abelian groups, their endomorphism rings, similarly on modules; naturally questions whose answer is known when we demand  $\aleph_1$ -free instead  $\aleph_n$ -free; see [GbT106]. But we feel those two cases can serve as a base for significant number of such problems (or we can immitate the proofs). Also these cases are reasonable for sorting out the set theoretical situation. Why not  $\theta = \aleph_\omega$  and higher cardinals? (there are more reasonable cardinals for which such results are not excluded), we do not fully know: note that also in previous questions historically this was harder.

Note that there is such an abelian group of cardinality  $\aleph_1$ , by [Sh:98, §4] and see more in Göbel-Shelah-Struüingman [GShS 785]. However, if MA then  $\aleph_2 < 2^{\aleph_0} \Rightarrow$  any  $\aleph_2$ -free abelian group of cardinality  $< 2^{\aleph_0}$  fail the question.

The groups we construct are in a sense complete, like  ${}^\omega\mathbb{Z}$ . They are close to the ones from [Sh 771, §5] but there  $S = \{0, 1\}$  as there we are interested in Borel abelian groups. See earlier [Sh 161], see representations of [Sh 161] in [Sh 523, §3], [EM02].

However we still like to have  $\theta = \aleph_\omega$ , i.e.  $\aleph_\omega$ -free abelian groups. Concerning this we continue in [Sh 898].

We thank Ester Sternfeld and Rüdiger Göbel for corrections.

We shall use freely the well known theorem saying

*0.1 Theorem.* A subgroup of a free abelian group is a free abelian group.

**0.2 Definition.** 1)  $\text{Pr}(\lambda, \kappa)$ : means that for some  $\bar{G}$  we have:

- (a)  $\bar{G} = \langle G_\alpha : \alpha \leq \kappa + 1 \rangle$
- (b)  $\bar{G}$  is an increasing continuous sequence of free abelian groups
- (c)  $|G_{\kappa+1}| \leq \lambda$ ,
- (d)  $G_{\kappa+1}/G_\alpha$  is free for  $\alpha < \kappa$ ,
- (e)  $G_0 = \{0\}$
- (f)  $G_\beta/G_\alpha$  is free if  $\alpha \leq \beta \leq \kappa$
- (g) some  $h \in \text{Hom}(G_\kappa; \mathbb{Z})$  cannot be extended to  $\hat{h} \in \text{Hom}(G_{\kappa+1}, \mathbb{Z})$ .

2) We let  $\text{Pr}^-(\lambda, \theta, \kappa)$  be defined as above, only replacing “ $G_{\kappa+1}/G_\alpha$  is free for  $\alpha < \kappa$ ” by “ $G_{\kappa+1}/G_\alpha$  is  $\theta$ -free”.

modified:2009-03-08

(883) revision:2009-03-05

§1 CONSTRUCTING  $\aleph_{k(*)+1}$ -FREE ABELIAN GROUPS

**1.1 Definition.** 1) We say  $\mathbf{x}$  is a combinatorial parameter if  $\mathbf{x} = (k, S, \Lambda) = (k^{\mathbf{x}}, S^{\mathbf{x}}, \Lambda^{\mathbf{x}})$  and they satisfy clauses (a)-(c)

- (a)  $k < \omega$
- (b)  $S$  is a set (in [Sh 771],  $S = \{0, 1\}$ ),
- (c)  $\Lambda \subseteq {}^{k+1}(\omega S)$  and for simplicity  $|\Lambda| \geq \aleph_0$  if not said otherwise.

1A) We say  $\mathbf{x}$  is an abelian group  $k$ -parameter when  $\mathbf{x} = (k, S, \Lambda, \mathbf{a})$  such that (a),(b),(c) from part (1) and:

- (d)  $\mathbf{a}$  is a function from  $\Lambda \times \omega$  to  $\mathbb{Z}$ .

2) Let  $\mathbf{x} = (k^{\mathbf{x}}, S^{\mathbf{x}}, \Lambda^{\mathbf{x}})$  or  $\mathbf{x} = (k^{\mathbf{x}}, S^{\mathbf{x}}, \Lambda^{\mathbf{x}}, \mathbf{a}^{\mathbf{x}})$ . A parameter is a  $k$ -parameter for some  $k$  and  $K_{k(*)}^{\text{cb}}/K_{k(*)}^{\text{gr}}$  is the class of combinatorial/abelian group  $k(*)$ -parameters.

3) We may write  $\mathbf{a}_{\bar{\eta}, n}^{\mathbf{x}}$  instead  $\mathbf{a}^{\mathbf{x}}(\eta, n)$ . Let  $w_{k, m} = w(k, m) = \{\ell \leq k : \ell \neq m\}$ .

4) We say  $\mathbf{x}$  is full when  $\Lambda^{\mathbf{x}} = {}^{k(*)}(\omega S)$ .

5) If  $\Lambda \subseteq \Lambda^{\mathbf{x}}$  let  $\mathbf{x} \upharpoonright \Lambda$  be  $(k^{\mathbf{x}}, S^{\mathbf{x}}, \Lambda)$  or  $(k^{\mathbf{x}}, S^{\mathbf{x}}, \Lambda, \mathbf{a} \upharpoonright (\Lambda \times \omega))$  as suitable. We may write  $\mathbf{x} = (\mathbf{y}, \mathbf{a})$  if  $\mathbf{a} = \mathbf{a}^{\mathbf{x}}, \mathbf{y} = (k^{\mathbf{x}}, S^{\mathbf{x}}, \Lambda^{\mathbf{x}})$ .

*1.2 Convention.* If  $\mathbf{x}$  is clear from the context we may write  $k$  or  $k(*)$ ,  $S, \Lambda, \mathbf{a}$  instead of  $k^{\mathbf{x}}, S^{\mathbf{x}}, \Lambda^{\mathbf{x}}, \mathbf{a}^{\mathbf{x}}$ .

A variant of the above is

**1.3 Definition.** 1) For  $\bar{S} = \langle S_m : m \leq k \rangle$  we define when  $\mathbf{x}$  is a  $\bar{S}$ -parameter:  $\bar{\eta} \in \Lambda^{\mathbf{x}} \wedge m \leq k^{\mathbf{x}} \Rightarrow \eta_m \in {}^{\omega}(S_m)$ .

2) We say  $\bar{\alpha}$  is a  $(\mathbf{x}, \bar{\chi})$ -black box or  $\bar{\alpha}$  witness  $\text{Qr}(\mathbf{x}, \bar{\chi})$  when:

- (a)  $\bar{\chi} = \langle \chi_m : m \leq k^{\mathbf{x}} \rangle$
- (b)  $\bar{\alpha} = \langle \bar{\alpha}_{\bar{\eta}} : \bar{\eta} \in \Lambda^{\mathbf{x}} \rangle$
- (c)  $\bar{\alpha}_{\bar{\eta}} = \langle \alpha_{\bar{\eta}, m, n} : m \leq k^{\mathbf{x}}, n < \omega \rangle$  and  $\alpha_{\bar{\eta}, m, n} < \chi_m$
- (d) if  $h_m : \Lambda_m^{\mathbf{x}} \rightarrow \chi_m$  for  $m \leq k^{\mathbf{x}}$  then for some  $\bar{\eta} \in \Lambda^{\mathbf{x}}$  we have:  $m \leq k^{\mathbf{x}} \wedge n < \omega \Rightarrow h_m(\bar{\eta} \upharpoonright \langle m, n \rangle) = \alpha_{\bar{\eta}, m, n}$ , see clause (a) of Definition 1.4 below on “ $\bar{\eta} \upharpoonright \langle m, n \rangle$ ” and  $\Lambda_m^{\mathbf{x}}$ .

2A) We may replace  $\bar{\chi}$  by  $\chi$  if  $\bar{\chi} = \langle \chi : \ell \leq k^{\mathbf{x}} \rangle$ . We may replace  $\mathbf{x}$  by  $\Lambda^{\mathbf{x}}$  (so say  $\text{Qr}(\Lambda^{\mathbf{x}}, \bar{\chi})$  or say  $\bar{\alpha}$  is a  $(\Lambda, \bar{\chi})$ -black box).

3) We say a parameter  $\mathbf{x}$  is  $\bar{S}$ -full or  $\mathbf{x}$  is a full  $(\bar{S}, k)$ -parameter when  $\Lambda^{\mathbf{x}} = \prod_{m \leq k} \omega(S_m)$ .

**1.4 Definition.** For a  $k(*)$ -parameter  $\mathbf{x}$  and for  $m \leq k(*)$  let

- (a)  $\Lambda_m^{\mathbf{x}} = \Lambda_{\mathbf{x}, m} = \{\bar{\eta} : \bar{\eta} = \langle \eta_\ell : \ell \leq k(*) \rangle$  and  $\eta_m \in \omega^> S$  and  $\ell \leq k(*) \wedge \ell \neq m \Rightarrow \eta_\ell \in \omega S$  and for some  $\bar{\eta}' \in \Lambda$  we have  $n < \omega, \bar{\eta} = \bar{\eta}' \upharpoonright \langle m, n \rangle\}$  where  $\bar{\eta} = \bar{\eta}' \upharpoonright \langle m, n \rangle$  means  $\eta_m = \eta'_m \upharpoonright n$  and  $\ell \leq k(*) \wedge \ell \neq m \Rightarrow \eta_\ell = \eta'_\ell\}$
- (b)  $\Lambda_{\leq k(*)}^{\mathbf{x}}$  is  $\cup\{\Lambda_m^{\mathbf{x}} : m \leq k(*)\}$
- (c)  $m(\bar{\eta}) = m$  if  $\bar{\eta} \in \Lambda_m^{\mathbf{x}}$ .

**1.5 Definition.** 1) We say a combinatorial  $k(*)$ -parameter  $\mathbf{x}$  is free when there is a list  $\langle \bar{\eta}^\alpha : \alpha < \alpha(*) \rangle$  of  $\Lambda^{\mathbf{x}}$  such that for every  $\alpha$  for some  $m \leq k(*)$  and some  $n < \omega$  we have

$$(*) \bar{\eta}_m^\alpha \upharpoonright \langle m, n \rangle \notin \{\eta_m^\beta \upharpoonright \langle m, n \rangle : \beta < \alpha\}.$$

2) We say a combinatorial  $k$ -parameter  $\mathbf{x}$  is  $\theta$ -free when  $\mathbf{x} \upharpoonright \Lambda = (k, S^{\mathbf{x}}, \Lambda)$  is free for every  $\Lambda \subseteq \Lambda^{\mathbf{x}}$  of cardinality  $< \theta$ .

*Remark.* 1) We can require in (\*) even  $(\exists^\infty n)[\eta_m^\alpha(n) \notin \cup\{\eta_\ell^\beta(n') : \ell \leq k, \beta < \alpha, n' < \omega\}]$ .

At present this seems an immaterial change.

**1.6 Definition.** For  $k(*) < \omega$  and an abelian group  $k(*)$ -parameter  $\mathbf{x}$  we define an abelian group  $G = G_{\mathbf{x}}$  as follows: it is generated by  $\{x_{\bar{\eta}} : m \leq k(*) \text{ and } \bar{\eta} \in \Lambda_m^{\mathbf{x}}\} \cup \{y_{\bar{\eta}, n} : n < \omega \text{ and } \bar{\eta} \in \Lambda^{\mathbf{x}}\} \cup \{z\}$  freely except the equations:

$$\boxtimes_{\bar{\eta}, n} (n!)y_{\bar{\eta}, n+1} = y_{\bar{\eta}, n} + \mathbf{a}_{\bar{\eta}, n}^{\mathbf{x}} z + \sum\{x_{\bar{\eta}' \upharpoonright \langle m, n \rangle} : m \leq k(*)\}.$$

*1.7 Explanation.* A canonical example of a non-free group is  $(\mathbb{Q}, +)$ . Other examples are related to it after we divide by something. The  $y$ 's here play the role of providing (hidden) copies of  $\mathbb{Q}$ . What about  $x$ 's? For  $\bar{\eta} \in \Lambda$  we consider  $\langle y_{\bar{\eta}, n} : n < \omega \rangle$ , as a candidate to represent  $(\mathbb{Q}, +), k(*) + 1$ , "opportunities" to avoid having  $(\mathbb{Q}, +)$  as a quotient, say by dividing  $K$  by a subgroup generated by some of the  $x$ 's.

This is used to prove  $G_{\mathbf{x}}$  is not free even not  $\aleph_{k(*)+2}$ -free, which is necessary. But for each  $m \leq k(*)$  if  $\langle x_{\bar{\eta}|(m,n)} : n < \omega \rangle$  are not in  $K$ , or at least  $x_{\bar{\eta}|(m,n)}$  for  $n$  large enough then  $\mathbb{Q}$  is not represented using  $\langle y_{\bar{\eta},n} : n < \omega \rangle$ ; so we have  $k(*) + 1$  “opportunities” to avoid having  $\langle y_{\bar{\eta},n} : n < \omega \rangle$  represent  $(\mathbb{Q}, +)$  in the quotient, one for each infinite cardinal  $\leq \aleph_{k(*)}$ . This helps us prove  $\aleph_{k(*)}$ -freeness. More specifically, for each  $m(*) \leq k(*)$  if  $H \subseteq G$  is the subgroup which is generated by  $X = \{x_{\bar{\eta}|<m,n>} : m \neq m(*) \text{ and } \bar{\eta} \in {}^{k(*)+1}(\omega S) \text{ and } m \leq k(*)\}$ , still in  $G/H$  the set  $\{y_{\bar{\eta},n} : n < \omega\}$  does not generate a copy of  $\mathbb{Q}$ , as witnessed by  $\{x_{\bar{\eta}|<m(*),n>} : n < \omega\}$ .

As a warm up we note:

**1.8 Claim.** For  $k(*) < \omega$  and  $k(*)$ -parameter  $\mathbf{x}$  the abelian group  $G_{\mathbf{x}}$  is an  $\aleph_1$ -free abelian group.

Now systematically

**1.9 Definition.** Let  $\mathbf{x}$  be a  $k(*)$ -parameter.

- 1) For  $U \subseteq {}^\omega S$  let  $G_U = G_U^{\mathbf{x}}$  be the subgroup of  $G$  generated by  $Y_U = Y_U^{\mathbf{x}} = \{z\} \cup \{y_{\bar{\eta},n} : \bar{\eta} \in \Lambda \cap {}^{k(*)+1}(U) \text{ and } n < \omega\} \cup \{x_{\bar{\eta}|<m,n>} : m \leq k(*) \text{ and } \bar{\eta} \in \Lambda \cap {}^{(k(*)+1)}(U) \text{ and } n < \omega\}$ . Let  $G_U^+ = G_U^{\mathbf{x},+}$  be the divisible hull of  $G_U$  and  $G^+ = G_{({}^\omega S)}^+$ .
- 2) For  $U \subseteq {}^\omega S$  and finite  $u \subseteq {}^\omega S$  let  $G_{U,u}$  be the subgroup<sup>1</sup> of  $G$  generated by  $\cup\{G_{U \cup (u \setminus \{\eta\})} : \eta \in u\}$ ; and for  $\bar{\eta} \in {}^{k(*)} \geq U$  let  $G_{U,\bar{\eta}}$  be the subgroup of  $G$  generated by  $\cup\{G_{U \cup \{\eta_k : k < \ell g(\bar{\eta}) \text{ and } k \neq \ell\}} : \ell < \ell g(\bar{\eta})\}$ .
- 3) For  $U \subseteq {}^\omega S$  let  $\Xi_U = \Xi_U^{\mathbf{x}} = \{\text{the equation } \boxtimes_{\bar{\eta},n} : \bar{\eta} \in \Lambda \cap {}^{k(*)+1}U \text{ and } n < \omega\}$ . Let  $\Xi_{U,u} = \Xi_{U,u}^{\mathbf{x}} = \cup\{\Xi_{U \cup (u \setminus \{\beta\})} : \beta \in u\}$ .

**1.10 Claim.** Let  $\mathbf{x} \in K_{k(*)}$ .

- 0) If  $U_1 \subseteq U_2 \subseteq {}^\omega S$  then  $G_{U_1}^+ \subseteq G_{U_2}^+ \subseteq G^+$ .
- 1) For any  $n(*) < \omega$ , the abelian group  $G_U^+$  (which is a vector space over  $\mathbb{Q}$ ), has the basis  $Y_U^{n(*)} := \{z\} \cup \{y_{\bar{\eta},n(*)} : \bar{\eta} \in \Lambda \cap {}^{k(*)+1}(U)\} \cup \{x_{\bar{\eta}|<m,n>} : m \leq k(*), \bar{\eta} \in \Lambda \cap {}^{k(*)+1}(U) \text{ and } n < \omega\}$ .
- 2) For  $U \subseteq {}^\omega S$  the abelian group  $G_U$  is generated by  $Y_U$  freely (as an abelian group) except the set  $\Xi_U$  of equations.
- 3) If  $m(*) < \omega$  and  $U_m \subseteq {}^\omega S$  for  $m < m(*)$  then the subgroup  $G_{U_0} + \dots + G_{U_{m(*)-1}}$  of  $G$  is generated by  $Y_{U_0} \cup Y_{U_1} \cup \dots \cup Y_{U_{m(*)-1}}$  freely (as an abelian group) except the equations in  $\Xi_{U_0} \cup \Xi_{U_1} \cup \dots \cup \Xi_{U_{m(*)-1}}$ .
- 3A) Moreover  $G/(G_{U_0} + \dots + G_{U_{m(*)+1}})$  is  $\aleph_1$ -free provided that

⊗ if  $\eta_0, \dots, \eta_{k(*)} \in \cup\{U_m : m < m(*)\}$  are such that

<sup>1</sup>note that if  $u = \{\eta\}$  then  $G_{U,u} = G_U$

$(\forall \ell \leq k(*))(\exists m < m(*))[\{\eta_0, \dots, \eta_{k(*)}\} \setminus \{\eta_\ell\} \subseteq U_m]$   
then for some  $m < m(*)$  we have  $\{\eta_0, \dots, \eta_{k(*)}\} \subseteq U_m$ .

- 4) If  $m(*) \leq k(*)$  and  $U_\ell = U \setminus U'_\ell$  for  $\ell < m(*)$  and  $\langle U'_\ell : \ell < m(*) \rangle$  are pairwise disjoint then  $\circledast$  holds.
- 5)  $G_{U,u} \subseteq G_{U \cup u}$  if  $U \subseteq {}^\omega S$  and  $u \subseteq {}^\omega S \setminus U$  is finite; moreover  $G_{U,u} \subseteq_{\text{pr}} G_{U \cup u} \subseteq_{\text{pr}} G$ .
- 6) If  $\langle U_\alpha : \alpha < \alpha(*) \rangle$  is  $\subseteq$ -increasing continuous then also  $\langle G_{U_\alpha} : \alpha < \alpha(*) \rangle$  is  $\subseteq$ -increasing continuous.
- 7) If  $U_1 \subseteq U_2 \subseteq U \subseteq {}^\omega S$  and  $u \subseteq {}^\omega S \setminus U$  is finite,  $|u| < k(*)$  and  $U_2 \setminus U_1 = \{\eta\}$  and  $v = u \cup \{\eta\}$  then  $(G_{U,u} + G_{U_2 \cup u}) / (G_{U,u} + G_{U_1 \cup u})$  is isomorphic to  $G_{U_1 \cup v} / G_{U_1, v}$ .
- 8) If  $U \subseteq {}^\omega S$  and  $u \subseteq {}^\omega S \setminus U$  has  $\leq k(*)$  members then  $(G_{U,u} + G_u) / G_{U,u}$  is isomorphic to  $G_u / G_{\emptyset, u}$ .

**1.11 Discussion:** For the reader's benefit we write what the group  $G_{\mathbf{x}}$  is for the case  $k(*) = 0$ . So, omitting constant indexes and replacing sequences of length one by the unique entry we get that it is generated by  $y_{\eta, n}$  (for  $\eta \in {}^\omega S, n < \omega$ ) and  $x_\nu$  (for  $\nu \in {}^{\omega >} S$ ) freely as an abelian group except the equations  $(n!)y_{\eta, n+1} = y_{\eta, n} + x_{\eta \upharpoonright n}$ . Note that if  $K$  is the countable subgroup generated by  $\{x_\nu : \nu \in {}^{\omega >} 2\}$  then  $G/K$  is a divisible group of cardinality continuum hence  $G$  is not free. So  $G$  is  $\aleph_1$ -free but not free.

Now we have the abelian group version of freeness, the positive results in 1.12, 1.13 and the negative results in 1.13.

**1.12 The Freeness Claim.** Let  $\mathbf{x} \in K_{k(*)}$ .

- 1) The abelian group  $G_{U \cup u} / G_{U, u}$  is free if  $U \subseteq {}^\omega S, u \subseteq {}^\omega S \setminus U$  and  $|u| \leq k \leq k(*)$  and  $|U| \leq \aleph_{k(*)-k}$ .
- 2) If  $U \subseteq {}^\omega S$  and  $|U| \leq \aleph_{k(*)}$ , then  $G_U$  is free.

- 1.13 Claim.** 1) If  $\mathbf{x}$  is a combinatorial  $k(*)$ -parameter then  $\mathbf{x}$  is  $\aleph_{k(*)+1}$ -free.  
 2) If  $\mathbf{x}$  is an abelian group  $k(*)$ -parameter and  $(k^{\mathbf{x}}, S^{\mathbf{x}}, \Lambda^{\mathbf{x}})$  is free, then  $G_{\mathbf{x}}$  is free.

*Proof.* 1) Easily follows by (2).

2) Similar and follows from 3.2 as easily  $G$  belongs to  $\mathcal{G}_{(k(*), S^{\mathbf{x}}, \Lambda^{\mathbf{x}})}$ , see Definition 3.3.

**1.14 Claim.** Assume  $\mathbf{x} \in K_{k(*)}^{\text{cb}}$  is full (i.e.  $\Lambda^{\mathbf{x}} = {}^{k(*)+1}(\omega(S^{\mathbf{x}}))$ ).

1) If  $U \subseteq \omega S$  and  $|U| \geq (|S| + \aleph_0)^{+(k(*)+1)}$ , the  $(k(*) + 1)$ -th successor of  $|S| + \aleph_0$ . Then  $G_U^{\mathbf{x}}$  is not free.

2) If  $|S^{\mathbf{x}}| \geq \aleph_{k(*)+1}$  then  $G_{\mathbf{x}}$  is not free.

3) Assume  $\mathbf{x} \in K_{k(*)}^{\text{cb}}$ ,  $|S_{\ell}^{\mathbf{x}}| + \lambda_{\ell} < \lambda_{\ell+1}$  for  $\ell < k(*)$  and  $|\Lambda^{\mathbf{x}}| \geq \lambda_{k(*)}$  and  $G \in \mathcal{G}_{\mathbf{x}}$  (see Definition 3.3) then  $G$  is not free.

*Proof.* 1) Let  $\aleph_{\alpha} = |S|$ . Assume toward contradiction that  $G_U$  is free and let  $\chi$  be large enough; for notational simplicity assume  $|U| = \aleph_{\alpha+k(*)+1}$ , this is O.K. as a subgroup of a free abelian group is a free abelian group. We choose  $N_{\ell}$  by downward induction on  $\ell \leq k(*)$  such that

- (a)  $N_{\ell}$  is an elementary submodel<sup>2</sup> of  $(\mathcal{H}(\chi), \in, <_{\chi}^*)$
- (b)  $\|N_{\ell}\| = |N_{\ell} \cap \aleph_{\alpha+k(*)}| = \aleph_{\alpha+\ell}$  and  $\{\zeta : \zeta \leq \aleph_{\alpha+\ell}\} \subseteq N_{\ell}$
- (c)  $\langle x_{\bar{\eta}} : \bar{\eta} \in \Lambda_{\leq k(*)}^{\mathbf{x}} \rangle, \langle y_{\bar{\eta}, n} : \bar{\eta} \in \Lambda^{\mathbf{x}} \text{ and } n < \omega \rangle, U$  and  $G_U$  belong to  $N_{\ell}$  and  $N_{\ell+1}, \dots, N_{k(*)} \in N_{\ell}$ .

Let  $G_{\ell} = G_U \cap N_{\ell}$ , a subgroup of  $G_U$ . Now

- (\*)<sub>0</sub>  $G_U / (\Sigma\{G_{\ell} : \ell \leq k(*)\})$  is a free (abelian) group [easy or see [Sh 52], that is: as  $G_U$  is free we can prove by induction on  $k \leq k(*)+1$  then  $G / (\Sigma\{G_{k(*)+1-\ell} : \ell < k\})$  is free, for  $k = 0$  this is the assumption toward contradiction, the induction step is by Ax VI in [Sh 52] for abelian groups and for  $k = k(*) + 1$  we get the desired conclusion.]

Now

- (\*)<sub>1</sub> letting  $U_{\ell}^1$  be  $U$  for  $\ell = k(*) + 1$  and  $\bigcap_{m=\ell}^{k(*)} (N_m \cap U)$  for  $\ell \leq k(*)$ ; we have:  $U_{\ell}^1$  has cardinality  $\aleph_{\alpha+\ell}$  for  $\ell \leq k(*) + 1$  [Why? By downward induction on  $\ell$ . For  $\ell = k(*) + 1$  this holds by an assumption. For  $\ell = k(*)$  this holds by clause (b). For  $\ell < k(*)$  this holds by the choice of  $N_{\ell}$  as the set  $\bigcap_{m=\ell+1}^{k(*)} (N_m \cap U)$  has cardinality  $\aleph_{\alpha+\ell+1} \geq \aleph_{\ell}$  and belong to  $N_{\ell}$  and clause (b) above.]

<sup>2</sup>  $\mathcal{H}(\chi)$  is  $\{x : \text{the transitive closure of } x \text{ has cardinality } < \chi\}$  and  $<_{\chi}^*$  is a well ordering of  $\mathcal{H}(\chi)$



(\*)<sub>2</sub>  $U_\ell^2 =: U_{\ell+1}^1 \setminus (N_\ell \cap U)$  has cardinality  $\aleph_{\alpha+\ell+1}$  for  $\ell \leq k(*)$   
 [Why? As  $|U_{\ell+1}^1| = \aleph_{\ell+1} > \aleph_\ell = \|N_\ell\| \geq |N_\ell \cap U|.$ ]

(\*)<sub>3</sub> for  $m < \ell \leq k(*)$  the set  $U_{m,\ell}^3 =: U_\ell^2 \cap \bigcap_{r=m}^{\ell-1} N_r$  has cardinality  $\aleph_{\alpha+m}$   
 [Why? By downward induction on  $m$ . For  $m = \ell - 1$  as  $U_\ell^2 \in N_m$  and  $|U_\ell^2| = \aleph_{\alpha+\ell+1}$  and clause (b). For  $m < \ell - 1$  similarly.]

Now for  $\ell = 0$  choose  $\eta_\ell^* \in U_\ell^2$ , possible by (\*)<sub>2</sub> above. Then for  $\ell > 0, \ell \leq k(*)$  choose  $\eta_\ell^* \in U_{0,\ell}^3$ . This is possible by (\*)<sub>3</sub>. So clearly

(\*)<sub>4</sub>  $\eta_\ell^* \in U$  and  $\eta_\ell^* \in N_m \cap U \Leftrightarrow \ell \neq m$  for  $\ell, m \leq k(*)$ .  
 [Why? If  $\ell = 0$ , then by its choice,  $\eta_\ell^* \in U_\ell^2$ , hence by the definition of  $U_\ell^2$  in (\*)<sub>2</sub> we have  $\eta_\ell^* \notin N_\ell$ , and  $\eta_\ell^* \in U_{\ell+1}^1$  hence  $\eta_\ell^* \in N_{\ell+1} \cap \dots \cap N_{k(*)}$  by (\*)<sub>1</sub> so (\*)<sub>4</sub> holds for  $\ell = 0$ . If  $\ell > 0$  then by its choice,  $\eta_\ell^* \in U_{0,\ell}^3$  but  $U_{m,\ell}^3 \subseteq U_\ell^2$  by (\*)<sub>3</sub> so  $\eta_\ell^* \in U_\ell^2$  hence as before  $\eta_\ell^* \in N_{\ell+1} \cap \dots \cap N_{k(*)}$  and  $\eta_\ell^* \notin N_\ell$ .

Also by (\*)<sub>3</sub> we have  $\eta_\ell^* \in \bigcap_{r=0}^{\ell-1} N_r$  so (\*)<sub>4</sub> really holds.]

Let  $\bar{\eta}^* = \langle \eta_\ell^* : \ell \leq k(*) \rangle$  and let  $G'$  be the subgroup of  $G_U$  generated by  $\{x_{\bar{\eta} \upharpoonright \langle m, n \rangle} : m \leq k(*) \text{ and } \bar{\eta} \in {}^{k(*)+1}U \text{ and } n < \omega\} \cup \{y_{\bar{\eta}, n} : \bar{\eta} \in {}^{k(*)+1}U \text{ but } \bar{\eta} \neq \bar{\eta}^* \text{ and } n < \omega\}$ . Easily  $G_\ell \subseteq G'$  recalling  $G_\ell = N_\ell \cap G_U$  hence  $\Sigma\{G_\ell : \ell \leq k(*)\} \subseteq G'$ , but  $y_{\bar{\eta}^*, 0} \notin G'$  hence

(\*)<sub>5</sub>  $y_{\bar{\eta}^*, 0} \notin \Sigma\{G_\ell : \ell \leq k(*)\}$ .

But for every  $n$

(\*)<sub>6</sub>  $\bar{n}! y_{\bar{\eta}^*, n+1} - y_{\bar{\eta}^*, n} = \Sigma\{x_{\bar{\eta}^* \upharpoonright \langle m, n \rangle} : m \leq k(*)\} \in \Sigma\{G_\ell : \ell \leq k(*)\}$ .  
 [Why?  $x_{\bar{\eta}^* \upharpoonright \langle m, n \rangle} \in G_m$  as  $\bar{\eta}^* \upharpoonright (k(*) + 1 \setminus \{m\}) \in N_m$  by (\*)<sub>4</sub>.]

We can conclude that in  $G_U / \Sigma\{G_\ell : \ell \leq k(*)\}$ , the element  $y_{\bar{\eta}^*, 0} + \Sigma\{G_\ell : \ell \leq k(*)\}$  is not zero (by (\*)<sub>5</sub>) but is divisible by every natural number by (\*)<sub>6</sub>.

This contradicts (\*)<sub>0</sub> so we are done.

2),3) Left to the reader.

□<sub>1.14</sub>

## §2 BLACK BOXES

**2.1 Claim.** 1) Assume  $k(*) < \omega$ ,  $\chi = \chi^{\aleph_0}$ ,  $\lambda = \beth_{k(*)}(\chi)$  and  $S = \lambda$ ,  $\Lambda = {}^{k(*)+1}(\omega S)$  or just  $S_\ell = \lambda_\ell = \chi_\ell$ ,  $\beth_\ell(\chi) = \lambda_\ell^{\aleph_0} = \chi_\ell$  for  $\ell \leq k(*)$  and  $\Lambda = \prod_{\ell \leq k(*)} \omega(S_\ell)$  and

$\mathbf{x} = (k(*), \lambda, \Lambda)$  so  $\mathbf{x}$  is a full combinatorial  $\langle S_\ell : \ell \leq k(*) \rangle$ -parameter. Then  $\Lambda$  has a  $\chi$ -black box, i.e.  $\text{Qr}(\Lambda_{\mathbf{x}^{k(*)}}, \chi)$ , see Definition 1.3.

2) Moreover,  $\mathbf{x}$  has the  $\langle \chi_\ell : \ell \leq k(*) \rangle$ -black box, i.e. for every  $\bar{B} = \langle B_{\bar{\eta}} : \bar{\eta} \in \Lambda_{\leq k(*)}^{\mathbf{x}} \rangle$  satisfying clause (c) below we can find  $\langle h_{\bar{\eta}} : \bar{\eta} \in \Lambda \rangle$  such that:

- (a)  $h_{\bar{\eta}}$  is a function with domain  $\{\bar{\eta} \upharpoonright \langle m, n \rangle : m \leq k(*), 2 \leq n < \omega\}$
- (b)  $h_{\bar{\eta}}(\bar{\eta} \upharpoonright \langle m, n \rangle) \in B_{\bar{\eta} \upharpoonright \langle m, n \rangle}$
- (c)  $B_{\bar{\eta} \upharpoonright \langle m, n \rangle}$  is a set of cardinality  $\chi_m$
- (d) if  $h$  is a function with domain  $\Lambda_{\leq k(*)}^{\mathbf{x}}$ , see Definition 1.4 such that  $h(\bar{\eta} \upharpoonright \langle m, n \rangle) \in B_{(\bar{\eta} \upharpoonright \langle m, n \rangle)}$  for  $\bar{\eta} \in \Lambda$ ,  $m \leq k(*), n < \omega$  and  $\alpha_\ell < \lambda_\ell$  for  $\ell \leq k(*)$  then for some  $\bar{\eta} \in \Lambda^{\mathbf{x}}$ ,  $h_{\bar{\eta}} \subseteq h$  and  $\eta_\ell(0) = \alpha_\ell$  for  $\ell \leq k(*)$ .

3) Assume  $\chi_\ell = \lambda_\ell^{\aleph_0}$ ,  $\chi_{\ell+1} = \chi_{\ell+1}^{\chi_\ell}$  for  $\ell \leq k(*)$ . If  $S_\ell = \lambda_\ell$  for simplicity, for  $\ell \leq k(*)$ ,  $\mathbf{x}$  is a full combinatorial  $(\bar{S}, k(*))$ -parameter, and  $|B_{\bar{\eta} \upharpoonright \langle m, n \rangle}| \leq \chi_{k(*)}$  for  $\bar{\eta} \in \Lambda^{\mathbf{x}}$  then we can find  $\langle h_{\bar{\eta}} : \bar{\eta} \in \Lambda^{\mathbf{x}} \rangle$  as in part (2), moreover such that:

- (e) if  $\bar{\eta} \in \Lambda$  then  $\eta_\ell$  is increasing
- (f) if  $\lambda_\ell$  is regular then we can in clause (d) above add: if  $E_\ell$  is a club of  $\lambda_\ell$  for  $\ell \leq k(*)$  then we can demand: if  $\bar{\eta} \in \Lambda^{\mathbf{x}}$  then for each  $\ell$  for some  $\alpha_\ell^* < \lambda_\ell$  we have  $\eta_\ell \in {}^\omega(E_\ell \cup \{\alpha_\ell^*\})$
- (g) if  $\lambda_\ell$  is singular of uncountable cofinality,  $\lambda_\ell = \Sigma\{\lambda_{\ell,i} : i < \text{cf}(\lambda_\ell)\}$ ,  $\text{cf}(\lambda_{i,\ell}) = \lambda_{i,\ell}$  increasing with  $i$  we can add: if  $u_\ell \subseteq \text{cf}(\lambda_\ell)$  is unbounded,  $E_{\ell,i}$  a club of  $\lambda_{\ell,i}$  then  $\eta_\ell \in {}^\omega(E_{i,\ell} \cup \{\alpha_\ell^*\})$  for some  $i \in u_\ell$ .

*Proof.* Part (1) follows from part (2) which follows from part (3), so let us prove part (3). To uniformize the notation in 2.1(1), i.e. 1.3(2) and 2.1(2),(3) we shall denote:

$$\odot_1 \quad h_{\bar{\eta}}(\bar{\eta} \upharpoonright \langle m, n \rangle) = \alpha_{\bar{\eta}, m, n}^{k(*)}.$$

Note that without loss of generality<sup>3</sup>  $\bar{\nu} \in \Lambda_{\leq k(*)}^{\mathbf{x}} \Rightarrow B_{\bar{\nu}} = |B_{\bar{\nu}}|$ , i.e. without loss of generality  $\bar{\eta} \in \Lambda^{\mathbf{x}} \wedge n < \omega \wedge m \leq k(*) \Rightarrow B_{\eta \upharpoonright \langle m, n \rangle} = \chi_n$  and we use  $\alpha_{\bar{\eta}, m, n}^{k(*)} = h_{\bar{\eta}}(\bar{\eta} \upharpoonright \langle m, n \rangle)$  for  $\bar{\eta} \in \Lambda_{\mathbf{x}}, m \leq k(*)$  and  $n < \omega$ . We prove part (3) by induction on  $k(*)$ . Let  $\Lambda_k = \Lambda^{\mathbf{x}}$  and without loss of generality  $S_\ell = \lambda_\ell$ .

Case 1:  $k(*) = 0$ .

By the simple black box, see [Sh 300, III,§4], or better [Sh:e, VI,§2], see below for details on such a proof.

Case 2:  $k(*) = k + 1$ .

Let

- ⊙<sub>2</sub>  $\alpha^k = \langle \alpha_{\bar{\eta}, m, n}^k : \bar{\eta} \in \Lambda_k, n < \omega, m \leq k \rangle$  witness parts (2),(3) for  $k$ , i.e. for  $\mathbf{x}^k$ , hence no need to assume  $\mathbf{x}^k$  is full.

So  $\lambda = \lambda_{k(*)}, \chi = \chi_{k(*)}$  and let  $\mathbf{H} = \{h : h \text{ is a function from } \Lambda_k \text{ to } \chi\}$ . So  $|\mathbf{H}| \leq (\lambda)^{\aleph_0} = \chi$ . By the simple black box, see below, we can find  $\langle \bar{h}_\eta : \eta \in {}^\omega \lambda \rangle$  such that

- ⊙<sub>3</sub> (α)  $\bar{h}_\eta = \langle h_{\eta, n} : n < \omega \rangle$  and  $h_{\eta, n} \in \mathbf{H}$  for  $\eta \in {}^\omega \lambda$
- (β) if  $\bar{f} = \langle f_\nu : \nu \in {}^\omega \lambda \rangle$  and  $f_\nu \in \mathbf{H}$  for every such  $\nu$  and  $\alpha < \lambda$  and  $\rho \in {}^\omega \lambda$  is increasing then for some increasing  $\eta \in {}^\omega \lambda$  we have  $\rho \triangleleft \eta$  and  $n < \omega \Rightarrow h_{\eta, n} = f_{\eta \upharpoonright n}$
- (γ) if  $\text{cf}(\lambda) > \aleph_0$  and  $E$  is a club of  $\lambda$  then we can add  $\cup \{\eta(n) : n < \omega\} \in E$ .

[Why? First assume  $\chi = \lambda$ . Let  $\langle \bar{g}_\alpha = \langle g_{\alpha, \ell} : \ell < n_\alpha \rangle : \alpha < \lambda \rangle$  enumerate  ${}^\omega \mathbf{H}$  such that for each  $\bar{g} \in {}^\omega \mathbf{H}$  the set  $\{\alpha < \lambda : \bar{g}_\alpha = \bar{g}\}$  is unbounded in  $\lambda$ . Now for  $\eta \in {}^\omega \lambda$  and  $n < \omega$  let  $h_{\eta, n} = g_{\eta(k), n}$  for every  $k$  large enough if well defined and  $g_{\eta \upharpoonright (n+1), n}$  otherwise. So clause (α) of ⊙<sub>3</sub> holds and as for clause (β) of ⊙<sub>3</sub>, let  $\bar{f} = \langle f_\nu : \nu \in {}^\omega \lambda \rangle$  be given,  $f_\nu \in \mathbf{H}$ .

Assume  $\rho \in {}^\omega \lambda$  is increasing. We choose  $\alpha_n$  by induction on  $n < \omega$  such that:

- ⊙<sub>4</sub> (α)  $\alpha_n = \rho(n)$  if  $n < \ell g(\rho)$

<sup>3</sup>Why? (As doubts were cast we shall elaborate.) For  $\bar{\eta} \in \Lambda_{\leq k(*)}$  let  $B'_{\bar{\eta}} = \{i : i < |B_{\bar{\eta}}|\}$  for  $\bar{\eta} \in \Lambda_{\leq k(*)}$  and let  $g_{\bar{\eta}}$  be a one-to-one function from  $B'_{\bar{\eta}}$  onto  $B_{\bar{\eta}}$ . Now assume that  $\langle h'_{\bar{\eta}} : \bar{\eta} \in \Lambda_{\leq k(*)} \rangle$  is as required in the claim for  $\langle B'_{\bar{\eta}} : \bar{\eta} \in \Lambda \rangle$  and define a function  $h_\eta$  with domain  $\text{Dom}(h'_\eta) = \{\bar{\eta} \upharpoonright \langle m, n \rangle : m \leq k(*) \text{ and } n < \omega\}$  such that  $h_{\bar{\eta}}(\bar{\eta} \upharpoonright \langle m, n \rangle) = g_{\bar{\eta}}(h'_\eta(\bar{\eta} \upharpoonright \langle m, n \rangle)) \in B_{\bar{\eta} \upharpoonright \langle m, n \rangle}$  for  $\bar{\eta} \in \Lambda, m \leq k(*), n < \omega$ . Define the function  $h'$  with domain  $\Lambda_{\leq k(*)}$  by  $h'(\bar{\eta}) = g_{\bar{\eta}}^{-1} \circ h$ , so  $h'$  is well defined with domain  $\Lambda_{\leq k(*)}$  such that  $h'(\bar{\eta}) \in B'_{\bar{\eta}}$ . By the choice of  $\langle h'_{\bar{\eta}} : \bar{\eta} \in \Lambda_{\leq k(*)} \rangle$  there is  $\bar{\eta} \in \Lambda$  such that  $m \leq k(*) \wedge n < \omega \Rightarrow h'(\bar{\eta} \upharpoonright \langle m, n \rangle) = h'(\bar{\eta} \upharpoonright \langle m, n \rangle)$ . But by the choice of  $h_{\bar{\eta}}, h'$  we have  $m \leq k(*) \wedge n < \omega \Rightarrow h_{\bar{\eta}}(\bar{\eta} \upharpoonright \langle m, n \rangle) = g_{\bar{\eta}}^{-1}(h'_{\bar{\eta}}(\bar{\eta} \upharpoonright \langle m, n \rangle)) = g_{\bar{\eta}}^{-1}(h'(\bar{\eta} \upharpoonright \langle m, n \rangle)) = h(\bar{\eta} \upharpoonright \langle m, n \rangle)$  as required.

modified:2009-03-08

(883) revision:2009-03-05

- ( $\beta$ )  $\alpha_n < \lambda$  and  $\alpha_n > \alpha_m$  if  $n = m + 1$   
 ( $\gamma$ ) if  $n \geq \ell g(\rho)$  then  $\alpha_n$  satisfies  $\bar{g}_{\alpha_n} = \langle f_{\langle \alpha_\ell : \ell < m \rangle} : m \leq n \rangle$ .

Now  $\eta =: \langle \alpha_n : n < \omega \rangle$  is as required in clause ( $\beta$ ) of  $\odot_3$ ; to get also clause ( $\gamma$ ) of  $\odot_3$  we should add in clause ( $\beta$ ) of  $\odot_4$  then  $\alpha_n > \min(E \setminus \alpha_m)$ .

Second, if  $\chi > \lambda$  but still  $\chi \leq \lambda^{\aleph_0}$ , let  $\langle \bar{g}_\alpha : \alpha < \chi^{\aleph_0} \rangle$  list  ${}^\omega \mathbf{H}$ , and let  $\mathbf{h}_n : \chi \rightarrow \lambda$  for  $n < \omega$  be such<sup>4</sup> that  $\alpha < \beta < \chi \Rightarrow (\forall^\infty n)(\mathbf{h}_n(\alpha) \neq \mathbf{h}_n(\beta))$  and let  $\text{cd} : \lambda \rightarrow {}^\omega \lambda$  be one to one onto. Now for  $\eta \in {}^\omega \lambda$  and  $n < \omega$  let  $h_{\eta,n}$  be  $g_\alpha$  where  $\alpha$  is the unique ordinal  $\alpha < \chi$  such that for every  $k < \omega$  large enough  $(\text{cd}(\eta(k)))(n) = \mathbf{h}_n(\alpha)$  so in particular  $\langle \ell g(\text{cd}(\eta(k)) : k < \omega) \rangle$  is going to infinity or  $h_{\eta,n}$  is not well defined; in fact, we can use only the case  $\ell g(\text{cd}(\eta(k))) = k$ ; stipulating  $h_{\eta,n} \in {}^\omega \{0\}$  when not defined. So we have defined  $\langle h_{\eta,n} : \eta \in {}^\omega \lambda, n < \omega \rangle$ . Now we immitate the previous argument: clause ( $\beta$ ) of  $\otimes_2$  holds.

Next we shall define  $\bar{\alpha}^{k(*)} = \langle \alpha_{\bar{\eta},m,n}^{k(*)} : \bar{\eta} \in \Lambda_{k+1}, m \leq k(*), n < \omega \rangle$  as required; so let  $\bar{\eta} = \langle \eta_\ell : \ell \leq k(*) \rangle \in \Lambda_{k(*)}$  we define  $\bar{\alpha}_{\bar{\eta}}^{k(*)} = \langle \alpha_{\bar{\eta},m,n}^{k(*)} : m \leq k(*), n < \omega \rangle$  as follows:

- $\odot_5$  if  $\eta_{k(*)} \in {}^\omega \lambda$  and  $\langle \eta_0, \dots, \eta_{k(*)-1} \rangle \in \Lambda_k$  then for  $m \leq k(*)$  and  $n < \omega$   
 ( $\alpha$ ) if  $m = k(*)$  then  $\alpha_{\bar{\eta},m,n}^{k(*)} = h_{\eta_{k(*)},n}(\langle \eta_0, \dots, \eta_{k(*)-1} \rangle) < \lambda_m$   
 ( $\beta$ ) if  $m < k(*)$ , i.e.  $m \leq k$  then  $\alpha_{\bar{\eta},m,n}^{k(*)} = \alpha_{\bar{\eta} \upharpoonright k(*),m,n}^k < \lambda_m$ .

Clearly  $\alpha_{\bar{\eta},m,n}^{k(*)} < \lambda_m$  in all cases, as required, (in clause (a),(b),(c) of 2.1(2) and (e) of 2.1(3)). But we still have to prove that  $\langle \bar{\alpha}_{\bar{\eta},m,n}^{k(*)} : \bar{\eta} \in \Lambda^{k+1}, m \leq k(*), n < \omega \rangle$  witness  $\text{QR}(\mathbf{x}^{k(*)}, \chi)$ , see Definition 1.3(2) this suffices for 2.1(2), little more is needed for 2.1(3); just using ( $\gamma$ ) of  $\odot_3$  and the induction hypothesis.

Why does this hold? Let  $h$  be a function with domain  $\Lambda_{\leq k(*)}^{\mathbf{x}^{k(*)}}$  as in part (3) and  $\alpha_\ell^* < \lambda_\ell$  for  $\ell \leq k(*)$ .

For  $\nu \in {}^\omega \lambda$  let  $f_\nu : \Lambda_k \rightarrow \lambda = \lambda_{k(*)}$  be defined by:  $f_\nu(\langle \eta_\ell : \ell \leq k \rangle) =: h(\langle \eta_\ell : \ell \leq k \rangle \hat{\ } \langle \nu \rangle)$ . So by  $\odot_3$  above for some increasing  $\eta_{k(*)}^* \in {}^\omega \lambda$  we have  $\eta_{k(*)}^*(0) = \alpha_{k(*)}^*$  and

$$\odot_6 \quad n < \omega \Rightarrow f_{\eta_{k(*)}^*} \upharpoonright n = h_{\eta_{k(*)}^*,n}.$$

Now substituting the definition of  $\bar{f}$  we have

$$\odot_7 \quad \langle \eta_0, \dots, \eta_k \rangle \in \Lambda_k \wedge n < \omega \Rightarrow h_{\eta_{k(*)}^*,n}(\eta_0, \dots, \eta_k) = h(\langle \eta_0, \dots, \eta_k, \eta_{\eta_{k(*)}^*}^* \upharpoonright n \rangle).$$

<sup>4</sup>recall  $(\forall^\infty N)$  means “for every large enough  $n < \omega$ ”

Substituting the definition of  $\bar{\alpha}^k$  we have

$$\odot_8 \text{ if } \langle \eta_0, \dots, \eta_k \rangle \in \Lambda_k \text{ and } n < \omega \text{ then } \alpha_{\langle \eta_0, \dots, \eta_k, \eta_{k(*)}^* \rangle}^{k(*)} = h(\langle \eta_0, \dots, \eta_k, \eta_{k(*)}^* \upharpoonright n \rangle).$$

Now we define a function  $h'$  with domain  $\Lambda_{\leq k}^{\mathbf{x}^k}$  by: if  $\bar{\eta} \in \Lambda_{\leq k}^{\mathbf{x}^k}$  then  $h'(\bar{\eta}) = h(\bar{\eta} \hat{\ } \langle \eta_{k(*)}^* \rangle)$ .

So by the choice of  $\bar{\alpha}^k$  in  $\odot_2$  we can find  $\langle \eta_0^*, \dots, \eta_k^* \rangle \in \Lambda_k$  with no repetitions such that  $\eta_\ell^*(0) = \alpha_\ell^*$  for  $\ell \leq k$  and in  $\odot_2$

$$\odot_9 \text{ } m \leq k \wedge n < \omega \Rightarrow \alpha_{\langle \eta_0^*, \dots, \eta_k^* \rangle, m, n}^k = h'(\langle \eta_0^*, \dots, \eta_k^* \rangle \upharpoonright (m, n)).$$

$$\text{Let } \bar{\eta}^* = \langle \eta_0^*, \dots, \eta_k^*, \eta_{k+1}^* \rangle, \bar{\eta}' = \langle \eta_0^*, \dots, \eta_i^* \rangle.$$

Note that

$$\odot_{10} \text{ if } m \leq k, n < \omega \text{ then } h'(\bar{\eta}' \upharpoonright \langle k, m \rangle) = h((\bar{\eta}' \upharpoonright \langle m, n \rangle) \hat{\ } \langle \eta_{k(*)}^* \rangle) = h(\bar{\eta}^* \upharpoonright \langle m, n \rangle).$$

Now by  $\odot_9 + \odot_{10}$  and  $\odot_5(\beta)$  this means

$$\odot_{11} \text{ if } m \leq k \text{ and } n < \omega \text{ then } \alpha_{\bar{\eta}^*, m, n}^{k(*)} = h(\bar{\eta}^* \upharpoonright \langle k, m \rangle).$$

So by putting together  $\odot_8 + \odot_{11}$  we are clearly done, i.e. we can check that  $\langle \eta_0^*, \dots, \eta_k^*, \eta_{k(*)}^* \rangle$  is as required.  $\square_{2.1}$

**2.2 Conclusion.** For every  $k < \omega$  there is an  $\aleph_{k+1}$ -free abelian group  $G$  of cardinality  $\beth_{k+1}$  and pure (non-zero) subgroup  $\mathbb{Z}z \subseteq G$  such that  $\mathbb{Z}z$  is not a direct summand of  $G$ .

*Proof.* Let  $\chi = 2^{\aleph_0}$  and  $\mathbf{x}$  be a combinatorial  $k$ -parameter as guaranteed by 2.1. Now by 2.3(2) below we can expand  $\mathbf{x}$  to an abelian group  $k$ -parameter, so  $G_{\mathbf{x}}$  is as required.

**2.3 Claim.** 1) If  $\mathbf{x}$  is a combinatorial  $k$ -parameter such that  $\text{Qr}(\mathbf{x}, 2^{\aleph_0})$  then for some  $\mathbf{a}, \mathbf{y} := (\mathbf{x}, \mathbf{a})$  is an abelian group  $k$ -parameter such that  $h \in \text{Hom}(G_{\mathbf{y}}, \mathbb{Z}) \Rightarrow h(z) = 0$ .

2) For every  $k$  there is an  $\aleph_{k+1}$ -free abelian group  $G$  of cardinality  $\beth_{k+1}$  and  $z \in G$  a pure  $z \in G$  as above.

*Proof.* 1) Let  $\bar{\alpha}$  witness  $\text{Qr}(\mathbf{x}, 2^{\aleph_0})$ . We define a function  $\iota: \text{Ord} \rightarrow \mathbb{Z}$  by:  $\iota(\alpha)$  in  $\alpha$  if  $\alpha < \omega$ , is  $-n$  if  $\alpha = \omega + n < \omega + \omega$  and is zero otherwise. For each  $\bar{\eta} \in \Lambda^{\mathbf{x}}$  we

shall choose a sequence  $\langle \mathbf{a}_{\bar{\eta},n} : n < \omega \rangle$  of integers such that for any  $b \in \mathbb{Z} \setminus \{0\}$  for no  $\bar{c} \in {}^\omega \mathbb{Z}$  do we have:

$\boxtimes_{\bar{\eta}}$  for each  $n < \omega$  we have

$$n!c_{n+1} = c_n + \mathbf{a}_{\bar{\eta},n}b + \Sigma\{\iota(\alpha_{\bar{\eta},m,n}) : m \leq k(*)\}.$$

This is easy: for each pair  $(b, c_0) \in \mathbb{Z} \times \mathbb{Z}$  the set of  $\langle \mathbf{a}_n : n < \omega \rangle \in {}^\omega \mathbb{Z}$  such that there is at least one sequence (and always at most one sequence)  $\langle c_0, c_1, c_2, \dots \rangle$  of integers such that  $\boxtimes_{\bar{\eta}}$  holds for them, is meagre, even no-where dense so the choice of  $\langle \mathbf{a}_{\bar{\eta},n} : n < \omega \rangle$  is possible.

Now toward contradiction assume that  $h$  is a homomorphism from  $G_{\mathbf{x}}$  to  $z\mathbb{Z}$  such that  $h(z) = bz, b \in \mathbb{Z} \setminus \{0\}$ . We define  $h' : \Lambda_{\leq k}^{\mathbf{x}} \rightarrow \chi$  by  $h'(\bar{\eta}) = n$  if  $n < \omega$  and  $h(x_{\bar{\eta}}) = nz$  and  $h'(y_{\bar{\eta}}) = \omega + n$  if  $n < \omega$  and  $h(x_{\bar{\eta}}) = (-n)z$ .

By the choice of  $\bar{\alpha}$ , for some  $\bar{\eta} \in \Lambda^{\mathbf{x}}$  we have:  $m \leq k \wedge n < \omega \Rightarrow h'(\bar{\eta} \upharpoonright \langle m, n \rangle) = \alpha_{\bar{\eta},m,n}$ . Hence  $h(x_{\bar{\eta} \upharpoonright \langle m, n \rangle}) = \iota(\alpha_{\bar{\eta},m,n})z$  for  $m \leq k, n < \omega$ .

Let  $c_n \in \mathbb{Z}$  be such that  $h(y_{\bar{\eta},n}) = c_n z$ . Now the equation  $\boxtimes_{\bar{\eta},n}$  in Definition 1.6 is mapped to the  $n$ -th equation in  $\boxtimes_{\bar{\eta}}$ , so an obvious contradiction.

2) By part (1) and 2.2.

$\square_{2.3}$

*2.4 Remark.* 1) We can replace  $\chi$  by a set of cardinality  $\chi$  in Definition 1.3. Using  $\mathbb{Z}z$  instead of  $\chi$  simplify the notation in the proof of 2.3.

2) We have not tried to save in the cardinality of  $G$  in 2.3(2), using as basic of the induction the abelian group of cardinality  $\aleph_0$  or  $\aleph_1$ .

**2.5 Claim.** 1) If  $\chi_0 = \chi_0^{\aleph_0}, \chi_{m+1} = 2^{\chi_m}$  and  $\lambda_m = \chi_m$  for  $m \leq k$  for the  $\bar{\chi}$ -full combinatorial  $k$ -parameter  $\mathbf{x}$ , the  $(\mathbf{x}, \bar{\chi})$ -black box exist.

*2.6 Conclusion.* Assume  $\mu_0 < \dots < \mu_{k(*)}$  are strong limit of cofinality  $\aleph_0$  (or  $\mu_0 = \aleph_0$ ),  $\lambda_\ell = \mu_\ell^+, \chi_\ell = 2^{\mu_\ell}$ .

Then in 2.1 for  $\bar{\eta} \in \Lambda^{\mathbf{x}}$  we can let  $h_{\bar{\eta},m}$  has domain  $\{\bar{\nu} \in \Lambda_m^{\mathbf{x}} : [\nu_\ell = \eta_\ell \text{ for } \ell = m+1, \dots, k(*)]\}$ .

§3 CONSTRUCTING ABELIAN GROUPS FROM COMBINATORIAL PARAMETERS

**3.1 Definition.** 1) We say  $F$  is a  $\mu$ -regressive function on a combinatorial parameter  $\mathbf{x} \in K_{k(*)}^{\text{cb}}$  when  $S^{\mathbf{x}}$  is a set of ordinals and:

- (a)  $\text{Dom}(F)$  is  $\Lambda^{\mathbf{x}}$
- (b)  $\text{Rang}(F) \subseteq [\Lambda^{\mathbf{x}} \cup \Lambda_{\leq k(*)}^{\mathbf{x}}]^{\leq \aleph_0}$
- (c) for every  $\bar{\eta} \in \Lambda^{\mathbf{x}}$  and  $m \leq k(*)$  we<sup>5</sup> have  $\sup \text{Rang}(\eta_m) > \sup(\cup\{\text{Rang}(\nu_m) : \bar{\nu} \in F(\bar{\eta})\})$ ; note  $\bar{\nu}_\ell \in \Lambda^{\mathbf{x}}$  or  $\bar{\nu} \in \Lambda_{\leq k(*)}^{\mathbf{x}}$  as  $F(\bar{\eta})$  is a set of such objects.

1A) We say  $F$  is finitary when  $F(\bar{\eta})$  is finite for every  $\bar{\eta}$ .

1B) We say  $F$  is simple if  $\eta_{k(*)}(0)$  determined  $F(\bar{\eta})$  for  $\bar{\eta} \in \Lambda^{\mathbf{x}}$ .

2) For  $\mathbf{x}, F$  as above and  $\Lambda \subseteq \Lambda^{\mathbf{x}}$  we say that  $\Lambda$  is free for  $(\mathbf{x}, F)$  when:  $\Lambda \subseteq \Lambda^{\mathbf{x}}$  and there is a sequence  $\langle \bar{\eta}^\alpha : \alpha < \alpha(*) \rangle$  listing  $\Lambda' = \Lambda \cup \bigcup\{F(\bar{\eta}) : \bar{\eta} \in \Lambda\}$  and sequence  $\langle \ell_\alpha : \alpha < \alpha(*) \rangle$  such that

- (a)  $\ell_\alpha \leq k(*)$
- (b) if  $\alpha < \alpha(*)$  and  $\bar{\eta}^\alpha \in \Lambda$  then  $F(\bar{\eta}^\alpha) \subseteq \{\bar{\eta}^\beta : \beta < \alpha\} \cup \{\bar{\eta}^\gamma \upharpoonright \langle m, n \rangle : \gamma < \alpha \text{ is such that } \bar{\eta}^\gamma \in \Lambda^{\mathbf{x}} \text{ and } n < \omega, m \leq k(*)\}$
- (c) if  $\alpha < \alpha(*)$  and  $\bar{\eta}^\alpha \in \Lambda$  then for some  $n < \omega$  we have  $\bar{\eta}^\alpha \upharpoonright \langle \ell_\alpha, n \rangle \notin \{\bar{\eta}^\beta \upharpoonright \langle \ell_\alpha, n \rangle : \beta < \alpha, \bar{\eta}^\beta \in \Lambda\} \cup \{\bar{\eta}^\beta : \beta < \alpha\}$ .

3) We say  $\mathbf{x}$  is  $\theta$ -free for  $F$  is  $(\mathbf{x}, F)$  is  $\mu$ -free when  $\mathbf{x}, F$  are as in part (1) and every  $\Lambda \subseteq \Lambda^{\mathbf{x}}$  of cardinality  $< \theta$  is free for  $(\mathbf{x}, F)$ .

**3.2 Claim.** 1) If  $\mathbf{x} \in K_{k(*)}^{\text{cb}}$  and  $F$  is a regressive function on  $\mathbf{x}$  then  $(\mathbf{x}, F)$  is  $\aleph_{k(*)+1}$ -free provided that  $F$  is finitary or simple.

2) In addition: if  $k \leq k(*)$ ,  $\Lambda \subseteq \Lambda^{\mathbf{x}}$  has cardinality  $\leq \aleph_k$  and  $\bar{u} = \langle u_{\bar{\eta}} : \bar{\eta} \in \Lambda \rangle$  satisfies  $u_{\bar{\eta}} \subseteq \{0, \dots, k(*)\}$ ,  $|u_{\bar{\eta}}| > k$ , then we can find  $\langle \bar{\eta}^\alpha : \alpha < \aleph_k \rangle, \langle \ell_\alpha : \alpha < \aleph_k \rangle, \langle n_\alpha : \alpha < \aleph_k \rangle$  such that:

- (a)  $\Lambda \subseteq \{\bar{\eta}^\alpha : \alpha < \aleph_k\}$
- (b) if  $\bar{\eta}_\alpha \in \Lambda^{\mathbf{x}}$  then  $\ell_\alpha \in u_{\bar{\eta}^\alpha}, n_\alpha < \omega$
- (c)  $\bar{\eta}^\alpha \upharpoonright \langle \ell_\alpha, n_\alpha \rangle \notin \{\bar{\eta}^\beta \upharpoonright \langle \ell_\alpha, n_\alpha \rangle : \beta < \alpha\} \cup \{\bar{\eta}^\beta : \beta < \alpha\}$ .

*Remark.* We may wonder:

---

<sup>5</sup>actually, suffice to have it for  $\ell = k(*)$

Ruedeger Question: Assume  $F(\bar{\eta}) \in [\Lambda_{\leq k(*)}]^{\leq \aleph_0}$  for  $\bar{\eta} \in \Lambda^{\mathbf{x}}$  is as in Definition 3.1. Is this O.K. in the proof of 3.2, particularly Case 1?

Answer: Seems not. Assume  $\bar{\nu} \neq \bar{\rho} \in \Lambda$  and

- (A)  $u_{\bar{\rho}} = \{\ell_1\}, F(\bar{\nu}) = \{\bar{\rho} \upharpoonright \langle \ell_1, n \rangle : n < \omega\}$   
 (B)  $u_{\bar{\nu}} = \{\ell_2\}, F(\bar{\rho}) = \{\bar{\nu} \upharpoonright \langle \ell_2, n \rangle : n < \omega\}$ .

So if  $(\nu, \bar{\rho}) = (\eta_{\alpha_4}, \eta_{\alpha_2})$ , we have  $\alpha_0 \neq \alpha_1$  as  $\bar{\nu} \neq \bar{\rho}, \neg(\alpha_1 < \alpha_2)$  by (B), and  $\neg(\alpha_2 < \alpha_1)$  by (A).

*Proof.* 1) Follows by part (2) for the case  $k = k(*), u_{\bar{\eta}} = \{0, \dots, k(*)\}$  for every  $\bar{\eta} \in \Lambda$ .

2) So we are assuming  $\mathbf{x} \in K_{k(*)}^{\text{cb}}, F$  is a regressive function on  $\mathbf{x}$  which is finitary or simple,  $k \leq k(*), \Lambda \subseteq \Lambda^{\mathbf{x}}$  has cardinality  $\leq \aleph_k$  and without loss of generality  $\Lambda$  is closed under  $\bar{\eta} \mapsto F(\bar{\eta}) \cap \Lambda^{\mathbf{x}}$ . We prove this by induction on  $k$ .

Case 1:  $k = 0$ .

Subcase 1A: Ignoring  $F$ .

Let  $\langle \bar{\eta}^\alpha : \alpha < |\Lambda| \rangle$  list  $\Lambda$  with no repetitions (so  $\alpha < |\Lambda| \Rightarrow \alpha < \aleph_k = \aleph_0$ ). Now  $\alpha < |\Lambda| \Rightarrow u_{\bar{\eta}^\alpha} \neq \emptyset$  and let  $\ell_\alpha = \min(u_{\bar{\eta}^\alpha}) \leq k(*)$ . Hence for each  $\alpha < |\Lambda|$  we know that  $\beta < \alpha \Rightarrow \bar{\eta}^\beta \neq \bar{\eta}^\alpha$ , hence for some  $n = n_{\alpha, \beta} < \omega$  we have  $\bar{\eta}^\beta \upharpoonright \langle \ell_\alpha, n_{\alpha, \beta} \rangle \neq \bar{\eta}^\alpha \upharpoonright \langle \ell_\alpha, n_{\alpha, \beta} \rangle$ .

Let  $n_\alpha = \sup\{n_{\alpha, \beta} : \beta < \alpha\}$ , it is  $< \omega$  as  $\alpha < \omega$ . Now  $\langle (\ell_\alpha, n_\alpha) : \alpha < |\Lambda| \rangle$  is as required.

Subcase 1B:  $\bar{\eta} \in \Lambda \Rightarrow F(\bar{\eta})$  is finite<sup>6</sup>.

Let  $\langle \bar{\eta}^\alpha : \alpha < |\Lambda| \rangle$  list  $\Lambda$ , we choose  $w_j$  by induction on  $j \leq j(*), j(*) \leq \omega$  such that:

- ⊗ (a)  $w_j \subseteq |\Lambda|$  is finite for  $j < \omega$   
 (b)  $j \in w_{j+1}$   
 (c) if  $\alpha \in w_j$  then  $F(\bar{\eta}^\alpha) \cap \Lambda \subseteq \{\bar{\eta}^\alpha : \beta \in w_j\}$   
 (d)  $w_{j(*)} = |\Lambda|$  and  $w_0 = \emptyset$   
 (e)  $w_j \subseteq w_{j+1}$   
 (f) if  $j(*) = \omega$  then  $w_{j(*)} = \cup\{w_j : j < j(*)\}$ .

<sup>6</sup>If we assume for  $\bar{\eta} \in \Lambda \Rightarrow F(\bar{\eta}) \subseteq \Lambda_{\leq k(*)}$  then any list  $\langle \bar{\eta}^\alpha : \alpha < |\Lambda| \rangle$  with no repetitions and  $\bar{\ell} = \langle \ell_\alpha : \alpha < |\Lambda| \rangle, \ell_\alpha \in u_{\bar{\eta}^\alpha}$  will do. Why? Because  $Y_\alpha := \cup\{F(\bar{\eta}^\beta) : \beta < \alpha\}$  is a finite subset of  $\Lambda_{\leq k(*)}$ . Now for  $\alpha < |\Lambda|$  the set  $u_\alpha^1 := \{n < \omega : \bar{\eta}^\alpha \upharpoonright \langle \ell_\alpha, n \rangle \text{ belongs to } Y_\alpha\}$  is finite, and also for each  $\beta < \alpha$  the set  $u_\alpha^r, Y_{\alpha, \beta} := \{n < \omega : \bar{\eta}^\alpha \upharpoonright \langle \ell_\alpha, n \rangle = \bar{\eta}^\beta \upharpoonright \langle \ell_\alpha, n \rangle\}$  is finite. As  $\alpha$  is finite we can find  $n = n_\alpha \in \omega \setminus Y_\alpha \setminus \cup\{Y_{\alpha, \beta} : \beta < \alpha\}$ . Now  $\langle n_\alpha : \alpha < |\Lambda| \rangle$  is as required.



No problem to do this; for clause (c) use “ $F$  is regressive, the ordinals well ordered but we elaborate. Assume that the finite  $w_j \subseteq |\Lambda|$  has been chosen. We define  $w_{j,m}$  by induction on  $m$  such that  $w_{j,m} \subseteq |\Lambda|$  is finite and  $\subseteq$ -increasing with  $m$ . For  $m = 0$  let  $w_{j,m} = w_j \cup \{\alpha\}$ . If  $w_{j,m}$  is defined let

$$w_{j,m+1} = w_{j,m} \cup \{\beta < |\Lambda| : \text{for some } \alpha \in w_{j,m} \text{ we have } \bar{\eta}^\beta \in F(\bar{\eta}^\alpha) \cap \Lambda\}.$$

As  $w_{j,m}$  is finite and  $\subseteq |\Lambda|$  and each  $F(\bar{\eta}^\alpha)$  is finite and  $\subseteq \{\bar{\eta}^\gamma : \gamma < |\Lambda|\}$  clearly  $w_{j,m+1}$  is finite  $\subseteq |\Lambda|$ .

Lastly, we let  $w_{j+1}$  be  $\cup\{w_{j,m} : m < \omega\}$ . If it is finite we have carried the inductive step on  $j$ . If not, then  $\langle w_{j,m} : m < \omega \rangle$  is  $\subset$ -increasing and we let  $\gamma_{j,m} = \sup\{\eta_{\alpha,0}(i) : i < \omega, \alpha \in w_{j,m+1} \setminus w_{j+m}\}$  and it suffices to prove

$$(*) \quad \gamma_{j,m} > \gamma_{j,m+1} \text{ (both are ordinals!).}$$

Why  $(*)$  is true? As by the definition of  $\gamma_{j,m+1}$  for some  $i_* < \omega$  and  $\beta_* \in w_{j,m+2} \setminus w_{j,m+1}$  we have  $\eta_{\beta_*,0}(i_*) = \gamma_{j,m+1}$ . By the definition of  $w_{j,m+2}$  as  $\beta_* \notin w_{j,m+1}$ , there is  $\alpha_* \in w_{j,m+1}$  such that  $\bar{\eta}^{\beta_*} \in F(\bar{\eta}^{\alpha_*}) \cap \Lambda$ .

As  $\beta_* \notin w_{j,m+1}$  necessarily  $\alpha_* \notin w_{j,m}$  hence by the definition of  $\gamma_{j,m}$  we know that  $(\forall i < \omega)(\eta_{\alpha_*,0}(i) < \gamma_{j,m})$ . By clause (c) of Definition 3.1(1) as  $\bar{\eta}^{\beta_*} \in F(\bar{\eta}^{\alpha_*})$  we know that  $\eta_{\beta_*,0}(i_*) < \sup\{\eta_{\alpha_*,0}(i) : i < \omega\}$ . By the last two sentences we are done proving  $(*)$ , so we are done defining  $w_{j+1}$  hence we finish justifying  $\circledast$ .

Now let  $\langle \beta(j,i) : i < i_j^* \rangle$  list  $w_{j+1} \setminus w_j$  such that: if  $i_1, i_2 < i_j^*$  and  $\bar{\eta}^{\beta(j,i_1)} \in F(\bar{\eta}^{\beta(j,i_2)})$  then  $i_1 < i_2$ ; we prove existence by  $F$  being regressive. Let  $\langle \bar{\nu}_{j,i} : i < i_j^{**} \rangle$  list  $\cup\{F(\bar{\eta}^\alpha) : \alpha \in w_{j+1} \setminus w_j\} \setminus \Lambda^{\mathbf{x}} \setminus \{F(\bar{\eta}^\alpha) : \alpha \in w_j\}$ .

Let  $\alpha_j^* = \Sigma\{i_{j(1)}^{**} + i_{j(1)}^* : j(1) < j\}$ . Now we choose  $\bar{\rho}_\varepsilon$  for  $\varepsilon < \alpha_j^*$  for  $j < j^*$  as follows:

- (a)  $\rho_{\alpha_j^*+i} = \nu_{j,i}$  if  $i < i_j^{**}$
- (b)  $\bar{\rho}_{\alpha_j^*+i_j^{**}+i} = \bar{\eta}^{\beta(j,i)}$  if  $i < i_j^*$ .

Lastly, we choose  $n_{\alpha_j^*+i} < \omega$  for  $i < i_j^*$  as in case 1A.

Now check.

Subcase 1C:  $F$  is simple.

Note that  $F(\bar{\eta})$  when defined is determined by  $\eta_{k(*)}(0)$  and is included in  $\{\bar{\nu} \in \Lambda_{\leq k(*)}^{\mathbf{x}} \cup \Lambda^{\mathbf{x}} : \sup \text{Rang}(\nu_{k(*)}) < \eta_{k(*)}(0)\}$ . So let  $u = \{\eta_{k(*)}(0) : \bar{\eta} \in \Lambda\}$  and  $u^* = u \cup \{\sup(u) + 1\}$  and for  $\alpha \in u$  let  $\Lambda_\alpha = \{\bar{\eta} \in \Lambda : \eta_{k(*)}(0) = \alpha\}$  and for  $\alpha \in u^*$

let  $\Lambda_{<\alpha} = \cup\{\Lambda_\beta : \beta \in u\}$ . Now by induction on  $\beta \in u^*$  we choose  $\langle (\bar{\eta}^\varepsilon, \ell_\varepsilon) : \varepsilon < \varepsilon_\beta \rangle$  such that it is a required for  $\Lambda_{<\beta}$ . For  $\beta = \min(u)$  this is trivial and if  $\text{otp}(u \cap \beta)$  is a limit ordinal this is obvious. So assume  $\alpha = \max(u \cap \beta)$ , we use Subcase 1A on  $\Lambda_\alpha$ , and combine them naturally promising  $\ell_\alpha = k(*) \Rightarrow n_\alpha > 1$ .

Case 2:  $k = k_* + 1$  and  $|\Lambda| = \aleph_k$ .

Let  $\langle \Lambda_\varepsilon : \varepsilon < \aleph_k \rangle$  be  $\subseteq$ -increasing continuous with union  $\Lambda$ ,  $|\Lambda_{1+\varepsilon}| = \aleph_{k_*}$ ,  $\Lambda_0 = \emptyset$ , each  $\Lambda_\varepsilon$  closed enough, mainly:

- ⊗<sub>1</sub> if  $\bar{\eta}^i \in \Lambda_\varepsilon$  for  $i < i(*) < \omega$ ,  $\bar{\rho} \in \Lambda$  and  $\{\rho_\ell : \ell \leq k(*)\} \subseteq \{\eta_\ell^i : \ell \leq k(*), i < i(*)\}$  then  $\bar{\rho} \in \Lambda_\varepsilon$
- ⊗<sub>2</sub>  $\Lambda_\varepsilon$  is closed under  $\bar{\eta} \mapsto F(\bar{\eta}) \cap \Lambda^\times$ .

Next

- ⊙ if  $\varepsilon < \aleph_k$ ,  $\bar{\eta} \in \Lambda_{\varepsilon+1} \setminus \Lambda_\varepsilon$  then  $u'_{\bar{\eta}} = \{\ell \in u_{\bar{\eta}} : \text{for every or just some } n < \omega \text{ for some } \bar{\nu} \in \Lambda_\varepsilon \text{ we have } \bar{\eta} \upharpoonright \langle \ell, n \rangle = \bar{\nu} \upharpoonright \langle \ell, n \rangle\}$  has at most one member.

[Why? So assume toward contradiction that  $\bar{\eta} \in \Lambda_{\varepsilon+1}$  and  $\ell(1) \neq \ell(2)$  belong to  $u'_{\bar{\eta}}$ . Hence by the definition of  $u'_{\bar{\eta}}$  there are  $\bar{\nu}^1, \bar{\nu}^2 \in \Lambda_\varepsilon$  and  $n_1, n_2 < \omega$  such that  $\bar{\eta} \upharpoonright \langle \ell_1, n_1 \rangle \in \bar{\nu}^1 \upharpoonright \langle \ell_1, n_1 \rangle$  and  $\bar{\eta} \upharpoonright \langle \ell_2, n_2 \rangle = \bar{\nu}^2 \upharpoonright \langle \ell_2, n_2 \rangle$ . Now  $m \leq k(*) \Rightarrow$  for some  $i \in \{1, 2\}$ ,  $m \leq \ell_i \Rightarrow$  for some  $i \in \{1, 2\}$ ,  $\eta_m$  is  $(\bar{\eta} \upharpoonright \langle \ell_i, n_i \rangle)_m \Rightarrow \eta_m \in \{\rho_\ell : \bar{\rho} \in \Lambda_\varepsilon\}$ . Hence  $\{\eta_\ell : \ell \leq k(*)\} \subseteq \{\rho_\ell : \ell \leq k(*) \text{ and } \bar{\rho} \in \Lambda_\varepsilon\}$ . So by ⊗<sub>1</sub> we have  $\bar{\eta} \in \Lambda_\varepsilon$ , so we are done.]

Apply the induction hypothesis to  $\Lambda_{\varepsilon+1} \setminus \Lambda_\varepsilon$  for each  $\varepsilon$  and get  $\langle (\bar{\eta}^{\varepsilon, \alpha}, \ell_{\varepsilon, \alpha, n_{\varepsilon, \alpha}}) : \alpha < \alpha(\varepsilon) \rangle$  such that  $\bar{\eta}^{\varepsilon, \alpha} \upharpoonright \langle \ell_{\varepsilon, \ell, n_{\varepsilon, \alpha}} \rangle \notin \{\bar{\eta}^{\varepsilon, \beta} \upharpoonright \langle \ell_{\varepsilon, \beta, n_{\varepsilon, \beta}} \rangle : \beta < \alpha\}$ .

Let  $\alpha_* = \Sigma\{\alpha(\varepsilon) : \varepsilon < |\Lambda|\}$  and  $\alpha = \Sigma\{\alpha(\zeta) : \zeta < \varepsilon\} + \beta, \beta < \alpha(\varepsilon)$  let  $\eta^\alpha = \eta^{\varepsilon, \beta}, \ell_\alpha = \ell_{\varepsilon, \beta}, \eta_\alpha = \eta_{\varepsilon, \beta}$ . I.e. we combine but for  $\Lambda_{\varepsilon+1} \setminus \Lambda_\varepsilon$  we use  $\langle u_{\bar{\eta}} \setminus u'_{\bar{\eta}} : \bar{\eta} \in \Lambda_{\varepsilon+1} \setminus \Lambda_\varepsilon \rangle$ , so  $|u_{\bar{\eta}} \setminus u'_{\bar{\eta}}| \geq k - 1 = k_*$ . □<sub>3.2</sub>

**3.3 Definition.** For a combinatorial parameter  $\mathbf{x}$  we define  $\mathcal{G}_{\mathbf{x}}$ , the class of abelian groups derived from  $\mathbf{x}$  as follows:  $G \in \mathcal{G}_{\mathbf{x}}$  if there is a simple (or finitary) regressive  $F$  on  $\Lambda^\times$  and  $G$  is generated by  $\{y_{\bar{\eta}, n} : \eta \in \Lambda^\times, n < \omega\} \cup \{x_{\bar{\eta}} : \bar{\eta} \in \Lambda_{\leq k(*)}^\times\}$  freely except

$$\boxtimes_{\bar{\eta}, n} (n!)y_{\bar{\eta}, n+1} = y_{\bar{\eta}, n} + b_{\bar{\eta}, n}z_{\bar{\eta}, n} + \sum\{x_{\bar{\eta} \upharpoonright \langle m, n \rangle} : m \leq k(*)\}$$

where

- ⊙ (a)  $b_{\bar{\eta}, n} \in \mathbb{Z}$
- (b)  $z_{\bar{\eta}, n}$  is a linear combination of

$$\{x_{\bar{\nu}} : \bar{\nu} \in F(\bar{\eta}) \setminus \Lambda^{\mathbf{x}}\} \cup \{y_{\bar{\eta},n} : \bar{\eta} \in F(\bar{\eta}) \cap \Lambda^{\mathbf{x}} \text{ and } (\forall m \leq k(*))(\bar{\eta} \upharpoonright \langle m, n \rangle) \in F(\bar{\eta})\}.$$

**3.4 Claim.** *If  $\mathbf{x} \in K_{k(*)}^{\text{cb}}$  and  $G \in \mathcal{G}_{\mathbf{x}}$  (i.e.  $G$  is an abelian group derived from  $\mathbf{x}$ ), then  $G$  is  $\aleph_{k(*)+1}$ -free.*

*Proof.* We use claim 3.2. So let  $H$  be a subgroup of  $G$  of cardinality  $\leq \aleph_{k(*)}$ . We can find  $\Lambda$  such that

- (\*) (a)  $\Lambda \subseteq \Lambda^{\mathbf{x}}$  has cardinality  $\leq \aleph_{k(*)}$
- (b) every equation which  $X_{\Lambda} = \{x_{\bar{\eta} \upharpoonright \langle m, n \rangle}, y_{\bar{\eta},n} : m \leq k(*), n < \omega, \bar{\eta} \in \Lambda\}$  satisfies in  $G$ , is implied by the equations from  $\Gamma_{\Lambda} = \cup \{\boxtimes_{\bar{\eta},n} : \bar{\eta} \in \Lambda\}$
- (c)  $H \subseteq G_{\Lambda} = \langle x_{\bar{\eta} \upharpoonright \langle m, n \rangle}, y_{\bar{\eta},n} : \bar{\eta} \in \Lambda, m \leq k(*), n < \omega \rangle_G$
- (d) if  $\bar{\eta} \in \Lambda$  then  $F(\bar{\eta})$  is included in  $\Lambda \cup \{\bar{\nu} \upharpoonright \langle \ell, n \rangle : \bar{\nu} \in \Lambda, \ell \leq k(*) \text{ and } n < \omega\}$ .

So it suffices to prove that  $G_{\Lambda}$  is a free (abelian) group.

Let the sequence  $\langle (\bar{\eta}^{\alpha}, \ell_{\alpha}) : \alpha < \alpha(*) \rangle$  be as proved to exist in 3.2. Let  $\mathcal{U} = \{\alpha < \alpha(*) : \bar{\eta}^{\alpha} \in \Lambda\} \cup \{\alpha(*)\}$  and for  $\alpha \leq \alpha(*)$  let  $X_{\alpha}^0 = \{x_{\bar{\eta}^{\beta} \upharpoonright \langle m, n \rangle} : \beta \in \alpha \cap \mathcal{U}, m \leq k(*) \text{ and } n < \omega\}$  and  $X_{\alpha}^1 = X_{\alpha}^0 \cup \{\bar{\eta}^{\beta} : \beta \in \alpha \setminus \mathcal{U}\}$ . So for each  $\alpha \in \mathcal{U}$  there is  $\bar{n}_{\alpha} = \langle n_{\alpha,\ell} : \ell \in v_{\alpha} \rangle$  such that:  $\ell_{\alpha} \in v_{\alpha} \subseteq \{0, \dots, k(*)\}, n_{\alpha,\ell} < \omega$  and  $X_{\alpha+1}^1 \setminus X_{\alpha}^1 = \{x_{\bar{\eta} \upharpoonright \langle \ell, n \rangle} : \ell \in v_{\alpha} \text{ and } n \in [n_{\alpha,\ell}, \omega)\}$ .

For  $\alpha \leq \alpha(*)$  let  $G_{\Lambda,\alpha} = \langle \{y_{\bar{\eta}^{\beta},n}, x_{\bar{\nu}} : \beta \in \mathcal{U} \cap \alpha \text{ and } \bar{\nu} \in X_{\beta}^1\} \rangle_{G_{\Lambda}}$ . Clearly  $\langle G_{\Lambda,\alpha} : \alpha \leq \alpha(*) \rangle$  is purely increasing continuous with union  $G_{\Lambda}$ , and  $G_{\Lambda,0} = \{0\}$ . So it suffices to prove that  $G_{\Lambda,\alpha+1}/G_{\Lambda,\alpha}$  is free. If  $\alpha \notin \mathcal{U}$  the quotient is trivially a free group, and if  $\alpha \in \mathcal{U}$  we can use  $\ell_{\alpha} \in v_{\alpha}$  to prove that it is free giving a basis. □<sub>3.4</sub>

**3.5 Conclusion.** For every  $k(*) < \omega$  there is an  $\aleph_{k(*)+1}$ -free abelian group  $G$  of cardinality  $\lambda = \beth_{k(*)+1}$  such that  $\text{Hom}(G, \mathbb{Z}) = \{0\}$ .

*Proof.* We use  $\mathbf{x}$  and  $\langle h_{\bar{\eta}} : \bar{\eta} \in \Lambda^{\mathbf{x}} \rangle$  from 2.1(3), and we shall choose  $G \in \mathcal{G}_{\mathbf{x}}$ . So  $G$  is  $\aleph_{k(*)+1}$ -free by 3.4.

Let  $\mathcal{S} = \{\langle (a_i, \bar{\eta}_i) : i < i_1 \rangle \wedge \langle (b_j, \bar{\nu}_j, n_j) : j < j_1 \rangle : i_1 < \omega, a_i \in \mathbb{Z}, \bar{\eta}_i \in \Lambda_{\leq k(*)}^{\mathbf{x}}$  and  $j_1 < \omega, b_j \in \mathbb{Z}, \nu_j \in \Lambda^{\mathbf{x}}, n_j < \omega\}$  (actually  $\mathcal{S} = \Lambda_{\leq k(*)}^{\mathbf{x}}$  suffice noting  $\bar{\nu}_j = \langle \nu_{j,\ell} : \ell \leq k(*) \rangle$ ).

So  $|\mathcal{S}| = \lambda_{k(*)}$  and let  $\bar{p}$  be such that:

- (a)  $\bar{p} = \langle p^\alpha : \alpha < \lambda \rangle$
- (b)  $\bar{p}$  lists  $\mathcal{S}$
- (c)  $p^\alpha = \langle (a_i^\alpha, \bar{\eta}_i^\alpha) : i < i_\alpha \rangle \wedge \langle (b_j^\alpha, \bar{\nu}_j^\alpha, n_j^\alpha) : j < j_\alpha \rangle$  so  $\bar{\nu}_j^\alpha = \langle \nu_{j,\ell}^\alpha : \ell \leq k(*) \rangle$
- (d)  $\sup \text{Rang}(\eta_{i,k(*)}^\alpha) < \alpha$ ,  $\sup \text{Rang}(\nu_{j,k(*)}^\alpha) < \alpha$  if  $i < i_\alpha, j < j_\alpha$ .

Now to apply Definition 3.3 we have to choose  $z_\alpha$  (for Definition 3.3) as  $\Sigma\{a_i^\alpha x_{\bar{\eta}_i} : i < i_\alpha\} + \Sigma\{b_j^\alpha y_{\bar{\nu}_j^\alpha, n_j^\alpha} : j < j_\alpha\}$  and  $z_{\bar{\eta}} = z_{\bar{\eta},n} = z_{\eta_{k(*)}(0)}$  for  $\bar{\eta} \in \Lambda^{\mathbf{x}}, n < \omega$  then for  $\bar{\eta} \in \Lambda^{\mathbf{x}}$  we choose  $\langle b_{\bar{\eta},n} : n < \omega \rangle \in {}^\omega \mathbb{Z}$  such that:

- ⊗ there is no function  $h$  from  $\{z_{\bar{\eta}}\} \cup \{y_{\bar{\eta},n} : n < \omega\} \cup \{x_{\bar{\eta} \upharpoonright \langle m,n \rangle} : m \leq k(*), n < \omega\}$  into  $\mathbb{Z}$  satisfying
  - ⊗ (a)  $h(z_{\bar{\eta}}) \neq 0$  and
  - (b)  $h(x_{\bar{\eta} \upharpoonright \langle m,n \rangle}) = h_{\bar{\eta}}(\bar{\eta} \upharpoonright \langle m,n \rangle)$  for  $m \leq k(*), n < \omega$
  - (c) for every  $n$  sn
    - (\*) $_n$   $n!h(y_{\bar{\eta},n+1}) = h(y_{\bar{\eta},n}) + b_{\bar{\eta},n}h(z_{\bar{\eta}}) + \Sigma\{\{x_{\bar{\eta} \upharpoonright \langle m,n \rangle}\} : m \leq k(*)\}$ .

E.g. for each  $\rho \in {}^\omega 2$  we can try  $b_n^\rho = \rho(n)$  and assume toward contradiction that for each  $\rho \in {}^\omega 2$  there is  $h_\rho$  as above. Hence for some  $c \in \mathbb{Z} \setminus \{0\}$  the set  $\{\rho \in {}^\omega 2 : h_\rho(z_{\bar{\eta}}) = c\}$  is uncountable. So we can find  $\rho_1 \neq \rho_2$  such that  $h_{\rho_1} = c = h_{\rho_2}(x_\nu)$  and  $\rho_1 \upharpoonright (|c| + 7) = \rho_2 \upharpoonright (|c| + 7)$ . So for some  $n \geq |c| + 7, \rho_1 \upharpoonright n = \rho_2 \upharpoonright n$  and  $\rho_1(n) \neq \rho_2(n)$ . Now consider the equation (\*) $_n$  for  $h_{\rho_1}$  and  $h_{\rho_2}$ , subtract them and get  $(\rho_1(n) - \rho_2(n))c$  is divisible by  $n!$ , clear contradiction.

So  $G \in \mathcal{G}_{\mathbf{x}}$  is well defined and is  $\aleph_{k(*)+1}$ -free by 3.4. Suppose  $h \in \text{Hom}(G, \mathbb{Z})$  is non-zero, so for some  $\alpha < \lambda_{k(*)}$ ,  $h(z_\alpha) \neq 0$  (actually as  $G^1 = \langle \{x_{\bar{\nu}} : \bar{\nu} \in \Lambda_{\leq k(*)}^{\mathbf{x}}\} \rangle_G$  is a subgroup such that  $G/G^1$  is divisible necessarily  $h \upharpoonright G^1$  is not zero hence in 2.1(2) for some  $\bar{\nu} \in \Lambda_{\leq k(*)}^{\mathbf{x}}$  we have  $h(x_{\bar{\nu}}) \neq 0$ ). So by the choice of  $\langle h_{\bar{\eta}} : \bar{\eta} \in \Lambda \rangle$  for some  $\bar{\eta} \in \Lambda^{\mathbf{x}}, \eta_{k(*)}(\bar{0}) = \alpha$  and we have  $h_{\bar{\eta}} = h \upharpoonright \{x_{\bar{\eta} \upharpoonright \langle m,n \rangle} : m \leq k(*), n < \omega\}$ . By ⊗ we clearly get a contradiction.  $\square_{3.5}$

*Remark.* We can give more details as in the proof of 2.3.

*3.6 Conclusion.* For every  $n \leq m < \omega$  there is a purely increasing sequence  $\langle G_\alpha : \alpha \leq \omega_n + 1 \rangle$  of abelian groups,  $G_\alpha, G_\beta/G_\alpha$  are free for  $\alpha < \beta \leq \omega_n$  and  $G_{\omega_n+1}/G_{\omega_n}$  is  $\aleph_n$ -free and for some  $h \in \text{Hom}(G_\kappa, \mathbb{Z})$  has no extension in  $\text{Hom}(G_{\omega_n+1}, \mathbb{Z})$ .

*Proof.* Let  $G, z$  be as in 2.2. So also  $G/\mathbb{Z}z$  is  $\aleph_n$ -free. Let  $G_\alpha = \langle \{z\} \rangle_G$  for  $\alpha \leq \omega_2, G_{\omega_n+1} = G$ .

## §4 APPENDIX 1

4.1 Notation. If  $\bar{\eta}^* \in \Lambda_m^{\mathbf{x}}$  and  $\bar{\eta} = \bar{\eta}^* \upharpoonright \{\ell \leq k(*) : \ell \neq m\}$  and  $\nu = \eta_m^*$  then let  $x_{m,\bar{\eta},\nu} := x_{\bar{\eta}^*}$ . (See proof of 1.12).

*Proof of 1.8.* Let  $U \subseteq {}^\omega S$  be countable (and infinite) and define  $G'_U$  like  $G$  restricting ourselves to  $\eta_\ell \in U$ ; by the Löwenheim-Skolem argument it suffices to prove that  $G'_U$  is a free abelian group. List  $\Lambda \cap {}^{k(*)+1}U$  without repetitions as  $\langle \bar{\eta}_t : t < t^* \leq \omega \rangle$ , and choose  $s_t < \omega$  by induction on  $t < \omega$  such that  $[r < t \ \& \ \bar{\eta}_r \upharpoonright k(*) = \bar{\eta}_t \upharpoonright k(*) \Rightarrow \emptyset = \{\eta_{t,k(*)} \upharpoonright \ell : \ell \in [s_t, \omega)\} \cap \{\eta_{r,k(*)} \upharpoonright \ell : \ell \in [s_r, \omega)\}]$ .

Let

$$Y_1 = \{x_{m,\bar{\eta},\nu} : m < k(*), \bar{\eta} \in {}^{k(*)+1} \setminus \{m\}U \text{ and } \nu \in {}^\omega > 2\}$$

$$Y_2 = \left\{ x_{m,\bar{\eta},\nu} : m = k(*), \bar{\eta} \in {}^{k(*)}U \text{ and for no } t < t^* \text{ do we have } \right. \\ \left. \bar{\eta} = \bar{\eta}_t \upharpoonright k(*) \ \& \ \nu \in \{\eta_{t,k(*)} \upharpoonright \ell : s_t \leq \ell < \omega\} \right\}$$

$$Y_3 = \{y_{\bar{\eta}_t, n} : t < t^* \text{ and } n \in [s_t, \omega)\}.$$

Now

$$(*)_1 \ Y_1 \cup Y_2 \cup Y_3 \cup \{z\} \text{ generates } G'_U.$$

[Why? Let  $G'$  be the subgroup of  $G'_U$  which  $Y_1 \cup Y_2 \cup Y_3$  generates. First we prove by induction on  $n < \omega$  that for  $\bar{\eta} \in {}^{k(*)}U$  and  $\nu \in {}^n S$  we have  $x_{k(*),\bar{\eta},\nu} \in G'$ . If  $x_{k(*),\bar{\eta},\nu} \in Y_2$  this is clear; otherwise, by the definition of  $Y_2$  for some  $\ell < \omega$  (in fact  $\ell = n$ ) and  $t < \omega$  such that  $\ell \geq s_t$  we have  $\bar{\eta} = \bar{\eta}_t \upharpoonright k(*), \nu = \eta_{t,k(*)} \upharpoonright \ell$ .

Now

$$(a) \ y_{\bar{\eta}_t, \ell+1}, y_{\bar{\eta}_t, \ell} \text{ are in } Y_3 \subseteq G'.$$

Hence by the equation  $\boxtimes_{\bar{\eta}, n}$  in Definition 1.6, clearly  $x_{k(*),\bar{\eta},\nu} \in G'$ . So as  $Y_1 \subseteq G' \subseteq G'_U$ , all the generators of the form  $x_{k(*),\bar{\eta},\nu}$  with each  $\eta_\ell \in U$  are in  $G'$ .

Next note that

$$(b) \ x_{m,\bar{\eta}_t \upharpoonright \{i \leq k(*) : i \neq m\}, \nu} \text{ belong to } Y_1 \subseteq G' \text{ if } m < k(*).$$

Now for each  $t < \omega$  we prove that all the generators  $y_{\bar{\eta}_t, n}$  are in  $G'$ . If  $n \geq s_t$  then clearly  $y_{\bar{\eta}_t, n} \in Y_3 \subseteq G'$ . So it suffices to prove this for  $n \leq s_t$  by downward induction on  $n$ ; for  $n = s_t$  by an earlier sentence, for  $n < s_t$  by  $\boxtimes_{\bar{\eta}, n}$ . Together all the generators are in this subgroup so we are done.]

(\*)<sub>2</sub>  $Y_1 \cup Y_2 \cup Y_3 \cup \{z\}$  generates  $G'_U$  freely.

[Why? Translate the equations, see more in [Sh 771, §5].]

□<sub>1.8</sub>

*Proof of 1.10.* 0), 1) Obvious.

2),3),4) Follows.

5) Let  $\langle \eta_\ell : \ell < m(*) \rangle$  list  $u, U_\ell = U \cup (u \setminus \{\eta_\ell\})$  so  $G_{U, u} = G_{U_0^+} \dots + G_{U_{m(*)-1}}$ . First,  $G_{U, u} \subseteq G_{U \cup u}$  follows by the definitions. Second, we deal with proving  $G_{U, u} \subseteq_{\text{pr}} G_{U \cup u}$ . So assume  $z^* \in G, a^* \in \mathbb{Z}$  and  $a^* z^*$  belongs to  $G_{U_0} + \dots + G_{U_{m(*)}}$  so it has the form  $\Sigma\{b_i x_{\bar{\eta}^i} : i < i(*)\} + \Sigma\{c_j y_{\bar{\eta}_j, n_j} : j < j(*)\} + az$  with  $i(*) < \omega, j(*) < \omega$  and  $a^*, b_i, c_j \in \mathbb{Z}$  and  $\nu_i, \bar{\eta}^i, \bar{\eta}_j$  are suitable sequences of members of  $U_{\ell(i)}, U_{\ell(i)}, U_{k(j)}$  respectively where  $\ell(i), k(j) < m(*)$ . We continue as in [Sh 771].

6) Easy.

7) Clearly  $U_1 \cup v = U_2 \cup u$  hence  $G_{U_1 \cup u} \subseteq G_{U_1 \cup v} = G_{U_2 \cup u}$  hence  $G_{U, u} + G_{U_1 \cup u}$  is a subgroup of  $G_{U, u} + G_{U_2 \cup u}$ , so the first quotient makes sense.

Hence  $(G_{U, u} + G_{U_2 \cup u}) / (G_{U, u} + G_{U_1 \cup u})$  is isomorphic to  $G_{U_2 \cup u} / (G_{U_2 \cup u} \cap (G_{U, u} + G_{U_1 \cup u}))$ . Now  $G_{U_1, v} \subseteq G_{U_1 \cup v} = G_{U_2 \cup v} \subseteq G_{U, u} + G_{U_2, u}$  and  $G_{U_1, v} \subseteq G_{U, v} = G_{U, v} \setminus U = G_{U, u} \subseteq G_{U, u} + G_{U_2, u}$ . Together  $G_{U_1, v}$  is included in their intersection, i.e.  $G_{U_2 \cup u} \cap (G_{U, u} + G_{U_1 \cup u})$  include  $G_{U_1, v}$  and using part (1) both has the same divisible hull inside  $G^+$ . But as  $G_{U_1, v}$  is a pure subgroup of  $G$  by part (5) hence of  $G_{U_1 \cup v}$ . So necessarily  $G_{U_1 \cup u} \cap (G_{U, u} + G_{U_1, u}) = G_{U_1, v}$ , so as  $G_{U_2 \cup u} = G_{U_1 \cup v}$  we are done.

8) See [Sh 771, §5].

□<sub>1.10</sub>

*Proof of 1.12.* 1) We prove this by induction on  $|U|$ ; without loss of generality  $|u| = k$  as also  $k' = |u|$  satisfies the requirements.

Case 1:  $U$  is countable.

So let  $\{v_\ell^* : \ell < k\}$  list  $u$  be with no repetitions, now if  $k = 0$ , i.e.  $u = \emptyset$  then  $G_{U \cup u} = G_U = G_{U, u}$  so the conclusion is trivial. Hence we assume  $u \neq \emptyset$ , and let  $u_\ell := u \setminus \{v_\ell^*\}$  for  $\ell < k$ .

Let  $\langle \bar{\eta}_t : t < t^* \leq \omega \rangle$  list with no repetitions the set  $\Lambda_{U, u} := \{\bar{\eta} \in \Lambda^{\times} \cap {}^{k(*)+1}(U \cup u) : \text{for no } \ell < k \text{ does } \bar{\eta} \in {}^{k(*)+1}(U \cup u_\ell)\}$ . Now comes a crucial point: let  $t < t^*$ , for each  $\ell < k$  for some  $r_{t, \ell} \leq k(*)$  we have  $\eta_{t, r_{t, \ell}} = v_\ell^*$  by the definition of  $\Lambda_{U, u}$ , so

$|\{r_{t,\ell} : \ell < k\}| = k < k(*) + 1$  hence for some  $m_t \leq k(*)$  we have  $\ell < k \Rightarrow r_{t,\ell} \neq m_t$  so for each  $\ell < k$  the sequence  $\bar{\eta}_t \upharpoonright (k(*) + 1 \setminus \{m_t\})$  is not from  $\langle \rho_s : s \leq k(*) \text{ and } s \neq m_t \rangle : \rho_s \in \omega(U \cup u_\ell)$  for every  $s \leq k(*)$  such that  $s \neq m_t$ .

For each  $t < t^*$  we define  $J(t) = \{m \leq k(*) : \text{the set } \{\eta_{t,s} : s \leq k(*) \text{ \& } s \neq m\}$  is included in  $U \cup u_\ell$  for no  $\ell \leq k\}$ . So  $m_t \in J(t) \subseteq \{0, \dots, k(*)\}$  and  $m \in J(t) \Rightarrow \bar{\eta}_t \upharpoonright \{j \leq k(*) : j \neq m\} \notin {}^{k(*)+1} \setminus \{m\}(U \cup u_\ell)$  for every  $\ell \leq k$ . For  $m \leq k(*)$  let  $\bar{\eta}'_{t,m} := \bar{\eta}_t \upharpoonright \{j \leq k(*) : j \neq m\}$  and  $\bar{\eta}'_t := \bar{\eta}'_{t,m_t}$ . Now we can choose  $s_t < \omega$  by induction on  $t < t^*$  such that

- (\*) if  $t_1 < t, m \leq k(*)$  and  $\bar{\eta}'_{t_1,m} = \bar{\eta}'_{t,m}$ , then  
 $\eta_{t,m} \upharpoonright s_t \notin \{\eta_{t_1,m} \upharpoonright \ell : \ell < \omega\}$ .

Let  $Y^* = \{x_{m,\bar{\eta},\nu} \in G_{U \cup u} : x_{m,\bar{\eta},\nu} \notin G_{U \cup u_\ell} \text{ for } \ell < k\} \cup \{y_{\bar{\eta},n} \in G_{U \cup u} : y_{\bar{\eta},n} \notin G_{U \cup u_\ell} \text{ for } \ell < k\}$ .

Let

$$Y_1 = \{x_{m,\bar{\eta},\nu} \in Y^* : \text{for no } t < t^* \text{ do we have } m = m_t \text{ \& } \bar{\eta} = \bar{\eta}'_t\}.$$

$$Y_2 = \{x_{m,\bar{\eta},\nu} \in Y^* : x_{m,\bar{\eta},\nu} \notin Y_1 \text{ but for no } t < t^* \text{ do we have } m = m_t \text{ \& } \bar{\eta} = \bar{\eta}'_t \text{ \& } \eta_{t,m_t} \upharpoonright s_t \trianglelefteq \nu \triangleleft \eta_{t,m_t}\}$$

$$Y_3 = \{y_{\bar{\eta},n} : y_{\bar{\eta},n} \in Y^* \text{ and } n \in [s_t, \omega) \text{ for the } t < t^* \text{ such that } \bar{\eta} = \bar{\eta}'_t\}.$$

Now the desired conclusion follows from

- (\*)<sub>1</sub>  $\{y + G_{U,u} : y \in Y_1 \cup Y_2 \cup Y_3\}$  generates  $G_{U \cup u} / G_{U,u}$   
 (\*)<sub>2</sub>  $\{y + G_{U,u} : y \in Y_1 \cup Y_2 \cup Y_3\}$  generates  $G_{U \cup u} / G_{U,u}$  freely.

*Proof of (\*)<sub>1</sub>.* It suffices to check that all the generators of  $G_{U \cup u}$  belong to  $G'_{U \cup u} =: \langle Y_1 \cup Y_2 \cup Y_3 \cup G_{U,u} \rangle_G$ .

First consider  $x = x_{m,\bar{\eta},\nu}$  where  $\eta \in {}^{k(*)+1}(U \cup u), m \leq k(*)$  and  $\nu \in {}^n S$  for some  $n < \omega$ . If  $x \notin Y^*$  then  $x \in G_{U,u_\ell}$  for some  $\ell < k$  but  $G_{U \cup u_\ell} \subseteq G_{U,u} \subseteq G'_{U \cup u}$  so we are done, hence assume  $x \in Y^*$ . If  $x \in Y_1 \cup Y_2 \cup Y_3$  we are done so assume  $x \notin Y_1 \cup Y_2 \cup Y_3$ . As  $x \notin Y_1$  for some  $t < t^*$  we have  $m = m_t \text{ \& } \bar{\eta} = \bar{\eta}'_t$ . As  $x \notin Y_2$ , clearly for some  $t$  as above we have  $\eta_{t,m_t} \upharpoonright s_t \trianglelefteq \nu \triangleleft \eta_{t,m_t}$ . Hence by Definition 1.6 the equation  $\boxtimes_{\bar{\eta}_t,n}$  from Definition 1.6 holds, now  $y_{\bar{\eta}_t,n}, y_{\bar{\eta}_t,n+1} \in Y_3 \subseteq G'_{U \cup u}$ . So in order to deduce from the equation that  $x = x_{\bar{\eta}'_t \upharpoonright \langle m_t, n \rangle}$  belongs to  $G'_{U \cup u}$ , it suffices to show that  $x_{\bar{\eta}'_t \upharpoonright \langle j, n \rangle} \in G'_{U \cup u}$  for each  $j \leq k(*)$ ,  $j \neq m_t$ . But each such  $x_{\bar{\eta}'_t \upharpoonright \langle j, n \rangle}$  belong to  $G'_{U \cup u}$  as it belongs to  $Y_1 \cup Y_2$ .



[Why? Otherwise necessarily for some  $r < t^*$  we have  $j = m_r, \bar{\eta}'_{t,j} = \bar{\eta}'_{r,m_r}$  and  $\eta_{r,m_r} \upharpoonright s_r \trianglelefteq \eta_t \upharpoonright n \triangleleft \eta_{r,m_r}$  so  $n \geq s_r$  and as said above  $n \geq s_t$ . Clearly  $r \neq t$  as  $m_r = j \neq m_t$ , now as  $\bar{\eta}'_{t,m_r} = \bar{\eta}'_{r,m_r}$  and  $\bar{\eta}_t \neq \bar{\eta}_r$  (as  $t \neq r$ ) clearly  $\eta_{t,m_r} \neq \eta_{r,m_r}$ . Also  $\neg(r < t)$  by (\*) above applied with  $r, t$  here standing for  $t_1, t$  there as  $\eta_{r,m_r} \upharpoonright s_r \trianglelefteq \eta_{t,j} \upharpoonright n \triangleleft \eta_{r,m_r}$ . Lastly for if  $t < r$ , again (\*) applied with  $t, r$  here standing for  $t_1, t$  there as  $n \geq m_t$  gives contradiction.]

So indeed  $x \in G'_{U \cup u}$ .

Second consider  $y = y_{\bar{\eta},n} \in G_{U \cup u}$ , if  $y \notin Y^*$  then  $y \in G_{U,u} \subseteq G'_{U \cup u}$ , so assume  $y \in Y^*$ . If  $y \in Y_3$  we are done, so assume  $y \notin Y_3$ , so for some  $t, \bar{\eta} = \bar{\eta}_t$  and  $n < s_t$ . We prove by downward induction on  $s \leq s_t$  that  $y_{\bar{\eta},s} \in G'_{U \cup u}$ , this clearly suffices. For  $s = s_t$  we have  $y_{\bar{\eta},s} \in Y_3 \subseteq G'_{U \cup u}$ ; and if  $y_{\bar{\eta},s+1} \in G'_{U \cup u}$  use the equation  $\boxtimes_{\bar{\eta}_t,s}$  from 1.6, in the equation  $y_{\bar{\eta},s+1} \in G'_{U \cup u}$  and the  $x$ 's appearing in the equation belong to  $G'_{U \cup u}$  by the earlier part of the proof (of  $(*)_1$ ) so necessarily  $y_{\bar{\eta},s} \in G'_{U \cup u}$ , so we are done.

*Proof of  $(*)_2$ .* We rewrite the equations in the new variables recalling that  $G_{U \cup u}$  is generated by the relevant variables freely except the equations of  $\boxtimes_{\bar{\eta},n}$  from Definition 1.6. After rewriting, all the equations disappear.

Case 2:  $U$  is uncountable.

As  $\aleph_1 \leq |U| \leq \aleph_{k(*)-k}$ , necessarily  $k < k(*)$ .

Let  $U = \{\rho_\alpha : \alpha < \mu\}$  where  $\mu = |U|$ , list  $U$  with no repetitions. Now for each  $\alpha \leq |U|$  let  $U_\alpha := \{\rho_\beta : \beta < \alpha\}$  and if  $\alpha < |U|$  then  $u_\alpha = u \cup \{\rho_\alpha\}$ . Now

- ⊙<sub>1</sub>  $\langle\langle G_{U,u} + G_{U_\alpha \cup u} \rangle\rangle / G_{U,u} : \alpha < |U|$  is an increasing continuous sequence of subgroups of  $G_{U \cup u} / G_{U,u}$ .  
[Why? By 1.10(6).]
- ⊙<sub>2</sub>  $G_{U,u} + G_{U_0 \cup u} / G_{U,u}$  is free.  
[Why? This is  $(G_{U,u} + G_{\emptyset \cup u}) / G_{U,u} = (G_{U,u} + G_u) / G_{U,u}$  which by 1.10(8) is isomorphic to  $G_u / G_{\emptyset,u}$  which is free by Case 1.]

Hence it suffices to prove that for each  $\alpha < |U|$  the group  $(G_{U,u} + G_{U_{\alpha+1} \cup u}) / (G_{U,u} + G_{U_\alpha \cup u})$  is free. But easily

- ⊙<sub>3</sub> this group is isomorphic to  $G_{U_\alpha \cup u_\alpha} / G_{U_\alpha, u_\alpha}$ .  
[Why? By 1.10(7) with  $U_\alpha, U_{\alpha+1}, U, \rho_\alpha, u$  here standing for  $U_1, U_2, U, \eta, u$  there.]
- ⊙<sub>4</sub>  $G_{U_\alpha \cup u_\alpha} / G_{U_\alpha, u_\alpha}$  is free.  
[Why? By the induction hypothesis, as  $\aleph_0 + |U_\alpha| < |U| \leq \aleph_{k(*)-(k+1)}$  and  $|u_\alpha| = k + 1 \leq k(*)$ .]

2) If  $k(*) = 0$  just use 1.8, so assume  $k(*) \geq 1$ . Now the proof is similar to (but easier than) the proof of case (2) inside the proof of part (1) above.

□<sub>1.12</sub>

## REFERENCES.

- [EM02] Paul C. Eklof and Alan Mekler. *Almost free modules: Set theoretic methods*, volume 65 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam, 2002. Revised Edition.
- [GbT106] Rüdiger Göbel and Jan Trlifaj. *Approximations and endomorphism algebras of modules*, volume 41 of *de Gruyter Expositions in Mathematics*. Walter de Gruyter, Berlin, 2006.
- [GShS 785] Rüdiger Göbel, Saharon Shelah, and Lutz Strüngmann. Almost-Free  $E$ -Rings of Cardinality  $\aleph_1$ . *Canadian Journal of Mathematics*, **55**:750–765, 2003. arxiv:math.LO/0112214.
- [Sh:e] Saharon Shelah. *Non-structure theory*, accepted. Oxford University Press.
- [Sh 52] Saharon Shelah. A compactness theorem for singular cardinals, free algebras, Whitehead problem and transversals. *Israel Journal of Mathematics*, **21**:319–349, 1975.
- [Sh:98] Saharon Shelah. Whitehead groups may not be free, even assuming CH. II. *Israel Journal of Mathematics*, **35**:257–285, 1980.
- [Sh 161] Saharon Shelah. Incompactness in regular cardinals. *Notre Dame Journal of Formal Logic*, **26**:195–228, 1985.
- [Sh 300] Saharon Shelah. Universal classes. In *Classification theory (Chicago, IL, 1985)*, volume 1292 of *Lecture Notes in Mathematics*, pages 264–418. Springer, Berlin, 1987. Proceedings of the USA–Israel Conference on Classification Theory, Chicago, December 1985; ed. Baldwin, J.T.
- [Sh 523] Saharon Shelah. Existence of Almost Free Abelian groups and reflection of stationary set. *Mathematica Japonica*, **45**:1–14, 1997. arxiv:math.LO/9606229.
- [Sh 771] Saharon Shelah. Polish Algebras, shy from freedom. *Israel Journal of Mathematics*, **181**:477–507, 2011. arxiv:math.LO/0212250.
- [Sh 898] Saharon Shelah. PCF and abelian groups. *Forum Mathematicum*, **25**:967–1038, 2013. arxiv:0710.0157.