

NO LIMIT MODEL IN INACCESSIBLE SH906

SAHARON SHELAH

Dedicated to Michael Makkai

ABSTRACT. Our aim is to improve the negative results i.e., non-existence of limit models, and the failure of the generic pair property from [5] to inaccessible λ as promised there. In [5], the negative results were obtained only for non-strong limit cardinals.

§ 0. INTRODUCTION

Let $\lambda = \lambda^{<\lambda} > \kappa$ be regular cardinals. A complete first order theory T may have (some variant of) (λ, κ) -limit model, which, if exists, is unique, see history in [5] and Definition 0.9. There we prove existence for the theory of linear order and non-existence for first order theories which are strongly independent and then just independent and even the parallel for $\kappa = 2$ (one direction of the so-called generic pair conjecture). Those non-existence results in [5] were for $\lambda = 2^\kappa$, here we deal with strongly inaccessible λ . In [8] there are existence results but for λ measurable, and we promise there the non-existence results for λ strongly inaccessible as complimentary results.

Let λ be strongly inaccessible ($> |T|$) such that $\lambda^+ = 2^\lambda$; this for transparency only.

Here in §1 we prove that for strongly independent T (see Definition 0.2), a strong version of the generic pair conjecture (see Definition 0.7(2)) holds. We also prove the non-existence of (λ, κ) -limit models, a related property (for all versions of “limit model”).

In §2, we prove this even for independent T . The use of $\lambda^+ = 2^\lambda$ is just to have a more transparent formulation of the conjecture. See more on the generic pair conjecture for dependent T in [4].

We thank Itay Kaplan for much helpful criticism.

- Notation* 0.1. 1) \mathcal{D}_λ is the club filter on λ for λ regular uncountable.
2) $S_\kappa^\lambda = \{\delta < \lambda : \text{cf}(\delta) = \kappa\}$.
3) For a limit ordinal δ let $\mathcal{P}^{\text{ub}}(\delta) = \{\mathcal{U} : \mathcal{U} \text{ is an unbounded subset of } \delta\}$.
4) T denotes a complete first order theory.

{0.4}

Date: October 29 , 2011.

1991 *Mathematics Subject Classification.* MSC 2010, Primary 03C45; Secondary: 03C55.

Key words and phrases. model theory, classification theory, dependent theorem, limit models, strongly inaccessible.

The author would like to thank the Israel Science Foundation for partial support of this research (Grant No.242/03). The author thanks Alice Leonhardt for the beautiful typing. First Typed - 06/Sept/12. Publ. 906.

- 5) For a model M , $\varphi(\bar{x}, \bar{y}) \in \mathbb{L}(\tau_M)$ and $\bar{d} \in {}^{\ell g(\bar{y})}M$, let $\varphi(M, \bar{d}) = \{\bar{c} \in {}^{\ell g(\bar{x})}M : M \models \varphi[\bar{c}, \bar{d}]\}$.
- 6) $\mathbf{S}^n(A, M) = \{\text{tp}(\bar{b}, A, N) : M \prec N \text{ and } \bar{b} \in {}^nN\}$ where $\text{tp}(\bar{c}, A, N) = \{\varphi(\bar{x}, \bar{a}) : \varphi(\bar{x}, \bar{y}) \in \mathbb{L}(\tau_M), \bar{a} \in {}^{\ell g(\bar{y})}A \text{ and } M \models \varphi[\bar{c}, \bar{a}]\}$.
- 7) $\mathbf{S}^n(M) = \mathbf{S}^n(M, M)$ and $\mathbf{S}^{<\omega}(M) = \cup\{\mathbf{S}^n(M) : n \in \omega\}$.
- 8) $\beth_\alpha(\lambda) = \lambda + \Sigma\{2^{\beth_\beta(\lambda)} : \beta < \alpha\}$ and $\beth_\alpha = \beth_\alpha(\aleph_0)$.

Recall (as in [5, 2.3])

{0.7}

Definition 0.2. 1) T has the strong independence property (or is strongly independent) when: some $\varphi(\bar{x}, \bar{y}) \in \mathbb{L}(\tau_T)$ has it, where:

2) $\varphi(\bar{x}, \bar{y}) \in \mathbb{L}(\tau_T)$ has the strong independence property for T when for every model M of T and pairwise disjoint finite $\mathbf{I}_1, \mathbf{I}_2 \subseteq {}^{\ell g(\bar{y})}(M)$ for some $\bar{a} \in {}^{\ell g(\bar{x})}M$ we have $\ell \in \{1, 2\} \wedge \bar{b} \in \mathbf{I}_\ell \Rightarrow M \models \varphi[\bar{a}, \bar{b}]^{\text{if}(\ell=2)}$.

Remark 0.3. 1) Elsewhere we use $\varphi(x, y)$, i.e. the x and y are singletons, but the proofs are not affected.

2) Also we may restrict ourselves to $\mathbf{I}_1, \mathbf{I}_2 \subseteq \psi(M, \bar{d})$ where $\psi \in \mathbb{L}(\tau_T)$ such that $\psi(M, \bar{d})$ is infinite, and we may restrict ourselves to $\mathbf{I}_1, \mathbf{I}_2$ such that every $\bar{b} \in \mathbf{I}_1 \cup \mathbf{I}_2$ realizes a fixed non-algebraic type $p \in \mathbf{S}^m(A, M)$ with M being $(|A|^+ + \aleph_0)$ -saturated. The results are not really affected.

{0q.37}

Question 0.4. 1) Assume $\lambda_2 = \lambda_2^{<\lambda_1} \geq \lambda_1 > |T|$, T a complete first order dependent theory. Is the theory $T_{\lambda_1, \lambda_2}^*$ a dependent theory or at least when is $T_{\lambda_1, \lambda_2}^*$ a dependent theory? where

(a) $T_{\lambda_1, \lambda_2}^* = \text{Th}(K_{\lambda_1, \lambda_2}^+)$ where

(b) $K_{\lambda_1, \lambda_2}^+ = \{(N, M) : M \text{ is a } \lambda_1\text{-saturated model of } T \text{ of cardinality } \lambda_2, N \text{ a } \lambda_2^+\text{-saturated elementary extension of } M\}$.

2) Similarly for other properties of $T_{\lambda_1, \lambda_2}^*$; note¹ that this theory is complete if $\lambda_1 = \lambda_2$.

3) When can we prove that $T_{\lambda_1, \lambda_2}^*$ does not depend on the cardinals at least for many pairs?

Remark 0.5. 1) Concerning failure of 0.4(1) see Kaplan-Shelah [3].

2) Any solution of the generic pair conjecture answers positively 0.4(3) for dependent T in the relevant cases.

3) It is known that in 0.4(1) if T extends PA or ZFC then in $T^* = \text{Th}(N, M)$ we can interpret the second order theory of λ_2 .

But may well be that as in Baldwin-Shelah [1]

{0q.44}

Question 0.6. Assume $|T| < \kappa \leq \lambda_1 \leq \lambda_2 = \lambda_2^{<\lambda_1}$, T a complete first order theory. For which T 's can we interpret in $M \in K_{\lambda_1, \lambda_2}^+$ a model of PA of cardinality $\geq \lambda_1$ by first order formula or just an $\mathbb{L}_{\infty, \kappa}(\tau_T)$ -formulas with parameters, the intention is that we assume λ_2 is enough larger than λ_1 which is large enough than $|T|$; if $2^\kappa \geq \lambda_1$ this is trivial.

Recall (from [5, 0.2])

¹let $(N_\ell, M_\ell) \in K_{\lambda_1, \lambda_2}^+$ for $\ell = 1, 2$ and let f_* be an isomorphism from M_1 onto M_2 and let $\mathcal{F} = \{f : f \text{ is a } (N_1, N_2)\text{-elementary mapping extending } f_* \text{ of cardinality } \leq \lambda_1\}$. Now we can prove that any $f \in \mathcal{F}$ preserve satisfaction for first order formulas.

{0.14}

Definition 0.7. 0) Let $\text{EC}_\lambda(T)$ be the class of models M of (the first order) T of cardinality λ . Let $\text{EC}_{\lambda,\kappa}(T)$ be the class of κ -saturated models $M \in \text{EC}_\lambda(T)$.

1) Assume $\lambda > |T|$, (we usually assume $\lambda = \lambda^{<\lambda}$) and $2^\lambda = \lambda^+$, $M_\alpha \in \text{EC}_\lambda(T)$ is \prec -increasing continuous for $\alpha < \lambda^+$ with $M = \cup\{M_\alpha : \alpha < \lambda^+\} \in \text{EC}_{\lambda^+}(T)$, and M is saturated. The generic pair property (for T, λ) says that for some club E of λ^+ for all pairs $\alpha < \beta$ of ordinals from E of cofinality λ , (M_β, M_α) has the same isomorphism type (we denote this property of T by $\text{Pr}_{\lambda,\lambda}^2(T)$).

2) The generic pair conjecture for $\lambda > \aleph_0$, (usually $\lambda = \lambda^{<\lambda}$) such that $2^\lambda = \lambda^+$ says that for any complete first order T of cardinality $< \lambda$, T is dependent iff it has the generic pair property for λ .

3) Let $\mathbf{n}_{\lambda,\kappa}(T)$ be $\min\{|\{M_\delta / \cong : \delta \in E \text{ has cofinality } \kappa\}| : E \text{ a club of } \lambda^+\}$ for λ and $\bar{M} = \langle M_\alpha : \alpha < \lambda^+ \rangle$ as above and $\kappa = \text{cf}(\kappa) \leq \lambda$; clearly the choice of \bar{M} is immaterial.

{0.22}

Remark 0.8. 1) Note that to say $\mathbf{n}_{\lambda,\kappa}(T) = 1$ is a way to say that T has (some variant of) a (λ, κ) -limit model, see 0.9 below. There are other variants of the Definition of limit.

2) Recall that we conjecture that for $\lambda = \lambda^{<\lambda} > \kappa = \text{cf}(\kappa) > |T|$, $2^\lambda = \lambda^+$ we have $\mathbf{n}_{\lambda,\kappa}(T) = 1 \Leftrightarrow \mathbf{n}_{\lambda,\kappa}(T) < 2^\lambda \Leftrightarrow T$ is dependent. The use of " $\lambda^+ = 2^\lambda$ " is just for clarity. See more in [5], [8], [4].

3) Recall that if $\lambda = \kappa = \lambda^{<\lambda}$, then $\mathbf{n}_{\lambda,\lambda}(T) = 1$ means T has a unique saturated model; (and parallelly if $\lambda > \text{cf}(\lambda) = \kappa$, λ strong limit). So we concentrate on the case $\mathbf{n}_{\lambda,\kappa}(T) = 1$ where $\kappa < \lambda$.

{0.26}

Definition 0.9. We define when M_* is a (λ, κ) -limit model of T where $\lambda \geq \kappa = \text{cf}(\kappa)$ and $\lambda \geq |T|$. In general it means that: letting $K_\lambda = \{M : M \text{ is a model of } T \text{ with universe an ordinal } \in [\lambda, \lambda^+)\}$, for some function \mathbf{F} with domain K and satisfying $M \prec \mathbf{F}(M) \in K$ we have:

$$\oplus \text{ if } M_\alpha \in K \text{ for } \alpha < \lambda^+ \text{ is } \prec\text{-increasing continuous and } \alpha < \lambda \Rightarrow \mathbf{F}(M_{\alpha+1}) \prec M_{\alpha+2} \text{ then for some club } E \text{ of } \lambda^+ \text{ we have:}$$

$$\delta \in E \wedge \text{cf}(\delta) = \kappa \Rightarrow M_\delta \cong M_*.$$

{0.29}

Remark 0.10. If $2^\lambda = \lambda^+$ we have: $\mathbf{n}_{\lambda,\kappa}(T) = 1$ iff T has a (λ, κ) -limit model.

§ 1. STRONGLY INDEPENDENT T

Context 1.1. 1) T is a fixed first order complete theory and $\mathfrak{C} = \mathfrak{C}_T$ a monster for it; for notational simplicity τ_T is relational.

2) We let λ be a regular uncountable cardinal $> |T|$; we deal mainly with strongly inaccessible λ .

Here for λ strongly inaccessible and (complete first order) T with the strong independence property (of cardinality $< \lambda$) we prove the non-existence of (λ, κ) -limit models for $\kappa = \text{cf}(\kappa) < \lambda$ (in Theorem 1.9) and the generic pair conjecture for λ and T , in Theorem 1.10 (which shows non-isomorphism). Recall that the generic pair property speaks on the isomorphism type of pairs of models.

Definition 1.2 gives us a more constructive invariant of $(N, M)/\cong$. Unfortunately it seemed opaque how to manipulate it so we shall use a related but different version, the one from Definition 1.4. Naturally it concentrates on types in one formula $\varphi(\bar{y}, \bar{x})$ witnessing the strong independence property. But mainly gives the pair (N, M) an invariant $\langle \mathcal{P}_\delta : \delta < \lambda \rangle / \mathcal{D}_\lambda$ where $\mathcal{P}_\delta \subseteq \mathcal{P}(\mathcal{P}(\delta))$. Now always $|\mathcal{P}_\delta| \leq 2^{|\delta|}$ and it is easily computable from one $\mathcal{P} \subseteq \mathcal{P}(\delta)$, in fact from the invariant $\text{inv}_4(M, N)$ from Definition 1.2, but in our proofs its use is more transparent. It has monotonicity property and we can increase it.

We need different but similar version for the proof of non-existence of (λ, κ) -limit models.

{inv.7}

Definition 1.2. 1) Let \mathcal{E}_T^* be the following two-place relation on $\{(M, \mathbf{P}) : M \models T \text{ and } \mathbf{P} \subseteq \mathbf{S}^{<\omega}(M)\}$; let $(M_1, \mathbf{P}_1) \mathcal{E}_T^*(M_2, \mathbf{P}_2)$ iff there is an isomorphism h from M_1 onto M_2 mapping \mathbf{P}_1 onto \mathbf{P}_2 .

2) For models $M \subseteq N$ we define (the important case is $M \prec N \models T$):

$$(a) \text{ inv}_1(M, N) = \{p \in \mathbf{S}^{<\omega}(M) : p \text{ is realized in } N\}$$

$$(b) \text{ inv}_2(M, N) = (M, \text{inv}_1(M, N)) / \mathcal{E}_T^*.$$

3) If $M \prec N$ are models of T such that the universe of N is $\subseteq \lambda$, recalling \mathcal{D}_λ is the club filter on λ , let:

(a) for any ordinal $\delta < \lambda$

$$\text{inv}_3(\delta, M, N) = (M \upharpoonright \delta, \{p \in \mathbf{S}^{<\omega}(M \upharpoonright \delta) : p \text{ is realized by some sequence from } N \upharpoonright \delta\}) / \mathcal{E}_T^*$$

$$(b) \text{ inv}_4(M, N) = \langle \text{inv}_3(\delta, M, N) : \delta < \lambda \rangle / \mathcal{D}_\lambda.$$

4) If $M \prec N$ are models of T of cardinality λ then $\text{inv}_4(M, N)$ is equal to $\text{inv}_4(f(M), f(N))$ for every one-to-one function f from N into λ (equivalently some f , see 1.3(1),(2) below)

{inv.8}

Observation 1.3. 0) In Definition 1.2(3) for a club of δ 's below λ we have $M \upharpoonright \delta \prec M$ and $N \upharpoonright \delta \prec N$ and so $M \upharpoonright \delta \prec N \upharpoonright \delta \models T$.

1) Concerning Definition 1.2(3), if $M \prec N$ are models of T of cardinality λ and f_1, f_2 are one-to-one functions from N into λ then $\text{inv}_4(f_1(M), f_1(N)) = \text{inv}_4(f_2(M), f_2(N))$ using the definition 1.2(3)(b).

2) Definitions 1.2(3), 1.2(4) are compatible and in 1.2(4), “some f such that f is a one-to-one function from N to λ ” is equivalent to “every f such that...”

Proof. Straight, e.g. (this argument will be used several times).

1) Let $E = \{\delta < \lambda : \delta \text{ is a limit ordinal such that } M \upharpoonright \delta \prec M, N \upharpoonright \delta \prec N \text{ and } \text{Rang}(f_\ell \upharpoonright \delta) = \text{Rang}(f_\ell) \cap \delta \text{ for } \ell = 1, 2\}$. So E is a club of λ and $\delta \in E \Rightarrow f_2 \circ f_1^{-1}$ is an isomorphism from $f_1(N \upharpoonright \delta)$ onto $f_2(N \upharpoonright \delta)$, mapping $f_1(M \upharpoonright \delta)$ onto $f_2(M \upharpoonright \delta)$. $\square_{1.3}$

{inv.10}

Definition 1.4. Assume $\varphi = \varphi(\bar{x}, \bar{y}) \in \mathbb{L}(\tau_T)$ and $N_1 \prec N_2$ are models of T of cardinality λ .

1) For one-to-one mapping f from N_2 to λ and $\delta < \lambda$ we define

$$\text{inv}_5^\varphi(\delta, f, N_1, N_2) = \{\mathcal{P} \subseteq \mathcal{P}(\delta) : \begin{array}{l} \text{there are } \bar{a}_\gamma \in {}^{\ell g(\bar{x})} N_2 \text{ satisfying } f(\bar{a}_\gamma) \in {}^{\ell g(\bar{x})} \delta \\ \text{for } \gamma < \delta \text{ such that for every} \\ \mathcal{U} \subseteq \delta \text{ the following are equivalent :} \\ (i) \quad \mathcal{U} \in \mathcal{P} \\ (ii) \quad \text{for some } \bar{b} \in {}^{\ell g(\bar{y})} N_1 \text{ we have} \\ \gamma < \delta \Rightarrow N_2 \models \varphi[\bar{a}_\gamma, \bar{b}]^{\text{if}(\gamma \in \mathcal{U})}\}. \end{array}$$

2) We let $\text{inv}_6^\varphi(N_1, N_2)$ be $\langle \text{inv}_5^\varphi(\delta, f, N_1, N_2) : \delta < \lambda \rangle / \mathcal{D}_\lambda$ for some (equivalently every) f as above.

{inv.11}

Claim 1.5. 1) In Definition 1.4(2) we have $\text{inv}_6^\varphi(N_1, N_2)$ is well defined.

2) In Definition 1.4, for $\delta, \lambda, N_1, N_2, \varphi(\bar{x}, \bar{y})$ as there

- (a) the set $\text{inv}_5^\varphi(\delta, f, N_1, N_2)$ has cardinality at most $2^{|\delta|}$
- (b) if π is a one-to-one function from $f(N_2)$ into λ mapping $f(N_2) \cap \delta$ onto $\pi(f(N_2)) \cap \delta$ then $\text{inv}_5^\varphi(\delta, \pi \circ f, N_1, N_2) = \text{inv}_5^\varphi(\delta, f, N_1, N_2)$.

Proof. Easy. $\square_{1.5}$

{inv.12}

Definition 1.6. 1) For $\varphi = \varphi(\bar{x}, \bar{y}) \in \mathbb{L}(\tau_T)$, a model N of T with universe λ , δ a limit ordinal $< \lambda$ and $\kappa < \lambda$ let

$$\text{inv}_{7,\kappa}^\varphi(\delta, N) = \{\mathcal{P} \subseteq \mathcal{P}(\delta) : \begin{array}{l} \text{we can find } \bar{a}_\gamma^i \in {}^{\ell g(\bar{x})} \delta \text{ for } \gamma < \delta, i < \kappa \text{ such that} \\ \text{the following conditions on } \mathcal{U} \subseteq \delta \text{ are equivalent :} \\ (i) \quad \mathcal{U} \in \mathcal{P} \\ (ii) \quad \text{for some } \bar{b} \in {}^{\ell g(\bar{y})} N \text{ we have :} \\ \text{for every } i < \kappa \text{ large enough for every} \\ \gamma < \delta \text{ we have } N \models \varphi[\bar{a}_\gamma^i, \bar{b}]^{\text{if}(\gamma \in \mathcal{U})}\}. \end{array}$$

2) For $\varphi = \varphi(\bar{y}, \bar{x}) \in \mathbb{L}(\tau_T)$ and a model N of T of cardinality λ let $\text{inv}_{8,\kappa}^\varphi(N) = \langle \text{inv}_7^\varphi(\delta, N') : \delta < \lambda \rangle / \mathcal{D}_\lambda$ for every, equivalently some model N' isomorphic to N with universe λ .

{inv.13}

Observation 1.7. 1) $\text{inv}_{8,\kappa}^\varphi(N)$ is well defined for $N \in \text{EC}_\lambda(T)$ when $|T| + \kappa < \lambda$.

2) In Definition 1.6(1) we have $|\text{inv}_{7,\kappa}^\varphi(\delta, N)| \leq 2^{|\delta| + \kappa}$.

Proof. Easy. $\square_{1.7}$

{inv.21}

Claim 1.8. Assume $\lambda > |T|$ is regular, $S \subseteq \lambda$ is stationary, $\varphi = \varphi(\bar{x}, \bar{y})$ and

- (a) $\langle N_i : i < \kappa \rangle$ is a \prec -increasing sequence
- (b) $N_i \in \text{EC}_\lambda(T)$

- (c) $N = \cup\{N_i : i < \kappa\}$
- (d) $\bar{\mathcal{P}} = \langle \mathcal{P}_\alpha : \alpha < \lambda \rangle$ where $\mathcal{P}_\alpha \subseteq \mathcal{P}(\alpha)$
- (e) f is a one-to-one function from N onto λ
- (f) for a club of δ 's below λ there are $\bar{a}_\gamma^j \in N_{j+1} \cap f^{-1}(\delta)$ for $\gamma < \delta$, $j < \kappa$ satisfying
 - (α) for every $\bar{c} \in {}^{\ell g(\bar{x})}(N_j)$ there is $\mathcal{U} \in \mathcal{P}_\delta$ such that $\gamma < \delta \Rightarrow N \models \varphi[\bar{a}_\gamma^j, \bar{c}]^{\text{if}(\gamma \in \mathcal{U})}$
 - (β) for every $\mathcal{U} \in \mathcal{P}_\delta$ for some $\bar{b} \in {}^{\ell g(\bar{y})}(N_\delta)$ we have $\gamma < \delta \Rightarrow N \models \varphi[\bar{a}_\gamma^j, \bar{b}]^{\text{if}(\gamma \in \mathcal{U})}$.

Then $\{\delta \in S : \mathcal{P}_\delta \in \text{inv}_{7,\kappa}^\varphi(\delta, f(N))\} \in \mathcal{D}_\lambda + S$.

Proof. Straight. □_{1.8}

{inv. 14} Now we come to the main two results of this section.

Theorem 1.9. For some club E of λ^+ , if $\delta_1 \neq \delta_2$ belong to $E \cap S_\kappa^{\lambda^+}$ then $M_{\delta_1}, M_{\delta_2}$ are not isomorphic, moreover $\text{inv}_{8,\kappa}^\varphi(M_{\delta_1}) \neq \text{inv}_{8,\kappa}^\varphi(M_{\delta_2})$ when:

- ⊠ (a) T has the strong independence property (see Definition 0.2)
- (b) $\lambda = \lambda^{<\lambda}$ is regular uncountable, $\lambda > |T|$, $\lambda > \kappa = \text{cf}(\kappa)$ and $\lambda^+ = 2^\lambda$
- (c) M is a saturated model of T of cardinality λ^+
- (d) $\langle M_\alpha : \alpha < \lambda^+ \rangle$ is \prec -increasing continuous sequence with union M , each of cardinality λ .

{inv. 24}

Theorem 1.10. Assume ⊠ of 1.9.

1) For some club E of λ^+ , if $\delta_1 < \delta_2 < \delta_3$ are from E and $\delta_\ell \in S_\kappa^{\lambda^+}$ for $\ell = 1, 2, 3$ then $(M_{\delta_2}, M_{\delta_1}) \not\cong (M_{\delta_3}, M_{\delta_1})$, moreover $\text{inv}_6^\varphi(M_{\delta_2}, M_{\delta_1}) \neq \text{inv}_6^\varphi(M_{\delta_3}, M_{\delta_1})$ for some φ .

2) If $M \prec N_0$ are models of T of cardinality λ , then for some elementary extension $N_1 \in \text{EC}_\lambda(T)$ of N_0 we have $N_1 \prec N_2 \in \text{EC}_\lambda(T) \Rightarrow (N_0, M) \not\cong (N_2, M)$.

Discussion 1.11. We shall below start with $M \in \text{EC}_\lambda(T)$ and a sequence $\langle b_i : i < \lambda \rangle$ of distinct members such that $\langle \varphi(\bar{b}_i, \bar{y}) : i < \lambda \rangle$ are independent, and like to find $N, \langle \bar{a}_i : i < \lambda \rangle$ such that $M \prec N \in \text{EC}_\lambda(T)$ and the $\langle b_i : i < \lambda \rangle$ has a real affect on the relevant φ -invariant, in the case of 1.10(1) this is $\text{inv}_6^\varphi(M, N)$: for a stationary set of δ 's below λ it adds something to the δ -th component in a specific representation, i.e. assuming $f : N \rightarrow \lambda$ is a one-to-one function and we deal with $\langle \text{inv}_5^\varphi(\delta, f, M, N) : \delta < \lambda \rangle$; we have freedom about $\varphi(\bar{a}_\alpha, \bar{b}_i)$ and we can assume $\bar{b} \in {}^{\ell g(\bar{y})}M \setminus \{b_i : i < \lambda\} \Rightarrow N \models \neg \varphi[\bar{a}_\alpha, \bar{b}]$.

But the relevant \mathcal{P}_δ is influenced not just by say $\langle \bar{b}_i : i \in [\delta, 2^{|\delta|}] \rangle$ but also by later \bar{b}_i 's (and earlier \bar{b}_i). To control this we use below $\langle \bar{a}_\alpha : \alpha < \lambda \rangle, S, E$ such that we deal with different $\delta \in S$ in an independent way to large extent; this is the reason for choosing the C_α^* 's.

Proof. Proof of 1.9 By the proof of [5, §2] without loss of generality λ is strongly inaccessible. Choose $\theta \in \text{Reg} \cap \lambda \setminus \{\aleph_0\}$, will be needed when we generalize the proof in §2.

Let $\langle \mathcal{U}_i : i < \kappa \rangle$ be a \subseteq -increasing sequence of subsets of λ such that the set $\mathcal{U}_i^- = \mathcal{U}_i \setminus \{\mathcal{U}_j : j < i\}$ has cardinality λ for each $i < \kappa$ and let $\mathcal{U}_\kappa = \cup\{\mathcal{U}_i : i < \kappa\}$

$\kappa\}$. Let $\varphi(\bar{x}, \bar{y}) \in \mathbb{L}(\tau_T)$ have the strong independence property, see Definition 0.2. We can choose $\langle C_\alpha^* : \alpha < \lambda^+ \rangle$ such that $C_\alpha^* \subseteq \text{nacc}(\alpha)$, $\text{otp}(C_\alpha^*) \leq \kappa$, $\beta \in C_\alpha^* \Rightarrow C_\beta^* = C_\alpha^* \cap \beta$ and $\lambda \mid \alpha \wedge \text{cf}(\alpha) = \kappa \Rightarrow \alpha = \sup(C_\alpha^*)$ and $\text{cf}(\alpha) \neq \kappa \Rightarrow \text{otp}(C_\alpha^*) < \kappa$.

[See [6] but for completeness we show this; by induction on $\alpha < \lambda^+$ we choose $\langle C_\varepsilon^* : \varepsilon < \lambda \rangle$ such that:

- (a) the relevant demand holds
- (b) if $\alpha = \beta + 1$, $C \subseteq \lambda\beta$, $(\forall i \in C)(C_i^* = C \cap i)$ and $\text{otp}(C) < \kappa$ then for some $i \in (\lambda\beta, \lambda\alpha)$ we have $C_{i+1}^* = C$.

As $\lambda = \lambda^{<\kappa}$ this is easy but we elaborate. For $\alpha = 0$ trivial for α limit obvious. Assume $\alpha = \beta + 1$ let $\alpha_* = \lambda\alpha$, $\beta_* = \lambda\beta$ and $\langle C_i^* : i < \beta_* \rangle$ has been defined.

First, we choose $C_{\beta_*}^*$. If $\text{cf}(\beta_*) \neq \kappa$ let $C_{\beta_*}^* = \emptyset$, so assume $\text{cf}(\beta_*) = \kappa$ then necessarily $\text{cf}(\beta) = \kappa$.

Let $\langle \alpha_\varepsilon : \varepsilon < \kappa \rangle$ be increasing with limit β , and choose $\beta_\varepsilon \in [\lambda\alpha_\varepsilon, \lambda\alpha_\varepsilon + \lambda)$ by induction on $\varepsilon < \kappa$ such that $C_{\beta_\varepsilon}^* = \{\beta_\zeta : \zeta < \varepsilon\}$.

Lastly, let $C_{\lambda\alpha}^* := \{\beta_\varepsilon : \varepsilon < \kappa\}$. So $C_{\beta_*}^*$ has been defined in any case.

Now let $\mathcal{C}_\alpha = \{C \subseteq \lambda\beta : \text{otp}(C) < \kappa \text{ and } \gamma \in C \Rightarrow C_\gamma^* = C \cap \gamma\}$, so $|\mathcal{C}| \leq \lambda$, also $\emptyset \in \mathcal{C}$, so let $\langle C_{\lambda\beta+i}^* : i \in (0, \lambda) \rangle$ list \mathcal{C}_α possibly with repetitions. So we have defined $\langle C_\varepsilon^* : \varepsilon < \lambda \rangle$, so have carried the induction.]

Let $S_* = \{\mu : \mu = \beth_{\alpha+\omega}$ for some $\alpha < \lambda\}$. Let E_* , $\langle C_\alpha : \alpha < \lambda \rangle$ be such that:

- ⊗₁ (a) $C_\alpha \subseteq \alpha \cap S_*$
- (b) $\beta \in C_\alpha \Rightarrow C_\beta = C_\alpha \cap \beta$
- (c) $\text{otp}(C_\alpha) \leq \theta$
- (d) E_* is the club of λ included in $\{\delta < \lambda : \theta < \delta = \beth_\delta\}$
- (e) $\text{otp}(C_\alpha) = \theta$ iff $\alpha \in E_* \cap S_\theta^\lambda$
- (f) if $\alpha \in S := E_* \cap S_\theta^\lambda$ then $\alpha = \sup(C_\alpha)$
- (g) if $\alpha \in E_*$ and $i < \kappa$ then $|\alpha \cap \mathcal{U}_i^-| = |\alpha|$.

[Why can we choose? By induction on the cardinal $\chi \in [\aleph_0, \lambda)$ we choose $\langle C_\alpha : \alpha < \beth_\chi \rangle$ and $E_\chi = E_* \cap \beth_\chi$ such that the relevant demands hold and: if $\chi = 2^{\lambda_1}$ and C is a subset of $S_* \cap \beth_{\chi_1}$ of order type $< \theta$ satisfying $\alpha \in C \Rightarrow C_\alpha = C \cap \alpha$ then for some $\alpha \in S_* \cap (\beth_{\chi_1}, \beth_\chi)$ we have $C_\alpha = C$. Why this extra induction hypothesis help? As arriving to $\alpha \in S$ so $\alpha = \beth_\alpha$ let $\langle \chi_i : i < \theta \rangle$ be an increasing sequence of cardinals with limit $\beth_\alpha = \alpha$ and we choose $\alpha_i \in (\beth_{\chi_i}, \beth_{2^{\chi_i}}) \cap S_*$ by induction on $i < \theta$ such that $C_{\alpha_i} = \{\alpha_j : j < i\}$ and the let $C_\chi = \{\alpha_i : i < \theta\}$.]

We shall prove that

- ⊗₂ if \square_2 below holds, then there is a β such that \odot_2 holds where:
- \square_2 (a) $\alpha < \lambda^+$, $i < \kappa$
- (b) f is a one-to-one function from M_α into $\mathcal{U}'_i = \cup\{\mathcal{U}_j : j < i\}$
- (c) $E \subseteq E_*$ is a club of λ such that $\delta \in E \Rightarrow f(M_\alpha) \upharpoonright \delta \prec f(M_\alpha)$
- (d) $\bar{\mathcal{P}} = \langle \mathcal{P}_\delta : \delta \in S \rangle$
- (e) $\mathcal{P}_\delta \subseteq \mathcal{P}(\delta)$ and $\emptyset \in \mathcal{P}_\delta$ and $\mathcal{P}_\delta \subseteq \bigcup_{\ell \leq 2} \mathcal{P}_\delta^{*,\ell}$ where²

²note that $\mathcal{P}_{\delta,i}^{*,1}, \mathcal{P}_{\delta,i}^{*,2}$ are the families of sets we like to ignore as they are influenced by our choices for $\delta_1 \in S \setminus \{\delta\}$, so we work to have them families of bounded subsets of δ .

- (α) $\mathcal{P}_\delta^{*,0} = \{A \subseteq \delta : \sup(A) = \delta \text{ and } A \subseteq \cup\{[\mu, 2^\mu) : \mu \in C_\delta\}\}$,
- (β) $\mathcal{P}_\delta^{*,1} = \cup\{\mathcal{P}_{\delta_1}^{*,0} : \delta_1 \in S \cap \delta\}$,
- (γ) $\mathcal{P}_\delta^{*,2} = \{A \subseteq \delta : \text{for some } \delta_1 \in \lambda \setminus (\delta + 1) \text{ we have } A \subseteq \cup\{[\partial, 2^\partial) : \partial \in C_{\delta_1} \cap \delta\}\}$
- (f) if $\delta_1 < \delta_2$ are from S then
 - (α) $A \in \mathcal{P}_{\delta_1} \Rightarrow A \in \mathcal{P}_{\delta_2}$
 - (β) $A \in \mathcal{P}_{\delta_2} \Rightarrow A \cap \delta_1 \in \mathcal{P}_{\delta_1}$,
 - (γ) for any $\delta \in S$ the family $\mathcal{P}_\delta^{*,1} \cup \mathcal{P}_\delta^{*,2}$ is a set of bounded subsets of δ ; (this follows)
- (g) $\bar{b}_{\delta, \mathcal{U}} \in M_\alpha$ for $\delta \in E \cap S, \mathcal{U} \in \mathcal{P}_\delta$ are such that

$$\bar{b}_{\delta_1, \mathcal{U}_1} = \bar{b}_{\delta_2, \mathcal{U}_2} \wedge \mathcal{U}_1 \in \mathcal{P}_{\delta_1} \wedge \mathcal{U}_2 \in \mathcal{P}_{\delta_2} \Rightarrow \delta_1 = \delta_2 \wedge \mathcal{U}_1 = \mathcal{U}_2$$
- \odot_2 (α) $\beta \in (\alpha, \lambda^+)$
- (β) there are $\bar{a}_\gamma \subseteq M_\beta$ for $\gamma < \lambda$ such that for a club of $\delta \in E$, if $\delta \in S$ then the following conditions on $\mathcal{U} \subseteq \delta$ are equivalent:
 - (i) $\mathcal{U} \in \mathcal{P}_\delta$
 - (ii) for some $\bar{b} \in {}^{\text{lg}(\bar{y})}M_\alpha$ we have: for every $\gamma < \delta$, $M_\beta \models \varphi[\bar{a}_\gamma, \bar{b}]$ iff $\gamma \in \mathcal{U}$
 - (iii) clause (ii) holds for $\bar{b} = \bar{b}_{\delta, \mathcal{U}}$ and $\mathcal{U} \in \mathcal{P}_\delta$

[Why? For each $\delta \in E \cap S$ let $\langle \mathcal{U}_{\delta, \varepsilon} : \varepsilon < |\mathcal{P}_\delta| \leq 2^{|\delta|} \rangle$ list \mathcal{P}_δ and let $\bar{b}_{\delta, \varepsilon} := \bar{b}_{\delta, \mathcal{U}_{\delta, \varepsilon}}$.
Let

$$\Gamma = \{\varphi(\bar{x}_\gamma, \bar{b}_{\delta, \varepsilon})^{\text{if } (\gamma \in \mathcal{U}_{\delta, \varepsilon})} : \gamma < \lambda, \delta \in E \text{ and } \varepsilon < |\mathcal{P}_{\delta, i}|\} \\ \cup \{\neg\varphi(\bar{x}_\gamma, \bar{b}) : \gamma < \lambda, \bar{b} \in {}^{\text{lg}(\bar{y})}M_\alpha \text{ and for no } \delta \in E, \varepsilon < |\mathcal{P}_\delta| \text{ do we have } \bar{b} = \bar{b}_{\delta, \varepsilon}\}.$$

As $\varphi(\bar{x}, \bar{y})$ has the strong independence property, recalling that by clause (g) of \square_2 the sequence $\langle \bar{b}_{\delta, \varepsilon} : \delta \in E \cap S \text{ and } \varepsilon < |\mathcal{P}_\delta| \rangle$ is with no repetitions, clearly Γ is finitely satisfiable in M_α , but M is λ^+ -saturated, $M_\alpha \prec M$ and $|\Gamma| = \lambda$ hence we can find $\bar{a}_\gamma \in {}^{\text{lg}(\bar{x})}M$ for $\gamma < \lambda$ such that the assignment $\bar{x}_\gamma \mapsto \bar{a}_\gamma$ ($\gamma < \lambda$) satisfies Γ in M . Lastly, choose $\beta \in (\alpha, \lambda^+)$ such that $\{\bar{a}_\gamma : \gamma < \lambda\} \subseteq M_\beta$.

Now check recalling $\emptyset \in \mathcal{P}_\delta$ for $\delta \in S$.]

Note

- \odot_3 in \odot_2 if h is a one-to-one mapping from M_β into \mathcal{U}_i extending f then for some club E of λ if for every $\delta \in S \cap E$ we have $(\forall \gamma < \lambda)(\gamma < \delta \rightarrow h(\bar{a}_\gamma) \in {}^{\text{lg}(\bar{x})}\delta)$ and so for every $\mathcal{U} \subseteq \delta$ the conditions (i), (ii), (iii) from \odot_2 are equivalent.

Next we can choose \bar{f} such that

- \otimes_3 (a) $\bar{f} = \langle f_\alpha : \alpha < \lambda^+ \rangle$
- (b) f_α is a one-to-one function from M_α into $\mathcal{U}_{\text{otp}(C_\alpha^*)}$
- (c) if $\alpha \in C_\beta^*$ then $f_\alpha \subseteq f_\beta$.

Now

- ⊗₄ for every $\alpha < \lambda^+$ there is $\bar{\mathcal{P}}^\alpha = \langle \mathcal{P}_\varepsilon^\alpha : \varepsilon \in S \rangle$ such that
- (i) $\mathcal{P}_\varepsilon^\alpha \subseteq \mathcal{P}(\varepsilon)$ are as in $\square_2(e), (f)$ above
 - (ii) for every $\beta \leq \alpha$, for a club of δ 's from S we have $\mathcal{P}_\delta^\alpha \notin \text{inv}_{7,\kappa}^\varphi(\delta, f_\beta(M_\beta))$.

[Why? For every $\beta \leq \alpha$ and $\delta \in (\kappa, \lambda)$ we have $\text{inv}_{7,\kappa}^\varphi(\delta, f_\beta(M_\beta))$ is a subset of $\mathcal{P}(\mathcal{P}(\delta))$ of cardinality $\leq 2^{|\delta|}$. As the number of β 's is $\leq \lambda$, by diagonalization we can do this: let $\alpha + 1 = \bigcup_{\varepsilon < \lambda} u_\varepsilon$ and $u_\varepsilon \in [\alpha + 1]^{<\lambda}$ increasing continuous for $\varepsilon < \lambda$; moreover, $\alpha < \lambda \Rightarrow u_\varepsilon = \alpha$ and $\alpha \geq \lambda \Rightarrow u_\varepsilon \cap \lambda \subseteq \varepsilon$ and $|u_\varepsilon| \leq |\varepsilon|$. By induction on $\varepsilon \in (\kappa, \lambda) \cap S$ choose $\mathcal{P}_\varepsilon^\alpha \subseteq \bigcup_{\ell < 3} \mathcal{P}_{\alpha_\varepsilon}^{*,\ell}$ which includes $\cup \{ \mathcal{P}_\zeta^\alpha : \zeta \in u_\varepsilon \cap S \} \cup \mathcal{P}_\varepsilon^{*,2}$ and satisfies $\mathcal{P}_\varepsilon^{*,0} \cap \mathcal{P}_\varepsilon^\alpha \in \mathcal{P}(\mathcal{P}_{\varepsilon,i}^{*,0}) \setminus \{ \mathcal{P} \cap \mathcal{P}_\varepsilon^{*,0} : \mathcal{P} \in \text{inv}_{7,\kappa}^\varphi(\varepsilon, f_\beta(M_\beta)), \beta \in u_\varepsilon \}$. Note that for each $\beta \leq \alpha$ the set $\{ \varepsilon < \lambda : \beta \in u_\varepsilon \}$ contains an end-segment of λ hence a club of λ as required.]

Now choose pairwise distinct $\bar{b}_{\delta,\mathcal{U}} \in {}^{\ell g(\bar{y})}(M_0)$ for $\delta \in E_*$, $\mathcal{U} \in \mathcal{P}_\delta^{*,0}$

- ⊗₅ for every $\alpha_* \leq \alpha < \lambda^+$ for some $\beta \in (\alpha, \lambda^+)$ and $\bar{a}_\gamma \in {}^{\ell g(\bar{x})}M_\beta$ for $\gamma < \lambda$ the condition in clause (γ) of \odot_2 holds with $\bar{\mathcal{P}}^{\alpha_*}$ here standing for $\bar{\mathcal{P}}$ there and the $\bar{b}_{\delta,\mathcal{U}}$ chosen above.

[Why? By ⊗₂.]

- ⊗₆ let $E = \{ \delta < \lambda^+ : \delta \text{ is a limit ordinal such that for every } \alpha_* \leq \alpha < \delta \text{ there is } \beta < \delta \text{ as in } \otimes_5 \}$.

Clearly E is a club of λ^+ .

- ⊗₇ if $\delta_1 < \delta_2$ are from $E \cap S_\kappa^{\lambda^+}$ then $M_{\delta_1}, M_{\delta_2}$ are not isomorphic.

[Why? Let $\alpha_* = \min(C_{\delta_2}^* \setminus \delta_1)$. We consider $\bar{\mathcal{P}}^{\alpha_*}$ which is from ⊗₄. On the one hand $\{ \varepsilon < \lambda : \mathcal{P}_\varepsilon^{\alpha_*} \notin \text{inv}_{7,\kappa}^\varphi(\varepsilon, f_{\delta_1}(M_{\delta_1})) \}$ contains a club by ⊗₄(ii). Note that $\langle f_\alpha : \alpha \in C_{\delta_2}^* \setminus \delta_1 \rangle$ is \subseteq -increasing sequence of functions with union f_{δ_2} . On the other hand choose an increasing $\langle \alpha_i : i < \kappa \rangle$ with limit δ_2 satisfying $\alpha_0 = 0, \alpha_1 = \delta_1$ such that $(\alpha_*, \alpha_{1+i}, \alpha_{1+i+1})$ are like $(\alpha_*, \alpha, \beta)$ in ⊗₅ for each $i < \kappa$ and $i \in (1, \kappa) \Rightarrow \alpha_i \in C_{\delta_2}^*$. Now by 1.8, $\{ \varepsilon < \lambda : \mathcal{P}_\varepsilon^{\alpha_*} \in \text{inv}_{7,\kappa}^\varphi(\varepsilon, f_{\delta_2}(M_{\delta_2})) \}$ contains a club. Hence by the last sentence and the end of the previous paragraph $M_{\delta_1} \not\cong M_{\delta_2}$ as required.]

So we are done. □_{1.9}

Remark 1.12. We can avoid using C_δ^* and also C_δ (e.g. using $A \in \mathcal{P}_\delta^{*,0} \Rightarrow \text{otp}(A) = \delta$) but this seems less transparent.

Proof. Proof of 1.10 Similar but easier (for λ regular not strong limit (but $2^\lambda > 2^{<\lambda}$) also easy), or see the proof of 2.9. □_{1.10}

§ 2. INDEPENDENT T

We would like to do something similar to §1, but our control on the relevant family of subsets of M is less tight. We control it to some extent by using the completion of a free Boolean algebra.

{2d.1}

Context 2.1. T a complete first order theory, $\varphi(x, \bar{y})$ has the independence property (of course the existence of such φ follows from the strong independence property but is weaker).

{2d.4}

We continue [5, 2.1-2.12], but we do not rely on it.

Definition 2.2. For a set I let

- (a) $\mathbb{B} = \mathbb{B}_I$ be the Boolean Algebra generated by $\langle e_t : t \in I \rangle$ freely,
- (b) \mathbb{B}_I^c is the completion of \mathbb{B}
- (c) for $J \subseteq I$ let $\mathbb{B}_{I,J}^c$ be the complete subalgebra of \mathbb{B}_I^c generated by $\{e_s : s \in J\}$
- (d) let $\text{uf}(\mathbb{B}_I^c)$ be the set of ultrafilters on I .

{2d.7}

Claim 2.3. Assume

- ⊗ (a) $M \models T$
- (b) $\bar{b}_t \in {}^{\ell g(\bar{y})}M$ for $t \in I$
- (c) $\langle \varphi(x, \bar{b}_t) : t \in I \rangle$ is an independent sequence of formulas.

Then there is a function F from ${}^{\ell g(\bar{y})}M$ to $\mathbb{B} = \mathbb{B}_I^c$ such that

- (α) $F(\bar{b}_t) = e_t$
- (β) for every ultrafilter D of \mathbb{B} there is $p = p_D = p_{F,D} \in \mathbf{S}_\varphi(M)$, in fact, a unique one, such that for every $\bar{b} \in {}^{\ell g(\bar{y})}M$ we have $\varphi(x, \bar{b}) \in p \Leftrightarrow F(\bar{b}) \in D$.

Remark 2.4. 1) Note that the mapping $D \mapsto p_D$ is not necessarily one to one, but $D_1 \cap \{e_t : t \in I\} \neq D_2 \cap \{e_t : t \in I\} \Rightarrow p_{D_1} \neq p_{D_2}$.

2) If $I = I_1 \cup I_2, I_1 \cap I_2 = \emptyset$ and $|I_2| = |I_1|^{\aleph_0}$ then we can find a mapping F from ${}^{\ell g(\bar{y})}M$ onto (not just into) $\mathbb{B} = \mathbb{B}_{I_1}^c$ such that clause (α), (β) are satisfied.

Proof. Clearly $\mathcal{P}(M)$ is a Boolean algebra and $\{\varphi(M, \bar{b}_t) : t \in M\}$ generates freely a subalgebra of $\mathcal{P}(M)$ which we call \mathbb{B}' . So there is a homomorphism h from \mathbb{B}' into \mathbb{B} mapping $\varphi(M, \bar{b}_t)$ to e_t (moreover h is unique and is an isomorphism from \mathbb{B}' onto $\mathbb{B}_I \subseteq \mathbb{B}_I^c$). So h is a homomorphism from $\mathbb{B}' \subseteq \mathcal{P}(M)$ into \mathbb{B}_I^c , which is a complete Boolean algebra hence there is a homomorphism h^+ from the Boolean algebra $\mathcal{P}(M)$ into \mathbb{B}^c extending h .

{2d.14}

Lastly, define $F : {}^{\ell g(\bar{y})}M \rightarrow \mathbb{B}^c$ by $F(\bar{b}) = h^+(\varphi(M, \bar{b}))$. Now check. □_{2.3}

Conclusion 2.5. Assume ⊗ from 2.3 and

- (a) $I = \lambda$ is regular uncountable
- (b) $|M| \subseteq \mathcal{U} \subseteq \lambda$
- (c) D_α is an ultrafilter of \mathbb{B}_I^c for $\alpha < \lambda$
- (d) $\mathcal{U} \setminus |M|$ is unbounded in λ .

Then we can find $\langle a_\alpha : \alpha < \lambda \rangle$ and N such that

- (α) $M \prec N$
- (β) $|N| \subseteq \mathcal{U}$
- (γ) $a_\alpha \in N$ for $\alpha < \lambda$
- (δ) a_α realizes $p_{D_\alpha} \in \mathbf{S}_\varphi(M)$.

Remark 2.6. Conclusion 2.5 is easy but intended to clarify how we shall use the ultrafilters, so is quoted toward the end of the section.

Proof. Should be clear. $\square_{2.5}$

Discussion 2.7. Note that compared to §1 instead $\bar{x}, \bar{y}, \bar{a}_\alpha, \bar{b}_\beta$ we have $x, \bar{y}, a_\alpha, \bar{b}_\beta$. Compared to §1, we have less control over $\{\text{tp}(a, M, N) : a \in N\}$. There, for the sequences \bar{b} of M which are not among $\{\bar{b}_\gamma : \gamma < \lambda\}$, we can demand $N \models \neg\varphi[\bar{a}_\gamma, \bar{b}]$ for $\gamma < \lambda$ so $\text{tp}_\varphi(\bar{a}_\gamma, M, N)$ can be clearly read. Here the complete Boolean Algebra \mathbb{B}_I^c is helping, a small price is that we need $\theta > \aleph_0$.

In order to try to keep track of what is going on we shall use only $\text{tp}(a_\gamma, M, N)$ of the form p_D for ultrafilter D on \mathbb{B}_I^c . Further, we better have, e.g. a nice function π from ${}^\lambda 2$ to $\text{uf}(\mathbb{B}_I^c)$ such that $(e_\alpha \in \pi(\eta)) \Leftrightarrow \eta(\alpha) = 1$.

A possible approach is: we define $\langle M_{\eta, u} : \eta \in \mathcal{T} \subseteq \text{des}(\lambda), u \in \mathcal{P}(n_\eta) \rangle$ as in [7, §3] and we define $D_\eta \in \text{uf}(\mathbb{B}^c \cap M)$ such that $\alpha \in M_\eta \cap \lambda \Rightarrow [e_\eta^{\eta(\alpha)} \in D_\eta]$ and $\bigcup_\eta D_\eta \in \text{uf}(\mathbb{B}^c)$.

We need some continuity so each “ $e \in D_\eta$ ” ($e \in \mathbb{B}^c$) depends on $\eta \upharpoonright u_e$ for some “small” $u_e \subseteq \lambda$. $\{3e.7\}$

Theorem 2.8. *In Theorem 1.9 it suffices to assume \boxtimes' which means clauses (b),(c),(d) of \boxtimes and*

- (a)' T has the independence property. $\{3e.14\}$

Theorem 2.9. *In Theorem 1.10 it suffices to assume \boxtimes' of 2.8.*

Proof. Proof of 2.8 Just combine the proofs of 1.9 from §1 and 2.9 below. $\square_{2.8}$

Proof. Proof of 2.9 As in the proof of 1.9 we can assume λ is strongly inaccessible though the proof is just easier otherwise. We let

- \otimes_1 (a) $E_* = \{\delta < \lambda : \delta = \beth_\delta\}$, a club of λ
- (b) $S_* = \{\beth_{\alpha+\omega} : \alpha < \lambda\}$
- (c) choose a regular uncountable $\theta < \lambda$

and let

- \otimes_2 (a) $S = \{\delta \in E_* : \text{cf}(\delta) = \theta\} = S_\theta^\lambda \cap E_*$
- (b) let \bar{C} be as in \otimes_1 of the proof of 1.9, in particular $\bar{C} = \langle C_\alpha : \alpha < \lambda \rangle, C_\alpha \subseteq S_*$, $\text{otp}(C_\alpha) \leq \theta, \alpha \in C_\beta \Rightarrow C_\alpha = C_\beta \cap \alpha$ and $\alpha \in S \Leftrightarrow \alpha = \sup(C_\alpha) \Leftrightarrow \text{otp}(C_\alpha) = \theta$ and $\alpha \in S \Rightarrow \sup(C_\alpha) = \alpha$
- (c) for $\mu \in S$ let $A_\mu^* = \cup\{\chi, 2^\chi : \chi \in C_\mu\}$.

Let \mathbb{D}_* be an ultrafilter of \mathbb{B}_λ^c such that $e_\alpha \notin \mathbb{D}_*$ for $\alpha < \lambda$.

Now for $\eta \in {}^\lambda 2$ we choose \mathbb{D}_η such that

- ⊗₃ (a) \mathbb{D}_η is an ultrafilter of \mathbb{B}_λ^c
- (b) if $e \in \mathbb{D}_* \subseteq \mathbb{B}_\lambda^c$ belongs to $\mathbb{B}_{\lambda, \eta^{-1}\{0\}}^c$ (see 2.2, the completion of the subalgebra of \mathbb{B}_λ^c generated by $\{e_\alpha : \eta(\alpha) = 0\}$) then $e \in \mathbb{D}_\eta$.
- (c) if $\alpha < \lambda$ and $\eta(\alpha) = 1$ then $e_\alpha \in \mathbb{D}_\eta$.

So

- ⊗₄ (a) if $\eta \in {}^\lambda 2$ is constantly zero then $\mathbb{D}_\eta = \mathbb{D}_*$
- (b) $e_\alpha \in \mathbb{D}_\eta \Leftrightarrow \eta(\alpha) = 1$ for $\alpha < \lambda, \eta \in {}^\lambda 2$.

Now let $\bar{\eta} = \langle \eta_\varepsilon : \varepsilon < \lambda \rangle$ be a sequence of members of ${}^\lambda 2$ and below we shall be interested mainly in the case $\alpha = \mu \in S$.

Define

- ⊗₅ for $e \in \mathbb{B}_\lambda^c$ and $\alpha \leq \lambda$ we let $Y_{\bar{\eta}, e}^\alpha := \{\varepsilon < \alpha : e \in \mathbb{D}_{\eta_\varepsilon}\}$
- ⊗₆ $\mathcal{P}_{\bar{\eta}, \alpha} := \{Y_{\bar{\eta}, e}^\alpha : e \in \mathbb{B}_\lambda^c\}$.

Now what can we say on $\mathcal{P}_{\bar{\eta}, \mu}$ for $\mu \in S$? As we can consider $e \in \{e_\alpha : \alpha \in [\mu, 2^\mu)\}$, clearly

- ⊗₇ $\{\{\varepsilon < \mu : \eta_\varepsilon(\alpha) = 1\} : \alpha \in [\mu, 2^\mu)\} \subseteq \mathcal{P}_{\bar{\eta}, \mu} \subseteq \mathcal{P}(\mu)$.

This may be looked at as a “lower bound” of $\mathcal{P}_{\bar{\eta}, \mu}$. Naturally we try to get also an “upper bound” to $\mathcal{P}_{\bar{\eta}, \mu}$; now note

- ⊗₈ if $e \in \mathbb{B}_\lambda^c$ then $Y_{\bar{\eta}, -e}^\mu = \mu \setminus Y_{\bar{\eta}, e}^\mu$.

Now define (recalling A_μ^* is from ⊗₂(c))

- ⊗₉ Ξ is the set of $\bar{\eta}$ of the form $\langle \eta_\varepsilon : \varepsilon < \lambda \rangle$ such that:
 - (a) $\eta_\varepsilon \in {}^\lambda 2$ for every $\varepsilon < \lambda$,
 - (b) if $\eta_\varepsilon(\alpha) = 1$ then $(\exists \mu \in S)[\mu \leq \alpha < 2^\mu \wedge \varepsilon \in A_\mu^*]$,
 - (c) if $\mu \in S$ and $u \subseteq [\mu, 2^\mu)$ is countable then $\{\varepsilon \in A_\mu^* : \text{if } \alpha \in u \text{ then } \eta_\varepsilon(\alpha) = 0\}$ is of cardinality μ .

Also (by our knowledge of the completion of a free Boolean algebra, \mathbb{B}_λ^c satisfies the c.c.c.) for every $e \in \mathbb{B}_\lambda^c$ we can choose u_e such that:

- ⊕₁ (a) $u_e \subseteq \lambda$ is countable
- (b) $e \in \mathbb{B}_{\lambda, u_e}^c$.

So by clause (b) of ⊗₃ clearly

- ⊕₂ if $\bar{\eta} \in \Xi, e \in \mathbb{B}_\lambda^c, \varepsilon < \mu \in S$ and $u_e \subseteq \eta_\varepsilon^{-1}\{0\}$ then $e \in \mathbb{D}_{\eta_\varepsilon} \Leftrightarrow e \in \mathbb{D}_*$

hence

- ⊕₃ if $\bar{\eta} \in \Xi, e \in \mathbb{B}_\lambda^c \cap \mathbb{D}_*$ and $\mu \in S$ then $Y_{\bar{\eta}, e}^\mu \supseteq \{\varepsilon < \mu : u_e \subseteq \eta_\varepsilon^{-1}\{0\}\}$.

Next

- ⊕₄ for $\bar{\eta} \in \Xi$
 - (a) let $\mathcal{D}_{\bar{\eta}, \mu}$ be the filter on μ generated by $\{A_\mu^*\} \cup \{\{\varepsilon < \mu : u \subseteq \eta_\varepsilon^{-1}\{0\} \text{ and } \varepsilon > \zeta\} : \zeta < \mu \text{ and } u \subseteq \lambda \text{ is countable}\}$
 - (b) let $\mathcal{I}_{\bar{\eta}, \mu}$ be the dual ideal.

Clearly

- ⊕₅ (a) if $\bar{\eta} \in \Xi$, $\mu \in S$ and $\alpha \in \lambda \setminus [\mu, 2^\mu)$ then $\{\varepsilon < \mu : \eta_\varepsilon(\alpha) \neq 0\}$ is a bounded subset of μ ,
- (b) if $\bar{\eta} \in \Xi$ and $\mu \in S$ then $\mathcal{D}_{\bar{\eta}, \mu}$ is a uniform \aleph_1 -complete filter on μ (recalling $\text{cf}(\mu) = \theta > \aleph_0$ as $\mu \in S$) and $\emptyset \notin \mathcal{D}_{\bar{\eta}, \mu}$.

[Why? See ⊗₉.]

Now by ⊕₃ we have $\bar{\eta} \in \Xi \wedge e \in \mathbb{B}_\lambda^c \cap \mathbb{D}_* \Rightarrow Y_{\bar{\eta}, e}^\mu \in \mathcal{D}_{\bar{\eta}, \mu}$ so recalling ⊗₈ we have $e \in \mathbb{B}_\lambda^c \setminus \mathbb{D}_* \Rightarrow Y_{\bar{\eta}, e}^\mu = \emptyset \pmod{\mathcal{D}_{\bar{\eta}, \mu}}$ hence

$$\oplus_6 \mathcal{D}_{\bar{\eta}, \mu} \subseteq \{X \subseteq \mu : X \in \mathcal{D}_{\bar{\eta}, \mu} \text{ or } \mu \setminus X \in \mathcal{D}_{\bar{\eta}, \mu}\}.$$

Now

- ⊙₁ if $\mu \in S$ then we can find \bar{A}_μ^ξ for $\xi < 2^{2^\mu}$ such that:
- (a) $\bar{A}_\mu^\xi = \langle A_\gamma^\xi : \gamma \in [\mu, 2^\mu) \rangle$
- (b) A_γ^ξ is an unbounded subset of A_μ^*
- (c) $\mathcal{D}_\mu^\xi, \mathcal{I}_\mu^\xi$ are well defined, i.e. $\emptyset \in \mathcal{D}_\mu^\xi$ when we let
- (α) \mathcal{D}_μ^ξ be the \aleph_1 -complete filter of subsets of μ generated by $\{A_\gamma^\xi \setminus \beta : \gamma \in [\mu, 2^\mu) \text{ and } \beta < \mu\}$ so $A_\mu^* \setminus \beta \in \mathcal{D}_\mu^\xi$ for $\beta < \mu$
- (β) $\mathcal{I}_\mu^\xi = \{\mu \setminus B : B \in \mathcal{D}_\mu^\xi\}$, i.e. the dual ideal
- (d) moreover if $\xi^1 \neq \xi^2$ are $< 2^{2^\mu}$, then

$$\{A_\mu^* \setminus A_\gamma^{\xi^1} : \gamma \in [\mu, 2^\mu)\} \not\subseteq \mathcal{D}_\mu^{\xi^2} \cup \mathcal{I}_\mu^{\xi^2}.$$

[Why ⊙₁ holds? As $|A_\mu^*| = |\mu|$ is a strong limit cardinal of cofinality $\theta > \aleph_0$ clearly $\mu = |A_\mu^*| = |A_\mu^*|^{\aleph_0}$ hence by [2] there is a sequence $\langle B_\gamma : \gamma \in [\mu, 2^\mu) \rangle$ of subsets of A_μ^* such that any non-trivial Boolean combination of countably many of them has cardinality μ . Let $\langle U_\xi : \xi < 2^{2^\mu} \rangle$ be a sequence of pairwise distinct subsets of $[\mu, 2^\mu)$ each of cardinality $2^{|\mu|}$ no one included in another and let $\langle A_\gamma^\xi : \gamma \in [\mu, 2^{|\mu|}) \rangle$ list $\{B_\gamma : \gamma \in U_\xi\}$.

Now check.]

- ⊙₂ in ⊙₁ it follows that
- (e) for every $\mathcal{P} \subseteq \mathcal{P}(\mu)$ for at most one $\xi < 2^{2^\mu}$ we have

$$\{A_\mu^* \setminus A_\gamma^\xi : \gamma \in [\mu, 2^\mu)\} \subseteq \mathcal{P} \subseteq \mathcal{D}_\mu^\xi \cup \mathcal{I}_\mu^\xi.$$

- ⊙₃ for every $\bar{\xi} = \langle \xi(\mu) : \mu \in S \rangle \in \Pi\{2^{2^\mu} : \mu \in S\}$ there is $\bar{\eta} = \bar{\eta}_{\bar{\xi}}$ such that:
- (a) $\bar{\eta}_{\bar{\xi}} \in \Xi$ so $\bar{\eta}_{\bar{\xi}} = \langle \eta_{\bar{\xi}, \varepsilon} : \varepsilon < \lambda \rangle$
- (b) if $\mu \in S$, $\gamma \in [\mu, 2^\mu)$ then $\{\varepsilon \in \lambda : \eta_{\bar{\xi}, \varepsilon}(\gamma) = 1\} = A_\mu^* \setminus A_\gamma^{\xi(\mu)}$.

[Why? Just read the definition of Ξ in ⊗₉ and \bar{A}_μ^ξ in ⊙₁.]

$$\oplus_4 \text{ if } \mu \in S \text{ then } \mathcal{D}_{\bar{\eta}_{\bar{\xi}}, \mu} \cup \mathcal{I}_{\bar{\eta}_{\bar{\xi}}, \mu} = \mathcal{D}_\mu^{\xi(\mu)} \cup \mathcal{I}_\mu^{\xi(\mu)}.$$

[Why? Easy, recalling ⊕₅(a).]

- ⊙₅ if $\gamma(*) < \lambda^+$ and $\bar{\mathbf{P}}^\gamma = \langle \mathbf{P}_\mu^\gamma : \mu \in S \rangle$, for $\gamma < \gamma(*)$ where $\mathbf{P}_\mu^\gamma \subseteq \mathcal{P}(\mathcal{P}(\mu))$ has cardinality $\leq 2^\mu$ for $\mu \in S$, $\gamma < \gamma(*)$ then we can find $\bar{\xi} = \langle \xi(\mu) : \mu \in S \rangle \in \Pi\{2^{2^\mu} : \mu \in S\}$ such that for every $\gamma < \gamma(*)$ the following set is not stationary: $S_{\bar{\eta}, \gamma} = \{\mu \in S : \text{for some } \mathcal{P} \in \mathbf{P}_\mu^\gamma \text{ we have } \{A_\gamma^{\xi(\mu)} : \gamma \in [\mu, 2^\mu]\} \subseteq \mathcal{P} \subseteq \mathcal{D}_{\bar{\eta}, \mu} \cup \mathcal{I}_{\bar{\eta}, \mu}\}$.

[Why? Let $\langle u_\alpha : \alpha < \lambda \rangle$ be an increasing continuous sequence of subsets of $\gamma(*)$ with union $\gamma(*)$ such that $|u_\alpha| \leq |\alpha|$ for $\alpha < \lambda$. Now for each $\mu \in S$, the family $\cup\{\mathbf{P}_\mu^\gamma : \gamma \in u_\mu\}$ is a family of $\leq |u_\mu| \times 2^\mu$ subsets of $\mathcal{P}(\mu)$.

Now by clause (e) of ⊙₁ for each $\mu \in S$, $\gamma \in u_\mu$, $\mathcal{P} \in \mathbf{P}_\mu^\gamma$ let $\xi_{\mu, \gamma, \mathcal{P}} < 2^{2^\mu}$ be such that: if for some $\xi < 2^{2^\mu}$ we have $\{A_\gamma^\xi : \gamma \in [\mu, 2^\mu]\} \subseteq \mathcal{P} \subseteq \mathcal{D}_\mu^\xi \cup \mathcal{I}_\mu^\xi$ then $\xi_{\mu, \gamma, \mathcal{P}}$ is the first such ξ . Choose $\xi(\mu) < 2^{2^\mu}$ which does not belong to $\{\xi_{\mu, \gamma, \mathcal{P}} : \gamma \in u_\mu \text{ and } \mathcal{P} \in \mathbf{P}_\mu^\gamma\}$.

So let $\bar{\eta} = \bar{\eta}_{\langle \xi(\mu) : \mu \in S \rangle} \in \Xi$ be as in ⊙₃ now $\bar{\eta}$ is as required by ⊙₂, ⊙₃, ⊙₄.

Let us elaborate, why is it as required in ⊙₅?

First, clearly $\eta_\varepsilon \in {}^\lambda 2$ for $\varepsilon < \lambda$. Second, fix $\gamma < \gamma(*)$, then there is $\alpha < \lambda$ such that $\gamma \in u_\alpha$, so it suffices to show that, for any $\mu \in S \setminus \alpha$, we have $\mu \notin S_{\bar{\eta}, \gamma}$. So assume $\mathcal{P} \in \mathbf{P}_\mu^\gamma$ satisfies clause (e) of ⊙₁, and we should prove that $\neg[\{A_\gamma^{\xi(\mu)} : \gamma \in [\mu, 2^\mu]\} \subseteq \mathcal{P} \subseteq \mathcal{D}_{\bar{\eta}, \mu} \cup \mathcal{I}_{\bar{\eta}, \mu}]$; but if for some $\xi < 2^{2^\mu}$ we have $\{A_\gamma^\xi : \gamma \in [\mu, 2^\mu]\} \subseteq \mathcal{P} \subseteq \mathcal{D}_{\bar{\eta}, \mu} \cup \mathcal{I}_{\bar{\eta}, \mu}$ then necessarily $\xi = \xi_{\mu, \gamma, \mathcal{P}} \neq \xi(\mu)$, contradiction to ⊙₁.]

- ⊙₆ if $\langle M_\gamma : \gamma \leq \gamma(*) \rangle$ is a \prec -increasing continuous and $M_\gamma \in \text{EC}_\lambda(T)$ and $\bar{b}_\alpha \in {}^{\ell g(\bar{y})}(M_0)$ for $\alpha < \lambda$ are such that $\langle \varphi(x, \bar{b}_\alpha) : \alpha < \lambda \rangle$ is independent, then we can find N such that
- (α) $M_{\gamma(*)} \prec N \in \text{EC}_\lambda(T)$
 - (β) if $N \prec N' \in \text{EC}_\lambda(T)$ and $\gamma < \gamma(*)$ then ³ we have $\text{inv}_6^\varphi(M_\gamma, N') \notin \{\text{inv}_6^\varphi(M_{\gamma_1}, M_{\gamma_2}) : \gamma \leq \gamma(*) \text{ and } \gamma_1 < \gamma_2 \leq \gamma(*)\}$.

[Why? Without loss of generality the universe of $M_{\gamma(*)}$ is $\mathcal{U}_1 \in [\lambda]^\lambda$ such that $\lambda \setminus \mathcal{U}_1$ has cardinality λ . Let $\langle u_\alpha : \alpha < \lambda \rangle$ be as in the proof of ⊙₅.

For $\delta < \lambda$ and $\gamma(1) < \gamma(2) \leq \gamma(*)$ let $\mathbf{P}_\delta^{\gamma(1), \gamma(2)} = \text{inv}_\varphi^5(\delta, \text{id}_{N_{\gamma(2)}}, M_{\gamma(1)}, M_{\gamma(2)})$, see Definition 1.4, clearly $\text{inv}_6^\varphi(M_{\gamma(1)}, M_{\gamma(2)}) = \langle \mathbf{P}_\delta^{\gamma(1), \gamma(2)} : \delta < \lambda \rangle / \mathcal{D}_\lambda$. So it is enough⁴ to find N and sequence $\langle a_\alpha : \alpha < \lambda \rangle$ of elements of N such that $M_{\gamma(*)} \prec N$, $|N| = \lambda$ and for each $\gamma(0) \leq \gamma(*)$, for every $\mu \in S$ except non-stationarily many, the family

$$\{\{\gamma < \mu : N \models \varphi[a_\gamma, \bar{b}]\} : \bar{b} \in {}^{\ell g(\bar{y})}(M_{\gamma(0)})\}$$

is not in $\mathbf{P}_\mu := \cup\{\mathbf{P}_\mu^{\gamma(1), \gamma(2)} : \gamma(1) < \gamma(2) \leq \gamma(*) \text{ are from } u_\mu\}$.

We choose $\bar{\xi} = \langle \xi(\mu) : \mu \in S \rangle$ as in ⊙₅; let $\bar{\eta} = \bar{\eta}_{\bar{\xi}}$, see ⊙₃, so recalling ⊙₃ clearly $\langle \mathbb{D}_{\eta_\varepsilon} : \varepsilon < \lambda \rangle$ is well defined. Now for each $\varepsilon < \alpha$ letting F be from 2.3 for the model $M_{\gamma(*)}$ and the sequence $\langle \varphi(x, \bar{b}_\alpha) : \alpha < \lambda \rangle$, let $p_\varepsilon \in \mathbf{S}_\varphi(M_{\gamma(*)})$ be such that for every $\bar{b} \in {}^{\ell g(\bar{y})}(M_{\gamma(*)})$ we have $\varphi(x, \bar{b}) \in p_\varepsilon \Leftrightarrow F(\bar{b}) \in \mathbb{D}_{\eta_\varepsilon}$ so $\neg\varphi(x, \bar{b}) \in p_\varepsilon \Leftrightarrow F(\bar{b}) \notin \mathbb{D}_{\eta_\varepsilon}$.

³really any pregiven set of $\leq \lambda$ “forbidden” inv_6^φ is O.K. and can make it work for $\text{inv}_6^\varphi(N_\gamma, N')$ for every $\gamma < \gamma(*)$.

⁴can demand $\alpha < \lambda \Rightarrow {}^\omega(\omega(\alpha + 1))$ if $(N \setminus M_{\gamma(*)}) \cap [\omega\alpha, \omega\alpha + \omega)$ is infinite for every $\alpha < \lambda$.

So by 2.5 we can find an elementary extension N of $M_{\gamma(*)}$ and $a_\alpha \in N$ for $\alpha < \lambda$ such that a_α realizes p_α , and without loss of generality N has universe $\subseteq \lambda$ such that $\lambda \setminus |N|$ has cardinality λ . Concerning inv_φ^6 our demand concerns what occurs for a club of $\delta < \lambda$ for this. Let $E \subseteq E_*$ be a club of λ such that $\gamma < \delta \in E \Rightarrow a_\gamma \in N \cap \delta$. Now in $\odot_6(\beta)$ we promise something (given N) on “every N' such that ...”, so let $N \prec N' \in \text{EC}_\lambda(T)$, and without loss of generality the universe of N' is $\subseteq \lambda$ and let $\delta \in S \cap E$. For any $\gamma \leq \gamma(*)$ by \odot_5 , i.e. by the choice of $\bar{\xi}, \bar{\eta}_\xi$ above there is a club $E_\gamma \subseteq E$ of λ such that for any $\mu \in S \cap E_\gamma$, the set $S_{\bar{\eta}, \mu}$ from \odot_5 is disjoint to E_γ , hence the set $\mathcal{P}_{\mu, \gamma} := \{\{\gamma < \mu : N' \models \varphi[a_\gamma, \bar{b}]\} : \bar{b} \in {}^{\ell g(\bar{y})}(M_\gamma)\}$ does not belong to $\cup\{\mathbf{P}_\mu^{\gamma(1), \gamma(n)} : \gamma(1) < \gamma(2) \leq \gamma(*) \text{ are from } u_\mu\}$ so we are done.]

\odot_7 if $\langle M_\alpha : \alpha < \lambda^+ \rangle$ is as in \boxtimes' from 2.8 then for some club E of λ^+ , we have:
if $\alpha_1 < \alpha_2, \beta_1 < \beta_2$ are from E and $\alpha_2 \neq \beta_2$ then

$$(M_{\alpha_2}, M_{\alpha_1}) \not\cong (M_{\beta_2}, M_{\beta_1}).$$

[Why? For every $\beta < \lambda^+$ we apply \odot_6 to $\langle M_\alpha : \alpha \leq \beta \rangle$ and get N_β as there so $M_\beta \prec N_\beta \in \text{EC}_\lambda(T)$. As $M = \cup\{M_\gamma : \gamma < \lambda^+\}$ is saturated, without loss of generality $N_\beta \prec M$ hence for some $\xi_\beta < \lambda^+$ we have $N_\beta \prec M_{\xi_\beta}$.

Let $E = \{\delta < \lambda^+ : \delta \text{ a limit ordinal such that } \beta < \delta \Rightarrow \xi_\beta < \delta\}$.

Let $\alpha_1 < \alpha_2, \beta_1 < \beta_2$ be from E such that $\alpha_2 \neq \beta_2$ and we shall prove that $(M_{\alpha_2}, M_{\alpha_1})$ is not isomorphic to $(M_{\beta_2}, M_{\beta_1})$. By symmetry without loss of generality $\alpha_2 < \beta_2$ and let $\gamma(*) = \max\{\alpha_2, \beta_1\}$ so $\gamma(*) < \beta_2$. Now we apply \odot_6 with $\langle M_\gamma : \gamma \leq \gamma(*) \rangle, N, N', \gamma, \beta_1, \alpha_2, \alpha_2$ here standing for $\langle M_\gamma : \gamma \leq \gamma(*) \rangle, N, N', \gamma(0), \gamma(1), \gamma(2)$ there so we are clearly done. $\square_{2.9}$

§ 3. PRIVATE APPENDIX

Definition 3.1. 1) We say that T has the PO-property when some $\varphi(\bar{x}, \bar{y})$ has it where:

2) We say $\varphi(\bar{x}, \bar{y})$ has the PO-property in T when: for every partial order I and for some model M of T and $\bar{a}_t \in {}^{\ell g(\bar{y})}M$ for $t \in I$ we have:

(a) if $s, t \in I$ has no common upper bound then $M \models \neg(\exists \bar{x})(\varphi(\bar{x}, \bar{a}_s) \wedge \varphi(\bar{x}, \bar{a}_t))$

(b) if $n < \omega$ and $t_0, \dots, t_{n-1} \in I$ has a common upper bound then $(\exists \bar{x}) \bigwedge_{\ell < n} \varphi(\bar{x}, \bar{a}_{t_\ell})$.

{3n.7}

Claim 3.2. In $\text{Th}(M, N)$ we can interpret by $\mathbb{L}_{|T|+, \omega}(\tau_T)$ -formulas with parameters, a Boolean algebra \mathbb{B} isomorphic to \mathbb{B}^* and can interpret qualification over ideals of \mathbb{B} when:

- ⊗ (a) $M \prec N$ are models of T
- (b) N is $\|M\|^+$ -saturated
- (c) \mathbb{B}^* is a complete Boolean Algebra of cardinality λ
- (d) $\langle \varphi(x, \bar{a}_\alpha) : \alpha < \lambda \rangle$ is an independent sequence of formulas in M
- (e) $\lambda = \lambda^{\aleph_0}$ (for simplicity).

Remark 3.3. Does this mean to imply interpretations of large models of PA?

Proof. Let $\mathbb{B} = \mathbb{B}_\lambda^c$ and let $\langle d_\alpha : \alpha < \lambda \rangle$ list the elements of \mathbb{B} , see 2.2. Let \mathbb{B}_M be the Boolean algebra of subsets of M definable by formulas with parameters, i.e. $\{\psi(M, \bar{b}) : \psi(x, \bar{y}) \in \mathbb{L}(\tau_T) \text{ and } \bar{b} \in {}^{\ell g(\bar{y})}M\}$ and let \mathbb{B}_M^- be the subalgebra generated by $\{\varphi(M, \bar{a}_\alpha) : \alpha < \lambda\}$. Let h be the unique homomorphism from \mathbb{B}_M^- into \mathbb{B}_λ^c mapping $\varphi(M, \bar{a}_\alpha)$ to d_α ; exists as $\langle \varphi(M, \bar{a}_\alpha) : \alpha < \lambda \rangle$ is an independent family of subsets of M . As \mathbb{B}_λ^c is a complete Boolean Algebra we can find an extension of h to a homomorphism from \mathbb{B}_M into (hence onto) \mathbb{B}_λ^c . \square

Moved from end of §1, pg.9:

Discussion: We may prefer to have an invariant which speaks only on one model.

Definition 3.4. Let $(\lambda$ be regular $\geq |T|$, $\varphi(x, \bar{y})$ as in xxx).

1) If $N \in \text{EC}_T(\lambda)$, $f : |N| \rightarrow \lambda$ and $\delta < \lambda$ and $\mathcal{P} \subseteq \mathcal{P}(\delta)$ we say (N, f, δ, κ) accept \mathcal{P} when: there is a witness $\langle a_{\eta, \varepsilon} : \eta \in \mathcal{I}, \varepsilon < \delta \rangle$ such that

- (a) $\mathcal{I} \in \mathbf{T}_\kappa$ where $\mathcal{I} \in \mathbf{T}_\kappa$ iff $\mathcal{I} \subseteq {}^{\omega > \kappa}$ is non-empty, closed under initial segments, with no infinite branch
- (b) for every $\bar{b} \in {}^{\ell g(\bar{y})}N$ for some $\mathcal{U} \in \mathcal{P}$ for almost every $\eta \in \mathcal{I}$ we have $M \models \varphi[a_{\eta, \varepsilon}, \bar{b}]^{\text{if } (\varepsilon \in \mathcal{U})}$.

FILL

Moved from Definition 1.2, pg.3: 3) If N is a model of T with universe λ let

$$\text{inv}_3(M) = \langle \text{inv}_2(M, N) : M \prec N \text{ has universe an ordinal } < \lambda \rangle$$

$$\text{inv}_4(M) = \text{inv}_3(M) / \mathcal{D}_\lambda$$

recalling \mathcal{D}_λ the club filter.

4) If N is a model of cardinality λ let $\text{inv}_4(N)$ be $\text{inv}_4(N')$ for any equivalent some model N' with universe λ isomorphic to λ .

We first deal with the explicit case (and use simplifying) cardinal arithmetic assumptions⁵. Moved from pgs.10,11:

Remark 3.5. Assume

- (a) λ is strongly inaccessible of cofinality κ
- (b) $S \subseteq \{\mu < \lambda : \mu = \mu^{\aleph_0} \text{ is strong limit}\}$ is stationary.

Then we can find $\bar{\mathbb{D}}$ such that

- (α) $D = \langle \mathbb{D}_B : B \subseteq \lambda \rangle$
- (β) \mathbb{D}_B is an ultrafilter of \mathbb{B}_λ^c
- (γ)₁ if $B_\varepsilon^\ell \subseteq \lambda$ for $\varepsilon < \lambda, \ell = 1, 2$ and $B_{\varepsilon_1}^1 \neq B_{\varepsilon_2}^2$ for $\varepsilon_1, \varepsilon_2 < \lambda$ then for a club of $\mu \in S$ we have:
 $\{\varepsilon < \mu : \{Y \subseteq \mu : \text{for some } e \in \mathbb{B}^\ell \text{ we have } (\forall \varepsilon < \mu)(\varepsilon \in Y \equiv e \in \mathbb{D}_{B_\varepsilon}^1)\}\} \neq \{y \subseteq \mu : \text{for some } e \in \mathbb{B}^c \text{ we have } (\forall \varepsilon < \mu)(\varepsilon \in y \equiv e \in \mathbb{D}_{B_\varepsilon}^2)\}$
- (γ)₂ if $\mathcal{P}_{\mu, \alpha} \subseteq \mathcal{P}(\mu)$ for $\alpha < 2^\mu$ then for some $\bar{\mathbb{D}} = \langle \mathbb{D}_\varepsilon : \varepsilon < \lambda \rangle$, for every $\mu \in S$ we have $\{\{\varepsilon < \mu : e \in \mathbb{D}_\varepsilon\} : e \in \mathbb{B}_\lambda^c\} \notin \{\mathcal{P}_{\mu, \alpha} : \alpha < 2^\mu\}$.

Proof. Choose \mathbb{D}_* such that

- \otimes_1 \mathbb{D}_* is an ultrafilter on \mathbb{B}_λ^c disjoint to $\{e_\alpha : \alpha < \lambda\}$.

Let $\bar{\pi}$ be such that

- \otimes_2 [??] (a) $\bar{\pi} = \langle \pi_{\mu_1, \mu_2} : \mu_1 < \mu_2 \text{ are from } S \rangle$
- (b) π_{μ_1, μ_2} is a mapping from $[\mu_2, 2^{\mu_2})$ to $[\mu_1, 2^{\mu_1})$
- (c) the π 's commute
- (d) if $(S \cap \mu)$ has not last inverse limit.

For each $\mu \in S$ we choose $\bar{\Lambda}_\mu$ such that

- \otimes_2' ?? (a) $\bar{\Lambda}_\mu = \langle \Lambda_{\mu, \alpha} : \alpha \in [\mu, 2^\mu) \rangle$
- (b) $\Lambda_{\mu, \alpha} \subseteq \mu^2$
- (c) if $\mu_1 < \mu_2$ are from S and $\pi_{\mu_1, \mu_2}(\alpha_2) = \alpha_1$ then $\Lambda_{\mu_1, \alpha_1} \subseteq \{\nu \upharpoonright \mu_1 : \nu \in \Lambda_{\mu_2, \alpha_2}\}$
- \otimes_2' let $\theta, S_*, E, S, \bar{c}$ be as in the proof of 1.9
- \otimes_3 if $\bar{\eta} = \langle \eta_\varepsilon : \varepsilon < \mu \rangle, \eta_\varepsilon \in {}^\lambda 2$ for $\varepsilon < \mu$ then let $\mathcal{G}_{\mu, \bar{\eta}}$ be the \aleph_1 -complete filter on μ generated by $\{\varepsilon < \mu : \eta_\varepsilon^{-1}\{0\} \supseteq u\} : u \subseteq \mu \text{ is countable}\}$.

Now we can choose $\bar{\eta}^*$ such that

- \boxplus_5 (a) $\bar{\eta}^* = \langle \eta_A^* : A \subseteq \lambda \rangle$
- (b) $\eta_A^* \in \mu^2$
- (c) if $\mu = \mu^{\aleph_0} < \lambda$ and $A_n \subseteq \lambda, \langle A_n \cap \mu : n < \omega \rangle$ are pairwise distinct then $\eta_{A_0}^{-1}\{1\} \not\subseteq \cup\{\eta_{1+n}^{-1}\{1\} : n < \omega\}$.

Hence

⁵if $2^\lambda > \lambda^+$ use the game formulation

⊗ if $A_\varepsilon \subseteq \lambda$ for $\varepsilon < \lambda$ and $B \in \mathcal{P}(\lambda) \setminus \{A_\varepsilon : \varepsilon < \lambda\}$ then $\eta_B^{-1}\{1\} \neq \emptyset, \lambda \bmod D_{\langle \eta_{A_\varepsilon} : \varepsilon < \lambda \rangle}$.

So $\langle \mathbb{D}_{\eta_B} : B \subseteq \lambda \rangle$ are as required in (γ) . □

Proof. Proof of $(\gamma)_2$ The proof is similar. FILL.

We choose $\bar{\Lambda}_\mu$ by induction on μ such that ... □

REFERENCES

- [1] John T. Baldwin and Saharon Shelah. Second-order quantifiers and the complexity of theories. *Notre Dame Journal of Formal Logic*, 26:229–303, 1985. Proceedings of the 1980/1 Jerusalem Model Theory year.
- [2] Ryszard Engelking and Monika Karłowicz. Some theorems of set theory and their topological consequences. *Fundamenta Math.*, 57:275–285, 1965.
- [3] Itay Kaplan and Saharon Shelah. Examples in dependent theories. *Journal of Symbolic Logic*, 49:585–619, 2014.
- [4] Saharon Shelah. Dependent dreams: recounting types. preprint. arxiv:1202.5795.
- [5] Saharon Shelah. Dependent T and Existence of limit models. *Tbilisi Mathematical Journal*, submitted. arxiv:math.LO/0609636.
- [6] Saharon Shelah. Remarks on squares. In *Around classification theory of models*, volume 1182 of *Lecture Notes in Mathematics*, pages 276–279. Springer, Berlin, 1986.
- [7] Saharon Shelah. Anti-homogeneous Partitions of a Topological Space. *Scientiae Mathematicae Japonicae*, 59, No. 2; (special issue:e9, 449–501):203–255, 2004. arxiv:math.LO/9906025.
- [8] Saharon Shelah. Dependent theories and the generic pair conjecture. *Communications in Contemporary Mathematics*, 17:1550004 (64 pps.), 2015. arxiv:math.LO/0702292.

EINSTEIN INSTITUTE OF MATHEMATICS, EDMOND J. SAFRA CAMPUS, GIVAT RAM, THE HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM, 91904, ISRAEL, AND, DEPARTMENT OF MATHEMATICS, HILL CENTER - BUSCH CAMPUS, RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY, 110 FRELINGHUYSEN ROAD, PISCATAWAY, NJ 08854-8019 USA

E-mail address: shelah@math.huji.ac.il

URL: <http://shelah.logic.at>