

NICE \aleph_1 GENERATED NON- P -POINTS, I

SAHARON SHELAH

ABSTRACT. We define a family of non-principal ultrafilters on \mathbb{N} which are, in a sense, very far from P -points. We prove the existence of such ultrafilters under reasonable conditions. In subsequent articles we shall prove that such ultrafilters may exist while no P -point exists. Though our primary motivations came from forcing and independence results, the family of ultrafilters introduced here should be interesting from combinatorial point of view too.

1. INTRODUCTION

One of important notions in general topology and set theory of the reals is that of a P -point. Recall that a P -point is a non-principal ultrafilter D on \mathbb{N} with the property that for any countable family $\mathcal{A} \subseteq D$ there is a $B \in D$ almost (modulo finite) included in all $A \in \mathcal{A}$ (see Definition 3.6(2)). Concerning these and other special ultrafilters on \mathbb{N} , their history and basic applications we refer the reader to the survey article by Blass [1].

In many applications it is important to preserve P -points by specific forcing notions and by a forcing iterated with countable supports. Recall that *preservation of an ultrafilter* means that the ultrafilter from the ground model \mathbf{V} generates an ultrafilter in the generic extension $\mathbf{V}[G]$ (see [2, Chapter VI]). We have a very good understanding of these questions and many relevant results have been presented in the literature. From our point of view the P -points are tractable for independence results because:

- ⊞₁ (A) there are quite many forcing notions preserving P -points,
- (B) a proper forcing notion \mathbb{Q} which preserves “ D is an ultrafilter” preserves its being a P -point,
- (C) the preservation of P -points is preserved in limits of CS iterations (together these give a well controlled way to have ultrafilters generated by $\aleph_1 < 2^{\aleph_0}$ sets).
- (D) We can destroy a P -point by forcing, i.e., ensure it has no extension to a P -point (and consequently we may prove the consistency of “there are no P -points”),
- (E) moreover, we can “split hairs”, i.e., destroy some P -points while preserving other, so we can have unique P -point up to isomorphisms.

For more details we refer the reader to [2, Ch.VI and Ch.XVIII,§4].

We may wonder if the theory developed for P -points can be repeated for other ultrafilters. We may ask:

1991 *Mathematics Subject Classification*. MSC 2010: Primary 03E05,54A24; Secondary: 03E35, 03E17, 54A35.

Key words and phrases. set theory, general topology, ultrafilters, P -point, forcing.

- Question 1.1.* (1) Are there other types of ultrafilters preserved by CS iterations of suitable forcing notions?
- (2) In particular, we are interested in preservation of our ultrafilters at limit stages of CS iterations: for a limit ordinal δ , having been preserved by \mathbb{P}_α for $\alpha < \delta$, does this hold for \mathbb{P}_δ when $\langle \mathbb{P}_\alpha, \mathbb{Q}_\beta : \alpha \leq \delta, \beta < \delta \rangle$ is a CS iteration of proper forcing notions?

We suggested this problem in [3, 3.13] and we speculated about it there. Note that ultrafilters as in Question 1.1 for CS iterations are naturally generated by \aleph_1 sets; moreover CS iterations are mainly interesting when we start with CH, and “preserve an ultrafilter” is meaningful only when we add reals, naturally \aleph_2 ones. We suspect this direction is related to the question on the existence of a point of van Dowen [4] (see $\boxplus_2(A)$ below), but at present we do not know neither if they are related nor how to answer it. Other specific problems that we have in mind when developing the theory for Question 1.1 are a problem of Nyikos and a problem of Dow:

- \boxplus_2 (A) [E. van Dowen] Is it consistent that: there is no ultrafilter D on \mathbb{Q} such that every $A \in D$ contains a member of D which is a closed set with no isolated points?
- (B) [P. Nyikos] Is it consistent to have some ultrafilter $D \in \beta^*(\mathbb{N}) \setminus \mathbb{N}$ of character \aleph_1 , but no P -point?
- (C) [A. Dow] Is it consistent to have $\mathfrak{u} = \aleph_1$, there is a P -point D , but no P -point D with $\chi(D) = \aleph_1$?

In the series of papers started here the main points are:

- \boxplus_3 (A) we have an involved family of sets (really well founded trees) appearing in the definition,
- (B) each ultrafilter has no P -point as a quotient,
- (C) they are related to a game,
- (D) such systems exists assuming, e.g., \diamond_{\aleph_1} ,
- (E) enough relevant forcing notions preserve such systems, in particular, some serving $\boxplus_1(C)$, so answering Question 1.1(1),
- (F) we have a preservation theorem for such ultrafilters under CS iterations.
- (G) As an application, we will solve Nyikos’ problem $\boxplus_2(B)$.

So problems 1.1 and $\boxplus_2(B)$ will be resolved by the methods we start developing here, but presently not $\boxplus_2(A,C)$ (a problem of van Dowen and a problem of Dow).

In the present article we define ultrafilters analogous to P -points but with no P -point as a quotient; this is done in Sections 2 and 3. In the fourth section we deal with basic connections to forcing that we will use in the independence results in subsequent papers.

In the second paper of the series (presently the first half of [5]) we present these ultrafilters in a more general framework and deal with sufficient conditions for such an ultrafilter to generate an ultrafilter in a suitable generic extension. For the limit case we continue the proof of preservation theorems in [2], in particular [2, Ch.VI,1.26,1.27] and Case A with transitivity of [2, Ch.XVIII,§3]. For the successor case we need that the relevant forcing preserves our ultrafilters. We will conclude with the proof for $\text{CON}(\mathfrak{u} = \aleph_1 + \text{no } P\text{-points})$.

In the third part (currently the second half of [5]) we note that the ultrafilters so far were really analogous to selective (i.e., Ramsey) ultrafilters and give a more general framework which also includes P -points.

Remark 1.2. (1) There may be P -point while $\mathfrak{d} > \aleph_1$, see Blass and Shelah [6] and references there, but the existence of ultrafilters in the direction here, far from P -point, implies $\mathfrak{d} = \aleph_1$, see the survey of Blass [1]. But note that the ultrafilter may be \aleph_1 -generated in a different sense: union of \aleph_1 families of the form $\text{fil}(B) \cap \mathcal{P}(\max(B))$.

- (2) Note that it may be harder (than in the P -point case) to build such ultrafilters as here which are μ -generated instead of \aleph_1 -generated because of the unbounded countable depth involved. We have not looked at this as well as at the natural variants of our definition (not to speak of generalization to reasonable ultrafilters, see [7] and Roslanowski and Shelah [8, 9]).

2. SYSTEM OF FILTERS USING WELL FOUNDED TREES

Notation 2.1. Here, $M = (M, <_M)$ is a partial order and B is a subset of M inheriting its order.

- (1) For $\eta \in B$ we let $B_{\geq \eta} = \{\nu \in B : \eta \leq_M \nu\}$ and similarly $B_{> \eta}$. We also define

$$\text{suc}_B(\eta) = \{\nu \in B : \eta <_M \nu \text{ but for no } \rho \in B \text{ do we have } \eta <_M \rho <_M \nu\}$$

$$\text{and } \max(B) = \{\nu \in B : B \cap M_{> \nu} = \emptyset\}.$$

- (2) We say that Y is a *front* of $B \subseteq M$ when $Y \subseteq B$ and every branch (maximal chain) of B meets Y and the members of Y are pairwise $<_M$ -incomparable.

Definition 2.2. Let $M = (M, <_M)$ be a partial order.

- (1) A set $B \subseteq M$ is a *CWT sub-tree* of M if the following conditions (a)–(f) are satisfied (CWT stands for *countable well-founded*).
- (a) B is a countable subset of M .
 - (b) B has a $<_M$ -minimal member called its root, $\text{rt}(B)$.
 - (c) B , i.e., $(B, <_M \upharpoonright B)$ is a tree with $\leq \omega$ levels and no ω -branch (so all chains in B are finite).
 - (d) For each $\nu \in B$ the set $\text{suc}_B(\nu)$ is either empty or infinite.
 - (e) If $\eta, \nu \in B$ are $<_M$ -incomparable, then they have no common \leq_M -upper bound (i.e., they are incompatible, $\eta \parallel_M \nu$).
 - (f) If $\nu \in B \setminus \max(B)$ and $F \subseteq M \setminus M_{\leq \nu}$ is finite, then for infinitely many $\varrho \in \text{suc}_B(\nu)$ we have $(\forall \rho \in F)(\rho \parallel_M \varrho)$.

The family of all CWT sub-trees of M is denoted by $\text{CWT}(M)$.

- (2) For $B \in \text{CWT}(M)$ let $\text{frt}(B)$ be the set of all fronts of B , which in this case means the family of all maximal sets of pairwise incomparable members of B .
- (2A) For antichains Y_1, Y_2 of M we say that Y_2 is above Y_1 when

$$(\forall \eta \in Y_2)(\exists \nu \in Y_1)[\nu \leq_M \eta]$$

and this will be used mainly for $Y_1, Y_2 \in \text{frt}(B)$, $B \in \text{CWT}(M)$.

- (2B) For Y_1, Y_2 as above let the projection h_{Y_1, Y_2} be the unique function $h : Y_2 \rightarrow Y_1$ such that $h(\eta) \leq_M \eta$ for $\eta \in Y_2$.

- (2C) If $Y_1, Y_2 \in \text{frt}(B)$ then Y_2 is almost above Y_1 when for some $B' \in \text{sb}(B)$, see (3) below, $B' \cap Y_2$ is above $B' \cap Y_1$. We also define the projection h_{Y_1, Y_2} as above, but its domain is not Y_2 but the set $\{\eta \in Y_2 : (\exists \nu \in Y_1)(\nu \leq_M \eta)\}$.
- (2D) The default value of $Y \in \text{frt}(B)$ is $\max(B) = \{\nu \in B : \nu \text{ is } <_M\text{-maximal in } B\}$.
- (3) Let $B \in \text{CWT}(M)$. We say that B' is an *exhaustive subtree* of B when:
- $B' \in \text{CWT}(M)$, $B' \subseteq B$, and
 - $\text{rt}(B') = \text{rt}(B)$, and
 - if $\nu \in B'$ then $\text{suc}_{B'}(\nu) \subseteq \text{suc}_B(\nu)$ and $\text{suc}_B(\nu) \setminus \text{suc}_{B'}(\nu)$ is finite.
- We let $\text{sb}(B)$ be the set of all exhaustive subtrees B' of B .
- (3A) For $B \in \text{CWT}(M)$ and $Y \in \text{frt}(B)$ let $E_{B, Y}$ be the filter on Y generated by the family
- $$\{Y \cap B' : B' \text{ is an exhaustive subtree of } B, \text{ i.e., } B' \in \text{sb}(B)\}.$$
- (3B) For B as above and $Y \subseteq B$ let $B[\leq Y] = \{\nu \in B : (\exists \eta \in Y)(\nu \leq_M \eta)\}$.
- (3C) In part (3), we say f witnesses “ $B' \in \text{sb}(B)$ ” if $f : B' \setminus \max(B) \rightarrow [B]^{<\aleph_0}$ satisfies

$$\nu \in B' \setminus \max(B) \Rightarrow \text{suc}_B(\nu) \setminus \text{suc}_{B'}(\nu) \subseteq f(\nu).$$

Note that for f being a witness only $f|B'$ matters; in fact only $f|B'[\leq Y]$ matters when we are interested in $D_{B, Y}$.

- (3D) For $B \in \text{CWT}(M)$ let $\text{psb}_M(B)$ (“p” stands for positive) be the set of positive subtrees B' of B which means (a),(b) as in part (3) above and (c)' if $\nu \in B' \setminus \max(B)$, then $\text{suc}_{B'}(\nu)$ is an infinite subset of $\text{suc}_B(\nu)$.
- (4) An antichain $Y \subseteq M$ is an *almost front of B* if for some $B' \in \text{sb}(B)$ the intersection $Y \cap B'$ is a front of B' . Let $\text{alm-frt}(B) = \text{alm-frt}_M(B)$ denote the set of all almost fronts of B .
- (4A) For $Y \in \text{alm-frt}_M(B)$ let
- $$\text{fil}_M(Y, B) = \{X \subseteq Y : \text{for some } B' \in \text{sb}(B) \text{ we have } X \supseteq B' \cap Y\}.$$
- (5) Let \leq_M^* be the following two-place relation (actually a partial order) on $\text{CWT}(M)$:
- $$B_1 \leq_M^* B_2 \text{ iff } (B_1, B_2 \in \text{CWT}(M) \text{ and } \text{rt}(B_1) = \text{rt}(B_2) \text{ and for some } B'_2 \in \text{sb}(B_2), \text{ we have}$$
- $B'_2 \cap B_1 \in \text{psb}_M(B_1)$, and
 - every almost front of $B'_2 \cap B_1$ is an almost front of B_2 .
- The tree B'_2 as above will be called *a witness for $B_1 \leq_M^* B_2$* .
- (6) For $B \in \text{CWT}(M)$, *the depth of B* is defined recursively by

$$\text{Dp}(B) = \sup\{\text{Dp}(B_{\geq \eta}) + 1 : \eta \in B \setminus \{\text{rt}(B)\}\}.$$

Remark 2.3. If $B, B' \in \text{CWT}(M)$, $B' \subseteq B$ and $\nu \in B'$, then $\text{suc}_B(\nu) \cap B' \subseteq \text{suc}_{B'}(\nu)$, but the two sets do not have to be equal. Note that in the definitions of both $B' \in \text{sb}(B)$ and $B' \in \text{psb}_M(B)$ we do require that

$$(*) \quad (\forall \nu \in B') (\text{suc}_B(\nu) \cap B' = \text{suc}_{B'}(\nu)).$$

The condition (*) implies that if $Y \subseteq B$ is a front of B , then $Y \cap B'$ is a front of B' .

Observation 2.4. Let M be a partial order and $B, B_1, B_2 \in \text{CWT}(M)$.

- We have that $B_1 \leq_M^* B_2$ if and only if every almost front of B_1 is an almost front of B_2 .

- (2) The relation \leq_M^* is a partial order on $\text{CWT}(M)$.
- (3) If $B_2 \in \text{psb}_M(B_1)$, then $B_1 \leq_M^* B_2$ and $\text{psb}_M(B_2) \subseteq \text{psb}_M(B_1)$.
- (4) If $B_2 \in \text{sb}(B_1)$, then $B_2 \in \text{psb}(B_1)$, $\text{sb}(B_2) \subseteq \text{sb}(B_1)$ and $B_1 \leq_M^* B_2 \leq_M^* B_1$.
- (5) For $B \in \text{CWT}(M)$, $\max(B)$ is a front of B and also $\{\text{rt}(B)\}$ is. If $B \neq \{\text{rt}(B)\}$, then $\text{suc}_B(\text{rt}(B))$ is a front of B .
- (6) Every front of $B \in \text{CWT}(M)$ is an almost front of B .
- (7) If $B \in \text{CWT}(M)$ then $\text{Dp}(B)$ is a countable ordinal and $B_{\geq \eta} \in \text{CWT}(M)$ for all $\eta \in B$.
- (8) If $Y \subseteq B \setminus \{\text{rt}(B)\}$ is a front of B , and $\eta \in \text{suc}_B(\text{rt}(B))$, then $Y \cap B_{\geq \eta}$ is a front of $B_{\geq \eta}$.
- (9) If Y is an almost front of B and an antichain Z is an almost front of $B_{\geq \eta}$ for every $\eta \in Y \cap B$, then Z is an almost front of B .
- (10) If $B_1 \leq_M^* B_2$ and Y is a front of B_1 , then there is $B'_2 \in \text{sb}(B_2)$ such that $Y \cap B'_2$ is a front of B'_2 and $(B_1)_{\geq \eta} \leq_M^* (B'_2)_{\geq \eta}$ for all $\eta \in Y \cap B'_2$.

Proof. Straightforward. \square

Definition 2.5. Let \mathbf{K} be the class of the objects $\mathbf{x} = \langle M_{\mathbf{x}}, <_{M_{\mathbf{x}}}, \overline{\mathcal{A}}_{\mathbf{x}}, \mathcal{A}_{\mathbf{x}}, \mathcal{B}_{\mathbf{x}}, \leq_{\mathbf{x}} \rangle$ satisfying the following properties (a)–(h).

- (a) $(M_{\mathbf{x}}, <_{M_{\mathbf{x}}}) = (M, <)$ is a partial order with the smallest element $\text{rt}_{\mathbf{x}} = \text{rt}(\mathbf{x})$. Let $M_{\mathbf{x}}^- = M_{\mathbf{x}} \setminus \{\text{rt}_{\mathbf{x}}\}$,
- (b) $\overline{\mathcal{A}}_{\mathbf{x}} = \overline{\mathcal{A}} = \langle \mathcal{A}_{\eta} : \eta \in M \rangle = \langle \mathcal{A}_{\eta}^{\mathbf{x}} : \eta \in M_{\mathbf{x}} \rangle$ and $\mathcal{A}_{\mathbf{x}} = \bigcup \{ \mathcal{A}_{\eta} : \eta \in M_{\mathbf{x}}^- \}$,
- (c) $\mathcal{A}_{\eta} \subseteq \text{CWT}(M)$, let $\mathcal{A}_{\eta}^- = \mathcal{A}_{\eta} \setminus \{ \{ \eta \} \}$,
- (d) $\text{rt}(B) = \eta$ for every $B \in \mathcal{A}_{\eta}$,
- (e) \mathcal{A}_{η} is not empty, in fact $\{ \eta \} \in \mathcal{A}_{\eta}$,
- (f) $\mathcal{B}_{\mathbf{x}} = \mathcal{A}_{\text{rt}_{\mathbf{x}}}^{\mathbf{x}} \setminus \{ \{ \text{rt}_{\mathbf{x}} \} \}$ and $\leq_{\mathbf{x}}$ is a directed partial order on $\mathcal{B}_{\mathbf{x}}$,
- (g) $B_1 \leq_{\mathbf{x}} B_2$ implies $B_1 \leq_M^* B_2$, see Definition 2.2(5) and, of course, $B_1, B_2 \in \mathcal{B}_{\mathbf{x}}$,
- (h) if $\nu \in B \in \mathcal{A}_{\eta}$ then $B \cap M_{\geq \nu} \in \mathcal{A}_{\nu}$.

When dealing with $M_{\mathbf{x}}, \overline{\mathcal{A}}_{\mathbf{x}}$ etc we may omit \mathbf{x} when clear from the context.

Definition 2.6. Let $\mathbf{x} \in \mathbf{K}$ and $\eta \in M_{\mathbf{x}}$.

- (1) Let $\text{frt}(\eta) = \text{frt}_{\mathbf{x}}(\eta) = \{ Y : Y \text{ is a front of } B \text{ for some } B \in \mathcal{A}_{\eta}^{\mathbf{x}} \}$ and instead of $\text{frt}(B)$ (see Definition 2.2(2)) we may write also $\text{frt}_{\mathbf{x}}(B)$. We let

$$\text{frt}^-(\eta) = \{ Y \in \text{frt}(\eta) : Y \neq \{ \eta \} \}.$$

Omitting η means $\eta = \text{rt}_{\mathbf{x}}$.

- (2) Similarly, using Definition 2.2(4), we define $\text{alm-frt}_{\mathbf{x}}(\eta)$ (and $\text{alm-frt}_{\mathbf{x}}$).
- (3) Let $B \in \mathcal{A}_{\eta}^{\mathbf{x}}$. We define

$$\text{Fin}(B) = \{ f : f \text{ is a function with domain } B \setminus \max(B) \text{ and such that } f(\nu) \in [\text{suc}_B(\nu)]^{< \aleph_0} \text{ for all } \nu \in B \setminus \max(B) \},$$

and for $f \in \text{Fin}(B)$ we set

$$A_f = A_{B,f} = \{ \eta \in B : (\forall \rho \in B \setminus \max(B)) (\forall \varrho \in \text{suc}_B(\rho)) (\varrho \leq_M \eta \Rightarrow \varrho \notin f(\rho)) \}.$$

(Recall Definition 2.2(3C).)

- (4) Assume that $Y \in \text{alm-frt}_{\mathbf{x}}$. We let $D_Y = D_Y^{\mathbf{x}}$ be the family

$$\{ Z \subseteq Y : \text{for some } B \in \mathcal{B}_{\mathbf{x}} \text{ and } B' \in \text{sb}(B) \text{ we have } Y \in \text{alm-frt}(B) \text{ and } B' \cap Y \subseteq Z \}.$$

- (5) If $B \in \mathcal{B}_x$, then $D_x(B) = D_{\max(B)}^x$.
- (6) We let $\text{Dp}_x(\eta) = \sup\{\text{Dp}(B) + 1 : B \in \mathcal{A}_\eta^x\}$ (recall Definition 2.2(6)).

If x is clear from the context, then we may omit the subscript/superscript x in the objects defined above.

Let us recall the definition of the Rudin—Keisler order on ultrafilters.

Definition 2.7. Let D_ℓ be an ultrafilter on \mathcal{U}_ℓ for $\ell = 1, 2$. We say $D_1 \leq_{\text{RK}} D_2$ iff there is a function h such that

- $\text{Dom}(h) \in D_2$, $\text{Rang}(h) \in D_1$ and
- $A \in D_1 \Leftrightarrow \{a \in \text{Dom}(h) : h(a) \in A\} \in D_2$ for every $A \subseteq \mathcal{U}_1$.

Observation 2.8. Assume $x \in \mathbf{K}$ and let $B, B_1, B_2 \in \mathcal{B}_x$.

- (1) $\{\text{rt}_x\} \in \text{frt}_x$ and $D_{\{\text{rt}_x\}}^x = \{\{\text{rt}_x\}\}$.
- (2) If $B_1 \leq_x B_2$, $f \in \text{Fin}(B_1)$ and $Y \in \text{alm-frt}(B_1)$, then $Y \in \text{alm-frt}(B_2)$ and there is $g \in \text{Fin}(B_2)$ such that $Y \cap A_{B_2, g} \subseteq Y \cap A_{B_1, f}$.
- (3) If $Y \in \text{alm-frt}(B_\ell)$, $f_\ell \in \text{Fin}(B_\ell)$ (for $\ell = 1, 2$), then there are $B^* \in \mathcal{B}_x$ and $g \in \text{Fin}(B^*)$ such that $B_1 \leq_x B^*$, $B_2 \leq_x B^*$ and

$$Y \cap A_{B^*, g} \subseteq Y \cap A_{B_1, f_1} \cap A_{B_2, f_2}.$$

- (4) If $Y \in \text{alm-frt}_x$, then D_Y^x is a filter on Y .
- (5) If $B_1 \leq_x B_2$, $Y_1 \in \text{alm-frt}(B_1)$, and $Y_2 = Y_1 \cap B_2$ (hence $Y_2 \in \text{alm-frt}(B_2)$), then $Y_2 \in D_{Y_1}^x$ and $D_{Y_2}^x = D_{Y_1}^x \upharpoonright Y_2$.
- (6) Assume that $Y_1, Y_2 \in \text{frt}(B)$ and Y_2 is above Y_1 . Let $h : Y_2 \xrightarrow{\text{onto}} Y_1$ be the projection, i.e.,

$$h(\nu_2) = \nu_1 \Leftrightarrow \nu_1 \in Y_1 \wedge \nu_2 \in Y_2 \wedge \nu_1 \leq_{M_x} \nu_2.$$

Then $h(D_{Y_2}) = D_{Y_1}$, i.e., $D_{Y_1} = \{A \subseteq Y_1 : h^{-1}[A] \in D_{Y_2}\}$ (so h witnesses $D_{Y_1} \leq_{\text{RK}} D_{Y_2}$).

- (7) If $B_1 \leq_x B_2$ and $Y_\ell = \text{suc}_{B_\ell}(\text{rt}_x)$ for $\ell = 1, 2$, then:
 - (a) Y_ℓ is a front of B_ℓ and Y_1 almost above Y_2 , see Definition 2.2(2C),
 - (b) if Y is a front of B_ℓ and it is not $\{\text{rt}_x\}$, then Y is above Y_ℓ .
- (8) $\max(B)$ is the maximal front of B which means that it is above any other.
- (9) If \mathbb{Q} is an ${}^\omega\omega$ -bounding forcing and $B \in \mathcal{B}_x$, then for any $B' \in \text{sb}(B)^{\mathbf{V}[\mathbb{Q}]}$ there is $B'' \in (\text{sb}(B))^{\mathbf{V}}$ such that $B'' \subseteq B'$.
- (10) If F is a finite subset of M_x^- , $B \in \mathcal{B}_x$, then there is a branch (i.e., a maximal chain) $C \subseteq B$ such that

$$(\forall \rho \in F)(\forall \sigma \in C)(\rho \not\leq_M \sigma).$$

- (11) If $B \in \mathcal{A}_\eta$ and $\nu \in B \setminus \max(B)$, then $\text{id}_x(\nu, B)$ is a proper ideal on $\text{suc}_B(\nu)$.

Proof. Straightforward. □

Definition 2.9. (1) For an (infinite) cardinal κ let $\mathbf{K}_{<\kappa}$ be the class of $x \in \mathbf{K}$ such that $\|x\| := |M_x| + \sum\{|\mathcal{A}_\eta^x| : \eta \in M_x\} < \kappa$, similarly $\mathbf{K}_{\leq\kappa}$.

- (2) $\leq_{\mathbf{K}}$ is the following two-place relation on \mathbf{K} (it is a partial order, see Observation 2.10 below): $x \leq_{\mathbf{K}} y$ if and only if

(a) $M_{\mathbf{x}} \subseteq M_{\mathbf{y}}$ (as partial orders) and, moreover, for any $\eta, \nu \in M_{\mathbf{x}}$ we have

$$\nu \parallel_{M_{\mathbf{x}}} \eta \quad \text{if and only if} \quad \nu \parallel_{M_{\mathbf{y}}} \eta,$$

and

- (b) $\eta \in M_{\mathbf{x}} \Rightarrow \mathcal{A}_{\eta}^{\mathbf{x}} \subseteq \mathcal{A}_{\eta}^{\mathbf{y}}$, and
(c) $\text{rt}_{\mathbf{y}} = \text{rt}_{\mathbf{x}}$ (actually follows from (d)), and
(d) $\leq_{\mathbf{x}} = \leq_{\mathbf{y}} \upharpoonright \mathcal{B}_{\mathbf{x}}$.

(3) If $\langle \mathbf{x}_{\alpha} : \alpha < \delta \rangle$ is a $\leq_{\mathbf{K}}$ -increasing sequence we define $\mathbf{x}_{\delta} = \bigcup \{\mathbf{x}_{\alpha} : \alpha < \delta\}$, the union of the sequence, by $M_{\mathbf{x}_{\delta}} = \bigcup \{M_{\mathbf{x}_{\alpha}} : \alpha < \delta\}$ as partial orders and $\mathcal{A}_{\eta}^{\mathbf{x}_{\delta}} = \bigcup \{\mathcal{A}_{\eta}^{\mathbf{x}_{\alpha}} : \alpha < \delta \text{ satisfies } \eta \in M_{\mathbf{x}_{\alpha}}\}$ and $\leq_{\mathbf{x}_{\delta}} = \bigcup \{\leq_{\mathbf{x}_{\alpha}} : \alpha < \delta\}$.

Observation 2.10. (1) $\leq_{\mathbf{K}}$ is really a partial order.

(2) If $\langle \mathbf{x}_{\alpha} : \alpha < \delta \rangle$ is $\leq_{\mathbf{K}}$ -increasing then \mathbf{x}_{δ} , the union of the sequence, is a $\leq_{\mathbf{K}}$ -lub of the sequence and $\|\mathbf{x}_{\delta}\| \leq \sum \|\{\mathbf{x}_{\alpha} : \alpha < \delta\}\|$.

Definition 2.11. Let $\mathbf{x} \in \mathbf{K}$. We say that \mathbf{x} is:

fat when: if $B \in \mathcal{B}_{\mathbf{x}}$ and $B' \in \text{sb}(B)$, then there is $B'' \in \text{sb}(B')$ such that $B'' \in \mathcal{B}_{\mathbf{x}}$ and $B \leq_{\mathbf{x}} B''$;

big when: if $B \in \mathcal{B}_{\mathbf{x}}$ and $\mathbf{c} : \max(B) \rightarrow \{0, 1\}$, then for some $B' \in \mathcal{B}_{\mathbf{x}}$ we have that

$$B' \in \text{psb}_{M_{\mathbf{x}}}(B) \cap \mathcal{B}_{\mathbf{x}}, \quad B \leq_{\mathbf{x}} B', \quad \text{and} \quad \mathbf{c} \upharpoonright \max(B') \text{ is constant,}$$

large when: if $B \in \mathcal{B}_{\mathbf{x}}$ and \mathbf{c} is a function with domain $\max(B)$, then for some $B' \in \text{psb}_{M_{\mathbf{x}}}(B) \cap \mathcal{B}_{\mathbf{x}}$ and a front Y of B' we have $B \leq_{\mathbf{x}} B'$ and

$$(\forall \eta, \nu \in \max(B')) (\mathbf{c}(\eta) = \mathbf{c}(\nu) \Leftrightarrow (\exists \rho \in Y) (\rho \leq_{M_{\mathbf{x}}} \eta \wedge \rho \leq_{M_{\mathbf{x}}} \nu)),$$

full when: if $B \in \mathcal{A}_{\eta}^{\mathbf{x}}$, $\eta \neq \text{rt}_{\mathbf{x}}$ and $B' \in \text{psb}_{M_{\mathbf{x}}}(B)$, then $B' \in \mathcal{A}_{\eta}^{\mathbf{x}}$.

3. CONSTRUCTION OF ULTRA-SYSTEMS

Lemma 3.1. $\mathbf{K}_{\leq \aleph_0}$ is non-empty.

Proof. Define \mathbf{x} so that $M_{\mathbf{x}} = \{\eta_*\}$, $\mathcal{A}_{\eta_*}^{\mathbf{x}} = \{\{\eta_*\}\}$, $\text{rt}_{\mathbf{x}} = \eta_*$. Now it is easy to check. \square

Lemma 3.2. If $\mathbf{x} \in \mathbf{K}$ and $\eta \in M_{\mathbf{x}}$ satisfies $|\mathcal{A}_{\eta}^{\mathbf{x}}| = 1$, i.e., $\mathcal{A}_{\eta}^{\mathbf{x}} = \{\{\eta\}\}$, then for some $\mathbf{y} \in \mathbf{K}$ we have $\mathbf{x} \leq_{\mathbf{K}} \mathbf{y}$, $|\mathcal{A}_{\eta}^{\mathbf{y}}| > 1$ and $\|\mathbf{y}\| \leq \|\mathbf{x}\| + \aleph_0$.

Proof. Let $\langle \eta_n : n < \omega \rangle$ be pairwise distinct objects not belonging to $M_{\mathbf{x}}$. We define \mathbf{y} by:

- (a) $M_{\mathbf{y}}$ has set of elements $M_{\mathbf{x}} \cup \{\eta_n : n < \omega\}$,
(b) $\nu <_{M_{\mathbf{y}}} \rho$ if and only if $\nu <_{M_{\mathbf{x}}} \rho$ or $\nu \leq_{M_{\mathbf{x}}} \eta \wedge (\exists n)(\rho = \eta_n)$,
(c) $\mathcal{A}_{\nu}^{\mathbf{y}}$ is:
 (α) $\mathcal{A}_{\nu}^{\mathbf{x}}$ when $\nu \in M_{\mathbf{x}} \setminus \{\eta\}$,
 (β) $\{\{\eta\}, \{\eta_n : n < \omega\} \cup \{\eta\}\}$ when $\nu = \eta$,
 (γ) $\{\{\eta_n\}\}$ when $\nu = \eta_n$,
(d) the order $\leq_{\mathbf{y}}$ is $\leq_{\mathbf{x}}$ if $\eta \neq \text{rt}_{\mathbf{x}}$, and it is determined by $\{\eta\} \leq_{\mathbf{y}} \{\eta_n : n < \omega\} \cup \{\eta\}$ if $\eta = \text{rt}_{\mathbf{x}}$.

Now check. \square

Lemma 3.3. (1) If $\mathbf{x} \in \mathbf{K}_{\leq \aleph_0}$ then for some $\mathbf{y} \in \mathbf{K}_{\leq \aleph_0}$ we have $\mathbf{x} \leq_{\mathbf{K}} \mathbf{y}$ and in $\mathcal{B}_{\mathbf{y}}$ there is a $\leq_{\mathbf{y}}$ -maximal member.

- (2) If $\mathbf{x} \in \mathbf{K}_{\leq \aleph_0}$ and some $B \in \mathcal{B}_{\mathbf{x}}$ is $\leq_{\mathbf{x}}$ -maximal then for some $\mathbf{y} \in \mathbf{K}_{\leq \aleph_0}$ and $B' \in \mathcal{B}_{\mathbf{y}}$ we have $\mathbf{x} \leq_{\mathbf{K}} \mathbf{y}$ and $B <_{\mathbf{y}} B'$.
- (3) If $\mathbf{x} \in \mathbf{K}_{\leq \aleph_0}$, $\eta \in M_{\mathbf{x}}$, $B_1 \in \mathcal{A}_{\eta}^{\mathbf{x}}$, $B_2 \in \text{psb}_{M_{\mathbf{x}}}(B_1)$ and
- $$\eta = \text{rt}_{\mathbf{x}} \Rightarrow B_1 \text{ is } \leq_{\mathbf{x}}\text{-maximal,}$$
- then there is $\mathbf{y} \in \mathbf{K}_{\leq \aleph_0}$ such that $\mathbf{x} \leq_{\mathbf{K}} \mathbf{y}$ and $B_2 \in \mathcal{A}_{\eta}^{\mathbf{y}}$.
- (4) If $\mathbf{x} \in \mathbf{K}_{\leq \aleph_0}$, $B_1 \in \mathcal{B}_{\mathbf{x}}$ and $B_2 \in \text{sb}(B_1)$, then there is $\mathbf{y} \in \mathbf{K}_{\leq \aleph_0}$ such that $\mathbf{x} \leq_{\mathbf{K}} \mathbf{y}$ and $B_2 \in \mathcal{B}_{\mathbf{y}}$.

Proof. 1) If in $(\mathcal{B}_{\mathbf{x}}, \leq_{\mathbf{x}})$ there is a maximal member then we let $\mathbf{y} = \mathbf{x}$. Otherwise, as it is directed (see clause (f) of Definition 2.5) and $\|\mathbf{x}\| \leq \aleph_0$ (because $\mathbf{x} \in \mathbf{K}_{\leq \aleph_0}$), there is a strictly $\leq_{\mathbf{x}}$ -increasing cofinal sequence $\langle B_n : n < \omega \rangle$. Let $Y_n = \text{suc}_{B_n}(\text{rt}_{\mathbf{x}})$.

Note that for each $m_1 < m_2$, $Y_{m_1} \cap B_{m_2}$ is an almost front of B_{m_2} (so also it is almost above Y_{m_2}). Hence for $m_1 < m_2 \leq n$ we have that $Y_{m_1} \cap B_n$ is an almost front of B_n which is almost above $Y_{m_2} \cap B_n$. Consequently we may choose $B_n^* \in \text{sb}(B_n)$ such that each $Y_{\ell} \cap B_n^*$ is a front of B_n^* and $Y_{\ell} \cap B_n^*$ is above $Y_{\ell+1} \cap B_n^*$ (for all $\ell < n$). Moreover, we may also require that

$$(\otimes)_0 \text{ for each } \ell < n \text{ and } \eta \in Y_{\ell} \cap B_n^* \text{ we have } (B_{\ell})_{\geq \eta} \leq_{M_{\mathbf{x}}}^* (B_n^*)_{\geq \eta}$$

(remember Observation 2.4(10)).

Fix a list $\langle \rho_{\ell} : \ell < \omega \rangle$ of all members of $M_{\mathbf{x}}$ (possibly with repetitions). By induction on $n < \omega$ choose ν_n such that

- (\otimes)₁ (a) $\nu_n \in Y_n \cap B_n^* = \text{suc}_{B_n^*}(\text{rt}_{\mathbf{x}})$,
 (b) if $\ell < n$, then ν_n, ν_{ℓ} are $<_{M_{\mathbf{x}}}$ -incompatible (i.e., $\nu_{\ell} \parallel_{M_{\mathbf{x}}} \nu_n$),
 (c) if $\ell < n$ and $\rho_{\ell} \neq \text{rt}_{\mathbf{x}}$, then $\rho_{\ell} \parallel_{M_{\mathbf{x}}} \nu_n$.

[Why is the choice possible? By the demand (f) of Definition 2.2(1) applied to $\nu = \text{rt}_{\mathbf{x}}$ and $F = \{\nu_{\ell}, \rho_{\ell} : \ell < n\} \setminus \{\text{rt}_{\mathbf{x}}\}$.]

We define

$$B^* = \{\text{rt}_{\mathbf{x}}\} \cup \bigcup \{B_n^* \cap (M_{\mathbf{x}})_{\geq \nu_n} : n < \omega\}.$$

Now,

$$(\otimes)_2 \ B^* \in \text{CWT}(M_{\mathbf{x}}) \text{ is a tree with root } \text{rt}_{\mathbf{x}} \text{ and } \text{suc}_{B^*}(\text{rt}_{\mathbf{x}}) = \{\nu_n : n < \omega\}.$$

[Why? It should be clear that conditions (a)–(d) of Definition 2.2(1) hold, $\text{rt}(B^*) = \text{rt}_{\mathbf{x}}$ and $\text{suc}_{B^*}(\text{rt}_{\mathbf{x}}) = \{\nu_n : n < \omega\}$. To verify clause (e) suppose $\eta, \nu \in B^*$ are $<_{M_{\mathbf{x}}}$ -incomparable. Then both $\eta \neq \text{rt}_{\mathbf{x}}$ and $\nu \neq \text{rt}_{\mathbf{x}}$, so $\eta, \nu \in \bigcup_{n < \omega} (B^*)_{\nu_n}$. If,

for some n , we have $\eta, \nu \in B_n^* \cap (M_{\mathbf{x}})_{\geq \nu_n}$, then they are $<_{M_{\mathbf{x}}}$ -incompatible as $B_n^* \subseteq B_n$ and B_n satisfies 2.2(1)(e). Otherwise, for some distinct ℓ, n we have $\eta \in B_{\ell}^* \cap (M_{\mathbf{x}})_{\geq \nu_{\ell}}$ and $\nu \in B_n^* \cap (M_{\mathbf{x}})_{\geq \nu_n}$. Now, if we could find $\rho \in M_{\mathbf{x}}$ such that $\rho \geq_{M_{\mathbf{x}}} \eta$ and $\rho \geq_{M_{\mathbf{x}}} \nu$, then ν_{ℓ}, ν_n would be compatible contradicting (\otimes)₁(b), so B^* indeed satisfies clause (e) of Definition 2.2(1). Finally, to verify (f) suppose $\nu \in B^* \setminus \max(B^*)$ and $F \subseteq M_{\mathbf{x}} \setminus (M_{\mathbf{x}})_{\leq \nu}$ is finite. If $\nu_n \leq_{M_{\mathbf{x}}} \nu$ for some n , then the properties of B_n^* apply. So suppose $\nu = \text{rt}_{\mathbf{x}}$. Choose m so that $F \subseteq \{\rho_{\ell} : \ell < m\}$ and use condition (\otimes)₁(c) to argue that for all $n \geq m$ and $\rho \in F$ we have $\nu_n \parallel_{M_{\mathbf{x}}} \rho$.]

Also,

$$(\otimes)_3 \ B \leq_{M_{\mathbf{x}}}^* B^* \text{ for all } B \in \mathcal{B}_{\mathbf{x}}.$$

[Why? Since $\leq_{M_{\mathbf{x}}}^*$ is a partial order and by the choice of B_n , it is enough to show that for each $n < \omega$ we have $B_n \leq_{M_{\mathbf{x}}}^* B^*$, i.e., that every almost front of B_n is an almost front of B^* . To this end suppose that $Z \subseteq B_n$ is an almost front of B_n for

some $n < \omega$. If $Z = \{\text{rt}_{\mathbf{x}}\}$, then there is nothing to do, so suppose $Z \subseteq B_n \setminus \{\text{rt}_{\mathbf{x}}\}$, i.e., $Z \subseteq \bigcup\{(B_n)_{\geq \rho} : \rho \in Y_n\}$. Plainly, the set

$$X = \{\rho \in Y_n : Z \text{ is not an almost front of } (B_n)_{\geq \rho}\}$$

is finite and hence for some $m > n$ we have $X \subseteq \{\rho_\ell : \ell < m\}$. Then for every $k > m$ we have:

- (a) ν_k is incompatible with every $\nu \in X$,
- (b) $Y_n \cap (B_k^*)_{\geq \nu_k}$ is a front of $(B_k^*)_{\geq \nu_k}$,
- (c) $(B_n)_{\geq \eta} \leq_{M_{\mathbf{x}}}^* (B_k^*)_{\geq \eta}$ for every $\eta \in Y_n \cap (B_k^*)_{\geq \nu_k}$ (by $(\otimes)_0$),
- (d) $Z \cap (B_n)_{\geq \eta}$ is an almost front of $(B_n)_{\geq \eta}$ for every $\eta \in Y_n \cap (B_k^*)_{\geq \nu_k}$, and thus
- (e) $Z \cap (B_k^*)_{\geq \eta}$ is an almost front of $(B_k^*)_{\geq \eta}$ for every $\eta \in Y_n \cap (B_k^*)_{\geq \nu_k}$.
- (f) Finally, Z is an almost front of $(B_k^*)_{\geq \nu_k}$ (by Observation 2.4(9) and (b)+(e)).

Since $\text{suc}_{B^*}(\text{rt}_{\mathbf{x}}) = \{\nu_k : k < \omega\}$, we know that $\{\nu_k : m < k < \omega\}$ is an almost front of B^* . Therefore, by Observation 2.4(9) and (f), we conclude that Z is an almost front of B^* .]

Lastly, we define \mathbf{y} :

- $(\otimes)_4$ (a) $(M_{\mathbf{y}}, <_{M_{\mathbf{y}}}) = (M_{\mathbf{x}}, <_{M_{\mathbf{x}}})$,
- (b) $\mathcal{A}_{\nu}^{\mathbf{y}} = \mathcal{A}_{\nu}^{\mathbf{x}}$ when $\nu \in M_{\mathbf{x}} \setminus \{\text{rt}_{\mathbf{x}}\}$, and $\mathcal{A}_{\text{rt}_{\mathbf{x}}}^{\mathbf{y}} = \mathcal{A}_{\text{rt}_{\mathbf{x}}}^{\mathbf{x}} \cup \{B^*\}$,
- (c) $B_1 \leq_{\mathbf{y}} B_2$ if and only if $B_1 \leq_{\mathbf{x}} B_2$ or $B_1 \in A_{\text{rt}_{\mathbf{x}}}^{\mathbf{y}} \wedge B_2 = B^*$.

It should be clear that $\mathbf{y} \in \mathbf{K}_{\leq \aleph_0}$ is as required.

(2,3,4) Straightforward; see also Lemmas 3.4, 3.5 below. \square

Lemma 3.4. *Assume that $\mathbf{x} \in \mathbf{K}_{\leq \aleph_0}$ and $B \in \mathcal{B}_{\mathbf{x}}$ is $\leq_{\mathbf{x}}$ -maximal. Then for some $\mathbf{y} \in \mathbf{K}_{\leq \aleph_0}$ and $B' \in \mathcal{B}_{\mathbf{y}}$ we have*

- (a) $\mathbf{x} \leq \mathbf{y}$, $M_{\mathbf{x}} = M_{\mathbf{y}} = M$, and
- (b) $B' \in \mathcal{B}_{\mathbf{y}}$ is $\leq_{\mathbf{y}}$ -maximal,
- (c) if $\nu \in B' \setminus \max(B')$ and $\rho \in M \setminus M_{\leq \nu}$, then for all but finitely many $\varrho \in \text{suc}_{B'}(\nu)$ we have $\rho \parallel_M \varrho$.

Proof. Fix a list $\langle \rho_\ell : \ell < \omega \rangle$ of all members of $M_{\mathbf{x}}$ (possibly with repetitions). For each $\eta \in B \setminus \max(B)$ by induction on $n < \omega$ we choose $\nu_{\eta,n}$ such that

- (\square) (a) $\nu_{\eta,n} \in \text{suc}_B(\eta)$,
- (b) $\nu_{\eta,n} \neq \nu_{\eta,k}$ for $k < n$ (and hence $\nu_{\eta,n} \parallel \nu_{\eta,k}$ for $k < n$),
- (c) if $k < n$ and $\rho_k \notin M_{\leq \eta}$, then $\rho_k \parallel \nu_{\eta,n}$.

Next, by downward induction on $\eta \in B$ we define

$$B_\eta = \bigcup \{B_{\nu_{\eta,n}} : n < \omega\} \cup \{\eta\}.$$

Lastly we define \mathbf{y} so that:

- $(M_{\mathbf{y}}, <_{\mathbf{y}}) = (M_{\mathbf{x}}, <_{\mathbf{x}})$,
- $\mathcal{A}_{\eta}^{\mathbf{y}} = \mathcal{A}_{\eta}^{\mathbf{x}}$ if $\eta \in M_{\mathbf{x}}$ but $\eta \notin B \setminus \max(B)$, and
- $\mathcal{A}_{\eta}^{\mathbf{y}} = \mathcal{A}_{\eta}^{\mathbf{x}} \cup \{B_\eta\}$ if $\eta \in B \setminus \max(B)$,
- $\mathcal{B}_{\mathbf{y}} = \mathcal{B}_{\mathbf{x}} \cup \{B_{\text{rt}_{\mathbf{x}}}\}$ and for $B', B'' \in \mathcal{B}_{\mathbf{y}}$ we let: $B' \leq_{\mathbf{y}} B''$ if and only if $B' \leq_{\mathbf{x}} B''$ or $B'' = B_{\text{rt}_{\mathbf{x}}}$. \square

Lemma 3.5. (1) *If $\mathbf{x} \in \mathbf{K}_{\leq \aleph_0}$, $Y \in \text{alm-frt}_{\mathbf{x}}$ and $Z \subseteq Y$ then for some $\mathbf{y} \in \mathbf{K}_{\leq \aleph_0}$ we have $\mathbf{x} \leq_{\mathbf{K}} \mathbf{y}$ and either $Z \in D_{\mathbf{y}}^{\mathbf{y}}$ or $(Y \setminus Z) \in D_{\mathbf{y}}^{\mathbf{y}}$.*

- (2) Moreover, if h is a function with domain Y , then above we can demand that for some $B \in \mathcal{B}_y$, $Y \cap B$ is a front of B and for some front Y' of B which is below Y and a one-to-one function h' with domain Y' we have

$$\rho \in Y' \wedge \varrho \in Y \cap B \wedge \rho \leq_{M_y} \varrho \quad \Rightarrow \quad h(\rho) = h'(\varrho).$$

(Note that possibly $Y' = \{\text{rt}_y\}$ and then $h \upharpoonright (Y \cap B)$ is constant.)

Proof. 1) By Lemma 3.3(1) without loss of generality there is $B \in \mathcal{B}_x$ such that B is \leq_x -maximal in \mathcal{B}_x ; clearly $Y \cap B$ is an almost front of B and so without loss of generality $Y \subseteq B$.

We know that $B[\leq Y] := \{\rho \in B : (\exists \nu)[\rho \leq_{M_x} \nu \in Y]\}$ has no ω -branch, so by $<_{M_x}$ -downward induction on $\nu \in B[\leq Y]$ we choose (\mathbf{t}_ν, Y_ν) such that (where $M = M_x$, of course):

- (a) $\mathbf{t}_\nu \in \{0, 1\}$ and
 - if $\mathbf{t}_\nu = 1$, then $Y_\nu \subseteq M_{\geq \nu} \cap Z$,
 - if $\mathbf{t}_\nu = 0$, then $Y_\nu \subseteq M_{\geq \nu} \cap (Y \setminus Z)$,
- (b) $Y_\nu = \max(B'_\nu)$ for some $B'_\nu \in \text{psb}_M(B_{\geq \nu})$,
- (c) if $\nu \in Y$ then $Y_\nu = \{\nu\}$ and $\mathbf{t}_\nu =$ (the truth value of $\nu \in Z$),
- (d) if $\nu \in B[\leq Y] \setminus Y$ then
 - (α) for every finite set $F \subseteq M \setminus M_{\leq \nu}$ there are infinitely many $\varrho \in \text{suc}_B(\nu)$ such that $(\forall \rho \in F)(\rho \parallel \varrho)$ and $\mathbf{t}_\varrho = \mathbf{t}_\nu$,
 - (β) $Y_\nu = \bigcup \{Y_\varrho : \varrho \in \text{suc}_B(\nu) \text{ and } \mathbf{t}_\varrho = \mathbf{t}_\nu\}$.

This is easily done and so \mathbf{t}_{rt_x} is well defined. For $\nu \in B[\leq Y]$ we let

$$B_\nu^* = \{\rho \in B_{\geq \nu} : \text{for some } \varrho \in Y_\nu \text{ we have } \varrho \leq_M \rho \vee \rho \leq_M \varrho\}.$$

Now define \mathbf{y} by adding B_ν^* to \mathcal{A}_ν^x for every $\nu \in B[\leq Y]$, and check.

- 2) First note that by Lemmas 3.3(1) and 3.4 we may assume that there is $B \in \mathcal{B}_x$ such that

- B is \leq_x -maximal,
- Y is a front of B , and
- if $\nu \in B \setminus \max(B)$ and $\rho \in M \setminus M_{\leq \nu}$, then for all but finitely many $\varrho \in \text{suc}_B(\nu)$ we have $\rho \parallel_M \varrho$.

Now note: if $h' : Y' \rightarrow A$, $Y' \in \text{frt}(B')$, $Z = \{\eta \in B' : \text{suc}_{B'}(\eta) \subseteq Y'\}$ is a front of B' and $h' \upharpoonright \text{suc}_{B'}(\eta)$ is one-to-one for all $\eta \in Z$, then we can find $B'' \in \text{psb}_M(B)$ such that $h' \upharpoonright B'' \cap Y'$ is one-to-one. So we may follow similarly as in part (1). \square

Let us recall the following definition.

- Definition 3.6.** (1) We say D is a Q -point when D is an ultrafilter on a countable set $\text{Dom}(D)$ such that if f is a finite-to-one function with domain $\text{Dom}(D)$, then $f \upharpoonright A$ is one-to-one for some $A \in D$.
- (2) A non-principal ultrafilter D on a countable set $\text{Dom}(D)$ is a P -point if for each sequence $\langle A_n : n < \omega \rangle$ of sets from D there is an $A \in D$ such that $A \setminus A_n$ is finite for each $n < \omega$.

We can conclude the main result of this section.

Theorem 3.7. *Assume CH. There is a $\mathbf{x} \in \mathbf{K}$ such that:*

- (a) (α) $\mathcal{A}_\eta^x \neq \{\{\eta\}\}$ for $\eta \in M_x$,

- (β) $\mathcal{B}_x = \mathcal{A}_{\text{rt}(x)}^x \setminus \{\{\text{rt}_x\}\}$ is \aleph_1 -directed under \leq_x ,
- (b) if $Y \in \text{frt}_x^-$, then
- (α) D_Y^x is a non-principal ultrafilter on Y , and
- (β) D_Y^x is a Q -point, see Definition 3.6(1),
- (c) if $B_1 \in \mathcal{B}_x$, then for some $B_2 \in \mathcal{B}_x$ we have $B_1 \leq_x B_2$ and $B_1 \cap \text{suc}_{B_2}(\text{rt}_x) = \emptyset$, moreover¹
- $$(\forall \varrho \in \text{suc}_{B_2}(\text{rt}_x))(\exists^\infty \rho \in \text{suc}_{B_1}(\text{rt}_x))[\varrho \leq_{M_x} \rho].$$
- (d) x is (see Definition 2.11): fat, big, large, and full.

Proof. We choose $x_\alpha \in \mathbf{K}_{\leq \aleph_0}$ by induction on $\alpha < \aleph_1$ so that

- (i) if $\beta < \alpha < \aleph_1$, then $x_\beta \leq_{\mathbf{K}} x_\alpha$,
- (ii) for each successor α , there is a \leq_{x_α} -maximal element in \mathcal{B}_{x_α} .

We use a bookkeeping device to ensure largeness and bigness and

- for $\alpha = 0$ we use Lemma 3.1,
- for α limit we use Definition 2.9(3) and Observation 2.10(2),
- if $\alpha = \beta + 1$, β is limit, then we use Lemma 3.5(1) (and the instructions from our bookkeeping device) to take care of the bigness,
- if $\alpha = \beta + 2$, β is limit, then we use Lemma 3.5(2) (and the instructions from our bookkeeping device) to take care of the largeness,
- if $\alpha = \beta + 3$, β is limit, then we use Lemma 3.3(3,4) (and the instructions from our bookkeeping device) to ensure that at the end x is fat and full,
- if $\alpha = \beta + k$, β is limit, $4 \leq k < \omega$, then we ensure clause (d).

In the end we let $x = \bigcup_{\alpha < \aleph_1} x_\alpha$. Then x is fat, big, large and \mathcal{B}_x is \aleph_1 -directed.

Note that clause (b)(β) follows from the largeness. \square

Definition 3.8. (1) We say that $x \in \mathbf{K}$ is *nice* if it satisfies conditions (a)–(d) of Theorem 3.7. The class of all nice x is denoted by \mathbf{K}_n .

- (2) An $x \in \mathbf{K}$ is *reasonable* if it satisfies (a), (c) of Theorem 3.7. Let \mathbf{K}_r be the set of all $x \in \mathbf{K}$ which are reasonable.
- (3) Let \mathbf{K}_u be the set of $x \in \mathbf{K}_r$ for which clause (b)(α) of Theorem 3.7 holds.
- (4) For $x \in \mathbf{K}$ we say that \mathcal{I} is *x -dense* when:
- (a) $\mathcal{I} \subseteq \mathcal{A}_x$ (see Definition 2.5(b)),
- (b) for every $B_1 \in \mathcal{B}_x$ there is B_2 such that
- (α) $B_1 \leq_x B_2 \in \mathcal{B}_x$, and
- (β) if $A \subseteq M_x \setminus \{\text{rt}_x\}$ is finite, then for some ν we have
- $$\nu \in \text{suc}_{B_2}(\text{rt}_x), \quad (B_2)_{\geq \nu} \in \mathcal{I}, \quad \text{and} \quad (\forall \rho \in A)(\rho \parallel \nu).$$

- (5) For $x \in \mathbf{K}$ we say \mathcal{I} is *x -open* if clause (a) from part (4) above and
- (c) if $B_1 \in \mathcal{I}$ then $\text{sb}(B_1) \cap \mathcal{A}_x \subseteq \mathcal{I}$.
- (6) Let \mathbf{K}_g be the class of $x \in \mathbf{K}_r$ which are *good*, which means: if \mathcal{I} is x -dense, x -open and $B_1 \in \mathcal{B}_x$ then for some $B_2 \in \mathcal{B}_x$ we have $B_1 \leq_x B_2$ and $(B_2)_{\geq \eta} \in \mathcal{I}$ for all but finitely many $\eta \in \text{suc}_{B_2}(\text{rt}_x)$.
- (7) We say that $x \in \mathbf{K}$ is *ultra* if it is both nice and good. Let \mathbf{K}_{ut} be the class of x which are ultra, i.e., $\mathbf{K}_{\text{ut}} = \mathbf{K}_g \cap \mathbf{K}_n$.

Theorem 3.9. Assume \diamond_{\aleph_1} . Then there exists an ultra $x \in \mathbf{K}$.

¹Not a serious addition. As always, the number of $\varrho \in \text{suc}_{B_2}(\text{rt}_x)$ failing this is finite.

Proof. We repeat the proof of Theorem 3.7 but at limit stages $\delta < \aleph_1$ we use additionally \diamond_{\aleph_1} to take care of the additional demand $\mathbf{x} \in \mathbf{K}_g$ here. So we are given:

- (A) $\delta < \aleph_1$ limit,
- (B) \mathcal{I} such that for some \mathbf{y}, \mathcal{I} we have
 - $\mathbf{x}_\delta \leq_{\mathbf{K}} \mathbf{y} \in \mathbf{K}_{\aleph_1}$,
 - $\mathcal{I} \subseteq \mathcal{A}_{\mathbf{y}}$ is dense open, and $\mathcal{I} = \mathcal{I} \cap \mathcal{A}_{\mathbf{x}_\delta}$,
 - for some countable elementary submodel $N \prec \mathcal{H}(\aleph_2)$, $(\mathbf{y}, \mathcal{I}) \in N$ and $(\mathbf{x}_\delta, \mathcal{I}) = (\mathbf{y} \upharpoonright N, \mathcal{I} \cap N)$, so $M_{\mathbf{x}_\delta} = M_{\mathbf{y}} \upharpoonright N$, etc.

Let $\langle B_\ell^0 : \ell < \omega \rangle$ be an increasing cofinal subset of $(\mathcal{B}_{\mathbf{x}_\delta}, \leq_{\mathbf{x}_\delta})$. For every ℓ there is $B_\ell^1 \in \mathcal{B}_{\mathbf{x}_\delta}$ such that $B_\ell^0 \leq_{\mathbf{x}_\delta} B_\ell^1$, and for every finite $A \subseteq M_{\mathbf{x}_\delta} \setminus \{\text{rt}(\mathbf{x}_\delta)\}$ there is $\nu \in \text{suc}_{B_\ell^1}(\text{rt}(\mathbf{x}_\delta))$ such that

$$(\forall \rho \in A)(\rho \parallel \nu) \quad \text{and} \quad (B_\ell^1)_{\geq \nu} \in \mathcal{I}.$$

Clearly, for every ℓ for some $k(\ell) > \ell$ we have $B_\ell^1 \leq_{\mathbf{x}_\delta} B_{k(\ell)}^0$. We can choose $\langle \ell_n : n < \omega \rangle$ so that $k(\ell_n) < \ell_{n+1}$. Let $B_n = B_{\ell_n}^1$. We continue as in Lemma 3.3(1) using the $\langle B_n : n < \omega \rangle$ and, when choosing ν_n , demanding additionally that $(B_n)_{\geq \nu_n} \in \mathcal{I}$. (Note that $(B_n)_{\geq \nu_n} \in \mathcal{I}$ implies $(B_n^*)_{\geq \nu_n} \in \mathcal{I}$ for B_n^* as there.) \square

Proposition 3.10. *Assume $\mathbf{x} \in \mathbf{K}_n$.*

- (α) *If $B \in \mathcal{B}_{\mathbf{x}}$ and $Y_1, Y_2 \in \text{frt}(B)$ and Y_2 is above Y_1 , then $h_{Y_2, Y_1}^{\mathbf{x}}$ exemplifies $D_{Y_1}^{\mathbf{x}} \leq_{\text{RK}} D_{Y_2}^{\mathbf{x}}$.*
- (β) *$\{D_Y^{\mathbf{x}} : Y \in \text{frt}_{\mathbf{x}}^-\}$ is \geq_{RK} -directed (even \aleph_1 directed).*
- (γ) *If $Y \in \text{alm-frt}_{\mathbf{x}}^-$, then \leq_{RK} -below $D_Y^{\mathbf{x}}$ there is no P -point.*

Proof. (α) Follows from Observation 2.8(6).

(β) By (α) and the directness of $\mathcal{B}_{\mathbf{x}}$.

(γ) Let $B_1 \in \mathcal{B}_{\mathbf{x}}$ be such that $B_1 \cap Y$ is an almost front of B_1 . Suppose that $h : Y \rightarrow \mathbb{N}$ is such that $h^{-1}[\{n\}] = \emptyset \pmod{D_Y^{\mathbf{x}}}$ for every n , hence there is $A_n \in \mathcal{B}_{\mathbf{x}}$ which witnesses this. Assume towards contradiction that $h(D_Y^{\mathbf{x}})$ is a P -point; without loss of generality h is onto \mathbb{N} . As $\mathcal{B}_{\mathbf{x}}$ is \aleph_1 -directed we may pick $B_2 \in \mathcal{B}_{\mathbf{x}}$ such that $A_n \leq_{\mathbf{x}} B_2$ (for all $n < \omega$) and $B_1 \leq_{\mathbf{x}} B_2$.

As \mathbf{x} is large, we may apply the Definition 2.11 of large to the pair (B_2, h') where $h'(\eta) = h(\nu)$ when $\nu \leq_{M_{\mathbf{x}}} \eta \in \max(B)$ and zero if there is no such ν . So there are B_3, Y_3 such that

- (i) $B_2 \leq_{\mathbf{x}} B_3$,
- (ii) Y_3 is a front of B_3 below $Y \cap B_3$,
- (iii) for $\eta, \nu \in Y \cap B_3$ we have: $h(\eta) = h(\nu) \Leftrightarrow (\exists \rho \in Y_3)(\rho \leq_{M_{\mathbf{x}}} \eta \wedge \rho \leq_{M_{\mathbf{x}}} \nu)$.

Let $Z = \text{suc}_{B_3}(\text{rt}_{\mathbf{x}})$. If $Y_3 = \{\text{rt}_{\mathbf{x}}\}$, then for some n we have $h^{-1}[\{n\}] \in D_Y^{\mathbf{x}}$, a contradiction. Therefore $Y_3 \neq \{\text{rt}_{\mathbf{x}}\}$ and thus $\text{rt}_{\mathbf{x}} \notin Y_3$, so Y_3 is above Z . Clearly, $D_Z^{\mathbf{x}} \leq_{\text{RK}} h(D_Y^{\mathbf{x}})$ and hence $D_Z^{\mathbf{x}}$ is a P -point.

By clauses (c) and (d) of Theorem 3.7 there is $B_4 \in \mathcal{B}_{\mathbf{x}}$ such that $B_3 \leq_{\mathbf{x}} B_4$, $B_4 \cap Z$ is a front of B_4 and

$$(\forall \rho \in \text{suc}_{B_4}(\text{rt}_{\mathbf{x}}))(\exists^\infty \rho \in \text{suc}_{B_3}(\text{rt}_{\mathbf{x}}))[\rho \leq_{M_{\mathbf{x}}} \rho].$$

For each $\rho \in \text{suc}_{B_4}(\text{rt}_{\mathbf{x}})$ let $Z_\rho = \{\rho \in Z : \rho \leq_{M_{\mathbf{x}}} \rho\}$, so $\langle Z_\rho : \rho \in \text{suc}_{B_4}(\text{rt}_{\mathbf{x}}) \rangle$ is a partition of Z , and $Z_\rho = \emptyset \pmod{D_Z^{\mathbf{x}}}$ for each ρ . But clearly there is no $Z' \in D_Z^{\mathbf{x}}$

such that $Z' \cap Z_\varrho$ is finite for every $\varrho \in \text{succ}_{B_4}(\text{rt}_x)$, contradiction to “ D_Z^x is a P -point” . \square

4. BASIC CONNECTIONS TO FORCING

Definition 4.1. (1) For a forcing notion \mathbb{Q} and $p \in \mathbb{Q}$ we define $\mathfrak{D}_{\text{sb}} = \mathfrak{D}_p^{\text{sb}} = \mathfrak{D}_{\mathbb{Q},p}^{\text{sb}}$, the strong bounding game between the null player NU and the bounding player BND as follows.

- (a) A play last ω moves, and
- (b) in the n -th move:
 - (α) first the NU player gives a (non-empty) tree \mathcal{T}_n with ω levels and no maximal node and a \mathbb{Q} -name \dot{F}_n of a function with domain \mathcal{T}_n such that

$$\eta \in \mathcal{T}_n \Rightarrow p \Vdash_{\mathbb{Q}} \text{“} \dot{F}_n(\eta) \in \text{succ}_{\mathcal{T}_n}(\eta) \text{”},$$

(β) then BND player chooses $\eta_n \in \mathcal{T}_n$.

- (c) In the end, the BND player wins the play $\langle \mathcal{T}_n, \eta_n : n < \omega \rangle$ iff there is $q \in \mathbb{Q}$ above p forcing that

$$\text{“} (\forall n < \omega) (\exists k < \text{level}(\eta_n)) (\dot{F}_n(\eta_n \upharpoonright k) \leq_{\mathcal{T}_n} \eta_n \wedge k \text{ is even}) \text{”},$$

where $\eta_n \upharpoonright k$ is the unique $\nu \leq_{\mathcal{T}_n} \eta_n$ of level k .

- (2) Omitting p means NU chooses it in his first move.
- (3) A forcing notion \mathbb{Q} is *strongly bounding* if for every condition $p \in \mathbb{Q}$ player BND has a winning strategy in the game $\mathfrak{D}_{\mathbb{Q},p}^{\text{sb}}$.

Definition 4.2. (1) We say $\mathcal{P} \subseteq [\mathbb{N}]^{\aleph_0}$ is big when for every $\mathbf{c} : \mathbb{N} \rightarrow \{0,1\}$ there is $A \in \mathcal{P}$ such that $\mathbf{c} \upharpoonright A$ is constant.

- (2) For $B \in \text{CWT}(\omega > \omega, \triangleleft)$ we say that a family $\mathcal{B} \subseteq \text{psb}(B)$ is big (in B) when for every $\mathbf{c} : \text{max}(B) \rightarrow \{0,1\}$ there is $B' \in \mathcal{B}$ such that $\mathbf{c} \upharpoonright \text{max}(B')$ is constant.

- (3) For $B \in \text{CWT}(\omega > \omega, \triangleleft)$ we say that a family $\mathcal{B} \subseteq \text{psb}(B)$ is large (in B) when for every function \mathbf{c} with domain $\text{max}(B)$ there is $B' \in \mathcal{B}$ and front Y of B' such that for every $\eta, \nu \in \text{max}(B')$ we have $\mathbf{c}(\eta) = \mathbf{c}(\nu) \Leftrightarrow (\exists \rho \in Y)(\rho \leq_B \nu \wedge \rho \leq_B \eta)$.

Theorem 4.3. *Assume that:*

- (a) $B \in \text{CWT}(M)$ for a partial order M , without loss of generality $M = (\omega > \omega, \triangleleft)$,
- (b) \mathbb{Q} is a strongly bounding forcing notion,
- (c) (α) forcing with \mathbb{Q} preserves some non-principal ultrafilter on \mathbb{N} , or just
 (β) $([\mathbb{N}]^{\aleph_0})^{\mathbf{V}}$ is big in $\mathbf{V}^{\mathbb{Q}}$, see Definition 4.2,
- (d) $p \Vdash \text{“} \dot{A} \subseteq \text{max}(B) \text{”}$.

Then there are $B' \in \text{psb}(B)$ and $q \in \mathbb{Q}$ such that $p \leq q$ and

$$q \Vdash \text{“} \text{max}(B') \subseteq \dot{\mathcal{T}} \text{”} \quad \text{or} \quad q \Vdash \text{“} \text{max}(B') \subseteq \text{max}(B) \setminus \dot{\mathcal{T}} \text{”}.$$

Proof. We prove this by induction on $\text{Dp}(B)$ (see Definition 2.2), for all such B 's. Let $\eta = \text{rt}(B)$.

Case 1: $\text{Dp}(B) = 0$

Trivial, as then $B = \{\eta\}$, i.e., B is a singleton so $B' = B$ can serve.

Case 2: $\text{Dp}_x(B) = 1$

Then $\text{Dp}(B_{\geq \nu}) = 0$ for all $\nu \in B \setminus \{\eta\}$. Now, $|B \setminus \{\eta\}| = \aleph_0$ and we just need to find $p' \in \mathbb{Q}$ above p such that $\{\nu \in B : \nu \neq \eta \text{ and } p' \text{ forces } \nu \in \underline{A} \text{ or forces } \nu \notin \underline{A}\}$ is infinite. As $\Vdash_{\mathbb{Q}} \text{“} ([\mathbb{N}]^{\aleph_0})^{\mathbf{V}}$ is big in $\mathbf{V}^{\mathbb{Q}}$ ” (see clause (c) of our assumptions) this is possible.

Case 3: $\alpha = \text{Dp}(B) > 1$

Let $Y = \text{suc}_B(\eta)$. Then for $\nu \in Y$ we have $\text{Dp}(B_{\geq \nu}) < \alpha$, hence the induction hypothesis applies to $B_{\geq \nu}$. We may assume that if ρ is not below η then for all but finitely many $\nu \in Y$ we have $\nu \parallel \rho$ (cf. the proof of Lemma 3.4). Let $\langle \nu_n : \nu \in \mathbb{N} \rangle$ list Y .

We simulate a play of $\mathfrak{D}_{\mathbb{Q}, p}^{\text{sb}}$ in which the BND player uses a winning strategy and the NU player acts so that in the n -th move:

- (*)₁ (a) $\mathcal{I}_n = \{\langle B_0, \dots, B_{k-1} \rangle : k \in \mathbb{N}, B_\ell \in \text{psb}(B_{\geq \nu_n}) \text{ for } \ell < k \text{ and } B_{\ell+1} \subseteq B_\ell \text{ if } \ell + 1 < k\}$,
- (b) $\langle \mathcal{I}_n \rangle$ is being an initial segment,
- (c) $F_n(\langle B_0, \dots, B_{k-1} \rangle)$ is $\langle B_0, \dots, B_{k-1}, B' \rangle$ for some $B' \in \text{psb}(B_{k-1}) \cap \mathbf{V}$ such that
 - either $\max(B') \subseteq \underline{A}$ or $\max(B') \cap \underline{A} = \emptyset$.

There is such a function F_n because of the induction hypothesis.

Clearly we can do this. As the player BND has used a winning strategy, BND has won the play so there is $q \in \mathbb{Q}$ stronger than p and such that

- (*)₂ $q \Vdash \text{“for every } n \text{ for some even } k < \text{level}_{\mathcal{I}_n}(\eta_n) \text{ we have } F_n(\eta_n \upharpoonright k) \leq_{\mathcal{I}_n} \eta_n \text{”}$.

Hence by the choice of (\mathcal{I}_n, F_n) , letting $\eta_n = \langle B_{n,0}, \dots, B_{n,k(n)} \rangle$ we have

- (*)₃ for some $\langle \mathbf{t}_n : n \in \mathbb{N} \rangle$
 - (α) $B_{n,k(n)} \in \text{psb}(B_{\geq \nu_n})$,
 - (β) \mathbf{t}_n is a \mathbb{Q} -name of the truth value,
 - (γ) $q \Vdash \text{“if } \mathbf{t}_n = 1 \text{ then } \max(B_{n,k(n)}) \subseteq \underline{A}, \text{ and if } \mathbf{t}_n = 0 \text{ then } \max(B_{n,k(n)}) \cap \underline{A} = \emptyset \text{”}$.

Now by clause (c) of our assumptions

- (*)₄ there is an infinite $\mathcal{U} \subseteq \mathbb{N}$, a truth value \mathbf{t} and a condition r such that $q \leq_{\mathbb{Q}} r$ and $r \Vdash \text{“}\mathbf{t}_n = \mathbf{t} \text{ for } n \in \mathcal{U} \text{”}$.

Lastly, let $B_* = \bigcup \{B_{n,k(n)} : n \in \mathcal{U}\} \cup \{\eta\}$ and clearly B_*, r are as required. \square

Remark 4.4. In the assumption (b) of Theorem 4.3 it is enough that the BND player does not lose the game $\mathfrak{D}_{\mathbb{Q}}^{\text{sb}}$, i.e., the NU player has no winning strategy.

Theorem 4.5. *Assume that*

- (a) \mathbb{Q} is an $\omega\omega$ -bounding proper forcing notion,
- (b) forcing with \mathbb{Q} preserves some P -point, and
- (c) $B \in \text{CWT}(\omega^{>\omega}, \triangleleft)$.

Then $(\text{psb}(B))^{\mathbf{V}}$ is big in $\mathbf{V}^{\mathbb{Q}}$; see Definition 4.2(2).

Proof. Let D be a P -point ultrafilter such that $\Vdash_{\mathbb{Q}} \text{“} D \text{ generates an ultrafilter”}$ and $p \in \mathbb{Q}$. Suppose that $p \Vdash \text{“} \mathcal{C} : \max(\underline{B}) \rightarrow \{0, 1\} \text{”}$. Let χ be a large enough regular cardinal and $N \prec (\mathcal{H}(\chi), \in)$ be a countable model with $B, \mathbb{Q}, p, \mathcal{C}, \dots \in N$. Let $q \in \mathbb{Q}$ be such that

- (a) $p \leq_{\mathbb{Q}} q$,
- (b) q is (N, \mathbb{Q}) -generic,

- (c) for some $g \in {}^\omega \omega^{\mathbf{V}}$ we have $q \Vdash$ “if $f \in {}^\omega \omega \cap N$, then $f <_{J_{\omega}^{\text{bd}}} g$ ”,
- (d) for some $A \in D$ we have $q \Vdash$ “if $B \in D \cap N$, then $A \subseteq^* B$ ”.

From (g, A) we can compute \mathbf{c} and $B' \in (\text{psb}(B))^{\mathbf{V}}$ such that $q \Vdash$ “ $\mathcal{C} \upharpoonright B'$ is constantly \mathbf{c} ”, so we are done. \square

Theorem 4.6. *Assume that $\mathbf{x} \in \mathbf{K}$ and*

- (A) \mathbb{Q} is a proper forcing notion,
- (B) D_* is a Ramsey ultrafilter in \mathbf{V} ,
- (C) $\Vdash_{\mathbb{Q}}$ “ $\text{fil}(D_*)$ is a Ramsey ultrafilter”,
- (D) $B \in \mathcal{B}_{\mathbf{x}}$.

Then $(\text{psb}(B))^{\mathbf{V}}$ is large in $\mathbf{V}^{\mathbb{Q}}$ (see Definition 4.2).

Proof. We prove this by induction on $\text{Dp}(B)$ for $B \in \mathcal{B}_{\mathbf{x}}$. Let $\mathbf{c} : \max(B) \rightarrow \mathbb{N}$ be from $\mathbf{V}^{\mathbb{Q}}$ and we should find (B', Y) as promised. We shall work in $\mathbf{V}^{\mathbb{Q}}$.

If $\text{Dp}(B) = 0$, i.e., $|B| = 1$ this is trivial.

If $\text{Dp}(B) = 1$ let $\langle \eta_n : n \in \mathbb{N} \rangle \in \mathbf{V}$ list $\text{suc}_B(\text{rt}_{\mathbf{x}})$: by assumption (C) in $\mathbf{V}^{\mathbb{Q}}$, for some $A \in \text{fil}(D_*)$ the sequence $\langle \mathbf{c}(\eta_n) : n \in A \rangle$ is constant or without repetitions. Without loss of generality $A \in D_* \subseteq \mathbf{V}$ and then $\{\text{rt}_{\mathbf{x}}\} \cup \{\eta_n : n \in A\}$ is as required.

So assume $\text{Dp}(B) > 1$. Without loss of generality $0 \notin \text{Rang}(\mathbf{c})$. For $\nu \in B \setminus \max(B)$ let $\langle \eta_{\nu, n} : n \in \mathbb{N} \rangle$ list $\text{suc}_B(\nu)$ so that the function $(\nu, n) \mapsto \eta_{\nu, n}$ belongs to \mathbf{V} . In $\mathbf{V}^{\mathbb{Q}}$, by downward induction on $\nu \in B$, we choose $k_{\nu} = k(\nu)$, $A_{\nu}, A_{\nu, \rho}$ and $\mathbf{t}_{\nu, \rho}$ so that the following requirements (a)-(d) are satisfied:

- (a) $k_{\nu} \in \mathbb{N}$, $A_{\nu} \in D_*$,
- (b) if $\nu \in \max(B)$, then $k_{\nu} = \mathbf{c}(\nu)$, so > 0 ,
- (c) if $\nu \notin \max(B)$ then $(\alpha)_{\nu}$ or $(\beta)_{\nu}$ where:
 - $(\alpha)_{\nu}$ $k_{\nu} = 0$ and $\langle k(\eta_{\nu, n}) : n \in A_{\nu} \rangle$ is with no repetitions, all non-zero,
 - $(\beta)_{\nu}$ $\langle k(\eta_{\nu, n}) : n \in A_{\nu} \rangle$ is constantly k_{ν} ,
- (d) for $\nu, \rho \in B \setminus \max(B)$ we have $A_{\nu, \rho} \in D_*$ and $\mathbf{t}_{\nu, \rho} \in \{0, 1\}$ and
 - either $\mathbf{t}_{\nu, \rho} = 1$ and $n \in A_{\nu, \rho} \Rightarrow k(\eta_{\rho, n}) = k(\eta_{\nu, n})$
 - or $\mathbf{t}_{\nu, \rho} = 0$ and $\{k(\eta_{\rho, n}) : n \in A_{\nu, \rho}\}$ is disjoint to $\{k(\eta_{\nu, n}) : n \in A_{\nu, \rho}\}$.

This is possible by assumption (C). By the same assumption, there is $A_* \in D_*$ such that

- (*)₁ (i) if $\nu \in B \setminus \max(B)$ then $A_* \subseteq^* A_{\nu}$,
- (ii) if $\nu, \rho \in B \setminus \max(B)$ then $A_* \subseteq^* A_{\nu, \rho}$.

Let $\langle \nu_n : n \in \mathbb{N} \rangle$ list $B \setminus \max(B)$ and let f_1 be the function with domain $B \setminus \max(B)$ such that

$$f_1(\nu) = \{\eta_{\nu, n} : n \in A_* \setminus A_{\nu} \text{ or for some } k < \ell \text{ we have } \nu = \nu_{\ell} \wedge n \in A_* \setminus A_{\nu_k, \nu_{\ell}}\}$$

(so $f_1(\nu) \in [\text{suc}_B(\nu)]^{< \aleph_0}$).

As the forcing \mathbb{Q} satisfies (C), it is bounding, so there is a function $f_2 \in \mathbf{V}$ with domain $B \setminus \max(B)$ such that $f_1(\nu) \subseteq f_2(\nu) \in [\text{suc}_B(\nu)]^{< \aleph_0}$. Clearly

- (*)₂ $B_1 = A_{B, f} = \{\nu \in B : \text{if } \rho \in B \text{ satisfies } \text{rt}_{\mathbf{x}} \leq_B \rho <_B \nu \text{ and } n \text{ is such that } \eta_{\rho, n} \leq_B \nu \text{ then } n \in A_* \text{ but } \eta_{\rho, n} \notin f_2(\nu)\} \in \text{psb}(B)^{\mathbf{V}}$.

Define

- (*)₃ $Y = \{\nu \in B_1 : k_{\nu} \neq 0 \text{ and } \rho <_B \nu \Rightarrow k_{\rho} = 0\}$.

Plainly,

- (*)₄ (a) Y is a front of B_1 , and

(b) if $\nu \in Y$ then $\mathbf{c} \upharpoonright (B_1)_{\geq \nu}$ is constantly k_ν .

Note that

(*)₅ if $\nu \in B_1$ and $k_\nu = 0$, then either $k_\eta = 0$ for all $\eta \in \text{suc}_{B_1}(\nu)$, or $k_\eta > 0$ for all $\eta \in \text{suc}_{B_1}(\nu)$.

Hence

(*)₆ if $\nu \in B_1 \setminus \max(B_1)$ and $\text{suc}_{B_1}(\nu)$ is not disjoint to Y , then $\text{suc}_{B_1}(\nu) \subseteq Y$.

If $Y = \{\text{rt}_{\mathbf{x}}\}$ we are done, so assume not. Let $Z = \{\eta \in B_1 : \eta \notin \max(B_1) \text{ and } \text{suc}_{B_1}(\eta) \subseteq Y\}$. So

(*)₈ (a) both Z and Y are fronts of B_1 ,
 (b) both Z and Y belong to \mathbf{V} ,
 (c) if $\nu \in Y$ then $\langle k_\rho : \rho \in \max((B_1)_{\geq \nu}) \rangle$ is constantly k_ν .

Also if $Z = \{\text{rt}_{\mathbf{x}}\}$ we are done so assume not and let $\langle \nu_n : n \in \mathbb{N} \rangle$ list Z . As $\text{fil}(D_*)$ is a Ramsey ultrafilter we can find \bar{n} such that

(*)₉ (a) $\bar{n} = \langle n(i) : i \in \mathbb{N} \rangle$ is an increasing enumeration of a member of D_* , hence $\bar{n} \in \mathbf{V}$,
 (b) if $\ell \leq i$ then $\eta_{\nu_\ell, n(i)} \in B_1$,
 (c) if $\ell < i$, $\mathbf{t}_{\nu_\ell, \nu_i} = 0$ and $\nu_\ell, \nu_i \in B_1[\leq Z]$, then $\{k(\eta_{\nu_i, n(j)}) : i \leq j\}$ is disjoint from $\{k(\eta_{\nu_\ell, n(j)}) : i \leq j\}$, moreover it is disjoint from $\{k(\eta_{\nu_\ell, n(j)}) : j \in \mathbb{N}\}$.

Lastly, as $\bar{n} \in \mathbf{V}$ we can find in \mathbf{V} a partition $\langle C_\ell : \ell \in \mathbb{N} \rangle$ of \mathbb{N} to (pairwise disjoint) infinite sets and let

$$B_2 = \{\varrho \in B_1 : \text{ if } \nu_\ell <_{B_1} \varrho \text{ and } \nu_\ell \in B_1[\leq Z], \\ \text{ then for some } i \in C_\ell \text{ we have } i > \ell \text{ and } \eta_{\nu_\ell, n(i)} \leq_{B_2} \varrho\}.$$

Easily $B_2 \in \mathbf{V}$, $B_2 \in \text{psb}(B_1)$ and it is as required. \square

Motivated by Definition 4.1 we introduce the following bounding games for a forcing notion \mathbb{Q} .

Definition 4.7. Let \mathbb{Q} be a forcing notion and $p \in \mathbb{Q}$

- (1) We define a game $\mathfrak{D}_{\text{bd}} = \mathfrak{D}_p^{\text{bd}} = \mathfrak{D}_{\mathbb{Q}, p}^{\text{bd}}$ by
 - (a) a play last ω -moves,
 - (b) in the n -th move
 - (α) first the NU player gives a \mathbb{Q} -name τ_n of a member of \mathbf{V} and then
 - (β) the BND player gives a finite set $w_n \subseteq \mathbf{V}$.
 - (c) In the end of the play the BND player wins the play iff there is $q \in \mathbb{Q}$ above p forcing “ $\tau_n \in w_n$ ” for every n .

Omitting p means that the player NU chooses it in his first move.

- (2) A game $\mathfrak{D}_{\text{ufbd}} = \mathfrak{D}_p^{\text{ufbd}} = \mathfrak{D}_{\mathbb{Q}, p}^{\text{ufbd}}$ is defined as follows.
 - (a) A play last ω -moves, and
 - (b) in the n -th move
 - (α) first the NU player chooses an ultrafilter E_n on some set I_n from \mathbf{V} and a \mathbb{Q} -name E_n^+ of an ultrafilter on I_n extending E_n and a \mathbb{Q} -name X_n of a member of E_n^+
 - (β) the BND player chooses $t_n \in I_n$.
 - (c) In the end of the play the BND player wins the play iff there is $q \in \mathbb{Q}$ above p forcing “ $t_n \in X_n$ ” for every n .

Omitting p means that the player NU chooses it in his first move.

- (3) A game $\mathfrak{D}_{\mathbb{Q},p}^{\text{vfbd}} = \mathfrak{D}_p^{\text{vfbd}} = \mathfrak{D}_{\mathbb{Q},p}^{\text{vfbd}}$ is defined like $\mathfrak{D}_p^{\text{vfbd}}$ in (2) but now

$$\Vdash_{\mathbb{Q}} \text{“} \dot{X}_n \in E_n \text{ or just includes a member of } E_n \text{”},$$

so \dot{E}_n^+ is redundant.

Basic relations between the games introduced above are given by the following result.

Proposition 4.8. *Let \mathbb{Q} be a forcing notion.*

- (1) *If BND wins in $\mathfrak{D}_{\mathbb{Q},p}^{\text{sb}}$ then BND wins in $\mathfrak{D}_{\mathbb{Q},p}^{\text{bd}}$ which implies that \mathbb{Q} is a bounding forcing.*
- (2) *The player BND wins in $\mathfrak{D}_{\mathbb{Q},p}^{\text{bd}}$ iff BND wins in $\mathfrak{D}_{\mathbb{Q},p}^{\text{vfbd}}$.*
- (3) *If the player BND wins in $\mathfrak{D}_{\mathbb{Q},p}^{\text{vfbd}}$ then BND wins in $\mathfrak{D}_{\mathbb{Q},p}^{\text{vfbd}}$.*
- (4) *We can replace in (1)–(3) above “wins” by “do not lose”.*

Proof. 1) The second implication is obvious, so we concentrate on the first. For every τ , a \mathbb{Q} -name of an ordinal we define a pair (T_τ, F_τ) as follows:

- (*)₁ (a) let $u = \{\alpha : \mathbb{K}_{\mathbb{Q}} \text{ “} \tau \neq \alpha \text{”}\}$, it is a non-empty set of $\leq |\mathbb{Q}|$ ordinals,
- (b) T_τ is the tree $\{\eta : \eta \in \omega^{>u}\}$, i.e., ordered by \triangleleft (being an initial segment),
- (c) $F_\tau(\eta) = \eta \hat{\ } \langle \tau \rangle$ for $\eta \in T_\tau$.

Clearly,

- (*)₂ (a) T_τ is in \mathbf{V} , a tree with ω levels,
- (b) F_τ is a \mathbb{Q} -name of a function with domain T_τ such that $\Vdash_{\mathbb{Q}} \text{“} F_\tau(\eta) \in \text{suc}_{T_\tau}(\eta) \text{”}$.
- (c) if $q \in \mathbb{Q}$ and $\eta \in T_\tau$ (so $\text{Rang}(\eta)$ is a finite subset of u) then the following are equivalent:
 - (i) $q \Vdash \text{“} \tau \in \text{Rang}(\eta) \text{”}$,
 - (ii) $q \Vdash \text{“for some } \nu \triangleleft \eta \text{ we have “} \nu \hat{\ } \langle F_\tau(\nu) \rangle \trianglelefteq \eta \text{”}$.

So playing the game $\mathfrak{D}_{\mathbb{Q},p}^{\text{bd}}$ we can “translate” it to a play of $\mathfrak{D}_{\mathbb{Q},p}^{\text{sb}}$ replacing the NU choice of τ_n by the choice of (T_τ, F_τ) . Thus every strategy st_1 of BND in $\mathfrak{D}_{\mathbb{Q},p}^{\text{sb}}$ translates it to a strategy st_2 of the player BND in $\mathfrak{D}_{\mathbb{Q},p}^{\text{bd}}$.

2) We now need two translations.

Translating $\mathfrak{D}_{\mathbb{Q},p}^{\text{vfbd}}$ to $\mathfrak{D}_{\mathbb{Q},p}^{\text{bd}}$:

So we are given a move $y = (I, E, \dot{X})$ of NU in a play of $\mathfrak{D}_{\mathbb{Q},p}^{\text{vfbd}}$ as in Definition 4.7(3), i.e.,

- $I \in \mathbf{V}$, E is an ultrafilter on I , in \mathbf{V} , and
- $\Vdash_{\mathbb{Q}} \text{“} \dot{X} \in E \text{ or just includes a member } \dot{X}' \text{ of } E \text{”}$.

Now,

- (*) if $q \Vdash \text{“} \dot{X}' \in \mathscr{W} \text{”}$ where $\mathscr{W} \subseteq E$ is finite (\mathscr{W} an object in \mathbf{V} not a name), then $\bigcap \{A : A \in \mathscr{W}\}$ is non-empty and $t \in \bigcap \{A : A \in \mathscr{W}\} \Rightarrow q \Vdash \text{“} t \in \dot{X}' \subseteq \dot{X} \text{”}$.

Translating $\mathfrak{D}_{\mathbb{Q},p}^{\text{bd}}$ to $\mathfrak{D}_{\mathbb{Q},p}^{\text{vfbd}}$:

Given $y = (I, \tau)$, τ a \mathbb{Q} -name of a member I of \mathbf{V} we define $I_y = [I]^{<\aleph_0} \in \mathbf{V}$ and choose $E_y \in \mathbf{V}$ an ultrafilter on I_y such that $u_* \in [I]^{<\aleph_0} \Rightarrow \{u \in [I]^{<\aleph_0} : u_* \subseteq$

$u\} \in E$; lastly we choose

$$\underline{X}_y = \{u \in [I]^{<\aleph_0} : \mathcal{T} \in u\}.$$

So $(I_y, E_y, \underline{X}_y)$ is a legal move in $\mathcal{D}_{\mathbb{Q},p}^{\text{vfbd}}$ and for a finite subset t of I :

(**) if $q \Vdash "t \in \underline{X}_y"$ then $q \Vdash "\mathcal{T} \in t"$.

3) Obvious.

4) The same proof. □

Acknowledgement: We thank Alan Dow for asking me about $\boxplus_2(B),(C)$ and for some comments and Andrzej Roslanowski for much help. We also thank Alice Leonhardt for the beautiful typing.

The author thanks the National Science Foundation, Grant 0600940 and the US-Israel Binational Science Foundation, Grant 2010405 for partial support of this work.

REFERENCES

- [1] A. Blass, Combinatorial Cardinal Characteristics of the Continuum, in: Handbook of Set Theory, edited by M. Foreman and A. Kanamori (Springer, 2010), pp. 395–490.
- [2] S. Shelah, Proper and improper forcing, Perspectives in Mathematical Logic (Springer, 1998).
- [3] S. Shelah, On what I do not understand (and have something to say:) Part I, Fundamenta Mathematicae **166**, 1–82 (2000).
- [4] E. K. van Douwen, The integers and topology, in: Handbook of Set-Theoretic Topology, edited by K. Kunen and J. E. Vaughan (Elsevier Science Publishers, 1984), pp. 111–167.
- [5] S. Shelah, Ultrafilters on \aleph_1 generated by few conditions, in preparation ([Sh:F1127]).
- [6] A. Blass and S. Shelah, There may be simple P_{\aleph_1} - and P_{\aleph_2} -points and the Rudin-Keisler ordering may be downward directed, Annals of Pure and Applied Logic **33**, 213–243 (1987).
- [7] S. Shelah, The combinatorics of reasonable ultrafilters, Fundamenta Mathematicae **192**, 1–23 (2006).
- [8] A. Roslanowski and S. Shelah, Generating ultrafilters in a reasonable way, Mathematical Logic Quarterly **54**, 202–220 (2008).
- [9] A. Roslanowski and S. Shelah, Reasonable ultrafilters, again, Notre Dame Journal of Formal Logic **52**, 113–147 (2011).
- [10] S. Shelah, Properness Without Elementaricity, Journal of Applied Analysis **10**, 168–289 (2004).

EINSTEIN INSTITUTE OF MATHEMATICS, EDMOND J. SAFRA CAMPUS, GIVAT RAM, THE HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM, 91904, ISRAEL, AND, DEPARTMENT OF MATHEMATICS, HILL CENTER - BUSCH CAMPUS, RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY, 110 FRELINGHUYSEN ROAD, PISCATAWAY, NJ 08854-8019 USA

E-mail address: shelah@math.huji.ac.il

URL: <http://shelah.logic.at>