

# COMPARING THE CLOSED ALMOST DISJOINTNESS AND DOMINATING NUMBERS

DILIP RAGHAVAN AND SAHARON SHELAH

ABSTRACT. We prove that if there is a dominating family of size  $\aleph_1$ , then there is are  $\aleph_1$  many compact subsets of  $\omega^\omega$  whose union is a maximal almost disjoint family of functions that is also maximal with respect to infinite partial functions.

## 1. INTRODUCTION

Recall that two infinite subsets  $a$  and  $b$  of  $\omega$  are *almost disjoint* or *a.d.* if  $a \cap b$  is finite. A family  $\mathcal{A}$  of infinite subsets of  $\omega$  is said to be *almost disjoint* or *a.d.* in  $[\omega]^\omega$  if its members are pairwise almost disjoint. A *Maximal Almost Disjoint family*, or *MAD family* in  $[\omega]^\omega$  is an infinite a.d. family in  $[\omega]^\omega$  that is not properly contained in a larger a.d. family.

Two functions  $f$  and  $g$  in  $\omega^\omega$  are said to be *almost disjoint* or *a.d.* if they agree in only finitely many places. We say that a family  $\mathcal{A} \subset \omega^\omega$  is *a.d. in  $\omega^\omega$*  if its members are pairwise a.d., and we say that an a.d. family  $\mathcal{A} \subset \omega^\omega$  is *MAD in  $\omega^\omega$*  if  $\forall f \in \omega^\omega \exists h \in \mathcal{A} [|f \cap h| = \aleph_0]$ . Identifying functions with their graphs, every a.d. family in  $\omega^\omega$  is also an a.d. family in  $[\omega \times \omega]^\omega$ ; however, it is never MAD in  $[\omega \times \omega]^\omega$  because any function is a.d. from the vertical columns of  $\omega \times \omega$ . MAD families in  $\omega^\omega$  that become MAD in  $[\omega \times \omega]^\omega$  when the vertical columns of  $\omega \times \omega$  are thrown in were considered by Van Douwen.

We say that  $p \subset \omega \times \omega$  is an *infinite partial function* if it is a function from some infinite set  $A \subset \omega$  to  $\omega$ . An a.d. family  $\mathcal{A} \subset \omega^\omega$  is said to be *Van Douwen* if for any infinite partial function  $p$  there is  $h \in \mathcal{A}$  such that  $|h \cap p| = \aleph_0$ .  $\mathcal{A}$  is Van Douwen iff  $\mathcal{A} \cup \{c_n : n \in \omega\}$  is a MAD family in  $[\omega \times \omega]^\omega$ , where  $c_n$  is the  $n$ th vertical column of  $\omega \times \omega$ . The first author showed in [3] that Van Douwen MAD families always exist.

Recall that  $\mathfrak{b}$  is the least size of an unbounded family in  $\omega^\omega$ ,  $\mathfrak{d}$  is the least size of a dominating family in  $\omega^\omega$ , and  $\mathfrak{a}$  is the least size of a MAD family in  $[\omega]^\omega$ . It is well known that  $\mathfrak{b} \leq \mathfrak{a}$ . Whether  $\mathfrak{a}$  could consistently be larger than  $\mathfrak{d}$  was an open question for a long time, until Shelah achieved a breakthrough in [4] by producing a model where  $\mathfrak{d} = \aleph_2$  and  $\mathfrak{a} = \aleph_3$ . However, it is not known whether  $\mathfrak{a}$  can be larger than  $\mathfrak{d}$  when  $\mathfrak{d} = \aleph_1$ ; this is one of the few major remaining open problems in the theory of cardinal invariants posed during the earliest days of the subject

---

*Date:* October 12, 2011.

*2010 Mathematics Subject Classification.* 03E35, 03E65, 03E17, 03E05.

*Key words and phrases.* maximal almost disjoint family, dominating family.

First author partially supported by Grants-in-Aid for Scientific Research for JSPS Fellows No. 23-01017.

Research partially supported by NSF grant DMS 1101597, and by the United States-Israel Binational Science Foundation (Grant no. 2006108). Publication 991.

(see [5] and [2]). In this note we take a small step towards resolving this question by showing that if  $\mathfrak{d} = \aleph_1$ , then there is a MAD family in  $[\omega]^\omega$  which is the union of  $\aleph_1$  compact subsets of  $[\omega]^\omega$ . More precisely, we will establish the following:

**Theorem 1.** *Assume  $\mathfrak{d} = \aleph_1$ . Then there exist  $\aleph_1$  compact subsets of  $\omega^\omega$  whose union is a Van Douwen MAD family.*

The cardinal invariant  $\mathfrak{a}_{closed}$  was recently introduced and studied by Brendle and Khomskii [1] in connection with the possible descriptive complexities of MAD families in certain forcing extensions of  $\mathbf{L}$ .

**Definition 2.**  $\mathfrak{a}_{closed}$  is the least  $\kappa$  such that there are  $\kappa$  closed subsets of  $[\omega]^\omega$  whose union is a MAD family in  $[\omega]^\omega$ .

Obviously,  $\mathfrak{a}_{closed} \leq \mathfrak{a}$ . Brendle and Khomskii showed in [1] that  $\mathfrak{a}_{closed}$  behaves differently from  $\mathfrak{a}$  by producing a model where  $\mathfrak{a}_{closed} = \aleph_1 < \aleph_2 = \mathfrak{b}$ . They asked whether  $\mathfrak{s} = \aleph_1$  implies that  $\mathfrak{a}_{closed} = \aleph_1$ . As  $\mathfrak{s} \leq \mathfrak{d}$ , our result in this paper provides a partial positive answer to their question.

## 2. THE CONSTRUCTION

Assume  $\mathfrak{d} = \aleph_1$  in this section. We will build  $\aleph_1$  many compact subsets of  $\omega^\omega$  whose union is a Van Douwen MAD family. To this end, we will construct a sequence  $\langle T_\alpha : \alpha < \omega_1 \rangle$  of finitely branching subtrees of  $\omega^{<\omega}$  such that  $\bigcup_{\alpha < \omega_1} [T_\alpha]$  has the required properties. Henceforth,  $T \subset \omega^{<\omega}$  will mean  $T$  is a subtree of  $\omega^{<\omega}$ .

**Definition 3.** Let  $T \subset \omega^{<\omega}$ . Let  $A \in [\omega]^\omega$  and  $p : A \rightarrow \omega$ . For any ordinal  $\xi$  and  $\sigma \in T$  define  $\text{rk}_{T,p}(\sigma) \geq \xi$  to mean

$$\forall \zeta < \xi \exists \tau \in T \exists l \in A [\tau \supset \sigma \wedge |\sigma| \leq l < |\tau| \wedge \tau(l) = p(l) \wedge \text{rk}_{T,p}(\tau) \geq \zeta].$$

Note that if  $\eta \leq \xi$  and  $\text{rk}_{T,p}(\sigma) \geq \xi$ , then  $\text{rk}_{T,p}(\sigma) \geq \eta$ , and that for a limit ordinal  $\xi$ , if  $\forall \zeta < \xi [\text{rk}_{T,p}(\sigma) \geq \zeta]$ , then  $\text{rk}_{T,p}(\sigma) \geq \xi$ . Also, for any  $\sigma, \tau \in T$ , if  $\sigma \subset \tau$  and  $\text{rk}_{T,p}(\tau) \geq \xi$ , then  $\text{rk}_{T,p}(\sigma) \geq \xi$ . Moreover, if  $\text{rk}_{T,p}(\sigma) \not\geq \xi$  and if  $\tau \in T$  and  $l \in A$  are such that  $\tau \supset \sigma$ ,  $|\sigma| \leq l < |\tau|$ , and  $p(l) = \tau(l)$ , then there is  $\zeta < \xi$  such that  $\text{rk}_{T,p}(\tau) \not\geq \zeta$ . Therefore, if there is  $f \in [T]$  with  $|f \cap p| = \aleph_0$ , and if  $\sigma \subset f$  and there is some ordinal  $\xi$  such that  $\text{rk}_{T,p}(\sigma) \not\geq \xi$ , then there is some  $\sigma \subset \tau \subset f$  and some ordinal  $\zeta < \xi$  such that  $\text{rk}_{T,p}(\tau) \not\geq \zeta$ , thus allowing us to construct an infinite, strictly descending sequence of ordinals. So if  $f \in [T]$  with  $|f \cap p| = \aleph_0$ , then for any  $\sigma \subset f$  and any ordinal  $\xi$ ,  $\text{rk}_{T,p}(\sigma) \geq \xi$ . On the other hand, suppose that  $\sigma \in T$  with  $\text{rk}_{T,p}(\sigma) \geq \omega_1$ . Then there is  $\tau \in T$  with  $\tau \supset \sigma$  and  $l \in A$  such that  $|\sigma| \leq l < |\tau|$ ,  $p(l) = \tau(l)$ , and  $\text{rk}_{T,p}(\tau) \geq \omega_1$ , allowing us to construct  $f \in [T]$  with  $\sigma \subset f$  such that  $|f \cap p| = \aleph_0$ .

**Definition 4.** Suppose  $T \subset \omega^{<\omega}$ ,  $A \in [\omega]^\omega$ , and  $p : A \rightarrow \omega$ . Assume that  $p$  is a.d. from each  $f \in [T]$ . Then define  $H_{T,p} : T \rightarrow \omega_1$  by  $H_{T,p}(\sigma) = \min\{\xi : \text{rk}_{T,p}(\sigma) \not\geq \xi + 1\}$ .

Note the following features of this definition

- (\*)  $\forall \sigma, \tau \in T [\sigma \subset \tau \implies H_{T,p}(\sigma) \geq H_{T,p}(\tau)]$
- (\*\*) for all  $\sigma, \tau \in T$  with  $\sigma \subset \tau$ , if there exists  $l \in A$  such that  $|\sigma| \leq l < |\tau|$  and  $p(l) = \tau(l)$ , then  $H_{T,p}(\tau) < H_{T,p}(\sigma)$ .

On the other hand, notice that if there is a function  $H : T \rightarrow \omega_1$  such that (\*) and (\*\*) hold when  $H_{T,p}$  is replaced with  $H$ , then  $p$  must be a.d. from  $[T]$ .

**Definition 5.**  $I$  is said to be an *interval partition* if  $I = \langle i_n : n \in \omega \rangle$ , where  $i_0 = 0$ , and  $\forall n \in \omega [i_n < i_{n+1}]$ . For  $n \in \omega$ ,  $I_n$  denotes the interval  $[i_n, i_{n+1})$ .

Given two interval partitions  $I$  and  $J$ , we say that  $I$  *dominates*  $J$  and write  $J \leq^* I$  if  $\forall^\infty n \in \omega \exists k \in \omega [J_k \subset I_n]$ .

It is well known that  $\mathfrak{d}$  is also the size of the smallest family of interval partitions dominating any interval partition. So fix a sequence  $\langle I^\alpha : \alpha < \omega_1 \rangle$  of interval partitions such that

- (1)  $\forall \alpha \leq \beta < \omega_1 [I^\alpha \leq^* I^\beta]$
- (2) for any interval partition  $J$ , there exists  $\alpha < \omega_1$  such that  $J \leq^* I^\alpha$ .

Fix an  $\omega_1$ -scale  $\langle f_\alpha : \alpha < \omega_1 \rangle$  such that  $\forall \alpha < \omega_1 \forall n \in \omega [f_\alpha(n) < f_\alpha(n+1)]$ . For each  $\alpha \geq 1$ , define  $e_\alpha$  and  $g_\alpha$  by induction on  $\alpha$  as follows. If  $\alpha$  is a successor, then  $e_\alpha : \omega \rightarrow \alpha$  is any onto function, and  $g_\alpha = f_\alpha$ . If  $\alpha$  is a limit, then let  $\{e_n : n \in \omega\}$  enumerate  $\{e_\xi : \xi < \alpha\}$ . Now, define  $e_\alpha : \omega \rightarrow \alpha$  and  $g_\alpha \in \omega^\omega$  such that

- (3)  $\forall n \in \omega [g_\alpha(n) \leq g_\alpha(n+1)]$
- (4)  $\forall n \in \omega \forall i \leq n \forall j \leq f_\alpha(n) \exists k < g_\alpha(n) [e_\alpha(k) = e_i(j)]$ .

Observe that such an  $e_\alpha$  must be a surjection. For each  $n \in \omega$ , put  $w_\alpha(n) = \{e_\alpha(i) : i \leq g_\alpha(n)\}$ .

Now fix  $\alpha < \omega_1$  and assume that  $T_\epsilon \subset \omega^{<\omega}$  has been defined for each  $\epsilon < \alpha$  such that each  $T_\epsilon$  is finitely branching and  $\bigcup_{\epsilon < \alpha} [T_\epsilon]$  is an a.d. family in  $\omega^\omega$ . Let  $\langle \epsilon_n : n \in \omega \rangle$  enumerate  $\alpha$ , possibly with repetitions. For a tree  $T \subset \omega^{<\omega}$  and  $l \in \omega$ ,  $T \upharpoonright l$  denotes  $\{\sigma \in T : |\sigma| \leq l\}$ , and  $T(l)$  denotes  $\{\sigma \in T : |\sigma| = l\}$ . We will define a sequence of natural numbers  $0 = l_0 < l_1 < \dots$  and determine  $T_\alpha \upharpoonright l_n$  by induction on  $n$ .  $T_\alpha \upharpoonright l_0 = \{0\}$ . Assume that  $l_n$  and  $T_\alpha \upharpoonright l_n$  are given. Suppose also that we are given a sequence of natural numbers  $\langle k_i : i < n \rangle$  such that

- (5)  $\forall i < i+1 < n [k_i < k_{i+1}]$
- (6)  $I_{k_i}^\alpha \subset [0, l_n)$ .

Let  $\sigma^*$  denote the member of  $T_\alpha(l_n)$  that is right most with respect to the lexicographical ordering on  $\omega^{l_n}$ . Suppose we are also given  $L_n : T_\alpha(l_n) \setminus \{\sigma^*\} \rightarrow W_n$ , an injection. Here  $W_n$  is the set of all pairs  $\langle p_0, \bar{h} \rangle$  such that

- (7) there are  $s \in [\omega]^{<\omega}$ , and numbers  $i_0 < j_0 \leq n$  such that
  - (a)  $s \subset \bigcup_{i \in [i_0, j_0)} I_{k_i}^\alpha$
  - (b) for each  $i \in [i_0, j_0)$ ,  $|s \cap I_{k_i}^\alpha| = 1$
  - (c)  $p_0 : s \rightarrow \omega$  such that  $\forall m \in s [p_0(m) \leq f_\alpha(m)]$
- (8) There is  $j_1 < n$  such that  $\bar{h} = \langle h_{\epsilon_i} : i \leq j_1 \rangle$  (if  $\alpha = 0$ , this means that  $\bar{h} = 0$ ). For each  $i \leq j_1$ ,  $h_{\epsilon_i} : T_{\epsilon_i} \upharpoonright \max(s) + 1 \rightarrow w_\alpha(\max(s) + 1)$  such that  $(*_1)$  and  $(*_2)$  hold when  $T$  is replaced there with  $T_{\epsilon_i} \upharpoonright \max(s) + 1$ ,  $H_{T,p}$  is replaced with  $h_{\epsilon_i}$ ,  $A$  with  $s$ , and  $p$  with  $p_0$ .

Assume that for each  $i < n$ , we are also given  $\sigma_i \in T_\alpha(l_i)$ , which we will call *the active node at stage  $i$* . Note that  $T_\alpha(l_0) = \{0\}$ , and so  $\sigma_0 = 0$ . For each  $\sigma \in T_\alpha(l_n)$ , let  $\Delta(\sigma) = \max(\{0\} \cup \{i < n : \sigma_i = \sigma \upharpoonright l_i\})$ . For,  $\sigma, \tau \in T_\alpha(l_n)$ , say  $\sigma \triangleleft \tau$  if either  $\Delta(\sigma) < \Delta(\tau)$  or  $\Delta(\sigma) = \Delta(\tau)$  and  $\sigma$  is to the left of  $\tau$  in the lexicographic ordering on  $\omega^{l_n}$ . Let  $\sigma_n$  be the  $\triangleleft$ -minimal member of  $T_\alpha(l_n)$ .  $\sigma_n$  will be active at stage  $n$ . The meaning of this is that none of the other nodes in  $T_\alpha(l_n)$  will be allowed to branch at stage  $n$ . Choose  $k_n$  greater than all  $k_i$  for  $i < n$  such that  $I_{k_n}^\alpha \subset [l_n, \infty)$ . Let  $V_n$  be the set of all pairs  $\langle p_1, \bar{h} \rangle$  such that

- (9) there exist  $s$  and a natural number  $i_1 \leq n$  such that

modified:2012-03-12

- (a)  $s \subset \bigcup_{i \in [i_1, n+1)} I_{k_i}^\alpha$
  - (b) for each  $i \in [i_1, n+1)$ ,  $|s \cap I_{k_i}^\alpha| = 1$
  - (c)  $p_1 : s \rightarrow \omega$  such that  $\forall m \in s [p_1(m) \leq f_\alpha(m)]$
- (10) There is  $j_2 \leq n$  such that  $\bar{\mathbf{h}} = \langle \mathbf{h}_{\epsilon_i} : i \leq j_2 \rangle$ . For each  $i \leq j_2$ ,  $\mathbf{h}_{\epsilon_i} : T_{\epsilon_i} \upharpoonright \max(s) + 1 \rightarrow w_\alpha(\max(s) + 1)$  such that  $(*_1)$  and  $(*_2)$  are satisfied when  $T$  is replaced with  $T_{\epsilon_i} \upharpoonright \max(s) + 1$ ,  $H_{T,p}$  is replaced with  $\mathbf{h}_{\epsilon_i}$ ,  $A$  with  $s$ , and  $p$  with  $p_1$ .

Note that  $V_n$  is always finite. Now, the construction splits into two cases.

Case I:  $\sigma_n \neq \sigma^*$ . Put  $\langle p_0, \bar{h} \rangle = L_n(\sigma_n)$ . Let  $i_0 < n$  be as in (7) above, and let  $j_1 < n$  be as in (8). Let

$$U_n = \{ \langle p_1, \bar{\mathbf{h}} \rangle \in V_n : p_0 \subset p_1 \wedge i_0 = i_1 \wedge j_1 < j_2 \wedge \forall i \leq j_1 [\mathbf{h}_{\epsilon_i} \upharpoonright \text{dom}(h_{\epsilon_i}) = h_{\epsilon_i}] \}.$$

Here  $i_1$  is as in (9), and  $j_2$  is as in (10) with respect to  $\langle p_1, \bar{\mathbf{h}} \rangle$ . Now choose  $l_{n+1} > l_n$  large enough so that  $I_{k_n}^\alpha \subset [l_n, l_{n+1})$  and so that it is possible to pick  $\{ \tau_x : x \in U_n \} \subset \omega^{l_{n+1}}$  and  $\{ \tau_\sigma : \sigma \in T_\alpha(l_n) \} \subset \omega^{l_{n+1}}$  such that the following conditions are satisfied.

- (11) for each  $x \in U_n$ ,  $\tau_x \supset \sigma_n$ , and for each  $\sigma \in T_\alpha(l_n)$ ,  $\tau_\sigma \supset \sigma$
- (12) for each  $x, y \in U_n$ , if  $x \neq y$ , then there exists  $m \in [l_n, l_{n+1})$  such that  $\tau_x(m) \neq \tau_y(m)$ . For each  $x \in U_n$ , there exists  $m \in [l_n, l_{n+1})$  such that  $\tau_x(m) \neq \tau_{\sigma_n}(m)$ . For  $x = \langle p_1, \bar{\mathbf{h}} \rangle \in U_n$ , if  $\{i^*\} = \text{dom}(p_1) \cap I_{k_n}^\alpha$ , then  $p_1(i^*) = \tau_x(i^*)$ .
- (13) for each  $x \in U_n$  and  $\sigma \in T_\alpha(l_n)$ ,  $\forall m \in [l_n, l_{n+1}) [\tau_x(m) \neq \tau_\sigma(m)]$ . For  $\sigma, \eta \in T_\alpha(l_n)$ , if  $\sigma \neq \eta$ , then  $\forall m \in [l_n, l_{n+1}) [\tau_\sigma(m) \neq \tau_\eta(m)]$ .
- (14) for each  $i \leq n$ ,  $\tau \in T_{\epsilon_i}(l_{n+1})$ ,  $\sigma \in T_\alpha(l_n)$  and  $m \in [l_n, l_{n+1})$ ,  $\tau(m) \neq \tau_\sigma(m)$ . For each  $x \in U_n$ ,  $i \leq j_2$ ,  $\tau \in T_{\epsilon_i}(l_{n+1})$  and  $m \in [l_n, l_{n+1})$ , if  $\tau_x(m) = \tau(m)$ , then  $m \in \text{dom}(p_1)$  and  $p_1(m) = \tau_x(m)$ .

Define  $L_{n+1}$  as follows. For any  $x \in U_n$ ,  $L_{n+1}(\tau_x) = x$ . For any  $\sigma \in T_\alpha(l_n) \setminus \{\sigma^*\}$ ,  $L_{n+1}(\tau_\sigma) = L_n(\sigma)$ . This finishes case I.

Case II:  $\sigma_n = \sigma^*$ . For each  $\sigma \in T_\alpha(l_n) \setminus \{\sigma_n\}$ , let  $\langle p_0(\sigma), \bar{h}(\sigma) \rangle = L_n(\sigma)$ . Let  $i_0(\sigma) < n$  witness (7) for  $L_n(\sigma)$  and let  $j_1(\sigma) < n$  witness (8) for  $L_n(\sigma)$ . Let  $U_n$  be the set of all  $\langle p_1, \bar{\mathbf{h}} \rangle \in V_n$  such that there is no  $\sigma \in T_\alpha(l_n) \setminus \{\sigma_n\}$  so that

$$p_0(\sigma) \subset p_1 \wedge i_0(\sigma) = i_1 \wedge j_1(\sigma) < j_2 \wedge \forall i \leq j_1(\sigma) [\mathbf{h}_{\epsilon_i} \upharpoonright \text{dom}(h_{\epsilon_i}) = h_{\epsilon_i}].$$

Here  $i_1 \leq n$  and  $j_2 \leq n$  witness (9) and (10) respectively with respect to  $\langle p_1, \bar{\mathbf{h}} \rangle$ . Choose  $l_{n+1} > l_n$  large enough so that  $I_{k_n}^\alpha \subset [l_n, l_{n+1})$  and so that it is possible to choose  $\{\tau^*\}$ ,  $\{\tau_x : x \in U_n\}$ , and  $\{\tau_\sigma : \sigma \in T_\alpha(l_n) \setminus \{\sigma_n\}\}$ , subsets of  $\omega^{l_{n+1}}$ , satisfying the following conditions.

- (15)  $\tau^* \supset \sigma_n$ . For each  $x \in U_n$ ,  $\tau_x \supset \sigma_n$ . For each  $\sigma \in T_\alpha(l_n) \setminus \{\sigma_n\}$ ,  $\tau_\sigma \supset \sigma$ .
- (16)  $\tau^*$  is the right most branch of  $T_\alpha(l_{n+1})$ . For each  $x \in U_n$ , there exists  $m \in [l_n, l_{n+1})$  such that  $\tau^*(m) \neq \tau_x(m)$ . For each  $x, y \in U_n$ , if  $x \neq y$ , then there is  $m \in [l_n, l_{n+1})$  so that  $\tau_x(m) \neq \tau_y(m)$ . For each  $x = \langle p_1, \bar{\mathbf{h}} \rangle \in U_n$ , if  $\{i^*\} = I_{k_n}^\alpha \cap \text{dom}(p_1)$ , then  $p_1(i^*) = \tau_x(i^*)$ .
- (17) For each  $x \in U_n$  and  $m \in [l_n, l_{n+1})$ ,  $\tau_x(m) \neq \tau^*(m)$ . For each  $\sigma \in T_\alpha(l_n) \setminus \{\sigma_n\}$  and for each  $m \in [l_n, l_{n+1})$ ,  $\tau^*(m) \neq \tau_\sigma(m)$ , and for each  $x \in U_n$ ,  $\tau_\sigma(m) \neq \tau_x(m)$ . For each  $\sigma, \eta \in T_\alpha(l_n) \setminus \{\sigma_n\}$ , if  $\sigma \neq \eta$ , then for all  $m \in [l_n, l_{n+1})$ ,  $\tau_\sigma(m) \neq \tau_\eta(m)$ .

- (18) For each  $i \leq n$ ,  $\tau \in T_{\epsilon_i}(l_{n+1})$ ,  $m \in [l_n, l_{n+1})$ , and  $\sigma \in T_\alpha(l_n) \setminus \{\sigma_n\}$ ,  $\tau^*(m) \neq \tau(m)$  and  $\tau_\sigma(m) \neq \tau(m)$ . For each  $x = \langle p_1, \bar{\mathbf{h}} \rangle \in U_n$ ,  $i \leq j_2$ ,  $\tau \in T_{\epsilon_i}(l_{n+1})$  and  $m \in [l_n, l_{n+1})$ , if  $\tau_x(m) = \tau(m)$ , then  $m \in \text{dom}(p_1)$  and  $p_1(m) = \tau_x(m)$ .

For each  $\sigma \in T_\alpha(l_n) \setminus \{\sigma_n\}$ , define  $L_{n+1}(\tau_\sigma) = L_n(\sigma)$ . For each  $x \in U_n$ , set  $L_{n+1}(\tau_x) = x$ . This completes the construction. We now check that it is as required.

**Lemma 6.** *For each  $f \in [T_\alpha]$ , there are infinitely many  $n \in \omega$  such that  $\sigma_n = f \upharpoonright l_n$ .*

*Proof.* For each  $n \in \omega$  put  $\Theta(n) = \min\{\Delta(\sigma) : \sigma \in T_\alpha(l_n)\}$ . It is clear from the construction that  $\Theta(n+1) \geq \Theta(n)$ . If the lemma fails, then there are  $m$  and  $\tau \in T_\alpha(l_{m+1})$  with the property that for infinitely many  $n > m+1$ , there is a  $\sigma \in T_\alpha(l_n)$  such that  $\Theta(n) = \Delta(\sigma) = m$  and  $\sigma \upharpoonright l_{m+1} = \tau$ . Let  $\tau$  be the left most node in  $T_\alpha(l_{m+1})$  with this property. Choose  $n_1 > n_0 > m+1$  and  $\sigma \in T_\alpha(l_{n_1})$  such that  $\Theta(n_1) = \Theta(n_0) = \Delta(\sigma) = m$ ,  $\sigma \upharpoonright l_{m+1} = \tau$ , and there is no  $\eta \in T_\alpha(l_{n_0})$  such that  $\Delta(\eta) = m$  and  $\eta \upharpoonright l_{m+1}$  is to the left of  $\tau$ . Note that  $\Delta(\sigma \upharpoonright l_{n_0}) = m$ . So  $\sigma_{n_0}$  is to the left of  $\sigma \upharpoonright l_{n_0}$ , and  $\sigma_{n_0} \upharpoonright l_{m+1}$  is not to the left of  $\tau$ , whence  $\sigma_{n_0} \upharpoonright l_{m+1} = \tau$ . But then there is some  $n \in [m+1, n_0)$  where  $\sigma \upharpoonright l_n$  was active, a contradiction.  $\dashv$

Note that Lemma 6 implies that for any  $\sigma \in T_\alpha$ , there is a unique minimal extension of  $\sigma$  which is active. Lemma 6 also implies that there are infinitely many  $n$  where case II occurs.

**Lemma 7.**  *$T_\alpha$  is finitely branching and  $\bigcup_{\epsilon \leq \alpha} [T_\epsilon]$  is a.d. in  $\omega^\omega$ .*

*Proof.* It is clear from the construction that  $T_\alpha$  is finitely branching. Fix  $f, g \in [T_\alpha]$ , with  $f \neq g$ . Let  $n = \max\{i \in \omega : f \upharpoonright l_i = g \upharpoonright l_i\}$ . It is clear from the construction that  $\forall m \geq l_{n+1} [f(m) \neq g(m)]$ .

Next, fix  $\epsilon < \alpha$ . Suppose  $\epsilon = \epsilon_i$ . Let  $h \in [T_{\epsilon_i}]$  and  $f \in [T_\alpha]$ , and suppose for a contradiction that  $|h \cap f| = \aleph_0$ . So there are infinitely many  $n \in \omega$  such that  $f \upharpoonright [l_n, l_{n+1}) \cap h \upharpoonright [l_n, l_{n+1}) \neq \emptyset$ . For any  $n \geq i$ , this can only happen if  $f \upharpoonright l_n = \sigma_n$  and  $f \upharpoonright l_{n+1} = \tau_{x_n}$  for some  $x_n \in U_n$ . This is because if  $n \geq i$  and  $f \upharpoonright [l_n, l_{n+1}) \cap h \upharpoonright [l_n, l_{n+1}) \neq \emptyset$ , then when case I occurs, (14) says that  $f \upharpoonright l_{n+1} \neq \tau_\sigma$  for any  $\sigma \in T_\alpha(l_n)$ , while when case II occurs, by (18),  $f \upharpoonright l_{n+1} \neq \tau^*$  and also  $f \upharpoonright l_{n+1} \neq \tau_\sigma$  for any  $\sigma \in T_\alpha(l_n) \setminus \{\sigma_n\}$ . So  $f \upharpoonright l_{n+1} = \tau_{x_n}$  for some  $x_n \in U_n$ , and  $f \upharpoonright l_n = \sigma_n$ . Now, put  $x_n = \langle p_{1,n}, \bar{\mathbf{h}}_n \rangle$ . Note that in this case  $L_{n+1}(f \upharpoonright l_{n+1}) = x_n$ . For such  $n$ , let  $j_2(n)$  be as in (10) with respect to  $x_n$ . So for infinitely many such  $n$ ,  $j_2(n) \geq i$ . But then for infinitely many such  $n$ ,  $\mathbf{h}_{\epsilon_i, n}(h \upharpoonright \max(\text{dom}(p_{1,n})) + 1) < \mathbf{h}_{\epsilon_i, n}(h \upharpoonright l_n)$ , producing an infinite strictly descending sequence of ordinals.  $\dashv$

**Lemma 8.** *For each  $A \in [\omega]^\omega$  and  $p : A \rightarrow \omega$ , there are  $\alpha < \omega_1$  and  $f \in [T_\alpha]$  such that  $|p \cap f| = \aleph_0$ .*

*Proof.* Suppose for a contradiction that there are  $A \in [\omega]^\omega$  and  $p : A \rightarrow \omega$  such that  $p$  is a.d. from  $[T_\alpha]$ , for each  $\alpha < \omega_1$ . Let  $M \prec H(\theta)$  be a countable elementary submodel containing everything relevant. Put  $\alpha = M \cap \omega_1$ . For each  $\epsilon < \alpha$ , let  $H_\epsilon$  denote  $H_{T_\epsilon, p}$ , and note that  $H_\epsilon$  and  $\text{ran}(H_\epsilon)$  are members of  $M$ . Let  $\xi_\epsilon = \sup(\text{ran}(H_\epsilon)) + 1 < \alpha$ . Find  $g \in M \cap \omega^\omega$  such that for  $n \in \omega$ ,  $H_\epsilon'' T_\epsilon \upharpoonright n \subset \{e_{\xi_\epsilon}(j) : j \leq g(n)\}$ . Since  $\forall^\infty n \in \omega [g(n) \leq f_\alpha(n)]$ , it follows from (4) that for all but finitely many  $n \in \omega$ , for all  $\sigma \in T_\epsilon \upharpoonright n$ ,  $H_\epsilon(\sigma) \in w_\alpha(n)$ . Now, find an infinite  $q \subset p$  such that  $\forall m \in \text{dom}(q) [q(m) \leq f_\alpha(m)]$  and  $\forall^\infty n \in \omega [|\text{dom}(q) \cap I_n^\alpha| = 1]$ .

Note that for any  $\epsilon < \alpha$ ,  $(*_1)$  and  $(*_2)$  are satisfied when  $T$  is replaced there with  $T_\epsilon$ ,  $H_{T,p}$  is replaced with  $H_{\epsilon_j}$ ,  $A$  with  $\text{dom}(q)$ , and  $p$  with  $q$ . But now, it follows from the construction that there is  $f \in [T_\alpha]$  such that for infinitely many  $n \in \omega$ , there is  $m \in [l_n, l_{n+1}) \cap \text{dom}(q)$  such that  $q(m) = f(m)$ . We describe how to find such a  $f \in [T_\alpha]$ .  $\forall^\infty n \in \omega$   $[|\text{dom}(q) \cap I_{k_n}^\alpha| = 1]$ . For each  $n \in \omega$  such that  $|\text{dom}(q) \cap I_{k_n}^\alpha| = 1$ , let  $m_n$  be the unique member of  $\text{dom}(q) \cap I_{k_n}^\alpha$ . We observed above that for any  $\epsilon < \alpha$ , for all but finitely many  $n \in \omega$ , for each  $\sigma \in T_\epsilon \upharpoonright n$ ,  $H_\epsilon(\sigma) \in w_\alpha(n)$ . It follows that for any  $i \in \omega$ , there is  $u_i \geq i$  such that for each  $j \leq i$  and each  $n \geq u_i$ ,  $m_n$  is defined and  $\forall \sigma \in T_{\epsilon_j} \upharpoonright m_n + 1$   $[H_{\epsilon_j}(\sigma) \in w_\alpha(m_n + 1)]$ . Choose  $n^* \geq u_0$  so that case II occurs at stage  $n^*$ . Put  $\eta_0 = \sigma_{n^*}$ . Define  $s_0 = \{m_{n^*}\}$  and  $q_0 = q \upharpoonright s_0$ . Put  $\bar{h}_0 = \langle h_0 \rangle$ , where  $h_0 = H_{\epsilon_0} \upharpoonright (T_{\epsilon_0} \upharpoonright \max(s_0) + 1)$ . Note that  $h_0$  is a map from  $T_{\epsilon_0} \upharpoonright \max(s_0) + 1$  to  $w_\alpha(\max(s_0) + 1)$ , and so  $x_0 = \langle q_0, \bar{h}_0 \rangle \in V_{n^*}$ . Since  $m_{n^*} \notin I_{k_i}^\alpha$  for any  $i < n^*$ , it follows that  $x_0 \in U_{n^*}$ . Put  $\eta_1 = \tau_{x_0} \supseteq \eta_0$ . Notice that  $\eta_1(m_{n^*}) = q(m_{n^*})$ . Notice also that  $\eta_1$  is not the right most branch of  $T_\alpha(l_{(n^*+1)})$ , and so if  $\sigma$  is any extension of  $\eta_1$  that happens to be active at a certain stage, then case I necessarily occurs at that stage. Finally, note that  $L_{n^*+1}(\eta_1) = x_0$ . Now, for each  $n > n^*$ , let  $s_n = \{m_j : n^* \leq j \leq n\}$ , and put  $q_n = q \upharpoonright s_n$ . For any  $i > 0$  and  $n > n^*$ , if  $n \geq u_i$ , then for each  $j \leq i$ , define  $h_j^n = H_{\epsilon_j} \upharpoonright (T_{\epsilon_j} \upharpoonright \max(s_n) + 1)$ . Put  $\bar{h}_i^n = \langle h_j^n : j \leq i \rangle$  and  $x_i^n = \langle q_n, \bar{h}_i^n \rangle$ . Note that for any  $i > 0$  and  $n > n^*$ , if  $n \geq u_i$ , then  $x_i^n \in V_n$ . Moreover, if at stage  $n$ , case I occurs and  $L_n(\sigma_n) = x_{i-1}^v$  for some  $v \in \omega$ , then  $x_i^n \in U_n$ ; here  $x_0^v = x_0$ , for all  $v \in \omega$ . Now, it is easy to see that there is a branch  $g \in [T_\alpha]$  such that  $\eta_1 \subset g$  and  $\forall n \geq n^* + 1$   $[L_n(g \upharpoonright l_n) = x_0]$ . This is because for any  $n \geq n^* + 1$ , given  $g \upharpoonright l_n$  such that  $\eta_1 \subset g \upharpoonright l_n$  and  $L_n(g \upharpoonright l_n) = x_0$ , if  $\sigma$  is the unique minimal extension of  $g \upharpoonright l_n$  that is active, then  $\tau_\sigma \supseteq g \upharpoonright l_n$  and  $L_{u+1}(\tau_\sigma) = x_0$ , where  $u$  is the stage at which  $\sigma$  is active. Applying Lemma 6 to  $g$ , find  $n^{**}$  such that  $n^{**} > n^*$ ,  $n^{**} \geq u_1$ , and  $\sigma_{n^{**}} = g \upharpoonright l_{n^{**}}$ . It follows that  $x_1^{n^{**}} \in U_{n^{**}}$ . Let  $\eta_2 = \tau_{x_1^{n^{**}}} \supseteq \eta_1$ . Note that  $\eta_2(m_{n^{**}}) = q(m_{n^{**}})$  and that  $L_{n^{**}+1}(\eta_2) = x_1^{n^{**}}$ . Continuing in this fashion, we get  $f = \bigcup_{n \in \omega} \eta_n \in [T_\alpha]$  with  $|f \cap q| = \omega$ .  $\dashv$

### 3. REMARKS AND QUESTIONS

The construction in this paper is very specific to  $\omega_1$ ; indeed, it is possible to show that  $\mathfrak{d}$  is not always an upper bound for  $\mathfrak{a}_{closed}$ . A modification of the methods of Section 4 of [4] shows that if  $\kappa$  is a measurable cardinal and if  $\lambda = \text{cf}(\lambda) = \lambda^\kappa > \mu = \text{cf}(\mu) > \kappa$ , then there is a c.c.c. poset  $\mathbb{P}$  such that  $|\mathbb{P}| = \lambda$ , and  $\mathbb{P}$  forces that  $\mathfrak{b} = \mathfrak{d} = \mu$  and  $\mathfrak{a} = \mathfrak{a}_{closed} = \mathfrak{c} = \lambda$ .

As mentioned in Section 1, we see the result in this paper as providing a weak positive answer to the following basic question, which has remained open for long.

**Question 9.** If  $\mathfrak{d} = \aleph_1$ , then is  $\mathfrak{a} = \aleph_1$ ?

There are also several open questions about upper and lower bounds for  $\mathfrak{a}_{closed}$ .

**Question 10** (Brendle and Khomskii [1]). If  $\mathfrak{s} = \aleph_1$ , then is  $\mathfrak{a}_{closed} = \aleph_1$ ?

**Question 11.** Is  $\mathfrak{h} \leq \mathfrak{a}_{closed}$ ?

Regarding Question 10, it is proved in Brendle and Khomskii [1] that if  $\mathbf{V}$  is any ground model satisfying CH, then any finite support iteration of Suslin c.c.c. posets in  $\mathbf{V}$  forces that  $\mathfrak{a}_{closed} = \aleph_1$ . It is well known that  $\mathbf{V}$  remains a splitting

family after such a finite support iteration of Suslin c.c.c. posets. Showing a positive answer to Question 10 would be an improvement of the result in this paper.

## REFERENCES

1. J. Brendle and Y. Khomskii,  $\aleph_1$ -perfect MAD families, In Preparation.
2. E. Pearl (ed.), *Open problems in topology. II*, Elsevier B. V., Amsterdam, 2007.
3. D. Raghavan, *There is a Van Douwen MAD family*, Trans. Amer. Math. Soc. **362** (2010), no. 11, 5879–5891.
4. S. Shelah, *Two cardinal invariants of the continuum ( $\mathfrak{d} < \mathfrak{a}$ ) and FS linearly ordered iterated forcing*, Acta Math. **192** (2004), no. 2, 187–223.
5. E. K. van Douwen, *The integers and topology*, Handbook of set-theoretic topology, North-Holland, Amsterdam, 1984, pp. 111–167.

GRADUATE SCHOOL OF SYSTEM INFORMATICS, KOBE UNIVERSITY, KOBE 657-8501, JAPAN.

*E-mail address:* `raghavan@math.toronto.edu`

*URL:* `http://www.math.toronto.edu/raghavan`