



# RAMSEY THEORY FOR HIGHLY CONNECTED MONOCHROMATIC SUBGRAPHS

J. BERGFALK<sup>1</sup>, M. HRUŠÁK<sup>1,\*</sup>,† and S. SHELAH<sup>2,3,‡</sup>

<sup>1</sup>Centro de Ciencias Matemáticas, Universidad Nacional Autónoma de México, Campus Morelia, Morelia, Michoacán 58089, México  
e-mail: jeffrey@matmor.unam.mx

<sup>2</sup>Einstein Institute of Mathematics, Edmond J. Safra Campus, The Hebrew University of Jerusalem, Givat Ram, Jerusalem, 91904, Israel  
e-mail: michael@matmor.unam.mx

<sup>3</sup>Department of Mathematics, Hill Center – Busch Campus, Rutgers, State University of New Jersey, 110 Frelinghuysen Road, Piscataway, NJ 08854-8019, USA

(Received February 29, 2020; revised April 9, 2020; accepted April 11, 2020)

**Abstract.** An infinite graph is highly connected if the complement of any subgraph of smaller size is connected. We consider weaker versions of Ramsey's Theorem asserting that in any coloring of the edges of a complete graph there exist large highly connected subgraphs all of whose edges are colored by the same color.

## 1. Introduction

Ramsey's celebrated theorem in its most basic infinite form is the following: for any partition of the collection of pairs of natural numbers into finitely many sets, there exists some infinite  $X \subseteq \mathbb{N}$  whose pairs all fall in one of those sets. A pithier rendering is by way of Erdős and Rado's arrow notation:

$$(\star) \quad \aleph_0 \rightarrow (\aleph_0)_k^2 \text{ for any finite } k.$$

Here the outer cardinals  $\aleph_0$ , 2, and  $k$  parametrize the sorts of partitions under consideration: letting  $[\mu]^\lambda$  denote the size- $\lambda$  subsets of  $\mu$ , the partitions

---

\* Corresponding author.

† The research of the second author was supported by a PAPIIT grant No. IN100317 and CONACyT grant No. A1-S-16164.

‡ The research of the third author was partially supported by NSF grant 136974 and by ERC grant 338821.

*Key words and phrases:* Ramsey theory,  $k$ -connected graph, highly connected graph, Mahlo cardinal, weakly compact cardinal.

*Mathematics Subject Classification:* 03E02, 03E10.

in question in the above relation are all of the form  $c: [\aleph_0]^2 \rightarrow k$ . The cardinal inside the parentheses records how large a homogeneous set we seek with respect to any such partition, and the arrow tells us we can always find one.

The following points are basic to the theory:

(1) The above relation “descends” to finite contexts (see [5, §1.5] for a direct deduction). More precisely, for any finite  $k$  and  $m$  there is some  $n$  such that

$$n \rightarrow (m)_k^2.$$

This least such  $n$  is often denoted  $R(m; k)$ .

(2) Extending the relation  $(\star)$  to higher cardinals is less straightforward. More precisely:

(a) The relation  $\aleph_1 \rightarrow (\aleph_1)_k^2$  fails in a very strong sense, for any  $k \geq 2$  (see [19]).

(b) More generally, for  $2 \leq \kappa < \mu$ , the relation  $\mu \rightarrow (\mu)_\kappa^2$  characterizes any uncountable cardinal  $\mu$  as *weakly compact*, that is, as a cardinal whose existence is a strictly stronger assumption than the ZFC axioms (see [9]).

(c) Item (1), on the other hand, does fully generalize: for any  $\kappa$  and  $\mu$  there is some least  $\nu$  such that

$$\nu \rightarrow (\mu)_\kappa^2$$

This follows from Erdős and Rado’s theorem that  $(2^\lambda)^+ \rightarrow (\lambda^+)_\lambda^2$  for any infinite  $\lambda$  (see [4]).

Each of these facts will figure in the following. Recall lastly the more pictorial framing of Ramsey’s relation in terms of edge-colorings of graphs:

$$\nu \rightarrow (\mu)_\lambda^2$$

if and only if every coloring of the edges of the complete graph on  $\nu$  by  $\lambda$  many colors contains some size- $\mu$  monochromatic subgraph which is complete. It is this framing we will have generally in mind — only our interest will be in subgraphs which are large in some finer sense than *complete*. Namely:

**DEFINITION 1.** A graph  $G$  is  $\kappa$ -*connected* if it remains connected after the deletion of any fewer than  $\kappa$  vertices.

Our question should at this point be clear. Where formality is necessary, we will denote a graph  $G$  as an ordered pair (*vertices, edges*). The *size* of a graph is the cardinality of its vertex-set. For cardinal numbers  $\kappa, \lambda, \mu, \nu$ , write

$$\nu \rightarrow_{\kappa\text{-c}} (\mu)_\lambda^2$$

if every coloring of the edges of the complete graph on  $\nu$  into  $\lambda$  many colors contains some size- $\mu$  monochromatic subgraph which is  $\kappa$ -connected. More formally:

DEFINITION 2.  $\nu \rightarrow_{\kappa-c} (\mu)_\lambda^2$  if and only if for every  $c: [\nu]^2 \rightarrow \lambda$  there exists a  $\xi < \lambda$  and  $X \in [\nu]^\mu$  such that the graph  $(X, c^{-1}(\xi) \cap [X]^2)$  is  $\kappa$ -connected.

QUESTION. For which cardinals does the relation  $\nu \rightarrow_{\kappa-c} (\mu)_\lambda^2$  hold?

## 2. Main results

Note at the outset that  $\kappa$ -connectedness is a well-studied notion, not least for its evident relevance to network design; it dates at least to Menger's 1927 [14] (see also [12]; see further [6], [1], [10], and [18] for interactions of the notion with vertex decompositions and chromatic number). Observe as well that it articulates a number of graph theory's most basic concerns:

- (1) A graph is 1-connected if and only if it is connected.
- (2) A graph with at least three vertices is 2-connected if and only if each pair of its vertices belongs to a cycle.
- (3) The only  $\mu$ -connected graph on any finite  $\mu$  is the complete one.

By this last point,

( $\star$ ) when  $\mu$  is finite,  $\nu \rightarrow_{\mu-c} (\mu)_\lambda^2$  is simply the Ramsey relation  $\nu \rightarrow (\mu)_\lambda^2$ .

Hence for any finite  $\lambda$  and  $\kappa \leq \mu$ ,

$$\nu \rightarrow_{\kappa-c} (\mu)_\lambda^2$$

for some  $\nu \leq R(\mu; \lambda)$ . See [13] for much finer bounds on the least such  $\nu$ .

For infinite  $\mu$ , a  $\mu$ -connected graph on  $\mu$  is no longer necessarily complete; such graphs nevertheless play a sufficiently critical role in the theory to merit a name and notation all their own:

DEFINITION 3. A graph  $G = (V, E)$  is *highly connected* if it remains connected after the deletion of any fewer than  $|V|$  vertices. Write  $\nu \rightarrow_{hc} (\mu)_\lambda^2$  if and only if  $\nu \rightarrow_{\mu-c} (\mu)_\lambda^2$ , i.e., if every coloring of the edges of the complete graph on  $\nu$  into  $\lambda$  many colors contains some size- $\mu$  monochromatic subgraph which is highly connected.

OBSERVATION 4.  $[\nu \rightarrow (\mu)_\lambda^2] \Rightarrow [\nu \rightarrow_{hc} (\mu)_\lambda^2] \Rightarrow [\nu \rightarrow_{\kappa-c} (\mu)_\lambda^2]$  for any  $\kappa \leq \mu$ .

In light of ( $\star$ ) and the following proposition, we might view  $\rightarrow_{hc}$  as a more satisfactory generalization of the positive Ramsey relations of ( $\star$ ) to the uncountably infinite:

PROPOSITION 5. *If  $\mu$  is an infinite cardinal and  $k$  is a natural number then  $\mu \rightarrow_{hc} (\mu)_k^2$ .*

PROOF. Given a coloring  $c: [\mu]^2 \rightarrow k$ , let  $\mathcal{D}$  be a uniform ultrafilter on  $\mu$  and define  $f: \mu \rightarrow k$  by

$$f(\alpha) = i \text{ if and only if } A_\alpha = \{\beta \in \mu : c(\{\alpha, \beta\}) = i\} \in \mathcal{D},$$

and let  $i < n$  be such that the set  $B = \{\alpha \in \mu : f(\alpha) = i\}$  is in  $\mathcal{D}$ . The set  $B$  is highly connected. This is because for any  $\alpha, \beta$  in  $B$ , the set  $A_\alpha \cap A_\beta \cap B$  is in  $\mathcal{D}$  and hence has cardinality  $\mu$ . Any  $\gamma$  in  $A_\alpha \cap A_\beta \cap B$  connects  $\alpha$  and  $\beta$  via the edge-colorings  $c(\{\alpha, \gamma\}) = c(\{\beta, \gamma\}) = i$ .  $\square$

The situation is considerably more complicated for infinitely many colors. Henceforth we will assume more set-theoretic background of the reader; we will focus as well on the relation  $\rightarrow_{hc}$ . This relation is subtle and significant in its own right, and we will tend to treat the finer relations  $\rightarrow_{\kappa-c}$  as secondary, as mainly grading its failure.

Perhaps the earliest result along these lines is Erdős and Kakutani's theorem [3] that the complete graph on an infinite cardinal  $\mu$  can be partitioned into  $\lambda$  many trees if and only if  $\mu \leq \lambda^+$ . In consequence, the relation  $\lambda^+ \rightarrow_{hc} (\lambda^+)_\lambda^2$  fails in the strongest possible respect:  $\lambda^+ \not\rightarrow_{2-c} (\lambda^+)_\lambda^2$ .

OBSERVATION 6.  $\mu \rightarrow_{1-c} (\mu)_\lambda^2$  holds for any cardinal  $\lambda$  less than the cofinality of  $\mu$ .

OBSERVATION 7. *The Erdős–Kakutani coloring shows even that  $\lambda^+ \not\rightarrow_{hc} (\mu)_\lambda^2$  for any  $\mu \geq 3$ .*

Alternately,  $\lambda^+ \not\rightarrow_{hc} (\lambda^+)_\lambda^2$  may be viewed as an instance of the following proposition, inspired by the *Sierpiński coloring* of [17]:

PROPOSITION 8. *If  $\mu \leq 2^\lambda$  then  $\mu \not\rightarrow_{hc} (\mu)_\lambda^2$ .*

PROOF. Let  $\{\eta_\alpha : \alpha < \mu\} \subseteq 2^\lambda$  be a collection of pairwise distinct functions. Given  $\alpha \neq \beta < \mu$  let

$$\Delta_{\alpha, \beta} = \min \{ \xi < \lambda : \eta_\alpha(\xi) \neq \eta_\beta(\xi) \}.$$

Then define  $c: [\mu]^2 \rightarrow \lambda \times 2$  by

$$c(\{\alpha, \beta\}) = (\Delta_{\alpha, \beta}, i) \text{ if and only if } \alpha < \beta \text{ and } \eta_\alpha(\Delta_{\alpha, \beta}) = i.$$

Aiming for a contradiction assume that  $A \subset \mu$  is highly connected in color  $(\xi, i)$ . Let  $\beta \in A$  be such that  $\eta_\beta(\xi) \neq i$  (there is such a  $\beta$  since  $A$  contains an edge). Let  $B = A \cap \beta$ . Then  $\beta$  has no adjacent edges in  $(A \setminus B, c^{-1}((\xi, i)) \cap [A \setminus B]^2)$ , a contradiction.  $\square$

By the following observation, Proposition 8 says even more.

LEMMA 9. Let  $\mu$  be the cofinality of  $\nu$ . Then  $\mu \not\rightarrow_{hc} (\mu)_\lambda^2$  implies that  $\nu \not\rightarrow_{hc} (\nu)_\lambda^2$ .

PROOF. Let  $c: [\mu]^2 \rightarrow \lambda$  witness that  $\mu \not\rightarrow_{hc} (\mu)_\lambda^2$ . Let  $\nu$  be the disjoint union of  $\mu$  many sets  $\nu_\alpha$ , each of strictly smaller cardinality than  $\nu$ . A coloring  $d: [\nu]^2 \rightarrow \lambda$  for which  $d(\xi, \eta) = c(\alpha, \beta)$  if  $\xi \in \nu_\alpha$  and  $\eta \in \nu_\beta$  and  $\alpha \neq \beta$  will witness that  $\nu \not\rightarrow_{hc} (\nu)_\lambda^2$ .  $\square$

The core relation in Proposition 8 is  $2^\lambda \not\rightarrow_{hc} (2^\lambda)_\lambda^2$ . This is sharp, in the sense that the relation  $(2^\lambda)^+ \rightarrow_{hc} (2^\lambda)_\lambda^2$  does hold:<sup>1</sup>

PROPOSITION 10. If  $\mu = \mu^\lambda$  then  $\mu^+ \rightarrow_{hc} (\mu)_\lambda^2$ .

PROOF. Observe first that by assumption, the cofinality of  $\mu$  is greater than  $\lambda$ .

Given a coloring  $c: [\mu^+]^2 \rightarrow \lambda$ , let  $\{M_\varepsilon : \varepsilon \leq \mu\}$  be a continuous  $\subseteq$ -chain of size- $\mu$  elementary submodels of some large enough  $H(\theta)$  such that

- (1)  $\mu + 1 \cup \{c\} \subseteq M_0$ ,
- (2)  $[M_{\varepsilon+1}]^\lambda \subseteq M_{\varepsilon+1}$  for every  $\varepsilon < \mu$ , and
- (3) for every  $\varepsilon < \mu$ , every formula  $\varphi \in L_{\lambda^+, \lambda^+}(\in)$  with parameters in  $M_{\varepsilon+1}$  that is satisfiable in  $H(\theta)$  is satisfiable in  $M_{\varepsilon+1}$ .

For each  $\varepsilon \leq \mu$  let  $\delta_\varepsilon = M_\varepsilon \cap \mu^+ \in \mu^+$ . The key observation is the following:

CLAIM 11. For co-boundedly many  $\varepsilon < \mu$  there is an  $i(\varepsilon) < \lambda$  such that  $c(\{\alpha, \delta_\mu\}) = c(\{\beta, \delta_\mu\}) = i(\varepsilon)$  for some  $\alpha, \beta < \delta_\varepsilon$  and such that for every such  $\alpha, \beta$ , the set

$$\{\gamma \in [\delta_\varepsilon, \delta_{\varepsilon+1}) : c(\{\alpha, \gamma\}) = c(\{\beta, \gamma\}) = c(\{\gamma, \delta_\mu\}) = i = i(\varepsilon)\}$$

is unbounded in  $\delta_{\varepsilon+1}$ .

Observe that we do not require the  $\alpha, \beta$  of the claim to be distinct.

PROOF. Adopt the convention that  $\min \emptyset = 0$  and let

$$\xi = \sup\{\min\{\alpha : \alpha < \delta_\mu \text{ and } c(\alpha, \delta_\mu) = i\} : i \in \lambda\}$$

and take  $\eta < \mu$  with  $\delta_\eta > \xi$ . Claim 11 will hold for all  $\varepsilon$  in the interval  $(\eta, \mu)$ . This we argue by contradiction: assume the claim fails for some  $\varepsilon \in (\eta, \mu)$ ; in other words, for each  $i < \lambda$  assume that one of the following holds:

- (a) There exist  $\alpha_i, \beta_i < \varepsilon$  such that  $c(\{\alpha_i, \delta_\mu\}) = c(\{\beta_i, \delta_\mu\}) = i$ , yet the set

$$\Gamma_i = \{\gamma \in [\delta_\varepsilon, \delta_{\varepsilon+1}) : c(\{\alpha_i, \gamma\}) = c(\{\beta_i, \gamma\}) = c(\{\gamma, \delta_\mu\}) = i\}$$

is bounded in  $\delta_{\varepsilon+1}$ .

<sup>1</sup>(Compare the Erdős–Rado relation  $(2^\lambda)^+ \rightarrow (\lambda^+)_\lambda^2$ . It too is sharp, in the sense that  $2^\lambda \not\rightarrow (\lambda^+)_\lambda^2$ , as witnessed by the Sierpiński coloring.)

(b) For no  $\alpha < \delta_\varepsilon$  does  $c(\alpha, \delta_\mu) = i$ . In this case let  $\Gamma_i$  be  $\emptyset$ .

As  $[M_{\varepsilon+1}]^\lambda \subseteq M_{\varepsilon+1}$ , the cofinality of  $\delta_{\varepsilon+1}$  is bigger than  $\lambda$ , hence some  $\zeta < \delta_{\varepsilon+1}$  bounds  $\bigcup_{i < \lambda} \Gamma_i$ . Denote the family of  $i$  satisfying (a) by  $I$ . Consider the conjunction  $\varphi$  of the formula  $\gamma > \zeta$  with the formulas  $c(\{\alpha_i, \gamma\}) = c(\{\beta_i, \gamma\}) = i$ , where  $i$  ranges through  $I$ . Then  $\delta_\mu$  witnesses that  $\varphi$  is satisfiable in  $H(\theta)$ , so by (3) some  $\gamma \in (\zeta, \delta_{\varepsilon+1})$  witnesses its satisfaction in  $M_{\varepsilon+1}$ . Let  $j = c(\{\gamma, \delta_\mu\})$ . Then  $j \in I$  and  $\gamma \in \Gamma_j$ , which contradicts the assumption that  $\Gamma_j$  is bounded by  $\zeta$ .  $\square$

Let  $i < \lambda$  be such that the set  $W = \{\varepsilon : i(\varepsilon) = i\}$  has size  $\mu$ , and let  $A = \bigcup \{A_\varepsilon : \varepsilon \in W\}$ , where

$$A_\varepsilon = \{\alpha \in [\delta_\varepsilon, \delta_{\varepsilon+1}) : c(\{\alpha, \delta_\mu\}) = i\}.$$

$A$  is then a subset of  $\mu^+$  of size  $\mu$  (in fact,  $A_\varepsilon$  has size  $\mu$  for every  $\varepsilon \in W$ ). We claim that it is highly connected in the color  $i$ . To see this it suffices to prove that if  $\alpha$  and  $\beta$  are distinct elements of  $A$  then the set

$$\{\gamma \in A : c(\{\alpha, \gamma\}) = c(\{\beta, \gamma\}) = i\}$$

has size  $\mu$ . To see this, let  $\varepsilon_1$  and  $\varepsilon_2$  be elements of  $W$  such that  $\alpha \in A_{\varepsilon_1}$  and  $\beta \in A_{\varepsilon_2}$ , and let  $\varepsilon$  be an element of  $W \setminus (\varepsilon_1 + 1 \cup \varepsilon_2 + 1)$ . We wish to find a  $\gamma \in [\delta_\varepsilon, \delta_{\varepsilon+1})$  such that

$$c(\{\alpha, \gamma\}) = c(\{\beta, \gamma\}) = c(\{\gamma, \delta_\mu\}) = i = i(\varepsilon).$$

Such a  $\gamma$  exists by Claim 11.  $\square$

A number of questions now come into focus. Most immediate among them is:

QUESTION 12. *For  $\lambda$  an infinite cardinal, what is the least cardinal  $\mu$  for which it is consistent with the ZFC axioms that  $\mu \rightarrow_{hc} (\mu)_\lambda^2$ ?*

By Observation 4,  $\mu \rightarrow_{hc} (\mu)_\lambda^2$  holds whenever  $\mu$  is weakly compact, for any  $\lambda < \mu$ . As we have seen, though,  $\rightarrow_{hc}$  holds in many cases where the classical arrow fails; hence we might reasonably hope for  $\mu \rightarrow_{hc} (\mu)_\lambda^2$  on much smaller  $\mu$ . Necessarily,  $2^\lambda$  must be smaller than any such  $\mu$ , by Proposition 8. Is this alone enough?

No. By the following, any instance of  $\mu \rightarrow_{hc} (\mu)_\lambda^2$  will involve large cardinal assumptions.

DEFINITION 13. For regular uncountable  $\mu$ , the principle  $\square(\mu)$  is the assertion that there exists a sequence  $\mathcal{C} = \langle C_\alpha \mid \alpha \in \mu \rangle$  such that

- $C_\alpha$  is a closed unbounded subset of  $\alpha$ , for each  $\alpha$ .
- $C_\beta \cap \alpha = C_\alpha$ , for every limit point  $\alpha$  of  $C_\beta$ .
- No club  $C \subseteq \mu$  satisfies  $C \cap \alpha = C_\alpha$  at every limit point  $\alpha$  of  $C$ .

If in addition the following holds, we will call  $\mathcal{C}$  a  $\lambda$ -stationary  $\square(\mu)$ -sequence:

- The set  $\{\alpha \in \mu : \text{otp}(C_\alpha) = \lambda\}$  is stationary in  $\mu$ .

The following is immediate from [8] together with [20, Lemma 7.2.2].

**THEOREM 14.** *If  $\lambda < \mu$  are infinite regular cardinals and  $\mu$  is not Mahlo in the constructible universe, then there exists a  $\lambda$ -stationary  $\square(\mu)$ -sequence.*

**PROPOSITION 15.** *If there exists a  $\lambda$ -stationary  $\square(\mu)$ -sequence, then  $\mu \not\rightarrow_{hc} (\mu)_\lambda^2$ .*

**PROOF.** The “bad” coloring  $c: [\mu]^2 \rightarrow \lambda$  will be  $c(\alpha, \beta) = \rho^\lambda(\alpha, \beta)$ , where  $\rho^\lambda$  is Todorcevic’s local rho function, defined in reference to some  $\lambda$ -stationary  $\square(\mu)$ -sequence  $\mathcal{C}$ . Readers are referred to [20] §7.2 for further information. The decisive features of  $\rho^\lambda$  for our purposes are the following: for all  $\alpha < \beta < \gamma < \mu$ ,

$$(1) \quad \rho^\lambda(\alpha, \beta) \leq \max \{ \rho^\lambda(\alpha, \gamma), \rho^\lambda(\beta, \gamma) \},$$

and

$$(2) \quad \rho^\lambda(\alpha, \gamma) \leq \max \{ \rho^\lambda(\alpha, \beta), \rho^\lambda(\beta, \gamma) \}$$

In consequence, for all  $\xi < \lambda$ , the relation

$$\alpha <_\xi^\lambda \beta \text{ iff } \alpha < \beta \text{ and } \rho^\lambda(\alpha, \beta) \leq \xi$$

is a tree-ordering on  $\mu$ . By our assumptions about  $\mathcal{C}$  and [20, Lemma 7.2.9], none of the orderings  $<_\xi^\lambda$  contains a chain of length  $\mu$ .

Now suppose towards contradiction that  $A \in [\mu]^\mu$  is highly connected in the color  $\xi$ . By the above, there exist  $\alpha < \beta$  in  $A$  with  $\alpha \not<_\xi^\lambda \beta$ . By highly-connectedness, some color- $\xi$  path  $\alpha = \alpha_0$  to  $\alpha_1$  to  $\dots$  to  $\alpha_j$  to  $\alpha_{j+1} = \beta$  must connect  $\alpha$  and  $\beta$  in  $A \setminus \alpha$ . It then follows from successive applications of (1) and (2) above that

$$\rho^\lambda(\alpha, \beta) \leq: \max_{i \leq j} \rho^\lambda(\alpha_i, \alpha_{i+1}) = \xi$$

This implies that  $\alpha <_\xi^\lambda \beta$ , a contradiction.  $\square$

**COROLLARY 16.** *It is consistent with the ZFC axioms — and even with ZFC+GCH — that  $\mu \not\rightarrow_{hc} (\mu)_\lambda^2$  for all infinite cardinals  $\lambda < \mu$ .*

**PROOF.** By Proposition 15, the relation  $\mu \not\rightarrow_{hc} (\mu)_\lambda^2$  holds for any infinite regular cardinals  $\lambda < \mu$  in a model of ZFC+(V=L)+“there exist no Mahlo cardinals.” It will hold then for any singular  $\mu$  by Lemma 9. Observe

finally that if  $\mu \not\rightarrow_{hc} (\mu)_\lambda^2$  failed for any singular  $\lambda$ , it would fail as well for some smaller regular  $\lambda$ , contradicting our premise.  $\square$

We now show in the opposite direction that, assuming the existence of a weakly compact cardinal above some  $\mu > \lambda$ , it is consistent with the ZFC axioms that  $2^\mu \rightarrow_{hc} (2^\mu)_\lambda^2$ .

Instrumental in the argument is the following “two-dimensional delta system lemma” of more general utility.

**DEFINITION 17.** A family of sets  $\mathcal{A}$  is a  $\Delta$ -system if there exists a fixed  $r$  such that  $a \cap b = r$  for every distinct  $a$  and  $b$  in  $\mathcal{A}$ . This  $r$  is called the *root* of the  $\Delta$ -system.

**LEMMA 18.** *Let  $\nu$  be weakly compact and let  $\mu$  be less than  $\nu$ . Then for any family  $\{u_{\alpha,\beta} : \alpha < \beta < \nu\} \subseteq [\nu]^{\leq \mu}$  there exists a  $B \in [\nu]^\nu$  such that:*

(1) *For each  $\alpha \in B$ , the set  $\{u_{\alpha,\beta} : \beta \in B \setminus (\alpha + 1)\}$  is a  $\Delta$ -system, with root  $V_\alpha^+$ .*

(2) *For each  $\beta \in B$ , the set  $\{u_{\alpha,\beta} : \alpha \in B \cap \beta\}$  is a  $\Delta$ -system, with root  $V_\beta^-$ .*

(3) *The sets  $\{V_\alpha^+ : \alpha \in B\}$  and  $\{V_\alpha^- : \alpha \in B\}$  and  $\{V_\alpha^- \cup V_\alpha^+ : \alpha \in B\}$  each form  $\Delta$ -systems.*

(4) *The elements of the set  $\{u_{\alpha,\beta} \setminus (V_\alpha^+ \cup V_\beta^-) : \alpha < \beta \text{ in } B\}$  are pairwise disjoint.*

In what follows, the relation  $\alpha < \beta$  will sometimes be left implicit; it is assumed to hold in any expression conjoining  $\alpha$  and  $\beta$ .

**PROOF.** By the weak compactness of  $\nu$ , we may begin by assuming all  $u_{\alpha,\beta}$  to be of the same order-type.

Define the coloring  $d: [\nu]^4 \rightarrow H(\mu^+)$  as follows: for any increasing  $a = \{\alpha_0, \alpha_1, \alpha_2, \alpha_3\}$  let

$$U_a = a \cup \bigcup_{j < k < 4} u_{\alpha_j, \alpha_k}.$$

and let

$$d(a) = \langle \text{otp}(U_a), \langle \xi_i : i < 4 \rangle, \langle v_{j,k} : j < k < 4 \rangle \rangle,$$

so that if  $h$  is the unique order-isomorphism between  $U_a$  and  $\text{otp}(U_a)$ , then

- $h(\alpha_i) = \xi_i$  for every  $i < 4$ , and
- $h[u_{\alpha_j, \alpha_k}] = v_{j,k}$  for every  $j < k < 4$ .

By the weak compactness of  $\nu$ , there exists a  $d$ -monochromatic  $A \in [\nu]^\nu$ . We argue most of the lemma for this set  $A$ , thinning to a  $B \in [A]^\nu$  only later if necessary.



Consider  $\alpha < \beta < \gamma < \delta < \varepsilon$  in  $A$ . Since  $d(\alpha, \beta, \gamma, \varepsilon) = d(\alpha, \beta, \delta, \varepsilon) = d(\alpha, \gamma, \delta, \varepsilon)$ ,

$$(\dagger) \quad \xi \in u_{\alpha, \beta} \cap u_{\alpha, \gamma} \Leftrightarrow \xi \in u_{\alpha, \beta} \cap u_{\alpha, \delta} \Leftrightarrow \xi \in u_{\alpha, \gamma} \cap u_{\alpha, \delta}$$

As  $\varepsilon$  was arbitrary, this implies item (1) of the lemma. We might usefully note more: a  $\xi$  as in  $(\dagger)$  must sit at the same relative height in each  $u_{\alpha, \beta}$ , for  $\beta \in A \setminus (\alpha + 1)$ . Pigeonhole arguments together with  $(\dagger)$  then ensure that any lesser elements of  $u_{\alpha, \beta}$  also fall in the root  $V_\alpha^+$  of the  $\Delta$ -system  $\{u_{\alpha, \beta} : \beta \in A \setminus (\alpha + 1)\}$ . In other words,  $V_\alpha^+$  is an initial segment of each such  $u_{\alpha, \beta}$ .

Item (2) of the lemma is similar. (Pigeonhole arguments are not available in this case, hence the root  $V_\beta^-$  is not so easily characterized.)

To see that  $\{V_\alpha^+ : \alpha \in A\}$  forms a  $\Delta$ -system with root  $r^+$ , observe that

$$\begin{aligned} \xi \in V_\beta^+ \cap V_\gamma^+ &\Rightarrow \xi \in u_{\beta, \varepsilon} \cap u_{\gamma, \varepsilon} \text{ for any } \varepsilon \in A \setminus (\gamma + 1) \\ &\Rightarrow \xi \in V_\varepsilon^- \\ &\Rightarrow \xi \in u_{\alpha, \varepsilon} \text{ for any } \alpha \in A \cap \varepsilon. \end{aligned}$$

As  $\varepsilon$  is arbitrary, this implies that  $\xi$  is in  $V_\alpha^+$ . As  $\alpha$  is arbitrary, this completes the argument.

The argument that  $\{V_\alpha^- : \alpha \in A\}$  forms a  $\Delta$ -system with root  $r^-$  is essentially identical (but may require the omission of the first two elements of  $A$ ).

Finally, note that  $\text{otp}(V_\alpha^+)$ ,  $\text{otp}(V_\beta^+)$ ,  $\text{otp}(V_\gamma^-)$ ,  $\text{otp}(V_\delta^-)$  are all legible from  $d(\alpha, \beta, \gamma, \delta)$ . In consequence:

(i)  $V_\beta^+ \cap V_\gamma^-$  is of the same order-type for all  $\beta < \gamma$  in  $A$ . Hence this intersection must be equal to  $r^+ \cap r^-$ . Thin  $A$  if necessary to a  $B \in [A]^\nu$  such that  $V_\beta^+ \setminus r^+ \cap V_\gamma^- \setminus r^- = \emptyset$  for all  $\beta \geq \gamma$  in  $B$ . Then  $\{V_\alpha^+ \cup V_\alpha^- : \alpha \in B\}$  forms a  $\Delta$ -system with root  $r = r^- \cup r^+$ .

(ii) The family  $\{u_{\alpha, \beta} \setminus (V_\alpha^+ \cup V_\beta^-) : \alpha < \beta \text{ in } B\}$  is pairwise disjoint. For by the homogeneity of  $B$ , any  $\xi$  in  $(u_{\beta, \gamma} \setminus V_\beta^+ \cup V_\gamma^-) \cap (u_{\alpha, \delta} \setminus V_\alpha^+ \cup V_\delta^-)$  is necessarily also in  $(u_{\beta, \gamma} \setminus V_\beta^+ \cup V_\gamma^-) \cap (u_{\alpha, \varepsilon} \setminus V_\alpha^+ \cup V_\varepsilon^-)$  for any  $\varepsilon \neq \delta$  in  $B$ . But this implies that  $\xi \in V_\alpha^+$ , a contradiction. Similarly for any other configuration of  $\alpha, \beta, \gamma$ , and  $\delta$ . These establish items (3) and (4) of the lemma.  $\square$

REMARK 19. Two further features of the above system will be useful below:

- $V_\alpha^+$  are all of the same order-type, for  $\alpha \in B$ . Similarly for  $V_\alpha^-$ .
- As each  $\alpha$  in  $B$  sits at the same distinguished relative location in  $V_\alpha := V_\alpha^- \cup V_\alpha^+$  we have  $|\{\text{otp}(V_\alpha) : \alpha \in B\}| = 1$  as well.

**THEOREM 20.** *Let  $\nu$  be a weakly compact cardinal, and let  $\lambda < \mu = \mu^{<\mu} < \nu$  be given. Then there is a cardinal-preserving forcing  $\mathbb{P}$  such that*

$$\Vdash_{\mathbb{P}} \text{“} 2^\mu = \nu \text{ and } \nu \rightarrow_{hc} (\nu)_\lambda^2 \text{”}.$$

**PROOF.** Let  $\mathbb{P}$  be the forcing for adding  $\nu$  many  $\mu$ -Cohen subsets of  $\nu$ , i.e.,

$$\mathbb{P} = \{p : p \text{ is a partial function from } \nu \text{ to } 2 \text{ of size less than } \mu\}$$

reverse-ordered by extension. By assumption,  $\mathbb{P}$  has the  $\mu^+$ -c.c.; it is evidently  $\mu$ -closed as well, and consequently preserves cardinals. By standard arguments,  $\Vdash_{\mathbb{P}} \text{“} 2^\mu = \nu \text{”}$ .

Let  $\dot{c}$  be a  $\mathbb{P}$ -name such that  $\Vdash_{\mathbb{P}} \text{“} \dot{c} : [\nu]^2 \rightarrow \lambda \text{”}$ . For every  $\alpha < \beta < \nu$  let  $\mathcal{A}_{\alpha,\beta} := \{p_{\alpha,\beta,\xi} : \xi < \mu\}$  be a maximal antichain in  $\mathbb{P}$  with corresponding  $\{i_{\alpha,\beta,\xi} : \xi < \mu\}$  such that  $p_{\alpha,\beta,\xi} \Vdash \text{“} \dot{c}(\{\alpha, \beta\}) = i_{\alpha,\beta,\xi} \text{”}$ . Let

$$u_{\alpha,\beta} = \{\alpha, \beta\} \cup \bigcup_{\xi < \mu} \text{dom}(p_{\alpha,\beta,\xi}).$$

Let  $\{\gamma_{\alpha,\beta,\eta} : \eta < \varepsilon_{\alpha,\beta}\}$  enumerate  $u_{\alpha,\beta}$  in increasing order. Define a relation  $E$  on  $[\nu]^2$  by declaring  $\{\alpha_1, \beta_1\} E \{\alpha_2, \beta_2\}$  if and only if

- (1)  $\varepsilon_{\alpha_1,\beta_1} = \varepsilon_{\alpha_2,\beta_2}$ ,
- (2)  $\alpha_1 = \gamma_{\alpha_1,\beta_1,\eta}$  if and only if  $\alpha_2 = \gamma_{\alpha_2,\beta_2,\eta}$ ,
- (3)  $\beta_1 = \gamma_{\alpha_1,\beta_1,\eta}$  if and only if  $\beta_2 = \gamma_{\alpha_2,\beta_2,\eta}$ ,
- (4)  $\{\eta : \gamma_{\alpha_1,\beta_1,\eta} \in \text{dom}(p_{\alpha_1,\beta_1,\xi})\} = \{\eta : \gamma_{\alpha_2,\beta_2,\eta} \in \text{dom}(p_{\alpha_2,\beta_2,\xi})\}$  for every  $\xi < \mu$ ,
- (5)  $p_{\alpha_1,\beta_1,\xi}(\gamma_{\alpha_1,\beta_1,\eta}) = p_{\alpha_2,\beta_2,\xi}(\gamma_{\alpha_2,\beta_2,\eta})$  for every  $\eta$  as in (4) and  $\xi < \mu$ , and
- (6)  $i_{\alpha_1,\beta_1,\xi} = i_{\alpha_2,\beta_2,\xi}$  for every  $\xi < \mu$ .

Clearly  $E$  is an equivalence relation on  $[\nu]^2$  with  $2^\mu < \nu$  many equivalence classes. As  $\nu$  is weakly compact, there exists an  $A \in [\nu]^\nu$  such that  $\{u_{\alpha,\beta} : \{\alpha, \beta\} \in [A]^2\}$  all falls in a single class. Further thin  $A$  to a  $B \subseteq A$  as in Lemma 18. Write  $\mathfrak{r}$  for the root of the  $\Delta$ -system  $\{V_\beta : \beta \in B\}$ .

Observe that in this context the key terms of Lemma 18 take on more particular meanings:  $V_\alpha^+$ , for example, records exactly those coordinates at which some  $p \in \mathcal{A}_{\alpha,\beta}$  and  $q \in \mathcal{A}_{\alpha,\gamma}$  may disagree. The argument now proceeds in two steps; the uniformities of the family  $B$  are important to each. In the first step, we extend any  $q \in \mathbb{P}$  to an  $s$  deciding the elements of some  $\dot{c}$ -homogeneous  $X \in [\nu]^\lambda$ . In the second step, genericity below  $q$  propagates that homogeneity to highly connect a cofinal  $Y \subseteq \nu$  in the forcing extension.

Therefore fix  $q \in \mathbb{P}$  and a  $W \subseteq B$  of order-type  $\lambda^+ + 1$  such that

$$\text{dom}(q) \cap \bigcup \{u_{\alpha,\beta} : \{\alpha, \beta\} \in [W]^2\} \subseteq \mathfrak{r}.$$

We will recursively define “colors”  $i_\eta \in \lambda$  and a sequence  $\langle \varepsilon_\eta : \eta \leq \lambda^+ \rangle \subseteq W$  and conditions  $q_\eta$  below  $q$  so that for all  $\eta < \lambda^+$ ,

- (1)  $\xi < \eta$  implies that  $q_\xi \geq q_\eta$ ,
- (2)  $q_\eta \Vdash \dot{c}(\{\varepsilon_\xi, \varepsilon_{\lambda^+}\}) = i_\xi$  for all  $\xi \leq \eta$ ,
- (3)  $q_\eta \Vdash \dot{c}(\{\varepsilon_\xi, \varepsilon_\zeta\}) = i_\xi$  for all  $\xi < \zeta \leq \eta$ , and
- (4)  $q_\eta = q \cup \left[ \bigcup_{\xi \leq \eta} p_{\varepsilon_\xi, \varepsilon_{\lambda^+}} \right] \cup \left[ \bigcup_{\xi < \zeta \leq \eta} p_{\varepsilon_\xi, \varepsilon_\zeta} \right]$ , where each  $p_{\varepsilon_\xi, \varepsilon_\zeta}$  is a member of the antichain  $\mathcal{A}_{\varepsilon_\xi, \varepsilon_\zeta}$ .

To begin the construction, let  $\varepsilon_0$  and  $\varepsilon_{\lambda^+}$  be the first and last elements of  $W$ , respectively, and let  $q_0$  be the union of  $q$  with any compatible element of  $\mathcal{A}_{\varepsilon_0, \varepsilon_{\lambda^+}}$ . Suppose now that for each  $\eta < \delta$  coordinates  $\varepsilon_\eta$  and conditions  $q_\eta$  have been selected which together satisfy (1)-(4). As  $\delta < \lambda^+$ , by Lemma 18 there exists an  $\varepsilon_\delta \in W$  such that  $u_{\varepsilon_\eta, \varepsilon_\delta} \setminus V_{\varepsilon_\eta}^+$  is disjoint from  $\bigcup_{\eta < \delta} q_\eta$  for each  $\eta < \delta$ . There also exists for each such  $\eta$  a  $p_{\varepsilon_\eta, \varepsilon_{\lambda^+}} \in \mathcal{A}_{\varepsilon_\eta, \varepsilon_{\lambda^+}}$  so that  $\bigcup_{\eta < \delta} p_{\varepsilon_\eta, \varepsilon_{\lambda^+}} \subseteq \bigcup_{\eta < \delta} q_\eta$ . Let  $p'_{\varepsilon_\eta, \varepsilon_\delta}$  denote their induced respective “copies” under the order-isomorphisms  $\pi_\eta : u_{\varepsilon_\eta, \varepsilon_{\lambda^+}} \rightarrow u_{\varepsilon_\eta, \varepsilon_\delta}$ . By arrangement,

$$q'_\delta := \left[ \bigcup_{\eta < \delta} q_\eta \right] \cup \left[ \bigcup_{\eta < \delta} p'_{\varepsilon_\eta, \varepsilon_\delta} \right]$$

is a function. Let  $q_\delta$  be the union of  $q'_\delta$  with any compatible element of  $\mathcal{A}_{\varepsilon_\delta, \varepsilon_{\lambda^+}}$ . The condition  $q_\delta$  is as desired; in this fashion the construction proceeds.

For some  $i \in \lambda$  the set  $\{\eta : i_\eta = i\}$  is unbounded in  $\lambda^+$ . Let  $X'$  collect its first  $\lambda$  elements. Let  $\bar{\eta} = \sup X'$  and let  $s = q_{\bar{\eta}}$  and let  $X = \{\varepsilon_\eta : \eta \in X'\}$ . Clearly  $s \Vdash \dot{c}''[X]^2 = \{i\}$ . This completes the first of the steps described above.

For the second step, let  $s_\varepsilon = s \upharpoonright V_\varepsilon$  for each  $\varepsilon \in X$ . Let  $V[G]$  be a forcing extension of  $V$  by a  $\mathbb{P}$ -generic filter  $G$ ; therein define the family

$$Y = \bigcup_{\varepsilon \in X} Y_\varepsilon$$

where

$$Y_\varepsilon = \left\{ \alpha \in B : \text{there exists a } p \in G \text{ with } \text{dom}(p) \subseteq V_\alpha \text{ such that} \right. \\ \left. \text{the order-isomorphism } \pi : V_\alpha \rightarrow V_\varepsilon \text{ sends } p \text{ to } s_\varepsilon \right\}.$$

Lastly, let  $q' = q \cup (s \upharpoonright \mathfrak{r})$ .

CLAIM 21.  $q' \Vdash \dot{c}^{-1}(i) \cap [\dot{Y}]^2$  is a highly connected graph of size  $\nu$ .

As  $q$  was arbitrary, this claim will establish the proposition.

PROOF OF CLAIM 21. Observe first that for each  $\alpha \in \nu$  and  $\varepsilon \in X$  the set

$$D_\alpha^\varepsilon := \{p : p \Vdash \text{“}\dot{Y}_\varepsilon \not\subseteq \alpha\text{”}\}$$

is dense below  $q'$ . Hence  $q'$  forces that each  $Y_\varepsilon$  is unbounded in  $\nu$ .

Suppose now that  $q'' \leq q'$  forces that  $\alpha$  and  $\beta$  are in  $Y$ . Without loss of generality,  $q''$  decides the witnesses to this fact as well, i.e., there exist some  $p_\alpha, p_\beta \subseteq q$  and  $\varepsilon(\alpha), \varepsilon(\beta) \in X$  such that the order-isomorphism  $\pi: V_\alpha \rightarrow V_{\varepsilon(\alpha)}$  sends  $p_\alpha$  to  $s_{\varepsilon(\alpha)}$ , and similarly for  $\beta$ . We'll show that for any  $\gamma \in \nu$  there exists a  $\delta > \gamma$  and  $q''' \leq q''$  such that

$$(\ddagger) \quad q''' \Vdash \text{“}\dot{c}(\alpha, \delta) = \dot{c}(\beta, \delta) = i\text{”}$$

This will establish the claim. To that end, take  $\varepsilon \in X \setminus (\max\{\varepsilon(\alpha), \varepsilon(\beta)\} + 1)$ . By Lemma 18 there exists a  $\delta \in B \setminus (\gamma + 1)$  such that  $u_{\alpha, \delta} \setminus V_\alpha$  and  $u_{\beta, \delta} \setminus V_\beta$  are disjoint from  $\text{dom}(q'')$ . Extend  $q''$  by “copying”  $s \upharpoonright (u_{\varepsilon(\alpha), \varepsilon} \cup u_{\varepsilon(\beta), \varepsilon})$  via the order-isomorphism to  $u_{\alpha, \delta} \cup u_{\beta, \delta}$ . Denote this extension by  $q'''$ . Our assumptions on  $\alpha$  and  $\beta$  ensure that  $q'''$  is in fact a condition, and our assumptions on  $E$  translate the relation  $s \Vdash \text{“}\dot{c}(\varepsilon(\alpha), \varepsilon) = \dot{c}(\varepsilon(\beta), \varepsilon) = i\text{”}$  to  $(\ddagger)$ , as desired.  $\square \square$

COROLLARY 22. *Assuming the existence of a weakly compact cardinal, it is consistent with the ZFC axioms that  $2^{\aleph_1} \rightarrow_{hc} (2^{\aleph_1})_{\aleph_0}^2$ .*

### 3. Main questions

We turn in conclusion to the most immediate instance of Question 12:

QUESTION 23. *What is the least cardinal  $\mu$  for which it is consistent with the ZFC axioms that  $\mu \rightarrow_{hc} (\mu)_{\aleph_0}^2$ ?*

Corollary 22 may be viewed as approximating to any of several possibilities. For example: the  $\mu = 2^{\aleph_1}$  of Corollary 22 falls, in the forcing extension of Proposition 20, well below any weakly compact  $\nu$ , but it remains a regular limit cardinal. Therefore we may ask:

QUESTION 24. *Must the least cardinal  $\mu$  for which it is consistent with the ZFC axioms that  $\mu \rightarrow_{hc} (\mu)_{\aleph_0}^2$  be weakly inaccessible?*

By Lemma 9, such a  $\mu$  is necessarily regular; hence Question 24 amounts to asking if such a  $\mu$  may be a successor cardinal. If indeed it may be, then Corollary 22 might be viewed instead as approximating to the following alternative to Question 24:

QUESTION 25. *Assuming whatever large cardinals may be necessary, is it consistent with the ZFC axioms that  $\aleph_2 \rightarrow_{hc} (\aleph_2)_{\aleph_0}^2$ ?*

Reasoning from [7] and [11] shows that if  $\mu$  is the successor of a regular cardinal and  $\lambda < \mu$ , then the consistency strength of “there exists no  $\lambda$ -stationary  $\square(\mu)$  sequence” is exactly a Mahlo cardinal. By Proposition 15, this gives a lower bound on the assumptions necessary to any affirmative answer to Question 25. Again by Proposition 15, answers to any of the above questions entail further questions of consistency strength; most generally:

QUESTION 26. *What is the consistency strength of the existence of an uncountable cardinal  $\mu$  such that  $\mu \rightarrow_{hc} (\mu)_{\aleph_0}^2$ ?*

By Proposition 8, positive relations  $\mu \rightarrow_{hc} (\mu)_\lambda^2$  will involve cardinal arithmetic assumptions as well. An affirmative answer to Question 25, for example, would imply the continuum hypothesis. It is unclear if weaker relations would also. More particularly, the continuum hypothesis implies that  $\aleph_2 \rightarrow_{hc} (\aleph_1)_{\aleph_0}^2$ , by Proposition 10. Is the reverse true? In other words:

QUESTION 27. *Is the continuum hypothesis equivalent to the assertion that  $\aleph_2 \rightarrow_{hc} (\aleph_1)_{\aleph_0}^2$ ?*

A question of a similar flavor is the following:

QUESTION 28. *Does  $\aleph_2 \rightarrow_{\aleph_1-c} (\aleph_2)_{\aleph_0}^2$ ?*

As this paper goes to press, the authors have the pleasure of referring the reader to Chris Lambie-Hanson’s recent <https://arxiv.org/abs/2005.10812> for partial progress on some of these and related questions.

**Acknowledgements.** The first author would like to thank Chris Lambie-Hanson for valuable discussion of the consistency strengths of failures of  $\lambda$ -stationary  $\square(\mu)$  sequences to exist. The authors would like to thank the referee for a prompt, alert, and intelligent reading. This paper is number 1157 in the third authors publication list.

## References

- [1] M. Elekes, D. Soukup, L. Soukup and Z. Szentmiklóssy, Decompositions of edge-colored infinite complete graphs into monochromatic paths, *Discrete Math.*, **340** (2017), 2053–2069.
- [2] P. Erdős, A. Hajnal, A. Máté and R. Rado, *Combinatorial Set Theory: Partition Relations for Cardinals*, North-Holland (Amsterdam, 1984).
- [3] P. Erdős and S. Kakutani, On non-denumerable graphs, *Bull. Amer. Math. Soc.*, **49** (1943).
- [4] P. Erdős and R. Rado, A partition calculus in set theory, *Bull. Amer. Math. Soc.*, **56** (1956).
- [5] R. L. Graham, B. L. Rothschild and J. H. Spencer, *Ramsey Theory*, John Wiley & Sons (New York, 1980).
- [6] A. Hajnal, P. Komjáth, L. Soukup and I. Szalkai, Decompositions of edge colored infinite complete graphs, in: *Combinatorics* (Eger, 1987), Colloq. Math. Soc. János Bolyai, **52** (1988).

- [7] L. Harrington and S. Shelah, Some exact equiconsistency results in set theory, *Notre Dame J. Formal Logic*, **26** (1985).
- [8] R. B. Jensen, The fine structure of the constructible hierarchy, *Ann. Math. Logic*, **4** (1972).
- [9] A. Kanamori, *The Higher Infinite: Large Cardinals in Set Theory from their Beginnings*, Springer-Verlag (Berlin, 1994).
- [10] P. Komjáth, On the connectivity of infinite graphs, *Acta Math. Hungar.*, **154** (2018).
- [11] C. Lambie-Hanson, Squares and covering matrices, *Ann. Pure Appl. Logic*, **165** (2014).
- [12] W. Mader, Connectivity and edge-connectivity in finite graphs, *Surveys in Combinatorics*, B. Bollobás (ed.), Cambridge University Press (Cambridge, 1979).
- [13] D. W. Matula, Ramsey theory for graph connectivity, *J. Graph Theory*, **7** (1983).
- [14] K. Menger, Zur allgemeinen Kurventheorie, *Fund. Math.*, **10** (1927).
- [15] F. P. Ramsey, On a problem of formal logic, *Proc. London Math. Soc.*, **30** (1930).
- [16] S. Shelah, Was Sierpiński right? *Israel J. Math.*, **62** (1988).
- [17] W. Sierpiński, Sur un problème de la théorie des relations, *Ann. Scuola Norm. Sup. Pisa*, **2** (1933).
- [18] D. Soukup, Trees, ladders, and graphs, *J. Combin. Theory, Series B*, **115** (2015).
- [19] S. Todorćević, Partitioning pairs of countable ordinals, *Acta Math.*, **159** (1987).
- [20] S. Todorćević, *Walks on Ordinals and their Characteristics*, Birkhäuser (Basel, 2007).