

Avoiding equal distances*

Ashutosh Kumar[†], Saharon Shelah[‡]

Abstract

We show that it is consistent that there is a non meager set of reals each of whose non meager subsets contains equal distances.

1 Introduction

In [1], Erdős and Kakutani showed that the continuum hypothesis (CH) is equivalent to the following statement: There is a partition of the set of reals \mathbb{R} into countably many rationally independent sets. It follows that, under CH, every non meager set of reals contains a non meager (in fact, everywhere non meager) subset avoiding equal distances. The aim of this note is to show that CH is needed here.

Theorem 1.1. *It is consistent that there is a non meager $X \subseteq \mathbb{R}$ such that for every non meager $Y \subseteq X$, there are $a < b < c < d \in Y$ such that $b - a = d - c$.*

Note that, by a result of Rado (Theorem 3.2 in [2]), we cannot require Y to avoid arithmetic progressions of length 3. Also by [1], X cannot have size \aleph_1 . So we start by adding \aleph_2 Cohen reals and consider the set X of their pairwise sums. We then make every small subset Y of X meager using a finite support product where Y is small if it avoids equal distances. To capture the new subsets of X that may appear later, we use a sigma ideal \mathcal{I}

*2010 *Mathematics Subject Classification*: Primary 03E35; Secondary 03E75. *Key words and phrases*: Forcing, Baire category

[†]Einstein Institute of Mathematics, The Hebrew University of Jerusalem, Edmond J Safra Campus, Givat Ram, Jerusalem 91904, Israel; email: akumar@math.huji.ac.il; Supported by a Postdoctoral Fellowship at the Einstein Institute of Mathematics funded by European Research Council grant 338821

[‡]Einstein Institute of Mathematics, The Hebrew University of Jerusalem, Edmond J Safra Campus, Givat Ram, Jerusalem 91904, Israel and Department of Mathematics, Rutgers, The State University of New Jersey, Hill Center-Busch Campus, 110 Frelinghuysen Road, Piscataway, NJ 08854-8019, USA; email: shelah@math.huji.ac.il; Partially supported by European Research Council grant 338821; Publication no. 1079

(see below). The rest of the work is in showing that X remains non meager in the final model. The dual problem for the null ideal will be dealt with in a forthcoming work.

2 Proof

On notation: We sometimes identify $x \in 2^\omega$ by a real whose binary expansion is x . Addition is always the usual addition in \mathbb{R} . We also sometimes interpret $y \in \mathbb{R}$ as a member of 2^ω which is the binary expansion of the fractional part of y . The relevant point here is that these transformations preserve meager sets.

Assume CH. Let $S_\star = [\omega_2]^2$. Let $\mathcal{I} = \{S \subseteq S_\star : (\exists c : S \rightarrow \omega)(\forall A \in [\omega_2]^{\aleph_1})([A]^2 \subseteq S \implies |c[[A]^2]| \geq 2)\}$. Note that, by Erdős-Rado theorem, \mathcal{I} is a **proper** sigma ideal over S_\star .

Let $\langle S_\gamma : \gamma < \gamma_\star \rangle$ be a one-one listing of \mathcal{I} . Let \mathbb{P} add \aleph_2 Cohen reals $\langle x_\alpha : \alpha < \omega_2 \rangle$. In $V^\mathbb{P}$, let $\mathbb{Q} = \prod \{\mathbb{Q}_\gamma : \gamma < \gamma_\star\}$ be the finite support product where $\mathbb{Q}_\gamma = \mathbb{Q}_{S_\gamma}$ is a sigma centered forcing making the set $\{x_\alpha + x_\beta : \{\alpha, \beta\} \in S_\gamma\}$ meager. For $S \subseteq S_\star$, \mathbb{Q}_S is defined as follows: $p \in \mathbb{Q}_S$ iff

- (1) $p = (F_p, \bar{n}_p, \bar{\sigma}_p, N_p) = (F, \bar{n}, \bar{\sigma}, N)$
- (2) $F \subseteq [S]^2$ is finite
- (3) $\bar{n} = \langle n_k : k \leq N \rangle$ is an increasing sequence of integers with $n_0 = 0$ and $n_{k+1} - n_k > 2^{n_k - n_{k-1}}$
- (4) $\bar{\sigma} = \langle \sigma_k : k < N \rangle$ where each $\sigma_k \in {}^{[n_k, n_{k+1})}2$

$p \leq q$ iff $F_p \subseteq F_q$, $\bar{n}_p \preceq \bar{n}_q$, $\bar{\sigma}_p \preceq \bar{\sigma}_q$ and for every $N_p \leq k < N_q$, for every $\{\alpha, \beta\} \in F_p$, $x_\alpha + x_\beta \upharpoonright [n_{q,k}, n_{q,k+1}) \neq \sigma_{q,k+1}$. It is clear that \mathbb{Q}_S is a sigma centered forcing adding a meager set covering $X_S = \{x_\alpha + x_\beta : \{\alpha, \beta\} \in S\}$. We write X for X_{S_\star} .

Note that the set of conditions $p = (p(0), p(1)) \in \mathbb{P} \star \mathbb{Q}$ such that for each $\gamma \in \text{dom}(p(1))$, $p(0)$ forces an actual value $p(1)(\gamma)$ and for every $\{\alpha, \beta\} \in F_{p(1)(\gamma)}$, $\{\alpha, \beta\} \subseteq \text{dom}(p(0))$ is dense in $\mathbb{P} \star \mathbb{Q}$. We will always assume that our conditions have this form.

Claim 2.1. *In $V^{\mathbb{P} \star \mathbb{Q}}$, whenever $Y \subseteq X$ is non meager, there are $y_1 < y_2 < y_3 < y_4$ in Y such that $y_2 - y_1 = y_4 - y_3$.*

Proof of Claim 2.1: Choose $S \subseteq S_*$ such that $Y = X_S$ is non meager and suppose p forces this. Let $S_1 = \{\{\alpha, \beta\} : (\exists p_{\alpha, \beta} \geq p)(p_{\alpha, \beta} \Vdash \{\alpha, \beta\} \in \mathring{S})\}$. So $S_1 \in \mathcal{I}^+$. Define an equivalence relation E on S_1 as follows: $\{\alpha_0, \beta_0\} E \{\alpha_1, \beta_1\}$ iff

- (a) $|\text{dom}(p_{\alpha_0, \beta_0}(i))| = |\text{dom}(p_{\alpha_1, \beta_1}(i))| = l_i$ for $i \in \{0, 1\}$. Let $\{\gamma_{\alpha_j, \beta_j, k}^i : k < l_i\}$ list $\text{dom}(p_{\alpha_j, \beta_j}(i))$ in increasing order for $i, j \in \{0, 1\}$
- (b) $p_{\alpha_0, \beta_0}(0)(\gamma_{\alpha_0, \beta_0, k}^0) = p_{\alpha_1, \beta_1}(0)(\gamma_{\alpha_1, \beta_1, k}^0)$ for each $k < l_0$.
- (c) $p_{\alpha_0, \beta_0}(1)(\gamma_{\alpha_0, \beta_0, k}^1)$ and $p_{\alpha_1, \beta_1}(0)(\gamma_{\alpha_1, \beta_1, k}^1)$ have the same $\bar{n}, \bar{\sigma}, N$ (but not necessarily F), for each $k < l_1$.

It is clear that E is an equivalence relation on S_1 with countably many equivalence classes. Since $S_1 \in \mathcal{I}^+$, we can choose $A \in [\omega_2]^{\aleph_1}$ such that $[A]^2 \subseteq S_1$ and for every $\{\alpha_0, \beta_0\}, \{\alpha_1, \beta_1\} \in [A]^2$, $\{\alpha_0, \beta_0\} E \{\alpha_1, \beta_1\}$. Let l_0, l_1 be the corresponding domain sizes. By Ramsey theorem, there is an infinite $A_1 \subseteq A$ such that whenever $\alpha_0 < \beta_0, \alpha_1 < \beta_1$ are from A_1 , for every $i \in \{0, 1\}$ and $k_0, k_1 < l_i$, the truth value of $\gamma_{\alpha_0, \beta_0, k_0}^i = \gamma_{\alpha_1, \beta_1, k_1}^i$ depends only on the order type of $\langle \alpha_0, \beta_0, \alpha_1, \beta_1 \rangle$. Choose $\alpha_1 < \alpha_2 < \alpha_3 < \alpha_4 \in A_1$ such that A_1 has at least two members between any two α_i 's. It is easy to check that the conditions $p_{\alpha_1, \alpha_3}, p_{\alpha_1, \alpha_4}, p_{\alpha_2, \alpha_3}, p_{\alpha_2, \alpha_4}$ have a least common extension q . For example, to see that p_{α_1, α_3} and p_{α_2, α_3} have a least common extension, choose some $\beta \in A_1 \cap (\alpha_2, \alpha_3)$ and use the fact that $\langle \alpha_1, \alpha_3, \beta, \alpha_3 \rangle$, $\langle \alpha_2, \alpha_3, \beta, \alpha_3 \rangle$ and $\langle \alpha_1, \alpha_3, \alpha_2, \alpha_3 \rangle$ have the same order type. Similarly, for p_{α_2, α_3} and p_{α_1, α_4} , choose $\beta_1 < \beta_2$ from $(\alpha_2, \alpha_3) \cap A_1$ and use that fact that $\langle \alpha_1, \alpha_4, \beta_1, \beta_2 \rangle$, $\langle \alpha_2, \alpha_3, \beta_1, \beta_2 \rangle$ and $\langle \alpha_1, \alpha_4, \alpha_2, \alpha_3 \rangle$ have the same order type. Now q forces that $x_{\alpha_1} + x_{\alpha_3}, x_{\alpha_1} + x_{\alpha_4}, x_{\alpha_2} + x_{\alpha_3}$ and $x_{\alpha_1} + x_{\alpha_4}$ are in Y and $(x_{\alpha_1} + x_{\alpha_3}) + (x_{\alpha_2} + x_{\alpha_4}) = (x_{\alpha_1} + x_{\alpha_4}) + (x_{\alpha_2} + x_{\alpha_3})$. \square

Claim 2.2. X is non meager in $V^{\mathbb{P} \star \mathbb{Q}}$.

Proof of Claim 2.2: Suppose not. Let $p_*, \langle \mathring{T}_m : m < \omega \rangle$ be such that $p_* \Vdash (\forall m)(\mathring{T}_m \subseteq {}^{<\omega}2$ is nowhere dense subtree) $\wedge \mathring{X} \subseteq \bigcup_m [\mathring{T}_m]$. Since $\mathbb{P} \star \mathbb{Q}$ is ccc, we can assume that each \mathring{T}_m is in $V^{\mathbb{P} \star \prod_{k \geq 1} \mathbb{Q}_k}$ where $\mathbb{Q}_k = \mathbb{Q}_{S_k}$ for some $S_k \in \mathcal{I}$. For each $\{\alpha, \beta\} \in S_*$, choose $p_{\alpha, \beta}, m(\alpha, \beta), k(\alpha, \beta), v(\alpha, \beta), l(\alpha, \beta), n(\alpha, \beta)$ etc. such that the following hold.

- (a) $p_{\alpha, \beta} \geq p_*, p_{\alpha, \beta} \Vdash (\mathring{x}_\alpha + \mathring{x}_\beta) \in [\mathring{T}_{m(\alpha, \beta)}]$
- (b) $p_{\alpha, \beta} = \langle p_{\alpha, \beta}(k) : k \leq k(\alpha, \beta) \rangle$ where $p_{\alpha, \beta}(0)$ is the Cohen part and $p_{\alpha, \beta}(k) \in \mathbb{Q}_k$

- (c) $\text{dom}(p_{\alpha,\beta}(0)) = v(\alpha, \beta)$, $|v(\alpha, \beta)| = l(\alpha, \beta)$ and $\alpha, \beta \in v(\alpha, \beta)$
- (c) For each $\gamma \in v(\alpha, \beta)$, $p_{\alpha,\beta}(0)(\gamma) \in {}^{n(\alpha,\beta)}2$
- (d) For each $1 \leq k \leq k(\alpha, \beta)$, $p_{\alpha,\beta}(k) = (F_{\alpha,\beta,k}, \bar{n}_{\alpha,\beta,k}, \bar{\sigma}_{\alpha,\beta,k}, N_{\alpha,\beta,k})$ where $F_{\alpha,\beta,k} \subseteq [v_{\alpha,\beta}]^2 \cap [S_k]^2$ and $n_{N_{\alpha,\beta,k}} = n(\alpha, \beta)$ does not depend on k

Let $\{\gamma_{\alpha,\beta,l} : l < l(\alpha, \beta)\}$ list $v(\alpha, \beta)$ in increasing order. Since $S_\star \in \mathcal{I}^+$, as before, we can choose $A \in [\omega_2]^{\aleph_0}$ such that for every $\alpha < \beta$ from A the following hold.

- (1) $\{\alpha, \beta\} \notin \bigcup_{k \geq 1} S_k$
- (2) $m(\alpha, \beta) = m_\star$, $k(\alpha, \beta) = k_\star$, $l(\alpha, \beta) = l_\star$, $n(\alpha, \beta) = n_\star$
- (3) For each $l < l_\star$, $p_{\alpha,\beta}(0)(\gamma_{\alpha,\beta,l}) = \eta_\star^l \in {}^{n_\star}2$
- (4) For each $1 \leq k \leq k_\star$, $\bar{n}_{\alpha,\beta,k} = \bar{n}_\star^k$, $\bar{\sigma}_{\alpha,\beta,k} = \bar{\sigma}_\star^k$, $N_{\alpha,\beta,k} = N_\star^k$
- (5) For all $\alpha_1 < \alpha_2$ and $\beta_1 < \beta_2$ from A and $l_1, l_2 < l_\star$, the truth value of $\gamma_{\alpha_1,\beta_1,l_1} = \gamma_{\alpha_2,\beta_2,l_2}$ depends only on the order type of $\langle \alpha_1, \alpha_2, \beta_1, \beta_2 \rangle$

Let $\langle \alpha_i : i < \omega \rangle$ be increasing members of A . Choose $n_{\star\star} > n_\star + (k_\star + l_\star + 10)!$. Choose $q_1 \geq p_{\alpha_0,\alpha_1}$ such that

- (i) $q_1 = \langle q_1(k) : k \leq k_\star \rangle$
- (ii) $\text{dom}(q_1(0)) = \text{dom}(p_{\alpha_0,\alpha_1})$ and for each $l < l_\star$, $q_1(0)(l) = \eta_\star^l \frown 0^{n_{\star\star} - n_\star}$
- (iii) For each $1 \leq k \leq k_\star$, $q_1(k) = (F_{\alpha_\star,\beta_\star,k}, \bar{n}_\star^{k \frown n_{\star\star}}, \bar{\sigma}_\star^{k \frown 01^{n_{\star\star} - n_\star - 2}0}, N_\star^k + 1)$

Since p_\star forces that $[T_{m_\star}^\circ]$ is nowhere dense, we can find $n_{\star\star\star} > n_{\star\star}$, $q_2 \geq q_1$ and $\rho \in [{}^{n_{\star\star\star}, n_{\star\star\star}}2$ such that the following hold.

- (a) $q_2 = \langle q_2(k) : k \leq K \rangle$ for some $k_\star \leq K < \omega$
- (b) For $1 \leq k \leq k_\star$, if $q_2(k) = (F_k, \bar{n}_k, \bar{\sigma}_k, N_k)$, then $n_{\star\star\star} < n_{k, N_k}$
- (c) $q_2 \Vdash (\forall x \in [T_{m_\star}^\circ])(x \upharpoonright [n_{\star\star\star}, n_{\star\star\star}) \neq \rho)$

For $j \geq 2$, consider the set $s_j = \{l < l_\star : (\exists l' < l_\star)(\gamma_{\alpha_0, \alpha_1, l} = \gamma_{\alpha_j, \alpha_{j+1}, l'})\}$. We claim that $s_j = s_\star$ is constant. To see this, suppose $2 \leq j_1 < j_2$. Choose j_3 much larger than j_2 and use the fact that the order types of $\langle \alpha_0, \alpha_1, \alpha_{j_i}, \alpha_{j_i+1} \rangle$, $\langle \alpha_0, \alpha_1, \alpha_{j_3}, \alpha_{j_3+1} \rangle$ and $\langle \alpha_{j_i}, \alpha_{j_i+1}, \alpha_{j_3}, \alpha_{j_3+1} \rangle$ are the same for $i \in \{1, 2\}$. It also follows that whenever $2 \leq j_1 < j_2 - 1$, $\{\gamma_{\alpha_{j_1}, \alpha_{j_1+1}, l} : l \in l_\star \setminus s_\star\} \cap \{\gamma_{\alpha_{j_2}, \alpha_{j_2+1}, l} : l \in l_\star \setminus s_\star\} = \emptyset$. So we can choose j large enough such that $(\text{dom}(q_2(0)) \cup \bigcup_{k \leq K} F_{q_2(k)}) \cap (\{\gamma_{\alpha_j, \alpha_{j+1}, l} : l \in l_\star \setminus s_\star\} \cup \{\alpha_j, \alpha_{j+1}\}) = \emptyset$. The next claim gives us the desired contradiction.

Claim 2.3. *For some q_3 , $q_3 \geq q_2$, $q_3 \geq p_{\alpha_j, \alpha_{j+1}}$ and $q_3 \Vdash \rho \subseteq \dot{x}_{\alpha_j} + \dot{x}_{\alpha_{j+1}}$*

Proof of Claim 2.3: Put $q_3 = \langle q_3(k) : k \leq K \rangle$ and $\text{dom}(q_3(0)) = \text{dom}(q_2(0)) \cup \{\gamma_{\alpha_j, \alpha_{j+1}, l} : l \in l_\star \setminus s_\star\}$. For each $l \in l_\star \setminus s_\star$, we would like to find $\eta_{\star\star}^l \succeq \eta_\star^l$ such that the following hold.

- (a) If $l_1 < l_2 \in l_\star \setminus s_\star$, $1 \leq k \leq k_\star$, $\{\gamma_{\alpha_j, \alpha_{j+1}, l_1}, \gamma_{\alpha_j, \alpha_{j+1}, l_2}\} \in S_k$ and $N_\star^k \leq i < N_{q_2(k)}$ then $(\eta_{\star\star}^{l_1} + \eta_{\star\star}^{l_2}) \upharpoonright [n_{q_2(k), i}, n_{q_2(k), i+1}] \neq \sigma_{q_2(k), i}$
- (b) If $l \in s_\star, l' \in l_\star \setminus s_\star$, $1 \leq k \leq k_\star$ and $N_\star^k \leq i < N_{q_2(k)}$ then $(q_2(0)(\gamma_{\alpha_0, \alpha_1, l}) + \eta_{\star\star}^{l'}) \upharpoonright [n_{q_2(k), i}, n_{q_2(k), i+1}] \neq \sigma_{q_2(k), i}$
- (c) If $\gamma_{\alpha_j, \alpha_{j+1}, l_1} = \alpha_j, \gamma_{\alpha_j, \alpha_{j+1}, l_2} = \alpha_{j+1}$ (so $l_1, l_2 \in l_\star \setminus s_\star$), then $(\eta_{\star\star}^{l_1} + \eta_{\star\star}^{l_2}) \upharpoonright [n_{\star\star}, n_{\star\star\star}] = \rho$

Here, for $\sigma, \tau \in 2^{<\omega}$, $m < n < \omega$ and $\rho : [m, n) \rightarrow 2$, by $(\sigma + \tau) \upharpoonright [m, n) \neq \rho$ we mean the following: For every $x \in [\sigma]$ and $y \in [\tau]$, $(x + y) \upharpoonright [m, n) \neq \rho$.

This would suffice since then we can let $q_3(0) = q_2(0) \cup \{(\gamma_{\alpha_j, \alpha_{j+1}, l}, \eta_{\star\star}^l) : l \in l_\star \setminus s_\star\}$ and for $1 \leq k \leq K$, $q_3(k) = (F_{q_2(k)} \cup F_{p_{\alpha_j, \alpha_{j+1}}(k)}, \bar{n}_{q_2(k)}, \bar{\sigma}_{q_2(k)}, N_{q_2(k)})$.

First put $\eta_{\star\star}^l \upharpoonright [n_\star, n_{\star\star}) = 0^{n_{\star\star} - n_\star}$ for every $l \in l_\star \setminus s_\star$. Next let $W = \{(k, i) : 1 \leq k \leq k_\star, N_\star^k + 1 \leq i < N_{q_2(k)}\}$. Note that for $(k, i) \in W$, $n_{q_2(k), i+1} - n_{q_2(k), i} > 2^{i - N_\star^k} (n_{\star\star} - n_\star) > 2^{i - N_\star^k} (k_\star + l_\star + 10)!$.

Inductively choose pairwise disjoint intervals $\langle I_{k,i} : (k, i) \in W \rangle$ such that each $I_{k,i} = [m_{k,i}, m_{k,i} + (l_\star + 5)!) \subseteq [n_{q_2(k), i}, n_{q_2(k), i+1})$. We claim that for each $(k, i) \in W$, we can choose $\langle \eta_{\star\star}^l \upharpoonright I_{k,i} : l \in l_\star \setminus s_\star \rangle$ such that the (k, i) -th instance of requirements (a), (b) are met. To see this, note that we have at most $\binom{l_\star - |s_\star|}{2} + |s_\star|(l_\star - |s_\star|)$ inequalities (coming from (a) and (b)) and one equality from (c) to satisfy and since $\{\alpha_j, \alpha_{j+1}\} \notin \bigcup \{S_k : k \geq 1\}$, there is no conflict between requirements (a) and (c). \square

References

- [1] P. Erdős, S. Kakutani, On non-denumerable graphs, Bull. Amer. Math. Soc., Vol. 49, Number 6 (1943), 457-461
- [2] P. Komjáth, Set theoretic constructions in Euclidean spaces, New Trends in Discrete and Computational Geometry (J. Pach, ed.), Springer, 1993, 303-325