

ON A CONJECTURE OF TARSKI ON PRODUCTS OF CARDINALS

THOMAS JECH¹ AND SAHARON SHELAH²

ABSTRACT^{3*}

We look at an old conjecture of A. Tarski on cardinal arithmetic and show that if a counterexample exists, then there exists one of length $\omega_1 + \omega$.

¹Supported partially by an NSF grant. I wish to express my gratitude to the Mathematical Institute of the Eidgenössische Technische Hochschule in Zürich for its hospitality during my visit.

²Publication #385. Supported partially by the B.S.F.

³AMS classification: 03E

*Keywords: Cardinal arithmetic, singular cardinals problem, *pcf*

In the early days of set theory, Hausdorff and Tarski established basic rules for exponentiation of cardinal numbers. In [T] Tarski showed that for every limit ordinal β , $\prod_{\xi < \beta} \aleph_\xi = \aleph_\beta^{|\beta|}$, and conjectured that

$$(1) \quad \prod_{\xi < \beta} \aleph_{\sigma_\xi} = \aleph_\alpha^{|\beta|}$$

holds for every ordinal β and every increasing sequence $\{\sigma_\xi\}_{\xi < \beta}$ such that $\lim_{\xi < \beta} \sigma_\xi = \alpha$. He remarked that (1) holds for every countable ordinal β .

Remarks. 1. The left hand side of (1) is less than or equal to the right hand side.

2. If β has $|\beta|$ disjoint cofinal subsets then the equality (1) holds. Thus the first limit ordinal that can be the length of a counterexample to (1) is $\omega_1 + \omega$.

[*Proof.* Let $\{A_i : i < |\beta|\}$ be disjoint cofinal subsets of β . Then

$$\prod_{\xi < \beta} \aleph_{\sigma_\xi} \geq \prod_{i < |\beta|} \prod_{\xi \in A_i} \aleph_{\sigma_\xi} \geq \prod_{i < |\beta|} \aleph_\alpha = \aleph_\alpha^{|\beta|}.]$$

It is not difficult to see that if one assumes the Singular Cardinals Hypothesis then (1) holds. With the hindsight given by results obtained in the last twenty years, it is also not difficult to find a counterexample to Tarski's conjecture. For instance, using the model described in [M], one can have an increasing sequence of cardinals of length $\beta = \omega_1 + \omega$ whose product does not satisfy (1). The purpose of this note is to show that if Tarski's conjecture fails then it fails in this specific way. Namely, if there is a counterexample then there is one of length $\omega_1 + \omega$.

The main result of this paper is the following:

Theorem. *A necessary and sufficient condition for Tarski's conjecture to fail is the existence of a singular cardinal \aleph_γ of cofinality \aleph_1 such that $\aleph_\gamma > \aleph_{\omega_1}^{\aleph_1}$ and $\aleph_\gamma^{\aleph_1} > \aleph_{\gamma+\omega}^{\aleph_0}$.*

If \aleph_γ is a cardinal that satisfies the condition then the sequence $\{\aleph_\xi\}_{\xi < \omega_1} \cup \{\aleph_{\gamma+n}\}_{n < \omega}$ is a counterexample to (1):

$$\prod_{\xi < \omega_1} \aleph_\xi \cdot \prod_{n < \omega} \aleph_{\gamma+n} = \aleph_{\omega_1}^{\aleph_1} \cdot \aleph_{\gamma+\omega}^{\aleph_0} < \aleph_{\gamma+\omega}^{|\omega_1+\omega|}.$$

Such a cardinal exists in one of Magidor's models, e.g. when $\aleph_\gamma = \aleph_{\omega_1+\omega_1}$ is a strong limit, $\aleph_{\omega_1+\omega_1}^{\aleph_1} = \aleph_{\omega_1+\omega_1+\omega+2}$ and $\aleph_{\omega_1+\omega_1+\omega}^{\aleph_0} = \aleph_{\omega_1+\omega_1+\omega+1}$.

Also, if $\lambda > \aleph_{\omega_1}$ is a strong limit singular cardinal of cofinality \aleph_1 such that $\lambda^{\aleph_1} > \lambda^{+(2^{\aleph_0})^+}$ then we have a counterexample as $(\lambda^{+\omega})^{\aleph_0} < \lambda^{+(2^{\aleph_0})^+}$ (by [ShA2, Ch. XIII, 5.1]).

The rest of this paper is devoted to the proof that the condition is necessary.

Assume that Tarski's conjecture fails, and let β be a limit ordinal for which there exists a sequence $\{\sigma_\xi\}_{\xi < \beta}$ that gives a counterexample:

$$(2) \quad \prod_{\xi < \beta} \aleph_{\sigma_\xi} < \aleph_\alpha^\kappa,$$

where

$$\kappa = |\beta| \text{ and } \alpha = \lim_{\xi < \beta} \sigma_\xi.$$

Lemma 1. *If (2) holds then $cf\beta < \kappa < \beta$, and there exists an ordinal $\gamma < \alpha$ such that $\aleph_\gamma^\kappa > \aleph_\alpha$.*

Proof. If (2) holds then β does not have $|\beta|$ disjoint cofinal subsets, and it follows that β is not a cardinal, and that $cf\beta < |\beta|$.

Assuming that $\aleph_\gamma^\kappa \leq \aleph_\alpha$ holds for all $\gamma < \alpha$, we pick a cofinal sequence $\{\alpha_i\}_{i < cf\beta}$ with limit α , and then

$$\aleph_\alpha^\kappa = \left(\sum_{i < cf\beta} \aleph_{\alpha_i} \right)^\kappa \leq \prod_{i < cf\beta} \aleph_{\alpha_i}^\kappa \leq \prod_{i < cf\beta} \aleph_\alpha = \aleph_\alpha^{cf\beta} = \prod_{i < cf\beta} \aleph_{\alpha_i} \leq \prod_{\xi < \beta} \aleph_{\sigma_\xi},$$

contrary to (2). \square

Now consider the shortest counterexample to Tarski's conjecture.

Lemma 2. *If β is the least ordinal for which (2) holds then $\beta = \kappa + \omega$ where κ is an uncountable cardinal.*

Proof. Without loss of generality, the sequence σ is continuous. (We can replace each σ_ξ by the limit of the sequence at ξ , for each limit ordinal ξ .)

Let $\kappa = |\beta|$. We claim that for every limit ordinal $\eta < \beta$, $\aleph_{\sigma_\eta}^\kappa < \aleph_\alpha$. If this were not true then, because $\beta > \kappa$, there would be a limit ordinal η such that $\kappa \leq \eta < \beta$ and that $\aleph_{\sigma_\eta}^{|\sigma_\eta|} \geq \aleph_\alpha^\kappa > \prod_{\xi < \eta} \aleph_{\sigma_\xi}$, which would make the sequence $\{\sigma_\xi\}_{\xi < \eta}$ a counterexample to Tarski's conjecture as well, contrary to the minimality of β .

Thus $\beta = \delta + \omega$ for some limit ordinal δ . It is clear that the sequence

$$\{\aleph_{\sigma_\xi} : \xi \leq \kappa \text{ or } \xi > \delta\}$$

of length $\kappa + \omega$ is also a counterexample, and by the minimality of β we have $\beta = \kappa + \omega$. \square

Now consider the least ordinal γ such that $\aleph_\gamma^\kappa > \aleph_\alpha$. We shall show that $cf\gamma = \kappa$ (and so κ is a regular uncountable cardinal). We also establish other properties of \aleph_γ .

Lemma 3. *If Tarski's conjecture fails, then there is a cardinal \aleph_γ of uncountable cofinality κ such that $\gamma > \kappa$, and that*

$$(3) \quad \text{for every } \nu < \gamma, \aleph_\nu^\kappa < \aleph_\gamma$$

$$(4) \quad \aleph_\gamma^\kappa > \aleph_{\gamma+\omega}^{\aleph_0}.$$

Proof. Let $\beta = \kappa + \omega$ be the least ordinal for which (2) holds, for some increasing continuous sequence $\{\sigma_\xi : \xi < \beta\}$ with limit α , and let γ be the least ordinal such that $\aleph_\gamma^\kappa > \aleph_\alpha$.

First we observe that for every $\nu < \gamma$, $\aleph_\nu^\kappa < \aleph_\gamma$. This is because if $\aleph_\nu^\kappa \geq \aleph_\gamma$ then $\aleph_\nu^\kappa \geq \aleph_\gamma^\kappa > \aleph_\alpha$, contradicting the minimality of γ .

As a consequence, we have $cf\gamma \leq \kappa$: otherwise, we would have $\aleph_\gamma^\kappa = \sum_{\nu < \gamma} \aleph_\nu^\kappa = \aleph_\gamma < \aleph_\alpha$, a contradiction. Also, if $\gamma = \lim_{i \rightarrow cf\gamma} \gamma_i$, then $\aleph_\gamma^\kappa = \left(\sum_{i < cf\gamma} \aleph_{\gamma_i} \right)^\kappa \leq \prod_{i < cf\gamma} \aleph_{\gamma_i}^\kappa \leq \prod_{i < cf\gamma} \aleph_\gamma = \aleph_\gamma^{cf\gamma}$ and so we have

$$\aleph_\gamma^{cf\gamma} = \aleph_\gamma^\kappa.$$

Since $\aleph_\alpha < \aleph_\gamma^\kappa$, we have $\aleph_\alpha^\kappa \leq \aleph_\gamma^\kappa = \aleph_\gamma^{cf\gamma} \leq \aleph_\alpha^{cf\gamma}$, and so $\aleph_\alpha^{cf\gamma} = \aleph_\alpha^\kappa$, and $\aleph_\alpha^{cf\gamma} > \prod_{\xi < \beta} \aleph_{\sigma_\xi}$. Hence the sequence

$$\{\aleph_{\sigma_\xi} : \xi \leq cf\gamma \text{ or } \xi > \kappa\}$$

of length $cf\gamma + \omega$ is also a counterexample, and it follows that $\kappa = cf\gamma$.

For every limit $\eta < \beta$ we have $\aleph_{\sigma_\eta}^\kappa < \aleph_\alpha$, and in particular $\aleph_{\sigma_\kappa}^\kappa < \aleph_\alpha$. Since $\aleph_\gamma^\kappa > \aleph_\alpha$, we have $\gamma > \kappa$. Finally,

$$\prod_{\xi < \beta} \aleph_{\sigma_\xi} = \prod_{\xi < \kappa} \aleph_{\sigma_\xi} \cdot \prod_{n < \omega} \aleph_{\sigma_{\kappa+n}} = \aleph_{\sigma_\kappa}^\kappa \cdot \aleph_\alpha^{\aleph_0} = \aleph_\alpha^{\aleph_0},$$

and because $\aleph_\gamma^\kappa = \aleph_\alpha^\kappa > \prod_{\xi < \beta} \aleph_{\sigma_\xi}$, we have $\aleph_\gamma^\kappa > \aleph_\alpha^{\aleph_0}$. Since $\alpha = \lim_{n \rightarrow \omega} \sigma_{\kappa+n} \geq \gamma + \omega$, we have

$$\aleph_\gamma^\kappa > \aleph_{\gamma+\omega}^{\aleph_0},$$

completing the proof. \square

The cardinal \aleph_γ obtained in Lemma 3 satisfies all the conditions stated in the Theorem except for the requirement that its cofinality be \aleph_1 . Thus the following lemma will complete the proof:

Lemma 4. *Let \aleph_γ be a singular cardinal of cofinality $\kappa > \aleph_1$ such that $\gamma > \kappa$ and that*

$$(5) \quad \text{for every } \nu < \gamma, \aleph_\nu^\kappa < \aleph_\gamma.$$

Assume further that for every $\delta, \omega_1 < \delta < \gamma$, of cofinality \aleph_1 ,

$$(6) \quad \text{if for every } \nu < \delta, \aleph_\nu^{\aleph_1} < \aleph_\delta, \text{ then } \aleph_\delta^{\aleph_1} \leq \aleph_{\delta+\omega}^{\aleph_0}.$$

Then $\aleph_\gamma^\kappa \leq \aleph_{\gamma+\omega}^{\aleph_0}$.

Lemma 4 implies that the least γ in Lemma 3 has cofinality \aleph_1 , and the theorem follows. The rest of the paper is devoted to the proof of Lemma 4. We use the second author's analysis of *pcf*.

Definition. If A is a set of regular cardinals, let

$$\Pi A = \{f : \text{dom } f = A \text{ and } f(\lambda) < \lambda \text{ for all } \lambda \in A\}.$$

If I is an ideal on A then $\Pi A/I$ is a partially ordered set under

$$f \leq_I g \text{ iff } \{\lambda : f(\lambda) > g(\lambda)\} \in I,$$

and similarly for filters on A . If D is an ultrafilter on A , then $\Pi A/D$ is a linearly ordered set, and $cf(\Pi A/D)$ denotes its cofinality. Let

$$pcf(A) = \{cf(\Pi A/D) : D \text{ an ultrafilter on } A\}.$$

It is clear that

$$A \subseteq pcf(A), A_1 \subseteq A_2 \text{ implies } pcf(A_1) \subseteq pcf(A_2), \text{ and} \\ pcf(A_1 \cup A_2) = pcf(A_1) \cup pcf(A_2),$$

and it is not difficult to show (using ultrapowers of ultrapowers) that

$$\text{if } |pcf(A)| < \min A \text{ then } pcf(pcf(A)) = pcf(A) \text{ and} \\ pcf(A) \text{ has a greatest element.}$$

Theorem (Shelah [Sh345]). *If $2^{|A|} < \min(A)$ then there exists a family $\{B_\nu : \nu \in pcf(A)\}$ of subsets of A such that*

$$(7) \quad \text{for every ultrafilter } D \text{ on } A, cf(\Pi A/D) = \text{the least } \nu \text{ such that } B_\nu \in D.$$

For every $\lambda \in pcf(A)$ there exists a family $\{f_\alpha : \alpha < \lambda\} \subseteq \Pi A$ such that

$$(8) \quad \alpha < \beta \text{ implies } f_\alpha < f_\beta \pmod{J_{<\lambda}}, \text{ where } J_{<\lambda} \text{ is the ideal generated by } \{B_\nu : \nu < \lambda\}, \text{ and the } f_\alpha \text{'s are cofinal in } \Pi B_\lambda \pmod{J_{<\lambda}}.$$

□

An immediate consequence of (7) is that $|pcf(A)| \leq 2^{|A|}$. The sets B_ν ($\nu \in pcf(A)$) are called *generators* for A . Note that $\max B_\nu = \nu$ when $\nu \in A$, and that $\max(pcf(B_\nu)) = \nu$ for all ν .

We shall use some properties of generators.

Lemma 5 [Sh345]. *Let B_ν be generators for A . For every $X \subseteq A$ there exists a finite set $F \subseteq pcf(X)$ such that $X \subseteq \bigcup\{B_\nu : \nu \in F\}$.*

Proof. Let $Y = pcf(X)$, and assume that the lemma fails. Then $\{X - B_\nu : \nu \in Y\}$ has the finite intersection property and so there is an ultrafilter D on A such that $X \in D$ and $B_\nu \notin D$ for all $\nu \in Y$. Let $\mu = cf(\Pi A/D)$. Then $\mu \in pcf(X)$ and by (7), $B_\mu \in D$. A contradiction. □

For each $X \subseteq A$, let $s(X)$ (a *support* of X) denote a finite set $F \subseteq pcf(X)$ with the property that $X \subseteq \bigcup_{\nu \in F} B_\nu$.

The set $pcf(A)$ has a set of generators that satisfy a transitivity condition:

Lemma 6 [Sh345]. *Assume that $2^{|A|} < \min(A)$ and let $\bar{A} = pcf(A)$. Then $pcf(\bar{A}) = \bar{A}$ and \bar{A} has a set of generators $\{B_\nu : \nu \in \bar{A}\}$ that satisfy, in addition to (7),*

$$(9) \quad \text{if } \xi \in B_\nu \text{ then } B_\xi \subseteq B_\nu.$$

□

We use the transitivity to prove the next lemma.

Lemma 7. *Assume that $2^{|A|} < \min(A)$, let $\bar{A} = pcf(A)$, let B_ν , $\nu \in \bar{A}$, be transitive generators for \bar{A} , and for each $X \subseteq \bar{A}$ let $s(X)$ be a support of X . If $A = \bigcup_{i \in I} A_i$, then*

$$\bar{A} = \bigcup \left\{ pcf(B_\nu) : \nu \in pcf \left(\bigcup_{i \in I} s(pcf(A_i)) \right) \right\}.$$

Corollary. $\max(\bar{A}) = \max pcf \bigcup_{i \in I} s(pcf(A_i))$.

[*Proof of Corollary.* Let $\lambda = \max(\bar{A})$; $\lambda \in pcf(B_\nu)$ for some ν in $pcf(\bigcup_i s(A_i))$. Since $\max(pcf(B_\nu)) = \nu$, we have $\lambda \leq \nu$.]

Proof. Let $X = \bigcup_{i \in I} s(pcf(A_i))$ and $F = s(X)$. We have

$$\begin{aligned} A &= \bigcup_{i \in I} A_i \subseteq \bigcup_{i \in I} pcf(A_i) \subseteq \bigcup_{i \in I} \bigcup \{B_\xi : \xi \in s(pcf(A_i))\} = \\ &= \bigcup \{B_\xi : \xi \in X\} \subseteq \bigcup \{B_\xi : \xi \in \bigcup_{\nu \in F} B_\nu\} \subseteq \bigcup_{\nu \in F} B_\nu \end{aligned}$$

(the last inclusion is a consequence of transitivity (9)). Therefore

$$\bar{A} = pcf(A) \subseteq pcf\left(\bigcup_{\nu \in F} B_\nu\right) = \bigcup_{\nu \in F} pcf(B_\nu) \subseteq \bigcup \{pcf(B_\nu) : \nu \in pcf(X)\}.$$

□

Toward the proof of Lemma 4, let $\{\gamma_i : i < \kappa\}$ be a continuous increasing sequence of limit ordinals of cofinality $< \kappa$, such that $\lim_{i \rightarrow \kappa} \gamma_i = \gamma$, $2^\kappa < \aleph_{\gamma_0}$, and that for all $i < \kappa$,

$$(10) \quad \text{for all } \nu < \gamma_i, \aleph_\nu^\kappa < \aleph_{\gamma_i}.$$

Lemma 8. *There is a closed unbounded set $C \subseteq \kappa$ such that for all $n = 1, 2, \dots$,*

$$(11) \quad \max pcf(\{\aleph_{\gamma_i+n} : i \in C\}) \leq \aleph_{\gamma+n}.$$

Proof. We show that for each n there exists a closed unbounded set $C_n \subseteq \kappa$ such that $\max pcf(\{\aleph_{\gamma_i+n} : i \in C_n\}) \leq \aleph_{\gamma+n}$. To prove this, let $n \geq 1$ be fixed and let $A = \{\aleph_{\gamma_i+n} : i < \kappa\}$. Let λ be the least element of $pcf(A)$ above $\aleph_{\gamma+n}$ (if there is none there is nothing to prove). Let $\{B_\nu : \nu \in pcf(A)\}$ be subsets of A that satisfy (7), and let $\{S_\nu : \nu \in pcf(A)\}$ be the subsets of κ such that $B_\nu = \{\aleph_{\gamma_i+n} : i \in S_\nu\}$. It suffices to prove that the set $S_{\aleph_{\gamma+1}} \cup \dots \cup S_{\aleph_{\gamma+n}}$ contains a closed unbounded set.

Thus assume that the set $S = \kappa - (S_{\aleph_{\gamma+1}} \cup \dots \cup S_{\aleph_{\gamma+n}})$ is stationary. Let $J_{<\lambda}$ be the ideal on A generated by $\{B_\nu : \nu < \lambda\}$. By Shelah's Theorem there exists a family $\{f_\alpha : \alpha < \lambda\}$ in ΠA such that $\alpha < \beta$ implies $f_\alpha < f_\beta \pmod{J_{<\lambda}}$. Since all the sets B_ν , $\nu < \aleph_\gamma$, are bounded, we get a family $\{g_\alpha : \alpha < \lambda\}$ of functions on S such that $g_\alpha(i) < \aleph_{\gamma_i+n}$ for all $i \in S$, and such that $\alpha < \beta$ implies that $g_\alpha(i) < g_\beta(i)$ for eventually all $i \in S$. This contradicts the results in [GH] by which, under the assumption (5), any family of almost disjoint functions in $\prod_{i \in S} \aleph_{\gamma_i+n}$ has size at most $\aleph_{\gamma+n}$. □

Proof of Lemma 4. Let γ be a singular cardinal of cofinality $\kappa > \aleph_1$ that satisfies (5) and (6). Let λ be a regular cardinal such that $\aleph_\gamma < \lambda \leq \aleph_\gamma^\kappa$. We shall prove that $\lambda \leq \aleph_{\gamma+\omega}^{\aleph_0}$.

Let $\{\gamma_i : i < \kappa\}$ be an increasing continuous sequence that satisfies (10), and let C be a closed unbounded subset of κ given by Lemma 8. Let

$$S = \{i \in C : cf \gamma_i = \aleph_1\}.$$

As $\kappa \geq \aleph_2$, S is a stationary subset of κ .

Lemma 9. *There exist regular cardinals λ_i , $i \in S$, such that for each $i \in S$, $\aleph_{\gamma_i} < \lambda_i \leq \aleph_{\gamma_i}^{\aleph_1}$, and an ultrafilter D on S such that $cf(\prod_{i \in S} \lambda_i / D) = \lambda$.*

Proof. Let I_0 be the nonstationary ideal on S . There are λ cofinal subsets X of ω_γ of size $|X| = \kappa$. For every such set X , let $F_X \in \prod_{i \in C} [\aleph_{\gamma_i}]^{\leq \kappa}$ be the function defined by $F(i) = X \cap \omega_{\gamma_i}$. Then when $X \neq Y$, F_X and F_Y are eventually distinct.

For every $i \in S$ we have $\aleph_{\gamma_i}^\kappa = \aleph_{\gamma_i}^{\aleph_1}$ (by (10)), and so there exist λ I_0 -distinct functions in $\prod_{i \in S} \aleph_{\gamma_i}^{\aleph_1}$. [f and g are I_0 -distinct if $\{i : f(i) = g(i)\} \in I_0$.]

Consider the partial ordering $f <_{I_0} g$ defined by $\{i : f(i) \geq g(i)\} \in I_0$; since I_0 is σ -complete, $<_{I_0}$ is well-founded. Let g be a $<_{I_0}$ -minimal function with the property that $g(i) \leq \aleph_{\gamma_i}^{\aleph_1}$ and that there are there are λ I_0 -distinct functions below g .

Let I be the extension of I_0 generated by all the stationary subsets X of S that have the property that g is not minimal on $I_0[X]$ (i.e. there is a function g' such that $g'(i) < g(i)$ almost everywhere on X and below g' there are λ I_0 -distinct functions).

Claim. I is a normal κ -complete ideal on S .

[*Proof.* Let X_i , $i < \kappa$, be sets in I , and let for each $i < \kappa$, $g_i < g$ on X_i and $\langle h_\xi^i : \xi < \lambda \rangle$ witness that $X_i \in I$. Then one constructs witnesses \bar{g} and $\langle \bar{h}_\xi : \xi < \lambda \rangle$ for $X = \{j \in \kappa : j \in \bigcup_{i < j} X_i\}$ by letting $\bar{g}(j) = g_i(j)$ and $\bar{h}_\xi(j) = h_\xi^i(j)$ where i is some $i < j$ such that $j \in X_i$.

For example, let us show that \bar{h}_ξ and \bar{h}_η are I_0 -distinct if $\xi \neq \eta$. Assume that $\bar{h}_\xi = \bar{h}_\eta$ on a stationary subset S_1 of S . Then on a stationary subset S_2 of S_1 the i less than $j \in S_2$ chosen such that $j \in X_i$ is the same i , and we have $h_\xi^i = h_\eta^i$ on S_2 , a contradiction.]

Let $\{h_\xi : \xi < \lambda\}$ be a family of I_0 -distinct functions below g .

Claim. For every $h <_I g$ there is some $\xi_0 < \lambda$ such that for all $\xi \geq \xi_0$, $h <_I h_\xi$.

[*Proof.* If there are λ many ξ 's such that $h \geq h_\xi$ on an I -positive set, then (because $2^\kappa < \lambda$) there is an I -positive set X such that $h \geq h_\xi$ on X for λ many ξ , but this contradicts the definition of I .]

Using this Claim, one can construct a $<_I$ -increasing λ -sequence (a subsequence of $\{h_\xi : \xi < \lambda\}$) of functions that is $<_I$ -cofinal in $\prod_{i \in S} g(i)$. Let $\lambda_i = cf g(i)$, for each $i \in S$. The product $\prod_{i \in S} \lambda_i$ has a $<_I$ -cofinal $<_I$ -increasing sequence of length λ , and since I is a normal ideal, we have $\lambda_i > \aleph_{\gamma_i}$ for I -almost all i . Now if D is any ultrafilter extending the dual of I , D satisfies $cf(\prod_{i \in S} \lambda_i / D) = \lambda$. \square

Back to the proof of Lemma 4. For each $i \in S$ we have a regular cardinal λ_i such that $\aleph_{\gamma_i} < \lambda_i \leq \aleph_{\gamma_i}^{\aleph_1}$. By the assumption (6) we have $\aleph_{\gamma_i}^{\aleph_1} \leq \aleph_{\gamma_i + \omega}^{\aleph_0}$, and so $\lambda_i \leq \aleph_{\gamma_i + \omega}^{\aleph_0}$. We use the following result:

Theorem (Shelah [ShA2], Chapter XIII, 5.1). *Let \aleph_δ be such that $\aleph_\delta^{\aleph_0} < \aleph_{\delta + \omega}$. Then for every regular cardinal μ such that $\aleph_\delta < \mu \leq \aleph_{\delta + \omega}^{\aleph_0}$ there is an ultrafilter U on ω such that $cf(\prod_{n \in \omega} \aleph_{\delta + n} / U) = \mu$. \square*

We apply the theorem to each \aleph_{γ_i} , and obtain for each $i \in S$ an ultrafilter U_i on ω such that $cf(\prod_{n \in \omega} \aleph_{\gamma_i + n} / U_i) = \lambda_i$. Combining the ultrafilters U_i with the ultrafilter D on S

from Lemma 9 we get an ultrafilter U on the set

$$A = \{\aleph_{\gamma_i+n} : i \in S, n = 1, 2, \dots\}$$

such that $cf(\Pi A/U) = \lambda$. Hence $\lambda \in pcf(A)$.

We shall now complete the proof of Lemma 4 by showing that $\max pcf(A) \leq \aleph_{\gamma+\omega}^{\aleph_0}$.

We have $A = \bigcup_{n=1}^{\infty} A_n$, where

$$A_n = \{\aleph_{\gamma_i+n} : i \in S\},$$

and since $2^{|A|} = 2^{\aleph} < \min(A)$, we apply the corollary of Lemma 7 and get

$$\max pcf(A) = \max pcf \bigcup_{n=1}^{\infty} s(pcf(A_n)),$$

where for each n , $s(pcf(A_n))$ is a finite subset of $pcf(pcf(A_n)) = pcf(A_n)$.

Let $E = \bigcup_{n=1}^{\infty} s(pcf(A_n))$. Since (by Lemma 8) $\max pcf(A_n) \leq \aleph_{\gamma+n}$ for each n , E is a countable subset of $\aleph_{\gamma+\omega}$. Hence $\max pcf(E) \leq \aleph_{\gamma+\omega}^{\aleph_0}$, and so

$$\lambda \leq \max pcf(A) = \max pcf(E) \leq \aleph_{\gamma+\omega}^{\aleph_0}.$$

□

REFERENCES

- [[**GH**]] Galvin and A. Hajnal, *Inequalities for cardinal powers*, Annals of Math. **101** (1975), 491–498.
- [[**M**]] I. Magidor, *On the singular cardinals problem I*, Israel J. Math. **28** (1977), 1–31.
- [[**ShA2**]] Shelah, *Proper Forcing*, Springer–Verlag Lecture Notes 940, 1982.
- [[**Sh345**]] —, *Products of regular cardinals and cardinals invariants of products of Boolean algebras*, Israel J. Math. **70** (1990), 129–187.
- [[**Sh355**]] —, $\aleph_{\omega+1}$ has a Jonsson algebra, preprint.
- [[**T**]] A. Tarski, *Quelques théorèmes sur les alephs*, Fundamenta Mathematicae **7** (1925), 1–14.

T. JECH, DEPARTMENT OF MATHEMATICS, THE PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA 16802

S. SHELAH, DEPARTMENT OF MATHEMATICS, THE HEBREW UNIVERSITY, JERUSALEM, ISRAEL