

**CHARACTERIZING AN \aleph_ϵ -SATURATED
MODEL OF SUPERSTABLE NDOP
THEORIES BY ITS $\mathbb{L}_{\infty, \aleph_\epsilon}$ -THEORY
SH401**

SAHARON SHELAH

The Hebrew University of Jerusalem
Einstein Institute of Mathematics
Edmond J. Safra Campus, Givat Ram
Jerusalem 91904, Israel

Department of Mathematics
Hill Center-Busch Campus
Rutgers, The State University of New Jersey
110 Frelinghuysen Road
Piscataway, NJ 08854-8019 USA

MSRI
Berkeley, CA USA

ABSTRACT. Assume a complete countable first order theory is superstable with NDOP. We knew that any \aleph_ϵ -saturated model of the theory is \aleph_ϵ -prime over a non-forking tree of “small” models and its isomorphism type can be characterized by its $\mathbb{L}_{\infty, \aleph_\epsilon}$ (dimension quantifiers)-theory; or if you prefer - appropriate cardinal invariants. We go here one step further providing cardinal invariants which are as finitary as seems reasonable.

Partially supported by the United States-Israel Binational Science Foundation and the NSF, and I thank Alice Leonhardt for the beautiful typing.

Revised with proofreading for the Journal

Done Oct/89

Publ No. 401

Latest Revision - 05/Dec/20

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{T}\mathcal{E}\mathcal{X}$

INTRODUCTION

After the main gap theorem was proved (see [Sh:c]), in discussion, Harrington expressed a desire for a finer structure - of finitary character (when we have a structure theorem at all). I point out that the logic $\mathbb{L}_{\infty, \aleph_0}(\text{d.q.})$ (where d.q. stands for dimension quantifier) does not suffice: suppose; e.g. for $T = \text{Th}(\lambda \times {}^\omega 2, E_n)_{n < \omega}$ where $(\alpha, \eta)E_n(\beta, \nu) =: \eta \upharpoonright n = \nu \upharpoonright n$ and for $S \subseteq {}^\omega 2$ define $M_S = M \upharpoonright \{(\alpha, \eta) : [\eta \in S \Rightarrow \alpha < \omega_1] \text{ and } [\eta \in {}^\omega 2 \setminus S \Rightarrow \alpha < \omega]\}$. Hence, it seems to me we should try $\mathbb{L}_{\infty, \aleph_\epsilon}(\text{d.q.})$ (essentially, in \mathfrak{C} we can quantify over sets which are included in the algebraic closure of finite sets, see below 1.1, 1.3), and Harrington accepts this interpretation. Here the conjecture is proved for \aleph_ϵ -saturated models.

I.e., the main theorem is $M \equiv_{\mathbb{L}_{\infty, \aleph_\epsilon}(\text{d.q.})} N \Leftrightarrow M \cong N$ for \aleph_ϵ -saturated models of a superstable countable (first order) theory T without dop. For this we analyze further regular types, define a kind of infinitary logic (more exactly, a kind of type of \bar{a} in M), “looking only up” in the definition (when thinking of the decomposition theorem). Recall that for as \aleph_ϵ -saturated model M of a superstable DNOP theory an \aleph_ϵ -decomposition is $\langle M_\eta, a_\eta : \eta \in \mathcal{I} \rangle$ where

- (a) $I \subseteq {}^\omega \text{ord}$ is nonempty closed under initial segments
- (b) $M_\eta \prec M$ is \aleph_ϵ -saturated
- (c) $\nu \triangleleft \eta \in I \Rightarrow M_\nu \prec M_\eta$
- (d) if $\nu = \eta \hat{\ } \langle \alpha \rangle \in I$ then M_ν is \aleph_ϵ -prime over $M_\eta \cup \{a_\nu\}$ and $\text{tp}(a_\eta, M_\eta)$ is orthogonal to M_ρ for $\rho \triangleleft \nu$ and (the last is not essential but clarifies)
- (e) $\langle M_\eta : \eta \in I \rangle$ is non-forking enough: for every $\nu \in I$ the set $\{a_\eta : \eta \in \text{Suc}_I(\nu)\} \subseteq M$ is independent over M_ν .

The point is that if $\eta = \nu \hat{\ } \langle \alpha \rangle$, M_{η_ν}, a_η are chosen then to a large extent $\langle M_\rho, a_\rho : \eta \triangleleft \rho \in I \rangle$ is determined. But the amount of “to a large extent” which suffices in [Sh:c], is not sufficient here, we need to find a finer understanding. In particular, we certainly do not like to “know” (M_ν, a_η) . So we consider a pair (A, B) where $A \subseteq M_\nu, A \cup \{a_\eta\} \subseteq B \subseteq M_\eta$, $\text{stp}_*(B, A) \vdash \text{stp}_*(B, M_\nu)$ and we try to define the type of such pairs in a way satisfying:

- (a) it can be impressed in our logic $\mathbb{L}_{\infty, \aleph_\epsilon}$
- (b) it expresses the essential information in $\langle M_\rho, a_\rho : \eta \triangleleft \rho \in I \rangle$.

To carry out the isomorphism proof we need: (1.27) the type of the sum is the sum of types (infinitary types) assuming first order independence. The main point of the proof is to construct an isomorphism between M_1 and M_2 when $M_1 \equiv_{\mathbb{L}_{\infty, \aleph_\epsilon}(\text{d.q.})} M_2, \text{Th}(M_\ell) = T$ where T and $\equiv_{\mathbb{L}_{\infty, \aleph_\epsilon}(\text{q.d.})}$ are as above. So by [Sh:c, X] it is

enough to construct isomorphic decompositions. The construction of isomorphic decompositions is by ω approximations, in stage n , $\sim n$ levels of the decomposition tree are approximated, i.e. we have $I_n^\ell \subseteq {}^{n \geq} \text{Ord}$ and $\bar{a}_\eta^{n,\ell} \in M_\ell$ for $\eta \in I_n, \ell = 1, 2$ such that $\text{tp}(\bar{a}_{\eta \upharpoonright 0}^{n,1} \hat{\ } \bar{a}_{\eta \upharpoonright 1}^{n,1} \hat{\ } \dots \hat{\ } \bar{a}_\eta^{n,1}, \emptyset, M) = \text{tp}(\bar{a}_{\eta \upharpoonright 0}^{n,2} \hat{\ } \bar{a}_{\eta \upharpoonright 1}^{n,2} \hat{\ } \dots \hat{\ } \bar{a}_\eta^{n,2}, \emptyset, M_2)$ with $\bar{a}_\eta^{n,\ell}$ being ε -finite, so in stage $n + 1$, choosing $\bar{a}_{\langle \rangle}^{n+1,\ell}$ we cannot take care of all types $\bar{a}_{\langle \rangle}^{n+1,\ell} \hat{\ } \bar{a}_{\langle \alpha \rangle}^{\eta,\ell}$ so addition theorem takes care. So though we are thinking on \aleph_ε -decompositions (i.e., the M_η 's are \aleph_ε -saturated), we get just a decomposition.

In the end of §1 (in 1.37) we point up that the addition theorem holds in fuller generalization. In the second section we deal with a finer type needed for shallow T , in the appendix we discuss how absolute is the isomorphism type.

Of course, we may consider replacing “ \aleph_ε -saturated models of an NDOP superstable countable T ” by “models of an NDOP \aleph_0 -stable countable T ”. But the use of ε -finite sets seems considerably less justifiable in this context, it seems more reasonable to use finite sets, i.e., $\mathbb{L}_{\infty, \aleph_0}(\text{d.q.})$. But subsequently Hrushovski and Bouscaren proved that even if T is \aleph_0 -stable, $\mathbb{L}_{\infty, \aleph_0}(\text{d.q.})$ is not sufficient to characterize models of T up to isomorphism. This is not sufficient even if one considers the class of all \aleph_ε -saturated models rather than all models. The first example is \aleph_0 -stable shallow of depth 3, and the second one is superstable (non \aleph_0 -stable), NOTOP, non-multi-dimensional.

If we deal with \aleph_ε -saturated models of shallow (superstable NDOP) theories T , we can bound the depth of the quantification $\gamma = DP(T)$; i.e. $\mathbb{L}_{\infty, \aleph_\varepsilon}^\gamma$ suffice.

We assume the reader has a reasonable knowledge of [Sh:c, V, §1, §2] and mainly [Sh:c, V, §3] and of [Sh:c, X].

Here is a slightly more detailed guide to the paper. In 1.1 we define the logic $\mathbb{L}_{\infty, \aleph_\varepsilon}$ and in 1.3 give a back and forth characterization of equivalence in this logic which is the operative definition for this paper.

The major tools are defined in 1.7, 1.11. In particular, the notion of tp_α defined in 1.5 is a kind of a depth α look-ahead type which is actually used in the final construction. In 1.28 we point out that equivalence in the logic $\mathbb{L}_{\infty, \aleph_\varepsilon}$ implies equivalence with respect to tp_α for all α . Proposition 1.14 contains a number of important concrete assertions which are established by means of Facts 1.16-1.23. In general these explain the properties of decompositions over a pair $\binom{B}{A}$. Claim 1.27 (which follows from 1.26) is a key step in the final induction. Definition 1.30 establishes the framework for the proof that two \aleph_ε -saturated structures that have the same tp_∞ are isomorphic. The induction step is carried out in 1.35.

I thank Baldwin for reading the typescript pointing needed corrections and writing down some explanations.

-1.1 Notation: The notation is of [Sh:c], with the following additions (or reminders).

If $\eta = \nu \hat{\ } \langle \alpha \rangle$ then we let $\eta^- = \nu$; for I a set of sequences ordinals we let

$\text{Suc}_T(\eta) = \{\nu : \text{for some } \alpha, \nu = \eta \hat{\ } \langle \alpha \rangle \in I\}$.

We work in \mathfrak{C}^{eq} and for simplicity every first order formula is equivalent to a relation.

- (1) \perp means orthogonal (so q is $\perp p$ means q is orthogonal to p), remember $p \perp A$ means p orthogonal to A ; i.e. $p \perp q$ for every $q \in S(\text{acl}(A))$ (in \mathfrak{C}^{eq})
- (2) \perp_a means almost orthogonal
- (3) \perp_w means weakly orthogonal
- (4) $\frac{\bar{a}}{B}$ and \bar{a}/B means $\text{tp}(\bar{a}, B)$
- (5) $\frac{A}{B}$ or A/B means $\text{tp}_*(A, B)$
- (6) $A + B$ means $A \cup B$
- (7) $\bigcup_A \{B_i : i < \alpha\}$ means $\{B_i : i < \alpha\}$ is independent over A
- (8) $A \bigcup_B C$ means $\{A, C\}$ is independent over B
- (9) $\{C_i : i < \alpha\}$ is independent over (B, A) means that¹
 $j < \alpha \Rightarrow \text{tp}_*(C_j, \bigcup \{C_i \cup B : i \neq j\})$ does not fork over A
- (10) regular type means stationary regular type $p \in S(A)$ for some A
- (11) for $p \in S(A)$ regular and C a set of elements realizing p , $\dim(C, p)$ is $\text{Max}\{|\mathbf{I}| : \mathbf{I} \subseteq C \text{ is independent over } A\}$
- (12) $\text{acl}(A) = \{c : \text{tp}(c, A) \text{ is algebraic}\}$
- (13) $\text{dcl}(A) = \{c : \text{tp}(c, A) \text{ is realized by one and only one element}\}$
- (14) $\text{Dp}(p)$ is depth (of a stationary type, see [Sh:c, X, Definition 4.3, p.528, Definition 4.4, p.529])
- (15) $\text{Cb}(p)$ is the canonical base of a stationary type p (see [Sh:c, III, 6.10, p.134])
- (16) B is \aleph_ε -atomic over A if for every finite sequence \bar{b} from A , for some find $A_0 \subseteq A$ we have $\text{stp}(\bar{b}, A_0) \vdash \text{stp}(\bar{b}, A)$, equivalently for some ε -finite $A_0 \subseteq \text{acl}(A)$ we have $\text{tp}(\bar{b}, A_0) \vdash \text{tp}(\bar{b}, \text{acl}(A))$.

¹Actually by the non-forking calculus this is equivalent to $\{C_i : i \leq \alpha\}$ is independent over A where we let $C_\alpha = B$.

§1 \aleph_ϵ -SATURATED MODELS

We first define our logic, but as said in §0, we shall only use the condition from 1.4. T is always superstable complete first order theory.

1.1 Definition. 1) The logic $\mathbb{L}_{\infty, \aleph_\epsilon}$ is slightly stronger than $\mathbb{L}_{\infty, \aleph_0}$, it consists of the set of formulas in $\mathbb{L}_{\infty, |T|^+}$ such that any subformula of ψ of the form $(\exists \bar{x})\varphi$ is actually the form

$$(\exists \bar{x}^0, \bar{x}^1) \left[\varphi_1(\bar{x}^1, \bar{y}) \ \& \ \bigwedge_{i < \ell g \bar{x}^1} (\theta_i(x_i^1, \bar{x}^0) \ \& \ (\exists^{< \aleph_0} z) \theta_i(z, \bar{x}^0)) \right],$$

with \bar{x}^0 finite, \bar{x}^1 not necessarily finite but of length $< |T|^+$; so φ “says” $\bar{x}^1 \subseteq acl(\bar{x}^0)$; note that always our final proof of the theorem uses $|T| \geq \aleph_0$.

2) $\mathbb{L}_{\infty, \aleph_\epsilon}(d.q.)$ is like $\mathbb{L}_{\infty, \aleph_\epsilon}$ but we have cardinality quantifiers and moreover dimensional quantifiers (as in [Sh:c, XIII,1.2,p.624]), see below.

3) The logic $\mathbb{L}_{\infty, \aleph_\epsilon}^\gamma$ consist of the formulas of $\mathbb{L}_{\infty, \aleph_\epsilon}$ such that φ has quantifier depth $< \gamma$ (but we start the inductive definition by defining the quantifier depth of all first order as zero).

4) $\mathbb{L}_{\infty, \aleph_\epsilon}^\gamma(d.q.)$ is like $\mathbb{L}_{\infty, \aleph_\epsilon}^\gamma$ but we have cardinality quantifiers and moreover dimensional quantifiers.

1.2 Remark. 1) In fact the dimension quantifier is used in a very restricted way (see Definition 1.9 and Claim 1.28 + Claim 1.30).

2) The reader may ignore this logic altogether and use just the characterization of equivalence in claim 1.4.

1.3 Convention. 1) T is a fixed first order complete theory, \mathfrak{C} is the “monster” model, as in [Sh:c], so is $\bar{\kappa}$ -saturated; \mathfrak{C}^{eq} is as in [Sh:c, III,6.2,p.131]. We work in \mathfrak{C}^{eq} so M, N vary on elementary submodels of \mathfrak{C}^{eq} of cardinality $< \bar{\kappa}$. We assume T is superstable with NDOP (countability is used only in the Proof of 1.5 for book-keeping, i.e., in the proof of 1.30 (and 1.29)).

Remember a, b, c, d denote members of \mathfrak{C}^{eq} , $\bar{a}, \bar{b}, \bar{c}, \bar{d}$ denote finite sequences of members of \mathfrak{C}^{eq} , A, B, C denote subsets of \mathfrak{C}^{eq} of cardinality $< \bar{\kappa}$.

Remember $acl(A)$ is the algebraic closure of A , i.e.

$\{b : \text{for some first order and } n < \omega, \varphi(x, \bar{y}) \text{ and } \bar{a} \subseteq A \text{ we have } \mathfrak{C}^{eq} \models \varphi[b, \bar{a}] \ \& \ (\exists \leq^n y) \varphi(y, \bar{a})\}$

and \bar{a} denotes $\text{Rang}(\bar{a})$ in places where it stands for a set (as in $acl(\bar{a})$). We write

$\bar{a} \in A$ instead of $\bar{a} \in {}^{\omega}A$.

2) A is ϵ -finite, if for some $\bar{a} \in {}^{\omega}A$, $A = acl(\bar{a})$. (So for stable theories a subset of an ϵ -finite set is not necessarily ϵ -finite but as T is superstable, a subset of an ϵ -finite set is ϵ -finite as if $B \subseteq acl(\bar{a})$, $\bar{b} \in B$ is such that $tp(\bar{a}, B)$ does not fork over \bar{b} , then trivially $acl(\bar{b}) \subseteq A$ and if $acl(\bar{b}) \neq B$, $tp_*(B, \bar{a} \wedge \bar{b})$ forks over B , hence ([Sh:c, III,0.1]) $tp(\bar{a}, B)$ forks over \bar{b} , a contradiction.

So if $acl(A) = acl(B)$, then A is ϵ -finite iff B is ϵ -finite).

3) When T is superstable by [Sh:c, IV, Table 1, p.169] for $\mathbf{F} = \mathbf{F}_{\aleph_0}^a$, all the axioms there hold and we write \aleph_ϵ instead of \mathbf{F} and may use implicitly the consequences in [Sh:c, IV, §3].

We may instead Definition 1.1 use directly the standard characterization from 1.4; as actually less is used we state the condition we shall actually use:

1.4 Claim. For models M_1, M_2 of T we have $M_1 \equiv_{\mathbb{L}_{\infty, \aleph_\epsilon}(\text{d.q.})} M_2$ iff

⊗ there is a non-empty family \mathcal{F} such that:

- (a) each $f \in \mathcal{F}$ is an (M_1, M_2) -elementary mapping, (so $\text{Dom}(f) \subseteq M_1$, $\text{Rang}(f) \subseteq M_2$)
- (b) for $f \in \mathcal{F}$, $\text{Dom}(f)$ is ϵ -finite (see 1.3(2)) above)
- (c) if $f \in \mathcal{F}$, $\bar{a}_\ell \in M_\ell$ ($\ell = 1, 2$) then for some $g \in \mathcal{F}$ we have:
 $f \subseteq g$ and $acl(\bar{a}_1) \subseteq \text{Dom}(f)$ and $acl(\bar{a}_2) \subseteq \text{Rang}(f)$
- (d) if $f \cup \{\langle a_1, a_2 \rangle\} \in \mathcal{F}$ and $tp(a_1, \text{Dom}(f))$ is stationary and regular then $\dim(\{a_1^1 \in M_1 : f \cup \{\langle a_1^1, a_2 \rangle\} \in \mathcal{F}\}, M_1)$
 $= \dim(\{a_2^1 \in M_2 : f \cup \{\langle a_1, a_2^1 \rangle\} \in \mathcal{F}\}, M_2)$.

Our main theorem is

1.5 Theorem. Suppose T is countable (superstable complete first order theory) with NDOP.

Then

- (1) the $\mathbb{L}_{\infty, \aleph_\epsilon}(\text{d.q.})$ theory of an \aleph_ϵ -saturated model characterizes it up to isomorphism.
- (2) Moreover, if M_1, M_2 are \aleph_ϵ -saturated models of T (so $M_\ell \prec \mathfrak{C}^{\text{eq}}$) and \otimes_{M_0, M_1} of 1.4 holds, then M_1, M_2 are isomorphic.

* * *

By 1.4, it suffices to prove part (2).

The proof is broken into a series of claims (some of them do not use NDOP, almost all do not use countability; but we assume T is superstable complete all the time (1.7(1)).

1.6 Discussion: Let us motivate the notation and Definition below.

Recall from the introduction that we are thinking of a triple (M, N, a) which may appear in \aleph_ε -decomposition $\langle M_\eta, a_\eta : \eta \in I \rangle$ of N , in the sense that for some $\eta \in I \setminus \{<\>\}$ we have $(M, M', a) = (M_{\eta^-}, M_\eta, a_\eta)$ so M, M' are \aleph_ε -saturated, $a_\eta \in M' \setminus M$, M' is \aleph_ε -prime over $M + a$ and $\text{tp}(a, M)$ is regular. But this is “too large for us” hence we consider an approximation (A, B) where $A \subseteq M (= M_{\eta^-})$, $A \subseteq B \subseteq M' (= M_\eta)$, $a = a_\eta \in B$ and $B/M (= B/M_{\eta^-})$ does not fork over A . We would like to define the α -type of (A, B) in N , which tries to say something on the decomposition above $(M, M', a) = (M_{\eta^-}, M_\eta, a_\eta)$, i.e., on $\langle M_\rho, a_\rho : \eta \triangleleft \rho \in I \rangle$. There are two natural “successor” of (A, B) we may choose in this context: the first 1.7 below replaces (A, B) to (A', B') such that $A \subseteq A' \subseteq M (= M_{\eta^-})$, $B \subseteq B' \subseteq M' (= M_\eta)$ and (as M' is \aleph_ε -prime over $M + a$) we have $\text{stp}_*(B', A' \cup B) \vdash \text{stp}(B', M)$, so $\text{tp}(B', A' \cup B)$ is almost orthogonal to A' ; we can think of this as “advancing in the same model”; in other words as A, B are ε -finite, we have to increase them in order to capture even (M, M') . This is formalized by \leq_a in Definition 1.7 below.

The second is to pass from (M_{η^-}, M_η, a) to (M_η, M_ν, a_ν) for some ν an immediate successor (in I) of $\eta \in I$. So the old B is included in the new A' and $B' = A' \cup \{a\}$ where $\text{tp}(a, A')$ is regular and is orthogonal to A (as in the decomposition we require $\text{tp}(a_\eta, M_{\eta^-})(M_\nu$ when $\nu \triangleleft \eta^-$). This is formalized by \leq_b in Definition 1.7 below.

1.7 Definition. 1) $\Gamma = \{(A, B) : A \subseteq B \text{ are } \varepsilon\text{-finite}\}$. Let

$\Gamma(M) = \{(A, B) \in \Gamma : A \subseteq B \subseteq M\}$.

2) For members (A, B) of Γ we may also write $\binom{B}{A}$; if $A \not\subseteq B$ we mean $\binom{B \cup A}{A}$.

3) $\binom{B_1}{A_1} \leq_a \binom{B_2}{A_2}$ (usually we omit a) iff (both are in Γ and)

$A_1 \subseteq A_2, B_1 \subseteq B_2, B_1 \cup A_2 \perp_a A_2$.

4) $\binom{B_1}{A_1} \leq_b \binom{B_2}{A_2}$ iff $A_2 = B_1, B_2 \setminus A_2 = \bar{b}$ and \bar{b}/A_2 is regular orthogonal to A_1 .

5) \leq^* is the transitive closure of $\leq_a \cup \leq_b$. (So it is a partial order, whereas in general $\leq_a \cup \leq_b$ and \leq_b are not).

6) We can replace A, B by sequences listing them (we do not always strictly distinguish).

Remark. The following observation may clarify.

1.8 Observation. If $\binom{B_1}{A_1} \leq^* \binom{B_2}{A_2}$ then we can find $\langle B'_\ell : \ell \leq n \rangle$ and

$\langle c_\ell : 1 \leq \ell < n \rangle$ for some $n \geq 1$, satisfying $\binom{B_1}{A_1} \leq_b \binom{B'_1}{B'_0}$, $c_\ell \in B'_{\ell+1}$, $\frac{c_\ell}{B'_\ell}$ regular, $\frac{B'_{\ell+1}}{c_\ell + B'_\ell} \perp_a B'_\ell$, $A_2 = B'_{n-1}$, $B_2 = B'_n$.

Remark. 1) Note that actually \leq_a is transitive. This means that in a sense \leq_b is enough, \leq_a inessential.

2) We may in 1.7(4) use $\bar{b} = \langle c \rangle$, does not matter.

Proof. By the definition of \leq^* there are $k < \omega$ and $\binom{B^\ell}{A^\ell}$ for $\ell \leq k$ such that: $\binom{B^\ell}{A^\ell} \leq_{x(\ell)} \binom{B^{\ell+1}}{A^{\ell+1}}$ for $\ell \leq k$ and $x(\ell) \in \{a, b\}$ and $\binom{B^0}{A^0} = \binom{B_1}{A_1}$, $\binom{B^k}{A^k} = \binom{B_2}{A_2}$ and without loss of generality $x(2\ell) = a, x(2\ell + 1) = b$. Let $N_0 \prec \mathfrak{C}$ be \aleph_ϵ -prime over \emptyset such that $A^0 \subseteq N_0$, $B_0 \cup A^0 \subseteq N_0$ and $f_0 = \text{id}_{A^0}$. We choose by induction on $\ell \leq k$, $N_{\ell+1}, f_{\ell+1}$ such that:

- (a) $\text{Dom}(f_{\ell+1}) = B^\ell$
- (b) $N_\ell \prec N_{\ell+1}$
- (c) if $x(\ell) = b$ then $f_{\ell+1}$ is an extension of f_ℓ (which necessarily has domain A_ℓ , check) with domain B^ℓ such that $f_\ell(B^\ell) \cup N_\ell \subseteq N_{\ell+1}$ and $N_{\ell+1}$ is \aleph_ϵ -prime over $N_\ell \cup f_\ell(B^\ell)$
- (d) if $x(\ell) = a$, then $f_{\ell+1}$ maps A^ℓ into $N_{\ell-1}$, B^ℓ into N_ℓ and $N_{\ell+1} = N_\ell$.

This is straightforward. Now on $\langle N_\ell : \ell \leq k + 1 \rangle$ we repeat the argument (of choosing $\langle B_\ell : \ell \leq n \rangle$) in the proof of 1.14(6) above, i.e., choose $B^\ell \subseteq N_\ell$ by downward induction on ℓ large enough as required. $\square_{1.8}$

1.9 Definition. 1) We define $\text{tp}_\alpha[\binom{B}{A}, M]$ (for $A \subseteq B \subseteq M$, A and B are ϵ -finite and α is an ordinal) and $\mathcal{S}_\alpha(\binom{B}{A}, M)$, $\mathcal{S}_\alpha(A, M)$ and $\mathcal{S}_\alpha^r(\binom{B}{A}, M)$, $\mathcal{S}_\alpha^r(A, M)$ by induction on α (we mean simultaneously; of course, we use appropriate variables):

- (a) $\text{tp}_0[\binom{B}{A}, M]$ is the first order type of $A \cup B$
- (b) $\text{tp}_{\alpha+1}[\binom{B}{A}, M] =$ the triple $\langle Y_{A,B,M}^{1,\alpha}, Y_{A,B,M}^{2,\alpha}, \text{tp}_\alpha(\binom{B}{A}, M) \rangle$
 where:

$$Y_{A,B,M}^{1,\alpha} =: \left\{ \text{tp}_\alpha[\binom{B'}{A'}, M] : \text{for some } A', B' \text{ we have } \binom{B}{A} \leq_a \binom{B'}{A'} \in \Gamma(M) \right\},$$

and $Y_{A,B,M}^{2,\alpha} =: \left\{ \langle \Upsilon, \lambda_{M,B}^\Upsilon \rangle : \Upsilon \in \mathcal{S}_\alpha^r(B, M) \right\}$

where $\lambda_{M,B}^\Upsilon = \dim \left[\left\{ d : \text{tp}_\alpha[(\begin{smallmatrix} B+d \\ B \end{smallmatrix}), M] = \Upsilon \right\}, B \right]$:

(c) for δ a limit ordinal, $\text{tp}_\delta[(\begin{smallmatrix} B \\ A \end{smallmatrix}), M] = \langle \text{tp}_\alpha[(\begin{smallmatrix} B \\ A \end{smallmatrix}), M] : \alpha < \delta \rangle$
(this includes $\delta = \infty$, really $\|M\|^+$ suffice).

(d) $\mathcal{S}_\alpha(A, M) = \left\{ \text{tp}_\alpha[(\begin{smallmatrix} B \\ A \end{smallmatrix}), M] : \text{for some } B \text{ such that } B \subseteq M, \right.$
 $\left. \text{and } (\begin{smallmatrix} B \\ A \end{smallmatrix}) \in \Gamma(M) \right\}$

(e) $\mathcal{S}_\alpha^r((\begin{smallmatrix} B \\ A \end{smallmatrix}), M) = \left\{ \text{tp}_\alpha[(\begin{smallmatrix} B+c \\ B \end{smallmatrix}), M] : \text{for some } c \in M \text{ we have} \right.$
 $\left. \frac{c}{B} \perp A \text{ and } \frac{c}{B} \text{ is regular} \right\}$

(f) $\mathcal{S}_\alpha^r(A, M) = \left\{ \text{tp}_\alpha[(\begin{smallmatrix} A+c \\ A \end{smallmatrix}), M] : c \in M \text{ and } \frac{c}{A} \text{ regular} \right\}$.

2) We define also $\text{tp}_\alpha[A, M]$, for A an ϵ -finite subset of M :

(a) $\text{tp}_0[A, M] =$ first order type of A

(b) $\text{tp}_{\alpha+1}[A, M]$ is the triple $\langle Y_{A,M}^{1,\alpha}, Y_{A,M}^{2,\alpha}, \text{tp}_\alpha[A, M] \rangle$ where
 $Y_{A,M}^{1,\alpha} =: \mathcal{S}_\alpha(A; M)$ and

$Y_{A,M}^{2,\alpha} =: \left\{ \langle \Upsilon, \dim\{d \in M : \text{tp}_\alpha[(\begin{smallmatrix} A+d \\ A \end{smallmatrix}), M] = \Upsilon\} \rangle : \Upsilon \in \mathcal{S}_\alpha^r(A, M) \right\}$

(c) $\text{tp}_\delta[A, M] = \langle \text{tp}_\alpha(A, M) : \alpha < \delta \rangle$

3) $\text{tp}_\alpha[M] = \text{tp}_\alpha[\emptyset, M]$.

1.10 Discussion: Clearly $\text{tp}[(\begin{smallmatrix} B \\ A \end{smallmatrix}), M]$ is intended, on the one hand, to be expressible by our logic and, on the other hand, to express the isomorphism type of M “in the direction of $(\begin{smallmatrix} B \\ A \end{smallmatrix})$ ”. To really say it we need to go back to the \aleph_ϵ -decompositions of M , a central notion of [Sh:c, Ch.X].

For the reader's benefit, by the referee request, let us review informally the proof in [Sh:c, Ch.X]. Let M be an \aleph_ϵ -saturated model, and we choose $\langle M_\eta : \eta \in I \cap {}^n\text{Ord} \rangle, \langle a_\eta : \eta \in I \cap {}^{n+1}\lambda \rangle$ by induction on n . For $n = 0$, of course, $I \cap {}^0\text{Ord} = \{ \langle \rangle \}$, we let $N_{\langle \rangle} \prec M$ be \aleph_ϵ -prime over \emptyset and let $\mathbf{I}_{\langle \rangle}$ a maximal subset of $\{c \in M : \text{tp}(c, N_{\langle \rangle}) \text{ regular}\}$ which is independent over $N_{\langle \rangle}$, let $\langle a_{\langle \rangle} : \alpha < |\mathbf{I}_{\langle \rangle}| \rangle$ list $\mathbf{I}_{\langle \rangle}$. Similarly for $n + 1, \eta \in I \cap {}^{n+1}\text{Ord}$ let $N_\eta \prec M$ be \aleph_ϵ -prime over $M_{\eta-} + a_\eta$, let \mathbf{I}_η be a maximal subset of $\{c \in M : \text{tp}(c, M_\eta) \text{ regular}\}$

orthogonal to M_{η^-} independent over N_η . Lastly, let $\langle c_{\eta \hat{<} \alpha} : \alpha < |\mathbf{I}_\eta| \rangle$ list \mathbf{I}_η and let $I \cap {}^{n+1}\text{Ord} = \{\eta \hat{<} \alpha : \eta \in I \cap {}^n\text{Ord} \text{ and } \alpha < |\mathbf{I}_\eta|\}$.

To carry this we use the existence of \aleph_ε -prime models (and the local character of independent). Also looking at the set $\cup\{M_\eta : \eta \in I\}$, its first order type is determined by the non-forking calculus. In fact, for any $\eta \in I \setminus \{\langle \rangle\}$, the set $\cup\{N_\nu : \eta \triangleleft \nu \in I\}, \cup\{N_\eta : \neg(\eta \leq \nu) \text{ and } \nu \in I\}$ are independent over N_η . Let $N \prec M$ be \aleph_ε -prime over $\cup\{N_\eta : \eta \in I\}$, now if $M = N$ we are done decomposing M , if not some $c \in M \setminus N$ realize a regular type (we use density of regular types). By NDOP, the $\text{tp}(c, N)$ is not orthogonal to some N_η . Choose η of minimal length hence $\nu \triangleleft \eta \Rightarrow \text{tp}(c, M_\eta) \perp N_\nu$. By properties of regular types, without loss of generality $\text{tp}(c, N)$ does not fork over N_η , so we get a contradiction to the maximality of $\{a_\nu : \nu \in \text{Suc}_I(\eta)\}$ (this explains the role of \mathcal{P} in Definition 1.11(5) below).

We are interested in the possible trees $\langle N_\nu : \eta \triangleleft \nu \in I \rangle$.

Now the tree determines M up to isomorphism, but there are “incidental” choices, so two trees may give isomorphic models (for investigating the number of non isomorphic models it is enough to find sufficiently pairwise far trees I).

Here we like to get exact information and in as finitary way as we can. So we replace $(M_{\eta^-}, M_\eta, a_\eta)$ by $\binom{B}{A}$, where $A \subseteq M_{\eta^-}, A + a_\eta \subseteq B \subseteq M_\eta$, $\text{tp}(B, M_{\eta^-})$ does not fork over A .

Now for $\eta \in I \setminus \{\langle \rangle\}$ we are interested in the possible trees $\langle N_\nu : \eta \triangleleft \nu \in I \rangle$, over $(N_{\eta_*}, N_\eta, a_\eta)$. But not only different trees may be equivalent (giving isomorphic \aleph_ε -prime models) but the other part of the tree, $\langle N_\nu : \nu \in I \text{ but } \neg(\eta \triangleleft \nu) \rangle$ may a priori cause non equivalent trees to contribute the same toward understanding M . This is done in [Sh:c, Ch.XII], but here we have to deal with ε -finite A, B .

The following claim 1.11 really does not add to [Sh:c, Ch.X], it just collects the relevant information which is proved there, or which follows immediately (particularly using the parameter (A, B)). We allow here a_η/M_η - to be not regular, but this is not serious: we can here deal exclusively with this case and we can omit this requirement in [Sh:c, Ch.X]; however, this does not eliminate the use of regular types (in the proof that M is \aleph_ε -prime over every \aleph_ε -decomposition of it).

1.11 Definition. 1) $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ε -decomposition inside M above (or for over) the pair $\binom{B}{A}$ (but we may omit the “ \aleph_ε - ”) if:

- (a) I a set of finite sequences of ordinals closed under initial segments
- (b) $\langle \rangle, \langle 0 \rangle \in I, \eta \in I \setminus \{\langle \rangle\} \Rightarrow \langle 0 \rangle \leq \eta$, let $I^- = I \setminus \{\langle \rangle\}$, really $a_{\langle \rangle}$ is meaningless
- (c) $A \subseteq N_{\langle \rangle}, B \subseteq N_{\langle 0 \rangle}, N_{\langle \rangle} \bigcup_A B$ and $dcl(a_{\langle 0 \rangle}) \subseteq dcl(B)$,
- (d) if $\nu = \eta \hat{<} \langle \alpha \rangle \in I$ then N_ν is \aleph_ε -primary over $N_\eta \cup \bar{a}_\nu, N_{\langle \rangle}$ is \aleph_ε -prime over A

- (e) for $\eta \in I$ such that $k = \ell g(\eta) > 1$ the type $a_\eta/N_{\eta \upharpoonright (k-1)}$ is orthogonal to $N_{\eta \upharpoonright (k-2)}$
- (f) $\eta \triangleleft \nu \Rightarrow N_\eta \prec N_\nu$
- (g) M is \aleph_ϵ -saturated and $N_\eta \prec M$ for $\eta \in I$
- (h) if $\eta \in I \setminus \{\langle \rangle\}$, then $\{a_\nu : \nu \in \text{Suc}_I(\eta)\}$ is (a set of elements realizing over N_η types orthogonal to N_{η^-} and is) an independent set over N_η .

2) We replace “inside M ” by “of M ” if in addition

- (i) in clause (h) the set is maximal.

3) $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition inside M if (a), (d), (e), (f), (g), (h) of part (1) holds and in clause (h) we allow $\eta = \langle \rangle$ (call this $(h)^+$). We add “over A ” if $A \subseteq M_{\langle \rangle}$.

4) $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition of M if in addition to 1.11(3) and we have the stronger version of clause (i) of 1.11(2) by including $\eta = \langle \rangle$, i.e. we have:

- (i)⁺ for $\nu \in I$, the set $\{a_\eta : \eta \in \text{Suc}_I(\nu)\}$ is a maximal subset of M independent over N_ν .

We may add “over A ” if $A \subseteq M$.

5) If $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition inside M we let

$$\mathcal{P}(\langle N_\eta, a_\eta : \eta \in I \rangle, M) = \left\{ p \in S(M) : p \text{ regular and for some } \eta \in I \setminus \{\langle \rangle\} \text{ we have} \right. \\ \left. p \text{ is orthogonal to } N_{\eta^-} \text{ but not to } N_\eta \right\}.$$

As said earlier it is natural to use regular types.

1.12 Definition. 1) We say that $\langle N_\eta, a_\eta : \eta \in I \rangle$, an \aleph_ϵ -decomposition inside M , is J -regular if $J \subseteq I$ and:

- (*) for each $\eta \in I \setminus J$ there² is c_η such that $a_\eta \in \text{acl}(N_{\eta^-} + c_\eta)$
 $\frac{c_\eta}{N_\eta}$ is regular and if $\eta \neq \langle \rangle$ then $\frac{a_\eta}{N_\eta + c_\eta} \perp_a N_{(\eta^-)}$.

2) We say “ $\langle N_\eta, a_\eta : \eta \in I \rangle$ is a regular \aleph_ϵ -decomposition inside M [of M]” if it is an \aleph_ϵ -decomposition inside M [of M] which is \emptyset -regular.

3) We say “ $\langle N_\eta, a_\eta : \eta \in I \rangle$ is a regular \aleph_ϵ -decomposition inside M [of M] over $\binom{B}{A}$ ” if it is an \aleph_ϵ -decomposition inside M [of M] over $\binom{B}{A}$ which is $\{\langle \rangle\}$ -regular.

²without loss of generality $c_\eta = a_\eta$

- 1.13 Claim.** 1) Every \aleph_ϵ -saturated model has an \aleph_ϵ -decomposition (i.e. of it).
 2) If M is \aleph_ϵ -saturated, $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition inside M , then for some J , and N_η, a_η for $\eta \in J \setminus I$ we have: $I \subseteq J$ and $\langle N_\eta, a_\eta : \eta \in J \rangle$ is an \aleph_ϵ -decomposition of M (even a $(J \setminus I)$ -regular one).
 3) If M is \aleph_ϵ -saturated, $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition of M then M is \aleph_ϵ -prime and \aleph_ϵ -minimal³ over $\bigcup_{\eta \in I} N_\eta$; if in addition $\langle N_\eta, a_\eta : \eta \in \{\langle \rangle, \langle 0 \rangle\} \rangle$ is an \aleph_ϵ -decomposition inside M above $\binom{B}{A}$, then $\langle N_\eta, a_\eta : \eta \in I \ \& \ (\eta \neq \langle \rangle \rightarrow \langle 0 \rangle \trianglelefteq \eta) \rangle$ is an \aleph_ϵ -decomposition of M above $\binom{B}{A}$.
 4) If $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition inside M above $\binom{B}{A}$, then it is an \aleph_ϵ -decomposition inside M .
 5) If $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition inside M [above $\binom{B}{A}$], $\eta \in I$, $[\eta \in I \setminus \{\langle \rangle\}]$, $\alpha = \text{Min}\{\beta : \eta \hat{\ } \langle \beta \rangle \notin I\}$, $\nu =: \eta \hat{\ } \langle \alpha \rangle$, $a_\nu \in M \setminus N_\eta$, $\frac{a_\nu}{N_\eta}$ is orthogonal to M_{η^-} if η^- if $\neq \langle \rangle$, $N_\nu \prec M$ is \aleph_ϵ -primary over $N_\eta + a_\nu$ and $a_\nu \uplus_{N_\eta} \left(\bigcup_{\rho \in I} N_\rho \right)$ (enough to demand $\{a_\rho : \rho^- = \eta \text{ and } \rho \in I\}$ is independent over a_ν/N_η) then $\langle N_\rho, a_\rho : \rho \in I \cup \{\nu\} \rangle$ is an \aleph_ϵ -decomposition inside M [over $\binom{B}{A}$].
 6) Assume $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition of M , if p is regular (stationary) and is not orthogonal to M (e.g. $p \in S(M)$) then for one and only one $\eta \in I$, there is a regular (stationary) $q \in S(N_\eta)$ not orthogonal to p such that: if η^- is well defined (i.e. $\eta \neq \langle \rangle$), then $p \perp N_{\eta^-}$.
 7) Assume $I = \bigcup_{\alpha < \alpha(*)} I_\alpha$, for each α we have $\langle N_\eta, a_\eta : \eta \in I_\alpha \rangle$ is an \aleph_ϵ -decomposition inside M [above $\binom{B}{A}$] and for each $\eta \in I$ for every $n < \omega$ and $\nu_\ell = \eta \hat{\ } \langle \beta_\ell \rangle \in I$ for $\ell < n$, for some α we have: $\{\nu_\ell : \ell < n\} \subseteq I_\alpha$ (e.g. I_α increasing). Then $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition inside M [above $\binom{B}{A}$].
 8) In (7), if $\eta \neq \langle \rangle$ and some ν_ℓ is not \triangleleft -maximal in I and $\frac{a_{\nu_\ell}}{N_\eta}$ is regular, it is enough:

$$\ell_1 < \ell_2 < n \Rightarrow \bigvee_{\alpha < \alpha(*)} [\{\nu_{\ell_1}, \nu_{\ell_2}\} \subseteq I_\alpha].$$

- 9) If $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition inside M , $I_1, I_2 \subseteq I$ are closed under initial segments and $I_0 = I_1 \cap I_2$ then $\left(\bigcup_{\eta \in I_1} N_\eta \right) \uplus_{\bigcup_{\eta \in I_0} N_\eta} \left(\bigcup_{\eta \in I_2} N_\eta \right)$.

³here we use NDOP

10) Assume that for $\ell = 1, 2$ that $\langle N_\eta^\ell, a_\eta^\ell : \eta \in I \rangle$ is an \aleph_ϵ -decomposition inside M_ℓ , and for $\eta \in I$ the function f_η is an isomorphism from N_η^1 onto N_η^2 and $\eta \triangleleft \nu \Rightarrow f_\eta \subseteq f_\nu$. Then $\bigcup_{\eta \in I} f_\eta$ is an elementary mapping; if in addition $\langle N_\eta^\ell, a_\eta^\ell : \eta \in I \rangle$ is an \aleph_ϵ -

decomposition of M_ℓ (for $\ell = 1, 2$) then $\bigcup_{\eta \in I} f_\eta$ can be extended to an isomorphism

from M_1 onto M_2 .

11) If $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition inside M (above $\binom{B}{A}$) and $M^- \prec M$ is \aleph_ϵ -prime over $\bigcup_{\eta \in I} N_\eta$ then $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition of M

(above $\binom{B}{A}$).

12) If $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition inside $M/$ of M [above $\binom{M}{A}$] and $a'_\eta \in N_\eta$ and N_η is \aleph_ϵ -prime over $N_{\eta^-} + a'_\eta$ for $\eta \in I \setminus \{ \langle \rangle \}$ [and $a'_{\langle 0 \rangle} = a_{\langle 0 \rangle}$ or at least $dcl(a'_{\langle 0 \rangle}) \subseteq dcl(B)$] then $\langle N_\eta, a'_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition inside $M/$ of M [above $\binom{B}{A}$].

Proof. 1), 2), 3), 5), 6), 9)-12). Repeat the proofs of [Sh:c, X]. (Note that here a_η/N_η is not necessarily regular, a minor change).

4), 7) Check.

8) As $Dp(p) > 0 \Rightarrow p$ is trivial, by [Sh:c, ChX,7.2,p.551] and [Sh:c, ChX,7.3]. $\square_{1.13}$

We shall prove:

1.14 Claim. 1) If M is \aleph_ϵ -saturated, $\binom{B}{A} \in \Gamma(M)$, then there is $\langle N_\eta, a_\eta : \eta \in I \rangle$, an \aleph_ϵ -decomposition of M above $\binom{B}{A}$.

2) Moreover if $\langle N_\eta, a_\eta : \eta \in I \rangle$ satisfies clauses (a) – (h) of Definition 1.11(1), we can extend it to satisfy clause (i) of 1.11(2), too.

3) If $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition of M above $\binom{B}{A}$, $M^- \prec M$ is \aleph_ϵ -prime over $\bigcup_{\eta \in I} N_\eta$ then:

(a) $\langle N_\eta : \eta \in I \rangle$ is a \aleph_ϵ -decomposition of M^-

(b) we can find an \aleph_ϵ -decomposition $\langle N_\eta, a_\eta : \eta \in J \rangle$ of M such that $J \supseteq I$ and $[\eta \in J \setminus I \Leftrightarrow (\eta \neq \langle \rangle \text{ and } \neg \langle 0 \rangle \triangleleft \eta)]$, moreover the last phrase follows from the previous ones.

4) If in 3)(b) the set $J \setminus I$ is countable (finite is enough for our applications), then necessarily M, M^- are isomorphic, even adding all members of an ϵ -finite subset of M^- as individual constants.

5) If $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition of M above $\binom{B}{A}$, $I \subseteq J$ and

$\langle N_\eta, a_\eta : \eta \in J \rangle$ is an \aleph_ϵ -decomposition of M , $M^- \prec M$ is \aleph_ϵ -prime over $\bigcup_{\eta \in I} N_\eta$

and $\binom{B}{A} \leq^* \binom{B_1}{A_1}$ and $B_1 \subseteq M$ and $c \in M$ and $\frac{c}{B_1} \perp A_1$ and $\frac{c}{B_1}$ is (stationary and) regular then

$$(\alpha) \frac{c}{B_1} \perp \frac{\cup\{N_\eta : \eta \in J \setminus I\}}{N_\langle \rangle}$$

(β) $\frac{c}{B_1}$ is not orthogonal to some $p \in \mathcal{P}(\langle N_\eta, a_\eta : \eta \in I \rangle, M^-)$.

6) If $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition of M above $\binom{B}{A}$, then the set $\mathcal{P} = \mathcal{P}(\langle N_\eta : \eta \in I \rangle, M)$ depends on $\binom{B}{A}$ and M only (and not on $\langle N_\eta : \eta \in I \rangle$ or M^- when $M^- \prec M$ is \aleph_ϵ -prime over $\cup\{N_\eta : \eta \in I\}$), recalling:

$$\mathcal{P} = \mathcal{P}(\langle N_\eta : \eta \in I \rangle, M) = \left\{ p \in S(M) : p \text{ regular and for some } \eta \in I \setminus \{ \langle \rangle \}, \text{ we have : } \right. \\ \left. p \text{ is orthogonal to } N_{\eta^-} \text{ but not to } N_\eta \right\}.$$

So let $\mathcal{P}(\binom{B}{A}, M) =: \mathcal{P}(\langle N_\eta : \eta \in I \rangle, M)$.

(7) If $\frac{B}{A}$ is regular of depth zero or just $\frac{b}{A} \leq_a \frac{B}{A}$, $\frac{b}{A}$ regular of depth zero and M is \aleph_ϵ -saturated and $B \subseteq M$ then

(a) for any α , we have $\text{tp}_\alpha(\binom{B}{A}, M)$ depend just on $\text{tp}_0(\binom{B}{A}, M)$

(b) if $\binom{B}{A} \leq^* \binom{B'}{A} \in \Gamma(M)$ then $\text{tp}_\alpha(\binom{B'}{A}, M)$ depends just on $\text{tp}_0(\binom{B}{A}, M)$ (and (A, B, A', B) but not on M).

8) For $\alpha < \beta$, from $\text{tp}_\beta(\binom{B}{A}, M)$ we can compute $\text{tp}_\alpha(\binom{B}{A}, M)$.

9) If f is an isomorphism from M_1 onto M_2 , $A_1 \subseteq B_1$ are ε -finite subsets of M_1 and $f(A_1) = A_2, f(B_1) = B_2$ then

$$\text{tp}_\alpha\left(\binom{B_1}{A_1}, M_1\right) = \text{tp}_\alpha\left(\binom{B_2}{A_2}, M_2\right)$$

(more pedantically $\text{tp}_\alpha(\binom{B_2}{A_2}, M_2) = f[\text{tp}_\alpha(\binom{B_1}{A_1}, M_1)]$ or considered the A_ℓ, B_ℓ as indexed sets).

We delay the proof (parts (1), (2), (3) are proved after 1.22, part (4), (6) after 1.23, and after it parts (5), (7), (8). Part (9) is obvious.

1.15 Definition. 1) If $\binom{B}{A} \in \Gamma(M)$, M is \aleph_ϵ -saturated let $\mathcal{P}_{\binom{B}{A}}^M$ be the set \mathcal{P} from Claim 1.14(6) above (by 1.14(6) this is well defined as we shall prove below).

2) Let $\mathcal{P}_{\binom{B}{A}} = \left\{ p : p \text{ is (stationary regular and) parallel to some } p' \in \mathcal{P}_{\binom{B}{A}}^{\text{ceq}} \right\}$.

1.16 Definition. If $\langle N_\eta^\ell, a_\eta : \eta \in J \rangle$ is a decomposition inside \mathfrak{C} for $\ell = 1, 2$ we say that $\langle N_\eta^1, a_\eta : \eta \in J \rangle \leq_{\text{direct}}^* \langle N_\eta^2, a_\eta : \eta \in J \rangle$ if:

(a) $N_{\langle \rangle}^1 \prec N_{\langle \rangle}^2$

(b) $N_{\langle \rangle}^2 \bigcup_{N_{\langle \rangle}^1} \{a_{\langle \alpha \rangle} : \langle \alpha \rangle \in J\}$

(c) for $\eta \in J \setminus \{\langle \rangle\}$, N_η^2 is \aleph_ϵ -prime over $N_\eta^1 \cup N_{\eta^-}$.

1.17 Claim. 1) M is \aleph_ϵ -prime over A iff M is \aleph_ϵ -primary over A iff M is \aleph_ϵ -saturated, $A \subseteq M$, M is \aleph_ϵ -atomic over A (see -1.1(16)) for every $\mathbf{I} \subseteq M$ indiscernible over A we have: $\dim(\mathbf{I}, M) \leq \aleph_0$ iff M is \aleph_ϵ -saturated, $A \subseteq M$, M is \aleph_ϵ -atomic over A and for every finite $B \subseteq M$ and regular (stationary) $p \in S(A \cup B)$, we have $\dim(p, M) \leq \aleph_0$.

2) If N_1, N_2 are \aleph_ϵ -prime over A , then they are isomorphic over A .

Proof. By [Sh:c, IV,4.18] (see Definition [Sh:c, IV,4.16], noting that we replace $\mathbf{F}_{\aleph_0}^a$ by \aleph_ϵ and that part (4) there disappears when we are speaking on $\mathbf{F}_{\aleph_0}^a$). $\square_{1.16}$

However, we need more specific information saying that “minor changes” preserve being \aleph_ϵ -prime; this is done in 1.18 below, parts of it are essentially done in [Sh 225] but we give full proof.

1.18 Fact. 0) If A is countable, N is \aleph_ϵ -primary over A then N is \aleph_ϵ -primary over \emptyset .

1) If N is \aleph_ϵ -prime over \emptyset , A countable, N^+ is \aleph_ϵ -prime over $N \cup A$ then N^+ is \aleph_ϵ -prime over \emptyset .

2) If $\langle N_n : n < \omega \rangle$ is increasing, each N_n is \aleph_ϵ -prime over \emptyset or just \aleph_ϵ -constructible over \emptyset and N_ω is \aleph_ϵ -prime over $\bigcup_{n < \omega} N_n$ then N_ω is \aleph_ϵ -prime over

\emptyset , (note that if each N_n is \aleph_ϵ -saturated then $N_\omega = \bigcup_{n < \omega} N_n$).

2A) If N is \aleph_ϵ -prime over C , $\bar{a} \wedge \bar{b} \subseteq N$, $\text{tp}(\bar{b}, \bar{a})$ is regular (stationary) and orthogonal to C then $\dim(\text{tp}(\bar{b}, \bar{a}), N) \leq \aleph_0$; also if $q \in S(C \cup \bar{a})$ is a non-forking extension

of $\text{tp}(\bar{b}, \bar{a})$ then $\dim(q, C \cup \bar{a}) = \dim(\text{tp}(\bar{b}, \bar{a}), N) = \aleph_0$.

2B) If $C \cup \bar{a} \hat{=} \bar{b} \subseteq N$ and \bar{a}/\bar{b} is a regular type orthogonal to C and $q \in S^{\text{lg}(\bar{a})}(N)$ is a non-forking extension of \bar{a}/\bar{b} then $\dim(p \upharpoonright (C + \bar{b}), N) \leq \dim(\bar{a}/\bar{b}, N) \leq \dim(p \upharpoonright (C + \bar{b}), N) + \aleph_0$; moreover, $\dim(p \upharpoonright (C + \bar{b}), N) \leq \dim(\bar{a}/\bar{b}, N) < \dim(p \upharpoonright (C + \bar{b}), N)^+ + \aleph_0$.

3) If $N_2 \amalg_{N_0} N_1$, each N_ℓ is \aleph_ϵ -saturated, N_2 is \aleph_ϵ -prime over $N_0 \cup \bar{a}$, and N_3 is

\aleph_ϵ -prime over $N_2 \cup N_1$ then N_3 is \aleph_ϵ -prime over $N_1 \cup \bar{a}$.

4) If $N_1 \prec N_2$ are \aleph_ϵ -primary over \emptyset then for some \aleph_ϵ -saturated $N_0 \prec N_1$ (necessarily \aleph_ϵ -primary over \emptyset) we have: N_1, N_2 are isomorphic over N_0 .

5) In part (4), if $A \subseteq N_1$ is ϵ -finite then we can demand $A \subseteq N_0$.

6) If M_0 is \aleph_ϵ -saturated, $A \amalg_{M_0} B, M_1$ is \aleph_ϵ -primary over $M_0 \cup A$ then $M_1 \amalg_{M_0} B$.

7) Assume $N_0 \prec N_1 \prec N_2$ are \aleph_ϵ -saturated, N_2 is \aleph_ϵ -primary over $N_1 + a$ and $\frac{a}{N_1} \perp N_0$ (and $a \notin N_1$). If $N'_0 \prec N_0, N'_0 \prec N'_1 \prec N_1, N'_1 \amalg_{N'_0} N_0$ and N_1 is \aleph_ϵ -

primary over $N_0 \cup N'_1, A_1^* \subseteq N'_1, A_2^* \subseteq N_2$ are ϵ -finite and $\text{tp}_*(A_2^*, N_1)$ does not fork over A_1^* , then we can find a', N'_2 such that: N'_2 is \aleph_ϵ -saturated, \aleph_ϵ -primary over $N'_1 + a', N'_1 \prec N'_2 \prec N_2, N_1 \amalg_{N'_1} N'_2$ and N_2 is \aleph_ϵ -primary over $N_1 \cup N'_2$ and $A_2^* \subseteq N'_2$.

8) Assume $N'_0 \prec N_0 \prec N_1$ and $a \in N_1$ and N_1 is \aleph_ϵ -prime over $N_0 + a$ and $\frac{a}{N_0} \perp N'_0$ and $A_0^* \subseteq N'_0, A_1^* \subseteq N_1$ are ϵ -finite and $\text{tp}_*(A_1^*, N_0)$ does not fork over A_0^* then we can find a', N'_1 such that $a' \in N', N'_0 \prec N'_1 \prec N_1, N'_1 \amalg_{N'_0} N_0, N'_1$ is \aleph_ϵ -prime over

$N'_0 + a$ and N_1 is \aleph_ϵ -prime over $N_0 + N'_1$ and $A_1^* \subseteq N'_1$.

9) If N_1 is \aleph_ϵ -prime over \emptyset and $A \subseteq B \subseteq N_1$ and A, B are ϵ -finite, then we can find N_0 such that: $A \subseteq N_0 \prec N_1, N_0$ is \aleph_ϵ -prime over $\emptyset, A \subseteq N_0, B \amalg_A N_0$, and N_1

is \aleph_ϵ -prime over $N_0 \cup B$.

10) If N_0 is \aleph_ϵ -prime over A and $B \subseteq N_0$ is ϵ -finite, then N_0 is \aleph_ϵ -prime over $A \cup B$ (and also over A' if $A \subseteq A' \subseteq \text{acl}(A)$).

1.19 Remark. In the proof of 1.18(1)-(6),(10) we do not use “ T has NDOP”.

Proof. 0) There is $\{a_\alpha : \alpha < \alpha^*\}$, a list of members of N in which every member of $N \setminus A$ appears such that for $\alpha < \alpha^*$ we have: $\text{tp}(a_\alpha, A \cup \{a_\beta : \beta < \alpha\})$ is \aleph_ϵ -isolated (which means just $\mathbf{F}_{\aleph_0}^\alpha$ -isolated).

[Why? by the definition of “ N is \aleph_ϵ -primary over A ”). Let $\{b_n : n < \omega\}$ list A (if $A = \emptyset$ the conclusion is trivial so without loss of generality $A \neq \emptyset$, hence we can find such a sequence $\langle b_n : n < \omega \rangle$). Now define $\beta^* = \omega + \beta$ and $b_{\omega+\alpha} = a_\alpha$

for $\alpha < \alpha^*$. So $\{b_\beta : \beta < \beta^*\}$ lists the elements of N (possibly with repetitions, remember $A \subseteq N$ and check). We claim that $\text{tp}(b_\beta, \{b_\gamma : \gamma < \beta\})$ is $\mathbf{F}_{\aleph_0}^\alpha$ -isolated for $\beta < \beta^*$.

(Why? if $\beta \geq \omega$, let $\beta' = \beta - \omega$ (so $\beta < \alpha^*$), now the statement above means $\text{tp}(a_{\beta'}, A \cup \{a_\gamma : \gamma < \beta'\})$ is $\mathbf{F}_{\aleph_0}^\alpha$ -isolated which we know; if $\beta < \omega$ this statement is trivial). By the definition of “ $\mathbf{F}_{\aleph_0}^\alpha$ -primary”, clearly $\langle b_\beta : \beta < \omega + \alpha \rangle$ exemplify that N is $\mathbf{F}_{\aleph_0}^\alpha$ -primary over \emptyset .

1) Note

(*)₁ if N is \aleph_ϵ -primary over \emptyset and $A \subseteq N$ is finite then N is \aleph_ϵ -primary over A [why? see [Sh:c, IV,3.12](3),p.180 (of course, using [Sh:c, IV, Table 1,p.169] for $\mathbf{F}_{\aleph_0}^\alpha$]

(*)₂ if N is \aleph_ϵ -primary over \emptyset , $A \subseteq N$ is finite and $p \in S^m(N)$ does not fork over A and $p \upharpoonright A$ is stationary then for some $\{\bar{a}_\ell : \ell < \omega\}$ we have: $\bar{a}_\ell \in N$ realize p , $\{\bar{a}_\ell : \ell < \omega\}$ is independent over A and $p \upharpoonright (A \cup \bigcup_{\ell < \omega} \bar{a}_\ell) \vdash p$

[why? [Sh:c, IV,proof of 4.18] (i.e. by it and [Sh:c, 4.9](3),4.11) or let N' be \aleph_ϵ -primary over $A \cup \bigcup_{\ell < \omega} \bar{a}_\ell$ and note: N' is \aleph_ϵ -primary over A (proof

like the one of 1.18(0)) but also N is \aleph_ϵ -primary over A so by uniqueness of the \aleph_ϵ -primary model N' is isomorphic to N over A , so without loss of generality $N' = N$; and easily N' is as required].

Now we can prove 1.18(1), for any $\bar{c} \in {}^{\omega>}A$, we can find a finite $B_{\bar{a}}^1 \subseteq N$ such that $\text{tp}(\bar{c}, N)$ does not fork over $B_{\bar{c}}^1$, let $\bar{b}_{\bar{c}} \in {}^{\omega>}N$ realize $\text{stp}(\bar{a}, B_{\bar{a}}^1)$ and let $B_{\bar{c}} = B_{\bar{c}}^1 \cup \bar{b}_{\bar{c}}$, so $\text{tp}(\bar{c}, N)$ does not fork over $B_{\bar{c}}$ and $\text{tp}(\bar{c}, B_{\bar{c}})$ is stationary, hence we can find $\langle \bar{a}_\ell^{\bar{c}} : \ell < \omega \rangle$ as in (*)₂ (for $\text{tp}(\bar{c}, B_{\bar{c}})$). Let

$A' = \cup\{B_{\bar{c}} : \bar{c} \in {}^{\omega>}A\} \cup \{\bar{a}_\ell^{\bar{c}} : \bar{c} \in {}^{\omega>}A \text{ and } \ell < \omega\}$, so A' is a countable subset of N and $\text{tp}_*(A, A') \vdash \text{tp}(A, N) = \text{stp}(A, N)$. As N is \aleph_ϵ -primary over \emptyset we can find a sequence $\langle d_\alpha : \alpha < \alpha^* \rangle$ and $\langle w_\alpha : \alpha < \alpha^* \rangle$ such that $N = \{d_\alpha : \alpha < \alpha^*\}$ and $w_\alpha \subseteq \alpha$ is finite and $\text{stp}(d_\alpha, \{d_\beta : \beta \in w_\alpha\}) \vdash \text{stp}(d_\alpha, \{d_\beta : \beta < \alpha\})$ and $\beta < \alpha \Rightarrow d_\beta \neq d_\alpha$.

We can find a countable set $W \subseteq \alpha^*$ such that $A' \subseteq \{d_\alpha : \alpha \in W\}$ and $\alpha \in W \Rightarrow w_\alpha \subseteq W$. Let $A'' = \{a_\alpha : \alpha \in W\}$. By [Sh:c, IV,§2,§3] without loss of generality W is an initial segment of α^* .

Easily

$$\alpha < \alpha^* \ \& \ \alpha \notin W \Rightarrow \text{stp}(d_\alpha, \{d_\beta : \beta \in w_\alpha\}) \vdash \text{stp}(d_\alpha, A \cup \{d_\beta : \beta < \alpha\}).$$

As N^+ is \aleph_ϵ -primary over $N \cup A$ we can find a list $\{d_\alpha : \alpha \in [\alpha^*, \alpha^{**}]\}$ of $N^+ \setminus (N \cup A)$ such that $\text{tp}(d_\alpha, N \cup A \cup \{d_\beta : \beta \in [\alpha^*, \alpha^{**}]\})$ is \aleph_ϵ -isolated. So

$\langle d_\alpha : \alpha \notin W, \alpha < \alpha^{**} \rangle$ exemplifies that N^+ is \aleph_ϵ -primary over $A \cup A''$, hence by 1.18(0) we know that N^+ is \aleph_ϵ -primary over \emptyset .

2) We shall use the characterization of “ N is $\mathbf{F}_{\aleph_0}^a$ -prime over A ” in 1.17, more exactly we use the last condition in 1.17(1) for $A = \emptyset, M = N_\omega$. Clearly N_ω is \aleph_ϵ -saturated (as it is \aleph_ϵ -prime over $\bigcup_{n < \omega} N_n$). Suppose $B \subseteq N_\omega$ is finite and $p \in S(B)$ is (stationary and) regular.

Case 1: p not orthogonal to $\bigcup_{n < \omega} N_n$.

So for some $n < \omega, p$ is not orthogonal to N_n , hence there is a regular $p_1 \in S(N_n)$ such that p, p_1 are not orthogonal. Let $A_1 \subseteq N_n$ be finite such that p_1 does not fork over A and $p_1 \upharpoonright A_1$ is stationary. So by [Sh:c, V, §2] we know $\dim(p, N_\omega) = \dim(p_1 \upharpoonright A_1, N_\omega)$, hence it suffices to prove that the latter is \aleph_0 . Now this holds by [Sh:c, V, 1.16](3), p.237 or immitate the proof of $(*)_2$ above.

Case 2: p is orthogonal to $\bigcup_{n < \omega} N_n$.

Note that if each N_n is \aleph_ϵ -prime then $\bigcup_{n < \omega} N_n$ is \aleph_ϵ -saturated hence $N = \bigcup_{n < \omega} N_n$ hence this case does not arise. Let $A = \bigcup_{n < \omega} N_n$, so $\dim(p, N) \leq \aleph_0$ follows from

2A) below.

Alternatively (and work even if we replace N_η by a set $A_n, \mathbf{F}_{\aleph_0}^a$ -constructible over \emptyset , see below).

2A) By 2B).

2B) The first inequality as immediate (as T is superstable and \bar{a}, \bar{b} are finite), so let us concentrate on the second. Let $B \subseteq C$ be a finite set such that $\text{tp}_*(\bar{a} \hat{\ } \bar{b}, C)$ does not fork over B and $\text{stp}_*(\bar{a} \hat{\ } \bar{b}, B) \vdash \text{stp}_*(\bar{a} \hat{\ } \bar{b}, C)$. Recall $q \in S(N)$ extend \bar{a}/\bar{b} and do not fork over \bar{b} , let $b^* \in \mathfrak{C}$ realize q and let $q_1 = \text{stp}(\bar{b}^*, B \cup \bar{b})$ and $q_2 = \text{stp}(\bar{b}^*, C \cup \bar{b})$. Now by the assumption of our case q_1 is orthogonal to $\text{tp}_*(C, B)$ hence (see [Sh:c, V, §3]) $q_1 \vdash q_2$ and let $\{a_\alpha : \alpha < \alpha^*\} \subseteq (q_1 \upharpoonright (\bar{b} \cup B))(N)$ be a maximal set independent over $C + \bar{b}$, so $|\alpha^*| \leq \dim(\bar{a}/(C + \bar{b}), N)$ and $q \upharpoonright (C \cup \bar{b} \cup \{a_\alpha : \alpha < \alpha^*\}) \vdash q$. Also clearly $\text{stp}_*(\{a_\alpha : \alpha < \alpha^*\}, \bar{b} \cup B) \vdash \text{stp}_*(\{a_\alpha : \alpha < \alpha^*\}, \bar{b} \cup C)$. Together $\dim(q_1, N) \leq |\alpha^*|$ and as $|B| < \aleph_0 = \kappa_r(T)$ clearly $\dim(\bar{a}/\bar{b}, N) < \aleph_0 + \dim(q_1, N)^+$, so we are done.

We can use a different proof for part (2), note:

\otimes_1 if $\kappa = \text{cf}(\kappa) \geq \kappa_r(T)$ and B_α is \mathbf{F}_κ^a -constructible over A for $\alpha < \delta, \delta \leq \kappa$

and $\alpha < \beta < \delta \Rightarrow B_\alpha \subseteq B_\beta$ then $\bigcup_{\alpha < \delta} B_\alpha$ is \mathbf{F}_κ^a -constructible over A

[why? see [Sh:c, IV,§3], [Sh:c, IV,5.6,p.207] for such arguments, assume $\mathcal{A}_\alpha = \langle A, \langle a_i^\alpha : i < i_\alpha \rangle, \langle B_i^\alpha : i < i_\alpha \rangle \rangle$ is an \mathbf{F}_κ^a -construction of B_α over A . Without loss of generality $i < j < i_\alpha \Rightarrow a_i^\alpha \neq a_j^\alpha$, and choose by induction on $\zeta, \langle u_\zeta^\alpha : \alpha < \delta \rangle$ such that: $u_\zeta^\alpha \subseteq i_\alpha, u_\zeta^\alpha$ increasing continuous in $i, u_0^\alpha = \emptyset, |u_{\zeta+1}^\alpha \setminus u_\zeta^\alpha| \leq \kappa, u_\zeta^\alpha$ is \mathcal{A}_α -closed and $\alpha < \beta < \delta$ implies $\{a_j^\alpha : j \in u_\zeta^\alpha\} \subseteq \{a_j^\beta : j \in u_\zeta^\beta\}$ and $\text{tp}_*(\{a_i^\beta : i \in u_\zeta^\beta\}, A \cup \{a_i^\alpha : i < i_\alpha\})$ does not fork over $A \cup \{a_i^\alpha : i \in u_\zeta^\alpha\}$. Now find a list $\langle a_j : j < j^* \rangle$ such that for each $\zeta, \{j : a_j \in a_i^\alpha : i \in u_\zeta^\alpha \text{ for some } \alpha < \delta, \varepsilon < \zeta\}$ is an initial segment β_ζ of j^* and $\beta_{\zeta+1} \leq \beta_\zeta + \kappa$.]

We use \otimes_1 for $\kappa = \aleph_0$.

So each N_n is \aleph_ε -constructible over \emptyset hence $\bigcup_{n < \omega} N_n$ is \aleph_ε -constructible over \emptyset and

also N_ω is \aleph_ε -constructible over $\bigcup_{n < \omega} N_n$ hence N_ω is \aleph_ε -constructible over \emptyset . But

N_ω is \aleph_ε -saturated hence N_ω is \aleph_ε -primary over \emptyset . Alternatively use: if B is \mathbf{F}_κ^a -constructible over $A, \kappa \geq \kappa_r(T)$ and \mathbf{I} is indiscernible over $A, |\mathbf{I}| > \kappa$ then for some $\mathbf{J} \subseteq \mathbf{I}$ of cardinality $\leq \kappa, \mathbf{I} \setminus \mathbf{J}$ is an indiscernible set over B .

3) Suppose N'_3 is \aleph_ε -saturated and $N_1 + \bar{a} \subseteq N'_3$. As N_2 is \aleph_ε -prime over $N_0 + \bar{a}$ and $N_0 + \bar{a} \subseteq N_1 + \bar{a} \subseteq N'_3$ we can find an elementary embedding f_0 of N_2 into N'_3 extending $\text{id}_{N_0 + \bar{a}}$. By [Sh:c, V,3.3], the function $f_1 = f_0 \cup \text{id}_{N_1}$ is an elementary mapping and clearly $\text{Dom}(f_1) = N_1 \cup N_2$. As N_3 is \aleph_ε -prime over $N_1 \cup N_2$ and f_1 is an elementary mapping from $N_1 \cup N_2$ into N'_3 which is an \aleph_ε -saturated model there is an elementary embedding f_3 of N_3 into N'_3 extending f_2 . So as for any such N'_3 there is such f_3 , clearly N_3 is \aleph_ε -prime over $N_1 + \bar{a}$, as required.

4) Let N_0 be \aleph_0 -prime over \emptyset and let $\{p_i : i < \alpha\} \subseteq S(N_0)$ be a maximal family of pairwise orthogonal regular types. Let $\mathbf{I}_i = \{\bar{a}_n^i : n < \omega\} \subseteq \mathfrak{C}$ be a set of elements realizing p_i independent over N_0 and let $\mathbf{I} = \bigcup_{i < \alpha} \mathbf{I}_i$ and N'_1 be $\mathbf{F}_{\aleph_0}^a$ -prime over $N_0 \cup \mathbf{I}$.

Now

(*) if $\bar{a}, \bar{b} \subseteq N'_1$ and \bar{a}/\bar{b} is regular (hence stationary), then $\dim(\bar{a}/\bar{b}, N'_1) \leq \aleph_0$.

[Why? If $\bar{a}/\bar{b} \perp N_0$ then $\dim(\bar{a}/\bar{b}, N'_1) \leq \aleph_0$ by part (2A) and the choice of the p_i and \mathbf{I}_i for $i < \alpha$. If $\bar{a}/\bar{b} \not\perp N_0$, then for some $\bar{b}' \hat{=} \bar{a}' \subseteq N_0$ realizing $\text{stp}(\bar{b} \hat{=} \bar{a}, \emptyset)$, we have $\bar{a}'/\bar{b}' \perp \bar{a}/\bar{b}$ hence $\dim(\bar{a}/\bar{b}, N'_1) = \dim(\bar{a}'/\bar{b}', N'_1)$, so without loss of generality $\bar{b} \hat{=} \bar{a} \subseteq N_0$, similarly without loss of generality there is $i(*) < \alpha$ such that $\bar{a}/\bar{b} \subseteq p_{i(*)}$ and $p_{i(*)}$ does not fork over \bar{b} now easily $\dim(\bar{a}/\bar{b}, N'_1) = \dim(\bar{a}/\bar{b}, N_0) + \dim(p_{i(*)}, N_0) \leq \aleph_0 + \aleph_0 = \aleph_0$ (see [Sh:c, V,1.6](3)). So we have proved (*).]

Now use 1.17(1) to deduce: N'_1 is $\mathbf{F}_{\aleph_\epsilon}^a$ -prime over \emptyset hence (by uniqueness of \aleph_ϵ -prime model, 1.17(2)) $N'_1 \cong N_1$.

By renaming without a loss of generality $N'_1 = N_1$. Now

- (**)(α) $(N_1, c)_{c \in N_0}, (N_2, c)_{c \in N_0}$ are \aleph_ϵ -saturated and
 (β) if $\bar{a} \in \mathfrak{C}, \bar{b} \in N_\ell, \bar{a}/\bar{b}$ a regular type and $\bar{a} \bigcup_{\bar{b}} (N_0 + \bar{b})$ (for $\ell = 1$ or $\ell = 2$),
 then
 $\dim(\bar{a}/(\bar{b} \cup N_0), N_\ell) = \aleph_0$.

[Why? Remember that we work in $(\mathfrak{C}^{\text{eq}}, c)_{c \in N_0}$. The “ \aleph_ϵ -saturated” follows from the second statement.

Note: $\dim(\bar{a}/(\bar{b} \cup N_0), N_\ell) \leq \dim(\bar{a}/\bar{b}, N_\ell) \leq \aleph_0$ (first inequality by monotonicity, second inequality by 1.17(1) and the assumption “ N_ℓ is \aleph_ϵ -prime over \emptyset ”). If \bar{a}/\bar{b} is not orthogonal to N_0 then for some $i < \alpha$ we have $p_i \pm (\bar{a}/\bar{b})$ so easily (using “ N_ℓ is \aleph_ϵ -saturated”) we have $\dim(\bar{a}/(\bar{b} \cup N_0), N_\ell) = \dim(p_i, N_\ell) \geq \|\mathbf{I}_i\| = \aleph_0$; so together with the previous sentence we get equality. Lastly, if $\bar{a}/\bar{b} \perp N_0$ by part (2B) of 1.18, we have $\dim(\bar{a}/(\bar{b} \cup N_0), N_\ell) < \aleph_0 \Rightarrow \dim(\bar{a}/\bar{b}, N_\ell) < \aleph_0$ which contradicts the assumption “ N_ℓ is \aleph_ϵ -saturated”.] So we have proved (**) hence by 1.17(1) we get “ N_1, N_2 are isomorphic over N_0 ” as required.

5) This is proved similarly as if N is \aleph_ϵ -prime over A and $B \subseteq N$ is ϵ -finite then N is \aleph_ϵ -prime over $A + B$ and also over A' if $A + B \subseteq A' \subseteq \text{acl}(A + B)$, see part (10).

6) By [Sh:c, V,3.2].

7) First assume that $A_2^* \subseteq N_1$ and a/N_1 is regular. As N_1 is \aleph_ϵ -prime over $N_0 \cup N'_1$ and as T has NDOP (i.e. does not have DOP) we know (by [Sh:c, X,2.1,2.2,p.512]) that N_1 is \aleph_ϵ -minimal over $N_0 \cup N'_1$ and $\frac{a}{N_1}$ is not orthogonal to N_0 or to N'_1 . But $a/N_1 \perp N_0$ by an assumption, so a/N_1 is not orthogonal to N'_1 hence there is a regular $p' \in S(N'_1)$ not orthogonal to $\frac{a}{N_1}$ hence (by [Sh:c, V,1.12,p.236]) p' is realized say by $a' \in N_2$. By [Sh:c, V,3.3], we know that N_2 is \aleph_ϵ -prime over $N_1 + a'$. We can find N'_2 which is \aleph_ϵ -prime over $N'_1 + a'$ and N''_2 which is \aleph_ϵ -prime over $N_1 \cup N'_2$ hence by part (3) of 1.18 we know that N''_2 is \aleph_ϵ -prime over $N_1 + a'$ so by uniqueness, i.e. 1.17(1), without loss of generality $N''_2 = N_2$ hence we are done. In general by induction on α choose $N'_{2,\alpha}$ such that $N'_{2,0}$ is \aleph_ϵ -prime over $N'_1 \cup A_2^*$, $N'_{2,\alpha}$ is increasing with α and $N_1 \bigcup N'_{2,\alpha}$. Easily for some α , $N'_{2,\alpha}$ is defined

$$N'_1$$

but not $N'_{2,\alpha+1}$. Necessarily N_2 is \aleph_ϵ -prime over $N'_1 \cup N'_{2,\alpha}$. Lastly let $a' \in N'_{2,\alpha}$ be such that $\text{tp}(a, N_1 \cup N'_{2,\alpha})$ does not fork over $N_1 + a'$. Easily $N'_{2,\alpha}$ is \aleph_ϵ -prime over $N'_1 + a'$ by (1.17(1)).

8) Similar easier proof.

9) Let N'_0 be \aleph_ϵ -prime over A such that $B \bigcup_A N'_0$, and let N'_1 be \aleph_ϵ -prime over

$N'_0 \cup B$. By 1.18(1), we know that N'_1 is \aleph_ϵ -prime over \emptyset , and by 1.18(10) below N'_1 is \aleph_ϵ -prime over $A \cup B$, hence by 1.17(2) we know that N'_1, N_1 are isomorphic over $A \cup B$ hence without loss of generality $N'_1 = N_1$ and so $N_0 = N'_0$ is as required.

10) By [Sh:c, IV,3.12](3),p.180. □_{1.18}

1.20 Fact. Assume $\langle N_\eta^1, a_\eta : \eta \in I \rangle \leq_{\text{direct}}^* \langle N_\eta^2, a_\eta : \eta \in I \rangle$ (see Definition 1.16) and $A \subseteq B \subseteq N_{<0}^1$ and $\bigwedge_{\eta \in I} N_\eta^2 \prec M$.

(1) If $\nu = \eta \hat{\ } \langle \alpha \rangle \in I$, then $N_\nu^2 \cup_{N_\eta^1} N_\nu^1$ and even $N_\eta^2 \cup_{N_\eta^1} \left(\bigcup_{\substack{\rho \in I \\ \neg \eta \triangleleft \rho}} N_\rho^1 \right)$; and

$\eta \triangleleft \nu \in I$ implies $N_\nu^2 \cup_{N_\eta^1} \left(\bigcup_{\substack{\rho \in I \\ \neg \eta \triangleleft \rho}} N_\rho^1 \right)$.

- (2) $\langle N_\eta^2, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition inside M above $\binom{B}{A}$ iff
 $\langle N_\eta^1, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition inside M above $\binom{B}{A}$.
 (3) Similarly replacing “ \aleph_ϵ -decomposition inside M above $\binom{B}{A}$ ” by
 “ \aleph_ϵ -decomposition of M above $\binom{B}{A}$ ”.

Proof. 1) We prove the first statement by induction on $lg(\eta)$. If $\eta = \langle \rangle$ this is clause (b) by the Definition 1.16 and clause (d) of Definition 1.11(1) (and [Sh:c, V,3.2]). If $\eta \neq \langle \rangle$, then $\frac{a_\nu}{N_\eta} \perp N_{(\eta^-)}^1$ (by condition (e) of Definition 1.11(1)). By the induction hypothesis $N_{(\eta^-)}^2 \cup_{N_{(\eta^-)}^1} N_\eta^1$ and we know N_η^2 is \aleph_ϵ -primary over

$N_{(\eta^-)}^2 \cup N_\eta^1$; we know this implies that no $p \in S(N_\eta^1)$ orthogonal to $N_{\eta^-}^1$ is realized in N_η^2 hence $\frac{a_\nu}{N_\eta^1} \perp \frac{N_\eta^2}{N_\eta^1}$, so $\frac{a_\nu}{N_\eta^1} \vdash \frac{a_\nu}{N_\eta^2}$ hence $\frac{N_\nu^1}{N_\eta^1} \perp \frac{N_\eta^2}{N_\eta^1}$ hence $N_\nu^1 \cup_{N_\eta^1} N_\eta^2$ as required.

The other statements hold by the non-forking calculus (remember if $\eta = \nu \hat{\ } \langle \alpha \rangle \in I$ then use $\text{tp}(\cup\{N_\rho^1 : \eta \trianglelefteq \rho \in I\}, N_\eta^1)$ is orthogonal to N_ν^1 or see details in the proof of 1.21(1)(α)).

2) By Definition 1.16, for $\ell = 1, 2$ we have: $\langle N_\eta^\ell, a_\eta : \eta \in I \rangle$ is a decomposition inside \mathfrak{C} and by assumption $\bigwedge_{\eta \in I} N_\eta^1 \prec N_\eta^2 \prec M$. So for $\ell = 1, 2$ we have to prove

“ $\langle N_\eta^\ell, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition inside M for $\binom{B}{A}$ ” assuming this holds for $1 - \ell$. We have to check Definition 1.11(1).

Clauses 1.5(1)(a),(b) for ℓ holds because they hold for $1 - \ell$.

Clause 1.5(1)(c) holds as by the assumptions $A \subseteq B \subseteq N_{<0>}^1 \prec N_{<0>}^2, A \subseteq N_{<>}^1$ and $N_{<0>}^1 \bigcup_{N_{<>}^1} N_{<>}^2$.

Clauses 1.5(1)(d),(e),(f),(h) holds as $\langle N_\eta^\ell, a_\eta : \eta \in I \rangle$ is a decomposition inside \mathfrak{C} (for $\ell = 1$ given, for $\ell = 2$ easily checked).

Clause 1.5(1)(g) holds as $\bigwedge_\eta N_\eta^1 \prec N_\eta^2 \prec M$ is given and M is \aleph_ϵ -saturated.

3) First we do the “only if” direction; i.e. prove the maximality of $\langle N_\eta^1, a_\eta : \eta \in I \rangle$ as an \aleph_ϵ -decomposition inside M for $\binom{B}{A}$ (i.e. condition (i) from 1.11(2)), assuming it holds for $\langle N_\eta^2, a_\eta : \eta \in I \rangle$. If this fails, then for some $\eta \in I \setminus \{<>\}$ and $a \in M, \{a_{\eta^{\langle \alpha \rangle}} : \eta^{\langle \alpha \rangle} \in I\} \cup \{a\}$ is independent over N_η^1 and $a \notin \{a_{\eta^{\langle \alpha \rangle}} : \eta^{\langle \alpha \rangle} \in I\}$ and $\frac{a}{N_\eta^1} \perp N_{\eta^-}^1$. Hence, if $\eta^{\langle \alpha_\ell \rangle} \in I$ for $\ell < k$ then $\bar{a} = \langle a \rangle^{\langle a_{\eta^{\langle \alpha_\ell \rangle}} : \ell < k \rangle}$ realizes over N_η^1 a type orthogonal to $N_{\eta^-}^1$, but $N_{\eta^-}^1 \prec N_\eta^1, N_{\eta^-}^1 \prec N_{\eta^-}^2$ and $N_\eta^1 \bigcup_{N_{\eta^-}^1} N_\eta^2$ (see 1.20(1), hence (by [Sh:c, V,2.8])

$\text{tp}(\bar{a}, N_\eta^2) \perp N_{\eta^-}^2$ hence $\{a\} \cup \{a_{\eta^{\langle \ell \rangle}} : \ell < k\}$ is independent over N_η^2 but $k, \eta^{\langle \alpha_\ell \rangle} \in I$ for $\ell < k$ were arbitrary so $\{a\} \cup \{a_{\eta^{\langle \alpha \rangle}} : \eta^{\langle \alpha \rangle} \in I\}$ is independent over N_η^2 contradicting condition (i) from Definition 1.11(2) for $\langle N_\eta^2, a_\eta : \eta \in I \rangle$.

For the other direction use: if the conclusion fails, then for some $\eta \in I \setminus \{<>\}$ and

$a \in M \setminus N_\eta^2 \setminus \{a_{\eta^{\langle \alpha \rangle}} : \eta^{\langle \alpha \rangle} \in I\}$ the set $\{a_{\eta^{\langle \alpha \rangle}} : \eta^{\langle \alpha \rangle} \in I\} \cup \{a\}$ is independent of N_η^2 and $\text{tp}(a, N_\eta^2)$ is orthogonal to $N_{\eta^-}^2$; let $N' \prec M$ be \aleph_ϵ -prime over $N_\eta^2 + a$. But N_η^2 is \aleph_ϵ -prime over $N_\eta^1 \cup N_{\eta^-}^2$ (by the definition of \leq -direct) so by NDOP $\text{tp}(a, N_\eta^2) \pm N_\eta^1$ hence there is a regular $q \in S(N_\eta^1)$ such that $q \pm \text{tp}(a, N_\eta^2)$. Hence some $a' \in N'$ realizes q , clearly $\{a_{\eta^{\langle \alpha \rangle}} : \eta^{\langle \alpha \rangle} \in I\} \cup \{a'\}$ is independent over N_η^2 (and $a' \notin \{a_{\eta^{\langle \alpha \rangle}} : \eta^{\langle \alpha \rangle} \in I\}$) hence over (N_η^2, N_η^1) and easily we get contradiction. $\square_{1.20}$

1.21 Fact. Assume $\langle N_\eta^1, a_\eta^1 : \eta \in I \rangle$ is an \aleph_ϵ -decomposition inside M .

1) If $N_{<>}^1 \prec N_{<>}^2 \prec M, N_\eta^2$ is \aleph_ϵ -prime over \emptyset and

$N_{<>}^2 \bigcup_{N_{<>}^1} \{a_{<\alpha>}^1 : <\alpha> \in I\}$ then

$$(\alpha) \left[N_{<>}^2 \bigcup_{N_{<>}^1} \bigcup_{\eta \in I} N_\eta^1 \right] \text{ and}$$

(β) we can find N_η^2 ($\eta \in I \setminus \{\langle \rangle\}$) such that $N_\eta^2 \prec M$, and
 $\langle N_\eta^1, a_\eta^1 : \eta \in I \rangle \leq_{\text{direct}}^* \langle N_\eta^2, a_\eta^1 : \eta \in I \rangle$.

2) If $\text{Cb} \frac{a_{\langle \alpha \rangle}^1}{N_{\langle \alpha \rangle}^1} \subseteq N_{\langle \alpha \rangle}^0 \prec N_{\langle \alpha \rangle}^1$ or at least $N_{\langle \alpha \rangle}^0 \prec N_{\langle \alpha \rangle}^1$ and $\frac{a_{\langle \alpha \rangle}^1}{N_{\langle \alpha \rangle}^1} \pm N_{\langle \alpha \rangle}^0$ whenever $\langle \alpha \rangle \in I$ then we can find $N_\eta^0 \prec M$ and $a_\eta^0 \in N_\eta$ (for $\eta \in I \setminus \{\langle \rangle\}$) such that $\langle N_\eta^0, a_\eta^0 : \eta \in I \rangle \leq_{\text{direct}}^* \langle N_\eta^1, a_\eta^0 : \eta \in I \rangle$.

3) In part (2), if in addition we are given $\langle B_\eta^* : \eta \in I \rangle$ such that B_η^* is an ε -finite subset of N_η , $\text{tp}_*(B_\eta^*, N_\eta)$ does not fork over $B_{\eta^-}^*$ and $B_{\langle \rangle}^* \subseteq N_{\langle \rangle}^0$ then we can demand in the conclusion that $\eta \in I \Rightarrow B_\eta^* \subseteq N_\eta^0$.

Proof. 1) For proving (α) let $\{\eta_i : i < i^*\}$ list the set I such that $\eta_i \triangleleft \eta_j \Rightarrow i < j$, so $\eta_0 = \langle \rangle$ and without loss of generality for some α^* we have

$\eta_i \in \{\langle \alpha \rangle : \langle \alpha \rangle \in I\} \Leftrightarrow i \in [1, \alpha^*)$. Now we prove by induction on $\beta \in [1, i^*)$ that $N_{\langle \alpha \rangle}^2 \bigcup \bigcup_{i < \beta} \{N_{\eta_i}^1\}$. For $\beta = 1$ this is assumed. For β limit use the local $N_{\langle \alpha \rangle}^1$

character of non-forking.

If $\beta = \gamma + 1 \in [1, \alpha^*)$, then by repeated use of [Sh:c, V,3.2] (as $\{a_{\eta_0} : j \in [1, \beta)\}$ is independent over $(N_{\langle \alpha \rangle}^1, N_{\langle \alpha \rangle}^2)$ and $N_{\langle \alpha \rangle}^1$ is \aleph_ε -saturated and $N_{\eta_j}^1$ ($j \in [1, \gamma)$) is \aleph_ε -prime over $N_{\langle \alpha \rangle}^1 + a_{\eta_j}$) we know that $\text{tp}(a_{\eta_\gamma}, N_{\langle \alpha \rangle}^2 \cup \bigcup_{i < \gamma} N_{\eta_i}^1)$ does not fork over $N_{\langle \alpha \rangle}^1$. Again by [Sh:c, V,3.2], the type $\text{tp}_*(N_{\eta_\gamma}^1, N_{\langle \alpha \rangle}^2 \cup \bigcup_{i < \gamma} N_{\eta_i}^1)$ does not fork over

$N_{\langle \alpha \rangle}^1$ hence $\bigcup_{i < \beta} N_{\eta_i}^1 \bigcup N_{\langle \alpha \rangle}^2$ and use symmetry.

Lastly, if $\beta \in \gamma + 1 \in [\alpha^*, i^*)$, $\text{tp}(a_{\eta_\gamma}, N_{\eta_\gamma})$ is orthogonal to $N_{\langle \alpha \rangle}^1$ and even to $N_{(\eta_\gamma)^-}^1$ so again by non-forking and [Sh:c, V,3.2] we can do it, so clause (α) holds.

For clause (β), we choose $N_{\eta_i}^2$ for $i \in [1, i^*)$ by induction on $i < i^*$ such that $N_{\eta_i}^2 \prec M$ is \aleph_ε -prime over $N_{\eta_i}^2 \cup N_{\eta_i}^1$. By the non-forking calculus we can check Definition 1.7.

2) We let $\{\eta_i : i < i^*\}$ be as above, now we choose $N_{\eta_i}^0, a_{\eta_i}^0$ by induction on $i \in [1, i^*)$ such that:

$$(*) \quad N_{\eta_i}^0 \prec N_{\eta_i}^1 \text{ and } N_{\eta_i}^1 \bigcup_{N_{\eta_i}^0} N_{\eta_i}^0 \text{ and } N_{\eta_i}^1 \text{ is } \aleph_\varepsilon\text{-prime over } N_{\eta_i}^0 \cup N_{\eta_i}^1$$

$$(**) \quad a_{\eta_i}^0 \in N_{\eta_i}^0 \text{ and } N_{\eta_i}^0 \text{ is } \aleph_\varepsilon\text{-prime over } N_{\eta_i}^0 + a_{\eta_i}^0.$$

The induction step has already been done: if $\ell g(\eta_i) > 1$ by 1.18(7) and if $\ell g(\eta_i) = 1$

by 1.18(8).

3) Similar.

□_{1.21}

1.22 Fact. 1) If $\langle N_\eta^1, a_\eta : \eta \in I \rangle \leq_{\text{direct}}^* \langle N_\eta^2, a_\eta : \eta \in I \rangle$ and both are \aleph_ϵ -decompositions of M above $\binom{B}{A}$, then

$$\mathcal{P}(\langle N_\eta^1, a_\eta^1 : \eta \in I \rangle, M) = \mathcal{P}(\langle N_\eta^2, a_\eta^2 : \eta \in I \rangle, M).$$

Proof. By Definition 1.11(5) it suffices to prove, for each $\eta \in I \setminus \{<>\}$ that

$$(*) \text{ for regular } p \in S(M) \text{ we have} \\ p \perp N_{\eta^-}^1 \ \& \ p \pm N_\eta^1 \Leftrightarrow p \perp N_{\eta^-}^2 \ \& \ p \pm N_\eta^2.$$

Now consider any regular $p \in S(M)$: first assume $p \perp N_{\eta^-}^1 \ \& \ p \pm N_\eta^1$ where $\eta \in I \setminus \{<>\}$ so $p \pm N_\eta^2$ (as $N_\eta^1 \prec N_\eta^2$ and $p \pm N_\eta^1$) and we can find a regular $q \in S(N_\eta^1)$ such that $q \pm p$; so as $p \perp N_{\eta^-}^1$ also $q \perp N_{\eta^-}^1$, now $q \perp N_{\eta^-}^2$ (as $N_\eta^1 \cup N_{\eta^-}^2$ and $q \perp N_\eta^1$ see [Sh:c, V,2.8]) hence $p \perp N_{\eta^-}^2$.

Second, assume $p \perp N_{\eta^-}^2 \ \& \ p \pm N_\eta^2$ where $\eta \in I \setminus \{<>\}$; remember $N_{\eta^-}^1, N_\eta^1, N_\eta^2, N_\eta^3$ are \aleph_ϵ -saturated, $N_\eta^1 \cup N_{\eta^-}^2$ and N_η^2 is \aleph_ϵ -prime over $N_\eta^1 \cup N_{\eta^-}^2$ and T does not have

DOP. Hence N_η^2 is \aleph_ϵ -minimal over $N_\eta^1 \cup N_{\eta^-}^2$ and every regular $q \in S(N_\eta^2)$ is not orthogonal to N_η^1 or to $N_{\eta^-}^2$. Also as $p \pm N_\eta^2$ there is a regular $q \in S(N_\eta^2)$ not orthogonal to p , so as $p \perp N_{\eta^-}^2$ also $q \perp N_{\eta^-}^2$; hence by the previous sentence $q \pm N_\eta^1$ hence $p \pm N_\eta^1$. Lastly, as $p \perp N_{\eta^-}^2$ and $N_{\eta^-}^1 \prec N_{\eta^-}^2$ clearly $p \perp N_{\eta^-}^1$, as required.

□_{1.22}

At last we start proving 1.14.

Proof of 1.14. 1) Let $N^0 \prec \mathfrak{C}$ be \aleph_ϵ -primary over A , without loss of generality $N^0 \cup B$ (but not necessarily $N^0 \prec M$), and let N^1 be \aleph_ϵ -primary over $N^0 \cup B$.

Now by 1.18(0) the model N^0 is \aleph_ϵ -primary over \emptyset and by 1.18(1) the model N^1 is \aleph_ϵ -primary over \emptyset hence (by 1.18(10)) is \aleph_ϵ -primary over B , hence without loss of generality $N^1 \prec M$. Let $N_{<>} =: N^0, N_{<0>} = N^1, I = \{<>, <0>\}$ and $a_{<0>} = B$. More exactly a_η is such that $\text{dcl}(\{a_\eta\}) = \text{dcl}(B)$. Clearly $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition inside M above $\binom{B}{A}$. Now apply part (2) of 1.14 proved below.

2) By 1.13(4) we know $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition inside M , by 1.18(2) we then find $J \supseteq I$ and N_η, a_η for $\eta \in J \setminus I$ such that $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition of M . By 1.18(3), $\langle N_\eta, a_\eta : \eta \in J' \rangle$ is an \aleph_ϵ -decomposition of M above $\binom{B}{A}$ where $J' =: \{\eta \in J : \eta = \langle \rangle \text{ or } \langle 0 \rangle \trianglelefteq \eta \in J\}$.

3) Part (a) holds by 1.13(2),(3). As for part (b) by 1.13(2) there is $\langle N_\eta, a_\eta : \eta \in J \rangle$, an \aleph_ϵ -decomposition of M with $I \subseteq J$; easily $[\langle 0 \rangle \trianglelefteq \eta \in J \Rightarrow \eta \in I]$.

□_{1.14(1),(2),(3)}

1.23 Fact. If $\langle N_\eta^\ell, a_\eta^\ell : \eta \in I^\ell \rangle$ are \aleph_ϵ -decompositions of M above $\binom{B}{A}$, for $\ell = 1, 2$ and $N_{\langle \rangle}^1 = N_{\langle \rangle}^2$ then $\mathcal{P}(\langle N_\eta^1, a_\eta^1 : \eta \in I^1 \rangle, M) = \mathcal{P}(\langle N_\eta^2, a_\eta^2 : \eta \in I^2 \rangle, M)$.

Proof. By 1.14(3)(b) we can find $J^1 \supseteq I^1$ and N_η^1, a_η^1 for $\eta \in J^1 \setminus I^1$ such that $\langle N_\eta^1, a_\eta^1 : \eta \in J^1 \rangle$ is an \aleph_ϵ -decomposition of M and moreover we have $\eta \in J^1 \setminus I^1 \Leftrightarrow \eta \neq \langle \rangle$ & $\neg(\langle 0 \rangle \triangleleft \eta)$. Let $J^2 = I^2 \cup (J^1 \setminus I^1)$ and for $\eta \in J^2 \setminus I^2$ let $a_\eta^2 =: a_\eta^1$, $N_\eta^2 =: N_\eta^1$. Easily $\langle N_\eta^2, a_\eta^2 : \eta \in J^2 \rangle$ is an \aleph_ϵ -decomposition of M . By 1.13(6) we know that for every regular $p \in S(M)$ there is (for $\ell = 1, 2$) a unique $\eta(p, \ell) \in J^\ell$ such that $p \pm N_{\eta(p, \ell)}$ & $p \perp N_{\eta(p, \ell)^-}$ (note $\langle \rangle^-$ – meaningless). By the uniqueness of $\eta(p, \ell)$, if $\eta(p, 1) \in J^1 \setminus I^1$ then as it can serve as $\eta(p, 2)$ clearly it is $\eta(p, 2)$ so $\eta(p, 2) = \eta(p, 1) \in J^1 \setminus I^1 = J^2 \setminus I^2$; similarly $\eta(p, 2) \in J^2 \setminus I^2 \Rightarrow \eta(p, 1) \in J^1 \setminus I^1$ and $\eta(p, 1) = \langle \rangle \Leftrightarrow \eta(p, 2) = \langle \rangle$. So

$$(*) \quad \eta(p, 1) \in I^1 \setminus \{\langle \rangle\} \Leftrightarrow \eta(p, 2) \in I^2 \setminus \{\langle \rangle\}.$$

But

$$(**) \quad \eta(p, \ell) \in I^\ell \setminus \{\langle \rangle\} \Leftrightarrow p \in \mathcal{P}(\langle N_\eta^\ell, a_\eta^\ell : \eta \in I^\ell \rangle, M).$$

Together we finish.

□_{1.23}

We continue proving 1.14.

Proof of 1.14(4). Let $A^* \subseteq M^-$ be ε -finite, so we can find an ε -finite $B^* \subseteq \cup\{N_\eta : \eta \in I\}$ such that $\text{stp}(A^*, B^*) \vdash \text{stp}(A^*, \cup\{N_\eta : \eta \in I\})$. Hence, there is a finite non empty $I^* \subseteq I$ such that $\langle \rangle \in I^*$, I^* is closed under initial segments and $B^* \subseteq \cup\{N_\eta : \eta \in I^*\}$, so of course $\text{stp}_*(A^*, \cup\{N_\eta : \eta \in I^*\}) \vdash \text{stp}(A^*, \cup\{N_\eta : \eta \in I\})$.

We can also find $\langle B_\eta^* : \eta \in I^* \rangle$ such that B_η^* is an ε -finite subset of N_η , $B_\eta^* = \text{acl}(B_\eta^*)$ and $B^* \subseteq \cup\{B_\eta^* : \eta \in I^*\}$, $\eta \neq \langle \rangle \Rightarrow a_\eta \in B_\eta^*$ and if $\eta \triangleleft \nu \in I^*$ then $B_\eta^* \subseteq B_\nu^*$ and $\text{tp}_*(B_\nu^*, N_\nu)$ does not fork over B_η^* . Without loss of generality $B \subseteq B_{\langle 0 \rangle}^*$. For $\eta \in I \setminus I^*$ let $B_\eta^* = B_{\eta \upharpoonright \ell}^*$ where $\ell < \ell g(\eta)$ is maximal such that $\eta \upharpoonright \ell \in I^*$, such ℓ exists as $\ell g(\eta)$ is finite and $\langle \rangle \in I^*$.

Let $N_\eta^1 = N_\eta$ and $a_\eta^1 = a_\eta$ for $\eta \in I$ and without loss of generality $J \neq I$ hence $J \setminus I \neq \emptyset$.

Let $N_{<>}^2 \prec M$ be \aleph_ϵ -prime over $\bigcup_{\nu \in J \setminus I} N_\nu$; letting $J \setminus I = \{\eta_i : i < i^*\}$ be such that $[\eta_i \triangleleft \eta_j \Rightarrow i < j]$ we can find $N_{<>,i}^2$ (for $i \leq i^*$) increasing continuous, $N_{<>,0}^2 = N_{<>}$ and $N_{<>,i+1}^2$ is \aleph_ϵ -prime over $N_{<>,i}^2 \cup N_{\eta_i}$ hence over $N_{<>,i}^2 + a_{\eta_i}$. Lastly, without loss of generality $N_{<>,i^*}^2 = N_{<>}^2$.

By 1.18(1),(2) we know $N_{<>}^2$ is \aleph_ϵ -primary over \emptyset and (using repeatedly 1.18(6) + finite character of forking) we have $N_{<>}^2 \bigcup_{N_{<>}^1} a_{<0>}$. By 1.18(4)

(with $N_{<>}^1, N_{<>}^2, B_{<>}^* \supseteq \text{Cb}(a_{<0>}/N_{<>}^1)$ here standing for N_1, N_2, A there) we can find a model $N_{<>}^0$ such that $a_{<0>} \bigcup_{N_{<>}^0} N_{<>}^1$ and $\text{Cb}(a_{<0>}/N_{<>}^1) \subseteq B_{<>}^* \subseteq N_{<>}^0$

$N_{<>}^0, N_{<>}^0 \prec N_{<>}^1, N_{<>}^0$ is \aleph_ϵ -primary over \emptyset and $N_{<>}^1, N_{<>}^2$ are isomorphic over $N_{<>}^0$. By 1.21(1) we can for $\eta \in I$ choose $N_\eta^2 \prec M$ with $N_\eta^1 \prec N_\eta^2$ and $\langle N_\eta^1, a_\eta^1 : \eta \in I \rangle \leq_{\text{direct}}^* \langle N_\eta^2, a_\eta^0 : \eta \in I \rangle$. Similarly, by 1.21(2) (here $\text{Suc}_I(<>) = \{\langle 0 \rangle\}$) we can choose an \aleph_ϵ -decomposition $\langle N_\eta^0, a_\eta^0 : \eta \in I \rangle$ with $\langle N_\eta^0, a_\eta^0 : \eta \in I \rangle \leq_{\text{direct}}^* \langle N_\eta^1, a_\eta^1 : \eta \in I \rangle$. Moreover, we can demand $\eta \in I^* \Rightarrow B_\eta^* \subseteq N_\eta^0$ using 1.21(3). By 1.13(12) + 1.14(3) we know that $\langle N_\eta^1, a_\eta^0 : \eta \in I \rangle$ is an \aleph_ϵ -decomposition of M^- and easily $\langle N_\eta^2, a_\eta^0 : \eta \in I \rangle$ is an \aleph_ϵ -decomposition of M . Now choose by induction on $\eta \in I$ an isomorphism f_η from N_η^1 onto N_η^2 over N_η^0 such that $\nu \triangleleft \eta \Rightarrow f_\nu \subseteq f_\eta$ and $\eta \in I^* \Rightarrow f_\eta \upharpoonright B_\eta^* = \text{id}_{B_\eta^*}$. For $\eta = <>$ we have chosen N_η^0 such that N_η^1, N_η^2 are isomorphic over N_η^0 . For the induction step note that $f_{(\eta^-)} \cup \text{id}_{N_\eta^0}$ is an elementary mapping as $N_{(\eta^-)}^2 \bigcup_{N_{(\eta^-)}^0} N_\eta^0$ and $f_{(\eta^-)} \cup \text{id}_{N_\eta^0}$ can

be extended to an isomorphism f_η from N_η^1 onto N_η^2 as N_η^ℓ is \aleph_ϵ -primary (in fact even \aleph_ϵ -minimal) over $N_{(\eta^-)}^\ell \cup N_\eta^0$ for $\ell = 1, 2$ (which holds easily). If $\eta \in I^*$ there is no problem to add $f_\eta \upharpoonright B_\eta^* = \text{id}_{B_\eta^*}$. Now by 1.13(3) the model M^- is \aleph_ϵ -saturated and \aleph_ϵ -primary and \aleph_ϵ -minimal over $\bigcup_{\eta \in J} N_\eta = \bigcup_{\eta \in I} N_\eta^1$; similarly M is

\aleph_ϵ -primary over $\bigcup_{\eta \in I} N_\eta^2$. Now $\bigcup_{\eta} f_\eta$ is an elementary mapping from $\bigcup_{\eta \in I} N_\eta^1$ onto

$\bigcup_{\eta \in I} N_\eta^2$ hence can be extended to an isomorphism f from M^- into M . Moreover

as $\text{stp}_*(A^*, \cup\{B_\eta^* : \eta \in I^*\}) \vdash \text{stp}(A^*, \{N_\eta^1 : \eta \in I\})$, by [Sh:c, CH.XII,§4] we have $\text{tp}_*(A^*, \cup\{B_\eta^* : \eta \in I^*\}) \vdash \text{tp}(A^*, \cup\{N_\eta^1 : \eta \in I\})$ hence $\text{tp}_*(A^*, \cup\{B_\eta^* : \eta \in I^*\})$ has a unique extension as a complete type over $\cup\{N_\eta^1 : \eta \in I\}$ hence over $\cup\{N_\eta^2 : \eta \in I\}$

so without loss of generality $f \upharpoonright A^* = \text{id}_{A^*}$. By the \aleph_ϵ -minimality of M over $\bigcup_{\eta \in I} N_\eta$ (see 1.13(3)), f is onto M , so f is as required. $\square_{1.14(4)}$

We delay the proof of 1.14(5).

Proof of 1.14(6). Let $\langle N_\eta^\ell, a_\eta^\ell : \eta \in I^\ell \rangle$ for $\ell = 1, 2$, be \aleph_ϵ -decompositions of M above $(\frac{B}{A})$, so $\text{dcl}(a_{<>}^\ell) = \text{dcl}(B)$. Let $p \in S(M)$, and assume that $p \in \mathcal{P}(\langle N_\eta^1, a_\eta^1 : \eta \in I^1 \rangle, M)$, i.e. for some $\eta \in I^1 \setminus \{<>\}$, $(p_\eta \perp N_{\eta^-})$ and $p_\eta \pm N_\eta$. We shall prove that the situation is similar for $\ell = 2$; i.e. $p \in \mathcal{P}(\langle N_\eta^2, a_\eta^2 : \eta \in I^2 \rangle, M)$; by symmetry this suffices.

Let $n = \text{lg}(\eta)$, choose $\langle B_\ell : \ell \leq n \rangle$ and d such that:

- (α) $A \subseteq B_0$,
- (β) $B \subseteq B_1$,
- (γ) $a_{\eta \upharpoonright \ell} \subseteq B_\ell \subseteq N_{\eta \upharpoonright \ell}^1$, for $\ell \leq n$
- (δ) $B_{\ell+1} \bigcup_{B_\ell} N_{\eta \upharpoonright \ell}^1$,
- (ϵ) $\frac{B_{\ell+1}}{B_\ell + a_{\eta \upharpoonright (\ell+1)}^1} \vdash \frac{B_{\ell+1}}{N_{\eta \upharpoonright \ell}^1 + a_{\eta \upharpoonright (\ell+1)}^1}$,
- (ζ) $d \in B_n$, $\frac{d}{B_n \setminus \{d\}}$ is regular $\pm p$, (hence $\perp B_{n-1}$)
- (η) B_ℓ is ϵ -finite.

[Why such $\langle B_\ell : \ell \leq n \rangle$ exists? We prove by induction on n that for any $\eta \in I$ of length n and ϵ -finite $B' \subseteq N_\eta$, there is $\langle B_\ell : \ell \leq n \rangle$ satisfying (α) – (ϵ), (η) such that $B' \subseteq B_n$. Now there is $p' \in S(N_\eta^1)$ regular not orthogonal to p , let $B^1 \subseteq N_\eta^1$ be an ϵ -finite set extending $\text{Cb}(p')$. Applying the previous sentence to η, B^1 we get $\langle B_\ell : \ell \leq n \rangle$, let $d \in N_\eta$ realize $p' \upharpoonright B_n$.

Now as $n > 0$, $\text{tp}(d, B_n) \perp N_{\eta^-}$ hence $\text{tp}(d, B_n) \perp B_{n-1}$, hence $\text{tp}(d, B_n) \perp \text{tp}_*(N_{\eta^-}, B_n)$, hence as $\text{tp}(d, B_n)$ is stationary, by [Sh:c, V,1.2](2),p.231, the types $\text{tp}(d, B_n), \text{tp}_*(N_{\eta^-}, B_n)$ are weakly orthogonal so $\text{tp}(d, B_n) \vdash \text{tp}(d, N_{\eta^-} \cup B_n)$ hence $\frac{B_n + d}{B_{n-1} + a_\eta^1} \vdash \frac{B_n + d}{N_{\eta^-} + a_\eta^1}$.

Now replace B_n by $B_n \cup \{d\}$ and we finish].

Note that necessarily

- (δ)⁺ $B_n \bigcup_{B_m} N_{\eta \upharpoonright m}^1$ for $m \leq n$.
[Why? By the non-forking calculus].
- (ϵ)⁺ $\frac{B_n}{B_m + a_{\eta \upharpoonright (m+1)}^1} \perp_a B_m$ for $m < n$.
[Why? As $N_{\eta \upharpoonright m}^1$ is \aleph_ϵ -saturated].

Choose $D^* \subseteq N_{\langle \rangle}^2$ finite such that $\frac{B_n}{N_{\langle \rangle}^2 + B}$ does not fork over $D^* + B$.

[Note: we really mean $D^* \subseteq N_{\langle \rangle}^2$, not $D^* \subseteq N_{\langle \rangle}^1$].

We can find $N_{\langle \rangle}^3, \aleph_\epsilon$ -prime over \emptyset such that $A \subseteq N_{\langle \rangle}^3 \prec N_{\langle \rangle}^2$ and $D^* \amalg_A N_{\langle \rangle}^3$ and $N_{\langle \rangle}^2$ is \aleph_ϵ -prime over $N_{\langle \rangle}^3 \cup D^*$ (by 1.18(9)). Hence $B_n \amalg_A N_{\langle \rangle}^3$ and $B_n \amalg_B N_{\langle \rangle}^3$ (by the non-forking calculus). As $\text{tp}_*(B, N_{\langle \rangle}^2)$ does not fork over $A \subseteq N_{\langle \rangle}^3 \subseteq N_{\langle \rangle}^2$ by 1.21(2) we can find N_η^3, a_η^3 (for $\eta \in I^2 \setminus \{\langle \rangle\}$), such that $\langle N_\eta^3, a_\eta^3 : \eta \in I \rangle$ is an \aleph_ϵ -decomposition inside M above $\binom{B}{A}$ and $\langle N_\eta^3, a_\eta^3 : \eta \in I^2 \rangle \leq_{\text{direct}}^* \langle N_\eta^2, a_\eta^3 : \eta \in I^2 \rangle$ and $a_{\langle 0 \rangle}^3 = a_{\langle 0 \rangle}^2$ (remember $\text{dcl}(a_{\langle 0 \rangle}^2) = \text{dcl}(B)$). By 1.20(2) we know $\langle N_\eta^3, a_\eta^3 : \eta \in I^2 \rangle$ is an \aleph_ϵ -decomposition of M above $\binom{B}{A}$.

By 1.22 it is enough to show $p \in \mathcal{P}(\langle N_\eta^3, a_\eta^3 : \eta \in I^2 \rangle, M)$. Let $N_{\langle \rangle}^4 \prec N_{\langle \rangle}^2 \prec M$ be \aleph_ϵ -prime over $N_{\langle \rangle}^3 \cup B_0$. Now by the non-forking calculus $B \amalg_A (N_{\langle \rangle}^3 \cup B_0)$ [why? because

- (a) as said above $B_n \amalg_B N_{\langle \rangle}^3$ but $B_0 \subseteq B_n$ so $B_0 \amalg_B N_{\langle \rangle}^3$, and
- (b) as $B \amalg_A N_{\langle \rangle}^1$ and $B_0 \subseteq N_{\langle \rangle}^1$ we have $B \amalg_A B_0$ so $B_0 \amalg_A B$

hence (by (a) + (b) as $A \subseteq B$)

- (c) $\frac{B_0}{N_{\langle \rangle}^3 + B}$ does not fork over A ,

also

- (d) $B \amalg_A N_{\langle \rangle}^3$ (as $A \subseteq N_{\langle \rangle}^3 \subseteq N_{\langle \rangle}^2$ and $\text{tp}(B, N_{\langle \rangle}^2)$ does not fork over A)

putting (c) and (d) together we get

- (e) $\amalg_A \{B_0, B, N_{\langle \rangle}^3\}$

hence the conclusion].

Hence $B \amalg_{N_{\langle \rangle}^3} B_0$ so $B \amalg_{N_{\langle \rangle}^3} N_{\langle \rangle}^4$ (by 1.18(6)) and so (as $N_{\langle 0 \rangle}^3$ is \aleph_ϵ -prime over $N_{\langle \rangle}^3 + \text{dcl}(a_{\langle \rangle}^3) = N_{\langle \rangle}^3 + \text{dcl}(B)$) we have $N_{\langle \rangle}^4 \amalg_{N_{\langle \rangle}^3} N_{\langle 0 \rangle}^3$ and by 1.21(1) we can choose $N_\eta^4 \prec M$ (for $\eta \in I^2 \setminus \{\langle \rangle\}$), such that $\langle N_\eta^4, a_\eta^3 : \eta \in I^2 \rangle \geq_{\text{direct}} \langle N_\eta^3, a_\eta^3 : \eta \in I^2 \rangle$. So by 1.20(1) $\langle N_\eta^4, a_\eta^3 : \eta \in I^2 \rangle$ is an \aleph_ϵ -decomposition of M above $\binom{B}{A}$ hence $a_{\langle 0 \rangle}^3 / N_\eta^4$ does not fork over A but $A \subseteq B_0 \subseteq N_{\langle \rangle}^4$ so $a_{\langle \rangle}^3 / N_\eta^4$ does not fork over B_0

and by 1.22 it is enough to prove $p \in \mathcal{P}(\langle N_\eta^4, a_\eta^3 : \eta \in I^2 \rangle, M)$. Now as said above $B \amalg_{N_{\langle \rangle}^3} N_{\langle \rangle}^4$ and $B \amalg_A N_{\langle \rangle}^3$ so together $B \amalg_A N_{\langle \rangle}^4$, also we have $A \subseteq B_0 \subseteq N_{\langle \rangle}^4$,

hence $B \amalg_{B_0} N_{\langle \rangle}^4$ and

$\frac{B_n}{B_0+B} \equiv \frac{B_n}{B_0+a_{\langle 0 \rangle}^3} \perp_a B_0$ (by $(\epsilon)^+$ above) but $a_{\langle \rangle}^3 \amalg_{B_0} N_{\langle \rangle}^4$ hence $\frac{B_n}{N_{\langle \rangle}^4+a_{\langle 0 \rangle}^3}$ is

\aleph_ϵ -isolated. Also letting $B'_n = B_n \setminus \{d\}$ we have $\frac{B'_n}{N_{\langle \rangle}^4+a_{\langle 0 \rangle}^3}$ is \aleph_ϵ -isolated and $\frac{d}{B'_n} \perp B_0$ (by clause (ζ)), and clearly $d \amalg_{B'_n} (N_{\langle \rangle}^4 \cup B'_n)$ so $\frac{d}{B'_n} \perp N_{\langle \rangle}^4$. Hence we can find

$\langle N_\eta^5, a_\eta^5 : \eta \in I^5 \rangle$ an \aleph_ϵ -decomposition of M above $\binom{B}{A}$ such that $N_{\langle \rangle}^5 = N_{\langle \rangle}^4$, $\text{dcl}(B) = \text{dcl}(a_{\langle 0 \rangle}^5, B_n \setminus \{d\}) \subseteq N_{\langle 0 \rangle}^5$ and $d = a_{\langle 0,0 \rangle}^5$ (on d see clause (ζ) above) so $d \amalg_{B_n} N_{\langle 0 \rangle}^5$.

By 1.23 it is enough to show $p \in \mathcal{P}(\langle N_\eta^5, a_\eta^5 : \eta \in I^5 \rangle, M)$ which holds trivially as $\text{tp}(d, B_n \setminus \{d\})$ witness. $\square_{1.14(6)}$

Proof of 1.14(5). By 1.8, with A, B, A_1, B_1 here standing for A_1, B_1, A_2, B_2 there, we know that there are $\langle B'_\ell : \ell \leq n \rangle, \langle c_\ell : 1 \leq \ell < n \rangle$ as there. By 1.18(9) we can choose $N_{\langle \rangle}^1$ such that $B_0 \subseteq N_{\langle \rangle}^1, N_{\langle 1 \rangle}^1 \amalg_{B_0} B_n, N_{\langle \rangle}^1$ is \aleph_ϵ -primary over \emptyset . Then

we choose $\langle N_\eta^1, a_\eta^1 : \eta \in \left\{ \langle \rangle, \langle 0 \rangle, \langle 0, 0 \rangle, \dots, \underbrace{\langle 0, \dots, 0 \rangle}_n \right\} \rangle$, (where

$\underbrace{\langle 0, \dots, 0 \rangle}_n \prec M$ and $\ell > 0 \Rightarrow \underbrace{\langle 0, \dots, 0 \rangle}_\ell = c_\ell, B'_{\ell g(\eta)} \subseteq N_\eta^1$ and we choose

N_η^1 by induction on $\ell g(\eta)$ being \aleph_ϵ -prime over $N_{\eta^-}^1 \cup a_\eta^1$ hence $a_\eta^1/N_{\eta^-}^1$ does not fork over $B'_{\ell g(\eta^-)}$ hence N_η^1 is \aleph_ϵ -prime also over $N_{\eta^-}^1 + B'_{\ell g(\eta)}$. So $\langle N_\eta^1, a_\eta^1 : \eta \in \{\langle \rangle, \dots\} \rangle$ is an \aleph_ϵ -decomposition inside M for $\binom{B_1}{A_1}$. Now apply first 1.14(2) and then 1.14(6).

Proof of 1.14(7). Should be easy. Note that

(*)₁ for no $\binom{B'}{A'}$ do we have $\binom{B}{A} \leq_b \binom{B'}{A'}$
 Why? By the definition of Depth zero.

(*)₂ if $\binom{B}{A} <_a \binom{B'}{A'}$, then also $\binom{B'}{A'}$ satisfies the assumption.

Hence

(**) for no $\binom{B_1}{A_1}, \binom{B_2}{A_2}$ do we have

$$\binom{B}{A} \leq_a \binom{B_1}{A_1} <_b \binom{B_2}{A_2}.$$

[Why? As also $\binom{B_1}{A_1}$ satisfies the assumption].

Now we can prove the statement by induction on α for all pairs $\binom{B}{A}$ satisfying the assumption. For $\alpha = 0$ the statement is a tautology. For α limit ordinal reread clause (c) of Definition 1.9(1). For $\alpha = \beta + 1$, reread clause (b) of Definition 1.9(1): on $\text{tp}_\beta(\binom{B}{A}, M)$ use the induction hypothesis also for computing $Y_{A,B,M}^{1,B}$ (and reread the definition of tp_0 , in Definition 1.9(1), clause (a)). Lastly $Y_{A,B,M}^{2,\beta}$ is empty by (*) above.

Proof of 1.14(8),(9). Read Definition 1.9.

□_{1.14(5),(7),(8),(9)}

Discussion. In particular, the following Claim 1.26 implies that if $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ε -decomposition of M above $\binom{B}{A}$ and M^- is \aleph_ε -prime over $\cup\{N_\eta : \eta \in I\}$ then $\binom{B}{A}$ has the same tp_α in M and M^- .

1.24 Claim. 1) Assume that $M_1 \prec M_2$ are \aleph_ε -saturated, $\binom{B}{A} \in \Gamma(M_1)$. Then the following are equivalent:

- (a) if $p \in \mathcal{P}(\binom{B}{A}, M_1)$
(see 1.14(6) for definition; so $p \in S(M_1)$ is regular) then p is not realized in M_2
- (b) there is an \aleph_ε -decomposition of M_1 above $\binom{B}{A}$, which is also an \aleph_ε -decomposition of M_2 above $\binom{B}{A}$
- (c) every \aleph_ε -decomposition of M_1 above $\binom{B}{A}$ is also an \aleph_ε -decomposition of M_2 above $\binom{B}{A}$.

2) If M is \aleph_ε -saturated, $\binom{B_1}{A_1} \leq^* \binom{B_2}{A_2}$ are both in $\Gamma(M)$ then $\mathcal{P}(\binom{B_2}{A_2}, M) \subseteq \mathcal{P}(\binom{B_1}{A_1}, M)$.

3) The conditions in 1.24(1) above implies

- (d) $p \in \mathcal{P}(\binom{B}{A}, M_2) \Rightarrow p \pm M_1$.

Proof. 1) $(c) \Rightarrow (b)$.

By 1.14(1) there is an \aleph_ε -decomposition of M_1 above $\binom{B}{A}$. By clause (c) it is also an \aleph_ε -decomposition of M_2 above $\binom{B}{A}$, just as needed for clause (b).

$(b) \Rightarrow (a)$

Let $\langle N_\eta, a_\eta : \eta \in I \rangle$ be as said in clause (b). By 1.14(3)(b) we can find $J_1, I \subseteq J_1$ and N_η, a_η (for $\eta \in J_1 \setminus I$) such that $\langle N_\eta, a_\eta : \eta \in J_1 \rangle$ is an \aleph_ε -decomposition of M_1 and $\nu \in J_1 \setminus I \Rightarrow \nu(0) > 0$. Then we can find $J_2, J_1 \subseteq J_2$ and N_η, a_η (for $\eta \in J_2 \setminus J_1$) such that $\langle N_\eta : \eta \in J_2 \rangle$ is an \aleph_ε -decomposition of M_2 (by 1.14(2)). By 1.14(3)(b), $\nu \in J_2 \setminus I \Rightarrow \nu(0) > 0$. So $\eta \in I \setminus \{\langle \rangle\} \Rightarrow \text{Suc}_{J_2}(\eta) = \text{Suc}_I(\eta)$, hence

(*) if $\eta \in I \setminus \{\langle \rangle\}$ and $q \in S(N_\eta)$ is regular orthogonal to N_{η^-} then the stationarization of q in $S(M_1)$ is not realized in M_2 .

Now if $p \in \mathcal{P}(\binom{B}{A}, M_1)$ then $p \in S(M_1)$ is regular and (see 1.14(1), 1.11(5)) for some $\eta \in I \setminus \{\langle \rangle\}$, $p \perp N_{\eta^-}$, $p \pm N_\eta$, so there is a regular $q \in S(N_\eta)$ not orthogonal to p . Now no $c \in M_2$ realizes the stationarization of q over M_1 (by (*) above), hence this applies to p , too.

$(a) \Rightarrow (c)$

Let $\langle N_\eta, a_\eta : \eta \in I \rangle$ be an \aleph_ε -decomposition of M_1 above $\binom{B}{A}$. We can find $\langle N_\eta, a_\eta : \eta \in J \rangle$ an \aleph_ε -decomposition of M_1 such that $I \subseteq J$ and $\nu \in J \setminus I \Rightarrow \nu(0) > 0$ (by 1.14(3)(b)), so M is \aleph_ε -prime over $\cup\{N_\eta : \eta \in J\}$. We should check that $\langle N_\eta : a_\eta : \eta \in I \rangle$ it is also an \aleph_ε -decomposition of M_2 above $\binom{B}{A}$, i.e. Definition 1.11(1),(2). Now in 1.11(1), clauses (a)-(h) are immediate, so let us check clause (i) (in 1.11(2)). Let $\eta \in I \setminus \{\langle \rangle\}$, now is $\{a_{\eta \hat{\ } \langle \alpha \rangle} : \eta \hat{\ } \langle \alpha \rangle \in I\}$ really maximal (among independent over N_η sets of elements of M_2 realizing a type from $\mathcal{P}_\eta = \{p \in S(N_\eta) : p \text{ orthogonal to } N_{\eta^-}\}$)?. This should be clear from clause (a) (and basic properties of dependencies and regular types).

2) By 1.14(5).

3) Left to the reader. □_{1.24}

1.25 Conclusion: Assume $M_1 \prec M_2$ are \aleph_ε -saturated and $\binom{B_1}{A_1} \leq^* \binom{B_2}{A_2}$ both in $\Gamma(M_1)$. If clause (a) (equivalently (b) or (c)) of 1.24 holds for $\binom{B_1}{A_1}, M_1, M_2$ then they hold for $\binom{B_2}{A_2}, M_1, M_2$.

Proof. By 1.24(2), clause (a) for $\binom{B_1}{A_1}, M_1, M_2$ implies clause (a) for $\binom{B_2}{A_2}, M_1, M_2$. □_{1.25}

1.26 Claim. *If $\left(\begin{smallmatrix} B_1 \\ A_1 \end{smallmatrix}\right) \in \Gamma(M)$ and $\langle N_\eta, a_\eta : \eta \in I \rangle$ is an \aleph_ϵ -decomposition of M above $\left(\begin{smallmatrix} B_1 \\ A_1 \end{smallmatrix}\right)$ and $M^- \subseteq M$ is \aleph_ϵ -saturated and $\bigcup_{\eta \in I} N_\eta \subseteq M^-$ and α is an ordinal then*

$$\text{tp}_\alpha \left[\left(\begin{smallmatrix} B_1 \\ A_1 \end{smallmatrix} \right), M \right] = \text{tp}_\alpha \left[\left(\begin{smallmatrix} B_1 \\ A_1 \end{smallmatrix} \right), M^- \right]$$

Proof. We prove this by induction on α (for all $B, A, \langle N_\eta, a_\eta : \eta \in I \rangle, I, M$ and M^- as above). We can find an \aleph_ϵ -decomposition $\langle N_\eta, a_\eta : \eta \in J \rangle$ of M with $I \subseteq J$ (by 1.13(4) + 1.13(2)) such that $\eta \in J \setminus I \Leftrightarrow \eta \neq \langle \rangle$ and $\neg \langle 0 \rangle \leq \eta$ and so M is \aleph_ϵ -prime over $\bigcup_{\eta \in J} N_\eta$ and also over $M^- \cup \{N_\eta : \eta \in J \setminus I\}$.

Case 0: $\alpha = 0$.

Trivial.

Case 1: α is a limit ordinal.

Trivial by induction hypothesis (and the definition of tp_α).

Case 2: $\alpha = \beta + 1$.

We can find $M^* \prec M^-$ which is \aleph_ϵ -prime over $\bigcup_{\eta \in I} N_\eta$, so as equality is transitive

it is enough to prove

$$\text{tp}_\alpha \left(\left(\begin{smallmatrix} B_1 \\ A_1 \end{smallmatrix} \right), M^* \right) = \text{tp}_\alpha \left(\left(\begin{smallmatrix} B_1 \\ A_1 \end{smallmatrix} \right), M^- \right)$$

and

$$\text{tp}_\alpha \left(\left(\begin{smallmatrix} B_1 \\ A_1 \end{smallmatrix} \right), M^* \right) = \text{tp}_\alpha \left(\left(\begin{smallmatrix} B_1 \\ A_1 \end{smallmatrix} \right), M \right).$$

By symmetry, this means that it is enough to prove the statement when M^- is \aleph_ϵ -prime over $\bigcup_{\eta \in I} N_\eta$.

Looking at the definition of $\text{tp}_{\beta+1}$ and remembering the induction hypothesis our problems are as follows:

First component of tp_α :

given $(\begin{smallmatrix} B_1 \\ A_1 \end{smallmatrix}) \leq_a (\begin{smallmatrix} B_2 \\ A_2 \end{smallmatrix})$, $B_2 \subseteq M$, it suffices to find $(\begin{smallmatrix} B_3 \\ A_3 \end{smallmatrix})$ such that:

- (*) there is $f \in \text{AUT}(\mathfrak{C})$ such that: $f \upharpoonright B_1 = \text{id}_{B_1}$, $f(A_2) = A_3$,
 $f(B_2) = B_3$ and $B_3 \subseteq M^-$ and $\text{tp}_\beta[(\begin{smallmatrix} B_2 \\ A_2 \end{smallmatrix}), M] = \text{tp}_\beta[(\begin{smallmatrix} B_3 \\ A_3 \end{smallmatrix}), M^-]$
 (pedantically we should replace B_ℓ, A_ℓ by indexed sets).

We can find J', M' such that:

- (i) $I \subseteq J' \subseteq J$, $|J' \setminus I| < \aleph_0$, J' closed under initial segments,
- (ii) $M' \prec M$ is \aleph_ϵ -prime over $M^- \cup \cup \{N_\eta : \eta \in J' \setminus I\}$
- (iii) $B_2 \subseteq M'$.

The induction hypothesis for β applies, and gives

$$\text{tp}_\beta[(\begin{smallmatrix} B_2 \\ A_2 \end{smallmatrix}), M] = \text{tp}_\beta[(\begin{smallmatrix} B_2 \\ A_2 \end{smallmatrix}), M'].$$

By 1.14(4) there is g , an isomorphism from M' onto M^- such that $g \upharpoonright B_1 = \text{id}$. So clearly $g(B_2) \subseteq M^-$ hence

$$\text{tp}_\beta[(\begin{smallmatrix} B_2 \\ A_2 \end{smallmatrix}), M'] = \text{tp}_\beta[(\begin{smallmatrix} g(B_2) \\ g(A_2) \end{smallmatrix}), M^-].$$

So $(\begin{smallmatrix} B_3 \\ A_3 \end{smallmatrix}) =: g(\begin{smallmatrix} A_2 \\ B_2 \end{smallmatrix})$ is as required.

Second component for tp_α :

So we are given Υ , a tp_β type, (and we assign the lower part as B) and we have to prove that the dimension in M and in M^- are the same, i.e.

$\dim(\mathbf{I}, M) = \dim(\mathbf{I}^-, M^-)$, where: $\mathbf{I} = \{c \in M : \Upsilon = \text{tp}_\beta((\begin{smallmatrix} c \\ B_1 \end{smallmatrix}), M)\}$ and
 $\mathbf{I}^- = \{c \in M^- : \Upsilon = \text{tp}_\beta((\begin{smallmatrix} c \\ B_1 \end{smallmatrix}), M^-)\}$.

Let p be such that: $\text{tp}_\beta((\begin{smallmatrix} c \\ B_1 \end{smallmatrix}), M) = \Upsilon \Rightarrow p = \frac{c}{B_1}$. Necessarily $p \perp A_1$ and p is regular (and stationary).

Clearly $\mathbf{I}^- \subseteq \mathbf{I}$, so without loss of generality $\mathbf{I} \neq \emptyset$ hence p is really well defined, now

- (*) for every $c \in \mathbf{I}$ for some $k < \omega$, $c'_\ell \in M^-$ realizing p for $\ell < k$ we have c depends on $\{c'_0, c'_1, \dots, c'_{k-1}\}$ over B_1 .

[Why? Clearly $p \perp N_{<>}$ (as $B_1 \cup \cup_{A_1} N_{<>}$ and $p \perp A_1$) hence

$$\text{tp}_*(\bigcup_{\eta \in J \setminus I} N_\eta, N_{<>}) \perp p \text{ hence}$$

$\text{tp}_*(\bigcup_{\eta \in J \setminus I} N_\eta, M^-) \perp p$, but M is \aleph_ε -prime over $M^- \cup \bigcup_{\eta \in J \setminus I} N_\eta$ hence by [Sh:c, V,3.2,p.250] for no $c \in M \setminus M^-$ is $\text{tp}(c, M^-)$ a stationarization of p hence by [Sh:c, V,1.16](3) clearly (*) follows].

If the type p has depth zero, then (by 1.14(7)):

$$\mathbf{I} = \{c \in M : \text{tp}(c, B) = p\} \text{ and}$$

$$\mathbf{I}^- = \{c \in M^- : \text{tp}(c, B) = p\}.$$

Now we have to prove $\dim(\mathbf{I}, A) = \dim(\mathbf{I}^-, A)$, as A is ε -finite and M, M^- are \aleph_ε -saturated and $\mathbf{I}^- \subseteq \mathbf{I}$ clearly $\aleph_0 \leq \dim(\mathbf{I}^-, A) \leq \dim(\mathbf{I}, A)$. Now the equality follows by (*) above.

So we can assume “ p has depth $>$ zero”, hence (by [Sh:c, X,7.2]) that the type p is trivial; hence, see [Sh:c, X7.3], in (*) without loss of generality $k = 1$ and dependency is an equivalence relation, so for “same dimension” it suffices to prove that every equivalence class (in M i.e. in \mathbf{I}) is representable in M^- i.e. in \mathbf{I}^- . By the remark on (*) in the previous sentence $(\forall d_1 \in \mathbf{I})(\exists d_2 \in \mathbf{I}^-)[\neg d_1 \bigcup_{B_1} d_2]$. So it

is enough to prove that:

⊗ if $d_1, d_2 \in M$ realize same type over B_1 , which is (stationary and) regular, and are dependent over B_1 and $d_1 \in M^-$ then there is $d'_2 \in M^-$ such that $\frac{d'_2}{B_1+d_1} = \frac{d_2}{B_1+d_1}$ and $\text{tp}_\beta\left[\left(\begin{smallmatrix} B_1+d_2 \\ B_1 \end{smallmatrix}\right), M\right] = \text{tp}_\beta\left[\left(\begin{smallmatrix} B_1+d'_2 \\ B_1 \end{smallmatrix}\right), M^-\right]$.

Let $M_0 = N_{\langle \rangle}$. There are J', M_1, M_1^+ such that

(*)₁(i) $J' \subseteq J$ is finite (and of course closed under initial segments)

(ii) $\langle \rangle \in J', \langle 0 \rangle \notin J'$

(iii) $M_1 \prec M$ is \aleph_ε -prime over $\cup\{N_\eta : \eta \in J'\}$

(iv) $M_1^+ \prec M$ is \aleph_ε -prime over $M_1 \cup M^-$ (and $M_1 \bigcup_{M_0} M^-$)

(v) $d_2 \in M_1^+$.

Now the triple $\left(\begin{smallmatrix} B_1+d_2 \\ B_1 \end{smallmatrix}\right), M_1, M$ satisfies the demand on $\left(\begin{smallmatrix} B_1 \\ A_1 \end{smallmatrix}\right), M^-, M$ (because $\left(\begin{smallmatrix} B_1 \\ A_1 \end{smallmatrix}\right) \leq^* \left(\begin{smallmatrix} B_1+d_2 \\ B_1 \end{smallmatrix}\right)$, by 1.25. Hence by the induction hypothesis we know that

$$\text{tp}\left[\left(\begin{smallmatrix} B_1+d_2 \\ B_1 \end{smallmatrix}\right), M\right] = \text{tp}_\beta\left[\left(\begin{smallmatrix} B_1+d_2 \\ B_1 \end{smallmatrix}\right), M_1^+\right].$$

By 1.29(4) there is an isomorphism f from M_1^+ onto M^- which is the identity on $B_1 + d_1$; let $d'_2 = f(d_2)$ so:

$$\text{tp}_\beta \left[\begin{pmatrix} B_1 + d_2 \\ B_1 \end{pmatrix}, M_1^+ \right] = \text{tp}_\beta \left[\begin{pmatrix} B_1 + d'_2 \\ B_1 \end{pmatrix}, M^- \right].$$

Together

$$\text{tp}_\beta \left[\begin{pmatrix} B_1 + d_2 \\ B_1 \end{pmatrix}, M \right] = \text{tp}_\beta \left[\begin{pmatrix} B_1 + d'_2 \\ B_1 \end{pmatrix}, M^- \right].$$

As $\{d_1, d_2\}$ is not independent over B_1 , also $\{f(d_1), f(d_2)\} = \{d_1, f(d_2)\}$ is not independent over B_1 , hence, as p is regular

(*) $\{d_2, f(d_2)\}$ is not independent over B_2 .

Together we have proved \bigoplus , hence finished proving the equality of the second component.

Third component: Trivial.

So we have finished the induction step, hence the proof. □_{1.26}

1.27 Claim. 1) Suppose M is \aleph_ϵ -saturated, $A \subseteq B \subseteq M$, $\binom{B}{A} \in \Gamma$, $\bigwedge_{\ell=1}^2 [A \subseteq A_\ell \subseteq M]$,

$A = \text{acl}(A)$, A_ℓ are ϵ -finite, $\frac{A_1}{A} = \frac{A_2}{A}$, $B \bigcup_A A_1$ and $B \bigcup_A A_2$.

Then $\text{tp}_\alpha \left[\binom{A_1 \cup B}{A_1}, M \right] = \text{tp}_\alpha \left[\binom{A_2 \cup B}{A_2}, M \right]$ for any ordinal α .

2) Suppose M is \aleph_ϵ -saturated, $B \subseteq M$, $\binom{B}{A} \in \Gamma$, $\bigwedge_{\ell=1}^2 [A \subseteq A_\ell \subseteq M]$, $A = \text{acl}(A)$,

$B = \text{acl}(B)$, $A_\ell = \text{acl}(A_\ell)$, A_ℓ is ϵ -finite, $\frac{A_1}{A} = \frac{A_2}{A}$, $B \bigcup_A A_1$, $B \bigcup_A A_2$, $f : A_1 \xrightarrow{\text{onto}} A_2$

an elementary mapping, $f \upharpoonright A = \text{id}_A$, $g \supseteq f \cup \text{id}_B$, g elementary mapping from $B_1 = \text{acl}(B \cup A_1)$ onto $B_2 = \text{acl}(B \cup A_2)$.

Then $g \left(\text{tp}_\alpha \left[\binom{B_1}{A_1}, M \right] \right) = \text{tp}_\alpha \left[\binom{B_2}{A_2}, M \right]$ for any ordinal α .

3) Assume that

(a) $A_\ell = \text{acl}(A_\ell) \subseteq B_\ell = \text{acl}(B_\ell) \subseteq M^\ell$ for $\ell = 1, 2$

(b) $A_\ell \subseteq A_\ell^+ \subseteq \text{acl}(A_\ell^+) \subseteq M^\ell$ for $\ell = 1, 2$

- (c) $B_\ell \bigcup_{A_\ell} A_\ell^+$ for $\ell = 1, 2$
- (d) f is an elementary mapping from A_1 onto A_2
- (e) g is an elementary mapping from A_1^+ onto A_2^+
- (f) $f \upharpoonright A_1 = g \upharpoonright A_1$
- (g) h is an elementary mapping from $B_1^+ = \text{acl}(B_1 \cup A_1^+)$ onto $B_2^+ = \text{acl}(B_2 \cup A_2^+)$ extending f and g
- (h) $f(\text{tp}_\alpha[(\frac{B_1}{A_1}), M_1]) = \text{tp}_\alpha[(\frac{B_2}{A_2}), M_2]$.

Then $h(\text{tp}_\alpha[(\frac{B_1^+}{A_1^+}), M_1]) = \text{tp}_\alpha[(\frac{B_2^+}{A_2^+}), M_2]$.

Proof. 1) Follows from part (2).

2) We can find $A_3 \subseteq M$ such that:

- (i) $\frac{A_3}{A} = \frac{A_1}{A}$
- (ii) $A_3 \bigcup_A (B \cup A_1 \cup A_2)$.

Hence without loss of generality $A_1 \bigcup_B A_2$ and even $\bigcup_A \{B, A_1, A_2\}$. Now we can find $N_{\langle \rangle}$, an \aleph_ϵ -prime model over \emptyset , $N_{\langle \rangle} \prec M$, $A \subseteq N_{\langle \rangle}$ and $(B \cup A_1 \cup A_2) \bigcup_A N_{\langle \rangle}$ (e.g. choose $\{A_1^\alpha \cup A_i^\alpha \cup B^\alpha : \alpha \leq \omega\} \subseteq M$ indiscernible over A , $A_1^\omega = A_1$, $A_2^\omega = A_2$, $B^\omega = B$ and let $N_{\langle \rangle} \prec M$ be \aleph_ϵ -primary over $\bigcup_{n < \omega} (A_1^n \cup A_2^n \cup B^n \cup A)$).

Now find $\langle N_\eta, a_\eta : \eta \in J \rangle$ an \aleph_ϵ -decomposition of M with $\text{dcl}(a_{\langle 0 \rangle}) = \text{dcl}(B)$, $\text{dcl}(a_{\langle 1 \rangle}) = \text{dcl}(A_1)$, $\text{dcl}(a_{\langle 2 \rangle}) = \text{dcl}(A_2)$.

Let $I = \{\eta \in J : \eta = \langle \rangle \text{ or } \langle 0 \rangle \trianglelefteq \eta\}$ and $J' = I \cup \{\langle 1 \rangle, \langle 2 \rangle\}$. Let $N_{\langle \rangle}^2 \prec M^*$ be \aleph_ϵ -prime over $N_{\langle 1 \rangle} \cup N_{\langle 2 \rangle}$. By 1.21 there is $\langle N_\eta^2, a_\eta : \eta \in I \rangle$ an \aleph_ϵ -decomposition of M above $(\frac{B}{A})$ such that $\langle N_\eta, a_\eta : \eta \in I \rangle \leq_{\text{direct}} \langle N_\eta^2, a_\eta : \eta \in I \rangle$. Let $M' \prec M$ be \aleph_ϵ -prime over $\bigcup_{\eta \in I} N_\eta^2$ and $M^- \prec M'$ be \aleph_ϵ -prime over $\bigcup_{\eta \in I} N_\eta$. So

$M^- \prec M' \prec M$ and M' is \aleph_ϵ -prime over $M^- \cup N_{\langle 1 \rangle} \cup N_{\langle 2 \rangle}$.

Now by 1.26 we have $\text{tp}_\alpha[(\frac{B}{A_\ell}), M] = \text{tp}_\alpha[(\frac{B}{A_\ell}), M']$ for $\ell = 1, 2$ hence it suffices to find an automorphism of M' extending g . Let $B^+ = \text{acl}(N_{\langle \rangle} \cup B)$, $A_\ell^* = \text{acl}(B \cup A_\ell)$, let $\bar{\mathbf{a}}_\ell$ list A_ℓ^* be such that $\bar{\mathbf{a}}_2 = g(\bar{\mathbf{a}}_1)$. Clearly $\text{tp}(\bar{\mathbf{a}}_\ell, B^+)$ does not fork over $A \subseteq B$ and $\text{acl}(B) = B$ and so $\text{stp}(\bar{\mathbf{a}}_1, B^+) = \text{stp}(\bar{\mathbf{a}}_2, B^+)$. Also $\text{tp}_*(A_2, B^+ \cup A_1)$ does not fork over A hence $\text{tp}(\bar{\mathbf{a}}_2, B^+ \cup \bar{\mathbf{a}}_1)$ does not fork over $A \subseteq B^+$ hence $\{\bar{\mathbf{a}}_1, \bar{\mathbf{a}}_2\}$ is independent over B^+ hence there is an elementary mapping g^+ from

$acl(B^+ \cup \bar{\mathbf{a}}_1)$ onto $acl(B^+ \cup \bar{\mathbf{a}}_2)$, $g^+ \supseteq id_{B^+} \cup g$ and even $g' = g^+ \cup (g^+)^{-1}$ is an elementary embedding.

Let $\bar{\mathbf{a}}'_1$ list $acl(N_{<>} \cup A_1)$ so clearly $\bar{\mathbf{a}}'_2 =: g^+(\bar{\mathbf{a}}'_1)$ list $acl(N_{<>} \cup A_2)$. Clearly $g' \upharpoonright (\bar{\mathbf{a}}'_1 \cup \bar{\mathbf{a}}'_2)$ is an elementary mapping from $\bar{\mathbf{a}}'_1 \cup \bar{\mathbf{a}}'_2$ onto itself. Now $N_{<>}^2$ is \aleph_ϵ -primary over $N_{<>} \cup A_1 \cup A_2$ and $N_{<>} \cup A_1 \cup A_2 \subseteq \bar{\mathbf{a}}'_1 \cup \bar{\mathbf{a}}'_2 \subseteq acl(N_{<>} \cup A_1 \cup A_2)$ so by 1.18(10) $N_{<>}^2$ is \aleph_ϵ -primary over $N_{<>} \cup \bar{\mathbf{a}}'_1 \cup \bar{\mathbf{a}}'_2$ hence we can extend $g' \upharpoonright (\bar{\mathbf{a}}'_1 \cup \bar{\mathbf{a}}'_2)$ to an automorphism $h_{<>}$ of $N_{<>}^2$ so clearly $h_{<>} \upharpoonright N_{<>} = id_{N_{<>}}$. Let $\bar{\mathbf{a}}_1^+$ list $acl(B^+ \cup A_1)$ and $\bar{\mathbf{a}}_2^+ = g^+(\bar{\mathbf{a}}_1^+)$. So $tp(\bar{\mathbf{a}}_\ell^+, N_{<>}^2)$ does not fork over $\bar{\mathbf{a}}'_1 (\subseteq N_{<>}^2)$ and $acl(\bar{\mathbf{a}}'_1) = \text{Rang}(\bar{\mathbf{a}}'_1) (= acl(N_{<>} \cup A_1))$ and $h_{<>} \upharpoonright \bar{\mathbf{a}}'_1 = g^+ \upharpoonright \bar{\mathbf{a}}'_1$ hence $h_{<>} \cup g^+$ is an elementary embedding. Remember g^+ is the identity on $B^+ = acl(N_{<>} \cup B)$, and $tp_*(N_{<0>}, N_{<>}^2)$ does not fork over $N_{<>}$ hence $tp_*(N_{<0>}, B^+ \cup N_{<>}^2)$ does not fork over B^+ , so as $acl(B^+) = B^+$ necessarily $(h_{<>} \cup g^+) \cup id_{N_{<0>}}$ is an elementary embedding. But this mapping has domain and range including $N_{<0>} \cup N_{<>}^2$ and included in $N_{<0>}^2$, but the latter is \aleph_ϵ -primary and \aleph_ϵ -minimal over the former. Hence $(h_{<>} \cup g^+) \cup id_{N_{<>}}$ can be extended to an automorphism of $N_{<0>}^2$ which we call $h_{<0>}$.

Now we define by induction on $n \in [2, \omega)$ for every $\eta \in I$ of length n , an automorphism h_η of N_η^2 extending $h_{\eta-} \cup id_{N_\eta}$, which exists as N_η^2 is \aleph_ϵ -primary over $N_{\eta-}^2 \cup N_\eta$ (and $N_{\eta-}^2 \cup N_\eta \subseteq \bigcup_{N_{\eta-}} N_\eta$). Now $\bigcup_{\eta \in I} h_\eta$ is an elementary mapping (as $\langle N_\eta^2 : \eta \in I \rangle$

is a non-forking tree; i.e. 1.13(10)), with domain and range $\bigcup_{\eta \in I} N_\eta^2$ hence can be

extended to an automorphism h^* of M' , (we can demand $h^* \upharpoonright M^- = id_{M^-}$ but not necessarily). So as h^* extends g , the conclusion follows.

3) Similarly to (2).

□_{1.27}

1.28 Claim. 1) For every $\Upsilon = tp_\delta \left[\begin{pmatrix} B \\ A \end{pmatrix}, M \right]$, and $\bar{\mathbf{a}}, \bar{\mathbf{b}}$ listing A, B respectively there is $\psi = \psi(\bar{x}_A, \bar{x}_B) \in \mathbb{L}_{\infty, \aleph_\epsilon}$ (q.d.) of depth δ such that:

$$tp_\delta \left[\begin{pmatrix} B \\ A \end{pmatrix}, M \right] = \Upsilon \Leftrightarrow M \models \psi[\bar{\mathbf{a}}, \bar{\mathbf{b}}].$$

2) Assume \otimes_{M_1, M_2} of 1.4 holds as exemplified by the family \mathcal{F} and $\begin{pmatrix} B \\ A \end{pmatrix} \in \Gamma(M_1)$ and $g \in \mathcal{F}$, $\text{Dom}(g) = B$; and α an ordinal then

$$tp_\alpha \left(\begin{pmatrix} B \\ A \end{pmatrix}, M \right) = tp_\alpha \left(\begin{pmatrix} g(B) \\ g(A) \end{pmatrix}, M_2 \right).$$

3) Similarly for $tp_\alpha([A], M), tp_\alpha[M]$.

Proof. Straightforward (remember we assume that every first order formula is equivalent to a predicate). □_{1.28}

1.29 Proof of Theorem 1.2. [The proof does not require that the M^ℓ are \aleph_ϵ -saturated, but only that 1.27, 1.28 hold except in constructing $g_{\alpha(*)}$ (see $\otimes_{14}, \otimes_{15}$ in 1.30(E), we could instead use NOTOP].

So suppose

$$(*)_0 \quad M^1 \equiv_{\mathbb{L}_{\infty, \aleph_\epsilon}(\text{d.q.})} M^2 \text{ or (at least) } \otimes_{M^1, M^2} \text{ from 1.4 holds.}$$

We shall prove $M^1 \cong M^2$. By 1.28 (i.e. by 1.28(1) if the first possibility in $(*)_0$ holds and by 1.28(2) if the second possibility in $(*)_0$ holds)

$$(*)_1 \quad \text{tp}_\infty[M^1] = \text{tp}_\infty[M^2].$$

So it suffices to prove:

1.30 Claim. *Assume that T is countable. If M^1, M^2 are \aleph_ϵ -saturated models (of T, T as in 1.5), then:*

$$(*)_1 \Rightarrow M^1 \cong M^2.$$

Proof. Let $\langle W_k, W'_k : k < \omega \rangle$ be a partition of ω to infinite sets (so pairwise disjoint).

1.31 Explanation: (If seems opaque, the reader may return to it after reading parts of the proof).

We shall now define an approximation to a decomposition. That is we are approximating a non-forking tree $\langle N_\eta^\ell, a_\eta^\ell : \eta \in I^* \rangle$ of countable elementary submodels of M^ℓ for $\ell = 1, 2$ and $\langle f_\eta^* : \eta \in I^* \rangle$ such that f_η^* an isomorphism from N_η^1 onto N_η^2 increasing with η such that M^ℓ is \aleph_ϵ -prime over $\bigcup_{\eta \in I^*} N_\eta^\ell$.

In the approximation Y we have:

- (α) I approximating I^* ,
[it will not be $I^* \cap^{n \geq} \text{Ord}$ but we may “discover” more immediate successors to each $\eta \in I$; as the approximation to N_η improves we have more regular types, but some member of I will be later will drop]
- (β) A_η^ℓ approximates N_η^ℓ and is ϵ -finite
- (γ) a_η^ℓ is the a_η^ℓ (if η survives, i.e. will not be dropped)

- (δ) $B_\eta^\ell, b_{\eta,m}^\ell$ expresses commitments on constructing A_η^ℓ : we “promise”
 $B_\eta^\ell \subseteq N_\eta^\ell$ and B_η^ℓ is countable; $b_{\eta,m}^\ell$ for $m < \omega$ list B_η^ℓ (so in the choice
 $B_\eta^\ell \subseteq M^\ell$ there is some arbitrariness)
- (ε) f_η approximate f_η^*
- (ζ) $p_{\eta,m}^\ell$ also expresses commitments on the construction.

Since there are infinitely many commitments that we must meet in a construction of length ω and we would like many chances to meet each of them, the sets W_k, W'_k are introduced as a further bookkeeping device. At stage n in the construction we will deal e.g. with the $b_{\eta,m}^\ell$ for η that are appropriate and for $m \in W_k$ for some $k < n$ and analogously for $p_{\eta,m}^\ell$ and the W'_k .

Note that while the A_η^ℓ satisfy the independence properties of a decomposition, the B_η^ℓ do not and may well intersect non-trivially. Nevertheless, a conflict arises if an $a_{\eta \hat{< i >}}^\ell$ falls into B_η^ℓ since the $a_{\eta \hat{< i >}}^\ell$ are supposed to represent independent elements realizing regular types over the model approximated by A_η^ℓ but now $a_{\eta \hat{< i >}}^\ell$ is in that model. This problem is addressed by pruning $\eta \hat{< i >}$ from the tree I .

1.32 Definition. An approximation Y to an isomorphism consist of:

- (a) natural numbers n, k^* and index set: $I \subseteq {}^{n \geq} \text{Ord}$
(and n minimal)
- (b) $\langle A_\eta^\ell, B_\eta^\ell, a_{\eta,m}^\ell, b_{\eta,m}^\ell : \eta \in I \text{ and } m \in \bigcup_{k < k^*} W_k \rangle$ for $\ell = 1, 2$ (this is an approxi-
mated decomposition)
- (c) $\langle f_\eta : \eta \in I \rangle$
- (d) $\langle p_{\eta,m}^\ell : \eta \in I \text{ and } m \in \bigcup_{k < k^*} W'_k \rangle$

such that:

- (1) I closed under initial segments
- (2) $\langle \rangle \in I$
- (3) $A_\eta^\ell \subseteq B_\eta^\ell \subseteq M^\ell$, A_η^ℓ is ϵ -finite, $\text{acl}(A_\eta^\ell) = A_\eta^\ell$, B_η^ℓ is countable,
 $B_\eta^\ell = \{b_{\eta,m}^\ell : m \in \bigcup_{k < k^*} W_k\}$
- (4) $A_\nu^\ell \subseteq A_\eta^\ell$ if $\nu \triangleleft \eta \in I$

- (5) if $\eta \in I \setminus \{\langle \rangle\}$, then $\frac{a_\eta^\ell}{A_{(\eta^-)}^\ell}$ is a (stationary) regular type and $a_\eta^\ell \in A_\eta^\ell$; in addition $\ell g(\eta) > 1$ then $\frac{a_\eta^\ell}{A_{(\eta^-)}^\ell} \perp A_{(\eta^-)}^\ell$ (note that we may decide $a_{\langle \rangle}^\ell$ be not defined or $\in A_{\langle \rangle}^\ell$)
- (6) $\frac{A_\eta^\ell}{A_{\eta^-+a_\eta}^\ell} \perp_a A_{\eta^-}^\ell$ if $\eta \in I, \ell g(\eta) > 0$
- (7) if $\eta \in I$, not \leftarrow -maximal in I , then the set $\{a_\nu^\ell : \nu \in I \text{ and } \nu^- = \eta\}$ is a maximal family of elements realizing over A_η^ℓ regular types $\perp A_{(\eta^-)}^\ell$ (when η^- is defined), independent over $(A_\eta^\ell, B_\eta^\ell)$, (and we can add: if $\nu_1^- = \nu_2^- = \eta$ and $\frac{a_{\nu_1}^\ell}{A_\eta} \pm \frac{a_{\nu_2}^\ell}{A_\eta}$ then $a_{\nu_1}^\ell/A_\eta = a_{\nu_2}^\ell/A_\eta$)
- (8) f_η is an elementary map from A_η^1 onto A_η^2
- (9) $f_{(\eta^-)} \subseteq f_\eta$ when $\eta \in I, \ell g(\eta) > 0$
- (10) $f_\eta(a_\eta^1) = a_\eta^2$
- (11) (α) $f_\eta \left(\text{tp}_\infty \left[\left(\frac{A_\eta^1}{A_{(\eta^-)}^1}, M^1 \right) \right] \right) = \text{tp}_\infty \left[\left(\frac{A_\eta^2}{A_{(\eta^-)}^2}, M^2 \right) \right]$ when $\eta \in I \setminus \{\langle \rangle\}$
- (β) $f_{\langle \rangle} \left(\text{tp}_\infty [A_{\langle \rangle}^1, M^1] \right) = \text{tp}_\infty [A_{\langle \rangle}^2, M^2]$
- (12) $B_\eta^\ell \prec M^\ell$ moreover, $B_\eta^\ell \subseteq_{\text{na}} M^\ell$, i.e., if $\bar{a} \subseteq N_\eta^\ell, b \in M^\ell \setminus B_\eta^\ell$ and $M^\ell \models \varphi(b, \bar{a})$ there for some $b' \in B_\eta^\ell, \models \varphi(b', \bar{a})$ and $b \notin \text{acl}(\bar{a}) \Rightarrow b' \notin \text{acl}(A)$
- (13) $\langle p_{\eta, m}^\ell : m \in \bigcup_{k < k^*} W_k' \rangle$ is a sequence of types over A_η^ℓ (so $\text{Dom}(p_{\eta, m}^\ell)$ may be a proper subset of A_η^ℓ).

1.33 Notation. We write $n = n_Y = n[Y], I = I_Y = I[Y], A_\eta^\ell = A_\eta^\ell[Y], B_\eta^\ell = B_\eta^\ell[Y], f_\eta = f_\eta^Y = f_\eta[Y], a_\eta^\ell = a_\eta^\ell[Y], b_\eta^\ell = b_\eta^\ell[Y], k^* = k_Y^* = k^*[Y]$ and $p_{\eta, m}^\ell = p_{\eta, k}^\ell[Y]$.

Remark. We may decide to demand: each $\frac{a_{\eta^{\langle i \rangle}}^\ell}{A_\eta}$ is strongly regular; also: if two such types are not orthogonal then they are equal (or at least have same witness φ for $(\varphi, \frac{a_{\eta^{\langle i \rangle}}^\ell}{A_\eta})$ regular). This is easy here as the models are \aleph_ϵ -saturated (so take $p' \pm p, \text{rk}(p')$ minimal).

1.34 Observation. $(*)_1$ implies that there is an approximation, (see 1.29).

Proof. Let $I = \{\langle \rangle\}$, $A_{\langle \rangle}^\ell = \text{acl}(\emptyset)$, $k^* = 1$ and then choose countable $B_{\langle \rangle}^\ell$ to satisfy condition (12) and then choose $f_\eta, p_k^\ell, b_{\eta,m}^\ell$ (for $k \in W'_0$ and $m \in W_0$) as required.

1.35 Main Fact. For any approximation $Y, i \in \bigcup_{k < k_Y^*} (W_k \cup W'_k)$ and $m \leq n_Y$ and $\ell(*) \in \{1, 2\}$ we can find an approximation Z such that:

- (\otimes)(α) $n_Z = \text{Max}\{m + 1, n_Y\}$, $I_Z \cap^{m \geq \text{Ord}} = I_Y \cap^{m \geq \text{Ord}}$,
 (we mean m not n_Y) and $k_Z^* = k_Y^* + 1$
 (β) (a) if $\eta \in I_Y, \ell g(\eta) < m$ then

$$A_\eta^\ell[Z] = A_\eta^\ell[Z],$$

$$a_\eta^\ell[Z] = a_\eta^\ell[Z]$$

$$B_\eta^\ell[Z] = B_\eta^\ell[Y]$$

- (b) if $\eta \in I_Y \cap I_Z, k < k_Y^*$ and $j \in W'_k$ then
 $p_{\eta,j}^\ell[Z] = p_{\eta,j}^\ell[Y]$
 (c) if $\eta \in I_Y \cap I_Z, k < k_Y^*$ and $j \in W_k$ then
 $b_{\eta,j}^\ell[Z] = b_{\eta,j}^\ell[Y]$

(γ)¹ if $\eta \in I_Y, \ell g(\eta) = m, k < k_Y^*$ and⁴ $i \in W_k$ and the element $b \in M^{\ell(*)}$ satisfies clauses (a), (b) below then for some such b we have: $A_\eta^{\ell(*)}[Z] = \text{acl}(A_\eta^{\ell(*)}[Y] \cup \{b\})$; where

(a) $b_{\eta,i}^{\ell(*)}[Y] \notin A_\eta^{\ell(*)}[Y]$ and $\ell g(\eta) > 0 \Rightarrow \frac{b}{A_\eta^{\ell(*)}[Y]} \perp_a A_{\eta^-}^{\ell(*)}[Y]$

(b) one of the conditions (i),(ii) listed below holds for b

(i) $b = b_{\eta,i}^{\ell(*)}[Y]$ and
 $\ell g(\eta) > 0 \Rightarrow \frac{b}{A_\eta^{\ell(*)}[Y]} \perp_a A_{\eta^-}^{\ell(*)}[Y]$ or

(ii) for no b is (i) satisfied (so $\ell g(\eta) > 0$) and $b \in M^{\ell(*)}$,

$$b_{\eta,i}^\ell \upharpoonright_{A_\eta^{\ell(*)}[Y]} b \text{ and } \ell g(\eta) > 0 \Rightarrow \frac{b}{A_\eta^{\ell(*)}[Y]} \perp_a A_{\eta^-}^{\ell(*)}[Y]$$

⁴recall that i is part of the information given in the main fact, and of course, k is uniquely determined by i

(γ)² assume $\eta \in I_Y$, $\ell g(\eta) = m$, $k < k_Y^*$ and $i \in W'_k$ then we have:

(a) if $p_{\eta,i}^{\ell(*)}$ is realized by some $b \in M^{\ell(*)}$ such that

$$\text{Rk} \left(\frac{b}{A_{\eta}^{\ell(*)}[Y]}, L, \infty \right) = \text{R} \left(p_{\eta,i}^{\ell(*)}, L, \infty \right) \text{ and}$$

$$\left[\ell g(\eta) > 0 \Rightarrow \frac{b}{A_{\eta}^{\ell(*)}[Y]} \perp_a A_{\eta^-}^{\ell(*)}[Y] \right]$$

then for some such b we have

$$A_{\eta}^{\ell(*)}[Z] = \text{acl} \left(A_{\eta}^{\ell(*)}[Y] \cup \{b\} \right)$$

(b) if the assumption of clause (a) fails but $p_{\eta,i}^{\ell(*)}$ is realized

by some $b \in M^{\ell(*)} \setminus A_{\eta}^{\ell(*)}$ such that

$$\left[\ell g(\eta) > 0 \Rightarrow \frac{b}{A_{\eta}^{\ell(*)}[Y]} \perp_a A_{\eta^-}^{\ell(*)}[Y] \right]$$

then for some such b we have

$$A_{\eta}^{\ell(*)}[Z] = \text{acl} \left[A_{\eta}^{\ell(*)}[Y] \cup \{b\} \right]$$

(δ) If $\eta \in I_Y$ and $\ell g(\eta) = m$, then $B_{\eta}^{\ell}[Z] = \{b_{\eta,j}^{\ell}[Y] : j \in \cup\{W_k : k < k_Z^*\}\}$ is a countable subset of M^{ℓ} , containing $\{B_{\nu}^{\ell}[Z] : \nu \leq \eta \text{ and } \nu \in Y\} \cup B_{\eta}^{\ell}[Y]$, with $B_{\eta}^{\ell}[Z] \prec M^{\ell}$ moreover $B_{\eta}^{\ell}[Z] \subseteq_{na} M^{\ell}$ i.e. if $\bar{a} \subseteq B_{\eta}^{\ell}[Z]$, $\varphi(x, \bar{y})$ is first order and $(\exists x \in M^{\ell} \setminus \text{acl}(\bar{a})) \varphi(x, \bar{a})$ then $(\exists x \in B_{\eta}^{\ell}[Z] \setminus \text{acl}(\bar{a})) \varphi(x, \bar{a})$ and $\{a_{\eta^{\wedge} < \alpha >}^{\ell}[Y] : \eta^{\wedge} < \alpha > \in I_Y \text{ and } a_{\eta^{\wedge} < \alpha >}^{\ell}[Y] \notin B_{\eta}^{\ell}[Z]\}$ is independent over $(B_{\eta}^{\ell}[Z], A_{\eta}^{\ell}[Y])$

(ϵ) if $\eta \in I_Y$, $\ell g(\eta) > m$, then $\eta \in I_Z \Leftrightarrow a_{\eta \upharpoonright (m+1)}^{\ell}[Y] \notin B_{\eta \upharpoonright m}^{\ell}[Z]$

(ζ) if $\eta \in I_Y \cap I_Z$, $\ell g(\eta) > m$ then $A_{\eta}^{\ell}[Z] = \text{acl}(A_{\eta}^{\ell}[Y] \cup A_{\eta \upharpoonright m}^{\ell}[Z])$ and $B_{\eta}^{\ell}[Z] = B_{\eta}^{\ell}[Y]$

(η) if $\eta \in I_Z \setminus I_Y$ then $\eta^- \in I_Y$ and $\ell g(\eta) = m + 1$

(θ) $\{p_{\eta,i}^{\ell}[Z] : i \in W'_{k_Z^* - 1}\}$ is “rich enough”, e.g. include all finite types over A_{η}^{ℓ}

(ι) $\{b_{\eta,i}^{\ell} : i \in W_{k_Z^* - 1}\}$ list $B_{\eta}^{\ell}[Z]$, each appearing infinitely often.

Proof. First we choose $A_\eta^{\ell(*)}[Z]$ for $\eta \in I$ of length m according to condition $(\gamma) = (\gamma)^1 + (\gamma)^2$. (Note: one of the clauses $(\gamma)^1, (\gamma)^2$ necessarily holds trivially as $\bigcup_k W_k \cap \bigcup_k W'_k = \emptyset$).

Second, we choose (for such η) an elementary mapping f_η^Z extending f_η^Y and a set $A_\eta^{3-\ell(*)}[Z] \subseteq M^{3-\ell(*)}$ satisfying “ f_η^Z is from $A_\eta^1[Z]$ onto $A_\eta^{3-\ell(*)}[Z]$ ” such that

(*)₂ if $m > 0$, then

$$f_\eta^Z(\text{tp}_\infty((A_\eta^1[Z], M_1))) = \text{tp}_\infty((A_\eta^2[Z], M_2))$$

(*)₃ if $m = 0$, then $f_\eta^Z(\text{tp}_\infty(A_\eta^1[Z], M_1)) = \text{tp}_\infty(A_\eta^2[Z], M_2)$.

[Why possible? If we ask just the equality of tp_α for an ordinal α , this follows by the first component of $\text{tp}_{\alpha+1}$. But (overshooting) for $\alpha \geq [(\|M_1\| + \|M_2\|)^{|T|}]^+$, equality of tp_α implies equality of tp_∞].

Third, we choose $B_\eta^\ell[Z]$ for $\eta \in I_Y, \ell g(\eta) = m$ according to condition (δ) (here we use the countability of the language, you can do it by extending it ω times) in both sides, i.e. for $\ell = 1, 2$.

Fourth, let $I' = \{\eta \in I : \text{if } \ell g(\eta) > m \text{ then } a_{\eta \upharpoonright (m+1)}^\ell[Y] \notin B_{\eta \upharpoonright m}^\ell[Z]\}$ (this will be $I_Y \cap I_Z$).

Fifth, we choose $A_\eta^\ell[Z]$ for $\eta \in I'$: if $\ell g(\eta) < m$, let $A_\eta^\ell[Z] = A_\eta^\ell[Y]$, if $\ell g(\eta) = m$ this was done, lastly if $\ell g(\eta) > m$, let $A_\eta^\ell[Z] = \text{acl}(A_\eta^\ell[Y] \cup A_{\eta \upharpoonright m}^\ell[Z])$.

Sixth, by induction on $k \leq n_Y$ we choose f_η^Z for $\eta \in I'$ of length k : if $\ell g(\eta) < m$, let $f_\eta^Z = f_\eta^Y$, if $\ell g(\eta) = m$ this was done, lastly if $\ell g(\eta) > m$ choose an elementary mapping from A_η^1 onto A_η^2 extending $f_\eta^Y \cup f_{\eta^-}^Z$ (possible as $f_\eta^Y \cup f_{\eta^-}^Z$ is an elementary mapping and $\text{Dom}(f_\eta^Y) \cap \text{Dom}(f_{\eta^-}^Z) = A_{\eta^-}^{\ell(*)}$, $\text{Dom}(f_\eta^Y) \bigcup_{A_{\eta^-}^{\ell(*)}} \text{Dom}(f_{\eta^-}^Z)$)

and $A_{\eta^-}^{\ell(*)} = \text{acl}(A_{\eta^-}^{\ell(*)})$.) Now f_η^Z satisfies clause (11) of Definition 1.32 when $\ell g(\eta) > m$ by applying 1.27(3).

Seventh, for $\eta \in I'$, of length $m < n_Z$, let $v_\eta =: \{\alpha : \eta \hat{<} \alpha \in I\}$, and we choose $\{a_{\eta \hat{<} \alpha}^1[Z] : \alpha \in u_\eta\}, [\alpha \in u_\eta \Rightarrow \eta \hat{<} \alpha \notin I]$, a set of elements of M^1 realizing (stationary) regular types over $A_\eta^1[Z]$, orthogonal to $A_{\eta^-}[Y]$ when $\ell g(\eta) > 0$, such that it is independent over $(\cup\{a_{\eta \hat{<} \alpha}^1[Y] : \eta \hat{<} \alpha \in I'\} \cup B_\eta^1[Z], A_\eta^1[Z])$ and maximal under those restrictions. Without loss of generality $\sup(v_\eta) < \min(u_\eta)$ and for $\alpha_1 \in v_\eta \cup u_\eta$ and $\alpha_2 \in u_\eta$ we have:

- (*)₁ if (for the given α_2 and η) α_1 is minimal such that $\frac{a_{\eta^\wedge <\alpha_1>}^1[Z]}{A_\eta^1[Z]} \pm \frac{a_{\eta^\wedge <\alpha_2>}^1[Z]}{A_\eta^1[Z]}$ then $\frac{a_{\eta^\wedge <\alpha_1>}^1[Z]}{A_\eta^1[Z]} = \frac{a_{\eta^\wedge <\alpha_2>}^1[Z]}{A_\eta^1[Z]}$
- (*)₂ if $\alpha_1 < \alpha_2$ and $a_{\eta^\wedge <\alpha_1>}^1[Z]/A_\eta^1[Z] = a_{\eta^\wedge <\alpha_2>}^1[Z]/A_\eta^1[Z]$ and for some $b \in M^1$ realizing $\frac{a_{\eta^\wedge <\alpha_2>}^1[Z]}{A_\eta^1[Z]}$ we have $b \uplus_{A_{\eta^\ell}^1[Z]} a_{\eta^\wedge <\alpha_2>}^1$ and $\text{tp}_\infty \left[\left(A_{\eta^\wedge <\alpha_2>}^1, M \right), M \right] = \text{tp}_\infty \left[\left(\frac{a_{\eta^\wedge <\alpha_1>}^1}{A_{\eta^\wedge <\alpha_2>}^1}, M \right), M \right]$ and α_1 is minimal (for the given α_2 and η) then $\text{tp}_\infty \left[\left(\frac{a_{\eta^\wedge <\alpha_2>}^1}{A_{\eta^\wedge <\alpha_2>}^1}, M \right), M \right] = \text{tp}_\infty \left[\left(\frac{a_{\eta^\wedge <\alpha_1>}^1}{A_{\eta^\wedge <\alpha_1>}^1}, M \right), M \right]$.

Easily (as in [Sh:c, X]) if $\alpha \in u_\eta$ and $\eta^\wedge \langle \beta \rangle \in I'$ then $\frac{a_{\eta^\wedge <\alpha>}^1[Z]}{A_\eta^1[Z]} \perp \frac{a_{\eta^\wedge <\beta>}^1[Y]}{A_\eta^1[Y]}$.

For $\alpha \in u_\eta$ let $A_{\eta^\wedge <\alpha>}^1[Z] = \text{acl} \left(A_\eta^1[Y] \cup \{a_{\eta^\wedge <\alpha>}^1[Z]\} \right)$.

Eighth, by the second component in the definition of $\text{tp}_{\alpha+1}$ (see Definition 1.9) we can choose (for $\alpha \in u_\eta$) $a_{\eta^\wedge <\alpha>}^2[Z]$, $A_{\eta^\wedge <\alpha>}^2[Z]$ and then $f_{\eta^\wedge <\alpha>}^Z$ as required (see (7) of Definition 1.32).

Ninth and lastly, we let $I_Z = I' \cup \{\eta^\wedge \langle \alpha \rangle : \eta \in I', \ell g(\eta) = m < n_Z \text{ and } \alpha \in u_\eta\}$ and we choose B_η^ℓ for $\eta \in I_Z \setminus I_Y$ and the $p_{\eta,i}^\ell, b_{\eta,j}^\ell$ as required (also in other case left).

□_{1.35}

1.36 Finishing the Proof of 1.11. We define by induction on $n < \omega$ an approximation $Y_n = Y(n)$. Let Y_0 be the trivial one (as in observation 1.30(C)).

Y_{n+1} is gotten from Y_n as in 1.35 for $m_n, i_n \leq n, \ell_n(*) \in \{1, 2\}$ defined by reasonable bookkeeping (so $i_n \in \bigcup_{k < k_Y^*(n)} (W_k \cup W'_k)$) such that any triples appear

infinitely often; without loss of generality: if $n_1 < n_2$ & $\eta \in I_{n_1}^\ell \cap I_{n_2}^\ell$ then $\eta \in \bigcap_{n=n_1}^{n_2} I_n$.

Let $I^* = I[*] = \lim(I_\ell^{Y(n)}) =: \{\eta : \text{for every large enough } n, \eta \in I_n\}$;

for $\eta \in I^*$ let: $A_\eta^\ell[*] = \bigcup_{n < \omega} A_\eta^\ell[Y_n], f_\eta^\ell[*] = \bigcup_{n < \omega} f_\eta^{Y(n)}$ and

$$B_\eta^\ell[*] = \bigcup_{n < \omega} B_\eta^\ell[Y_n].$$

Easily

- \oplus_0 $\langle \rangle \in I^*$ and $I^* \subseteq \omega$ Ord is closed under initial segments
- \oplus_1 for $\eta \in I^*$, $\langle B_\eta^\ell[Y_n] : n < \omega \text{ and } \eta \in I[Y_n] \rangle$ is an increasing sequence of \subseteq_{na} -elementary submodels of M^ℓ
[Why? By clause (12) of Definition 1.32, Main Fact 1.35, clauses $(\beta)(a)$, (δ) , (ζ) .]

hence

$$\oplus_2 \text{ for } \eta \in I^*, B_\eta^\ell[*] \subseteq_{\text{na}} M^\ell.$$

Also

$$\oplus_3 \nu \triangleleft \eta \in I^* \Rightarrow B_\nu^\ell[*] \subseteq B_\eta^\ell[*].$$

[Why? Because for infinitely many n , $m_n = \ell g(\eta)$ and clause (δ) of Main Fact 1.35].

$$\oplus_4 \text{ if } \eta \in I[Y_{n_1}] \cap I^*, \eta^- = \nu \text{ and } n_1 \leq n_2 \text{ then}$$

$$A_\eta^\ell[Y_{n_1}] \cup A_\nu^\ell[Y_{n_2}].$$

[Why? Prove by induction on n_2 (using the non-forking calculus), for $n_2 = n_1$ this is trivial, so assume $n_2 > n_1$. If $m_{(n_2-1)} > \ell g(\nu)$ we have $A_\nu^\ell[Y_{n_2}] = A_\nu^\ell[Y_{n_2-1}]$ (see 1.35, clause $(\beta)(a)$ and we have nothing to prove. If $m_{(n_2-1)} < \ell g(\nu)$ then we note that $A_\nu^\ell[Y_{n_2}] = \text{acl}(A_\nu^\ell[Y_{n_2-1}] \cup A_{\nu \upharpoonright m_{(n_2-1)}}^\ell[Y_{n_2}])$ and $A_\nu^\ell[Y_{n_2-1}] \cup A_{\nu \upharpoonright m_{(n_2-1)}}^\ell[Y_{n_2}]$

(as $\nu \in I[Y_{n_2}]$, by 1.35 clause (δ) last phrase) and now use clauses (5), (6) of Definition 1.35. Lastly if $m_{(n_2-1)} = \ell g(\nu)$ again use $\nu \in I[Y_{n_2}]$ by 1.35, clause (δ) , last phrase].

$$\oplus_5 \text{ if } \eta \in I[Y_{n_1}] \cap I^*, \eta^- = \nu \text{ and } n_1 \leq n_2 \text{ then}$$

$$\frac{A_\eta^\ell[*]}{A_\nu^\ell[*] + a_\eta^\ell[*]} \perp_a A_\nu^\ell.$$

[Why? By clause (6) of Definition 1.32, and orthogonality calculus].

- \oplus_6 if $\eta \in I^*$, then $A_\eta^\ell[*] \subseteq B_\eta^\ell[*] \prec M^\ell$ moreover
- \otimes_7 $A_{\eta(*)}^\ell[*] \subseteq_{\text{na}} B_\eta^\ell[*] \subseteq_{\text{na}} M^\ell$.

[Why? The second relation holds by \otimes_2 . The first relation we prove by induction on $lg(\eta)$; clearly $A_\eta^\ell[*] = acl(A_\eta^\ell[*])$ because $A_\eta^\ell[Y_n]$ increases with n by 1.35 and $A_\eta^\ell[Y_n] = acl(A_\eta^\ell[Y_n])$ by clause (3) of Definition 1.32. We prove “ $A_{\eta(*)}^\ell[*] \subseteq_{na} B_\eta^\ell[*]$ ” by induction on $m = lg(\eta)$, so suppose this is true for every $m' < m, m = lg(\eta), \eta \in I^*$, let $\varphi(x)$ be a formula with parameters in $A_\eta^\ell[*]$ realized in M^ℓ as above say by $b \in M^\ell$. As $\langle A_\eta^\ell[Y_n] : n < \omega, \eta \in Y_n \rangle$ is increasing with union $A_\eta^\ell[*]$, clearly for some n we have $b \bigcup_{A_\eta^\ell[Y_n]} A_\eta^\ell[*]$.

So $\{\varphi(x)\} = p_{\eta,i}^\ell$ for some i and for some $n' > n$ defining $Y_{n'+1}$ we have used 1.35 with $(\ell(*), i, m)$ there being $(\ell, i, lg(\eta))$ here, hence we consider clause $(\gamma)^2$ of 1.35. So the case left is when the assumption of both clauses (a) and (b) of $(\gamma)^2$ fail, so we have $lg(\eta) > 0$ and

$$b' \notin A_\eta^\ell[Y_{n'}], b' \in M^\ell \models \varphi[b'] \Rightarrow \frac{b'}{A_\eta^\ell[Y_{n'}]} \pm A_{\eta^-}^\ell[Y_{n'}].$$

We can now use the induction hypothesis (and [BeSh 307, 5.3,p.292]).]

\oplus_8 if $\eta \in I^*$ and $\ell = 1, 2$, then
 $\{a_{\eta \hat{<} \alpha}^\ell[*] : \eta \hat{<} \alpha \in I^*\}$ is a maximal subset of

$$\{c \in M_\ell : \frac{c}{A_\eta^\ell[*]} \text{ regular, } c \bigcup_{A_\eta^\ell[*]} B_\eta^\ell[*] \text{ and } lg(\eta) > 0 \Rightarrow \frac{c}{A_\eta^\ell[*]} \perp A_{\eta^-}^\ell[*]\}$$

independent over $(A_\eta^\ell[*], B_\eta^\ell[*])$.

[Why? Note clause (7) of Definition 1.32 and clause (δ) of Main Fact 1.35].

\otimes_9 $A_{<>}^\ell[*] = B_{<>}^\ell[*]$

[Why? By the bookkeeping every $b \in B_{<>}^\ell[*]$ is considered for addition to $A_{<>}^\ell[*]$ see 1.35, clause $(\gamma)^1$, subclause (b)(i) and for $\langle \rangle$ there is nothing to stop us].

\otimes_{10} if $\eta \in I^* \setminus \langle \rangle$ and $p \in S(A_\eta^\ell[*])$ is regular orthogonal to $A_{\eta^-}^\ell[*]$ then $\frac{B_\eta^\ell[*]}{A_\eta^\ell[*]} \perp p$.

[Why? If not, as $A_\eta^\ell[*] \subseteq_{na} B_\eta^\ell[*]$ by [BeSh 307, Th.B,p.277] there is

$c \in B_\eta^\ell[*] \setminus A_\eta^\ell[*]$ such that: $\frac{c}{A_\eta^\ell[*]}$ is p . As $c \in B_\eta^\ell[*] = \bigcup_{n < \omega} B_\eta^\ell[Y_n]$, for every $n < \omega$ large enough $c \in B_\eta^\ell[Y_n]$, and p does not fork over $A_\eta^\ell[Y_n]$. So for some such n

the triple (i_n, ℓ_n, m_n) is such that $\ell_n = \ell, m_n = \ell g(\eta)$ and $b_{\eta, i_n}^\ell = c$, so by clause $(\gamma)^1(b)(ii)$ of 1.35 we have $c \in A_\eta^\ell[Y_n] \subseteq A_\eta^\ell[*]$.

\otimes_{11} if $\eta \in I^*, \ell \in \{1, 2\}$ then $\{a_{\eta^\wedge \langle \alpha \rangle} : \eta^\wedge \langle \alpha \rangle \in I^*\}$ is a maximal subset of $\{c \in M^\ell : \frac{c}{A_\eta^\ell[*]} \text{ regular, } \perp A_{\eta^-}^\ell[*] \text{ when meaningful}\}$ independent over $A_\eta^\ell[*]$.

[Why? If not, then for some $c \in M, \{a_{\eta^\wedge \langle \alpha \rangle} : \eta^\wedge \langle \alpha \rangle \in I^*\} \cup \{c\}$ is independent over $A_\eta^\ell[*]$ and $\text{tp}(c, A_\eta^\ell[*])$ is regular (and stationary). Hence by \otimes_{10} we have $\{a_\eta^\ell[Y_n] : \eta^\wedge \langle \alpha \rangle \in I^*\} \cup \{c\}$ is independent over $(A_\eta^\ell[*], B_\eta^\ell[*])$. Now for large enough n we have $c \not\perp A_\eta^\ell[*]$ and by \otimes_{10} we have $c \not\perp B_\eta^\ell[*]$, hence

$c \not\perp B_\eta^\ell[Y_n]$, and so by Definition 1.32(7) $\{c\} \cup \{a_{\eta^\wedge \langle \alpha \rangle}^\ell[Y_n] : \eta^\wedge \langle \alpha \rangle \in I[Y_n]\}$ is $A_\eta^\ell[*]$

not independent over $(A_\eta^\ell[Y_n], B_\eta^\ell[Y_n])$, but $\{a_{\eta^\wedge \langle \alpha \rangle}^\ell[Y_n] : \eta^\wedge \langle \alpha \rangle \in I[Y_n]\}$ is independent over $(A_\eta^\ell[Y_n], B_\eta^\ell[Y_n])$. So there is a finite set w of ordinals such that $\alpha \in w \Rightarrow \eta^\wedge \langle \alpha \rangle \in I[Y_n]$ and $\{c\} \cup \{a_{\eta^\wedge \langle \alpha \rangle}^\ell[Y_n] : \alpha \in w\}$ is not independent over $(A_\eta^\ell[Y_n], B_\eta^\ell[Y_n])$, and without loss of generality w is minimal. Let $n_1 \in [n, \omega)$ be such that $\alpha \in w \ \& \ a_{\eta^\wedge \langle \alpha \rangle}^\ell \in B_\eta^\ell[*] \Rightarrow a_\eta^\ell \in B_\eta^\ell[Y_{n_1}]$; clearly exist as w is finite and let $u = \{\alpha \in w : a_{\eta^\wedge \langle \alpha \rangle}^\ell \notin B_\eta^\ell[*]\}$; clearly $\alpha \in u \Rightarrow \eta^\wedge \langle \alpha \rangle \in I^*$. Now $\{a_{\eta^\wedge \langle \alpha \rangle}^\ell[*] : \eta^\wedge \langle \alpha \rangle \in I^*\} \cup B_\eta^\ell[*]$ includes $\{a_{\eta^\wedge \langle \alpha \rangle}^\ell[Y_n] : \alpha \in w\}$, easy contradiction to the second sentence above.]

$\oplus_{12} f_\eta^* = \bigcup_{m < \omega} f_\eta[Y_m]$ (for $\eta \in I^*$) is an elementary map from $A_\eta^1[*]$ onto $A_\eta^2[*]$.

[Easy].

$\oplus_{13} f^* =: \bigcup_{\eta \in I^*} f_\eta^*$ is an elementary mapping from $\bigcup_{\eta \in I^*} A_\eta^1[*]$ onto $\bigcup_{\eta \in I^*} A_\eta^2[*]$.

[Clear using by $\otimes_5 + \otimes_6 + \otimes_{12}$ and non-forking calculus].

\oplus_{14} We can find $\langle d_\alpha^\ell : \alpha < \alpha(*) \rangle$ such that:

(a) $d_\alpha^\ell \in M^\ell, \beta < \alpha \Rightarrow d_\beta^\ell \neq d_\alpha^\ell$
 $\text{tp}(d_\alpha^\ell, \bigcup_{\eta \in I^*} A_\eta^\ell[*] \cup \{d_\beta^\ell : \beta < \alpha\})$ is \aleph_ϵ -isolated and $\mathbf{F}_{\aleph_0}^\ell$ -isolated, and

(b) $g_\alpha = \bigcup_{\eta \in I^*} f_\eta^* \cup \{\langle (d_\alpha^1, d_\alpha^2) : \alpha < \alpha(*) \rangle\}$ is an elementary mapping,

- (c) $\alpha(*)$ is maximal, i.e., we cannot find $d_{\alpha(*)}^1$ such that the demand in (a) holds for $\alpha(*) + 1$.

[Why? We can try to choose by induction on α , a member d_α^1 of $M^1 \setminus \bigcup_{\eta \in I[*]} \cup \{d_\beta^1 : \beta < \alpha\}$ such that $\text{tp}(d_\alpha^1, \bigcup_{\eta \in I[*]} A_\eta^\ell[*] \cup \{d_\beta^1 : \beta < \alpha\})$ is \aleph_ε -isolated and $\mathbf{F}_{\aleph_0}^\ell$ -isolated.

So for some $\alpha(*)$, d_α^1 is well defined iff $\alpha < \alpha(*)$ (as $\beta < \alpha \Rightarrow d_\beta^1 \neq d_\alpha^1 \in M^1$). Now choose by induction on $\alpha < \alpha(*)$, $d_\alpha^2 \in M^2$ as required above, possible by “ M_i^2 being \aleph_ε -saturated (see [Sh:c, XII,2.1,p.591], [Sh:c, IV,3.10,p.179].]

\otimes_{15} $\text{Dom}(g_{\alpha(*)})$, $\text{Rang}(g_{\alpha(*)})$ are universes of elementary submodels of M^1, M^2 respectively called M'_1, M'_2 respectively.

[Why? See [Sh:c, XII,1.2](2),p.591 and the proof of \otimes_{14} .
Alternatively, choose a formula $\psi(x, \bar{a})$ such that:

- (a) $\bar{a} \subseteq \text{Dom}(g_{\alpha(*)})$ and $\models \exists x \psi(x, \bar{a})$ but no $b \in \text{Dom}(g_{\alpha(*)})$ satisfy $\varphi(x, \bar{a})$
- (b) under clause (a), $\text{Rk}(\psi(x, \bar{a}), \mathbb{L}_{\tau|T|}, \infty)$ is minimal (or just has no extension in $S(\text{Dom}(g_{\alpha(*)}))$) forking over \bar{a}).

Let $\{\varphi_\ell(x, \bar{y}_\ell) : \ell < \omega\}$ list that $\mathbb{L}_{\tau(T)}$ -formulas and we choose by induction on ℓ as formula $\psi_n(x, \bar{a}_n)$ such that:

- (i) $\bar{a} \subseteq \text{Dom}(g_{\alpha(*)})$
- (ii) $\models (\exists x) \psi_n(x, \bar{a}_n)$
- (iii) $\psi_{n+1}(x, \bar{a}_{n+1}) \vdash \psi_n(x, \bar{a}_n)$
- (iv) $\psi_0(x, \bar{a}_0) = \psi(x, \bar{a})$
- (v) for any formula $\psi'(x, \bar{a}')$ satisfying the demands on $\psi_{n+1}(x, \bar{a}_{n+1})$ we have $\text{Rk}(\psi_{n+1}(x, \bar{a}_{n+1}), \{\varphi_n(x, \bar{y}_n)\}, 2) < \text{Rk}(\psi'(x, \bar{a}'), \{\varphi_n(x, \bar{y})\}, 2)$ (on this rank see [Sh:c, II,§2]).

So $p = \{\psi_n(x, \bar{a}_n) : n < \omega\}$ has an extension in $S(\text{Dom}(g_{\alpha(*)}))$ call it q . Now q is \aleph_ε -isolated because $\psi(x, \bar{a}) \in q \in S(\text{Dom}(g_{\alpha(*)}))$. For every n , $\psi_{n+1}(x, \bar{a}_n) \vdash q \upharpoonright \{\varphi_n(x, \bar{y}_n)\}$ by clause (v) above so as $\psi_{n+1}(x, \bar{a}_n) \in q$ and this holds for every n clearly q is $\mathbf{F}_{\aleph_0}^\ell$ -isolated.

\otimes_{16} If $M^\ell \neq M'_\ell$ then for some $d \in M_\ell \setminus M'_\ell$, $\frac{d}{M'_\ell}$ is regular.

[Why? By [BeSh 307, Th.5.9,p.298] as $N_\eta^\ell \subseteq_{\text{na}} M^\ell$ by \otimes_7].

\otimes_{17} if $M^\ell \neq M'_\ell$ then for some $\eta \in I^*$, there is $d \in M^\ell \setminus M'_\ell$ such that $\frac{d}{A_\eta^\ell[*]}$ is regular, $d \cup \bigcup_{A_\eta^\ell[*]} M'_\ell$ and $\left[\ell g(\eta) > 0 \Rightarrow \frac{d}{A_\eta^\ell[*]} \perp A_{\eta^-}^\ell[*] \right]$.

[Why? By [Sh:c, XII,1.4,p.529] every non-algebraic $p \in S(M'_\ell)$ is not orthogonal to some $A_\eta^\ell[*]$ so by \otimes_{16} we can choose $\eta \in I^*$ and $d \in M^\ell \setminus M'_\ell$ such that $\frac{d}{M'_\ell}$ is regular $\perp A_\eta^\ell[*]$; without loss of generality $\ell g(\eta)$ is minimal, now $A_\eta^\ell[*] \subseteq_{\text{na}} M^\ell$ and by [BeSh 307, 4.5,p.290] without loss of generality $d \cup \bigcup_{A_\eta^\ell[*]} M'_\ell$; the last clause is by $A_\eta^\ell[*]$

“ $\ell g(\eta)$ minimal”].

\oplus_{18} $M_\ell = M'_\ell$.

[Why? By $\oplus_{11} + \oplus_{17}$].

\oplus_{19} there is an isomorphism from M_1 onto M_2 extending $\bigcup_{\eta \in I^*} f_\eta^*$.

[Why? By $\oplus_{14} + \otimes_{15}$ we have $M'_1 \cong M'_2$, so by \otimes_{18} we are done]. $\square_{1.36} \square_{1.30}$

1.37 Lemma. *Assume $B \cup C, A = \text{acl}(A) = B \cap C$ and A, B, C are ϵ -finite, $A \cup B \cup C \subseteq M, M$ an \aleph_ϵ -saturated model of T . For notational simplicity make A a set of individual constants.*

Then $\text{tp}_{\mathbb{L}_\infty, \aleph_\epsilon}(d.q.)(B + C; M) = \text{tp}_{\mathbb{L}_\infty, \aleph_\epsilon}(d.q.)(B; M) + \text{tp}_{\mathbb{L}_\infty, \aleph_\epsilon}(d.q.)(C; M)$ where

1.38 Definition. 1) For any logic \mathbb{L} and \bar{b} a sequence from a model M , let

$$\text{tp}_{\mathcal{L}}(\bar{b}; M) = \left\{ \varphi(\bar{x}) : M \models \varphi[B], \varphi \text{ a formula in the vocabulary of } M, \right. \\ \left. \begin{array}{l} \text{from the logic } \mathcal{L} \text{ (with free variables from} \\ \bar{x}, \text{ where } \bar{x} = \langle x_i : i < \ell g(\bar{b}) \rangle) \end{array} \right\}.$$

2) Replacing \bar{b} by a set B means we use the variables $\langle x_b : b \in B \rangle$.

3) Saying $p_1 = p_2 + p_3$ in 1.37 means that we can compute p_1 from p_2 and p_3 (and the knowledge how the variables fit and the knowledge of T , of course).

Proof of the Lemma 1.37.

It is enough to prove:

1.39 Claim. *Assume*

- (a) M^1, M^2 are \aleph_ϵ -saturated and
- (b) $A_1^i \cup_{A_0^i} A_2^i$ for $i = 1, 2$
- (c) $A_0^i = \text{acl}(A_0^i)$ and A_m^i is ϵ -finite for $i = 1, 2$ and $m < 3$
- (d) for $m = 0, 1, 2$ we have $f_m : A_m^1 \xrightarrow{\text{onto}} A_m^2$ is an elementary mapping preserving tp_∞ (in M^1, M^2 respectively) and
- (e) $f_0 \subseteq f_1, f_2$.

Then there is an isomorphism from M^1 onto M^2 extending $f_1 \cup f_2$.

Proof of 1.39. Repeat the proof of 1.5, but starting with Y_0 such that $A_{\langle \rangle}^\ell[Y_0] = A_0^\ell, A_{\langle \rangle}^\ell[Y_0] = A_1^\ell, A_{\langle 1 \rangle}^\ell[Y_0] = A_2^\ell, f_{\langle \rangle}^{Y_0} = f_0, f_{\langle 0 \rangle}^{Y_0} = f_1, f_{\langle 1 \rangle}^{Y_0} = f_2$ and that $\langle \rangle, \langle 0 \rangle, \langle 1 \rangle$ belongs to all $I[Y_0]$. During the construction we preserve $\langle 0 \rangle, \langle 1 \rangle \in I[Y_n]$ and for helping to preserve this we add also the demand

$$\textcircled{*}_{2,m} B_{\langle \rangle}^\ell[Y_n] \cup_{A_0^\ell} A_1^\ell \cup A_2^\ell.$$

During the proof, when we have to increase $B_{\langle \rangle}^\ell$, we use 1.18(1) + 1.16(1).

□_{1.39}

Discussion: A natural version of 1.39 to say is the conclusion only that $\text{tp}_\alpha[(\frac{A_1^1 \cup A_2^1}{A_0^1}), M^1] = \text{tp}_\alpha[(\frac{A_1^2 \cup A_2^2}{A_0^2}), M^2]$ and to prove this by induction on α . The case $\alpha = 0$ and α limit are obvious. If $\alpha = \beta + 1$, for the condition of \leq_a , we use the induction hypothesis and claim 1.27(1). The condition involving \leq_b is similar but harder. □_{1.39}

§2 FINER TYPES

We shall use here alternative types showing us probably a finer way to manipulate tp.

2.1 Convention. T is superstable, NDOP

M, N are \aleph_ϵ -saturated $\prec \mathfrak{C}^{\text{eq}}$.

2.2 Definition. $\Gamma_3 = \left\{ \binom{\bar{b}}{\bar{a}} : \bar{a} \subseteq \bar{b} \text{ are } \epsilon\text{-finite} \right\}$

$\Gamma_1 = \left\{ \binom{p}{\bar{a}} : \bar{a} \text{ is } \epsilon\text{-finite, } p \in S(\bar{a}) \text{ is regular (so stationary)} \right\}$

$\Gamma_2 = \left\{ \binom{p, r}{\bar{a}} : \bar{a} \text{ is } \epsilon\text{-finite, } p \text{ is a regular type of depth } > 0, \right.$
 $p \pm \bar{a}$ (really only the equivalence class p/\pm matters),
 $r = r(x, \bar{y}) \in S(\bar{a})$ is such that for (c, \bar{b}) realizing r ,
 $c/(\bar{a} + \bar{b})$ is regular $\pm p$, and $\frac{\bar{b}}{\bar{a}} = (r \upharpoonright \bar{y}) \perp p \left. \right\}$.

We may add (to Γ_x) superscripts:

- (α) f if \bar{a} (or $\bar{a} \hat{=} \bar{b}$) is finite
- (β) s : for Γ_3 if $\frac{\bar{b}}{\bar{a}}$ is stationary, for Γ_1 if p is stationary which holds always and for Γ_2 if r is stationary and every automorphism of \mathfrak{C} over \bar{a} fix p/\pm
- (γ) c if \bar{a} (or \bar{a}, \bar{b}) are algebraically closed.

2.3 Claim. *If p is regular of depth > 0 and $p \pm \bar{a}$ and \bar{a} is ϵ -finite then for some $\bar{a}', \bar{a} \subseteq \bar{a}' \subseteq \text{acl}(\bar{a})$ and for some q we have $\binom{p, q}{\bar{a}'} \in \Gamma_2^s$.*

Proof. Use, e.g., [Sh:c, V,4.11,p.272], assume $\frac{\bar{b}}{\bar{a}} \pm p$; we can define inductively equivalence relations E_n , with parameters from $\text{acl}(\bar{a}^\ell)$, $\bar{a}^\ell = \bar{a} \hat{=} (\bar{b}/E_0) \hat{=} \dots \hat{=} (\bar{b}/E_{n-1})$, such that $\text{tp}(\bar{b}/E_n, \text{acl}(\bar{a}^n))$ is semi-regular. By superstability this stop for some n hence $\bar{b} \subseteq \text{acl}(\bar{a}^n)$. For some first m $\text{tp}(\bar{b}/E_m, \text{acl}(\bar{a}^n))$ is $\pm p$, by [Sh:c, X,7.3](5),p.552 the type is regular (as because p is trivial having depth > 0 ; see [Sh:c, X,7.2,p.551]).

□_{2.3}

2.4 Definition. We define by induction on an ordinal α the following (simultaneously): note — if a definition of something depends on another which is not well defined, neither is the something)

$$\begin{aligned} \text{tp}_\alpha^1 \left[\begin{pmatrix} p \\ \bar{a} \end{pmatrix}, M \right] & \quad \text{for} \quad \begin{pmatrix} p \\ \bar{a} \end{pmatrix} \in \Gamma_1, \bar{a} \subseteq M \\ \text{tp}_\alpha^2 \left[\begin{pmatrix} p, r \\ \bar{a} \end{pmatrix}, M \right] & \quad \text{for} \quad \begin{pmatrix} p, r \\ \bar{a} \end{pmatrix} \in \Gamma_1, \bar{a} \subseteq M \\ \text{tp}_\alpha^3 \left[\begin{pmatrix} \bar{b} \\ \bar{a} \end{pmatrix}, M \right] & \quad \text{for} \quad \begin{pmatrix} \bar{b} \\ \bar{a} \end{pmatrix} \in \Gamma_3^c, \bar{a} \subseteq \bar{b} \subseteq M \end{aligned}$$

Case A $\alpha = 0$: $\text{tp}_\alpha^1 \left[\begin{pmatrix} p \\ \bar{a} \end{pmatrix}, M \right]$ is $\text{tp}((c, \bar{a}), \emptyset)$ for any c realizing p .

$\text{tp}_\alpha^2 \left[\begin{pmatrix} p, r \\ \bar{a} \end{pmatrix}, M \right]$ is $\text{tp}((c, \bar{b}, \bar{a}), \emptyset)$ for any (c, \bar{b}) realizing r

$\text{tp}_\alpha^3 \left[\begin{pmatrix} \bar{b} \\ \bar{a} \end{pmatrix}, M \right]$ is $\text{tp}((\bar{b}, \bar{a}), \emptyset)$

(i.e., the type and the division of the variables between the sequences).

Case B $\alpha = \beta + 1$:

(a) $\text{tp}_\alpha^1 \left[\begin{pmatrix} p \\ \bar{a} \end{pmatrix}, M \right]$ is:

Subcase a1: if p has depth zero, it is $w_p(M/\bar{a})$ (the p -weight, equivalently, the dimension)

Subcase a2: if p has depth > 0 (hence is trivial), then it is $\{ \langle y, \lambda_{\bar{a}, p}^y \rangle : y \}$ where

$$\begin{aligned} \lambda_{\bar{a}, p}^y &= \dim(\mathbf{I}_{\bar{a}, p}^y[M], a) \text{ where } \mathbf{I}_{\bar{a}, p}^y[M] = \\ & \left\{ c \in M : c \text{ realize } p \text{ and } y = \text{tp}_\beta^3 \left[\begin{pmatrix} \text{acl}(\bar{a} + c) \\ \text{acl}(\bar{a}) \end{pmatrix}, M \right] \right. \\ & \quad \left. \text{where } \bar{a}^* \text{ list } \text{acl}(\bar{a}) \text{ and } \bar{c}^* \text{ list } \text{acl}(\bar{a} + c) \right\} \end{aligned}$$

an alternative probably more transparent and simpler in use is:

$$\lambda_{\bar{a},p}^y = \dim \left\{ c \in M : c \text{ realizes } p \text{ and} \right.$$

$$y = \left\{ \text{tp}_\beta^3 \left[\begin{pmatrix} \text{acl}(\bar{a} + c') \\ \text{acl}(\bar{a}) \end{pmatrix}, M \right] : c' \in p(M) \text{ and } c' \upharpoonright_{\bar{a}} \perp c \right\}$$

pedantically $y = \left\{ \text{tp}_\beta^3 \left(\begin{matrix} \langle c' \rangle \\ \bar{a} \hat{\ } \bar{a}^* \end{matrix}, M \right) : \bar{a}^* \text{ list } \text{acl}(\bar{a}) \text{ and} \right.$

$$\left. \bar{c}^* \text{ list } \text{acl}(\bar{a} + c'), c' \in p(M) \text{ and } c' \upharpoonright_{\bar{a}} \perp c \right\}.$$

(b) $\text{tp}_\alpha^2 \left[\begin{pmatrix} p,r \\ \bar{a} \end{pmatrix}, M \right]$ is:

$\text{tp}_\alpha^1 \left[\begin{pmatrix} c/\bar{b}^+ \\ \bar{b}^+ \end{pmatrix}, M \right]$ for any (c, \bar{b}) realizing r , $\bar{b}^+ = \text{acl}(\bar{a} + \bar{b})$, i.e., \bar{b}^+ lists $\text{acl}(\bar{a} + \bar{b})$ (so not well defined if we get at least two different cases; so remember $c/\bar{b}^+ \in S(\bar{b}^+)$).

(c) $\text{tp}_\alpha^3 \left[\begin{pmatrix} \bar{b} \\ \bar{a} \end{pmatrix}, M \right]$ is $\left\{ \langle p, \text{tp}_\alpha^2 \left[\begin{pmatrix} p,r \\ \bar{b} \end{pmatrix}, M \right] \rangle : \begin{pmatrix} p,r \\ \bar{b} \end{pmatrix} \in \Gamma_2^s \text{ and } p \perp \bar{a} \right\}$.

Case C: α limit: For any $\ell \in \{1, 2, 3\}$ and suitable object OB :

$$\text{tp}_\alpha^\ell[OB, M] = \langle \text{tp}_\beta^\ell[OB, M] : \beta < \alpha \rangle.$$

2.5 Definition. 1) For $\begin{pmatrix} p \\ \bar{a} \end{pmatrix} \in \Gamma_1$ where $\bar{a} \in M$, let (remembering 1.14(8)):

$$\mathcal{P}_{\begin{pmatrix} p \\ \bar{a} \end{pmatrix}}^M = \left\{ q \in S(M) : q \text{ regular and } : q \pm p \text{ or for some} \right.$$

$$\left. c \in p(M) \text{ we have } q \in \mathcal{P}_{\begin{pmatrix} c \\ \bar{a} \end{pmatrix}}^M \right\}.$$

2) For $\begin{pmatrix} p,r \\ \bar{a} \end{pmatrix} \in \Gamma_2$ let

$$\mathcal{P}_{\begin{pmatrix} p,r \\ \bar{a} \end{pmatrix}}^M = \left\{ q \in S(M) : q \text{ regular and } : q \pm p \text{ or for some} \right.$$

$$\left. (c, \bar{b}) \in r(M), q \in \mathcal{P}_{\begin{pmatrix} c \\ \bar{a} + \bar{b} \end{pmatrix}}^M \right\}.$$

3) For a set \mathcal{P} of (stationary) regular types not orthogonal to M_1 , let $M_1 \leq_{\mathcal{P}} M_2$ means $M_1 \prec M_2$ and for every $p \in \mathcal{P}$ and $\bar{c} \in M_2$, $\frac{\bar{c}}{M_1} \perp p$.

4) If (in (3)), $\mathcal{P} = \mathcal{P}_{\left(\frac{p}{\bar{a}}\right)}^{M_1}$ we may write $\left(\frac{p}{\bar{a}}\right)$ instead \mathcal{P} , similarly if $\mathcal{P} = \mathcal{P}_{\left(\frac{p,r}{\bar{a}}\right)}^{M_1}$ we may write $\left(\frac{p,r}{\bar{a}}\right)$.

2.6 Claim. :

1) From $\text{tp}_{\alpha}^1\left[\left(\frac{p}{\bar{a}}\right), M\right]$ we can compute $\text{tp}_{\infty}^1\left[\left(\frac{p}{\bar{a}}\right), M\right]$ if $\text{Dp}(p) < \alpha$.

2) From $\text{tp}_{\alpha}^2\left[\left(\frac{p,q}{\bar{a}}\right), M\right]$ we can compute $\text{tp}_{\infty}^2\left[\left(\frac{p,q}{\bar{a}}\right), M\right]$ if $\text{Dp}(p) < \alpha$.

3) From $\text{tp}_{\alpha}^3\left[\left(\frac{\bar{b}}{\bar{a}}\right), M\right]$ we can compute $\text{tp}_{\infty}^3\left[\left(\frac{\bar{b}}{\bar{a}}\right), M\right]$ if $\text{Dp}(\bar{b}/\bar{a}) < \alpha$.

4) If Definition 2.5(2) we can replace “some $(c, \bar{b}) \in r(M)$ ” by “every $(c, \bar{b}) \in r(M)$ ”.

Proof. 1),2),3) We prove this by induction on α . By the definition.

4) Left to the reader.

2.7 Observation. From $\text{tp}_{\alpha}^{\ell}(OB, M)$ we can compute $\text{tp}_{\beta}^{\ell}[OB, M]$ and $\text{tp}_{\beta}^{\ell}[OB, M]$ is well defined if $\beta \leq \alpha$ and the former is well defined.

2.8 Lemma. For every ordinal α the following holds:

- (1) tp_{α}^1 is well defined⁵
- (2) tp_{α}^2 is well defined.
- (3) tp_{α}^3 is well defined.
- (4) If $\bar{a} \in M_1$, $\left(\frac{p}{\bar{a}}\right) \in \Gamma_1$, $M_1 \leq_{\left(\frac{p}{\bar{a}}\right)} M_2$ then
 $\text{tp}_{\alpha}^1\left[\left(\frac{p}{\bar{a}}\right), M_1\right] = \text{tp}_{\alpha}^1\left[\left(\frac{p}{\bar{a}}\right), M_2\right]$.
- (5) If $\bar{a} \in M_1$, $\left(\frac{p,r}{\bar{a}}\right) \in \Gamma_2^s$, $M_1 \leq_{\left(\frac{p}{\bar{a}}\right)} M_2$ then
 $\text{tp}_{\alpha}^2\left[\left(\frac{p,r}{\bar{a}}\right), M_1\right] = \text{tp}_{\alpha}^2\left[\left(\frac{p,r}{\bar{a}}\right), M_2\right]$.
- (6) If $\bar{a} \subseteq \bar{b} \subseteq M_1$, $\left(\frac{\bar{b}}{\bar{a}}\right) \in \Gamma_3^c$, $M_1 \leq_{\left(\frac{\bar{b}}{\bar{a}}\right)} M_2$ then
 $\text{tp}_{\alpha}^3\left[\left(\frac{\bar{b}}{\bar{a}}\right), M_1\right] = \text{tp}_{\alpha}^3\left[\left(\frac{\bar{b}}{\bar{a}}\right), M_2\right]$.

⁵i.e. in all the cases we have tried to define it in Definition 2.9

Proof. We prove it, by induction on α , simultaneously (for all clauses and parameters).

If α is zero, they hold trivially by the definition.

If α is limit, they hold trivially by the definition and induction hypothesis. So for the rest of the proof let $\alpha = \beta + 1$.

Proof of (1) $_\alpha$. If p has depth zero — check directly.

If p has depth > 0 - by (3) $_\beta$

(i.e. induction hypothesis) no problem.

Proof of (2) $_\alpha$. Like 1.27 (and (4) $_\alpha$).

Proof of (3) $_\alpha$. Like (2) $_\alpha$.

Proof of (4) $_\alpha$. Like 1.26 (and (3) $_\beta$, (6) $_\beta$).

Proof of (5) $_\alpha$. By (2) $_\alpha$ we can look only at (c, \bar{b}^+) in M_1 , then use (4) $_\alpha$.

Proof of (6) $_\alpha$. By (5) $_\alpha$.

□_{2.8}

2.9 Lemma. *For an ordinal α restricting ourselves to the cases (the types p, p_1 being) of depth $< \alpha$:*

(A1) Assume $(\frac{p}{\bar{a}}) \in \Gamma_1, \bar{a} \subseteq \bar{a}_1 \subseteq M, \bar{a}_1$ is ϵ -finite, $\frac{\bar{a}_1}{\bar{a}} \perp p$ and p_1 is the stationarization of p over \bar{a}_1 .

Then from $\text{tp}_\alpha^1[(\frac{p}{\bar{a}}), M]$ we can compute $\text{tp}_\alpha^1[(\frac{p_1}{\bar{a}_1}), M]$.

(A2) Under the assumption of (A1) also the inverse computations are O.K.

(A3) Assume $(\frac{p_\ell}{\bar{a}}) \in \Gamma_1$ for $\ell = 1, 2, \bar{a} \subseteq M$ and $p_1 \pm p_2$.

Then from $\text{tp}_\alpha^1[(\frac{p_1}{\bar{a}}), M]$ (and $\text{tp}((\bar{a}, c_1, c_2), \emptyset)$ where c_1, c_2 realizes p_1, p_2 respectively, of course) we can compute $\text{tp}_\alpha^1[(\frac{p_2}{\bar{a}}), M]$.

(B1) Assume $(\frac{p_\ell, r_\ell}{\bar{a}}) \in \Gamma_2^{sc}$ for $\ell = 1, 2, \bar{a} \in M$ and $p_1 \pm p_2$.

Then (from the first order information on $\bar{a}, p_1, p_2, r_1, r_2$, of course, and) $\text{tp}_\alpha^2[(\frac{p_1, r_1}{\bar{a}_1}), M]$ we can compute $\text{tp}_\alpha^2[(\frac{p_2, r_2}{\bar{a}}), M]$.

(B2) Assume $\bar{a} \subseteq \bar{a}_1 \subseteq M, \frac{\bar{a}_1}{\bar{a}} \perp p, (\frac{p, r}{\bar{a}}) \in \Gamma_2^{cs}, r \subseteq r_1 \in S(\bar{a}_1), r_1$ does not fork over $\bar{a},$ (so $(\frac{p, r_1}{\bar{a}_1}) \in \Gamma_2$).

Then from $\text{tp}_\alpha^2[(\frac{p, r_1}{\bar{a}_1}), M]$ we can compute $\text{tp}_\alpha^2[(\frac{p, r_2}{\bar{a}}), M]$.

(B3) Under the assumption of (B2), the inverse computation are O.K.

(C1) Assume $(\bar{b}) \in \Gamma_3^c, \bar{a} \subseteq \bar{b} \subseteq M, \bar{a} \subseteq \bar{a}_1, \bar{b} \underset{\bar{a}}{\cup} \bar{a}_1, \bar{b}_1 = \text{acl}(\bar{a}_1 + \bar{b})$.

Then from $\text{tp}_\alpha^3[(\bar{b}), M]$ we can compute $\text{tp}_\alpha^3[(\bar{b}_1), M]$.

(C2) Under the assumptions of (C1) the inverse computation is O.K.

(C3) Assume $(\bar{b}) \in \Gamma_3, \bar{b} \subseteq b^*, \bar{b}^* \perp_a \bar{a}, \bar{b}^* = \text{acl}(\bar{b}^*)$. Then from $\text{tp}_\alpha^3[(\bar{b}), M]$ we can compute

$$\left\{ \text{tp}_\alpha^3[(\bar{b}'), M] : \bar{b} \subseteq b' \subseteq M \text{ and } \frac{\bar{b}'}{\bar{b}} = \frac{\bar{b}^*}{b^*} \right\}.$$

Proof. We prove it, simultaneously, for all clauses and parameters, by induction on α and the order of the clauses.

For $\alpha = 0$: easy.

For α limit: very easy.

So assume $\alpha = \beta + 1$.

Proof of (A1) $_\alpha$. As p is stationary $\perp \frac{\bar{a}_1}{\bar{a}}$, for every $c \in p(M), \frac{c}{\bar{a}} \vdash \frac{c}{\bar{a}_1}$, which necessarily is p_1 , hence $p(M) = p_1(M)$. Also the dependency relation on $p(M)$ is the same over \bar{a}_1 , hence dimension. So it suffices to show:

$$(*) \text{ for } c \in p(M), \text{ from } \text{tp}_\beta^3\left[\left(\begin{smallmatrix} \text{acl}(\bar{a}+c) \\ \text{acl } \bar{a} \end{smallmatrix}\right), M\right] \text{ we can compute } \text{tp}_\beta^3\left[\left(\begin{smallmatrix} \text{acl}(\bar{a}_1+c) \\ \text{acl } \bar{a}_1 \end{smallmatrix}\right), M\right].$$

But this holds by (C1) $_\beta$.

Proof of (A2) $_\alpha$. Similar using (C2) $_\beta$.

Proof of (A3) $_\alpha$. If p_1 (equivalently p_2) has depth zero — the dimensions are equal. Assume they have depth > 0 hence are trivial and dependency over \bar{a} is an equivalence relation on $p_1(M) \cup p_2(M)$.

Now for $c_1 \in p_1(M)$, from $\text{tp}_\beta^3\left[\left(\begin{smallmatrix} \text{acl}(a+c_1) \\ \text{acl}(\bar{a}) \end{smallmatrix}\right), M\right]$ we can compute for every complete type over $\text{acl}(\bar{a} + c_1)$ not forking over \bar{a} , and \bar{d} realizing r , $\text{tp}_\beta^3\left[\left(\begin{smallmatrix} \text{acl}(\bar{a}+\bar{d}+c_1) \\ \text{acl}(\bar{a}+\bar{d}) \end{smallmatrix}\right), M\right]$ - by (C1) $_\beta$, then we can compute for each such r, \bar{d} ,

$$\left\{ \text{tp}_\beta^3 \left[\left(\begin{array}{c} \text{acl}(\bar{a} + \bar{d} + c_2) \\ \text{acl}(\bar{a} + \bar{d}) \end{array} \right), M \right] : c_2 \in p_2(M) \text{ and } \frac{c_2}{\text{acl}(\bar{a} + \bar{d} + c_1)} \perp_a (\bar{a} + \bar{d}) \right. \\ \left. (\text{necessarily } c_2 \upharpoonright_{\bar{a}} \bar{d}) \right\}$$

(this by $(C3)_\beta$).

Proof of $(B1)_\alpha$. As in earlier cases we can restrict ourselves to the case $\text{Dp}(p_\ell) > 0$. We can find $(c_\ell, \bar{b}_\ell) \in r_\ell(M)$, $\bar{b}_1 \upharpoonright_{\bar{a}} \bar{b}_2, c_1 \bar{b}_1 \upharpoonright_{\bar{a}} \bar{b}_2$ (by [Sh:c, X,7.3](6)). By 2.8(2) (and the definition) from $\text{tp}_\alpha^2 \left[\left(\begin{array}{c} p_1, r_1 \\ \bar{a} \end{array} \right), M \right]$ we can compute that it is equal to $\text{tp}_\alpha^1 \left[\left(\begin{array}{c} c_1 / \text{acl}(\bar{a} + \bar{b}_1) \\ \text{acl}(\bar{a} + \bar{b}_1) \end{array} \right), M \right]$.

By $(A1)_\alpha$ we can compute $\text{tp}_\alpha^1 \left[\left(\begin{array}{c} c_1 / \text{acl}(\bar{a} + \bar{b}_1 + \bar{b}_2) \\ \text{acl}(\bar{a} + \bar{b}_1 + \bar{b}_2) \end{array} \right), M \right]$ hence by $(A3)_\alpha$ we can compute $\text{tp}_\alpha^1 \left[\left(\begin{array}{c} c_2 / \text{acl}(\bar{a} + \bar{b}_1 + \bar{b}_2) \\ \text{acl}(\bar{a} + \bar{b}_1 + \bar{b}_2) \end{array} \right), M \right]$.

Now use $(A2)_\alpha$ to compute $\text{tp}_\alpha^1 \left[\left(\begin{array}{c} c_2 / \text{acl}(\bar{a} + \bar{b}_2) \\ \text{acl}(\bar{a} + \bar{b}_2) \end{array} \right), M \right]$ and by 2.8(2), 2.4(2) it is equal to $\text{tp}_\alpha^2 \left[\left(\begin{array}{c} p, r \\ \bar{a} \end{array} \right), M \right]$.

Proof of $(B2)_\alpha$. Choose $(c, \bar{b}) \in r(M)$ such that $c\bar{b} \upharpoonright_{\bar{a}} \bar{a}_1$.

From $\text{tp}_\alpha^2 \left[\left(\begin{array}{c} p, r_1 \\ \bar{a} \end{array} \right), M \right]$ we can compute $\text{tp}_\alpha^1 \left[\left(\begin{array}{c} c / (\bar{a} + \bar{b}) \\ \bar{a} + \bar{b} \end{array} \right), M \right]$ (just — see 2.8(2) and Definition 2.4), from it we can compute $\text{tp}_\alpha^1 \left[\left(\begin{array}{c} c / (\bar{a} + \bar{b} + \bar{a}_1) \\ \bar{a} + \bar{b} + \bar{a}_1 \end{array} \right), M \right]$ (by $(A1)_\alpha$), from it we can compute $\text{tp}_\alpha^2 \left[\left(\begin{array}{c} p, r_2 \\ \bar{a}_2 \end{array} \right), M \right]$ (see 2.8(2) and Definition 2.4).

Proof of $(B3)_\alpha$. Let $\left(\begin{array}{c} p, r \\ \bar{b}_1 \end{array} \right) \in \Gamma_r^s, p \perp \bar{a}_1$ be given. So necessarily $\frac{\bar{a}_1}{\bar{a}} \pm p$ (this to enable us to use $(B2, 3)$). It suffices to compute $\text{tp}_\alpha^2 \left[\left(\begin{array}{c} p, r \\ \bar{b}_1 \end{array} \right), M \right]$ and we can discard the case $\text{Dp}(p) = 0$.

So p is regular $\pm \bar{b}_1, \perp \bar{a}_1$, hence $p \pm \bar{b}, p \perp \bar{a}$, and as $\bar{a} \subseteq \bar{b}, \bar{b} = \text{acl}(\bar{b})$ we can find $r, \left(\begin{array}{c} p, r_1 \\ \bar{b} \end{array} \right) \in \Gamma_2$, (see 2.3) and we know $\text{tp}_\alpha^2 \left[\left(\begin{array}{c} p, r_1 \\ \bar{b} \end{array} \right), M \right]$, and we can find r_2 , a complete type over \bar{b}_1 extending r_1 which does not fork over \bar{b}_1 . From $\text{tp}_\alpha^2 \left[\left(\begin{array}{c} p, r_1 \\ \bar{b} \end{array} \right), M \right]$ we can compute $\text{tp}_\alpha^2 \left[\left(\begin{array}{c} p, r_2 \\ \bar{b}_1 \end{array} \right), M \right]$ by $(B2)_\alpha$, and from it $\text{tp}_\alpha^2 \left[\left(\begin{array}{c} p, r \\ \bar{b}_1 \end{array} \right), M \right]$ by $(B1)_\alpha$.

Proof of $(C2)_\alpha$. Similar, use $(B3)_\alpha$ instead of $(B2)_\alpha$.

Proof of (C3) $_{\alpha}$. Without loss of generality $\frac{\bar{b}^*}{\bar{b}}$ is semi regular, let p^* be a regular type not orthogonal to it and without loss of generality $\text{Dp}(p^*) > 0 \Rightarrow \frac{\bar{b}^*}{\bar{b}}$ regular (as in 2.3).

If p^* has depth zero, then the only p appearing in the definition $\text{tp}_{\alpha}^3([\frac{\bar{b}}{\bar{a}}], M)$ is p^* (up to \pm) and this is easy. Then tp_{α}^2 is just the dimension and we have no problem.

So assume p^* has depth > 0 . We can by $(B1)_{\alpha}, (B2)_{\alpha}$ compute $\text{tp}_{\alpha}^2([\frac{p', q'}{\bar{b}^*}], M)$ when $p' \pm \bar{b}, p' \pm p^*$ (regardless of the choice of \bar{b}^*). Next assume $p' \pm p^*$; by $(B1)_{\alpha}$ without loss of generality q' does not fork over \bar{b} . As $\text{Dp}(p^*) > 0$, it is trivial (and we assume $w_p(\bar{b}^*, \bar{b}) = 1$) hence \bar{b}^*/\bar{b} is regular so in $\text{tp}_{\alpha}^2([\frac{p', q'}{\bar{b}^*}], M)$ we just lose a weight 1 for one specific tp_{β}^3 type: the one \bar{b}^* realizes concerning which we have a free choice. We are left with the cases $p' \pm \bar{b}, p' \pm p^*$; well we know tp_{β}^3 but we have to add tp_{α}^3 ? Use Claim 2.6(3) (and $(A1)_{\alpha}$ as we add a parameter). $\square_{2.9}$

2.10 Claim. $\text{tp}_{\gamma}^3([\frac{\bar{b}}{\bar{a}}], M), \text{tp}_{\gamma}^3[\bar{a}, M], \text{tp}_{\gamma}^3[M]$ are expressible by formulas in $\mathbb{L}_{\infty, \aleph_{\epsilon}}^{\gamma}(\text{d.q.})$.

By 2.9 we have

2.11 Conclusion. If $\text{Dp}(T) < \infty$ then:

- 1) From $\text{tp}_{\infty}^3([\frac{B}{A}], M)$ we can compute $\text{tp}_{\infty}([\frac{B}{A}], M)$ (the type from §1).
- 2) Similarly from $\text{tp}_{\infty}^3[A, M]$ we can compute $\text{tp}_{\infty}[(A), M]$.

From 2.6, 2.10, 2.11 and 1.30 we get

2.12 Corollary. *If $\gamma = \text{Dp}(T)$ and M, N are \aleph_{ϵ} -saturated, then*

$$M \cong N \Leftrightarrow \text{tp}_{\gamma}^3[M] = \text{tp}_{\gamma}^3[N] \Leftrightarrow M \cong_{\mathbb{L}_{\infty, \aleph_{\epsilon}}^{\gamma}(\text{d.q.})} N.$$

APPENDIX

The following clarifies several issues raised by Baldwin. A consequence of

- ⊗ the existence of nice invariants for characterization up to isomorphism (or characterization of the models up to isomorphism by their \mathcal{L} -theory for suitable logic \mathcal{L})

naturally give absoluteness, e.g. extending the universe say by nice forcing preserve non-isomorphism. So negative results for

- (*) is non-isomorphism (of models of T) preserved by forcing by “nice forcing notions”?

implies that we cannot characterize models up to isomorphism by their \mathcal{L} -theory when the logic \mathcal{L} is “nice”, i.e. when $Th_{\mathcal{L}}(M)$ preserved by nice forcing notions. So coding a stationary set by the isomorphism type can be interpreted as strong evidence of “no nice invariants”, see [Sh 220]. Baldwin, Laskowski, Shelah [BLSh 464] show that not only for every unsuperstable; but also for some quite trivial superstable (with NDOP, NOTOP) countable T , there are non-isomorphic models which can be made isomorphic by some ccc (even σ -centered) forcing notion. This shows that the lack of a really finite characterization is serious.

Can we still get from the characterization in this paper an absoluteness result? Note that for preserving \aleph_ϵ -saturation (for simplicity for models of countable T) we need to add no reals⁶, and in order not to erase distinction of dimensions we want not to collapse cardinals, so the following questions is natural, for a first order (countable) complete T :

- (*) $_T$ assume $\mathbf{V}_1 \subseteq \mathbf{V}_2$ are transitive models of ZFC with the same cardinals and reals, the theory $T \in \mathbf{V}_1$. If the models M_1, M_2 are from \mathbf{V}_1 and they are models of T not isomorphic in \mathbf{V}_1 ; must they still be not isomorphic in V_2 ⁷
- (*) $_{T,\kappa}$ like (*) $_T$ we assume in addition $\mathcal{P}(\kappa)^{\mathbf{V}_1} = \mathcal{P}(\kappa)^{\mathbf{V}_2}$.

2.13 Theorem. 1) For countable first order complete T the answer to (*) $_T$ and (*) $_{T,\kappa}$ for any κ is negative except when possibly T is superstable, NDOP, NOTOP. 2) For any first order complete T for the class of \aleph_ϵ -saturated models, the answer to (*) $_{T,|T|}$ is negative except possible when T is superstable with NDOP.

⁶(the set of $\{acl(\bar{a}) : \bar{a} \in {}^\omega M\}$ is absolute but the set of their enumeration and of the $\{f \upharpoonright (acl(\bar{a})) : f \in \text{AUT}(\mathfrak{C}), f(\bar{a}) = \bar{a}\}$ is not).

⁷Note we did not say they have the same ω -sequences of ordinals, e.g. if $V_2 = V_1^P$, P Prikry forcing, then the assumption of (*) $_T$ holds though a new ω -sequence of ordinals was added. So for $V_1 \subseteq V_2$ as in (*) $_T$, the $\mathcal{L}_{\infty, \aleph_1}$ -theory is not necessarily preserved.

Proof. By quoting.

So we restrict ourselves to these. It should be quite transparent that $\mathbb{L}_{\infty, \aleph_\epsilon}(q.d.)$ -theory is preserved from \mathbf{V}_1 to \mathbf{V}_2 (as well as the set of sentences in the logic).

Hence

2.14 Theorem. *For the class of \aleph_ϵ -saturated models of superstable NDOP, NO-TOP theory T the answer to $(*)_{T, |T|}$ is yes.*

Proof. In \mathbf{V}_1 for $\ell = 1, 2$ let $\mathcal{A}_\ell = \{A \subseteq M_\ell^{\text{eq}} : A \text{ is } \epsilon\text{-finite and } \text{acl}(A) = A\}$. Clearly the same definition gives \mathcal{A}_ℓ in \mathbf{V}_2 and M_1, M_2 are \aleph_ϵ -saturated also in \mathbf{V}_2 .

$A \in \mathcal{A}_\ell \Rightarrow |A| \leq |T|$ and let $\mathcal{F}^* = \{f : \text{for some } A_1 \in \mathcal{A}_2 \text{ and } A_2, f \text{ is a one-to-one function from } A_1 \text{ onto } A_2 \text{ which is an } (M_1, M_2)\text{-elementary mapping}\}$. Again this definition gives the same set. We can define a rank function $\text{rk} : \mathcal{F}^* \rightarrow \text{Ord} \cup \{\infty\}$ such that $\text{rk}(f) = \infty$ iff $(M_1, a)_{a \in \text{Dom}(f)} \equiv_{\mathbb{L}_{\infty, \aleph_\epsilon}, (d.q.)} (M_2, f(a))_{a \in \text{Dom}(f)}$ and it too is absolute.

Easily in both universes

$$(a) \quad M_1 \cong M_2 \text{ iff } M_1 \equiv_{\mathbb{L}_{\infty, \aleph_\epsilon}, (d.q.)} M_2.$$

[Why? By Theorem 1.5.]

$$(b) \quad M_2 \equiv_{\mathbb{L}_{\infty, \aleph_0}, (d.q.)} M_2 \text{ iff there is } \mathcal{F} \subseteq \mathcal{F}^* \text{ as in } \otimes_{M_1, M_2} \text{ from 1.4.}$$

[Why? By 1.4.]

$$(c) \quad \text{there is } \mathcal{F} \subseteq \mathcal{F}^* \text{ as in } \otimes_{M_1, M_2} \text{ from 1.4 iff } \infty \text{ belongs to the range of the rank function.}$$

[Why? Well known.]

REFERENCES.

- [BLSh 464] John T. Baldwin, Michael C. Laskowski, and Saharon Shelah. Forcing Isomorphism. *Journal of Symbolic Logic*, **58**:1291–1301, 1993.
- [BeSh 307] Steven Buechler and Saharon Shelah. On the existence of regular types. *Annals of Pure and Applied Logic*, **45**:277–308, 1989.
- [Sh 220] Saharon Shelah. Existence of many $L_{\infty, \lambda}$ -equivalent, nonisomorphic models of T of power λ . *Annals of Pure and Applied Logic*, **34**:291–310, 1987. Proceedings of the Model Theory Conference, Trento, June 1986.
- [Sh 225] Saharon Shelah. On the number of strongly \aleph_ϵ -saturated models of power λ . *Annals of Pure and Applied Logic*, **36**:279–287, 1987. See also [Sh:225a].
- [Sh:c] Saharon Shelah. *Classification theory and the number of nonisomorphic models*, volume 92 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, xxxiv+705 pp, 1990.