

PARTITION THEOREMS FOR EXPANDED TREES
1176

SAHARON SHELAH

ABSTRACT. We look for partition theorems for large subtrees for suitable uncountable trees and colourings.

We concentrate on sub-trees of $\kappa^{\geq 2}$ expanded by a well ordering of each level. Unlike earlier works, we do not ask the embedding to preserve the height of the tree but the equality of levels is preserved. We get consistency results without large cardinals.

The intention is to apply it to model theoretic problems.

Date: November 8, 2022.

2010 Mathematics Subject Classification. Primary: 03E02; Secondary: 03E35 .

Key words and phrases. Ramsey theory, partition theorems, uncountable trees.

The author thanks Alice Leonhardt for the beautiful typing up to 2019. In later versions, the author thanks typing services generously funded by an individual who wishes to remain anonymous.. We thank the ISF (Israel Science Foundation) grant 1838(19) for partially supporting this research. First typed November 14, 2018. References like [Shea, Th0.2=Ly5] means the label of Th.0.2 is y5. The reader should note that the version in my website is usually more updated than the one in the mathematical archive. Publication number 1176 in the author list of publications.

Annotated Content

§0 Introduction, pg. 3

§(0A) Background, pg. 3

§(0B) Preliminaries, (label z) pg. 4

§1 Partition Theorems, (label a), pg.10

§1A The definitions, pg.6

[We consider here partition theorems for trees. Here we deal with the case we waive the equality of cardinals as needed for the model theory application. This is used in §1B.

In 1.1 define \mathbf{T}

In 1.5 state the partition relation.]

§1B Forcing in ZFC, (label b) pg. 10

[We work on a partition theorem for trees, for the main case here, we get consistency without the large cardinal. Naturally the price is having to vary the size of the cardinals (parallel to the Erdős-Rado theorem).]

§ 0. INTRODUCTION

§ 0(A). **Background and Results.**

We continue two lines of research. One is set theoretic: pure partition relations on trees and the other is model theoretic: Hanf numbers and non-deniability of well ordering, in particular related to ω_1 . This is related to the existence of GEM (generalized Ehrenfuecht-Mostowski) for suitable templates (see [Sheb]), and applications to descriptive set theory.

Halpern-Levy [HL71] had proved a milestone theorem on independence of versions of the axiom of choice: in ZF, AC is strictly stronger than the maximal prime ideal theorem (i.e. every Boolean algebra has a maximal ideal).

This work isolated a partition theorem¹ on the tree ${}^\omega > 2$, necessary for the proof. This partition theorem was subsequently proved by Halpern-Lauchli [HL66] and was a major and early theorem in Ramsey theory, (so the proof above relies on it). See more in Laver [Lav71], [Lav73] and [She78b, AP,§2] and Milliken [Mil79], [Mil81].

The [HL66] proof uses induction, later Harrington found a different proof using forcing: adding many Cohen reals and a name of a (non-principal) ultrafilter on \mathbb{N} . Earlier, (on using adding many reals and a partition theorem) see Silver's proof of π_1^1 -equivalence relations, in [Sil80].

Now [She92, §4] turn to uncountable trees, i.e. for some $\kappa > \aleph_0$, we consider trees \mathcal{T} which are sub-trees of $({}^\kappa > 2, \triangleleft)$, such that (as in [HL66]) for every level $\varepsilon < \kappa$, either $(\forall \eta \in \mathcal{T} \cap {}^\varepsilon 2)(\eta \hat{\ } \langle 0 \rangle, \eta \hat{\ } \langle 1 \rangle \in \mathcal{T})$ or $(\forall \eta \in \mathcal{T} \cap {}^\varepsilon 2)(\exists ! \iota < 2)[\eta \hat{\ } \langle \iota \rangle \in \mathcal{T}]$; (and of course the first occurs unboundedly often). But a new point is that we have to use a well ordering of $\mathcal{T} \cap {}^\varepsilon 2$ for $\varepsilon < \kappa$.

Naturally we add “is closed enough (that is under unions of increasing sequences of length $< \kappa$)”. Also colouring with infinite number of colours, the proof uses “measurable κ which remains so when we add λ many κ -Cohens for appropriate λ ”; it generalizes Harrington's proof. This was continued in several works, see Dobrinen-Hathaway [DH17] and references there.

We are here mainly interested in a weaker version which is enough for the model theoretic applications we have in mind, we start with a large tree and get one of smaller cardinality, in a sense this is solving the “equations” $X / (\text{Erdős-Rado theorem}) = [\text{She92}] / (\text{the partition relation of a weakly compact cardinal}) = [\text{HL66}] / (\text{Ramsey theorem})$. On other consistent partition relation see Boney-Shelah [S⁺a], in preparation.

Turning to model theory see [She75], [She76] and Dzamonja-Shelah [DS04] where such indiscernibility is considered in a model theoretic context.

A central direction in model theory in the sixties were two cardinal theorems. For infinite cardinals $\mu > \lambda$, let $K_{\mu,\lambda}$ be the class of models M such that M is of cardinality μ and P^M of cardinality λ . The main problems were transfer, compactness and completeness. For connection to partition theorems, Morley's proof of [Vau65], the Vaught far apart two cardinal theorem used Erdős-Rado theorem; generally see [She71b], [She71a], [She78a] and the survey [DS79]. Jensen's celebrated gap n two

¹Using not splitting to 2 but other finite splitting make a minor difference; similarly here.

cardinal theorem solve those problems for e.g. (\aleph_n, \aleph_0) when $\mathbf{V} = \mathbf{L}$. But can we get a nice picture in different universes?

Note that by [She89], [Sheb], consistently we have GEM (generalized Ehrenfuecht-Mostowski) models for ordered graphs as index models, even omitting types.

On a different direction Douglas Ulrich has asked me on $(*)_n$ below (and told me it has descriptive set theoretic consequences, see [SU19]). We intend to prove (in the sequel [S⁺b]) that for $n < \omega$:

- $(*)_n$ consistently
- (a) if $\psi \in \mathbb{L}_{\aleph_1, \aleph_0}$ has a model M of cardinality \beth_{n+1} with $(P^M, <^M)$ having order type ω_1 then ψ has a model N of cardinality \beth_{n+1} and $(P^N, <^N)$ is not well ordered,
 - (b) moreover, it is enough that M will have cardinality $\aleph_\delta, \delta \geq \beth_n^{++}$,
 - (c) of course, preferably not using large cardinals.

This requires consistency of many cases of partition relations on trees and more complicated structures, analysing GEM models. Much earlier we have intended (mentioned in [She00, 1.15]) to prove the parallel for first order logic; and (\beth_n, \aleph_0) , using (many Cohen's indestructible) measurables $\kappa_1 < \dots < \kappa_n$ as in [She92, §4] and forcing by blowing 2^{\aleph_0} to κ_1 , 2^{κ_1} to κ_2 etc relying on [She92]; but have not carried out that.

In preparation are also solutions to the two cardinal problems above and

- $(*)$ (a) $\alpha_\bullet < \omega_1, \beth_{\alpha+1} = (\beth_\alpha)^{+\omega_1+1}$ for $\alpha < \alpha_\bullet$ and well ordering of ω_1 is not definable in $\{\text{EC}_\psi(\beth_{\alpha_\bullet}): \psi \in \mathbb{L}_{\aleph_1, \aleph_0}\}$ or at least,
- (b) as above but for $\beth_{\alpha+1} = \aleph_{\beth_\alpha^{++}}$,
 - (c) parallel results replacing \aleph_0 by μ .

Contrary to the a priori expectation no large cardinal is used.

In a sequel ([S⁺c]) we intend also to deal with other partition relations and with weakly compact cardinals.

We thank Shimoni Garti and Mark Poór for many helpful comments.

§ 0(B). Preliminaries.

Definition 0.1. If $\mu = \mu^{<\kappa}$ then “for a (μ, κ) -club of $u \subseteq X$ we have $\varphi(u)$ ” means that: for some χ such that $\mu, u \in \mathcal{H}(\chi)$ and e.g. $\beth_3(\mu + |u|) < \chi$ and some $x \in \mathcal{H}(\chi)$, if $x \in \mathcal{B} \prec (\mathcal{H}(\chi), \in)$, $\|\mathcal{B}\| = \mu$, $[\mathcal{B}]^{<\kappa} \subseteq \mathcal{B}$ and $\mu + 1 \subseteq \mathcal{B}$, then the set $u = \mathcal{B} \cap X$ satisfies $\varphi(u)$; there are other variants.

Definition 0.2. For κ regular (usually $\kappa = \kappa^{<\kappa}$) and an ordinal γ , the forcing $\mathbb{P} = \text{Cohen}(\kappa, \gamma)$ of adding γ many κ -Cohen reals is defined as follows:

- (A) $p \in \mathbb{P}$ iff:
 - (a) p is a function with domain from $[\gamma]^{<\kappa}$,
 - (b) if $\alpha \in \text{dom}(p)$ then $p(\alpha) \in {}^\kappa 2$,
- (B) $\mathbb{P} \models p \leq q$ iff:
 - (a) $p, q \in \mathbb{P}$,
 - (b) $\text{dom}(p) \subseteq \text{dom}(q)$,
 - (c) if $\alpha \in \text{dom}(p)$ then $p(\alpha) \leq q(\alpha)$.
- (C) for $\alpha < \gamma$ let $\eta_\alpha = \bigcup \{p(\alpha): p \in \mathbb{G}_{\mathbb{P}} \text{ satisfies } \alpha \in \text{dom}(p)\}$, so $\Vdash_{\mathbb{P}} \text{“}\eta_\alpha \in {}^\kappa 2\text{”}$,

(D) for $u \subseteq \gamma$ let $\mathbb{P}_u := \{p \in \mathbb{P} : \text{dom}(p) \subseteq u\}$, so $\mathbb{P}_u \triangleleft \mathbb{P}$ and $\bar{\eta}_u = \langle \eta_\alpha : \alpha \in u \rangle$ is generic for \mathbb{P}_u .

Notation 0.3. 1) We denote infinite cardinals by $\kappa, \lambda, \mu, \chi, \theta, \partial$, and σ denotes a possibly finite cardinal.

2) We denote ordinals by $\alpha, \beta, \gamma, \delta, \varepsilon, \zeta, \xi$ and sometimes i, j .

3) We denote natural numbers by k, ℓ, m, n and sometimes i, j .

4) Instead e.g. a_i we may write $a[i]$, particularly in sub-script; also $\kappa(+)$ means κ^+ .

§ 1. PARTITION THEOREMS

§ 1(A). **The definitions.**

Here we consider partition on trees. For uncountable trees we find the need to consider a well ordering of each level, still preserving equality of level. We may consider embeddings where levels are not preserved, see Dzamonja-Shelah [DS04] (in the web version). Also we may waive completeness of the tree but usually still like to have many branches.

We intend to deal with an intermediate ones (and with the weakly compact cardinal) in a sequel [S⁺c].

Definition 1.1. 1) Let \mathbf{T} be the class of structures \mathcal{T} such that:

- (a) $\mathcal{T} = (u, <_*, E, <, \cap, S, R_0, R_1) = (u_{\mathcal{T}}, <^*_{\mathcal{T}}, E_{\mathcal{T}}, <_{\mathcal{T}}, \cap_{\mathcal{T}}, S_{\mathcal{T}}, R_{\mathcal{T}}^0, R_{\mathcal{T}}^1)$ but we may write $s \in \mathcal{T}$ instead of $s \in u$,
- (b) $(u, <_*)$ is a well ordering, so linear, u non-empty,
- (c) $<_{\mathcal{T}}$ is a partial order included in $<_*$,
- (d) $(u, <_{\mathcal{T}})$ is a tree, i.e. if $t \in \mathcal{T}$ then $\{s : s <_{\mathcal{T}} t\}$ is linearly ordered by $<_{\mathcal{T}}$; and is even well ordered, the level of t is the order type of this set; the tree is with $\text{ht}(\mathcal{T})$ levels,
- (e) E is an equivalence relation on u , convex under $<_*$,
- (f) (α) each E -equivalence class is the set of $t \in \mathcal{T}$ of level ε for some ε , so the set of E -equivalence classes is naturally well ordered,
 - (β) we denote the ε -th equivalence class by $\mathcal{T}_{[\varepsilon]}$,
 - (γ) E has no last E -equivalence class if not said otherwise,
 - (δ) let $\text{lev}_{\mathcal{T}}(s) = \text{lev}(s, \mathcal{T})$ be ε when $s \in \mathcal{T}_{[\varepsilon]}$, equivalently $\{t : t <_{\mathcal{T}} s\}$ has order type ε under the order $<_{\mathcal{T}}$,
 - (ε) so $\text{ht}(\mathcal{T})$ is $\bigcup\{\text{lev}(s) + 1 : s \in \mathcal{T}\}$.
- (g) if $s \in u, \text{lev}_{\mathcal{T}}(s) < \zeta < \text{ht}(\mathcal{T})$ then there is $t \in \mathcal{T}_{[\zeta]}$ which is $<_{\mathcal{T}}$ -above s ,
- (h) each $s \in \mathcal{T}$ has exactly two immediate successors by $<_{\mathcal{T}}$,
- (i) for $s \in \mathcal{T}$ we let:
 - ₁ $\mathcal{T}_{\geq s} = \{t \in \mathcal{T} : s \leq_{\mathcal{T}} t\}$,
 - ₂ $\text{succ}_{\mathcal{T}}(s) = \{t : t \in \mathcal{T}_{[\text{lev}(s)+1]} \text{ satisfies } s <_{\mathcal{T}} t\}$,
- (j) let $s = t|\varepsilon$ mean that $\text{lev}_{\mathcal{T}}(s) = \varepsilon \leq \text{lev}_{\mathcal{T}}(t) \wedge (s \leq_{\mathcal{T}} t)$,
- (k) for $t_1, t_2 \in \mathcal{T}, t_1 \cap_{\mathcal{T}} t_2$ is the maximal common lower bound of t_1, t_2 so we demand it always exists, i.e. $(\mathcal{T}, <)$ is normal,
- (l) for $\ell = 0, 1$ we have $R_{\ell} \subseteq \{(s, t) : s \in \mathcal{T} \text{ and } s <_{\mathcal{T}} t\}$ and if $s \in \mathcal{T}$ then for some $t_0 \neq t_1$ we have $\text{succ}_{\mathcal{T}}(s) = \{t_0, t_1\}$ and $\ell < 2 \Rightarrow (\forall t)(s R_{\ell} t \text{ iff } t_{\ell} \leq_{\mathcal{T}} t)$; so $s R_{\ell} t$ is the parallel to $\eta^{\wedge}(\ell) \trianglelefteq \nu$; we may think of $\{t : s R_{\ell} t\}$ as a division to the left side and the right side of the set of the t 's above s .

1A) For $\mathcal{T} \in \mathbf{T}, <_{\text{lex}} := <_{\mathcal{T}}^{\text{lex}}$ is the lexicographic order, e.g.,

$$\eta <_{\text{lex}} \nu \text{ iff } (\exists \rho)(\rho R_0 \nu \wedge \eta R_1 \rho) \text{ or } (\eta < \nu \wedge \eta R_1 \nu) \text{ or } (\nu < \eta \wedge \nu R_0 \eta).$$

2) Let $\mathbf{T}_{\theta, \kappa} = \{\mathcal{T} \in \mathbf{T} : \text{the tree } \mathcal{T} \text{ has } \delta \text{ levels, for some ordinal } \delta \text{ of cofinality } \kappa \text{ and for every } \varepsilon < \delta \text{ we have } \theta > |\{s \in \mathcal{T} : s \text{ of level } \leq \varepsilon\}|\}$.

3) Let $\mathcal{T}_1 \subseteq \mathcal{T}_2$ mean:

- (a) $\mathcal{T}_1, \mathcal{T}_2 \in \mathbf{T}$,
- (b) $<_{\mathcal{T}_1} := <_{\mathcal{T}_2} \upharpoonright u_{\mathcal{T}_1}$,

- (c) if $\mathcal{T}_1 \models \text{“}\eta \cap \nu = \rho\text{”}$ then $\mathcal{T}_2 \models \text{“}\eta \cap \nu = \rho\text{”}$,
- (d) $R_{\mathcal{T}_1, \ell} = R_{\mathcal{T}_2, \ell} \upharpoonright u_{\mathcal{T}_1}$ for $\ell = 0, 1$,
- (e) $\langle^*_{\mathcal{T}_1} := \langle^*_{\mathcal{T}_2} \upharpoonright u_{\mathcal{T}_1}$;
- (f) $E_{\mathcal{T}_1} := E_{\mathcal{T}_2} \upharpoonright u_{\mathcal{T}_1}$,

4) For $s \in \mathcal{T}$ and $\ell \in \{0, 1\}$, let $\text{succ}_{\mathcal{T}, \ell}(s)$ be the unique immediate successor of s in \mathcal{T} such that $(s, t) \in R_\ell^{\mathcal{T}}$.

5) We say $\mathcal{T}_1, \mathcal{T}_2 \in \mathbf{T}$ are *neighbors* when they are equal except that for each $t \in \mathcal{T}_1$ we can change the order $\langle^*_{\mathcal{T}_1} \upharpoonright (t/E_{\mathcal{T}_1})$ to $\langle^*_{\mathcal{T}_2} \upharpoonright (t/E_{\mathcal{T}_2})$.

Definition 1.2. 1) We say f is a \subseteq -embedding of $\mathcal{T}_1 \in \mathbf{T}$ into $\mathcal{T}_2 \in \mathbf{T}$ when: f is an isomorphism from \mathcal{T}_1 onto \mathcal{T}'_1 where $\mathcal{T}'_1 \subseteq \mathcal{T}_2$.

2) For any ordinal α (limit, if not said otherwise) and sequence $\bar{\zeta} = \langle \langle \beta : \beta < \alpha \rangle \rangle$, with $\langle \beta$ a well ordering of ${}^\beta 2$ we define $\mathcal{T} = \mathcal{T}_{\alpha, \bar{\zeta}}$ as follows (omitting $\bar{\zeta}$ means “for some”):

- (a) universe ${}^{\alpha > 2}$,
- (b) $\langle_{\mathcal{T}}$ is $\triangleleft \upharpoonright {}^{\alpha > 2}$,
- (c) $E_{\mathcal{T}} := \{(\eta, \nu) : \eta, \nu \in {}^\beta 2 \text{ for some } \beta < \alpha\}$,
- (d) $\langle^*_{\mathcal{T}} := \{(\eta, \nu) : \eta, \nu \in {}^{\alpha > 2} \text{ and } \ell g(\eta) < \ell g(\nu) \text{ or } (\exists \beta < \alpha)(\ell g(\eta) = \beta = \ell g(\nu) \wedge \eta <_{\beta} \nu)\}$,
- (e) $R_\ell := \{(\eta, \nu) : \eta \hat{\ } \langle \ell \rangle \trianglelefteq \nu \in \mathcal{T}\}$,
- (f) $\eta \cap_{\mathcal{T}} \nu := \eta \cap \nu$.

3) For $\mathcal{T} \in \mathbf{T}$ and $\zeta < \text{ht}(\mathcal{T})$, let $\langle_{\mathcal{T}, \zeta}$ be $\langle^*_{\mathcal{T}} \upharpoonright \mathcal{T}_{[\zeta]}$.

Claim 1.3. *If $\theta = \sup\{(2^{|\alpha|})^+ : \alpha < \kappa\}$ and $\bar{\zeta} = \langle \langle \beta : \beta < \kappa \rangle \rangle$ as above, then $\mathcal{T}_{\kappa, \bar{\zeta}}$ is well defined and belongs to $\mathbf{T}_{\theta, \kappa}$.*

Proof. It is clear. □_{1.3}

Definition 1.4. 1) For $\mathcal{T} \in \mathbf{T}$ let $\text{eseq}_n(\mathcal{T})$ be the set of sequences \bar{a} such that:

- (a) \bar{a} is an $\langle^*_{\mathcal{T}}$ -increasing sequence of length n of members of \mathcal{T} ,
- (b) $k < \ell < n \Rightarrow a_k \cap a_\ell \in \{a_m : m < n\}$,
- (c) $k, \ell < n \wedge \text{lev}(a_k) \leq \text{lev}(a_\ell) \Rightarrow a_\ell \upharpoonright \text{lev}(a_k) \in \{a_m : m < n\}$,

1A) For $\mathcal{U} \subseteq \mathcal{T}$ we let $\text{eseq}_n(\mathcal{U}, \mathcal{T})$ be $\text{eseq}_n(\mathcal{T}) \cap ({}^n \mathcal{U})$, similarly for part (2).

2) Let $\text{eseq}(\mathcal{T}) = \text{eseq}_{< \omega}(\mathcal{T}) = \cup \{\text{eseq}_n(\mathcal{T}) : n < \omega\}$.

2A) For finite $A \subseteq \mathcal{T}$ we define the sequence $\bar{b} = \text{cl}(A) = \text{cl}_{\mathcal{T}}(A) = \text{cl}(A, \mathcal{T})$ as the unique \bar{b} such that:

- (a) $\bar{b} \in \text{eseq}(\mathcal{T})$,
- (b) $A \subseteq \text{Rang}(\bar{b})$,
- (c) $\text{Range}(\bar{b})$ is minimal under those restrictions.

Also let $\text{pos}(A) = \text{pos}(A, \mathcal{T})$ be the unique function h from A into $\text{lg}(\bar{b})$ such that for every $a \in A$ we have: $i = h(a)$ iff $b_i = a$.

2B) We may replace above A by a finite sequence \bar{a} , and let $\text{pos}_{\mathcal{T}}(\bar{a})$ be $\text{pos}_{\mathcal{T}}(\text{rang}(\bar{a}))$.

3) We say $\bar{a}, \bar{b} \in \text{eseq}(\mathcal{T})$ are \mathcal{T} -similar or $\bar{a} \sim_{\mathcal{T}} \bar{b}$ when for some n we have:

- (a) $\bar{a}, \bar{b} \in \text{eseq}_n(\mathcal{T})$,
- (b) for any $k, i, m < n$ we have:

- ₁ $a_k \leq_{\mathcal{T}} a_i$ iff $b_k \leq_{\mathcal{T}} b_i$,
- ₂ $(a_k, a_i) \in R_{\ell}^{\mathcal{T}}$ iff $(b_k, b_i) \in R_{\ell}^{\mathcal{T}}$ for $\ell = 0, 1$,
- ₃ $a_k \cap_{\mathcal{T}} a_{\ell} = a_m$ iff $b_k \cap_{\mathcal{T}} b_{\ell} = b_m$, actually follows,
- ₄ $a_k = a_{\ell} \upharpoonright \text{lev}(a_m)$ iff $b_k = b_{\ell} \upharpoonright \text{lev}(b_m)$, actually follows,
- ₅ $(a_k \cap a_m) R_{\mathcal{T}, \ell} a_i$ iff $(b_k \cap b_m) R_{\mathcal{T}, \ell} b_i$ for $\ell = 0, 1$; actually follows,
- ₆ $\text{lev}_{\mathcal{T}}(a_k) \leq \text{lev}_{\mathcal{T}}(a_{\ell})$ iff $\text{lev}_{\mathcal{T}}(b_k) \leq \text{lev}_{\mathcal{T}}(b_{\ell})$; actually follows.

3A) We say that $\bar{a}, \bar{b} \in {}^n \mathcal{T}$ are \mathcal{T} -similar when $\bar{a}' = \text{cl}(\bar{a}), \bar{b}' = \text{cl}(\bar{b})$ are \mathcal{T} -similar and $a'_{\ell} = a_k \Leftrightarrow b'_{\ell} = b_k$ for any $\ell < \text{lg}(\bar{a}'), k < n$.

4) For $\bar{a} \in {}^n \mathcal{T}$ let $\text{Lev}(\bar{a})$ be the set $\{\text{lev}_{\mathcal{T}}(a_{\ell}) : \ell < \text{lg}(\bar{a})\}$.

5) We say that $\mathcal{T} \in \mathbf{T}$ is weakly \aleph_0 -saturated when:

- (*) for every $\varepsilon < \text{ht}(\mathcal{T})$ and s_0, \dots, s_{n-1} from $\mathcal{T}_{[\varepsilon]}$, there are $\zeta \in (\varepsilon, \text{ht}(\mathcal{T}))$ and $t_0 <_{\mathcal{T}}^* \dots <_{\mathcal{T}}^* t_{n-1}$ from $\mathcal{T}_{[\zeta]}$ satisfying $k < n \Rightarrow s_k <_{\mathcal{T}} t_k$,

6) For $\mathcal{T} \in \mathbf{T}$ let:

- (a) $\text{incr}_n(\mathcal{T})$ be the set of $<_{\mathcal{T}}^*$ -increasing $\bar{a} \in {}^n \mathcal{T}$ and let $\text{seq}_n(\mathcal{T}) = {}^n \mathcal{T}$ that is the set sequences of length n from \mathcal{T} ,
- (b) $\text{incr}(\mathcal{T}) = \cup \{\text{incr}_n(\mathcal{T}) : n < \omega\}$ and $\text{seq}(\mathcal{T}) = \cup \{\text{seq}_n(\mathcal{T}) : n < \omega\}$.

7) For $\mathcal{T} \in \mathbf{T}$:

- (a) for $\bar{t} \in \text{incr}(\mathcal{T})$ or just $\bar{t} \in {}^n \mathcal{T}$ let $\text{sim} - \text{tp}(\bar{t}, \mathcal{T})$ be the pair (the similarity type of $\text{cl}(\bar{t}, \mathcal{T}), \text{pos}(\bar{t}, \mathcal{T})$), that is all the information from part (3) (of 1.4) (including pos),
- (b) if in addition, $\mathcal{U} \subseteq \mathcal{T}$ then we let $\text{sim} - \text{tp}(\bar{t}, \mathcal{U}, \mathcal{T})$ be the function mapping $\bar{s} \in \omega^{>} \mathcal{U}$ to $\text{sim} - \text{tp}(\bar{t} \hat{\ } \bar{s}, \mathcal{T})$.

8) Let \mathbb{S}^n be the set of similarity types of sequences of length n in some $\mathcal{T} \in \mathbf{T}$, so the sequences are not necessarily increasing.

9) Naturally $\mathbb{S} = \cup \{\mathbb{S}^n : n < \omega\}$.

Now comes the main property.

Definition 1.5. 1) For $\mathcal{T}_1, \mathcal{T}_2 \in \mathbf{T}$ and $n < \omega$ and a cardinal σ let $\mathcal{T}_1 \rightarrow (\mathcal{T}_2)_{\sigma}^n$ mean:

- (*) if $\mathbf{c} : \text{eseq}_n(\mathcal{T}_1) \rightarrow \sigma$, then there is a \subseteq -embedding g of \mathcal{T}_2 into \mathcal{T}_1 such that the colouring $\mathbf{c} \circ g$ is homogeneous for \mathcal{T}_2 , which means:
 - if $\bar{a}, \bar{b} \in \text{eseq}_n(\mathcal{T}_2)$ are \mathcal{T}_2 -similar, then $\mathbf{c}(g(\bar{a})) = \mathbf{c}(g(\bar{b}))$.

2) For $\mathcal{T}_1, \mathcal{T}_2 \in \mathbf{T}$, $k < \omega$ and σ , let $\mathcal{T}_1 \rightarrow (\mathcal{T}_2)_{\sigma}^{\text{end}(k)}$ mean that:

- (*) if $\mathbf{c} : \text{eseq}(\mathcal{T}_1) \rightarrow \sigma$ then there is an embedding g of \mathcal{T}_2 into \mathcal{T}_1 such that the colouring $\mathbf{c}' = \mathbf{c} \circ g$ (see below) satisfies $\mathbf{c}'(\bar{\eta})$ does not depend on the last k levels, that is:
 - ₁ the meaning of $\mathbf{c}' = \mathbf{c} \circ g$ is that for every $s_0, \dots, s_{m-1} \in \mathcal{T}_2$ we have $\mathbf{c}'(\langle s_0, \dots, s_{m-1} \rangle) = \mathbf{c}'(\langle g(s_0), \dots, g(s_{m-1}) \rangle)$,
 - ₂ if $n < \omega$ and $\bar{a}, \bar{b} \in \text{eseq}_n(\mathcal{T}_2)$ are \mathcal{T}_2 -similar and

$$\ell < n \wedge (k \leq |\text{Lev}(\bar{a}) \setminus \text{lev}(a_{\ell})|) \Rightarrow b_{\ell} = a_{\ell},$$

then $\mathbf{c}'(\bar{a}) = \mathbf{c}'(\bar{b})$.

3) Let $\mathcal{T}_1 \rightarrow (\mathcal{T}_2)_\sigma^{\text{end}(k,m)}$ is defined as in part (2), but we restrict in \bullet_2 demanding that \bar{a}, \bar{b} satisfy $m \geq |\{\ell < n : k \leq |\text{Lev}(\bar{a}) \setminus \text{lev}(a_\ell)|\}|$.

4) We define $\mathcal{T}_1 \rightarrow' (\mathcal{T}_2)_\sigma^n$ as in part (1), but $\bar{a}, \bar{b} \in \text{seq}_n(\mathcal{T}_2)$. We define similarly $\mathcal{T}_1 \rightarrow (\mathcal{T}_2)^{\leq n}$ and $\mathcal{T}_1 \rightarrow' (\mathcal{T}_2)^{\leq n}$.

We may mention some implications among the \rightarrow ,

Remark 1.6. Of course, the equality $\mathbf{c}(g(\bar{a})) = \mathbf{c}(g(\bar{b}))$ is required only if \bar{a} and \bar{b} are \mathcal{T}_2 -similar since this is the best possible homogeneity, as one can define a coloring according to similarity types.

Claim 1.7. *Let $\mathcal{T} \in \mathbf{T}$.*

1) *If $A \subseteq \mathcal{T}$ is finite non-empty with m elements then:*

- (a) *For some $n \leq (2m - 1)m^2$ and $\bar{a} \in \text{eseq}_n(\mathcal{T})$ we have $A \subseteq \text{Rang}(\bar{a})$; moreover $\max\{\text{lev}_{\mathcal{T}}(a) : a \in A\} = \max\{\text{lev}_{\mathcal{T}}(a_\ell) : \ell < n\}$; in fact $\bar{a} = \text{cl}_{\mathcal{T}}(A)$,*
- (b) *If $\mathcal{T} \in \mathbf{T}$ and finite $A \subseteq \mathcal{T}$ then $\text{cl}(A, \mathcal{T}), \text{pos}(A, \mathcal{T})$ are well defined.*

2) *The number of quantifier free complete n -types realized in some $\mathcal{T} \in \mathbf{T}$ by some $\bar{a} \in \text{eseq}_n(\mathcal{T})$ is, e.g. $\leq 2^{2n^2+n}$ but $\geq n$.*

3) *If $\mathcal{T} \in \mathbf{T}$ is weakly \aleph_0 -saturated then \mathcal{T} realizes all possible such types, i.e. each type realized in some $\mathcal{T}' \in \mathbf{T}$; here “ht(\mathcal{T}) is a limit ordinal” follows.*

4) *Assume $\mathcal{T} \in \mathbf{T}$ and \mathcal{U} is a subset of \mathcal{T} closed under $<_{\mathcal{T}}$, (that is $s <_{\mathcal{T}} t \in \mathcal{U} \Rightarrow s \in \mathcal{U}$). Let $\text{lev}(\mathcal{U}, \mathcal{T}) = \sup\{\text{lev}(s, \mathcal{T}) + 1 : s \in \mathcal{U}\}$.*

If $\text{lev}(\mathcal{U}, \mathcal{T}) < \text{lev}(t, \mathcal{T})$ and $A \subseteq \mathcal{U}$ is finite then $\bar{b} = \text{cl}(A \cup \{t\})$ has the form $\bar{c} \hat{\ } (t)$ with $\bar{c} \in \text{eseq}(\mathcal{T}) \cap {}^{\omega} \mathcal{U}$.

Proof. Clearly (4) holds and we shall use it freely.

1) Let $B_1 = \{\eta \cap_{\mathcal{T}} \nu : \eta, \nu \in A\}$ and note that $\eta \in A \Rightarrow \eta = \eta \cap \eta \in B_1$. Now by induction on $|A|$ easily $|B_1| \leq 2m - 1$. Let $B_2 := \{\eta \upharpoonright \text{lev}_{\mathcal{T}}(\nu) : \eta \in A, \nu \in B_1 \text{ and } \text{lev}_{\mathcal{T}}(\eta) \geq \text{lev}_{\mathcal{T}}(\nu)\}$.

Easily $B_2 = \text{cl}(A, \mathcal{T})$, also $|B_2| \leq m|B_1| = m(2m - 1)$.

We may improve the bound but this does not matter here; similarly below.

2) Considering the class of such pairs (\bar{a}, \mathcal{T}) (fixing n) the number of $E_{\bar{a}} = \{(k, i) : a_k E_{\mathcal{T}} a_i\}$ is $\leq 2^{n^2}$ and the number of $<_{\bar{a}} = \{(k, i) : a_k <_{\mathcal{T}} a_i\}$ is $\leq 2^{n^2}$ and the number of $\{(a_k, a_i) : (a_k, a_i) \in R_1^{\mathcal{T}} \text{ and for no } j, a_k <_{\mathcal{T}} a_j <_{\mathcal{T}} a_i\}$ is $\leq 2^n$.

Lastly, from those we can compute $\{(a_k, a_i) : (a_k, a_i) \in R_0^{\mathcal{T}}\}$ as $\{(a_k, a_i) : (a_k \cap_{\mathcal{T}} a_i = a_k) \wedge ((a_k, a_i) \notin R_1) \wedge a_k \neq a_i\}$, so together the number is $\leq 2^{2n^2+n}$.

Clearly we can get a better bound, e.g. letting $m_n^\bullet(\mathcal{T}) = |\{\text{tp}_{\text{qf}}(\bar{a} \upharpoonright n, \emptyset, \mathcal{T}) : \bar{a} \in \text{eseq}(\mathcal{T}) \text{ has length } \geq n\}|$ then:

- (*)₁ \bullet_1 $m_n^\bullet(\mathcal{T}) = 1$ for $n = 0, 1$,
- \bullet_2 $m_{n+1}^\bullet(\mathcal{T}) \leq 4n(m_n^\bullet(\mathcal{T}))$,
- \bullet_3 hence $m_n^\bullet(\mathcal{T}) \leq 4^{n-1}(n-1)!$.

[Why? e.g. for \bullet_2 notice that $\text{tp}_{\text{qf}}(\bar{a} \upharpoonright (n+1), \emptyset, \mathcal{T})$ is determined by $q = \text{tp}_{\text{qf}}(\bar{a} \upharpoonright n, \emptyset, \mathcal{T})$ and the unique triple $(m, \iota, \ell) \in n \times 2 \times 2$ such that:

- (*)_{1.1} (a) $m < n$ is such that $\text{lev}(a_m \cap a_n)$ is maximal, hence $a_m <_{\mathcal{T}} a_n$,

- (b) $a_m R_i a_n$,
- (c) ℓ iff $\text{Lev}_{\mathcal{T}}(a_n) > \text{Lev}_{\mathcal{T}}(a_{n-1})$.

As there are $\leq 4n$ possibilities we are done].

It suffices to consider the case \mathcal{T} is weakly \aleph_0 -saturated (see 1.4(5), 1.7(3)) and then we can get exact values.

Now for $n \geq k \geq 1$ let,

$$m_{n,k}^*(\mathcal{T}) := |\{\text{tp}_{\text{qf}}(\bar{a}, \emptyset, \mathcal{T}) : \bar{a} \in \text{eseq}_n(T) \text{ such that } |\{\ell : \text{lev}(a_\ell) = \max(\text{Lev}(\bar{a}))\}| = k\}|.$$

So,

- $m_{1,1}^*(\mathcal{T}) = 1$, $m_{1,0}^*(\mathcal{T}) = 0$ and stipulate $m_{0,k}^*(\mathcal{T}) = 0$,
- if $n = k \geq 1$, then $m_{n,k}^*(\mathcal{T}) = 1$,
- if $2k - 1 > n \geq k \geq 1$, then $m_{n,k}^*(\mathcal{T}) = 0$,
- if $n \geq 1$, then $m_{n+1,1}^*(\mathcal{T}) = \Sigma\{2k \cdot m_{n,k}^*(\mathcal{T}) : k \in [1, n]\}$,

and more generally,

- if $n > k \geq 1$, then

$$m_{n+k,k}^* = \sum \left\{ \ell! \cdot \binom{\ell}{\ell_1} \cdot \binom{\ell - \ell_1}{\ell_2} \cdot 2^{\ell} \cdot m_{n,\ell}^*(\mathcal{T}) : \ell, \ell_0, \ell_1, \ell_2 \in [0, n], \ell = \ell_0 + \ell_1 + \ell_2 \right\}.$$

[Why? Considering $p = \text{tp}_{\text{qf}}(\bar{a} \upharpoonright (n+k), \emptyset, \mathcal{T})$ we fix $q = \text{tp}(\bar{a} \upharpoonright n, \emptyset, \mathcal{T})$, let ℓ be maximal such that $n - \ell \leq i < n \Rightarrow \text{lev}(a_i) = \text{lev}(a_{n-1})$ (equivalently $\text{lev}(a_{n-\ell}) = \text{lev}(a_{n-1})$). For $\iota = 0, 1, 2$, let $S_\iota = \{m : n - \ell \leq m < n \text{ and } \iota = |\{j < k : a_m <_{\mathcal{T}} a_j\}|\}$, so (S_0, S_1, S_2) is a partition of $[n - \ell, n)$. Let $S_1^\bullet = \{m \in S_1 : \text{if } j < k \text{ then } a_m R_1 a_j\}$. Fixing ℓ the number of possible q 's in $m_{n,\ell}^*(\mathcal{T})$ and fixing q (and so ℓ) the freedom left is choosing $\ell_0, \ell_1, \ell_2 \geq 0$ such that $\ell_1 + 2\ell_2 = k$ and then choosing the partition (S_0, S_1, S_2) which have $\binom{\ell}{\ell_1} \binom{\ell - \ell_1}{\ell_2}$ possibilities we have 2^{ℓ_1} possible choices of S_1^\bullet and lastly k possible linear orders of $\{a_i : i \in [n, n+k]\}$ clearly we are done.]

3), 4), Clear. □_{1.7}

Claim 1.8. *Let² $\sigma \geq \aleph_0$*

- 1) *If $\mathcal{T} \rightarrow (\mathcal{T})_\sigma^{\text{end}(1)}$, then $\mathcal{T} \rightarrow (\mathcal{T})_\sigma^{\text{end}(k)}$ for every $k < \omega$.*
- 2) *If $\mathcal{T} \in \mathbf{T}$ and $\mathcal{T} \rightarrow (\mathcal{T})_\sigma^{\text{end}(1)}$, then $\mathcal{T} \rightarrow (\mathcal{T})_\sigma^n$ for every $n < \omega$.*
- 3) *If $k \geq 1$ and $\mathcal{T}_\ell \in \mathbf{T}$ for $\ell = 0, \dots, k$ and $\mathcal{T}_{\ell+1} \rightarrow (\mathcal{T}_\ell)_\sigma^{\text{end}(1,m)}$ for $\ell < k$, then $\mathcal{T}_k \rightarrow (\mathcal{T}_0)_\sigma^{\text{end}(k,m)}$, hence $\mathcal{T}_k \rightarrow (\mathcal{T}_0)_\sigma^{\leq m}$.*

Proof. 1.8 Clear. □

§ 1(B). Forcing in ZFC.

Remark 1.9. Concerning the choice of $m(*)$ in 1.10 below (given \mathbf{m}), its value is immaterial for the model theoretic results.

In [She92, §4] we use Erdős-Radó theorem for $m = 2n$, but what we really need in essence is the canonical version, see [EHMR84], [DH17] and references there.

²If $\sigma < \aleph_0$ we have parallel results depending decreasing σ in the [conclusion](#) on the bounds from [a4](#), that is, for part (2): if $\mathcal{T} \rightarrow (\mathcal{T})_{\sigma(1)}^{\text{end}(1,m(1))}$, then $\mathcal{T} \rightarrow (\mathcal{T})_{\sigma(2)}^{\text{end}(n,m(2))}$.

Claim 1.10. *In $\mathbf{V}^{\mathbb{P}}$ we have $\mathcal{T}_2 \rightarrow (\mathcal{T}_1)_\sigma^{\text{end}(1,m)}$ when for a suitable $m(*)$:*

- (a) $\kappa = \kappa^{<\kappa}$, $\lambda \rightarrow (\kappa_1)_\sigma^{m(*)}$, $\sigma = 2^\kappa$ and $\kappa_1 \rightarrow (\kappa^+)_\kappa^m$,
- (b) $\mathbb{P} = \text{Cohen}(\kappa, \lambda)$, noting δ^* has cofinality κ because $u_* = \mathcal{B} \cap \lambda$, $[\mathcal{B}]^{<\kappa} \subseteq \mathcal{B}$ and $\|\mathcal{B}\| = \kappa$,
- (c) $\mathcal{T}_2 \in \mathbf{T}$ expands $(\kappa^{(+)>2}, \triangleleft)$ in $\mathbf{V}^{\mathbb{P}}$,
- (d) In $\mathbf{V}^{\mathbb{P}}$, $\mathcal{T}_1 \in \mathbf{T}_{\kappa,\kappa}$ and $\mathcal{T}_1 \subseteq \mathcal{T}_1^+$, where \mathcal{T}_1^+ expands $(\kappa^{>2}, \triangleleft)$ and so $\text{otp}(\mathcal{T}_{1,[\alpha]}, <\mathcal{T}_{1,\alpha}) < \kappa$ for $\alpha < \kappa$, see 1.2(3),
- (e) let $\mathfrak{A}_1 = \mathfrak{A}_0[\mathbf{G}_{u_*}]$ and $\mathfrak{A}_2 = \mathfrak{A}_0[\mathbf{G}]$.

Proof. First,

\square_1 Without loss of generality, $\mathcal{T}_1 \in \mathbf{V}$ is an object (not just a \mathbb{P} -name).

[Why? Let \mathcal{T}_1 be a \mathbb{P} -name, then for some $u \in [\lambda]^{\leq \kappa}$ we have \mathcal{T}_1 is a \mathbb{P}_u -name. We can force by \mathbb{P}_u , so as $\mathbb{P}/\mathbb{P}_u = \text{Cohen}(\kappa, \lambda)$, we are done.]

So $\kappa, \mathcal{T}_1, \mathcal{T}_2$ are well defined (\mathcal{T}_2 a \mathbb{P} -name). Let \mathfrak{c} be a \mathbb{P} -name, $\mathfrak{c}: \text{eseq}(\mathcal{T}_2) \rightarrow \sigma$, without loss of generality be such that:

\square_2 if $\bar{t} \in \text{eseq}(\mathcal{T}_2)$, then from $\mathfrak{c}(\bar{t})$ we can compute the similarity type of \bar{t} in \mathcal{T}_2 and of $\mathfrak{c}(\bar{t} \upharpoonright u)$ when $u \subseteq \text{dom}(\bar{t})$, $\bar{t} \upharpoonright u \in \text{eseq}(\mathcal{T}_2)$.

Let $\bar{\eta} = \langle \eta_\alpha : \alpha < \lambda \rangle$ be the generic of \mathbb{P} , so $\Vdash \eta_\alpha \in {}^\kappa 2$ and let $\bar{\eta}_u = \langle \eta_\alpha : \alpha \in u \rangle$ for $u \subseteq \lambda$.

Next, in \mathbf{V} , we choose:

- (*)₁ (a) let $\chi > \lambda$ and $<_\chi^*$ a well ordering of $\mathcal{H}(\chi)$,
- (b) let $\mathfrak{B} \prec \mathfrak{A}_0 = (\mathcal{H}(\chi), \in, <_\chi^*)$ be of cardinality κ such that $[\mathfrak{B}]^{<\kappa} \subseteq \mathfrak{B}$ and $\lambda, \kappa, \mu, \sigma, \mathcal{T}_1, \mathcal{T}_2, \mathfrak{c} \in \mathfrak{B}$,
- (c) let $u_* = \mathfrak{B} \cap \lambda \in [\lambda]^\kappa$,
- (d) let $\mathbf{G}_{u_*} \subseteq \mathbb{P}_{u_*}$ be generic over $\mathbf{V}_0 = \mathbf{V}$, $\mathbf{G} \subseteq \mathbb{P}$ be generic over \mathbf{V}_0 such that $\mathbf{G}_{u_*} \subseteq \mathbf{G}$,
- (e) let $\bar{\eta}_u = \langle \eta_\alpha[\mathbf{G}_u] : \alpha \in u \rangle$, for $u \subseteq \lambda$,
- (f) let $\mathbf{V}_1 = \mathbf{V}_0[\mathbf{G}_{u_*}] = \mathbf{V}_0[\bar{\eta}_{u_*}]$,
- (g) let $\mathbf{V}_2 = \mathbf{V}_0[\bar{\eta}_\lambda] = \mathbf{V}_1[\bar{\eta}_{\lambda \setminus u_*}]$.
- (*)₂ (a) let \mathcal{T}_0 be the sub-structure of \mathcal{T}_2 with set of elements $\{\eta : \eta \text{ is a canonical } \mathbb{P}\text{-name of a member of } \mathcal{T}_2 \text{ and this name belongs to } \mathfrak{B}\}$,
- (b) let $\delta_* = \delta(*)$ be $\min(\lambda \setminus u_*) = \min(\kappa^+ \setminus u_*) = \kappa^+ \cap u_*$,
- (c) let $\langle \delta_\varepsilon : \varepsilon < \kappa \rangle$ be increasing continuous with limit δ_* in \mathbf{V}_0 ,
- (d) $\mathfrak{B}_2 = \mathfrak{B}[\mathbf{G}_\lambda]$, $\mathfrak{B}_1 = \mathfrak{B}_2 \upharpoonright \{\mathcal{T}[\mathbf{G}_\lambda] : \mathcal{T} \text{ is a } \mathbb{P}_u\text{-name from } \mathfrak{B} \text{ for some } u \in [\lambda]^{\leq \kappa} \cap \mathfrak{B}\}$, $\mathfrak{B}_0 = \mathfrak{B}$, so $\mathfrak{B}_2 \prec \mathfrak{A}_2 = \mathcal{H}(\chi)[\mathbf{G}_\lambda]$ and $\mathfrak{B}_1 \cap \kappa^{\geq \lambda} = \mathfrak{B}_2 \cap \kappa^{\geq \lambda}$.

Clearly,

- (*)₃ (a) $\Vdash \mathcal{T}_0 \subseteq \mathcal{T}_2$ is closed under initial segments, is of cardinality κ and has δ_* levels and is closed under unions of increasing chains of length $< \kappa$ and $\nu \in \mathcal{T}_0 \Rightarrow \nu \wedge \langle 0 \rangle, \nu \wedge \langle 1 \rangle \in \mathcal{T}_0$, and $\alpha < \delta_* \Rightarrow (\forall \nu \in \mathcal{T}_0)(\exists \rho)[\nu \triangleleft \rho \in \mathcal{T}_0 \wedge \text{lg}(\rho) \geq \alpha]$, so $\mathcal{T}_0 \in \mathbf{T}$,
- (b) \mathcal{T}_0 is actually a \mathbb{P}_{u_*} -name and we can use $\delta_* = \mathfrak{B} \cap \kappa^+$ as its set of levels.

- (*)₄ (a) let $\mathcal{T}_0 = \mathcal{T}_0[\mathbf{G}_{u_*}]$, $\mathbf{c}_0 = \mathbf{c} \upharpoonright \text{eseq}(\mathcal{T}_0)$ so they are from \mathbf{V}_1 ,
 (b) let $\mathbb{P}_* = \mathbb{P}/\mathbf{G}_{u_*} = \mathbb{P}_{\lambda \setminus u_*}$.
- (*)₅ (a) for each $\alpha \in \lambda \setminus u_*$, in $\mathbf{V}_1[\eta_\alpha]$ there is $\eta_\alpha^\bullet \in \lim_{\delta_*}(\mathcal{T}_0)^{\mathbf{V}_1[\eta_\alpha]}$ with $\text{lg}(\eta_\alpha^\bullet) = \delta_*$ such that $\varepsilon < \delta_* \Rightarrow \eta_\alpha^\bullet \upharpoonright \varepsilon \in \mathcal{T}_0$ and η_α^\bullet is a generic δ_* -branch of \mathcal{T}_0 over \mathbf{V}_1 ,
 (b) Clearly $\lim_{\delta_*}(\mathcal{T}_0)^{\mathbf{V}_1[\eta_\alpha]} \subseteq \mathcal{T}_2[G_\lambda]$.

We shall work in \mathbf{V}_1 .

- (*)₆ (in \mathbf{V}_1) for $\alpha \in \lambda \setminus u_*$, let η_α^\bullet be a $\mathbb{P}_{\{\alpha\}}$ -name of η_α^\bullet , so without loss of generality for some κ -Borel function $\mathbf{B}: \kappa^2 \rightarrow \delta^{(*)}2$, from \mathbf{V}_1 , we have $\Vdash \eta_\alpha^\bullet = \mathbf{B}(\eta_\alpha)$ is as above in (*)₅(a)". [Saharon 2022-11-06, recheck]
- (*)₇ (a) Recall we had in (*)₂(c) fix an increasing sequence $\langle \delta(\varepsilon) := \delta_\varepsilon : \varepsilon < \kappa \rangle$ so that $\delta(*) = \bigcup_{\varepsilon < \kappa} \delta(\varepsilon)$
 (b) Let $\langle \mathbf{B}_\varepsilon : \varepsilon < \kappa \rangle$ be such that \mathbf{B}_ε is a κ -Borel function from κ^2 to $\delta(\varepsilon)2$ from \mathbf{V}_1 and $\Vdash \eta_\alpha^\bullet \upharpoonright \delta_\varepsilon = \mathbf{B}_\varepsilon(\eta_\alpha)$ ",
 (c) Let $\mathbf{B}, \mathbf{B}_\varepsilon$ be \mathbb{P}_{u_*} -names forced to be as above can be considered as \mathbb{P}_{u_*} -name.

Without loss of generality (using the freedom in choosing \mathfrak{B}).

- (*)₈ If $\varepsilon < \kappa$, $\nu \in \kappa^2$, $\mathbf{B}_\varepsilon(\nu) = \rho$, then for every large enough $\zeta < \kappa$, we have $\nu \upharpoonright \zeta \triangleleft \varrho \in \kappa^2 \Rightarrow \rho \triangleleft \mathbf{B}(\varrho)$ and we write $\mathbf{B}_\varepsilon(\nu \upharpoonright \zeta) = \rho$.

Recalling we work in \mathbf{V}_1 there are \mathcal{U}, \bar{N} such that:

- (*)₉ (a) $\mathcal{U} \subseteq \lambda \setminus u_*$,
 (b) $\text{otp}(\mathcal{U})$ satisfies $\text{otp}(\mathcal{U}) \rightarrow (\kappa^+)_\kappa^m$
 (c) $\bar{N} = \langle N_u : u \in [\mathcal{U}]^{\leq m} \rangle$,
 (d) $N_u \cap N_v \subseteq N_{u \cap v}$,
 (e) $\kappa, \mathcal{T}_1, \mathcal{T}_2, \mathfrak{c}, \mathfrak{B}, \mathfrak{B}_1 \in N_u \prec \mathfrak{A}_1, \|N_u\| = \kappa, [N_u]^{< \kappa} \subseteq N_u$,
 (f) if $u, v \in [\mathcal{U}]^{\leq m}$ and $|u| = |v|$, then there is a unique isomorphism $g_{u,v}$ from N_v onto N_u which is the identity on $(\kappa + 1) \cup \{\mathfrak{c}, \eta, \mathbf{B}\}$ hence on $\mathcal{T}_2, \langle \eta_\alpha, \eta_\alpha^\bullet : \alpha < \lambda \setminus u_* \rangle$ and maps v onto u ,
 (g) $\langle \eta_\alpha^\bullet : \alpha \in \mathcal{U} \rangle$ is $<_{\mathcal{T}_2}$ -increasing.

[Why? As in [She89], if $m(*) = 2n$; for smaller $m(*)$ this is close to canonical partitions, see [EHMR84]; [DH17] and references there; for clause (g) recall that $<_{\mathcal{T}_2}^*$ is a linear well ordering.] [2222-11-06, Saharon elaborate +, see $\boxplus_{\mathfrak{a}}^0$.]

(*)₁₀ notation:

- (a) in \mathbf{V}_2 for $\bar{s} \in \text{eseq}(\mathcal{T}_2)$, let $\varepsilon(\bar{s}) := \max\{\text{lev}_{\mathcal{T}_2}(s_i) : i < \text{lg}(\bar{s})\}$, $w(\bar{s}) = \{i < \text{lg}(\bar{s}) : \text{lev}_{\mathcal{T}_2}(s_i) = \varepsilon(\bar{s})\}$ and $u(\bar{s}) = \{s_i : i \in w(\bar{s})\}$ and $v(\bar{s}) = \{s_i : i < \text{lg}(\bar{s}) \text{ and } i \notin u(\bar{s})\}$,
- (b) for finite $u \subseteq \mathcal{T}_{0, [\varepsilon]}$ for some $\varepsilon < \delta_*$, let $\mathbf{H}_u := \{h : h \text{ is a one-to-one function from } u \text{ into } \mathcal{U}\}$,
- (c) If in \mathbf{V}_2 , $\bar{s} \in \text{eseq}(\mathcal{T}_0)$, $u = u[\bar{s}] \in [\mathcal{T}_{0, [\varepsilon]}]^{\leq n}$ and $h \in \mathbf{H}_u$, then we let $\bar{s}^{[h]}$ be the $\bar{t} \in \text{eseq}(\mathcal{T}_2)$ such that $\text{lg}(\bar{t}) = \text{lg}(\bar{s})$, $i \in \text{lg}(\bar{s}) \setminus u \Rightarrow t_i = s_i$ and $i \in w(\bar{s}) \Rightarrow t_i = s_i \hat{\wedge} \eta_{h(s_i)}^\bullet \upharpoonright [\varepsilon(\bar{s}), \delta_*)$.

We define $\text{AP} := \bigcup_{\varepsilon < \kappa} \text{AP}_\varepsilon$, where AP_ε is the set of objects \mathfrak{a} which consists of (so $\varepsilon = \varepsilon_{\mathfrak{a}}, \bar{v} = \bar{v}_{\mathfrak{a}}$, etc).

- $\boxplus_{\mathbf{a}}^0$
- (a) $\varepsilon < \kappa$,
 - (b) $\bar{\eta} = \langle \eta_\rho : \rho \in \mathcal{T}_{1, [\varepsilon]} \rangle$ is with no repetitions,
 - (c) $\bar{\nu} = \langle \nu_\rho : \rho \in \mathcal{T}_{1, [\varepsilon]} \rangle$,
 - (d) $\eta_\rho \in \mathcal{T}_{0, [\delta(\zeta)]}$ and $\nu_\rho \in \kappa^{>2}$ for some $\zeta = \zeta_{\mathbf{a}} < \kappa$ but $\zeta < \varepsilon$ (for all $\rho \in \mathcal{T}_{1, [\varepsilon]}$),
 - (e) $\mathbf{B}_\zeta^*(\nu_\rho) = \eta_\rho$,
 - (f) $\bar{p} = \langle p_{u,h} : u \in [^\varepsilon 2]^{\leq n}, h \in \mathbf{H}_u \rangle$, where $p_{u,h} \in \mathbb{P}_{\lambda \setminus u_*}$,
 - (g) $p_{u,h} \in N_{h[u]}$ and $[\rho \in u \wedge h(\rho) = \alpha \Rightarrow p_{u,h}(\alpha) = \nu_\rho]$,
[Saharon: **incomp** 2022-11-06 **was** hence $\eta_{h(\rho)}^\bullet = s_i$ in $(*)_{10}$ (c) above,
Shimoni **object**]
 - (h) if $h_1, h_2 \in \mathbf{H}_u$ and $h_1[u] = h_2[u]$ then $\alpha \in \text{dom}(p_{u,h_1}) \setminus h_1[u] \Rightarrow p_{u,h_1}(\alpha) = p_{u,h_2}(\alpha)$,
 - (i) if $h_1, h_2 \in \mathbf{H}_u$ and $\rho_1, \rho_2 \in u \Rightarrow h_1[\rho_1] < h_1[\rho_2] \equiv h_2[\rho_1] < h_2[\rho_2]$,
then $g_{h_2(u), h_1(u)}$ maps p_{u,h_1} to p_{u,h_2} ,
 - (j) $p_{u,h}$ forces a value to $\mathbf{c}(\bar{s}^{[h]})$ when $\bar{s} \in \text{eseq}(\mathcal{T}_0)$ and $u = u[\bar{s}]$,
 - (k) if $u_1 \subseteq u_2 \in [^\varepsilon 2]^{\leq n}$, $h_\ell \in \mathbf{H}_{u_\ell}$, $h_1 \subseteq h_2$, then $p_{u_1, h_1} \leq_{\mathbb{P}_*} p_{u_2, h_2}$.

Let further.

- $\boxplus_{\mathbf{a}}^1$ $\text{AP}^+ = \bigcup \{ \text{AP}_\varepsilon^+ : \varepsilon < \kappa \}$, where AP_ε^+ is the set of $\mathbf{a} \in \text{AP}_\varepsilon$ such that:
- (1) for some $\zeta \in [\varepsilon, \zeta_{\mathbf{a}}]$ if (α) then (β) , where:
 - (α) $\bar{s} \in \text{eseq}(\mathcal{T}_1)$, $u = u[\bar{s}]$ and $h \in \mathbf{H}_u$,
 - (β) \bullet_1 for every $\rho, \varrho \in u[\bar{s}]$, we have that $\eta_\rho \leq_{\mathcal{T}_{1, \zeta}} \eta_\varrho \Leftrightarrow p_{u,h} \Vdash \eta_{h(\rho)}^\bullet \upharpoonright \zeta <_{\mathcal{T}_{2, \delta(*)}} \eta_{h(\varrho)}^\bullet \upharpoonright \zeta$,
 - \bullet_2 $p_{u,h} \Vdash \mathbf{c}(\bar{s}^{[h]}) = j$ iff $\mathbf{c}(\bar{t}) = j$, where $\bar{t} \in \text{eseq}(\mathcal{T}_0)$, $\text{lg}(\bar{t}) = \text{lg}(\bar{s})$, $i \in \text{lg}(\bar{s}) \setminus u \Rightarrow t_i = s_i$ and $i \in u \Rightarrow t_i = \eta_{s_i} \upharpoonright \zeta$ noting $i \in u \Rightarrow s_i \in \mathcal{T}_{1, [\varepsilon]}$.
- \boxplus_2 we define the two-place relation \leq_{AP} as follows: $\mathbf{a}_1 \leq_{\text{AP}} \mathbf{a}_2$ iff:
- (a) $\mathbf{a}_1, \mathbf{a}_2 \in \text{AP}$,
 - (b) $\varepsilon_1 = \varepsilon_{\mathbf{a}_1} \leq \varepsilon_{\mathbf{a}_2} = \varepsilon_2$,
 - (c) if $\iota \in \{0, 1\}$, $\rho_1 \in {}^{\varepsilon(1)} 2$, $\rho_1 \hat{\ } \langle \iota \rangle \trianglelefteq \rho_2 \in {}^{\varepsilon(2)} 2$, then $\eta_{\mathbf{a}_1, \rho_1} \hat{\ } \langle \iota \rangle \trianglelefteq \eta_{\mathbf{a}_2, \rho_2}$ and $\nu_{\mathbf{a}_1, \rho_1} \hat{\ } \langle \iota \rangle \trianglelefteq \nu_{\mathbf{a}_2, \rho_2}$,
 - (d) if $m < \omega$, $u_1 \in [{}^{\varepsilon(1)} 2]^m$, $u_2 \in [{}^{\varepsilon(2)} 2]^m$, $u_1 = \{ \rho \upharpoonright \varepsilon(1) : \rho \in u_2 \}$, $h_\ell \in \mathbf{H}_{u_\ell}$ and $\rho \in u_2 \Rightarrow h_2(\rho) = h_1(\rho \upharpoonright \varepsilon(1))$, then $p_{u_1, h_1} \leq_{\mathbb{P}_*} p_{u_2, h_2}$.

\boxplus_3 $(\text{AP}, <_{\text{AP}})$ is a partial order.

[Why? Read the definitions.]

\boxplus_4 for $\varepsilon = 0$ there is $\mathbf{a} \in \text{AP}_\varepsilon$.

[Why? Trivial.]

\boxplus_5 if $\varepsilon < \kappa$ is a limit ordinal and $\mathbf{a}_\zeta \in \text{AP}_\zeta$ for $\zeta < \varepsilon$ is \leq_{AP} increasing, then there is $\mathbf{a}_\varepsilon \in \text{AP}_\varepsilon$ such that $\zeta < \varepsilon \Rightarrow \mathbf{a}_\zeta <_{\text{AP}} \mathbf{a}_\varepsilon$.

[Why? Straightforward.]

\boxplus_6 if $\varepsilon < \kappa$ and $\mathbf{a} \in \text{AP}_\varepsilon$ there is \mathbf{b} such that:

- (a) $\mathbf{b} \in \text{AP}_\varepsilon$,
- (b) $\mathbf{a} \leq_{\text{AP}} \mathbf{b}$,
- (c) $\varepsilon_{\mathbf{a}} = \varepsilon_{\mathbf{b}} + 1$,

[Why? Straightforward.]

\boxplus_7 if $\mathbf{a} \in \text{AP}_\varepsilon$, then there is $\mathbf{b} \in \text{AP}_\varepsilon^+$ such that $\mathbf{a} \leq_{\text{AP}} \mathbf{b}$.

Why? Noting,

- $\boxplus_{7.1}$ if $\bar{\eta}' = \langle \eta'_\rho : \rho \in \mathcal{T}_{1, [\varepsilon]} \rangle$ and $\eta_{\mathbf{a}, \rho} \leq \eta'_\rho \in \mathcal{T}_1$ for $\rho \in \mathcal{T}_{1, [\varepsilon]}$, then:
- (a) there is $\mathbf{b} \in \text{AP}_\varepsilon$ such that $\mathbf{a} \leq_{\text{AP}} \mathbf{b}$ and $\eta'_\rho \triangleleft \eta_{\mathbf{b}, \rho}$,
 - (b) moreover there is \mathbf{b} as above such that for some $\zeta \in (\zeta_{\mathbf{a}}, \zeta_{\mathbf{b}})$ such that:
 - if $\rho, \varrho \in \mathcal{T}_{1, [\varepsilon]}$, then $\rho <_{\mathcal{T}_1}^* \varrho \Leftrightarrow \eta_{\mathbf{b}, \rho} \upharpoonright \zeta <_{\mathcal{T}_0}^* \eta_{\mathbf{b}, \varrho} \upharpoonright \zeta$,
 - $\langle \eta_{\mathbf{b}, \rho} \upharpoonright \zeta : \rho \in \mathcal{T}_{1, [\varepsilon]} \rangle$ is as in clause (1) (see $\boxplus_{\mathbf{a}}^1$).

[Why? Let $\alpha_\rho \in \mathcal{U}$ for $\rho \in T_{1, [\varepsilon]}$ be such that $\rho <_{\mathcal{T}_{1, \varepsilon}} \varrho \Rightarrow \alpha_\rho < \alpha_\varrho$. Let $\bar{h} := \langle h_u : u \subseteq T_{1, [\varepsilon]} \text{ is finite} \rangle$ be such that $\rho \in u \in [\mathcal{T}_{1, [\varepsilon]}]^{< \aleph_0} \Rightarrow h_u(\rho) = \alpha_\rho$. Let $q_* = \bigcup \{p_{u, h_u} : u \subseteq \mathcal{T}_{1, [\varepsilon]} \text{ is finite}\}$, so $q_* \in \mathbb{P}_*$.

Clearly there is $p_* \in \mathbf{G}_0$ such that $p_* \cup q_* \in \mathbb{P}_\lambda$ forces all the relevant information related to \mathbf{a} . Now, as $\mathfrak{B}_0 \prec \mathfrak{A}_0$, there is a 1-to-1 function \mathbf{g} from $\text{dom}(p_* \cup q_*)$ such that $(p_* \cup q_*) = \mathbf{g}(p_* \cup q_*)$ satisfies all the relevant information; necessarily $q_\bullet \in \mathbb{P}_{u_*}$; without loss of generality $q_\bullet \in \mathbf{G}_0$ and the rest follows.]

Now \boxplus_7 follows because $\langle \eta_\alpha : \alpha \in \mathcal{U} \rangle$ is generic over \mathbf{V}_1 for $\mathbb{P}_{\mathcal{U}}$.

\boxplus_8 we can choose $\mathbf{a}_\varepsilon \in \text{AP}_\varepsilon$ by induction on $\varepsilon < \kappa$ such that $\zeta < \varepsilon \Rightarrow \mathbf{a}_\zeta \leq_{\text{AP}} \mathbf{a}_\varepsilon$ and $\varepsilon = \zeta + 1 \Rightarrow \mathbf{a}_\varepsilon \in \text{AP}_\varepsilon^+$.

[Why? Use \boxplus_4 for $\varepsilon = 0$, use \boxplus_5 for ε a limit ordinal and $\boxplus_6 + \boxplus_7$ for $\varepsilon = \zeta + 1$.]

To finish let $\zeta_\varepsilon \in [\varepsilon, \zeta_{\mathbf{a}_{\varepsilon+1}})$ be as in clause (j) of $\boxplus_{\mathbf{a}}^1$. Let $g: \kappa \rightarrow \kappa$ be increasing such that $\zeta_{(\mathbf{a}_{g(\varepsilon)})} < g(\varepsilon + 1)$.

Now define $h: \mathcal{T}_1 \rightarrow \mathcal{T}_2$ by $\rho \in \mathcal{T}_{1, [\varepsilon]} \Rightarrow h(\rho) = \eta_{\mathbf{a}_{g(\varepsilon)+1} \upharpoonright \zeta_{g(\varepsilon)}}$ and check. The indiscernibility holds by $\boxplus_{\mathbf{a}}^1(\ell)(\beta)$ not just $\boxplus_{\mathbf{a}}^0(\mathbf{g})$. $\square_{1.10}$

Remark 1.11. 1) Using the end of §1A we get the desired conclusions.

2) In 1.10 we may state and prove the variant with the square bracket. In more details,

- (A) We say $\bar{a}, \bar{b} \in \text{eseq}_m(\mathcal{T})$ are weakly \mathcal{T} -similar as before but omitting “ $a_\ell <_{\mathcal{T}}^* a_k \Leftrightarrow b_\ell <_{\mathcal{T}}^* b_k$ ”; that is, when $\text{lg}(\bar{a}) = \text{lg}(\bar{b})$ and for some permutation π of $\text{lg}(\bar{a})$ for $k, \ell, m < \text{lg}(\bar{a})$, we have:
 - (a) $\text{lev}_{\mathcal{T}}(a_\ell) = \text{lev}_{\mathcal{T}}(a_k) \Rightarrow \text{lev}_{\mathcal{T}}(b_{\pi(\ell)}) = \text{lev}_{\mathcal{T}}(b_{\pi(k)})$,
 - (b) $a_\ell <_{\mathcal{T}} a_k \Leftrightarrow b_{\pi(\ell)} <_{\mathcal{T}} b_{\pi(k)}$.
- (B) We replace “ $\mathcal{T}_2 \rightarrow (\mathcal{T}_1)_\sigma^{\text{end}(k, m)}$ ” by “ $\mathcal{T}_2 \rightarrow [\mathcal{T}_1]_{\sigma, j}^{\text{end}(k, m)}$ ” for suitable finite j which means that 1.5(1)(*) \bullet_2 is replaced by:
 - $_2$ if $n < \omega$ and $\bar{a} \in \text{eseq}_n(\mathcal{T}_2)$, then the following set has at most j elements:
$$\{\mathbf{c}'(\bar{b}) : \bar{b} \in \text{eseq}_n(\mathcal{T}_2) \text{ is weakly } \mathcal{T}_1\text{-similar to } \bar{a} \text{ and } \ell < n \wedge (k \leq |\text{Lev}(\bar{a})| \setminus \text{lev}(a_\ell)) \Rightarrow b_\ell = a_\ell\}.$$

3) This will be enough for the model theory and if we use minimal \mathbf{j} (well, depends on $\bar{s} \upharpoonright v(\bar{s})$) we get back to 1.10.

Conclusion 1.12. Assume $\chi \leq \infty$ is limit and $\theta < \chi \Rightarrow 2^\theta = \theta^+$.

We can find a forcing notion \mathbb{P} such that:

- (a) \mathbb{P} collapses no cardinal, changes no cofinality,
- (b) $2^\theta < \theta^{+\omega}$, $2^{2^\theta} = (2^\theta)^+$ for $\theta < \chi$,
- (c) in $\mathbf{V}^{\mathbb{P}}$, if $\theta^{+\omega} \leq \chi$, then for some $n < \omega$, for every \mathcal{T}_1 expanding (${}^\theta > 2, \triangleleft$) there is \mathcal{T}_2 expanding (${}^{\theta(+n)} > 2, \triangleleft$) such that $\mathcal{T}_2 \rightarrow (\mathcal{T}_1)_\theta^{\text{end}(n)}$.

Remark 1.13. Concerning the order on each level.

Discussion 1.14. We may like to replace ${}^{\kappa} > 2$ by ${}^{\kappa} > I$ and even use creature tree forcing, see Roslanowski-Shelah [RS99], [RS07], Goldstern-Shelah [GS05]) but (in second thought, for $\kappa = \aleph_0$ maybe see the paper with Zapletal [SZ11]). That is, for $\kappa > \aleph_0$ in each node we have a forcing notion which is quite complete, but of cardinality $< \kappa = \text{set of levels}$.

So we do not have a tree but a sequence of creatures, $\langle \mathfrak{c}_\varepsilon : \varepsilon < \text{ht}(\mathcal{T}) \rangle$, such that for a colouring we like to find $\mathfrak{d}_\varepsilon \in \Sigma(\mathfrak{c}_\varepsilon)$ for $\varepsilon < \kappa$, which induces a sub-tree in which the colouring is 1-end-homogeneous. Alternatively we have $\langle \mathfrak{c}_\eta : \eta \in \mathcal{T} \rangle$ where \mathfrak{c}_η is a creature a with set of possible values being in $\text{suc}_{\mathcal{T}}(\eta)$, see [RS07].

Clearly the answer is that we can, but it is not clear how interesting it is. We can just,

- replace 2 by $\Upsilon \in [2, \kappa)$ and ${}^{\kappa} > 2$ by ${}^{\kappa} > \Upsilon$; in the Definition 1.1 replace $R_{\mathcal{T}, \ell}$ ($\ell < 2$) by $R_{\mathcal{T}, \ell}$ ($\ell < \Upsilon$) and add: if $s \in \mathcal{T}$, then $\text{suc}_{\mathcal{T}}(s)$ is either a singleton or is $\{s_\ell : \ell < \Upsilon\}$, where $s_\ell R_{\mathcal{T}, \ell} s$ for $\ell < \Upsilon$.

REFERENCES

- [DH17] Natasha Dobrinen and Dan Hathaway, *The Halpern-Läuchli theorem at a measurable cardinal*, J. Symb. Log. **82** (2017), no. 4, 1560–1575.
- [DS79] Keith J. Devlin and Saharon Shelah, *A note on the normal Moore space conjecture*, Canadian J. Math. **31** (1979), no. 2, 241–251. MR 528801
- [DS04] Mirna Džamonja and Saharon Shelah, *On \triangleleft^* -maximality*, Ann. Pure Appl. Logic **125** (2004), no. 1-3, 119–158, arXiv: math/0009087. MR 2033421
- [EHMR84] Paul Erdős, Andras Hajnal, A. Maté, and Richard Rado, *Combinatorial set theory: Partition relations for cardinals*, Studies in Logic and the Foundation of Math., vol. 106, North-Holland Publ. Co, Amsterdam, 1984.
- [GS05] Martin Goldstern and Saharon Shelah, *Clones from creatures*, Trans. Amer. Math. Soc. **357** (2005), no. 9, 3525–3551, arXiv: math/0212379. MR 2146637
- [HL66] J. D. Halpern and H. Läuchli, *A partition theorem*, Trans. Amer. Math. Soc. **124** (1966), 360–367.
- [HL71] J. D. Halpern and A. Levy, *The boolean prime ideal theorem does not imply the axiom of choice*, Axiomatic set theory (Providence, R.I.), Proceedings of symposia in pure mathematics, 'vol. 13, part 1, American Mathematical Society, 1967, 1971, pp. 83–134.
- [Lav71] Richard Laver, *On fraïssé's order type conjecture*, Annals of Mathematics **93** (1971), 89–111.
- [Lav73] ———, *An order type decomposition theorem*, Annals of Mathematics **98** (1973), 96–119.
- [Mil79] Keith R. Milliken, *A Ramsey theorem for trees*, J. Combin. Theory Ser. A **26** (1979), no. 3, 215–237.
- [Mil81] ———, *A partition theorem for the infinite subtrees of a tree*, Trans. Amer. Math. Soc. **263** (1981), no. 1, 137–148.
- [RS99] Andrzej Roslanowski and Saharon Shelah, *Norms on possibilities. I. Forcing with trees and creatures*, Mem. Amer. Math. Soc. **141** (1999), no. 671, xii+167, arXiv: math/9807172. MR 1613600
- [RS07] ———, *Sheva-Sheva-Sheva: large creatures*, Israel J. Math. **159** (2007), 109–174, arXiv: math/0210205. MR 2342475
- [S⁺a] S. Shelah et al., *Tba*, In preparation. Preliminary number: Sh:F1822.

- [S⁺b] ———, *Tba*, In preparation. Preliminary number: Sh:F2060.
- [S⁺c] ———, *Tba*, In preparation. Preliminary number: Sh:F2064.
- [Shea] Saharon Shelah, *Dependent dreams: recounting types*, arXiv: 1202.5795.
- [Sheb] ———, *General non-structure theory and constructing from linear orders; to appear in Beyond first order model theory II*, arXiv: 1011.3576 Ch. III of The Non-Structure Theory” book [Sh:e].
- [She71a] ———, *Two cardinal and power like models: compactness and large group of automorphisms*, Notices Amer. Math. Soc. **18** (1971), no. 2, 425, 71 T-El5.
- [She71b] ———, *Two cardinal compactness*, Israel J. Math. **9** (1971), 193–198. MR 0302437
- [She75] ———, *A two-cardinal theorem*, Proc. Amer. Math. Soc. **48** (1975), 207–213. MR 357105
- [She76] ———, *A two-cardinal theorem and a combinatorial theorem*, Proc. Amer. Math. Soc. **62** (1976), no. 1, 134–136 (1977). MR 434800
- [She78a] ———, *Appendix to: “Models with second-order properties. II. Trees with no undefined branches”* (Ann. Math. Logic **14** (1978), no. 1, 73–87), Ann. Math. Logic **14** (1978), 223–226. MR 506531
- [She78b] ———, *Classification theory and the number of nonisomorphic models*, Studies in Logic and the Foundations of Mathematics, vol. 92, North-Holland Publishing Co., Amsterdam-New York, 1978. MR 513226
- [She89] ———, *Consistency of positive partition theorems for graphs and models*, Set theory and its applications (Toronto, ON, 1987), Lecture Notes in Math., vol. 1401, Springer, Berlin, 1989, pp. 167–193. MR 1031773
- [She92] ———, *Strong partition relations below the power set: consistency; was Sierpiński right? II*, Sets, graphs and numbers (Budapest, 1991), Colloq. Math. Soc. János Bolyai, vol. 60, North-Holland, Amsterdam, 1992, arXiv: math/9201244, pp. 637–668. MR 1218224
- [She00] ———, *On what I do not understand (and have something to say), model theory*, Math. Japon. **51** (2000), no. 2, 329–377, arXiv: math/9910158. MR 1747306
- [Sil80] Jack H. Silver, *Counting the number of equivalence classes of Borel and coanalytic equivalence relations*, Ann. Math. Logic **18** (1980), no. 1, 1–28. MR 568914
- [SU19] Saharon Shelah and Danielle Ulrich, *Torsion-free abelian groups are consistently \aleph_2 -complete*, Fund. Math. **247** (2019), no. 3, 275–297, arXiv: 1804.08152. MR 4017015
- [SZ11] Saharon Shelah and Jindřich Zapletal, *Ramsey theorems for product of finite sets with submeasures*, Combinatorica **31** (2011), no. 2, 225–244. MR 2848252
- [Vau65] R. L. Vaught, *A löwenheim-skolem theorem for cardinals far apart*, The Theory of Models (J. V. Addison, L. A. Henkin, and A. Tarski, eds.), North-Holland Publishing Company, 1965, pp. 81–89.

EINSTEIN INSTITUTE OF MATHEMATICS, EDMOND J. SAFRA CAMPUS, GIVAT RAM, THE HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM, 9190401, ISRAEL, AND, DEPARTMENT OF MATHEMATICS, HILL CENTER - BUSCH CAMPUS, RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY, 110 FRELINGHUYSEN ROAD, PISCATAWAY, NJ 08854-8019 USA

Email address: shelah@math.huji.ac.il

URL: <http://shelah.logic.at>