

# THERE ARE JUST FOUR SECOND-ORDER QUANTIFIERS

BY

SAHARON SHELAH

## ABSTRACT

Among the second-order quantifiers ranging over relations satisfying a first-order sentence, there are four for which any other one is bi-interpretable with one of them: the trivial, monadic, permutational, and full second order.

## Introduction

The problem of elementary theories of permutation groups was discussed in Vazhenin and Rasin [12], McKenzie [5], Pinus [7], and essentially solved in Shelah [11]. It became clear that this is equivalent to the problem of the expressive power of the quantifier  $Q_p$ , ranging over permutations. (Of course in rich enough languages it is equivalent to the second-order quantifier, so the interesting case is of languages with no nonlogical symbols.) After examining [11], J. Stavi doubted the naturality of this quantifier, whereas I was convinced that there are no new quantifiers of this kind. At last he suggested, as explication of “this kind”, the family of quantifiers  $Q_\psi$ , where  $\psi = \psi(r)$  is a first-order sentence with the single predicate  $r$ , and  $(Q_\psi r)\phi$  means: “There is a relation  $r$  satisfying  $\psi$  such that  $\phi$ ”... Here we prove that up to bi-interpretability there are really only four such quantifiers. It seems that this justifies the preoccupation with  $Q_p$ . We define interpretability in a way even weaker than in [11]:  $Q_{\psi_1}$  is interpretable in  $Q_{\psi_2}$  if there is a *first-order* formula  $\theta(\bar{x}, y_1 \dots, r_1, \dots)$  such that for any *infinite* set  $A$ , and relation  $R$  over it,  $A \models \psi_1[R]$ , there are elements  $a_1, \dots \in A$  and relations  $S_1, \dots$  over  $A$ ,  $A \models \psi_2[S_i]$ , such that  $A \models (\forall \bar{x}) [R(\bar{x}) \equiv \theta(\bar{x}, a_1, \dots, S_1, \dots)]$ .

Our proofs give somewhat more than what is required. If  $Q_X$  is one of those four quantifiers (see Theorem 2 for details) and  $Q_\psi, Q_X$  are bi-interpretable, then

---

Received October 11, 1972 and in revised form March 5, 1973.

there is a  $\theta(\bar{x}, \bar{y}, r_1, \dots, r_n)$  interpreting  $Q_X$  by  $Q_\psi$  with bounded  $n$  (that is the bound on  $n$  is absolute). No attempt has been made to determine a minimal bound, but notice that if  $Q_\psi, Q_M$  are bi-interpretable ( $Q_M$ —the monadic quantifier) then by Claim 5H, some  $\theta(x, y, r)$  interprets  $Q_M$  by  $Q_\psi$ .

There are several ways in which we can try to generalize our results and most directions were not investigated.

We can quantify over a pair of relations, e.g. two operations defining a field; but this can be reduced to the previous case.

We can permit finite models, but then we can find a quantifier very strong for models with an even number of elements, and trivial for models with an odd number of elements.

We can have quantifiers ranging over pseudo-elementary classes. That is,  $(Q_{\psi(r,s)}r) \dots$ , means “there is an  $r$  such that for some  $s$ ,  $\psi(r, s)$  holds, and  $r$  satisfies  $\dots$ ”. In this case, our proofs give similar classification, but the equivalence classes of  $Q_M, Q_P$  are divided into infinitely many equivalence classes. It is not so difficult to give a complete picture. If we want to find which cardinals can be characterized by a sentence with such quantifiers but with no nonlogical symbol, we are stuck by the independence of, e.g., the function  $2^{\aleph_\alpha}$ .

Another direction is multi-sorted models. Here the classification depends on  $n$ -cardinal theorems (see e.g. [1]) but modulo these, it seems possible to give a classification.

Still another direction is to replace first-order logic by the infinitary logic  $L_{\omega_1, \omega}$  (or  $L_{\lambda, \omega}$ ). Here it is reasonable to ignore models of cardinality  $< \beth_{\omega_1}$ . In this case we have a quantifier  $Q_{II}^\lambda$  ranging over all two-place relations of cardinality  $< \lambda$ , where there is  $\psi \in L_{\omega_1, \omega}$  which has a model of cardinality  $\mu$  iff  $\mu < \lambda$ . We also have the quantifiers ranging over equivalence relations with  $< \lambda$  equivalence classes or with equivalence classes of power  $\leq \mu < \lambda$  for some  $\mu$ , where  $\lambda$  satisfies the condition mentioned for  $Q_{II}^\lambda$ . It is easy to define when a quantifier  $Q_\psi$  is interpretable by a set of quantifiers and hence when a quantifier and set of quantifiers, or two such sets, are bi-interpretable.

**CONJECTURE.** *Any  $Q_\psi$  is bi-interpretable with a finite set consisting of quantifiers mentioned above.*

The following conjecture seems to imply all others. Let  $A$  be a fixed infinite set. For each  $m$ -place relation  $R$  over  $A$  define “ $(Q_R)r \dots$ ” to mean “there is a relation  $r$  over  $A$ ,  $(A, R) \cong (A, r)$  such that  $\dots$ ”

CONJECTURE. Any quantifier  $(Q_R r)$  is bi-interpretable with a finite set of quantifiers  $\{(Q_{E_i} r) : i < n\}$  where  $E_i$  is an equivalence relation over  $A$ .

NOTATION. Let  $r, s, t$  denote predicates (= variables over relations);  $R, S, T$  (the corresponding) relations;  $x, y, z$  individual variables; and  $a, b, c, d$  elements. A bar on any one of them means that it is a finite sequence of this sort. Let  $\phi, \psi, \theta, \chi$  denote formulae, first-order if not stated otherwise.  $\phi = \phi(x_1, \dots, r_1, \dots)$  means that  $x_1, \dots$  include all the free variables of  $\phi$ , and  $r_1, \dots$  include all the predicates in  $\phi$ .  $L$  denotes first-order language (always with equality). Let  $\psi = \psi(r)$  always,  $r$  have  $n(\psi)$  places, and  $L_\psi = L(Q_\psi)$  be language  $L$  with the added second-order quantifier  $(Q_\psi r) \dots$  which means "there is an  $r$  which satisfies  $\psi$  such that  $\dots$ ". Let  $R_\psi(A) = \{R : R \text{ an } n(\psi)\text{-ary relation over } A, A \models \psi[R]\}$  ( $\models$  denotes satisfaction). Let  $(Q_\psi \bar{r})$  mean  $(Q_\psi r_1) \dots (Q_\psi r_n)$ , where  $\bar{r} = \langle r_1, \dots, r_n \rangle$ . We shall write  $\bar{a} \in A$  instead of  $\bar{a} = \langle a_1, \dots, a_n \rangle$ ,  $a_i \in A$ . For any  $\bar{a}$ ,  $l(\bar{a})$  is its length, and  $\bar{a}_i$  or  $a_i$  its  $i$ 'th element, so  $\bar{a} = \langle a_1, \dots, a_{l(\bar{a})} \rangle$ .

Let  $i, j, k, l, m, n$  range over natural numbers,  $i, j, \alpha, \beta, \gamma, \delta$  over ordinals, and  $\lambda, \mu, \kappa$  over cardinals.

A sequence  $\bar{a}$  is without repetitions if  $i \neq j$  implies  $\bar{a}_i \neq \bar{a}_j$ , and  $\bar{a}, \bar{b}$  are disjoint if  $\bar{a}_i \neq \bar{b}_j$  for any  $i, j$ . Let  $\text{Eq}_\lambda(A) [\text{Eq}_\lambda^*(A)]$  be the set of equivalence relations over  $A$ , with each equivalence class having  $< \lambda[\lambda]$  elements. Let  $e$  denote an equivalence relation.

DEFINITION 1.  $Q_{\psi_1}$  is interpretable in  $Q_{\psi_2}$  if there is a formula  $\phi(\bar{x}, \bar{y}, \bar{r})$ ,  $l(\bar{x}) = n(\psi_1)$  such that for any infinite  $A$  and  $R_1 \in R_{\psi_1}(A)$  there are  $\bar{a} \in A$ ,  $\bar{R} \in R_{\psi_2}(A)$  such that

$$A \models (\forall \bar{x}) [R_1(\bar{x}) \equiv \phi(\bar{x}, \bar{a}, \bar{R})].$$

DEFINITION 2.  $Q_{\psi_1}$  and  $Q_{\psi_2}$  are equivalent if each is interpretable in the other.

LEMMA 1. If  $Q_{\psi_1}$  is interpretable in  $Q_{\psi_2}$ , then there is a recursive function  $F$  from the formulae of any language  $L_{\psi_1}$  into those of  $L_{\psi_2}$  such that for any infinite model  $M$  and sentence  $\theta \in L_{\psi_1}$  (not necessarily first-order)

$$M \models \theta \text{ iff } M \models F(\theta).$$

PROOF. We define  $F(\theta)$  for formulae  $\theta$ , by induction on  $\theta$ . The only nontrivial case is  $\theta = (Q_{\psi_1} r)\chi$ . Without loss of generality no variable occurs both in  $\theta$  and in the interpreting formula  $\phi$  (otherwise change names). Replace in  $F(\chi)$  and in

$\psi_1$  every occurrence of  $r(\bar{z})$  by  $\phi(\bar{z}, \bar{y}, \bar{r})$ , call the results  $\chi^*, \psi_1^*$  and let  $F(\theta) = (\exists \bar{y})(Q_{\psi, \bar{r}})(\chi^* \wedge \psi_1^*)$ .

Our main result is

**THEOREM 2.** *Each  $Q_\psi$  is equivalent to exactly one of the following quantifiers:*

A)  $Q_I$ —the trivial quantifier, i.e.,  $Q_{\psi_I}, \psi_I = r, n(\psi_I) = 0$ , so  $L_{\psi_I}$  is just first-order logic

B)  $Q_M$ —the monadic second-order quantifier, i.e.,  $Q_{\psi_M}, \psi_M = (\forall x)[r(x) \equiv r(x)]$ ,  $n(\psi_M) = 1$ ,

C)  $Q_P$ —the permutational second-order quantifier, ranging over permutations of the universe of order two, i.e.  $Q_{\psi_P}$ ,

$$\psi_P = (\forall x)[f(f(x)) = x]$$

(of course we can quantify over functions instead of relations; equivalently we can quantify over  $\text{Eq}_3(A)$ )

D)  $Q_{II}$ —the (full) second-order quantifier i.e.,  $Q_{\psi_{II}}, \psi_{II} = (\forall xy)[r(x, y) \equiv r(x, y)]$ ,  $n(\psi_{II}) = 2$ .

The proof is broken into a series of lemmas and claims.

**LEMMA 3.**  $Q_I$  can be interpreted in  $Q_M$ ,  $Q_M$  can be interpreted in  $Q_P$ , and  $Q_P$  can be interpreted in  $Q_{II}$ . However, none of the converses holds. (In fact, in the negative parts, also the conclusion of Lemma 1 fails.)

**PROOF.** The positive statements are immediate. As for the negative statements, let  $L$  be a language with no predicates or function symbols (except equality, of course), and  $L_{ord}$  be the language of models of order.

We know that in  $L_{ord}(Q_I)$  there is no formula (with parameters) defining the class of well-ordering but that there is one in  $L_{ord}(Q_M)$ . Hence  $Q_M$  cannot be interpreted by  $Q_I$ .

We know that for every sentence  $\phi \in L(Q_M)$ , either every infinite model satisfies it or no infinite model satisfies it. As in McKenzie [5] (or Pinus [7], Shelah [11]) this is not true for  $L(Q_P)$ ;  $Q_P$  cannot be interpreted by  $Q_M$ .

By Shelah [11], if a sentence  $\phi \in L(Q_P)$  has a model of cardinality  $\geq \aleph_{\Omega^\omega}$  ( $\Omega = (2^{\aleph_0})^+$ ) then  $\phi$  has models of arbitrarily high power. Of course  $L(Q_{II})$  does not satisfy this, hence  $Q_{II}$  is not interpretable by  $Q_P$ .

**LEMMA 4.** *If  $Q_\psi$  is not interpretable by  $Q_I$  then  $Q_M$  is interpretable by  $Q_\psi$ .*

**CLAIM 4A.**  $Q_M$  is interpretable by  $Q_\psi$  if there is a formula  $\phi = \phi(x, \bar{y}, \bar{r})$ ,

and a set  $A$ ,  $\bar{a} \in A$ ,  $\bar{R} \in R_\psi(A)$  such that  $\phi(y, \bar{a}, \bar{R})$  divides  $A$  into two infinite sets, that is  $|\phi(A, \bar{a}, \bar{R})| \geq \aleph_0$ ,  $|\neg \phi(A, \bar{a}, \bar{R})| \geq \aleph_0$ , where  $\phi(A, \bar{a}, \bar{R}) = \{b \in A : A \models \phi[b, \bar{a}, \bar{R}]\}$ .

PROOF OF CLAIM 4A. Assuming the existence of such  $\phi$ , by the compactness and Lowenheim-Skolem theorems, for every infinite  $B$  there are  $\bar{a} \in B$ ,  $\bar{R} \in R_\psi(B)$  such that  $|B| = |\phi(B, \bar{a}, \bar{R})| = |\neg \phi(B, \bar{a}, \bar{R})|$ . By applying a permutation of  $B$  for every  $B_1 \subseteq B$ ,  $|B_1| = |B - B_1| = |B|$ , there are  $\bar{a} \in A$ ,  $\bar{R} \in R_\psi(B)$  such that  $\phi(B, \bar{a}, \bar{R}) = B_1$ . Now for every  $C \subseteq B$  there are  $B_i \subseteq B$   $i = 1, \dots, 4$  such that  $|B_i| = |B - B_i| = |B|$  and  $C = (B_1 \cap B_2) \cup (B_3 \cap B_4)$ . Let

$$\theta = \theta(x, \bar{y}^*, \bar{r}^*) = [\phi(x, \bar{y}^1, \bar{r}^1) \wedge \phi(x, \bar{y}^2, \bar{r}^2)] \vee [\phi(x, \bar{y}^3, \bar{r}^3) \wedge \phi(x, \bar{y}^4, \bar{r}^4)].$$

Then as the  $\bar{a}_i^*$  range over  $B$ , and the  $\bar{R}_i^*$  range over  $R_\psi(B)$ ,  $\theta(B, \bar{a}^*, \bar{R}^*)$  ranges over the subsets of  $B$ .

DEFINITION 3.

A) The sequences  $\bar{a}^1, \bar{a}^2$  are similar over  $B$  if  $\bar{a}^i = \langle \dots, \bar{a}_j^i, \dots \rangle_{j < k}$  and (i)  $\bar{a}_j^1 = \bar{a}_j^2$  iff  $\bar{a}_j^1 = \bar{a}_j^2$ ; (ii) for  $b \in B$ ,  $\bar{a}_j^1 = b$  iff  $\bar{a}_j^2 = b$ .

B) The sequences  $\bar{a}^1, \bar{a}^2$  are similar over  $\bar{b}$  iff they are similar over  $\{\dots, \bar{b}_i, \dots\}$ .

CLAIM 4B. If  $Q_M$  is not interpretable by  $Q_\psi$  then for every formula  $\phi(\bar{x}, \bar{y}, \bar{r})$  there is a formula  $\theta(z, \bar{y}, \bar{r})$  and  $n < \omega$  such that for any  $A$ ,  $\bar{b} \in A$ ,  $\bar{R} \in R_\psi(A)$

(i)  $A \models (\exists^{\leq n} z)\theta(z, \bar{b}, \bar{R})$  that is  $|\theta(A, \bar{b}, \bar{R})| \leq n$

(ii) if  $\bar{a}^1, \bar{a}^2$  are similar over

$$\{\dots, \bar{b}_i, \dots\} \cup \theta(A, \bar{b}, \bar{R}) \text{ then } A \models \phi(\bar{a}^{-1}, \bar{b}, \bar{R}) \equiv \phi(\bar{a}^2, \bar{b}, \bar{R}).$$

REMARK. In the induction step, only the validity of our claim for the previous case is needed.

PROOF OF CLAIM 4B. We shall prove it by induction on  $l(\bar{x})$ .

For  $l(\bar{x}) = 1$  by Claim 4A (and compactness) for some  $m$ ,

$$\theta_m = [(\exists^{\leq m} x)\phi(x, \bar{y}, \bar{r}) \rightarrow \phi(z, \bar{y}, \bar{r})] \wedge [(\exists^{\leq m} x)\neg \phi(x, \bar{y}, \bar{r}) \rightarrow \neg \phi(z, \bar{y}, \bar{r})]$$

satisfies our demands.

Suppose we have proved it for  $l(\bar{x}) \leq l$ , and we shall prove it for the case  $l(\bar{x}) = l + 1$ . Choose any  $A$ ,  $\bar{b} \in A$ ,  $\bar{R} \in R_\psi(A)$  and  $\bar{x} = \langle x_1, \dots, x_{l+1} \rangle$ ,  $\bar{x}^1 = \langle x_1, \dots, x_l \rangle$ ,  $\bar{y}^1 = \langle x_{l+1}, \bar{y}_1, \dots \rangle$ . For  $\phi(\bar{x}^1, \bar{y}^1, \bar{r})$  we have proved the claim, and let  $\theta(z, \bar{y}^1, \bar{r})$ ,  $n$  be as mentioned there. Now for any  $a \in A$  let  $Ex(a) = \theta(A, a, \bar{b}, \bar{R}) - \{a, \dots, \bar{b}_i, \dots\}$ . Thus  $|Ex(a)| \leq n$  always.

Let us show that  $\cup_{a \in A} Ex(a)$  is finite. If not, define by induction on  $i < \omega$ ,  $a_i \in A - \{a_j : j < i\}$ ,  $c_i$  such that  $Ex(a_i) \not\subseteq \cup_{j < i} Ex(a_j)$ , and  $c_i \in Ex(a_i) - \cup_{j < i} Ex(a_j)$ . By Ramsey's theorem we can assume (by replacing the sequence of  $a_i$ 's and  $c_i$ 's by a subsequence) that the truth value of  $c_i \in Ex(a_j)$  depends only on whether  $i = j$ ,  $i < j$  or  $i > j$ . Clearly  $c_i \in Ex(a_i)$ , and for  $j > i$ ,  $j < \omega$ ,  $c_j \notin Ex(a_i)$ . Since  $|Ex(a_j)| \leq n$ , clearly there is an  $i < n + 2$  such that  $c_i \notin Ex(a_{n+2})$ . Hence  $c_i \in Ex(a_j)$  iff  $i = j$ . Similarly  $c_i = c_j$  iff  $i = j$ ; and  $a_i \neq c_j$ . As the  $a_i$ 's and  $c_i$ 's are distinct, we can assume that none of them appear in  $\bar{b}$ . Let  $f$  be a permutation of  $A$  which interchanges  $c_{3i+1}$  with  $c_{3i+2}$ , and takes the other elements of  $A$  to themselves. Let  $\bar{R}^*$  be the image of  $\bar{R}$  by  $f$  (so  $f$  is an isomorphism from  $(A, \bar{R})$  onto  $(A, \bar{R}^*)$ ). Clearly  $A \models (\forall x) [\theta(x, a_i, \bar{b}, \bar{R}) \equiv (x, a_i, \bar{b}, \bar{R}^*)]$  iff  $f$  takes the set  $\theta(A, a_i, \bar{b}, \bar{R})$  onto itself iff  $i$  is divisible by three; thus

$$\chi(y, \bar{b}, \bar{R}, \bar{R}^*) = (\forall x) [\theta(x, y, \bar{b}, \bar{R}) \equiv \theta(x, y, \bar{b}, \bar{R}^*)]$$

satisfies the conditions mentioned in Claim 4A, a contradiction. Hence  $C = \cup_{a \in A} Ex(a)$  is finite. Let  $C = \{c_1, \dots, c_j\}$ ,  $\bar{c} = \langle c_1, \dots, c_j \rangle$ .

DEFINITION 4. Let us call  $\chi(\bar{z})$  complete if it is a conjunction such that for every  $i, j$ ,  $z_i = z_j$  or  $z_i \neq z_j$  is a conjunct (and all the conjuncts are of this form).

Let  $\chi_i(\bar{x}^1, x, \bar{y}, \bar{z})$   $i = 1, \dots, k$  be a list of all complete formulae in the displayed variables. By definition of  $Ex$  for every  $i$ , and  $a \in A$

$$(i) A \models (\forall \bar{x}^1) [\chi_i(\bar{x}^1, a, \bar{b}, \bar{c}) \rightarrow \phi(\bar{x}^1, a, \bar{b}, \bar{R})]$$

or

$$(ii) A \models (\forall \bar{x}^1) [\chi_i(\bar{x}^1, a, \bar{b}, \bar{c}) \rightarrow \neg \phi(\bar{x}^1, a, \bar{b}, \bar{R})].$$

For each  $a$  let  $I(a)$  be the set of  $i$ 's for which (i) holds.

By Claim 4A, except for finitely many  $a$ 's, all  $I(a)$  are equal (to I). Let  $C^1$  be the set of exceptional  $a$ 's. It is easy to check that:

(\*) if  $\bar{a}^1, \bar{a}^2$  are similar over  $C^2 = \{\dots, \bar{b}_i, \dots\} \cup C \cup C^1$ , then  $A \models \phi[\bar{a}^1, \bar{b}, \bar{R}] \equiv \phi[\bar{a}^2, \bar{b}, \bar{R}]$ ;  $C^2$  is finite.

Without loss of generality we cannot replace  $C^2$  by a set of smallest cardinality satisfying (\*). Let  $n_1 = |C^2|$ , and let  $\theta_1 = \theta_1(z, \bar{y}, \bar{r})$  say that there are  $z_2, \dots, z_{n_1}$ , such that if  $\bar{x}^1, \bar{x}^2$  are similar over  $\{z, z_2, \dots, z_{n_1}, \dots, \bar{y}_i, \dots\}$ , then  $\phi(\bar{x}^1, \bar{y}, \bar{r}) \equiv \phi(\bar{x}^2, \bar{y}, \bar{r})$ .

SUBCLAIM 4C.  $\theta_1(A, \bar{b}, \bar{R})$  is finite.

PROOF OF SUBCLAIM 4C. If not, there are distinct  $C_i^2$ ,  $i < \omega$  satisfying (\*),  $|C_i^2| = n_1$ .

Now w.l.o.g. there is a  $C^*$ ,  $|C^*| < n_1$ , such that for any  $i < j < \omega$ ,  $C_i^2 \cap C_j^2 = C^*$ ; this follows by Erdős and Rado [2], but we can also prove it directly. Let  $C_i^2 = \{c_{i,1}^2, \dots, c_{i,n_1}^2\}$ , and by Ramsey's theorem [9] there is an infinite  $I \subseteq \omega$ , such that for  $1 \leq l, k \leq n_1$ ,  $i < j \in I$ , the truth value of  $c_{i,l}^2 = c_{j,k}^2$  does not depend on the particular  $i, j$ . Without loss of generality  $I = \omega$ . Let

$$C^* = \{c_{0,k}^2 : c_{0,k}^2 = c_{1,k}^2, 1 \leq k \leq n_1\}.$$

By definition of  $I$ ,  $C^* \subseteq C_i^2$  for every  $i$ . As  $C_0^2 \neq C_1^2$ ,  $|C^*| < n_1$ . Let  $i < j < \omega$ . Then clearly  $C^* \subseteq C_i^2 \cap C_j^2$ ; if equality does not hold let  $c \in C_i^2 \cap C_j^2 - C^*$ . Thus  $c = c_{i,k}^2 = c_{j,l}^2$ ; since  $i < j$ , this implies  $c_{0,k}^2 = c_{2,l}^2$ ,  $c_{1,k}^2 = c_{2,l}^2$ ,  $c_{0,k}^2 = c_{j,l}^2$ . Hence  $c_{0,k}^2 = c_{1,k}^2 = c_{j,l}^2 = c$ ,  $c_{0,k}^2 \in C^*$ , and  $c \in C^*$ , a contradiction. So it is proved that w.l.o.g. there is such a  $C^*$ , but if  $\bar{a}^1, \bar{a}^2$  are similar over  $C^*$  then they are similar over all  $C_i^2$  except finitely many, and this contradicts the definition of  $n_1$ . Thus Subclaim 4C is proved.

CONTINUATION OF THE PROOF OF CLAIM 4B. Let  $|\theta_1(A, \bar{b}, \bar{R})| = n_2$ .

So  $\theta_1(z, \bar{y}, \bar{r})$ ,  $n_2$  satisfy the demands in Claim 4B except that they depend on  $A, \bar{b}, \bar{R}$ . By the compactness theorem there are  $\theta^i(z, \bar{y}, \bar{r})$ ,  $n^i$   $i = 1, \dots, k (< \omega)$  such that for any  $A, \bar{b} \in A$ ,  $\bar{R} \in R_\psi(A)$  there is an  $i$  such that  $\theta^i, n^i$  satisfy the demands of the claim. Let  $\theta^* = \theta^*(z, \bar{y}, \bar{r}) = \bigvee_i [(\exists \leq n^i u) \theta^i(u, \bar{y}, \bar{r}) \wedge \theta^i(z, \bar{y}, \bar{r})]$ . Clearly this is the right one, so Claim 4B is proved.

PROOF OF LEMMA 4. Assume  $Q_M$  is not interpretable by  $Q_\psi$ . Use Claim 4B for  $\phi(\bar{x}, r) = r(\bar{x})$ , and let  $\theta, n$  be the  $\theta, n$  whose existence is proved there. Let  $\chi_i(\bar{x}, \bar{z})$  ( $l(\bar{z}) = n$ )  $i = 1, \dots, k$  be the complete formulae mentioned in the proof of Claim 4B. Let  $I_1, \dots, I_{2^k}$  be the subsets of  $\{1, \dots, k\}$ .

Let

$$\phi^*(\bar{x}, \bar{y}, \bar{z}) = \bigwedge_j [y_{2j} = y_{2j+1} \rightarrow \bigvee_{i \in I_j} \chi_i(\bar{x}, \bar{z})].$$

For an infinite  $A$ , for every  $\bar{R} \in R_\psi(A)$  let  $\{c_1, \dots, c_n\} \cong \theta(A, \bar{R})$ .

Let  $I = \{i : (\exists \bar{x}) [\chi_i(\bar{x}, \bar{c}) \wedge r(\bar{x})]\}$ ,  $j$  be such that  $I = I_j$ . Define  $\bar{b}$  such that  $\bar{b}_{2p} = \bar{b}_{2p+1}$  iff  $p = j$ . Then

$$A \models \phi^*(\bar{x}, \bar{b}, \bar{c}) \equiv r(\bar{x}),$$

a contradiction. Thus Lemma 4 is proved.

LEMMA 5. If  $Q_\psi$  is not interpretable by  $Q_M$  then  $Q_P$  is interpretable by  $Q_\psi$ .

PROOF. Clearly  $Q_\psi$  is a *fortiori* not interpretable by  $Q_I$ , hence by Lemma 4,  $Q_M$  is interpretable by  $Q_\psi$ .

CLAIM 5A.  $Q_P$  is interpretable by  $Q_\psi$  if there is a formula  $\phi(x, y, \bar{z}, \bar{r})$ , a set  $A$ ,  $\bar{c} \in A$ ,  $\bar{R} \in R_\psi(A)$ ,  $B \subseteq A$  such that  $\phi(x, y, \bar{c}, \bar{R})$  defines on  $B$  an equivalence relation with infinitely many equivalence classes with  $\geq 2$  elements.

PROOF OF CLAIM 5A. The proof is similar to that of Claim 4A. By replacing  $B$  by a subset, we may assume that each equivalence class has exactly two elements and that  $A - B$  is infinite. Now for every infinite  $A$ , by the compactness and the Lowenheim-Skolem theorems, there are  $B \subseteq A$ ,  $\bar{a} \in A$ ,  $\bar{R} \in R_\psi(A)$ , such that  $|B| = |A - B| = |A|$ , and  $\phi(x, y, \bar{a}, \bar{R})$  defines on  $B$  a relation  $\in \text{Eq}_2^*(B)$ . We can easily find  $\bar{b} \in A$ ,  $\bar{S} \in R_\psi(A)$  such that  $\phi(x, y, \bar{b}, \bar{S})$  defines on  $A - B$  an equivalence relation from  $\text{Eq}_2^*(A - B)$ . Also there is a formula  $\phi^*(x, \bar{c}, \bar{T})$   $\bar{c} \in A$ ,  $\bar{T} \in R_\psi(A)$ , which defines  $B$ . So

$$\begin{aligned} \theta(x, y, \bar{a}, \bar{b}, \bar{c}, \bar{R}, \bar{S}, \bar{T}) = & [\phi^*(x, \bar{c}, \bar{T}) \equiv \phi^*(y, \bar{c}, \bar{T})] \\ & \wedge [\phi^*(x, \bar{c}, \bar{T}) \rightarrow \phi(x, y, \bar{a}, \bar{R})] \wedge [\neg \phi^*(x, \bar{c}, \bar{T}) \rightarrow \phi(x, y, \bar{b}, \bar{S})] \end{aligned}$$

defines a relation from  $\text{Eq}_2^*(A)$ .

Clearly for every  $e \in \text{Eq}_2^*(A)$  there are  $\bar{a}', \bar{b}', \bar{c}' \in A$ ,  $\bar{R}', \bar{S}', \bar{T}' \in R_\psi(A)$ , such that

$$A \models (\forall xy) [\theta(x, y, \bar{a}, \dots) \equiv xey].$$

Since we can interpret  $Q_M$  in  $Q_\psi$ , by a small change in  $\theta$  we can have the same for  $e \in \text{Eq}_3(A)$ . This proves the claim.

DEFINITION 5. We call  $\phi = \phi(x_1, \dots, x_n, r)$  atomic if  $\phi = [x_i = x_j]$  or  $\phi = r(x_{i_1}, \dots, x_{i_{n(\psi)}})$ .

DEFINITION 6. For every  $A$ ,  $B \subseteq A$ ,  $R \in R_\psi(A)$ , define the equivalence relation  $e = e(R, B, A)$  over  $B$  by  $bec$  iff  $b, c \in B$ , and for every atomic  $\phi(x_1, \dots, x_n)$  and  $a_2, \dots, a_n \in A - B$ ,  $A \models \phi[b, a_2, \dots, R] \equiv \phi[c, a_2, \dots, R]$ .

CLAIM 5B.  $e(R, B, A)$  is defined by a formula in  $A$  (with  $R$  and  $B$  as parameters).

PROOF. Immediate.

CLAIM 5C. If  $Q_P$  is not interpretable by  $Q_\psi$ , then for every  $A$ ,  $B \subseteq A$ ,  $R \in R_\psi(A)$ ,  $e(R, B, A)$  has finitely many equivalence classes.

PROOF. Suppose  $e(R, B, A)$  has infinitely many equivalence classes. By Claim 5A, only finitely many of them have  $\geq 2$  elements. But if we replace  $B$  by a smaller set,  $e(R, B, A)$  becomes finer (i.e., the equivalence classes become smaller). Hence



w.l.o.g. each equivalence class of  $e(R, B, A)$  has one element, and of course  $B$  is infinite.

Let  $f$  be a permutation of order two of  $A$ , such that  $f(a) = a \leftrightarrow a \notin B$ . Define

$$R_1 = \{\langle a_1, \dots \rangle : a_1, \dots \in A, \langle f(a_1), \dots \rangle \in R\}.$$

Let

$$e_1 = \{\langle c, b \rangle : b, c \in B, \text{ for every atomic } \phi(x, \bar{y}, r) \text{ and} \\ \text{every } \bar{a} \in (A - B); A \models \phi[c, \bar{a}, R] \equiv \phi[b, \bar{a}, R_1] \\ A \models \phi[b, \bar{a}, R] \equiv \phi[c, \bar{a}, R_1]\}.$$

It is easy to see that  $c = f(b)$ ,  $c, b \in B$  implies  $\langle c, b \rangle \in e_1$ . It is easy to check that  $\langle c, b \rangle \in e_1$  implies  $\langle c, f(b) \rangle \in e(R_1, B, A)$  but this implies  $c = f(b)$ .

Hence  $[\langle x, y \rangle \in e_1] \vee x = y$  defines an equivalence relation of  $\text{Eq}_2^*(B)$ , and clearly it is definable by a formula. By Claim 5A this leads to a contradiction, hence 5C is proved.

**CLAIM 5D.** *If  $Q_P$  is not interpretable by  $Q_\psi$ , then there is a formula  $\phi(x, y, r)$  such that for every  $A, R \in R_\psi(A)$ .*

(i)  $\phi(x, y, R)$  defines an equivalence relation with finitely many equivalence classes.

(ii)  $A \models \phi[a, b, R]$  implies that there is a finite  $B$  such that  $\langle a, b \rangle \in e(R, B, A)$ .

**PROOF.** Define for  $A, R \in R_\psi(A)$   $n < \omega$  the relation

$$e_n(R, A) = \{\langle c, b \rangle : c, b \in A, \text{ there is } B \subseteq A, |B| \leq n$$

such that  $\langle c, b \rangle \in e(R, B, A)\}$ .

Define  $\phi_n(x, y, r)$  such that  $A \models \phi_n[c, b, R]$  iff  $\langle c, b \rangle \in e_n(R, A)$ ,  $R \in R_\psi(A)$ . Note that  $\phi_{n+1}(x, y, r) \rightarrow \phi_n(x, y, r)$  always.

Clearly  $e^*(R, A) = \cup_{n < \omega} e_n(R, A)$  is an equivalence relation over  $A$ . Moreover it has only finitely many equivalence classes. Otherwise choose nonequivalent  $a_i$   $1 \leq i < \omega$ . By Claim 5C and the compactness theorem, there is  $n_0 < \omega$  such that  $e(R^1, B, A)$  always has  $\leq n_0$  equivalence classes, for  $B \subseteq A$ ,  $R^1 \in R_\psi(A)$ . Let  $B = \{a_i : 1 \leq i \leq n_0 + 1\}$ . Then  $e(R, B, A)$  has  $n_0 + 1$  equivalence classes (by the choice of the  $a_i$ 's and the definition of  $e^*$ ). We prove in fact that  $e^*(R, A)$  has  $\leq n_0$  equivalence classes for any  $R \in R_\psi(A)$ . Hence in

$$\Gamma = \{\psi(r)\} \cup \{\neg \phi_n(x_i, x_j, r) : n < \omega, 1 \leq i < j \leq n_0 + 1\}$$

there is a contradiction.

Thus for some  $n_1 < \omega$  there is a contradiction in

$$\{\psi(r)\} \cup \{\neg \phi_n(x_i, x_j, r) : n < n_1, 1 \leq i < j \leq n_0 + 1\}.$$

The closure of  $\phi_{n_1}(x, y, r)$  to an equivalence relation is

$$\phi(x, y, r) = \text{df} (\exists z_1, \dots, z_m) \left[ \bigwedge_{i=1}^m \phi_{n_1}(z_i, z_{i+1}, r) \wedge z_0 = x \wedge z_m = y \right]$$

where  $m = 3n_0$  is sufficient. This is because for every  $A, R \in R_\psi(A)$  there is a maximal set  $\{a_i : 1 \leq i < i_0\}$  such that  $i < j < i_0$  implies  $A \vDash \neg \phi_{n_1}(a_i, a_j, R)$ ; hence  $i_0 \leq n_0$  by the definition of  $n_1$ . By the maximality of the set, for every  $a \in A$  for at least one  $i$   $A \vDash \phi_{n_1}(a, a_i, R)$ . Now if  $b, c$  are equivalent in the closure of  $e_{n_1}(R, A)$  then there are  $d_1, \dots, d_m$ ,  $d_1 = b$ ,  $d_m = c$  and  $\langle d_i, d_{i+1} \rangle \in e_{n_1}(R, A)$ . Choose such  $d_i$ 's with minimal  $m$ ; we should show  $m \leq 3n_0$ . For this it suffices to prove there are no four  $d_i$  from one  $\phi_{n_1}(A, a_i, R)$ . Let  $1 \leq i_1 < i_2 < i_3 < i_4 \leq m$ ,  $d_{i_1}, \dots, d_{i_4} \in \phi_{n_1}(A, a_j, R)$ . Then  $\langle d_{i_1}, a_j \rangle, \langle a_j, d_{i_4} \rangle \in e_{n_1}(R, A)$ , hence also  $d_1, \dots, d_{i_1}, a_j, d_{i_4}, \dots, d_m$  is a suitable sequence, and it has smaller length, a contradiction.

Since  $e^*(R, A)$  is an equivalence relation, it refines the closure of  $e_{n_1}(R, A)$ . Hence  $R \in R_\psi(A)$ ,  $A \vDash \phi[b, c, R]$  implies that there is a finite  $B \subseteq A$  such that  $\langle b, c \rangle \in e(R, B, A)$ .

CLAIM 5E. In Claim 5D we conclude also that there are  $\theta(z, x, y, r)$ ,  $n_2 < \omega$  such that for any  $A, R \in R_\psi(A)$ ,  $b, c \in A$ ,

- (i)  $A \vDash (\forall xy)(\exists^{\leq n_2} z) \theta(z, x, y, R)$
- (ii)  $A \vDash (\forall xyz)[\theta(z, x, y, R) \rightarrow z \neq x \wedge z \neq y]$
- (iii)  $A \vDash \phi[b, c, R]$  implies  $\langle b, c \rangle \in e(R, B, A)$  where  $B = \theta(A, b, c, R) \cup \{b, c\}$
- (iv)  $A \vDash \neg \phi(b, c, R)$  implies  $A \vDash (\forall z) \neg \theta(z, b, c, R)$ .

PROOF. By the compactness theorem and Claim 5D, there is an  $n_3 < \omega$  such that  $R \in R_\psi(A)$ ,  $A \vDash \phi[b, c, R]$  implies  $\langle b, c \rangle \in e_{n_3}(R, A)$ .

Let  $\theta(z, x, y, r)$  say “ $\phi(x, y, r)$ ,  $z \neq x$ ,  $z \neq y$  and for some  $n \leq n_3$  there are no  $z_1, \dots, z_{n-1}$  such that  $\langle x, y \rangle \in e(r, \{x, y, z_1, \dots, z_{n-1}\})$ , but there are  $z_1, \dots, z_n$  such that  $\langle x, y \rangle \in e(r, \{x, y, z_1, \dots, z_n\})$ , and  $z = z_1$ ”. As in the proof of Claim 4C for all  $R \in R_\psi(A)$ ,  $b, c \in A$ ,  $\theta(A, b, c, R)$  is finite, and so clearly the claim holds.

CLAIM 5F. In the conclusion of Claim 5E we can add

- (v) there is  $n_4 < \omega$  such that for  $R \in R_\psi(A)$

$$A \vDash (\exists^{\leq n_4} z) (\exists xy) \theta(z, x, y, R).$$

For this it suffices to prove Claim 5G (by applying Claim 5G twice we get Claim 5F).

CLAIM 5G. *If  $Q_P$  is not interpretable by  $Q_\psi$ , and for any  $R \in R_\psi(A)$ ,  $A \models (\forall \bar{x})(\forall y)(\exists^{\leq m_1} z) \theta(z, y, \bar{x}, R)$  and  $\theta(z, y, \bar{x}, r) \rightarrow z \neq y$ , then for some  $m_2 < \omega$ , for every  $R \in R_\psi(A)$*

$$A \models (\forall \bar{x})(\exists^{\leq m_2} z)(\exists y) \theta(z, y, \bar{x}, R).$$

PROOF. If not, by the compactness theorem, there are  $A, R \in R_\psi(A)$ ,  $\bar{a} \in A$  such that

$$(1) \quad A \models (\forall y)(\exists^{\leq m_1} z) \theta(z, y, \bar{a}, R)$$

$$(2) \text{ for every finite } B \subseteq A \text{ there are } b \in A, c \in A - B, A \models \theta(c, b, \bar{a}, R).$$

Define by induction on  $n$ ,  $b_n \in A, c_n \in A - \{c_i : i < n\}$  such that  $A \models \theta[c_n, b_n, \bar{a}, R]$ .

By Ramsey's theorem [9] we can assume that the truth value of  $A \models \theta[c_m, b_n, \bar{a}, R]$ ,  $b_n = c_m$  depends only on whether  $m = n$ ,  $m < n$  or  $m > n$ . Since,  $A \models (\exists^{\leq m_1} z) \theta(z, b_n, \bar{a}, R)$  clearly  $A \models \theta[c_m, b_n, \bar{a}, R]$  if  $m = n$  (recall that the  $c_n$ 's are distinct); therefore,  $b_n$ 's are distinct. Also  $b_n \neq c_m$  because (1) if  $n = m$ , this holds by the assumption on  $\theta$ , (2) if  $n < m$ , then  $c_1 = b_0 = c_2$ , a contradiction, and (3) if  $n > m$ ,  $c_1 = b_3 = c_2$ , a contradiction.

Also w.l.o.g.  $b_n \neq \bar{a}_i, c_n \neq \bar{a}_i, \models \neg \theta[c_n, c_m, \bar{a}, R] \wedge \neg \theta(b_n, b_m, \bar{a}, R]$  for  $n \neq m$  (otherwise omit finitely many  $\langle c_i, b_i \rangle$ 's). Let

$$B = \{b_n : n < \omega\} \cup \{c_n : n < \omega\}.$$

Now the formula  $y = z \vee \theta(z, y, \bar{a}, R) \vee \theta(y, z, \bar{a}, R]$  defines on  $B$  a relation of  $\text{Eq}_2^*(B)$ , a contradiction. Thus Claim 5G, and hence Claim 5F are proved.

CLAIM 5H. *If  $Q_P$  is not interpretable by  $Q_\psi$ , then for every  $A, R \in R_\psi(A)$ ,  $e^+(R, A) = \{\langle a, b \rangle : a, b \in A, \text{ the permutation } f (f(a) = b, f(b) = a, f(c) = c \text{ for } c \neq a, b) \text{ is an automorphism of } (A, R)\}$  is an equivalence relation with finitely many equivalence classes.*

PROOF. Define by induction on  $n, 1 \leq n < \omega$ , formulae

$$\phi_n(x, y, r), \theta_n(z, r) \text{ such that}$$

1) for any  $R \in R_\psi(A)$ ,  $\phi_n(x, y, R)$  is an equivalence relation with  $< k_1(n) < \omega$  equivalence classes

2) for any  $R \in R_\psi(A)$ ,  $|\theta_n(A, R)| \leq k_2(n) < \omega$

3) for any  $R \in R_\psi(A)$ ,  $a, b \in A$ ,  $A \models \phi_n[a, b, R]$  implies  $\langle a, b \rangle \in e(R, (B_n - B_{n-1}) \cup \{a, b\}, A)$

4) for any  $1 \leq n \leq m < \omega$ ,  $\theta_n(A, R) \subseteq \theta_m(A, R)$  where  $B_0 = \emptyset$ ,  $B_n = \theta_n(A, R)$ .

For  $n = 1$  the existence of  $\phi_1, \theta_1$  follows from Claims 5D, 5E, and 5F and the compactness theorem. (Take  $\phi_1 = \phi$ ,  $\theta_1 = (\exists xy)\theta(z, x, y, r)$ .)

Suppose  $\phi_n \theta_n$  are defined. Let  $c_1, \dots, c_k [k = \sum_{i=1}^n k_2(l)]$  be individual constants, and replace  $\psi(r)$  by

$$\psi(r) \wedge (\forall z) \left[ \bigvee_{i=1}^n \theta_i(z, r) \equiv \bigvee_{i=1}^k z = c_i \right].$$

Now repeat the proof of Claims 5D, E and F (the change from  $r$  to  $r$  and  $c$ 's is technical; just add more atomic formulae). Hence we get  $\phi_{n+1} \theta_{n+1}$  as we got  $\phi_1 \theta_1$ . Clearly (1), (2) and (3) hold.

Now for any  $R \in R_\psi(A)$  define

$$e' = \{ \langle a, b \rangle : (\forall n < \omega) A \models \phi_n[a, b, R] \}.$$

Clearly  $e'$  is an equivalence relation with  $\leq 2^{\aleph_0}$  equivalence classes.

It is also clear that  $e^+(R, A)$  is an equivalence relation. We shall now show that if  $a e' b$ ,  $a, b \notin \cup_n B_n$  and their  $e'$ -equivalence class is infinite, then  $a e^+(R, A) b$ .

This implies that  $e^+(R, A)$  has  $\leq 2^{\aleph_0}$  equivalence classes, hence by the compactness theorem this is sufficient. For proving that the permutation interchanging  $a, b$  is an automorphism, it suffices to prove that if  $\phi(x, y, z_1, \dots, z_m; r)$  is atomic,  $c_1, \dots, c_m \in A - \{a, b\}$ ,  $\models \phi(a, b, c_1, \dots, c_m, r) \equiv \phi(b, a, c_1, \dots, c_m)$ . We can choose  $n$  such that  $(B_{n+1} - B_n) \cap \{c_1, \dots, c_m, a, b\} = \emptyset$  and  $a_1$  such that  $a_1 e' a$ ,  $a_1 \notin B_{n+1} \cup \{c_1, \dots, c_m, a, b\}$ . By (3)

$$\models \phi[a, b, c_1, \dots, c_m, r] \equiv \phi[a_1, b, c_1, \dots, c_m, r],$$

$$\models \phi[a_1, b, c_1, \dots, c_m, r] \equiv \phi[a_1, a, c_1, \dots, c_m, r] \text{ and also}$$

$$\models \phi[a_1, a, c_1, \dots, c_m, r] \equiv \phi[b, a, c_1, \dots, c_m, r]. \text{ Combining we get the result.}$$

**PROOF OF LEMMA 5.** From Claim 5H and the compactness theorem, it follows that if  $Q_p$  is not interpretable by  $Q_\psi$  then there is some  $n_5 < \omega$  such that for any  $A, R \in R_\psi(A)$ ,  $e^+(R, A)$  has  $\leq n_5$  equivalence classes. Let us show that this implies that  $Q_\psi$  is interpretable by  $Q_M$ . This implies that for every  $A, R \in R_\psi(A)$ , there are sets  $B_1, \dots, B_{n_5}$  (the  $e^+(R, A)$  equivalence classes) such that the truth value of  $R[a_1, \dots, a_{n_5}]$  ( $a_i \in A$ ) depends only on the truth values of  $a_i = a_j$ ,  $a_i \in B_k$ ; hence there is a (quantifier free) formula  $\phi$  such that

$$A \models (\forall \bar{x}) [R(\bar{x}) \equiv \phi(\bar{x}, B_1, \dots, B_{n_5})].$$

From the construction, the number of possible  $\phi$ 's is finite, and let them be  $\phi_1, \dots, \phi_{n_s}$ . Let

$$\phi^* = \bigwedge_{i=1}^n [y_0 = y_i \rightarrow \phi_i(\bar{x}_1, X_1, \dots, X_{n_s})]$$

( $X_i$ -variables over sets).

Hence for every infinite  $A$ , and  $R \in R_\psi(A)$  there are  $c_0, \dots, c_{n_s}, B_1, \dots, B_{n_s}$  such that

$$A \models (\forall \bar{x}) [R(\bar{x}) = \phi^*(\bar{x}, \bar{c}, B_1, \dots)].$$

Thus the proof of Lemma 5 is complete.

LEMMA 6. *If  $Q_\psi$  is not interpretable by  $Q_P$  then  $Q_{II}$  is interpretable by  $Q_\psi$ .*

PROOF. As  $Q_\psi$  is not interpretable by  $Q_P$ , it is obviously not interpretable by  $Q_M$ ; hence by Lemma 5,  $Q_P$  is interpretable by  $Q_\psi$ .

DEFINITION 7.

1) A family of sequences of length  $n$  is pseudofinite if there is a finite set such that in every sequence of the family appears an element from the finite set.

2) A family  $F$  of sequences of length  $n$  from a model  $(A, \bar{R})$  is  $\phi(\bar{x}, \bar{y}, \bar{r})$ -minimal in  $(A, \bar{R})$  ( $l(\bar{x}) = n$ ) if it is not pseudo-finite, but for any  $\bar{a} \in A$ ,  $\{\bar{b} \in F: A \models \phi[\bar{b}, \bar{a}, \bar{R}]\}$  is pseudo-finite or  $\{\bar{b} \in F: A \models \neg \phi(\bar{b}, \bar{a}, \bar{R})\}$  is pseudo-finite.

3)  $\phi(x, \bar{a}, \bar{R})$  is algebraic (in  $(A, \bar{R})$ ) if  $|\phi(A, \bar{a}, \bar{R})| < \aleph_0$ .

4)  $\phi(\bar{x}, \bar{a}, \bar{R})$  is pseudo-algebraic (in  $(A, \bar{R})$ ) if  $\{\bar{b} \in A: A \models \phi[\bar{b}, \bar{a}, \bar{R}]\}$  is pseudo-finite.

5)  $a(\bar{a})$  is (pseudo-) algebraic over  $B$  in  $(A, \bar{R})$  if for some (pseudo-)algebraic  $\phi(x, \bar{b}, \bar{R})$  ( $\phi(\bar{x}, \bar{b}, \bar{R})$ ),  $A \models \phi[a, \bar{b}, \bar{R}]$  ( $A \models \phi[\bar{a}, \bar{b}, \bar{R}]$ ) and  $\bar{b} \in B$ .

6) The type of  $\bar{b}$  over  $B$  in  $(A, \bar{R})$  is  $\{\phi(\bar{x}, \bar{c}, \bar{R}): \bar{c} \in B, A \models \phi[\bar{b}, \bar{c}, \bar{R}]\}$ .

CLAIM 6A.  *$Q_{II}$  is interpretable by  $Q_\psi$  if there are  $\phi(\bar{x}, \bar{y}, \bar{z}, \bar{r})$  [ $l(\bar{x}) = l(\bar{y}) = n$ ],  $A, \bar{R} \in R_\psi(A)$ ,  $\bar{c} \in A$ ,  $B \subseteq A$  such that  $\phi(\bar{x}, \bar{y}, \bar{c}, \bar{R})$  defines over  ${}^n B = \{\bar{b}: \bar{b} \in B, l(\bar{b}) = n\}$  an equivalence relation, with infinitely many non-pseudo-finite equivalence classes.*

PROOF. For  $n = 1$ , we can show as in Claim 4A, Claim 5A that we can interpret the quantifier over equivalence relations. By Rabin [8], it then follows that we can interpret  $Q_{II}$ .

Now we shall reduce the case  $n > 1$  to  $n = 1$ , using the interpretability of  $Q_P$  by  $Q_\psi$ .

Choose by induction on  $\max\{i, j\}$  sequences  $\bar{a}^{i,j}, j < \omega$  such that

- 1)  $\bar{a}^{i,j} \in B$
- 2)  $A \models \phi[\bar{a}^{i,j}, \bar{a}^{l,k}, \bar{c}, \bar{R}]$  iff  $i = l$
- 3) for  $\langle i, j \rangle \neq \langle l, k \rangle$ ,  $\bar{a}^{i,j}, \bar{a}^{l,k}$  are disjoint, and  $\bar{a}^{i,j}, \bar{c}$  are disjoint.

For  $m = 1, n$ , define  $f_m$  as the permutation of  $A$  (of order two) interchanging  $\bar{a}_1^{i,j}$  with  $\bar{a}_m^{i,j}$  for  $i, j < \omega$ , and taking any other  $b \in A$  to itself.

Let  $B^* = \{\bar{a}_1^{i,j} : i, j < \omega\}$ .

Now the formula

$$\phi^*(x, y, \bar{z}, \bar{R}, f_1, \dots, f_n) = \phi(f_1(x), f_2(x), \dots, f_n(x), f_1(y), f_2(y), \dots, f_n(y), \bar{c}, \bar{R})$$

defines on  $B^*$  an equivalence relation with infinitely many infinite equivalence classes. This proves Claim 6A.

**CLAIM 6B.**  $Q_{II}$  is interpretable by  $Q_\psi$  if there are  $\phi(\bar{x}, \bar{y}, r)$ ,  $A, R \in R_\psi(A)$  and  $\bar{a}^n \in A (n < \omega)$ , such that for every  $n < \omega$ ,  $\theta_n = \bigwedge_{m < n} \phi(\bar{x}, \bar{a}^m, R) \wedge \neg \phi(\bar{x}, \bar{a}^n, R)$  is not pseudo-algebraic.

**PROOF.** By the compactness theorem we can assume that each formula  $\theta_n$  is satisfied by  $> 2^{\aleph_0}$  pairwise disjoint sequences. Let

$$\begin{aligned} B &= \{\bar{a}_i^m : m < \omega, 1 \leq i \leq l(\bar{a}^m)\}, \quad e = \{\langle \bar{b}, \bar{c} \rangle : \bar{b}, \bar{c} \in A, l(\bar{b}) = l(\bar{c})\} \\ &= l(\bar{x}), (\forall \bar{a} \in B) A \models \phi[\bar{b}, \bar{a}, R] \equiv \phi[\bar{c}, \bar{a}, R]. \end{aligned}$$

Then  $e$  is an equivalence relation over  ${}^{l(\bar{a}^m)}A$ . The set of sequences which satisfies  $\theta_n$  is split into at most  $2^{\aleph_0}$  equivalence classes (as  $|B| = \aleph_0$ ), so at least one of them contains  $> 2^{\aleph_0}$  pairwise disjoint sequences, hence is not pseudo-finite. Clearly for  $n \neq m$ , a sequence satisfying  $\theta_n$  and a sequence satisfying  $\theta_m$  are not equivalent. Thus we get our result by Claim 6A.

**CLAIM 6C.** If  $Q_{II}$  is not interpretable by  $Q_\psi$  then for every  $\phi(\bar{x}, \bar{y}, r)$  there are  $m(\phi) < \omega$ , and  $\chi_{\phi,i}(\bar{x}, \bar{z}, r)$   $i = 1, \dots, m(\phi)$  such that

for any  $A, R \in R_\psi(A)$  there is  $\bar{c} \in A$  which satisfies

- 1)  $A \models (\forall \bar{x}) \bigvee_{i=1}^{m(\phi)} \chi_{\phi,i}(\bar{x}, \bar{c}, R)$
- 2)  $A \models \neg (\exists \bar{x}) [\chi_{\phi,i}(\bar{x}, \bar{c}, R) \wedge \chi_{\phi,j}(\bar{x}, \bar{c}, R)]$  for  $i \neq j$
- 3) the sets  $S_i = \{\bar{a} : A \models \chi_{\phi,i}[\bar{a}, \bar{c}, R]\}$  are  $\phi(\bar{x}, \bar{y}, r)$ -minimal; moreover for some fixed  $m_1(\phi) < \omega$ , for no  $S_i$  and no  $\bar{b} \in A$ , do both  $\{\bar{a} \in S_i : A \models \phi[\bar{a}, \bar{b}, R]\}$  and

$\{\bar{a} \in S_i : A \models \neg \phi[\bar{a}, \bar{b}, R]\}$  contain  $m_1(\phi)$  pairwise disjoint sequences (we call this property “ $(\phi, m_1(\phi))$ -minimality”).

PROOF. By Claim 6B and the compactness theorem, there is an  $m_1(\phi) < \omega$  such that we cannot find  $A, R \in R_\psi(A)$ , sequences  $\bar{a}^n \in A$  for  $n < m_1(\phi)$ , and a formula  $\phi^* \in \{\phi(\bar{x}, \bar{y}, r), \neg \phi(\bar{x}, \bar{y}, r)\}$  such that for each  $n < m_1(\phi)$ ,  $\bigwedge_{m < n} [\phi^*(\bar{x}, \bar{a}^m, R) \wedge \neg \phi^*(\bar{x}, \bar{a}^n, R)]$  is satisfied by  $\geq m_1(\phi)$  pairwise disjoint sequences.

Now let  $\eta$  denote a sequence of ones and zeros. Define by induction on  $l$ , sequences  $\bar{a}_\eta l(\eta) \leq l$  and formulae  $\chi_\eta = \chi_\eta(\bar{x}, \bar{b}_\eta, R)$ .

For  $l = 0$ ,  $\eta$  the empty sequence,  $\chi_\eta = (\forall x)(x = x)$ .

Suppose we have made the definitions for  $l$ ; let us do so for  $l+1$ . Let  $l(\eta) = l$ . If there is an  $\bar{a}_\eta \in A$  such that both  $\chi_\eta(\bar{x}, \bar{b}_\eta, R) \wedge \phi(\bar{x}, \bar{a}_\eta, R)$ ,  $\chi_\eta(\bar{x}, \bar{b}_\eta, R) \wedge \neg \phi(\bar{x}, \bar{a}_\eta, R)$  are satisfied by  $\geq m_1(\phi)$  pairwise disjoint sequences, then choose such  $\bar{a}_\eta$ ; otherwise choose  $\bar{a}_\eta$  arbitrarily.

Then if  $l(\eta) = l + 1$ , define  $\chi_\eta(\bar{x}, \bar{b}_\eta, R)$  as follows:  $\eta = \langle i(1), \dots, i(l+1) \rangle$ ; then if  $i(l+1) = 0$ ,

$$\chi_\eta(\bar{x}, \bar{b}_\eta, R) = \chi_{\langle i(1), \dots, i(l) \rangle}(\bar{x}, \bar{b}_{\langle i(1), \dots, i(l) \rangle}, R) \wedge \phi(\bar{x}, \bar{a}_{\langle i(1), \dots, i(l) \rangle}, R)$$

and if  $i(l+1) = 1$ , it is the same with  $\neg \phi$  instead of  $\phi$ .

By the definition of  $m_1(\phi)$ , if, e.g.,  $l(\eta) = 2m_1(\phi) + 2$ , then  $\chi_\eta(\bar{x}, \bar{b}_\eta, R)$  is  $(\phi, m_1(\phi))$ -minimal. Clearly the  $\chi_\eta(\bar{x}, \bar{b}_\eta, R)$ ,  $l(\eta) = 2m_1(\phi) + 2$  form a partition; and the choice of  $\chi_\eta(\bar{x}, z, r)$  does not depend on the particular model. Thus Claim 6C is proved.

CLAIM 6D. Suppose  $Q_{II}$  is not interpretable by  $Q_\psi$ . If  $A$  is an infinite  $R \in R_\psi(A)$ ,  $B \subseteq A$ ,  $\bar{a}, \bar{b} \in A$ , and  $\bar{a}$  is pseudo-algebraic over  $B \cup \{\dots, \bar{b}_i, \dots\}$  but not over  $B$ , then  $\bar{b}$  is pseudo-algebraic over  $B \cup \{\dots, \bar{a}_i, \dots\}$ .

PROOF. Suppose the conclusion fails. There are  $\bar{c} \in B$ , and  $\phi(\bar{x}, \bar{y}, \bar{z}, r)$  such that  $A \models \phi[\bar{a}, \bar{b}, \bar{c}, R]$ , and  $\phi(\bar{x}, \bar{b}, \bar{c}, R)$  is pseudo-algebraic. Say there do not exist  $m$  pairwise disjoint sequences in  $\phi[A, \bar{b}, \bar{c}, R]$ . Let  $\theta(\bar{x}, \bar{y}, \bar{z}, R)$  say that  $\phi(\bar{x}, \bar{y}, \bar{z}, R)$  and there do not exist  $m$  pairwise disjoint sequences in  $\phi(A, \bar{y}, \bar{z}, R)$ . Since  $A \models \theta[\bar{a}, \bar{b}, \bar{c}, R]$ ,  $\theta[\bar{a}, \bar{y}, \bar{c}, R]$  is not pseudo-algebraic. For each  $n < \omega$ , let  $\chi_n(\bar{x}, \bar{z}, R)$  say that there are  $n$  disjoint sequences  $\bar{d}$  such that  $\theta(\bar{x}, \bar{d}, \bar{z}, R)$  is satisfied. Thus  $A \models \chi_n[\bar{a}, \bar{c}, R]$  for all  $n$ , and hence  $\chi_n(\bar{x}, \bar{c}, R)$  is not pseudo-algebraic.

Now, by the compactness theorem, we can assume that there are  $\bar{a}^i, \bar{b}^{i,j} \in A$  for  $i, j < \omega$  such that

$$A \models \theta[\bar{a}^i, \bar{b}^{i,j}, \bar{c}, R] \text{ for all } i, j,$$

and  $\bar{a}^k, \bar{a}^l$  (likewise  $\bar{b}^{i,k}, \bar{b}^{i,l}$ ) are disjoint for  $k \neq l$ . By rejecting some  $\bar{b}^{i,j}$ , we can assume that  $\bar{b}^{i,j}, \bar{b}^{k,l}$  are disjoint unless  $\langle i, j \rangle = \langle k, l \rangle$ , and also that

$$A \models \theta[\bar{a}^i, \bar{b}^{j,k}, \bar{c}, R] \equiv \theta[\bar{a}^i, \bar{b}^{j,l}, \bar{c}, R]$$

when  $i \leq j$ . Further, by Ramsey's theorem, we arrange that the truth value of  $\theta[\bar{a}^i, \bar{b}^{j,k}, \bar{c}, R]$  for  $i < j$  is independent of  $i, j$ .

Now since there are no  $m$  pairwise disjoint sequences in  $\theta[A, \bar{b}^{m,0}, \bar{c}, R]$ , it follows that for all  $i, j, k$ , with  $i \leq j$ ,  $A \models \theta[\bar{a}^i, \bar{b}^{j,k}, \bar{c}, R]$  if and only if  $i = j$ . Thus we get a contradiction as in Claim 6B.

**CLAIM 6E.** *If  $\bar{a} = \langle \bar{a}_1, \dots, \bar{a}_n \rangle$  is pseudo-algebraic over  $B \subseteq A$  in  $(A, R)$ , then some  $a_i$  is algebraic over  $B$  in  $(A, R)$ .*

**PROOF.** Since  $\bar{a}$  is pseudo-algebraic over  $B$ , there is a pseudo-algebraic  $\phi(\bar{x}, \bar{b}, R)$  ( $\bar{b} \in B$ ),  $A \models \phi[\bar{a}, \bar{b}, R]$ . Hence there is a finite set  $C = \{c_1, \dots, c_n\}$  such that for any  $\bar{a}^1 \in A$ ,  $A \models \phi[\bar{a}^1, \bar{b}, R]$  implies  $\{\bar{a}_1^1, \dots\}$  and  $C$  are not disjoint. Without loss of generality  $n$  is minimal. Let

$$\theta^1(z_1, \dots, z_n, \bar{y}, r) = (\forall \bar{x}) [\phi(\bar{x}, \bar{y}, r) \rightarrow \bigvee_{i,j} \bar{x}_i = z_j]$$

$$\theta^2(z, \bar{y}, r) = (\exists z_2, \dots, z_n) \theta^1(z, z_2, \dots, z_n, r).$$

Clearly for some  $i$ ,  $A \models \theta^2[\bar{a}_i, \bar{b}, R]$ . As in Claim 4C we can show that  $\theta^2(z, \bar{b}, R)$  is algebraic.

**CLAIM 6F.** *Assume  $Q_{II}$  is not interpretable by  $Q_\psi$ . Let  $R \in R_\psi(A)$ , and for every formula  $\phi$ , let  $\chi_{\phi,i}$   $i = 1, \dots, m(\phi)$ ,  $\bar{c}^\phi$  be as in Claim 6C. Let  $C = \{\bar{c}_i^\phi: \phi, i\} \cup \{\text{elements algebraic over some } \bar{c}^h\}$ .*

*If  $\bar{a}, \bar{b} \in A$ ,  $l(\bar{a}) = l(\bar{b}) = n$  and if the following conditions are met:*

1) *if  $\bar{a}_{i_2}, \dots, \bar{a}_{i_l}$  are algebraic over  $C \cup \{\bar{a}_{i_1}\}$ , then  $\langle \bar{a}_{i_1}, \dots, \bar{a}_{i_l} \rangle, \langle \bar{b}_{i_1}, \dots, \bar{b}_{i_l} \rangle$  realize the same type over  $C$  in  $(A, R)$ ,*

2) *as in (1), interchanging  $\bar{a}, \bar{b}$ ,*

*then  $\bar{a}, \bar{b}$  realize the same type over  $C$ .*

**PROOF.** We prove by induction on  $n$ .

For  $n = 1$ , (1) for  $l = 1$  is the conclusion.

Suppose we have proved the claim for  $n$ ; we shall prove it for  $n + 1$ . Let  $\phi = \phi(x, \bar{y}, \bar{z}, r)$  be a formula,  $\bar{c} \in C$ .

If each  $\bar{a}_i$  is algebraic over  $\bar{a}_1$  we are finished. By renaming the  $\bar{a}_i$ 's we can



assume that  $\bar{a}_2, \dots, \bar{a}_l$  are algebraic over  $C \cup \{a_1\}$ , but  $a_{l+1}, \dots, \bar{a}_{n+1}$  are not;  $l \leq n$ . Let

$$\bar{a}^1 = \langle \bar{a}_1, \dots, \bar{a}_l \rangle, \quad \bar{a}^2 = \langle \bar{a}_{l+1}, \dots, \bar{a}_{n+1} \rangle,$$

$$\bar{b}^1 = \langle \bar{b}_1, \dots, \bar{b}_l \rangle, \quad \bar{b}^2 = \langle \bar{b}_{l+1}, \dots, \bar{b}_{n+1} \rangle.$$

By (1) and (2),  $\bar{b}_2, \dots, \bar{b}_l$  are algebraic over  $\bar{b}_1$ , but  $b_{l+1}, \dots, \bar{b}_{n+1}$  are not. By Claim 6E,  $\bar{a}^2, \bar{b}^2$  are not pseudo-algebraic over, respectively,  $\bar{a}^1 \cup C, \bar{b}^1 \cup C$ .

We must prove that for any  $\bar{c} \in C$ ,  $\phi(\bar{x}, \bar{y}, \bar{z}, r)$ ,  $A \models \phi[\bar{a}^1, \bar{a}^2, \bar{c}, R] \equiv \phi[\bar{b}^1, \bar{b}^2, \bar{c}, R]$ . By the induction hypothesis,  $\bar{a}^i, \bar{b}^i$  realize the same type over  $C$ . Now we apply the definition of  $\bar{c}^\psi$  for  $\psi(\bar{y}, \bar{x}, \bar{z}, R) = \phi(\bar{x}, \bar{y}, \bar{z}, R)$  (see Claim 6C).

By Claim 6C (1) there is an  $i$  such that  $A \models \chi_{\psi,i}[\bar{a}^2, \bar{c}^\psi, R]$ .

By Claim 6C (2) one of

$$\chi_{\psi,i}(\bar{y}, \bar{c}^\psi, R) \wedge \phi(\bar{a}^1, \bar{y}, \bar{c}, R)$$

$$\chi_{\psi,i}(\bar{y}, \bar{c}^\psi, R) \wedge \neg \phi(\bar{a}^1, \bar{y}, \bar{c}, R)$$

(w.l.o.g. the second), is not satisfied by  $\geq m_1(\psi)$  pairwise disjoint sequences. As  $\bar{a}^2$  is not pseudo-algebraic over  $\bar{a}^1 \cup C$ , clearly

$$A \models \phi[\bar{a}^1, \bar{a}^2, \bar{c}, R].$$

Since  $\bar{a}^2$  and  $\bar{b}^2$  have the same type over  $C$ ,  $A \models \chi_{\psi,i}[\bar{b}^2, \bar{c}^\psi, R]$ , and since  $\bar{a}^1, \bar{b}^1$  have the same type over  $C$ ,  $\chi_{\psi,i}[\bar{y}, \bar{c}^\psi, R] \wedge \neg \phi(\bar{b}^1, \bar{y}, \bar{c}, R)$  is not satisfied by  $\geq m_1(\psi)$  pairwise disjoint sequences. Hence the above reasoning gives that

$$A \models \phi[\bar{b}^1, \bar{b}^2, \bar{c}, R]$$

which completes the proof.

**CLAIM 6G.** Suppose  $Q_{II}$  cannot be interpreted by  $Q_\psi$ . Then there are  $n_0, n_1 < \omega$ ,  $\phi(x, y, \bar{z}, r)$ ,  $\chi_i(\bar{x}^i, \bar{z}, r)$   $i < n_1$   $l(\bar{x}^i) = n^i$  such that  $(\exists^{\leq n_0} x) \phi(x, y, \bar{z}, r)$  and  $\phi(x, x, \bar{z}, r)$  and  $(\exists^{\leq n_1} y) \phi(x, y, \bar{z}, r)$  hold and for any  $A, R \in R_\psi(A)$  there is a  $\bar{c} \in A$ , such that if  $\bar{a}, \bar{b} \in A$   $l(\bar{a}) = l(\bar{b}) = n(\psi)$  and if the following conditions are met

1) if  $\models \phi[\bar{a}_{i_1}, \bar{a}_{i_1}, \bar{c}, R]$  for  $l = 2, \dots, k$  and  $n^i = k$  then  $A \models \chi_i[\bar{a}_{i_1}, \dots, \bar{a}_{i_k}, \bar{c}, R] \equiv \chi_i[\bar{b}_{i_1}, \dots, \bar{b}_{i_k}, \bar{c}, R]$ ,

2) as in (1), interchanging  $\bar{a}$  and  $\bar{b}$ ,

then  $A \models r[\bar{a}] \equiv r[\bar{b}]$ .

PROOF. It follows from Claim 6D and 6F and the compactness theorem. (Note that in Claim 6F, we can choose any  $\bar{c}^\phi$ , as long as it satisfies a first-order condition which expresses (1), (2), and (3) of Claim 6C, when we are interested in the formula  $r(\bar{x})$  only. We can have one  $\phi$  because the disjunction of algebraic formulae is algebraic and if  $a$  is algebraic over  $B$ , then for some  $n, \phi, \bar{b} \in B$ ,  $A \models (\exists^{\leq n} x)\phi(x, \bar{b}, R)$ ; hence  $a$  satisfies  $\theta^1(x, \bar{b}, R) = (\exists^{\leq n} y)\theta(y, \bar{b}, R) \wedge \theta(x, \bar{b}, R)$ , and  $(\exists^{\leq n} x)\theta^1(x, \bar{b}, R)$  holds.)

PROOF OF LEMMA 6. Assume  $Q_{II}$  cannot be interpreted by  $Q_\psi$ , and we shall interpret  $Q_\psi$  by  $Q_P$ . We use the results and notation of Claim 6G.

Call  $a, b$   $n$ -connected (in  $(A, R)$ ,  $R \in R_\psi(A)$ ,  $\bar{c}$  as in Claim 6G if there are  $a = c^0, c^2, \dots, c^n = b$  such that  $A \models \phi[c^i, c^{i+1}, \bar{c}, R] \vee \phi[c^{i+1}, c^i, \bar{c}, R]$  for  $1 \leq i < n$ . By the remark above, the number of  $b$ 's  $n$ -connected to  $a$  is  $\leq k(n) < \omega(k(n))$  depends only on  $\phi, \psi$  and  $n$ ).

Now choose inductively  $A_n \subseteq A$ ,  $n \geq 1$  such that  $A_n$  is a maximal subset of  $A - \cup_{i < n} A_i$  with no two 2-connected elements. For  $n \geq k(2) + 2$ ,  $A_n$  is empty, because if  $a \in A_n$ , then by the definition of  $A_i$ , ( $i < n$ ) there is a  $b_i \in A_i$  such that  $a, b_i$  are 2-connected. So  $> k(2)$  elements are two-connected to  $A$ , a contradiction. Now for any  $a \neq b \in A_n$ ,  $\phi(A, a, \bar{c}, R)$ ,  $\phi(A, b, \bar{c}, R)$  are disjoint (because if  $c$  is in the intersection, then  $c, a$  and  $c, b$  are 1-connected, hence  $a, b$  are 2-connected).

Now it is clear how to define  $r$  by permutations and sets. By dividing the  $A_i$ 's according to  $|\phi(A, a, \bar{c}, R)|$ , we get  $A = \cup_{i < m} A_i$ ,  $a \neq b \in A_i$  implies  $\phi(A, a, \bar{c}, R) \cap \phi(A, b, \bar{c}, R) = \emptyset$ , and  $|\phi(A, a, \bar{c}, R)| = m(i)$ . For each  $i$  choose permutations of order two  $f_1^i, \dots, f_{m(i)}^i$  such that

$$\phi(A, a, \bar{c}, R) = \{f_j^i(a) : 1 \leq j \leq m(i)\}.$$

In view of Claim 6G, we thus represent  $R[\in R_\psi(A)]$  by the permutations  $f_j^i$ , the sets  $A_i$ , and the additional sets

$$A_{i,k,1,1,\dots} = \{a \in A_i : A \models \chi_k[f_{1,1}^i(a), \dots, R]\}.$$

In fact there are only finitely many such possible representations, so by adding a sequence of elements, we can encode, by equalities, the proper case.

#### REFERENCES

1. Bell, J. L. and A. B. Slomson, *Models and Ultraproducts*, North Holland, 1969.
2. P. Erdős and R. Rado, *Intersection theorems for systems of sets*, J. London Math. Soc. 44 (1969), 467-479.

3. Ju. L. Ershov, *Undecidability of theories of symmetric and simple finite groups*, Dokl. Akad. Nauk SSSR **158**, (1964) 777–779.
4. Ju. L. Ershov, *New examples of undecidability of theories*, Algebra i Logika **5** (1966), 37–47.
5. R. McKenzie, *On elementary types of symmetric groups*, Algebra Universalis **1** (1971), 13–20.
6. M. D. Morley and R. L. Vaught, *Homogeneous universal models*, Math. Scand. **11** (1962), 37–57.
7. A. G. Pinus, *On elementary definability of symmetric group and lattices of equivalences*, Algebra Universalis, to appear.
8. M. O. Rabin, *A simple method for undecidability proofs*, Proc. 1964 Int. Congress for Logic, North Holland, 1965, pp. 58–68.
9. F. D. Ramsey, *On a problem of formal logic*, Proc. London Math. Soc. **30** (1929), 338–384.
10. S. Shelah, *There are just four possible second-order quantifiers and on permutation groups*, Notices Amer. Math. Soc. **19** (1972), A-717.
11. S. Shelah, *First order theory of permutations groups*, Israel J. Math. **14** (1973), 149–162; and *Errata to "First order theory of permutations groups"*, Israel J. Math. **15** (in press).

INSTITUTE OF MATHEMATICS  
THE HEBREW UNIVERSITY OF JERUSALEM  
JERUSALEM, ISRAEL