



ELSEVIER

Annals of Pure and Applied Logic 79 (1996) 289–316

ANNALS OF
PURE AND
APPLIED LOGIC

Saturated filters at successors of singulars, weak reflection and yet another weak club principle

Mirna Džamonja^{a,*} Saharon Shelah^b

^a *Department of Mathematics, University of Wisconsin–Madison, Madison, WI 53706, USA*

^b *Institute of Mathematics, Hebrew University of Jerusalem, 91904 Givat Ram, Israel*

Received 1 February 1995

Communicated by A. Nerode

Abstract

Suppose that λ is the successor of a singular cardinal μ whose cofinality is an uncountable cardinal κ . We give a sufficient condition that the club filter of λ concentrating on the points of cofinality κ is not λ^+ -saturated.¹ The condition is phrased in terms of a notion that we call weak reflection. We discuss various properties of weak reflection.

We introduce a weak version of the \clubsuit -principle, which we call \clubsuit^* , and show that if it holds on a stationary subset S of λ , then no normal filter on S is λ^+ -saturated. Under the above assumptions, $\clubsuit^*(S)$ is true for any stationary subset S of λ which does not contain points of cofinality κ . For stationary sets S which concentrate on points of cofinality κ , we show that $\clubsuit^*(S)$ holds modulo an ideal obtained through the weak reflection.

0. Introduction

Suppose that $\lambda = \mu^+$ and μ is an infinite cardinal of cofinality κ . We revisit the classical question of whether a normal filter on λ can be λ^+ -saturated. We are in particular concerned with the case of κ uncountable and less than μ . We are mostly interested in the club filter on λ .

While the richness of the literature on the subject provides us with a strong motivation for a further study, it also prevents us from giving a complete history

* Corresponding author. E-mail: dzamonja@math.wisc.edu.

¹ Added in proof: M. Gitik and S. Shelah have subsequently and by a different technique shown that the club filter on such λ is never saturated.

and bibliography involved. We shall give a list of those references which are most directly connected or used in our results, and for further reading we can suggest looking at the references mentioned in the papers that we refer to.

It is well known that for no regular $\theta > \aleph_0$ the club filter on θ can be θ -saturated, but modulo the existence of huge or other large cardinals it is consistent that \aleph_1 carries an \aleph_2 -saturated normal filter [5], or in fact that any uncountable regular \aleph_α carries an $\aleph_{\alpha+1}$ -saturated normal filter [3]. In these arguments the saturated filter obtained is not the club filter. As a particular case of [13, Th. 14] or [14, Th. 6], if $\sigma = \rho^+$, then the club filter \mathcal{D}_σ restricted to the elements of σ of a fixed cofinality $\theta \neq \text{cf}(\rho)$ is not σ^+ -saturated. It is consistent that \mathcal{D}_{\aleph_1} is \aleph_2 -saturated, as is shown in [4] and also in [21].

If $\sigma < \rho$ and ρ is a regular cardinal, we use S_ρ^σ to denote the set of elements of σ which have cofinality ρ . Let λ be as above.

In the first section of the present paper, we give a sufficient condition that $\mathcal{D}_\lambda \upharpoonright S_\kappa^\lambda$ is not λ^+ -saturated. Here $\aleph_0 < \kappa = \text{cf}(\mu) < \mu$. The condition is a reflection property, which we shall call *weak reflection*, as we show that it is weaker than some known reflection properties. We discuss the properties of weak reflection in more detail in Section 1. Of course, the main part of the section is to show that the appropriate form of the weak reflection indeed suffices for $\mathcal{D}_\lambda \upharpoonright S_\kappa^\lambda$ not to be λ^+ -saturated, which is done in 1.13. In 1.15 the argument is generalized to some other normal filters on λ , and [11], Section 3 is revisited.

In the second section we introduce the combinatorial principle \clubsuit^* which has the property that, if $\clubsuit^*(S)$ holds, then no normal filter on S is λ^+ -saturated. Here S is a stationary subset of λ . The \clubsuit^* is a weak form of \clubsuit . The \clubsuit -principle was first introduced for \aleph_1 in Ostaszewski [8], and later investigated in a more general setting in [10] and elsewhere.

If λ is the successor of the singular cardinal μ whose cofinality is κ , then $\clubsuit^*(\lambda \setminus S_\kappa^\lambda)$ is true just in *ZFC*. As a corollary of this we obtain an alternative proof of a part of the result of [13, Th. 14] or [14, Th. 6]. On the other hand, $\clubsuit^*(S_\kappa^\lambda)$ only holds modulo an ideal defined through the weak reflection (in Section 1), so this gives a connection with the results of the first section. For the case of μ being a strong limit, \clubsuit^* just becomes the already known \clubsuit . Trying to apply to \clubsuit^* arguments which work for the corresponding version of \diamond , we came up with a question we could not answer, so we pose it in 2.6.

Before we proceed to present our results, we shall introduce some notation and conventions that will be used throughout the paper.

Notation 0.0. (0) Suppose that $\gamma \geq \theta$ and θ is a regular cardinal. Then

$$S_{<\theta}^\gamma = \{\delta < \gamma : \aleph_0 \leq \text{cf}(\delta) < \theta\}, \quad \text{if } \theta > \aleph_0.$$

$$S_\theta^\gamma = \{\delta < \gamma : \text{cf}(\delta) = \theta\}.$$

More generally, we use $S_{r\theta}^\gamma$ for $r \in \{<, \leq, =, \neq, >, \geq\}$ to describe

$$S_{r\theta}^\gamma = \{\delta < \gamma : \aleph_0 \leq \text{cf}(\delta) \& \text{cf}(\delta) r \theta\}.$$

(1) *SING* denotes the class of singular ordinals, that is, all ordinals δ with $\text{cf}(\delta) < \delta$. *LIM* denotes the class of all limit ordinals.

(2) For λ a cardinal with $\text{cf}(\lambda) > \aleph_0$, we denote by \mathcal{D}_λ the club filter on λ . The ideal of non-stationary subsets of λ is denoted by J_λ .

(3) If $C \subseteq \lambda$, then

$$\text{acc}(C) = \{\alpha \in C : \alpha = \sup(C \cap \alpha)\} \quad \text{and} \quad \text{nacc}(C) = C \setminus \text{acc}(C).$$

We now go on to the first section of the paper.

1. Saturated filters on the successor of a singular cardinal of uncountable cofinality

Suppose that μ is a cardinal with the property $\mu > \text{cf}(\mu) = \kappa > \aleph_0$, and that $\lambda = \mu^+$. We wish to discuss the saturation of $\mathcal{D}_\lambda \upharpoonright S_\kappa^\lambda$ and some other normal filters on λ . For the reader's convenience, we recall some relevant definitions and notational conventions. The cardinals λ , μ and κ as above will be fixed throughout this section.

Notation 1.0. (0) Suppose that \mathcal{D} is a filter on the set A . The *dual ideal* \mathcal{I} of \mathcal{D} is defined as

$$\mathcal{I} = \{a \subseteq A : A \setminus a \in \mathcal{D}\}.$$

A set $a \subseteq A$ is \mathcal{D} -positive or \mathcal{D} -stationary if $a \notin \mathcal{I}$. The family of all \mathcal{D} -stationary sets is denoted by \mathcal{D}^+ . A subset of A which is not \mathcal{D} -stationary is referred to as \mathcal{D} -non-stationary.

(1) Suppose that σ is a cardinal. A filter \mathcal{D} on a set A is σ -saturated iff there are no σ sets which are all \mathcal{D} -stationary, but no two of them have a \mathcal{D} -stationary intersection. If we denote the dual ideal of \mathcal{D} by \mathcal{I} , then this is equivalent to saying that $\mathcal{P}(A)/\mathcal{I}$ has σ -ccc.

For a cardinal ρ , the filter \mathcal{D} is ρ -complete if it is closed under taking intersections of $< \rho$ of its elements.

(2) A σ -complete filter \mathcal{D} on a cardinal σ is *normal* if, for any sequence $X_\alpha (\alpha < \sigma)$ of elements of \mathcal{D} , the diagonal intersection

$$\Delta_{\alpha < \sigma} X_\alpha \stackrel{\text{def}}{=} \{\beta \in \sigma : \alpha < \beta \Rightarrow \beta \in X_\alpha\}$$

is an element of \mathcal{D} , and \mathcal{D} contains all final segments of σ .

(3) Suppose that \mathcal{D} is a filter on A and $S \subseteq A$ is \mathcal{D} -stationary. We use $\mathcal{D} \upharpoonright S$ to denote

$$\mathcal{D} \upharpoonright S \stackrel{\text{def}}{=} \{X \cap S : X \in \mathcal{D}\}.$$

We have fixed cardinals λ , μ and κ at the beginning of this section. By [14, Th. 6] or [13, Th. 14], we know that $\mathcal{D}_\lambda \upharpoonright S_{\neq \kappa}^\lambda$ is not λ^+ -saturated. We shall now introduce

a sufficient condition, under which we can prove that $\mathcal{D}_\lambda \upharpoonright S_\kappa^\lambda$ is not λ^+ -saturated. In Theorem 1.15, we extend the result to a somewhat larger class of normal filters on λ .

Definition 1.1. Suppose $\chi > \aleph_0$ is a regular cardinal and $\eta > \chi$ is an ordinal. We say that η has the strong non-reflection property for χ , if there is a function $h : \eta \rightarrow \chi$ such that for every $\delta \in S_\chi^\eta$, there is a club subset C of δ with $h \upharpoonright C$ strictly increasing. In such a case we say that h witnesses the strong non-reflection of η for χ .

If η does not have the strong non-reflection property for χ , then we say that η has the weak reflection for χ , or that η is weakly reflective for χ .

If $h : \eta \rightarrow \chi$ is a function, we define

$$\text{ref}(h) \stackrel{\text{def}}{=} \{\delta \in S_\chi^\eta : h \upharpoonright C \text{ is not strictly increasing for any club } C \text{ of } \delta\}.$$

We shall first make some general remarks about Definition 1.1.

Observation 1.2. (1) If η is weakly reflective for $\chi > \aleph_0$, and $\zeta > \eta$, then ζ is weakly reflective for χ .

(2) If $\eta > \chi = \text{cf}(\chi) > \aleph_0$, then η has the strong non-reflection property for χ iff there is an $h : \eta \rightarrow \chi$ such that for all $\delta \in S_\chi^\eta$, there is a club C of δ with the property that $h \upharpoonright C$ is 1–1. In fact, the two sides of the equivalence can be witnessed by the same function h .

(3) Suppose that χ is a given uncountable regular cardinal such that there is an η which weakly reflects at χ . Then the minimal such η , which we shall denote by $\theta^*(\chi)$, is a regular cardinal $> \chi$. Consequently, ζ weakly reflects at χ iff there is a regular cardinal $\theta \leq \zeta$ such that θ weakly reflects at χ .

(4) Suppose that χ is a regular cardinal, and $\eta = \theta^*(\chi)$ or η is an ordinal of cofinality $\geq \theta^*(\chi)$, so that $S_{\theta^*(\chi)}^\eta$ is stationary in η . (This makes sense, as by (3), if $\theta^*(\chi)$ is defined, then it is a regular cardinal.) Then not only is η weakly reflective for χ , but for every $h : \eta \rightarrow \chi$, the set $\text{ref}(h)$ is stationary.

Proof. (1) Follows from the definition.

(2) If η has the strong non-reflection for χ , the other side of the above equivalence is obviously true.

In the other direction, suppose that h satisfies the conditions on the right hand side, and fix a $\delta \in S_\chi^\eta$. Let C be a club of δ on which h is 1–1. In particular note that $\text{ran}(h)$ is cofinal in χ and that $\text{otp}(C) = \chi$. By induction on $\gamma < \chi$, define

$$\beta_\gamma = \text{Min} \left\{ \alpha \in C : (\forall \beta \in (C \setminus \alpha)) (\forall \zeta < \gamma) (h(\beta) > h(\beta_\zeta)) \right\}.$$

Then $D = \{\beta_\gamma : \gamma < \chi\}$ is a club of δ and $h \upharpoonright D$ is strictly increasing.

(3) Let $\eta = \theta^*(\chi)$ be the minimal ordinal which weakly reflects at χ , so obviously $\eta > \chi$. If η is not a regular cardinal, we can find an increasing continuous sequence of ordinals $\langle \eta_i : i < \zeta \rangle$ which is cofinal in η , and such that $\zeta < \eta$. We can also assume that $\eta_0 = 0$ and $\eta_1 > \chi$. In addition, for every $i < \zeta$, if i is a successor, we can assume that η_i is also a successor.

Then, for every $i < \zeta$, there is an $h_{i+1} : \eta_{i+1} \rightarrow \chi$ which exemplifies that η_{i+1} has the strong non-reflection property for χ . There is also a $g : \zeta \rightarrow \chi$ which witnesses that ζ has the strong non-reflection property for χ . Define $h : \eta \rightarrow \chi$ by

$$h(\alpha) = \begin{cases} h_{i+1}(\alpha) & \text{if } \alpha \in (\eta_i, \eta_{i+1}), \\ g(i) & \text{if } \alpha = \eta_i. \end{cases}$$

Since η is weakly reflective for χ , there must be a $\delta \in S_\chi^\eta$ such that for no club C of δ is $h \upharpoonright C$ strictly increasing. We can distinguish two cases:

Case 1: $\delta \in (\eta_i, \eta_{i+1}]$ for some $i < \zeta$. Then there is a club C of δ on which h_{i+1} is strictly increasing. But then $h \upharpoonright (C \setminus (\eta_i + 1)) = h_{i+1} \upharpoonright (C \setminus (\eta_i + 1))$ is strictly increasing, and $C \setminus (\eta_i + 1)$ is a club of δ . A contradiction.

Case 2: $\delta = \eta_i$ for some limit $i < \zeta$. Notice that $D = \{\eta_j : j < i\}$ is a club of η_i . We know that there is a club C of i on which g is increasing. Then setting $E = C \cap D$, we conclude that $h \upharpoonright E$ is strictly increasing. Therefore, a contradiction.

(4) We first assume that $\eta = \theta^*(\chi)$. By (2), $\theta^*(\chi)$ is necessarily a regular cardinal $> \chi$. Let $h : \theta^*(\chi) \rightarrow \chi$ be given. Suppose that $\text{ref}(h)$ is non-stationary in $\theta^*(\chi)$ and fix a club E in $\theta^*(\chi)$ such that $E \cap \text{ref}(h) = \emptyset$. Without loss of generality, $\text{otp}(E) = \theta^*(\chi)$. Let us fix an increasing enumeration $E = \{\alpha_i : i < \theta^*(\chi)\}$. Without loss of generality, $\alpha_0 = 0$ and $\alpha_i > \chi$ and α_{i+1} is a successor for every i . So, for each i , there is a function $h_i : \alpha_{i+1} \rightarrow \chi$ which witnesses that α_{i+1} is strongly non-reflective at χ .

But now we can use h and h_i for $i < \theta^*(\chi)$ to define a function which will contradict that $\theta^*(\chi)$ weakly reflects at χ , similarly to the proof of (3).

The other case is that $\eta > \theta^*(\chi)$, so $S_{\theta^*(\chi)}^\eta$ is stationary in η . Let $h : \eta \rightarrow \chi$ be given.

If $\delta \in S_{\theta^*(\chi)}^\eta$, then let us fix a club C_δ of δ such that $\text{otp}(C_\delta) = \theta^*(\chi)$. Let $C_\delta = \{\alpha_i : i < \theta^*(\chi)\}$ be an increasing enumeration. Then $h \upharpoonright C_\delta$ induces a function $g_\delta : \theta^*(\chi) \rightarrow \chi$, given by $g_\delta(i) = h(\alpha_i)$.

We wish to show that $\text{ref}(h)$ is stationary in η . So, let C be a club of η . As we know that $S_{\theta^*(\chi)}^\eta$ is stationary in η , we can find a $\delta \in S_{\theta^*(\chi)}^\eta$ which is an accumulation point of C . Then $C \cap C_\delta$ is a club of δ , so $E \stackrel{\text{def}}{=} \{i < \theta^*(\chi) : \alpha_i \in C\}$ is a club in $\theta^*(\chi)$. Therefore, there is an $i \in E \cap \text{ref}(g_\delta)$, by the first part of this proof. Then $\alpha_i \in \text{ref}(h) \cap C$. \square

Remark 1.2a. One can ask the question of 1.2(3) in the opposite direction: suppose that σ reflects at some η , what can we say about the first such η ? This is an independence question, for more on this see [2].

We find it convenient to introduce the following

Definition 1.3. For ordinals $\eta > \chi > \aleph_0$, where χ is a regular cardinal, we define

$$\mathcal{S}[\eta, \chi] = \{A \subseteq \eta : \text{there is a function } h : \eta \rightarrow \chi \text{ such that for every } \delta \in A \cap S_\chi^\eta, \text{ there is a club } C \text{ of } \delta \text{ with } h \upharpoonright C \text{ strictly increasing}\}.$$

Saying that η has the strong non-reflection property for χ is equivalent to claiming that $\eta \in \mathcal{S}[\eta, \chi]$ or $S_\chi^\eta \in \mathcal{S}[\eta, \chi]$. In such a case, we say that $\mathcal{S}[\eta, \chi]$ is *trivial*.

$I(\sigma, \chi)$ is the statement: There is an $\eta \in (\chi, \sigma)$ with $\mathcal{S}[\eta, \chi]$ non-trivial.

Theorem 1.4. *Suppose that $\eta > \chi > \aleph_0$ and χ is a regular cardinal. Then $\mathcal{S}[\eta, \chi]$ is a χ -complete ideal on η .*

Proof. $\mathcal{S}[\eta, \chi]$ is obviously non-empty and downward closed.

Suppose that A_i ($i < i^* < \chi$) are sets from $\mathcal{S}[\eta, \chi]$, and that $h_i : \eta \rightarrow \chi$ witnesses that $A_i \in \mathcal{S}[\eta, \chi]$ for $i < i^*$. Let $A = \bigcup_{i < i^*} A_i$, and we shall see that $A \in \mathcal{S}[\eta, \chi]$. This will be exemplified by the function $h : \eta \rightarrow \chi$ defined by $h(\beta) \stackrel{\text{def}}{=} \sup\{h_i(\beta) : i < i^*\}$. Let $\delta \in A \cap S_\chi^\eta$ be given.

Then $\delta \in A_i$ for some $i < i^*$, and therefore there is a club C of δ such that $h_i \upharpoonright C$ is strictly increasing. Then it must be that $\text{otp}(C) = \text{cf}(\delta) = \chi$, and we can enumerate C increasingly as $\{\beta_\varepsilon : \varepsilon < \chi\}$. Therefore, the sequence $\langle h_i(\beta_\varepsilon) : \varepsilon < \chi \rangle$ is a strictly increasing sequence in χ , and $\chi = \sup_{\varepsilon < \chi} (h_i(\beta_\varepsilon))$.

On the other hand, for every ε , we have that $h(\beta_\varepsilon) < \chi$, so there is a minimal $\zeta(\varepsilon) < \chi$ such that $h(\beta_\varepsilon) < h_i(\beta_{\zeta(\varepsilon)})$ for some $i < i^*$. Let

$$E \stackrel{\text{def}}{=} \{\zeta < \rho : \bigwedge_{\varepsilon < \zeta} \zeta(\varepsilon) < \zeta\},$$

so E is a club of χ and $C^* \stackrel{\text{def}}{=} \{\beta_\zeta : \zeta \in E\}$ is a club of δ .

But then, for $\varepsilon_1 < \varepsilon_2 \in E$ we have $\zeta(\varepsilon_1) < \varepsilon_2$, so for some $i < i^*$ we have $h(\beta_{\varepsilon_1}) < h_i(\beta_{\zeta(\varepsilon_1)}) < h_i(\beta_{\varepsilon_2}) \leq h(\beta_{\varepsilon_2})$, so $h \upharpoonright C^*$ is strictly increasing. \square

If there is a square on σ^+ , then $I(\sigma^+, \chi)$ is false. As there are various notations in use, to make this statement precise, we state the definition of the \square principle that we use. Note that what we refer to as $\square(\sigma^+)$, some authors regard as \square_σ .

Definition 1.5. Suppose that χ and σ are cardinals. $\square_{(\chi, \sigma)}$ denotes the following statement: $\chi < \sigma$ and there is a sequence $\langle C_\delta : \delta \in (\text{SING} \cap \text{LIM}) \ \& \ \chi < \delta < \sigma \rangle$ such that:

- (1) C_δ is a club in δ .
- (2) $\text{otp}(C_\delta) < \delta$.
- (3) If δ is an accumulation point of C_α , then C_δ is defined and $C_\delta = \delta \cap C_\alpha$. In particular, $\delta > \chi$.

We use $\square(\chi^+)$ as a shorthand for $\square_{(\chi, \chi^+)}$. The sequence as above is called a $\square_{(\chi, \sigma)}$ -sequence, and its subsequences are called partial $\square_{(\chi, \sigma)}$ -sequences.

Observation 1.6. (0) Note that by a closed unbounded set of ω , we simply mean an unbounded subset of ω . Similarly, any δ with $\text{cf}(\delta) = \aleph_0$ will have a club subset consisting of an unbounded ω -sequence in δ . So, $\square(\omega_1)$ trivially holds.

- (1) If $\chi_1 \leq \chi_2 < \sigma$, then $\square_{(\chi_1, \sigma)} \Rightarrow \square_{(\chi_2, \sigma)}$. If $\chi < \sigma_1 \leq \sigma_2$, then $\square_{(\chi, \sigma_2)} \Rightarrow \square_{(\chi, \sigma_1)}$.

Theorem 1.7. *If $\theta > \chi$ is regular, then $\square_{(\chi, \theta)}$ implies that θ has the strong non-reflection property for χ .*

Proof. From [17, VII 1.7], we recall the following Fact 1.7a. For the reader's convenience, we also include the proof.

Fact 1.7a. *Assume that $\square_{(\chi, \theta)}$ holds. Then, there is a partial $\square_{(\chi, \theta)}$ -sequence*

$$\langle C_\delta : \delta \in (SING \cap LIM) \ \& \ \chi < \delta < \theta \ \& \ cf(\delta) < \chi \rangle,$$

such that for each δ for which C_δ is defined,

- (i) C_δ is a club subset of δ ,
- (ii) $otp(C_\delta) < \delta \ \& \ cf(\delta) < \chi \Rightarrow otp(C_\delta) < \chi$,
- (iii) $\gamma \in acc(C_\delta) \ \& \ C_\gamma \text{ defined} \Rightarrow C_\gamma = C_\delta \cap \gamma$.

Proof of Fact 1.7a. We start with a sequence $\langle D_\delta : \delta \in (SING \cap LIM) \ \& \ \chi < \delta < \theta \rangle$ which exemplifies $\square_{(\chi, \theta)}$.

We can without loss of generality assume that each D_δ satisfies $D_\delta \cap \chi = \emptyset$. For each δ for which D_δ is defined, we can define a 1–1 onto function $f_\delta : D_\delta \rightarrow otp(D_\delta)$ by

$$f_\delta(\alpha) \stackrel{\text{def}}{=} otp(D_\delta \cap \alpha).$$

Note that $\gamma \in acc(D_\delta) \Rightarrow f_\delta \upharpoonright D_\gamma = f_\gamma$. Now we define C_δ for $\delta < \theta$ with $cf(\delta) < \chi$ by induction on $\delta < \theta$.

If D_δ is not defined, then C_δ is not either.

If D_δ is defined, and $D_{otp(D_\delta)}$ is not defined, we set $C_\delta = D_\delta$. Note that $otp(D_\delta) < \chi$ in this case.

Finally, suppose that both D_δ and $D_{otp(D_\delta)}$ are defined. Then $\delta > otp(D_\delta) > \chi$ and $cf(otp(D_\delta)) = cf(\delta) < \chi$. So $C_{otp(D_\delta)}$ is already defined and we can define

$$C_\delta \stackrel{\text{def}}{=} \{ \alpha \in D_\delta : f_\delta(\alpha) \in C_{otp(D_\delta)} \}.$$

We can check that

$$\bar{C} \stackrel{\text{def}}{=} \langle C_\delta : \delta \in (SING \cap LIM) \ \& \ \chi < \delta < \theta \ \& \ cf(\delta) < \chi \rangle$$

is as required. One thing to note is that if $\gamma \in acc(C_\delta)$ and $D_{otp(D_\delta)}$ is defined, then $D_{otp(D_\gamma)}$ must be defined too. \square

Fixing a sequence \bar{C} like in Fact 1.7a, the following defines a function $h : \theta \rightarrow \chi$:

$$h(\delta) = \begin{cases} otp(C_\delta) & \text{if } \delta \in S_{< \chi}^\theta \setminus \chi, \\ 0 & \text{otherwise.} \end{cases}$$

Now, if $\delta \in S_\chi^\theta$, we can choose a club E_δ of δ which consists only of elements of $S_{< \chi}^\theta \setminus \chi$.

We let

$$D_\delta = \text{acc}(C_\delta) \cap E_\delta.$$

Then $h \upharpoonright D_\delta$ is increasing and D_δ is a club in δ . \square

We can also show, for example, that $I(\sigma, \chi^+)$ is consistently true if $\sigma > \chi^+$ and χ is regular. This follows from Fact 1.10 below. This fact also explains why we choose to call the properties under consideration “weak reflection” and “strong non-reflection”. There are other reflection properties that imply the weak reflection, like the reflections considered in [1] and elsewhere, so Fact 1.10 can be used as an example of a proof that the weak reflection is weaker than other reflection principles. Let us first recall the notion of stationary reflection.

Definition 1.8. (0) Suppose that S is a stationary subset of a cardinal θ with $\text{cf}(\theta) > \aleph_0$. We say that S *reflects* at $\delta \in \theta$ if $\text{cf}(\delta) > \aleph_0$ and $S \cap \delta$ is stationary in δ . We say that S is *reflecting* if there is $\delta \in \theta$ such that S reflects at δ . Otherwise, S is said to be *non-reflecting*.

(1) A regular cardinal θ is *reflecting* iff for every regular χ such that $\chi < \chi^+ < \theta$, every stationary $S \subseteq S_\chi^\theta$ is reflecting.

(2) For a regular cardinal θ , notation $REF(\theta)$ means that θ is reflecting. REF is the statement denoting that for every $\theta > \aleph_1$ which is a regular cardinal, $REF(\theta)$ holds.

(3) Suppose that $\lambda > \theta, \kappa$ are regular cardinals and $\kappa > \aleph_0$. We define the statement $\text{Ref}(\lambda, \kappa, \theta)$ to mean: For every $S \subseteq S_\delta^\lambda$ which is stationary, there is a δ of cofinality κ such that S reflects at δ .

Remark 1.9. (0) If δ is an ordinal of uncountable cofinality, then δ has a club subset consisting only of elements of cofinality $< \text{cf}(\delta)$. Therefore, if S reflects at δ , then S has to have elements of cofinality $< \text{cf}(\delta)$. This explains the gap of one between χ and θ in the definition of $REF(\theta)$.

REF is consistent modulo the existence of infinitely many supercompact cardinals [15]. The consistency of $\text{Ref}(\lambda, \kappa, \theta)$ has also been extensively studied, starting with a result of Baumgartner in [1] that $CON(\text{Ref}(\aleph_2, \aleph_1, \aleph_0))$ follows from the existence of a weakly compact cardinal.

(1) M. Magidor points out the following equivalent definition of the weak reflection, from which it is easy to see that it is weaker than what is usually meant by reflection: We can say that $\theta > \kappa$ weakly reflects at κ iff for any partition

$$S_{<\kappa}^\theta = \bigcup_{i < \kappa} S_i,$$

there is an $i < \kappa$ and an $\alpha \in S_\kappa^\theta$ such that $S_i \cap \alpha$ is stationary in α .

Fact 1.10. (0) Suppose that χ is a successor cardinal. Then

$REF(\chi^+) \Rightarrow \chi^+$ weakly reflects at χ .

(1) $Ref(\lambda, \kappa, \theta) \Rightarrow \lambda \notin \mathcal{S}[\lambda, \kappa]$.

Proof. (0) Let $\chi = \sigma^+$. So, suppose for contradiction that $h : \chi^+ \rightarrow \chi$ is a function such that for every $\delta \in S_\chi^{\chi^+}$, there is a club of C_δ with $h \upharpoonright C_\delta$ 1–1.

Now, h is regressive on (χ, χ^+) . Since $(\chi, \chi^+) \cap S_\sigma^{\chi^+}$ is stationary in χ^+ , there is a stationary $S \subseteq (\chi, \chi^+) \cap S_\sigma^{\chi^+}$ such that $h \upharpoonright S$ is a constant.

By $REF(\chi^+)$, there is a $\delta \in \chi^+$ such that $S \cap \delta$ is stationary in δ . As in 1.9, we conclude that $cf(\delta) > \sigma$. Therefore $cf(\delta) = \chi$ and C_δ is defined. But then $S \cap C_\delta$ is stationary in δ , and $h \upharpoonright (S \cap C_\delta)$ is constant.

(1) Similar. \square

We now go on to present the last two facts before we proceed to the Main Theorem.

Fact 1.11. Suppose that $cf(\delta) > \aleph_0$ and $f : \delta \rightarrow \delta$ is a function which is not increasing on any club of δ . Then there is a stationary set S in δ such that $\alpha \in S \Rightarrow f(\alpha) < \text{Min}(S)$. (Hence, there is also a stationary subset of δ on which f is a constant.)

Proof. We fix an increasing continuous sequence of ordinals $\langle \alpha_\varepsilon : \varepsilon < cf(\delta) \rangle$ which is cofinal in δ . Let $T \stackrel{\text{def}}{=} \{\varepsilon : f(\alpha_\varepsilon) < \alpha_\varepsilon\}$. So $T \subseteq cf(\delta)$.

We shall see that T is stationary in $cf(\delta)$. Let us first assume that it is true, and define for $\varepsilon \in T \setminus \{0\}$,

$$g(\varepsilon) \stackrel{\text{def}}{=} \text{Min}\{\zeta : f(\alpha_\varepsilon) < \alpha_{\zeta+1}\}.$$

Therefore, g is regressive, and we can find a stationary $T_1 \subseteq T$ such that $g \upharpoonright T_1$ is constantly equal to some ζ^* .

Let $S \stackrel{\text{def}}{=} \{\alpha_\varepsilon : \varepsilon \in T_1 \setminus (\zeta^* + 2)\}$. We can check that this S is as required.

It remains to be seen that T is stationary in $cf(\delta)$. Suppose not. Then we can find a club C in $cf(\delta)$ such that $C \cap T = \emptyset$. Let

$$E \stackrel{\text{def}}{=} \{\varepsilon : \varepsilon \in LIM \cap C \ \& \ (\forall \zeta < \varepsilon) (f(\alpha_\zeta) < \alpha_\varepsilon)\}.$$

Then E is also a club of $cf(\delta)$. But then $D \stackrel{\text{def}}{=} \{\alpha_\varepsilon : \varepsilon \in E\}$ is a club of δ and $f \upharpoonright D$ is strictly increasing, contradicting our assumption. \square

Remark 1.12. As a remark on the side: we cannot improve the previous result to conclude, from the assumptions given above, that there is a club C of δ such that $f \upharpoonright C$ is constant. Namely, if we take a stationary costationary set S in δ , and define

f on δ by

$$f(\alpha) = \begin{cases} 1 & \text{if } \alpha \in S, \\ 0 & \text{otherwise,} \end{cases}$$

then this f is neither increasing nor constant on any club of δ . In the following Fact 1.12a we recall a more general result along the lines of Fact 1.11. This fact is not used in the proof of the Main Theorem 1.13, and a reader who is in a hurry may without loss of continuity proceed directly to 1.13.

Fact 1.12a. *Suppose that δ is an ordinal with $\text{cf}(\delta) > \aleph_0$ and f is a function from δ to the ordinals. Then, there is a stationary $S \subseteq \delta$ such that*

- either $f \upharpoonright S$ is constant,
or $f \upharpoonright S$ is strictly increasing.

Proof. Let $\theta = \text{cf}(\delta) > \aleph_0$, and $\langle \alpha_\varepsilon : \varepsilon < \theta \rangle$ a strictly increasing enumeration of a club of δ . Let $E \stackrel{\text{def}}{=} \theta \cap \text{LIM}$. We define a partial function $g : E \rightarrow \theta$ as follows:

$$g(\varepsilon) = \zeta \text{ if } \begin{cases} \text{(a) } f(\alpha_\varepsilon) \leq f(\alpha_\zeta), \\ \text{(b) } (\forall \xi < \varepsilon)(f(\alpha_\varepsilon) \leq f(\alpha_\xi) \Rightarrow f(\alpha_\xi) \leq f(\alpha_\zeta)), \\ \text{(c) } \zeta < \varepsilon \text{ is minimal under (a) and (b).} \end{cases}$$

Note that $g(\varepsilon) < \varepsilon$ for all $\varepsilon \in \text{Dom}(g)$. Now we consider three cases:

Case 1: $S_0 \stackrel{\text{def}}{=} E \setminus \text{Dom}(g)$ is stationary in θ . Then $S \stackrel{\text{def}}{=} \{\alpha_\varepsilon : \varepsilon \in S_0\}$ is stationary in δ . We claim that $f \upharpoonright S$ is strictly increasing on S . Otherwise, there would be an $\varepsilon \in S$ such that there is a $\zeta_0 < \varepsilon$ with $f(\alpha_\varepsilon) \leq f(\alpha_{\zeta_0})$, which contradicts the fact that $\varepsilon \notin \text{Dom}(g)$.

Case 2: $\text{Dom}(g)$ is stationary in θ and $S_1 \stackrel{\text{def}}{=} \{\varepsilon \in \text{Dom}(g) : f(\alpha_\varepsilon) = f(\alpha_{g(\varepsilon)})\}$ is stationary in θ . Since $g \upharpoonright S_1$ is regressive, there is a stationary $S_2 \subseteq S_1$ such that $g \upharpoonright S_2$ is constant. Then $S \stackrel{\text{def}}{=} \{\alpha_\varepsilon : \varepsilon \in S_2\}$ is stationary in δ , and $f \upharpoonright S$ is constant.

Case 3: $\text{Dom}(g)$ is stationary in θ , but S_1 is not stationary. Then $S_3 \stackrel{\text{def}}{=} \text{Dom}(g) \setminus S_1 = \{\varepsilon \in \text{Dom}(g) : f(\alpha_\varepsilon) < f(\alpha_{g(\varepsilon)})\}$ is stationary, and there is a stationary $S_4 \subseteq S_3$ such that $g \upharpoonright S_4$ is a constant. Let $S = \{\alpha_\varepsilon : \varepsilon \in S_4\}$, so S is a stationary subset of δ . We claim that $f \upharpoonright S$ is strictly increasing.

Otherwise, there are $\varepsilon_1 < \varepsilon_2 \in S_4$ such that $f(\alpha_{\varepsilon_2}) \leq f(\alpha_{\varepsilon_1})$. On the other hand, $f(\alpha_{\varepsilon_1}) < f(\alpha_{g(\varepsilon_1)}) = f(\alpha_{g(\varepsilon_2)})$, since both ε_1 and ε_2 are members of S_4 . This contradicts the definition of $g(\varepsilon_2)$, since $\varepsilon_1 < \varepsilon_2$. \square

We now present our main result.

Main Theorem 1.13. *Assume that $\lambda = \mu^+$ and $\mu > \text{cf}(\mu) = \kappa > \aleph_0$. In addition, for some $\theta \in (\kappa, \lambda)$, we know that θ has the weak reflection property for κ . Then $\mathcal{D}_\lambda \upharpoonright S_\kappa^\lambda$ is not λ^+ -saturated.*

Proof. By 1.2(3), without loss of generality, θ is a regular cardinal, so $\theta < \mu$.

Fact 1.13a. *There is a stationary subset S of S_θ^λ such that: For some $S^+ \supseteq S$ and $S^+ \subseteq \lambda \setminus \kappa$, there is a sequence*

$$\bar{C} = \langle C_\alpha : \alpha \in S^+ \rangle,$$

such that, for every $\alpha \in S^+$:

- (1) C_α is a subset of $\alpha \setminus \kappa$ and $\text{otp}(C_\alpha) \leq \theta$.
- (2) α is a limit ordinal $\Rightarrow \sup(C_\alpha) = \alpha$.
- (3) $\beta \in C_\alpha \Rightarrow (\beta \in S^+ \text{ and } C_\beta = C_\alpha \cap \beta)$.
- (4) $\text{cf}(\alpha) = \theta$ iff $\alpha \in S$.
- (5) For every club E in λ , there are stationarily many $\delta \in S$ such that for all α, β ,

$$(\alpha < \beta \ \& \ \alpha, \beta \in C_\delta) \Rightarrow (\alpha, \beta] \cap E \neq \emptyset.$$

Proof of Fact 1.13a. Since $\theta^+ < \lambda$ we can apply [16, 1.5], so there is a stationary subset S_1 of $S_\theta^\lambda \setminus \{\theta\}$ with $S_1 \in I[\lambda]$. By [16, 1.2], this means that there is a sequence $\langle D_\alpha : \alpha < \lambda \rangle$ such that:

- (a) D_α is a closed subset of α .
- (b) $\alpha^* \in \text{nacc}(D_\alpha) \Rightarrow D_{\alpha^*} = D_\alpha \cap \alpha^*$.
- (c) For some club E of λ , for every $\delta \in S_1 \cap E$,

$$\delta = \sup(D_\delta) \quad \& \quad \text{otp}(D_\delta) = \theta.$$

- (d) $\text{nacc}(D_\alpha)$ is a set of successor ordinals.

Observation: We do not lose generality if we in addition require that for each α , $\text{otp}(D_\alpha) \leq \theta$.

[Why? Let E be the club guaranteed by (c). Since S_1 is stationary, so is $S_1 \cap E$, so we can define for $\alpha \in \lambda$

$$D_\alpha^\dagger = \begin{cases} D_\alpha & \text{if } \alpha \in S_1 \cap E \\ & \text{or } \exists \beta > \alpha (\beta \in S_1 \cap E \ \& \ \alpha \in \text{nacc}(D_\beta)), \\ \emptyset & \text{otherwise.} \end{cases}$$

Then we can set $S_2 \stackrel{\text{def}}{=} S_1 \cap E$, so the sequence $\langle D_\alpha^\dagger : \alpha < \lambda \rangle$ will satisfy (a)–(d), with S_1 replaced by S_2 , and $\text{otp}(D_\alpha^\dagger) \leq \theta$ will hold for each α .

Continuation of the proof of Fact 1.13a: Now, for any club F of λ , we let

$$\bar{C}[F] \stackrel{\text{def}}{=} \langle C_\delta[F] : \delta \in S^+[F] \stackrel{\text{def}}{=} S_1 \cap E \cap \text{acc}(F) \cup (\text{successors}) \setminus \kappa \rangle,$$

where

$$C_\delta[F] \stackrel{\text{def}}{=} \{ \gamma \in \text{nacc}(D_\delta) \setminus \kappa : F \cap [\sup(\gamma \cap D_\delta), \gamma) \neq \emptyset \}.$$

We claim that \bar{C} can be set to be equal to $\bar{C}(F)$ for some club F , with $S = S[F] \stackrel{\text{def}}{=} E \cap S_1 \cap \text{acc}(F) \setminus \kappa$ and $S^+ = S^+[F]$. We are using the ideas of [19, Section 2].

It is easily checked that (1)–(4) are satisfied for any club F . So, let us suppose that (5) is not satisfied for any choice of F , and we shall obtain a contradiction.

By induction on $\zeta < \theta^+$, we define a club F_ζ of λ .

If $\zeta = 0$, we let $F_\zeta = LIM$.

If ζ is a limit ordinal, we let $F_\zeta = \bigcap_{\xi < \zeta} F_\xi$. This is still a club of λ , as $\zeta < \theta^+ < \lambda$.

If $\zeta = \xi + 1$, we have assumed that there is a club F_ζ such that the set

$$G_\xi[F_\zeta] \stackrel{\text{def}}{=} \{\delta \in S[F_\zeta] : \forall \alpha, \beta \in C_\delta[F_\xi] (\alpha < \beta \Rightarrow (\alpha, \beta] \cap F_\zeta \neq \emptyset)\}$$

is non-stationary. Without loss of generality, we assume that $F_\zeta \subseteq F_\xi$.

At the end, let us let $C = \bigcap_{\zeta < \theta^+} F_\zeta$. This is still a club of λ , as $\theta^+ < \lambda$. Since $S \stackrel{\text{def}}{=} S[C]$ is stationary in λ , and for every $\zeta < \theta^+$, we have that $S \subseteq S[F_\zeta]$, we conclude that the set

$$T \stackrel{\text{def}}{=} C \cap S \setminus \left(\bigcup_{\zeta < \theta^+} G_\zeta[F_{\zeta+1}] \right)$$

is stationary. So, let us take a $\delta \in T \cap LIM$. Then D_δ is a club of δ , since $T \cap LIM \subseteq S_1 \cap E$.

For $\beta \in D_\delta$, we consider the sequence

$$\langle \sup(\beta \cap F_\zeta) : \zeta < \theta^+ \rangle.$$

This is a non-increasing sequence of ordinals $\leq \beta$, so there must be a $\zeta_\beta < \theta^+$ and $\gamma_\beta \leq \beta$ such that

$$\zeta_\beta \leq \zeta < \theta^+ \Rightarrow \sup(\beta \cap F_\zeta) = \gamma_\beta.$$

Notice that $\zeta^* \stackrel{\text{def}}{=} \sup\{\zeta_\beta : \beta \in D_\delta\} < \theta^+$, since $\text{otp}(D_\delta) = \theta$.

Since $\delta \notin G_{\zeta^*}[F_{\zeta^*+1}]$, by the definition of $G_{\zeta^*}[F_{\zeta^*+1}]$, there are $\alpha < \beta \in C_\delta[F_{\zeta^*}]$ such that

$$(\alpha, \beta] \cap F_{\zeta^*+1} = \emptyset.$$

On the other hand, $\beta \in C_\delta[F_{\zeta^*}] \Rightarrow [\sup(\beta \cap D_\delta), \beta) \cap F_{\zeta^*} \neq \emptyset \Rightarrow \sup(\beta \cap D_\delta) \leq \sup(\beta \cap F_{\zeta^*}) = \gamma_\beta = \sup(\beta \cap F_{\zeta^*+1})$. But $\alpha < \beta$ and $\alpha \in \text{nacc}(D_\delta)$, so $\sup(\beta \cap D_\delta) \geq \alpha$. Therefore $\alpha \leq \gamma_\beta \leq \beta$. Note now that γ_β must be a limit ordinal, since $F_{\zeta^*+1} \subseteq LIM$. But α is a successor, by (d) in the definition of D_δ . So, $\alpha < \gamma_\beta \leq \beta \in F_{\zeta^*+1}$ and $(\alpha, \beta] \cap F_{\zeta^*+1} = \emptyset$, by the choice of F_{ζ^*} . A contradiction. \square

Continuation of the proof of Main Theorem 1.13: Let us fix S, S^+ and $\bar{C} \stackrel{\text{def}}{=} \langle C_\alpha : \alpha \in S^+ \rangle$ as in Fact 1.13a. Denote by $\text{cl}(\bar{C})$ the following:

$$\text{cl}(\bar{C}) = \{C : C \subseteq S^+ \wedge \forall \beta \in C (C_\beta = C \cap \beta)\}.$$

Now fix the following enumerations: For $\alpha \in S^+$, let

$$C_\alpha = \{\gamma(\alpha, \varepsilon) : \varepsilon < \text{otp}(C_\alpha)\},$$

such that $\gamma(\alpha, \varepsilon)$ is increasing in ε . For each ε , let

$$\gamma^*(\alpha, \varepsilon) \stackrel{\text{def}}{=} \bigcup_{\xi \leq \varepsilon} \gamma(\alpha, \xi).$$

Let $\mu = \sum_{i < \kappa} \mu_i$ be such that $\langle \mu_i : i \in \kappa \rangle$ is a continuous increasing sequence of cardinals, and, for simplicity, $\mu_0 > \theta$.

Claim 1.13b. *Given enumerations as above, we can for each $\alpha < \lambda$, find sets $a_i^\alpha (i < \kappa)$ such that*

- (I) $\alpha = \bigcup_{i < \kappa} a_i^\alpha$.
- (II) $|a_i^\alpha| \leq \mu_i$.
- (III) $i < j \Rightarrow a_i^\alpha \subseteq a_j^\alpha$ and for i a limit ordinal $< \kappa$,

$$a_i^\alpha = \bigcup_{j < i} a_j^\alpha.$$

- (IV) $\beta \in a_i^\alpha \Rightarrow a_i^\beta \subseteq a_i^\alpha$.
- (V) $\beta \in a_i^\alpha \cap S^+ \Rightarrow C_\beta \subseteq a_i^\alpha$.

Proof of Claim 1.13b. It is easy to see that we can choose a_i^α for $\alpha < \lambda$ and $i < \kappa$ such that (I)–(III) are satisfied. Suppose we have done so. Then define by induction on $\alpha < \lambda$, and then by induction on $j < \kappa$, sets

$$a_i^{\alpha^\dagger} \stackrel{\text{def}}{=} a_i^\alpha \cup \{a_i^{\beta^\dagger} : \beta \in a_i^\alpha \cup \bigcup \{C_\beta : \beta \in a_i^\alpha \cap S^+\}\} \cup \bigcup \{C_\beta : \beta \in a_i^\alpha \cap S^+\}.$$

Now we can check that, by renaming $a_i^\alpha = a_i^{\alpha^\dagger}$, we satisfy the claim. \square

Continuation of the proof of Main Theorem 1.13: So we now fix a_i^α as in Claim 1.13b.

Similarly, for each $\xi < \lambda^+$, we can let $\xi = \bigcup_{\alpha < \lambda} b_\alpha^\xi$, where b_α^ξ are \subseteq -increasing in α , while $|b_\alpha^\xi| < \lambda$ and, if $\zeta \in b_\alpha^\xi$, then $b_\alpha^\zeta \subseteq b_\alpha^\xi$. We require in addition that, if $\xi = \zeta + 1$, then $\zeta \in b_0^\xi$, and if $\text{cf}(\xi) < \lambda$, then $\xi = \sup(b_0^\xi)$.

We now define, for $\xi < \lambda^+$, a function $h_\xi : \lambda \rightarrow \lambda$. This function is given by $h_\xi(\alpha) \stackrel{\text{def}}{=} \text{otp}(b_\alpha^\xi)$.

Clearly, each h_ξ is non-decreasing, and

$$\zeta \in b_\alpha^\xi \Rightarrow h_\zeta(\alpha) < h_\xi(\alpha). \tag{a}$$

Now, fix a sequence $\bar{E} = \{E_\eta : \eta \in S_\kappa^\lambda\}$, where each E_η is a club subset of η with $\text{otp}(E_\eta) = \kappa$ and consisting only of elements of cofinality $< \kappa$. We prove the following claim, with the intention of applying it later on the functions $h_\xi (\xi < \lambda^+)$.

Claim 1.13c. *Suppose that $h : \lambda \rightarrow \lambda$ is non-decreasing, and \bar{E} is given as above. Then, there is an $i = i(h) < \kappa$ such that for stationarily many $\eta \in S_\kappa^\lambda$, there is a*

$C = C(\eta) \in \text{cl}(\bar{C})$ with:

- (i) $C \subseteq \eta = \sup(C)$.
- (ii) $C \subseteq \bigcup_{\beta \in E_\eta} a_i^\beta$ (hence $\bigcup_{\gamma \in C} a_i^\gamma \subseteq \bigcup_{\beta \in E_\eta} a_i^\beta$, by the choice of a 's).
- (iii) $\eta = \sup\{\alpha \in C : h(\alpha) \in \bigcup_{\beta \in E_\eta} a_i^\beta \text{ and } h(\sup(C \cap \alpha)) \in \bigcup_{\beta \in E_\eta} a_i^\beta\}$.

Proof of Claim 1.13c. Recall the definition of S and C_δ 's from Claim 1.13a. In particular, for all $\delta \in S \subseteq S_\theta^\zeta$,

$$C_\delta = \{\gamma(\delta, \varepsilon) : \varepsilon < \text{otp}(C_\delta) = \theta\}.$$

Let us first fix a $\delta \in S$ and suppose that for all $\alpha \in C_\delta$,

$$h(\alpha) < \text{Min}(C_\delta \setminus (\alpha + 1)). \quad (\text{b})$$

We define a function $f = f_\delta : \theta \rightarrow \kappa$ by:

If $\varepsilon \in (\theta \setminus S_{<\kappa}^\theta)$, $f(\varepsilon) = 0$.

If $\varepsilon \in S_{<\kappa}^\theta$, $f(\varepsilon) = \text{Min}\{i < \kappa : h(\gamma^*(\delta, \varepsilon)), h(\gamma(\delta, \varepsilon)) \in a_i^{\gamma(\delta, \varepsilon+1)} \text{ and } a_i^{\gamma^*(\delta, \varepsilon)} \cap C_\delta \text{ is unbounded in } \gamma^*(\delta, \varepsilon)\}$.

Remember that all $\varepsilon \in S_{<\kappa}^\theta$ satisfy $\aleph_0 \leq \text{cf}(\gamma^*(\delta, \varepsilon)) < \kappa$. By the definition of $\gamma^*(\delta, \varepsilon)$, we have that $C_\delta \cap \gamma^*(\delta, \varepsilon)$ is unbounded in $\gamma^*(\delta, \varepsilon)$. Then, as $\langle a_i^{\gamma^*(\delta, \varepsilon)} : i < \kappa \rangle$ is a \subseteq -increasing sequence of sets with union $\gamma^*(\delta, \varepsilon)$, it must be that $C_\delta \cap a_i^{\gamma^*(\delta, \varepsilon)}$ is eventually unbounded in $\gamma^*(\delta, \varepsilon)$.

On the other hand, since h is non-decreasing and $\gamma^*(\delta, \varepsilon) \leq \gamma(\delta, \varepsilon) < \gamma(\delta, \varepsilon + 1)$, we have by (b) (at the beginning of this proof) that $h(\gamma^*(\delta, \varepsilon)), h(\gamma(\delta, \varepsilon)) < \gamma(\delta, \varepsilon + 1)$, so $h(\gamma^*(\delta, \varepsilon)), h(\gamma(\delta, \varepsilon)) \in a_i^{\gamma(\delta, \varepsilon+1)}$ for every large enough $i < \kappa$.

So, $f(\varepsilon)$ is well defined.

We have assumed that θ is weakly reflective for κ . So, by Observation 1.2(4), for stationarily many $\varepsilon \in S_\kappa^\theta$, f is not strictly increasing on any club of $\gamma^*(\delta, \varepsilon)$. If we take any such ε , then $\text{cf}(\gamma^*(\delta, \varepsilon)) > \aleph_0$ and $f \upharpoonright \gamma^*(\delta, \varepsilon) : \gamma^*(\delta, \varepsilon) \rightarrow \gamma^*(\delta, \varepsilon)$, as $\gamma^*(\delta, \varepsilon) > \kappa$. So Fact 1.11 applies and f is constant on some stationary subset of $\gamma^*(\delta, \varepsilon)$. Let us denote that constant value by $i_\varepsilon = i_\delta(\varepsilon)$. Let $\eta \stackrel{\text{def}}{=} \gamma^*(\delta, \varepsilon)$.

Observation: The set $\bigcup_{\alpha \in E_\eta} a_i^\alpha$ is an unbounded (even stationary) subset of $C_\delta \cap \gamma^*(\delta, \varepsilon)$.

[Why? We can check that $e_\varepsilon \stackrel{\text{def}}{=} \{\zeta < \varepsilon : \gamma^*(\delta, \zeta) \in E_\eta\}$ is a club of ε , so $s_\varepsilon \stackrel{\text{def}}{=} \{\zeta \in e_\varepsilon : f(\zeta) = i_\varepsilon\} \subseteq e_\varepsilon$ is stationary, and we conclude that $\{\gamma(\delta, \zeta) : \zeta \in s_\varepsilon\} \subseteq \gamma^*(\delta, \varepsilon)$ is stationary in $\gamma^*(\delta, \varepsilon)$. Note that for each $\zeta \in e_\varepsilon$, by the choice of \bar{E} , we have that $\text{cf}(\gamma^*(\delta, \zeta)) < \kappa$.

Now we take any $\zeta \in s_\varepsilon$. Since $\text{cf}(\gamma^*(\delta, \zeta)) < \kappa$, by the definition of f , we have that $a_{i_\varepsilon}^{\gamma^*(\delta, \zeta)} \cap C_\delta$ is unbounded in $\gamma^*(\delta, \zeta)$. Then

$$\bigcup_{\alpha \in E_\eta} a_i^\alpha \cap C_\delta \supseteq \bigcup_{\gamma^*(\delta, \zeta) \in s_\varepsilon} a_{i_\varepsilon}^{\gamma^*(\delta, \zeta)} \cap C_\delta,$$

and since s_ε is stationary in $\gamma^*(\delta, \varepsilon)$, we derive the desired conclusion.]

Continuation of the proof of Claim 1.13c: So, by (V) in the choice of a 's and (3) of Fact 1.13a, we conclude that $\bigcup_{\alpha \in E_\eta} a_\alpha^x$ contains the entire $C_\delta \cap \gamma^*(\delta, \varepsilon)$. Hence, by (IV) of 1.13b,

$$\bigcup_{\alpha \in E_\eta} a_\alpha^x \supseteq \bigcup \{a_{i_\varepsilon}^\beta : \beta \in C_\delta \cap \gamma^*(\delta, \varepsilon)\}.$$

Now, there must be some $j = j_\delta < \kappa$, such that $\{\varepsilon \in S_\kappa^\theta : i_\varepsilon \text{ is well defined and } = j\}$ is a stationary subset of θ .

If we let δ vary, then j_δ is defined for every $\delta \in S_\theta^\lambda$ which satisfies (b). We show that the set of such δ is stationary in S_θ^λ .

We now get to use (5) from 1.13a. Namely,

$$E = \{\delta < \lambda : (\forall \gamma < \delta)(h(\gamma) < \delta)\}$$

is a club in λ . Therefore,

$$T = T_h \stackrel{\text{def}}{=} \{\delta \in S : \forall \alpha < \beta (\alpha, \beta \in C_\delta \Rightarrow (\alpha, \beta] \cap E \neq \emptyset)\}$$

is stationary. It is easily seen that every element of T satisfies (b), so the set of all $\delta \in S$ for which j_δ is defined is stationary.

Now, for some $i(*) < \kappa$, the set $\{\delta \in S : j_\delta = i(*)\}$ is stationary. Therefore,

$$\{\eta = \gamma^*(\delta, \varepsilon) : \varepsilon \in S_\kappa^\lambda \text{ \& } i_\delta(\varepsilon) \text{ is defined and } = i(*)\}$$

is stationary in λ . For every such $\eta = \gamma^*(\delta, \varepsilon)$, we define $C = C(\eta)$ by $C \stackrel{\text{def}}{=} C_\delta \cap \gamma^*(\delta, \varepsilon)$.

We can easily check that this is a well posed definition and that $i = i(*)$ is as required. \square

Continuation of the proof of Main Theorem 1.13: Now we apply the previous claim to h_ζ ($\zeta < \lambda^+$). For every $\zeta < \lambda^+$, we fix $i(\zeta) < \kappa$ as guaranteed by the claim. Then note that for some $i(*) < \kappa$, the set

$$W = \{\zeta < \lambda^+ : i(\zeta) = i(*)\}$$

is unbounded in λ^+ . Of course, we can in fact assume W to be stationary, but we only need it to be unbounded.

We now define a new family of functions, based on the h_ζ 's.

For every $\zeta \in W$ and $\eta \in S_\kappa^\lambda$ let $h_{\zeta, \eta}$ be the function with domain $a_\eta^* \stackrel{\text{def}}{=} \bigcup_{\alpha \in E_\eta} a_{i(*)}^\alpha$, defined by

$$h_{\zeta, \eta}(\beta) = \text{Min}(a_\eta^* \setminus h_\zeta(\beta)).$$

Observe that $a_\eta^* \subseteq \eta$ and $\eta = \sup(a_\eta^*)$. We have noted before that for $\zeta < \xi < \lambda$, we can fix an $\alpha_{\zeta, \xi} \in \lambda$ such that

$$h_\zeta \upharpoonright [\alpha_{\zeta, \xi}, \lambda) < h_\xi \upharpoonright [\alpha_{\zeta, \xi}, \lambda).$$

So, if $\zeta \leq \xi \in W$, then

$$\alpha \in [\alpha_{\zeta, \xi}, \lambda) \Rightarrow h_{\zeta, \eta}(\alpha) \leq h_{\xi, \eta}(\alpha).$$

Also, if $\zeta < \xi \in W$, then

$$h_{\zeta}(\beta) < h_{\xi}(\beta) \wedge h_{\zeta, \eta}(\beta) = h_{\xi, \eta}(\beta) \Rightarrow [h_{\zeta}(\beta), h_{\xi}(\beta)) \neq \emptyset \ \& \ [h_{\zeta}(\beta), h_{\xi}(\beta)) \cap a_{\eta}^* = \emptyset.$$

Using the above functions, we define the following sets $A_{\zeta, \xi}$ for $\zeta < \xi \in W$.

$$A_{\zeta, \xi} \stackrel{\text{def}}{=} \{\eta \in S_{\kappa}^{\lambda} : \text{for unboundedly } \beta \in a_{\eta}^*, \text{ we have } h_{\zeta, \eta}(\beta) < h_{\xi, \eta}(\beta)\}.$$

We now show that these sets witness that $\mathcal{D}_{\lambda} \upharpoonright S_{\kappa}^{\lambda}$ is not λ^+ -saturated.

Note:

(A) *If $\zeta < \xi \in W$, then $A_{\zeta, \xi}$ is a stationary subset of λ .*

[Why? To see this, fix such $\zeta < \xi$ and suppose that E is a club in λ which misses $A_{\zeta, \xi}$.

Suppose that $\eta > \alpha_{\zeta, \xi}$ and $\eta \in S_{\kappa}^{\lambda} \cap E$. Then $\eta \notin A_{\zeta, \xi}$, so there is $\beta_0 < \sup(a_{\eta}^*) = \eta$, such that for all $\beta \in (a_{\eta}^* \setminus \beta_0)$,

$$h_{\zeta, \eta}(\beta) = h_{\xi, \eta}(\beta),$$

or, equivalently,

$$[h_{\zeta}(\beta), h_{\xi}(\beta)) \cap a_{\eta}^* = \emptyset.$$

We can also assume $\beta_0 \geq \alpha_{\zeta, \xi}$, so $[h_{\zeta}(\beta), h_{\xi}(\beta)) \neq \emptyset$. In particular, $h_{\zeta}(\beta) \notin a_{\eta}^*$. We can further assume that η satisfies the conclusion of Claim 1.13c, with $h = h_{\zeta}$ and $i = i(*)$ (since $\zeta \in W$). Let C be as there.

But then the conclusion of Claim 1.13c tells us that we can find a β in C which is greater than β_0 and such that $h_{\zeta}(\beta) \in a_{\eta}^*$.]

(B) *If $\zeta_1 \leq \zeta_2 \leq \xi_2 \leq \xi_1 \in W$, then $A_{\zeta_2, \xi_2} \setminus A_{\zeta_1, \xi_1}$ is bounded.*

[Why? This follows easily by the remarks after the definition of $h_{\zeta, \eta}$.]

Now assume, for contradiction, that $\mathcal{D}_{\lambda} \upharpoonright S_{\kappa}^{\lambda}$ is λ^+ -saturated. Then, by (B):

(C) *For each $\zeta \in W$, we have $\langle A_{\zeta, \xi} / \mathcal{D}_{\lambda} : \xi \in (\zeta, \lambda^+) \cap W \rangle$ is eventually constant, say for $\xi \in [\xi_{\zeta}, \lambda^+)$.*

So, $A_{\zeta} \stackrel{\text{def}}{=} A_{\zeta, \xi_{\zeta}}$ satisfies $\zeta_1 < \zeta_2 \in W \Rightarrow A_{\zeta_1} \supseteq A_{\zeta_2} \pmod{\mathcal{D}_{\lambda}}$ (again by (B)). Hence, again by the λ^+ -saturation of $\mathcal{D}_{\lambda} \upharpoonright S_{\kappa}^{\lambda}$,

(D) *$\langle A_{\zeta} / \mathcal{D}_{\lambda} : \zeta \in W \rangle$ is eventually constant, say for $\zeta \geq \zeta^*$.*

Choose $\zeta_{\varepsilon} \in W \setminus \zeta^*$ for $\varepsilon < \mu_{i(*)}^+$ such that

$$\varepsilon(1) < \varepsilon(2) \Rightarrow \zeta_{\varepsilon(1)} < \zeta_{\varepsilon(2)},$$

which is possible since W is unbounded. By (a) after the definition of h_{ξ} , we can find an $\alpha^* < \lambda$ such that

$$\varepsilon(1) < \varepsilon(2) < \mu_{i(*)}^+ \wedge \alpha^* \leq \alpha < \lambda \Rightarrow h_{\zeta_{\varepsilon(1)}}(\alpha) < h_{\zeta_{\varepsilon(1)}}(\alpha) < h_{\zeta_{\varepsilon(2)}}(\alpha). \quad (\oplus)$$

By clauses (C) and (D), for all $\zeta, \xi \in W$ and $\varepsilon < \mu_{i(*)}^+$,

$$A_{\zeta^*} = A_{r_\varepsilon} = A_{r_{\varepsilon, \zeta}} = A_{r_{\varepsilon, \zeta_{\varepsilon+1}}} \neq \emptyset \pmod{\mathcal{D}_\lambda}.$$

By (C) and (D),

$$\bigcap_{\varepsilon < \mu_{i(*)}^+} A_{r_{\varepsilon, \zeta_{\varepsilon+1}}} = A_{\zeta^*} \pmod{\mathcal{D}_\lambda}.$$

So,

$$A \stackrel{\text{def}}{=} \bigcap_{\varepsilon < \mu_{i(*)}^+} A_{r_{\varepsilon, \zeta_{\varepsilon+1}}} \neq \emptyset \pmod{\mathcal{D}_\lambda}.$$

Now we can choose an $\eta \in A \setminus (\alpha^* + 1)$, hence by \oplus ,

$$\varepsilon(1) < \varepsilon(2) < \mu_{i(*)}^+ \wedge \beta \in a_\eta^* \Rightarrow h_{\eta, \varepsilon(1)}(\beta) \leq h_{\eta, \varepsilon(2)}(\beta). \quad (\oplus_1)$$

For each $\beta \in a_\eta^* \setminus (\alpha^* + 1)$, the sequence $\langle h_{\eta, \varepsilon}(\beta) : \varepsilon < \mu_{i(*)}^+ \rangle$ is non-decreasing. Hence, since $|a_\eta^*| \leq \mu_{i(*)}$, for some $\varepsilon(\beta) < \mu_{i(*)}^+$, the sequence $\langle h_{\eta, \varepsilon}(\beta) : \varepsilon(\beta) \leq \varepsilon < \mu_{i(*)}^+ \rangle$ is constant. Let $\varepsilon(*) \stackrel{\text{def}}{=} \sup_{\beta \in a_\eta^* \setminus (\alpha^* + 1)} (\varepsilon(\beta)) < \mu_{i(*)}^+$. But $\eta \in A_{r_{\varepsilon(*), \zeta_{\varepsilon(*)+1}}}$, hence by the definition of $A_{r_{\zeta, \xi}}$'s, there is a $\beta \in a_\eta^* \setminus (\alpha^* + 1)$ such that

$$h_{\zeta_{\varepsilon(*), \eta}}(\beta) < h_{\zeta_{\varepsilon(*)+1, \eta}}(\beta). \quad (\oplus_2)$$

So we get a contradiction. \square

A similar argument can be applied to other normal filters \mathcal{D} on λ , under certain conditions. In addition, we shall see that under some assumptions on the cardinal arithmetic, the fact that \mathcal{D} is not λ^+ -saturated is strongly witnessed, by the existence of a \diamond on \mathcal{D} . That is, we obtain $\diamond_{\mathcal{D}}^*(S_\kappa^\lambda)$. This notation is explained in the following

Definition 1.14. If \mathcal{D} is a normal filter on λ , and S is \mathcal{D} -stationary, then $\diamond_{\mathcal{D}}(S)$ means: There is a sequence $\langle A_\alpha : \alpha \in S \rangle$ such that each $A_\alpha \subseteq \alpha$ and for every $A \in [\lambda]^\lambda$,

$$\{\alpha \in S : A \cap \alpha = A_\alpha\} \in \mathcal{D}^+.$$

The statement $\diamond_{\mathcal{D}}^*(S)$ means: There is a sequence $\langle \mathcal{P}_\alpha : \alpha \in S \rangle$ such that each $\mathcal{P}_\alpha \subseteq \mathcal{P}(\alpha)$ and $|\mathcal{P}_\alpha| \leq \alpha$, and for every $A \subseteq \lambda$, there is a $C \in \mathcal{D}$ such that for all α ,

$$\alpha \in S \cap C \Rightarrow A \cap \alpha \in \mathcal{P}_\alpha.$$

By a well-known result of Kunen (see [6]),

$$\diamond_{\mathcal{D}}^*(S) \Rightarrow \diamond_{\mathcal{D}}(S).$$

It is easily seen that $\diamond_{\mathcal{D}}(S)$ implies the existence of an almost disjoint family of \mathcal{D} -stationary subsets of λ , of size 2^λ . Therefore, if $\diamond_{\mathcal{D}}(S)$ holds, \mathcal{D} is not 2^λ -saturated.

Looking back at the proof of Theorem 1.13, there are two important facts that we were using. The first is that there is a $\theta \in (\kappa, \lambda)$ such that θ has the weak reflection property for κ . The other important ingredient of the proof is Fact 1.13c. Dealing with filters other than \mathcal{D}_λ , to obtain the corresponding version of 1.13c, we have to strengthen our assumptions. Here, $\mathcal{I}[\theta, \kappa]$ is as in Definition 1.3.

Theorem 1.15. *Assume $\mu > \theta = \text{cf}(\theta) > \kappa = \text{cf}(\mu) > \aleph_0$, while $\lambda = \mu^+$ and $S_\kappa^\theta \notin \mathcal{I}[\theta, \kappa]$. Suppose that \mathcal{D} is a normal filter on λ such that $S_\kappa^\lambda \in \mathcal{D}$. If, for some S ,*

(*) *For every $A \in \mathcal{D}$,*

$$\left\{ \delta \in S : \forall C \text{ a club in } \delta \left(\{ \text{otp}(C \cap \gamma) : \gamma \in C \cap A \} \neq \emptyset \text{ mod } \mathcal{I}[\theta, \kappa] \right) \right\}$$

is a \mathcal{D} -stationary set in λ , then:

(1) *Claim 1.13c holds for any \tilde{C} and a_i^δ as in the assumptions of Claim 1.13c, with “stationary $\subseteq \lambda$ ” replaced by “ \mathcal{D} -stationary $\subseteq \lambda$ ”.*

(2) *$\mathcal{D} \upharpoonright S_\kappa^\lambda$ is not λ^+ -saturated.*

(3) *If $2^\mu = \lambda$ and $\mu^{[\kappa]} = \mu$, then $\diamond_{\mathcal{D}}^*(S_\kappa^\lambda)$ holds.*

We remind the reader of the notation $\mu^{[\theta]}$ for the revised cardinal power, from [20].

Definition 1.16. Suppose that $\nu > \chi$ are infinite cardinals and χ is regular. Then

$$\nu^{[\chi]} = \text{Min}\{|\mathcal{P}| : \mathcal{P} \subseteq [v]^\chi \text{ and } \forall A \in [v]^\chi (A \subseteq \text{the union of } < \chi \text{ elements of } \mathcal{P})\}.$$

Proof of Theorem 1.15. (1) and (2) are easily adjusted from Theorem 1.13.

(3) The conclusion is the same as that of [11], Section 3, and the proof is the same. The assumptions on the cardinal arithmetic are here the same as in [11, Section 3], with χ from there equal to our κ . The only difference is that the proof in [11, Section 3], started from $\square(\lambda)$, but we can use Fact 1.13a instead. \square

2. The $\clubsuit_{-\bar{\theta}}^*(S)$ principle

Suppose that S is a stationary subset of a regular uncountable cardinal λ . The aim of this section is to formulate a combinatorial principle which suffices to show that certain normal filters on λ cannot be λ^+ -saturated. The principle, $\clubsuit_{-\bar{\theta}}^*(S)$, is a form of $\clubsuit(S)$, where $\bar{\theta}$ is a sequence of ordinals. Our interest in this comes from two facts presented in 2.8. Firstly, if λ is a successor of a singular cardinal μ of uncountable cofinality κ , then $\clubsuit_{-\mu}^*(\lambda \setminus S_\kappa^\lambda)$ is always true. This can be used to obtain an alternative proof of that part of the result from [13, Th. 14] = [14, Th. 6] which states that no normal filter concentrating on $\lambda \setminus S_\kappa^\lambda$ is λ^+ -saturated. Secondly, if $\mathcal{I}[\lambda, \kappa] \upharpoonright S_\kappa^\lambda$ contains only non-stationary sets, $\clubsuit_{-\mu}^*(S_\kappa^\lambda)$ is true, so by 2.5. we can conclude that $\mathcal{D} \upharpoonright S_\kappa^\lambda$ is not λ^+ -saturated. The key to the proof of 2.8. is the combinatorial Lemma 2.7.

We commence by recalling the definition of some versions of \clubsuit .

Definition 2.0. Let λ be an uncountable regular cardinal and $S \subseteq \lambda$ stationary. Then:

(0) $\clubsuit(S)$ means: There is a sequence $\bar{A} = \langle A_\alpha : \alpha \in S \cap LIM \rangle$ such that:

- (i) For each $\alpha \in S \cap LIM$, the set A_α is an unbounded subset of α .
- (ii) For all $A \in [\lambda]^\lambda$, the set

$$G_{\clubsuit(S)}^{\bar{A}}[A] \stackrel{\text{def}}{=} \{\alpha \in S \cap LIM : A_\alpha \subseteq A\}$$

is non-empty.

(1) $\clubsuit^*(S)$ means: There is a sequence $\bar{\mathcal{P}} = \langle \mathcal{P}_\alpha : \alpha \in S \cap LIM \rangle$ such that

- (i) For each $\alpha \in S \cap LIM$, we have $|\mathcal{P}_\alpha| \leq |\alpha|$.
- (ii) If $B \in \mathcal{P}_\alpha$, then B is an unbounded subset of α .
- (iii) For every $A \in [\lambda]^\lambda$, there is a club C_A of λ such that

$$G_{\clubsuit^*(S)}^{\bar{\mathcal{P}}}[A] \stackrel{\text{def}}{=} \{\alpha \in S \cap LIM : \exists B \in \mathcal{P}_\alpha (B \subseteq A)\} \supseteq C_A \cap S \cap LIM.$$

Many authors use a different definition of \clubsuit , in which every unbounded set is required to be “guessed” stationarily many times. The following well-known fact shows that the two definitions are equivalent. We also include some other easy observations about Definition 2.0.

Fact 2.1. Assume that λ and S are as in Definition 2.0.

(0) If $\bar{A} = \langle A_\alpha : \alpha \in S \cap LIM \rangle$ is a $\clubsuit(S)$ -sequence, then, for every $A \in [\lambda]^\lambda$, the set $G_{\clubsuit(S)}^{\bar{A}}[A]$ is stationary.

(1) If $\clubsuit(S)$ holds, then there is a $\clubsuit(S)$ -sequence $\langle A_\alpha : \alpha \in S \cap LIM \rangle$ such that for each $\alpha \in S \cap LIM$, we have that $\text{otp}(A_\alpha) = \text{cf}(\alpha)$.

(3) Suppose that \mathcal{D} is a normal filter on λ and $S \in \mathcal{D}^+$ is such that $\bar{\mathcal{P}}$ exemplifies that $\clubsuit^*(S)$ holds. Then, for every $A \in [\lambda]^\lambda$, the set $G_{\clubsuit^*(S)}^{\bar{\mathcal{P}}}[A]$ is \mathcal{D} -stationary.

Proof. (0) Otherwise, we could find an $A \in [\lambda]^\lambda$ and a club C in λ such that for all α in $C \cap S \cap LIM$, the set A_α is not a subset of A . Then set $A^\dagger = \{\text{Min}(A \setminus \alpha) : \alpha \in C\}$, so $A^\dagger \in [\lambda]^\lambda$. But if $A_\alpha \subseteq A^\dagger$, then A_α is also a subset of A . On the other hand, since A^\dagger is unbounded in α (as A_α is), also C is unbounded in α , therefore $\alpha \in C$. This is a contradiction.

(1) Suppose that $\langle B_\alpha : \alpha \in S \cap LIM \rangle$ exemplifies $\clubsuit(S)$ and define for each $\alpha \in S \cap LIM$, the set A_α to be any cofinal subset of B_α with $\text{otp}(A_\alpha) = \text{cf}(\alpha)$.

(3) We simply remind the reader of the following elementary

Observation: For every club C of λ , we have $C \in \mathcal{D}$ (so the set $S \cap C$ is \mathcal{D} -stationary).

[Why? Suppose that C is a club of λ such that $C \notin \mathcal{D}$, so $S \setminus C \in \mathcal{D}^+$. We define the following function, for $\alpha \in S \setminus C$:

$$f(\alpha) \stackrel{\text{def}}{=} \sup(C \cap \alpha)$$

and we note that f is regressive on $S \setminus C$. Then we can find a $T \subseteq S \setminus C$ which is \mathcal{D} -stationary and such that $f \upharpoonright T$ is a constant. Then T must be bounded, which is a contradiction.] \square

Now we introduce the version of the \clubsuit^* principle that will mainly interest us. The guessing requirement is weaker, while the order type of the sets entering each family in the \clubsuit^* -sequence is controlled by a sequence of ordinals.

Definition 2.2. Let λ be a regular uncountable cardinal and $S \subseteq \lambda$ stationary. Let $\bar{\theta} = \langle \theta_\alpha : \alpha \in S \rangle$ be a sequence of ordinals. Then

$\clubsuit_{-\bar{\theta}}^*(S)$ means: There is a sequence $\bar{\mathcal{P}} = \langle \mathcal{P}_\alpha : \alpha \in S \cap LIM \rangle$ such that:

- (i) For each $\alpha \in S \cap LIM$, the family \mathcal{P}_α consists of $\leq |\alpha|$ unbounded subsets of α .
- (ii) For each $\alpha \in S \cap LIM$ and $B \in \mathcal{P}_\alpha$, we have $\text{otp}(B) < \theta_\alpha$.
- (iii) For $A \in [\lambda]^\lambda$, there is a club C_A of λ such that

$$G_{\clubsuit_{-\bar{\theta}}^*(S)}^{\bar{\mathcal{P}}}[A] \stackrel{\text{def}}{=} \{ \alpha : (\exists B \in \mathcal{P}_\alpha) (\alpha = \sup(B \cap A)) \} \supseteq C_A \cap S \cap LIM.$$

If for each $\alpha \in S$, $\theta_\alpha = \text{cf}(\alpha) + 1$, we omit $\bar{\theta}$ in the above notation.

If for some μ we have that $\theta_\alpha = \mu$ for all $\alpha \in S \cap LIM$, then we write $\clubsuit_{-\mu}^*(S)$ rather than $\clubsuit_{-\bar{\theta}}^*(S)$.

We make some easy remarks on Definition 2.2.

Observation 2.3. Assume that λ , S and $\bar{\theta}$ are as in Definition 2.2.

(0) If the set of all $\alpha \in S \cap LIM$ for which $\theta_\alpha > \text{cf}(\alpha)$ is non-stationary, then $\clubsuit_{-\bar{\theta}}^*(S)$ is false. Otherwise $\clubsuit^*(S) \Rightarrow \clubsuit_{-\bar{\theta}}^*(S)$.

(1) If $\sum_{\gamma < \theta_\alpha} |\gamma|^{\text{cf}(\alpha)} \leq |\alpha|$, for each $\alpha \in S$, then

$$\clubsuit_{-\bar{\theta}}^*(S) \Rightarrow \clubsuit^*(S).$$

(2) Suppose that \mathcal{D} is a normal filter on λ , while S is \mathcal{D} -stationary and $\bar{\mathcal{P}}$ exemplifies $\clubsuit_{-\bar{\theta}}^*(S)$. Then, for every $A \in [\lambda]^\lambda$ the set $G_{\clubsuit_{-\bar{\theta}}^*(S)}^{\bar{\mathcal{P}}}[A]$ is \mathcal{D} -stationary.

Proof. (0) Obvious.

(1) If $\bar{\mathcal{P}} = \langle \mathcal{P}_\alpha : \alpha \in S \cap LIM \rangle$ exemplifies $\clubsuit_{-\bar{\theta}}^*(S)$, define

$$\mathcal{P}_\alpha^\dagger = \begin{cases} \left\{ \left\{ B : \exists A \in \mathcal{P}_\alpha (B \subseteq A \wedge B \text{ cofinal in } \alpha \wedge \text{otp}(B) = \text{cf}(\alpha)) \right\} \right\} & \text{if } \theta_\alpha > \text{cf}(\alpha), \\ \{ \alpha \} & \text{otherwise.} \end{cases}$$

Then $|\mathcal{P}_\alpha^\dagger| = |\mathcal{P}_\alpha|^{\text{cf}(\alpha)} \leq \alpha$ and, by (0), $\langle \mathcal{P}_\alpha^\dagger : \alpha \in S \cap LIM \rangle$ exemplifies $\clubsuit^*(S)$.

(2) Like 2.1.3. \square

We can consider also \clubsuit -sequences whose failure to guess is always confined to a set in a given ideal on λ . In this context, for example $\clubsuit(S)$ will mean $\clubsuit(S)/J_\lambda$.

Definition 2.4. Let λ and S be as above, while \mathcal{I} is an ideal on λ . Then

$\clubsuit(S)/\mathcal{I}$ means: There is a sequence $\bar{A} = \langle A_\alpha : \alpha \in S \rangle$ such that each A_α is a cofinal subset of α , and this sequence has the following property. For every $A \in [\lambda]^\lambda$, the set

$$G_{\clubsuit(S)/\mathcal{I}}^{\bar{A}}[A] \stackrel{\text{def}}{=} \{\alpha : A_\alpha \subseteq A\}$$

satisfies $G_{\clubsuit(S)/\mathcal{I}}^{\bar{A}}[A] \notin \mathcal{I}$.

If \mathcal{D} is the dual filter of the ideal \mathcal{I} , then $\clubsuit(S)/\mathcal{D}$ means the same as $\clubsuit(S)/\mathcal{I}$. We extend this definition in the obvious way to the other mentioned versions of the \clubsuit principle.

Theorem 2.5. *Suppose that $\lambda = \mu^+$ and μ is a limit cardinal.*

(1) *Assume that $\clubsuit_{\mu}^*(S)$ holds for a stationary $S \subseteq \lambda$. Then no normal filter \mathcal{D} on S is λ^+ -saturated.*

(2) *If $S \subseteq \lambda$ is stationary and \mathcal{D} is a normal filter on S such that $\clubsuit_{\mu}^*(S)/\mathcal{D}$ holds, then \mathcal{D} is not λ^+ -saturated.*

Proof. Let us fix a sequence $\bar{\mathcal{P}} = \langle \mathcal{P}_\delta : \delta \in S \cap LIM \rangle$ which exemplifies $\clubsuit_{\mu}^*(S)$. Therefore, for each $\delta \in S \cap LIM$, we have a family $\mathcal{P}_\delta = \{P_{\delta,i} : i < i^\delta \leq \delta\}$ such that each $P_{\delta,i}$ is a cofinal subset of δ of order type $\text{otp}(P_{\delta,i}) < \mu$.

We fix a 1–1 onto pairing function $\text{pr} : \lambda \times \lambda \rightarrow \lambda \setminus \omega$ such that for each $\alpha, \beta \in \lambda$, we have

$$\text{Max}\{\alpha, \beta\} \leq \text{pr}(\alpha, \beta) < (|\alpha| + |\beta|)^+.$$

(Here, we use the convention that $n^+ = \aleph_1$ for $n \in \omega$.)

We shall also fix a club C of $\lambda \setminus \mu$ such that

$$\alpha, \beta < \gamma \ \& \ \gamma \in C \Rightarrow \text{pr}(\alpha, \beta) < \gamma.$$

Now, we choose sets b_ζ^ζ for $\zeta < \lambda^+$ and $\alpha < \lambda$, and the functions $h_\zeta : \lambda \rightarrow \lambda$ for $\zeta < \lambda^+$ as in the proof of 1.13, and with the same properties as there. For $\zeta < \lambda^+$, we define the unbounded subset X_ζ of λ by

$$X_\zeta \stackrel{\text{def}}{=} \{\text{pr}(\alpha, h_\zeta(\alpha)) : \alpha < \lambda\},$$

and partial functions g_ζ by

$$g_\zeta(\delta) \stackrel{\text{def}}{=} \text{Min}\{i < \delta : \sup(X_\zeta \cap P_{\delta,i}) = \delta\}.$$

In fact, the domain of each g_ζ is exactly the set $G_{\clubsuit_{\mu}^*(S)}^{\bar{\mathcal{P}}}[X_\zeta]$, so $\text{Dom}(g_\zeta)$ is \mathcal{D} -stationary (by 2.3.2). For $\delta \in \text{Dom}(g_\zeta) \cap C$, we can define

$$f_\zeta(\delta) \stackrel{\text{def}}{=} \text{pr}(g_\zeta(\delta), |P_{\delta, g_\zeta(\delta)}|^+).$$

Notice that g_ζ is regressive on its domain, so if $\delta \geq \mu$ and $f_\zeta(\delta)$ is defined, then $f_\zeta(\delta) < \delta$ (as $\delta \in C$). Therefore, there is a \mathcal{D} -stationary set B_ζ such that $f_\zeta \upharpoonright B_\zeta$ is constantly equal to $\text{pr}(i_\zeta, \theta_\zeta)$ for some i_ζ and a regular cardinal $\theta_\zeta < \mu$. Then there are $i^* < \lambda$ and a regular cardinal $\theta^* < \mu$ such that the set

$$W \stackrel{\text{def}}{=} \{\zeta < \lambda^+ : (i_\zeta, \theta_\zeta) = (i^*, \theta^*)\}$$

is unbounded.

Let us define for $S \cap \text{acc}(C) \cap LIM$ with $i^* < i^\delta$ the set

$$d_\delta \stackrel{\text{def}}{=} \{\alpha : (\exists \beta)(\text{pr}(\alpha, \beta) \in P_{\delta, i^*})\} \cup \{\beta : (\exists \alpha)(\text{pr}(\alpha, \beta) \in P_{\delta, i^*})\}.$$

By the definition of pr , we have $d_\delta \subseteq \delta$. In fact, since $\delta \in \text{acc}(C)$, the set d_δ is an unbounded subset of δ . Like in 1.13, we can define for $\zeta \in W$ and δ for which d_δ is defined, a function $h_{\zeta, \delta}$ on d_δ given by

$$h_{\zeta, \delta}(\beta) = \begin{cases} \text{Min}(d_\delta \setminus (h_\zeta(\beta) + 1)) & \text{if } d_\delta \setminus (h_\zeta(\beta) + 1) \neq \emptyset, \\ \lambda & \text{otherwise.} \end{cases}$$

Therefore

$$h_{\zeta, \delta} : d_\delta \rightarrow d_\delta \cup \{\lambda\}.$$

For $\zeta, \xi \in W$ we define sets

$$A_{\zeta, \xi} = \{\delta \in S : d_\delta \text{ is defined and for unboundedly } \beta \in d_\delta$$

$$\text{we have } h_{\zeta, \delta}(\beta) < h_{\xi, \delta}(\beta)\}.$$

Assume now that \mathcal{D} is λ^+ -saturated, and let us make some simple observations about the just defined sets:

(a) $A_{\zeta, \xi} / \mathcal{D}$ increase with ξ and decrease with ζ .

(b) Since \mathcal{D} is λ^+ -saturated, for any fixed $\zeta \in W$, the sequence $\langle A_{\zeta, \xi} / \mathcal{D} : \xi < \lambda^+ \rangle$ is eventually constant, let us say for $\xi \in [\xi_\zeta, \lambda^+) \cap W$.

Similarly,

(c) $\langle A_\zeta \stackrel{\text{def}}{=} A_{\zeta, \xi_\zeta} / \mathcal{D} : \zeta < \lambda^+ \rangle$ are eventually constant, say for $\zeta \in W \setminus \zeta^*$.

We choose by induction on $\varepsilon < \theta^*$ ordinals ζ_ε such that:

- (i) $\zeta_\varepsilon \in (\zeta^*, \lambda^+)$.
- (ii) $\zeta_\varepsilon \in W$.
- (iii) ζ_ε are strictly increasing with ε .
- (iv) $\zeta_{\varepsilon+1} > \xi_{\zeta_\varepsilon}$.

Therefore

$$\varepsilon(1) < \varepsilon(2) < \theta^* \Rightarrow A_{\zeta_{\varepsilon(1)}, \zeta_{\varepsilon(2)}} / \mathcal{D} = A_{\zeta_{\varepsilon(1)}, \xi_{\zeta_{\varepsilon(1)}}} / \mathcal{D} = A_{\zeta^*} / \mathcal{D}.$$

Let $\gamma^* < \lambda$ be such that for all $\alpha \in (\gamma^*, \lambda^+)$, the sequence $\langle h_{\zeta_\varepsilon}(\alpha) : \varepsilon < \theta^* \rangle$ is strictly increasing. (Recall that for $\zeta < \xi < \lambda^+$, we know that h_ζ is eventually strictly less than h_ξ .) We can as well assume that $\gamma^* > \omega$.

By the above and the fact that $\theta^* < \lambda$, we can find a set $E \in \mathcal{D}$ such that

$$\varepsilon(1) < \varepsilon(2) < \theta^* \Rightarrow A_{\zeta_{\varepsilon(1)}, \zeta_{\varepsilon(2)}} \cap E = A_{\zeta_{\varepsilon(1)}, \xi_{\varepsilon(1)}} \cap E = A_{\zeta(*)} \cap E.$$

We can assume that $\text{Min}(E) > \gamma(*)$. Without loss of generality, we can also add that $E \subseteq \text{acc}(C)$ and

$$\varepsilon < \theta^* \ \& \ \delta \in E \cap S \cap \text{LIM} \Rightarrow h_{\zeta_\varepsilon}'' \delta \subseteq \delta,$$

since $\theta^* < \lambda$.

Now we discuss two possible cases, the first of which corresponds to the situation in 1.13.

Case 1. For some $\bar{\zeta} \geq \zeta(*)$, we have $\bar{\zeta} \in W$ and $A_{\zeta(*)} \cap B_{\bar{\zeta}}$ is \mathcal{D} -stationary. We choose a $\delta \in E \cap B_{\bar{\zeta}} \cap A_{\zeta(*)}$. In particular, d_δ is defined. We consider $\langle h_{\zeta_\varepsilon, \delta} : \varepsilon < \theta^* \rangle$. By the choice of E , each $h_{\zeta_\varepsilon, \delta}$ is a function from d_δ to itself.

For any $\alpha \in d_\delta \setminus \gamma(*)$, the sequence $\langle h_{\zeta_\varepsilon, \delta}(\alpha) : \varepsilon < \theta^* \rangle$ is non-decreasing. As $\delta \in B_{\bar{\zeta}}$ and $\bar{\zeta} \in W$, we know that $|d_\delta| \leq |P_{\delta, i^*}| < \theta^*$, so the above sequence is eventually constant.

Similarly, $|d_\delta \setminus \gamma(*)| < \theta^*$, so there is some $\varepsilon_0 < \theta^*$ and some function h such that for all $\varepsilon \in (\varepsilon_0, \theta^*)$, we have $h_{\zeta_\varepsilon, \delta} \upharpoonright (d_\delta \setminus \gamma(*) = h$. But $\delta \in A_{\zeta(*)} \cap E$, so $\delta \in A_{\zeta_\varepsilon, \zeta_{\varepsilon+1}}$ for all $\varepsilon < \theta^*$, and therefore for any such ε ,

$$\{\alpha \in d_\delta : h_{\zeta_\varepsilon, \delta}(\alpha) < h_{\zeta_{\varepsilon+1}, \delta}(\alpha)\}$$

is unbounded in δ . This is a contradiction.

Case 2. $A_{\zeta(*)} \cap B_{\bar{\zeta}}$ is not \mathcal{D} -stationary for any $\bar{\zeta} \in W$ with $\bar{\zeta} \geq \zeta(*)$. Then for all $\zeta \in W$ and $\bar{\zeta} \in W \setminus \zeta(*)$, the set $A_{\zeta, \bar{\zeta}} \cap B_{\bar{\zeta}}$ is not \mathcal{D} -stationary, as $A_{\zeta, \bar{\zeta}} \subseteq A_{\zeta(*)} / \mathcal{D}$. Similarly, since for $\zeta < \xi \in W$, we have that $A_{\zeta, \xi} \subseteq A_{\zeta, \bar{\zeta}} / \mathcal{D}$, we conclude that $A_{\zeta, \xi} \cap B_{\bar{\zeta}}$ is not \mathcal{D} -stationary for any $\zeta < \xi \in W$ and $\bar{\zeta} \in W \setminus \zeta(*)$.

On the other hand, as each B_ζ for $\zeta \in W$ is \mathcal{D} -stationary, and we are assuming that \mathcal{D} is λ^+ -saturated, there are $\zeta < \xi \in W \setminus \zeta(*)$ such that $B_\zeta \cap B_\xi$ is \mathcal{D} -stationary. We fix such ζ and ξ .

Let us choose a $\delta \in (E \cap B_\zeta \cap B_\xi) \setminus A_{\zeta, \xi}$. Without loss of generality, we can assume that there is an $\alpha_{\zeta, \xi} < \delta$ such that $\alpha \geq \alpha_{\zeta, \xi} \Rightarrow h_\zeta(\alpha) < h_\xi(\alpha)$. Note that d_δ is defined. Then, by the definition of $A_{\zeta, \xi}$, we can find a $\gamma_1 \in (\alpha_{\zeta, \xi}, \delta)$ such that

$$h_{\zeta, \delta} \upharpoonright (d_\delta \setminus \gamma_1) = h_{\xi, \delta} \upharpoonright (d_\delta \setminus \gamma_1).$$

Note now that there is a club E_ξ such that

$$\alpha \in E_\xi \Rightarrow h_\xi'' \alpha \subseteq \alpha,$$

and a similarly defined club E_ζ . We can without loss of generality assume that $\delta \in E_\zeta \cap E_\xi$, so both h_ζ and h_ξ are functions from d_δ to itself. Now, since $\delta \in B_\zeta \cap B_\xi$, we know that $g_\zeta(\delta) = g_\xi(\delta) = i(*)$. By the definition of X_ξ , we can find an $\alpha \in (\gamma_1, \delta)$ such that $\text{pr}(\alpha_1, h_\xi(\alpha_1)) \in d_\delta$, so by the definition of $h_{\xi, \delta}$ we have

$$h_{\xi, \delta}(\alpha_1) > h_\xi(\alpha_1).$$

On the other hand, $h_{\zeta}(\alpha_1) < h_{\zeta}(\alpha_1) \in d_{\delta}$, so $h_{\zeta,\delta}(\alpha_1) \leq h_{\zeta}(\alpha_1)$. Therefore, $h_{\zeta,\delta}(\alpha_1) \neq h_{\zeta,\delta}(\alpha_1)$, which is a contradiction to $\alpha_1 > \gamma_1$.

(2) Follows from the proof of (1). \square

Like \diamond and by the same proof, the usual \clubsuit principle on λ can be used for guessing not just unbounded subsets of λ , but any other structure which can be coded by the unbounded subsets of λ . With the \clubsuit^* principle, this does not seem to be the case, or at least the \diamond -like proof fails. In particular, we do not know if the following is true:

Question 2.6. Suppose that λ is a regular uncountable cardinal and S a stationary subset of λ such that $\clubsuit^*(S)$ holds. Is it true that there is a sequence $\langle \mathcal{F}_{\alpha} : \alpha \in S \cap LIM \rangle$ with the following properties:

(i) Each \mathcal{F}_{α} consists of partial functions from α to α , each of which has an unbounded subset of α as its domain.

(ii) $|\mathcal{F}_{\alpha}| \leq \alpha$.

(iii) For every function $f : \lambda \rightarrow \lambda$, there is a club C_f of λ with the property

$$\alpha \in C_f \cap S \cap LIM \Rightarrow (\exists g \in \mathcal{F}_{\alpha})(\sup\{\beta \in \text{Dom}(g) : g(\beta) = f(\beta)\} = \alpha)?$$

We note that a positive answer to 2.6 would quite simplify the proof of 2.5.

Our next goal is to prove that $\clubsuit^*_{-\mu}(\lambda \setminus S_{\kappa}^{\lambda})$ is true for any λ which is the successor of a singular cardinal μ of uncountable cofinality κ . The key is the following

Lemma 2.7. Assume that $\lambda = \mu^+$ and $\aleph_0 < \kappa = \text{cf}(\mu) < \mu$. Also $\mu = \sum_{i < \kappa} \mu_i$, where $\langle \mu_i : i < \kappa \rangle$ is an increasing continuous sequence of cardinals and, for simplicity, $\mu_0 > \kappa$. Then there is a sequence

$$\langle \langle a_i^{\alpha} : i < \kappa, \alpha < \lambda \rangle \rangle$$

such that for every $\alpha < \lambda$, the sets $a_i^{\alpha} (i < \kappa)$ are subsets of α which are \subseteq -increasing in i , with $|a_i^{\alpha}| \leq \mu_i$, and such that:

For any $f : \lambda \rightarrow \lambda$, if

$$A_i^f = \left\{ \alpha < \lambda : \alpha = \sup\{\zeta \in a_i^{\alpha} : f(\zeta) \in a_i^{\alpha}\} \right\},$$

then:

(A) A_i^f are \subseteq -increasing in i .

(B) $\lambda \setminus S_{\kappa}^{\lambda} \subseteq \bigcup_{i < \kappa} A_i^f \pmod{\mathcal{D}_{\lambda}}$.

(C) If $\gamma \in S_{\kappa}^{\lambda}$, while $i < \kappa$ and A_i^f reflects on γ , then $\gamma \in A_i^f$. (In fact, this is true for any $\gamma < \lambda$ with $\kappa \geq \text{cf}(\gamma) > \aleph_0$.) So:

(C') If $S_f \stackrel{\text{def}}{=} S_{\kappa}^{\lambda} \setminus \bigcup_{i < \kappa} A_i^f$, then for no $i < \kappa$ does A_i^f reflect in any $\delta \in S_f$.

(D) There is a non-stationary set N such that $S_f \setminus N \in \mathcal{I}[\lambda, \kappa]$.

Proof. We describe the choice of sets a_i^{α} , and then we check that all claims of the lemma are satisfied.

First we fix a sequence $\langle e_\gamma : \gamma \in LIM \cap \lambda \rangle$ such that e_γ is a club of γ with $\text{otp}(e_\gamma) = \text{cf}(\gamma)$. Then we define by induction on $\alpha < \lambda$ sets a_i^α for all $i < \kappa$, requiring that for all limit γ

$$\mu_i \geq \text{cf}(\gamma) \Rightarrow \bigcup_{\beta \in e_\gamma} a_i^\beta \subseteq a_i^\gamma,$$

in addition to the requirements that we already mentioned in the statement of the lemma.

Now we suppose that $f : \lambda \rightarrow \lambda$ is given, and check claims (A)–(D).

(A) This follows from the fact that for every $\alpha < \lambda$,

$$i < j < \kappa \Rightarrow a_i^\alpha \subseteq a_j^\alpha.$$

(B) Let

$$E \stackrel{\text{def}}{=} \left\{ \alpha \in \lambda : \alpha = \sup\{\zeta < \alpha : f(\zeta) < \alpha\} \right\}.$$

Then E is a club of λ . We shall show that

$$(\lambda \setminus S_\kappa^\lambda) \cap \text{acc}(E) \subseteq \bigcup_{i < \kappa} A_i^f.$$

So, let us take a $\gamma \in \text{acc}(E)$ such that $\text{cf}(\gamma) \neq \kappa$. Then $e_\gamma \cap E$ is a club of γ , and $\text{otp}(e_\gamma \cap E) = \text{cf}(\gamma)$. Let $e_\gamma \cap E = \{\beta_\varepsilon : \varepsilon < \text{cf}(\gamma)\}$ be an increasing enumeration. So, for all ε , we have

$$f(\beta_\varepsilon) < \beta_{\varepsilon+1} < \gamma.$$

Since $\text{cf}(\gamma) \neq \kappa$, there must be an $i < \kappa$ such that for some unbounded $c \subseteq \text{cf}(\gamma)$, we have $\{\beta_\varepsilon, f(\beta_\varepsilon) : \varepsilon \in c\} \subseteq a_i^\gamma$ (note that we are using the fact that a_i^α are increasing with i). Then $\gamma \in A_i^f$.

(C) Suppose that $\kappa \geq \text{cf}(\gamma) > \aleph_0$ and A_i^f reflects on γ . In particular, $A_i^f \cap e_\gamma$ is stationary in γ . Given an $\alpha < \gamma$, we can find a $\zeta \in A_i^f \cap e_\gamma$ such that $\alpha < \zeta$. Then, there is a $\xi \in a_i^\zeta$ such that $\alpha < \xi$ and $f(\xi) \in a_i^\zeta$, by the definition of A_i^f . Since $\text{cf}(\gamma) \leq \kappa \leq \mu_i$, we have that $a_i^\zeta \subseteq a_i^\gamma$, and we are done.

(C') This follows immediately by (C).

(D) Define $h : S_{<\kappa}^\lambda \rightarrow \kappa$ by $h(\alpha) \stackrel{\text{def}}{=} \text{Min}\{i : \alpha \in A_i^f\}$. Then, if $\delta \in S_f \cap \text{acc}(E)$, the function h is defined on $S_{<\kappa}^\lambda \cap \delta$ and not constant on any stationary set of δ , by (C'). By 1.11, h is increasing on some club of δ . \square

Theorem 2.8. (1) Suppose that $\lambda = \mu^+$, and $\mu > \text{cf}(\mu) = \kappa > \aleph_0$. Then, $\clubsuit_{-\mu}^*(\lambda \setminus S_\kappa^\lambda)$ holds and $\clubsuit_{-\mu}^*(S_\kappa^\lambda) / \mathcal{S}[\lambda, \kappa]$ holds.

(2) If in (1) we in addition assume that μ is a strong limit, then $\clubsuit^*(\lambda \setminus S_\kappa^\lambda)$ holds and $\clubsuit^*(S_\kappa^\lambda) / \mathcal{S}[\lambda, \kappa]$ holds.

Proof. Let us fix sets a_i^α ($i < \kappa$) for $\alpha < \lambda$ as guaranteed by Lemma 2.7. We fix for all $\alpha \in \lambda$ a cofinal subset P_α of α such that $\text{otp}(P_\alpha) = \text{cf}(\alpha)$.

(1) We shall define for $\alpha \in \lambda$,

$$\mathcal{P}_\alpha = \begin{cases} \{a_i^\alpha : i < \kappa \text{ \& sup}(a_i^\alpha) = \alpha\} \cup \{P_\alpha\} & \text{if } |\alpha| \geq \kappa, \\ \{\alpha\} & \text{otherwise.} \end{cases}$$

Now, certainly each \mathcal{P}_α is a family of $\leq |\alpha|$ subsets of α .

Suppose that $A \in [\lambda]^\lambda$ is given, and let f be an increasing enumeration of A . In particular, $f(\zeta) \geq \zeta$ for all $\zeta \in \lambda$. The set

$$C \stackrel{\text{def}}{=} \{\alpha < \lambda : \forall \zeta \in \alpha (f(\zeta) < \alpha)\} \setminus \kappa$$

is a club of λ . Let us take any $\alpha \in C \cap S_{\neq \kappa}^\lambda$. By (B) of Lemma 2.7, there is an $i < \kappa$ such that $\alpha \in A_i^f$, where A_i^f is as defined in Lemma 2.7. Then $\alpha = \sup\{\zeta \in a_i^\alpha : f(\zeta) \in a_i^\alpha\}$, so $\alpha = \sup(A \cap a_i^\alpha)$.

This proves $\clubsuit_{-\mu}^*(\lambda \setminus S_\kappa^\lambda)$. With the same definition of $\bar{\mathcal{P}} \stackrel{\text{def}}{=} \langle \mathcal{P}_\alpha : \alpha < \lambda \rangle$, let us start again from an $A \in [\lambda]^\lambda$, and f and C as above. Let S_f and N be as in Lemma 2.7, so $S_f \setminus N \in \mathcal{I}[\lambda, \kappa]$. If $\alpha \in C \cap S_\kappa^\lambda \setminus S_f$, then we argue as above, to conclude that $\alpha \in G_{\clubsuit_{-\mu}^*(S_\kappa^\lambda)/\mathcal{I}[\lambda, \kappa]}^{\bar{\mathcal{P}}}[A]$.

(2) With the same notation as above, we define

$$\mathcal{P}_\alpha \stackrel{\text{def}}{=} \begin{cases} \{\text{all cofinal sequences of } \alpha \text{ included in } a_i^\alpha : i < \kappa\} \cup \{P_\alpha\} & \text{if } |\alpha| \geq \kappa, \\ \{\alpha\} & \text{otherwise.} \end{cases}$$

Then $|\mathcal{P}_\alpha| \leq 2^{|\alpha|} + \kappa < \mu$.

We argue similarly on S_κ^λ , using the set S_f as above. \square

As a consequence, we obtain another proof of (a part of) a theorem from [13, Th. 14] and [14, Th. 6], as well as some other statements.

Corollary 2.9. *Suppose that $\lambda = \mu^+$ and $\mu > \text{cf}(\mu) = \kappa > \aleph_0$. Then:*

- (1) *No normal filter on $\lambda \setminus S_\kappa^\lambda$ is λ^+ -saturated.*
- (2) *If \mathcal{D} is a normal filter on S_κ^λ such that $\mathcal{I}[\lambda, \kappa] \upharpoonright S_\kappa^\lambda$ contains only non-stationary sets, then \mathcal{D} is not λ^+ -saturated.*
- (3) *If $\clubsuit(S)$ holds, then $\mathcal{D}_\lambda \upharpoonright S$ is not λ^+ -saturated.*

Proof. (1) $\clubsuit_{-\mu}^*(\lambda \setminus S_\kappa^\lambda)$ holds, by 2.8. By 2.5, \mathcal{D} cannot be λ^+ -saturated.

(2) Similar.

(3) See 2.10(2) below. \square

Concluding Remarks 2.10. (0) Another useful version of the \clubsuit -principle on λ for a stationary $S \subseteq \lambda$ is $\clubsuit^-(S)$, which says that: There is a sequence $\bar{\mathcal{P}} = \langle \mathcal{P}_\alpha : \alpha \in S \cap LIM \rangle$ such that \mathcal{P}_α is a family of $\leq \alpha$ unbounded subsets of α with the property that for all unbounded subsets A of λ ,

$$G_{\clubsuit^-(S)}^{\bar{\mathcal{P}}}[A] \stackrel{\text{def}}{=} \{\alpha \in S : \exists B \in \mathcal{P}_\alpha (B \subseteq A \cap \alpha)\}$$

is non-empty.

As opposed to the situation with \diamond , it does not have to be true that $\clubsuit^-(S) \Rightarrow \clubsuit(S)$. Of course, still $\clubsuit^*(S) \Rightarrow \clubsuit^-(S)$, so we do not consider \clubsuit^- here.

(1) We can also consider versions of \clubsuit^* or \clubsuit^- for which the size of each \mathcal{P}_α is determined by some cardinal μ_α , not necessarily equal to $|\alpha|$. Also, we can combine this idea with the idea of $\clubsuit_{-\bar{\theta}}$, so also the order type of sets in \mathcal{P}_α is controlled by some prescribed sequence $\bar{\theta}$.

(2) If we now define $\clubsuit_{-\mu}^-(S)$ in the obvious way, then it follows from the proof of 2.5 that for $\lambda = \mu^+$ and μ singular, $\clubsuit_{-\mu}^-(S)$ is enough to guarantee that $\mathcal{D}_\lambda \upharpoonright S$ is not λ^+ -saturated. Therefore, in particular, $\clubsuit(S)$ suffices.

For λ the successor of a strong limit, most “reasonable” versions of \clubsuit coincide.

(3) After hearing our lecture at the Logic Seminar in Jerusalem, Fall 1994, M. Magidor showed us an alternative proof of 2.5 using elementary embeddings and ultrapowers and not requiring μ to be a limit cardinal.

(4) The assumptions of 1.13 and 2.5 seem similar, but we point out that they are in fact different. The existence of a $\theta < \lambda$ which weakly reflects at κ is not the same as the assumption that $\mathcal{S}[\lambda, \kappa] \upharpoonright \mathcal{S}_\kappa^\lambda$ contains only bounded sets.

Acknowledgements

Authors partially supported by the Basic Research Foundation Grant Number 0327398 administered by the Israel Academy of Sciences and Humanities. For easier future reference, note that this is publication [DjSh 545] in Shelah’s bibliography. The research for this paper was done in the period between January and August 1994. The most sincere thanks to James Cummings for his numerous comments and corrections, and to Uri Abraham, Menachem Magidor and other participants of the Logic Seminar in Jerusalem, Fall 1994, for their interest. The first author also wishes to thank the Hebrew University and the Lady Davis Foundation for the Forchheimer Postdoctoral Fellowship.

References

- [1] J.E. Baumgartner, A new class of order types, *Ann. Math. Logic* 9 (1976) 187–222.
- [2] J. Cummings, M. Džamonja and S. Shelah, A consistency result on weak reflection, in *Fund. Math.* 148 (1995) 91–100.
- [3] M. Foreman, More saturated ideals, *Cabal Seminar* 79–81, *Pro. Caltech-UCLA Logic Seminar 1979–81*, A.S. Kekris, D.A. Martin and Y.N. Moschovakis, eds., *Lecture Notes in Mathematics*, Vol. 1019 (Springer, Berlin, 1981) 1–27.
- [4] M. Foreman, M. Magidor and S. Shelah, Martin’s maximum, saturated ideals and nonregular ultrafilters I, *Ann. of Math. (Second Series)* 27 (1988) 1–47.
- [5] K. Kunen, Saturated ideals, *J. Symbolic Logic* 43 (1978) 65–76.
- [6] K. Kunen, *Set Theory, An Introduction to Independence Proofs* (North-Holland, Amsterdam, 1980).
- [7] A.H. Mekler and S. Shelah, The consistency strength of “every stationary set reflects”, *Israel J. Math.* 67 (1989) 353–366.
- [8] A.J. Ostaszewski, On countably compact perfectly normal spaces, *J. London Math. Soc.* 14 (1975) 505–516.

- [9] S. Shelah, On successors of singular cardinals, in: M. Boffa, D. van Dalen and K. McAloon, eds., *Logic Colloquium '78* (North-Holland, Amsterdam, 1979) 357–380.
- [10] S. Shelah, Whitehead groups may not be free, even assuming CH , *Israel J. Math.* 35 (1980) 257–285.
- [11] S. Shelah, Diamonds, uniformization, *J. Symbolic Logic* 49 (1984) 1022–1033.
- [12] S. Shelah, Appendix: on stationary sets (to “Classification of nonelementary classes. II. Abstract elementary classes”), in: J.T. Baldwin, ed., *Classification Theory* (Chicago, IL, 1985), Proc. USA–Israel Conference on Classification Theory, Chicago, December 1985; *Lecture Notes in Mathematics*, Vol. 1292 (Springer, Berlin, 1987) 483–495.
- [13] S. Shelah, The existence of coding sets, in: *Around Classification Theory of Models*, *Lecture Notes in Mathematics*, Vol. 1182 (Springer, Berlin, 1986) 188–202.
- [14] S. Shelah, More on stationary coding, in: *Around Classification Theory of Models*, *Lecture Notes in Mathematics*, Vol. 1182 (Springer, Berlin, 1986) 224–246.
- [15] S. Shelah, Reflecting stationary sets and successors of singular cardinals, *Arch. Math. Logic* 31 (1991) 25–53.
- [16] S. Shelah, Advances in cardinal arithmetic, in: N. Sauer et al., eds., *Finite and Infinite Combinatorics in Sets and Logic* (Kluwer Academic, Dordrecht, 1993) 355–383.
- [17] S. Shelah, *Cardinal Arithmetic* (Oxford Univ. Press, Oxford, 1994).
- [18] S. Shelah, $\aleph_{\omega+1}$ has a Jonsson algebra, in: *Cardinal Arithmetic*, Ch. II (Oxford Univ. Press, Oxford, 1994).
- [19] S. Shelah, Jonsson algebras in inaccessible cardinals, in: *Cardinal Arithmetic*, Ch. III (Oxford Univ. Press, Oxford, 1994).
- [20] S. Shelah, The generalized continuum hypothesis revisited, *Israel J. Math.*, submitted.
- [21] J.R. Steel and R. van Wesep, Two consequences of determinacy consistent with choice, *Trans. AMS* 272(1) (1982) 67–85.