



On Invariants for ω_1 -Separable Groups

Author(s): Paul C. Eklof, Matthew Foreman and Saharon Shelah

Source: *Transactions of the American Mathematical Society*, Vol. 347, No. 11 (Nov., 1995), pp. 4385-4402

Published by: [American Mathematical Society](#)

Stable URL: <http://www.jstor.org/stable/2155042>

Accessed: 28/06/2014 10:35

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



American Mathematical Society is collaborating with JSTOR to digitize, preserve and extend access to *Transactions of the American Mathematical Society*.

<http://www.jstor.org>

ON INVARIANTS FOR ω_1 -SEPARABLE GROUPS

PAUL C. EKLOF, MATTHEW FOREMAN, AND SAHARON SHELAH

ABSTRACT. We study the classification of ω_1 -separable groups by using Ehrenfeucht-Fraïssé games and prove a strong classification result assuming PFA, and a strong non-structure theorem assuming \diamond .

INTRODUCTION

An ω_1 -separable (or \aleph_1 -separable) group is an abelian group such that every countable subset is contained in a free direct summand of the group. In particular, therefore, an ω_1 -separable group is \aleph_1 -free, i.e., every countable subgroup is free. The structure of ω_1 -separable groups of cardinality \aleph_1 was investigated in [1] and [8]; most of the results proved there required set-theoretic assumptions beyond ZFC. (See also [2, Chapter VIII] for an exposition of these results.) Specifically, assuming Martin's Axiom (MA) plus \neg CH or the stronger Proper Forcing Axiom (PFA), one can prove nice structure and classification results; these results are not theorems of ZFC since counterexamples exist assuming CH or "prediction principles" like \diamond . In [1, Remark 3.3] it is asserted that a construction given there under the assumption of CH (or even $2^{\aleph_0} < 2^{\aleph_1}$) of two non-isomorphic ω_1 -separable groups

"is strong evidence for the claim that in a model of CH there is no possible meaningful classification of all ω_1 -separable groups. It is difficult to see what conceivable scheme of classification could distinguish between [the groups constructed here]."

But, in fact, the Helsinki school of model theory provides a scheme for distinguishing between such groups. It is our aim here to use the methodology of the Helsinki school—which involves Ehrenfeucht-Fraïssé games (cf. [9], [11] or [12])—to strengthen the dichotomy referred to above: that is, to obtain strong classification results assuming PFA, and a strong "non-structure theorem" assuming \diamond .

Received by the editors October 13, 1994.

1991 *Mathematics Subject Classification.* Primary 03C55, 20K20; Secondary 03E35, 03C75.

Key words and phrases. Ehrenfeucht-Fraïssé games, \aleph_1 -separable abelian groups, almost free groups.

Thanks to Rutgers University for its support of this research through its funding of the first and third authors' visits to Rutgers.

The second author thanks the NSF, Grant No. DMS-9203726, for partial support.

The third author was partially supported by Basic Research Fund, Israeli Academy of Sciences. Publ. No. 520.

©1995 American Mathematical Society

We begin by describing the Ehrenfeucht-Fraïssé (or EF) games, after which we can state our results more precisely. If α is an ordinal and A and B are any structures, the game $\text{EF}_\alpha(A, B)$ is played between two players \forall and \exists who take turns choosing elements of $A \cup B$ through α rounds. Specifically, in each round \forall picks first an element of either A or B ; and then \exists picks an element of the other structure. The result is, at the end, two sequences $(a_\nu)_{\nu < \alpha}$ and $(b_\nu)_{\nu < \alpha}$ of elements of, respectively, A and B . Player \exists wins if and only if the function f which takes a_ν to b_ν is a partial isomorphism; otherwise \forall wins. If A and B have cardinality κ , \exists has a winning strategy for $\text{EF}_\kappa(A, B)$ if and only if A and B are isomorphic. (Let \forall list all the elements of $A \cup B$ during his moves.)

We consider variations of these games defined using trees. Given any tree T , we define the game $\text{EF}(A, B; T)$: the game is played as before except that player \forall must also, whenever it is his turn, pick a node of the tree strictly above his previous choices (thus his successive choices will form a branch—a linearly ordered subset—of the tree). The game ends when \forall can no longer pick a node above his previous choices. The criterion for winning is as before; that is, \exists wins if and only if the function f defined by the play is a partial isomorphism. We write $A \equiv^T B$ if \exists has a winning strategy in the game $\text{EF}(A, B; T)$. For the purposes of motivation consider first the case $\alpha = \omega$. (Our interest is in the case $\alpha = \omega_1$.) In this case, we consider only well-founded trees, i.e., trees without infinite branches; then for every such T , each play of the game $\text{EF}(A, B; T)$ is finite. (So $\text{EF}(A, B; T)$ may be regarded as an approximation to the game $\text{EF}_\omega(A, B)$.) Scott's Theorem implies that for each countable A there is a countable ordinal β such that if T_β is any tree of rank β , then for any countable B , B is isomorphic to A if and only if $A \equiv^{T_\beta} B$. In terms of infinitary languages, A is determined up to isomorphism (among countable structures) by a sentence of $L_{\infty\omega}$ of rank β .

For structures of cardinality \aleph_1 , it is natural to look at approximations to the EF game of length ω_1 and use trees of cardinality \aleph_1 which may have countably infinite branches, but do not have branches of cardinality \aleph_1 ; we call these *bounded trees*. For such T , each play of the game $\text{EF}(A, B; T)$ will end after countably many moves. We will say A is *T-equivalent* to B if $A \equiv^T B$. This relation provides a possible way of distinguishing between the ω_1 -separable groups constructed in [1] under the assumption of CH (cf. the remark after the quotation above).

By a theorem of Hyttinen [3], the entire class of bounded trees determines A up to isomorphism; that is, assuming CH if A and B are of cardinality \aleph_1 and $A \equiv^T B$ for all bounded trees, then A is isomorphic to B . The structure of the class of bounded trees is much more complicated than that of the class of well-founded trees (cf. [12]). However, in contrast to the situation for countable structures, there is not always a single tree which suffices to describe A up to isomorphism. Specifically, Hyttinen and Tuuri [4] proved (assuming CH) that there is a linear order A of cardinality \aleph_1 such that for every bounded tree T there is a linear order B_T of cardinality \aleph_1 such that $A \equiv^T B_T$ but A is not isomorphic to B_T . They call this result a *non-structure theorem* for A . It can be translated in terms of infinitary languages and says that there is no complete description of A by a sentence of a certain strong language $M_{\omega_2\omega_1}$ (which we shall not define here).

A similar non-structure theorem for p -groups was proved by Mekler and Oikkonen [10]; their theorem is proved by carrying over to p -groups, by means of a Hahn power construction, the result of Hyttinen and Tuuri. Whether the analogous result for \aleph_1 -free groups is a theorem of ZFC + CH remains open, but when we consider the question for \aleph_1 -separable groups, we obtain an independence result, which is the subject of this paper. In the first section we prove (with the help of the structural results referred to above) that assuming PFA

if A and B are ω_1 -separable groups of cardinality \aleph_1 such that $A \equiv^{\omega^2 + \omega} B$, then they are isomorphic (where $\omega^2 + \omega$ is the countable ordinal regarded as a—linearly ordered—tree).

See Theorem 2. Thus a single, simple tree contains enough information to classify any ω_1 -separable group—in the precise sense that a single sentence of M_{ω_2, ω_1} of “tree rank” $\omega^2 + \omega$ completely describes A .

In Section 2 we show, assuming \diamond , that not only does $\omega^2 + \omega$ not have the property above, but for any bounded tree T , there are non-isomorphic ω_1 -separable groups A^T and B^T of cardinality \aleph_1 which cannot be separated by T , in the sense that $A^T \equiv^T B^T$. (See Theorem 7.) The construction in Section 2 is strengthened in Section 3 to obtain a non-structure theorem (Theorem 8.):

there is an ω_1 -separable group A of cardinality \aleph_1 such that for every bounded tree T there is an ω_1 -separable group B^T of cardinality \aleph_1 which is not isomorphic to A but is T -equivalent to A .

(Note that A does not depend on T .)

We shall make use, at times, of the following simple lemma, where A^* denotes the dual of A , i.e. $\text{Hom}(A, \mathbb{Z})$.

Lemma 1. *Suppose $A \subseteq B$ and $A' \subseteq B' \subseteq C'$ where C'/B' is \aleph_1 -free, B/A is countable and $(B/A)^* = 0$. If $\theta : B \rightarrow C'$ such that $\theta[A] \subseteq A'$, then $\theta[B] \subseteq B'$.*

Proof. θ induces a homomorphism: $B/A \rightarrow C'/A'$. By the hypotheses, the composition of this map with the canonical surjection: $C'/A' \rightarrow C'/B'$ must be zero; that is, $\theta[B] \subseteq B'$. \square

1. A STRUCTURE THEOREM

An \aleph_1 -separable group A of cardinality \aleph_1 is characterized by the property that it has a *filtration*, that is, a continuous chain $\{A_\nu : \nu < \omega_1\}$ of countable free subgroups whose union is A and is such that $A_0 = 0$ and for all ν , $A_{\nu+1}$ is a direct summand of A . We say that two \aleph_1 -separable groups A and B are *quotient-equivalent* if and only if they have filtrations, $\{A_\nu : \nu < \omega_1\}$ and $\{B_\nu : \nu < \omega_1\}$, respectively, such that for every $\alpha < \omega_1$, $A_{\alpha+1}/A_\alpha$ is isomorphic to $B'_{\alpha+1}/B'_\alpha$. We say that A and B are *filtration-equivalent* if and only if they satisfy the stronger condition that for every $\alpha < \omega_1$ there is a *level-preserving isomorphism* $\theta_\alpha : A_{\alpha+1} \rightarrow B_{\alpha+1}$, i.e., an isomorphism such that, for every $\nu \leq \alpha$, $\theta[A_\nu] = B_\nu$. Under the assumption of MA + \neg CH, filtration-equivalence implies isomorphism.

In [8] (see also [2, Chapter VIII]) it is proved under the hypothesis of the Proper Forcing Axiom, PFA, that \aleph_1 -separable groups of cardinality \aleph_1 have a nice structure theory. More precisely, it is shown that, under PFA, every \aleph_1 -separable group of cardinality \aleph_1 is *in standard form*. (Roughly, this means that they have a “classical” construction. We will give a definition below.) Our goal in this section is to use that theory to prove the following:

Theorem 2. (PFA) $\omega^2 + \omega$ is a universal equivalence tree for the class of \aleph_1 -separable abelian groups of cardinality \aleph_1 . That is, any two \aleph_1 -separable abelian groups of cardinality \aleph_1 which are $\omega^2 + \omega$ -equivalent are isomorphic.

We shall see in the next section that this is not a theorem of ZFC. We begin with a weaker result.

Proposition 3. If A and A' are strongly \aleph_1 -free groups of cardinality \aleph_1 which are ω^2 -equivalent, then they are quotient equivalent.

Proof. Suppose that τ is a w.s. for \exists . Let C be a cub such that if $\alpha \in C$, then for any $n \in \omega$, as long as the first n moves of \forall are in $A_\alpha \cup A'_\alpha$, the replying moves of \exists given by τ are also in $A_\alpha \cup A'_\alpha$. If A and A' are not quotient-equivalent, there exists $\alpha \in C$ such that $A_{\alpha+1}/A_\alpha \oplus \mathbb{Z}^{(\omega)}$ is not isomorphic to $A'_{\alpha+1}/A'_\alpha \oplus \mathbb{Z}^{(\omega)}$. Now let \forall play the game so that during the first ω moves he makes sure that all elements of $A_\alpha \cup A'_\alpha$ are played; the result, since τ is a w.s., is that an isomorphism $f: A_\alpha \rightarrow A'_\alpha$ is obtained.

Then in the next ω moves, \forall plays so that all, and only, the elements of $A_\beta \cup A'_\beta$ are played for some $\beta \geq \alpha+1$. This is possible by using a bijection of ω with $\omega \times \omega$. The result is an extension of f to an isomorphism $f': A_\beta \rightarrow A'_\beta$. Then, since $A_\beta/A_{\alpha+1}$ and $A'_\beta/A'_{\alpha+1}$ are free, we have $A_\beta/A_\alpha \cong A_{\alpha+1}/A_\alpha \oplus A_\beta/A_{\alpha+1}$ and similarly on the other side. Since f' induces an isomorphism of A_β/A_α with A'_β/A'_α , we obtain a contradiction of the choice of α . \square

Suppose A is an \aleph_1 -separable group of cardinality \aleph_1 with a filtration $\{A_\nu : \nu \in \omega_1\}$, and let $E = \{\delta : A_\delta \text{ is not a direct summand of } A\}$; A is said to be *in standard form* if:

(1) it has a coherent system of projections $\{\pi_\nu : \nu \notin E\}$, i.e., projections $\pi_\nu : A \rightarrow A_\nu$ with the property that, for all $\nu < \tau$ in $\omega_1 \setminus E$, $\pi_\nu \circ \pi_\tau = \pi_\nu$; and

(2) for every $\delta \in E$ there is a ladder η_δ on δ and a subset Y_δ of $A_{\delta+1}$ such that $A_{\delta+1} = A_\delta + \langle Y_\delta \rangle$ and

$$(\dagger) \quad \text{for all } y \in \langle Y_\delta \rangle \text{ and all } \nu < \delta \text{ with } \nu \notin E, \pi_\nu(y) = \sum_{\alpha \in S} (\pi_{\alpha+1}(y) - \pi_\alpha(y)) \text{ where } S = \{\alpha \in \text{rge}(\eta_\delta) : \alpha < \nu\}.$$

(Here a ladder on δ means a strictly increasing function $\eta_\delta : \omega \rightarrow \delta$ with $\text{rge}(\eta_\delta) \subseteq \omega_1 \setminus E$ and $\sup \text{rge}(\eta_\delta) = \delta$.) This property is actually stronger than the usual definition of standard form (because of the assertion about the ladder); it can be shown that the Proper Forcing Axiom (PFA) implies that every strongly \aleph_1 -free group of cardinality \aleph_1 has this property (by essentially the same proof as in [2, Theorem VIII.3.3]).

Let $K_\alpha = \ker(\pi_\alpha)$ and let $K_{\alpha, \alpha+1} = K_\alpha \cap A_{\alpha+1}$. Notice that we can replace any y in Y_δ by $y + u$ where $u \in K_{\alpha, \alpha+1}$ for some $\alpha \in \delta \setminus E$, and we will still have a generating set of $A_{\delta+1}$ over A_δ which satisfies (\dagger) . Also we can, and will, assume that $A_{\nu+1}/A_\nu$ has infinite rank for every $\nu \notin E$.

Lemma 4. *Suppose A is in standard form. Then there is a filtration $\{A_\nu : \nu \in \omega_1\}$ of A and for each $\delta \in E = \{\delta : A_\delta \text{ is not a direct summand of } A\}$, there are: a ladder η_δ on δ ; and a subset $\bar{y}_\delta = \{y_{\delta,i} : i \in I\}$ of $A_{\delta+1}$ which is linearly independent mod A_δ such that if $\beta_n = \eta_\delta(n)$:*

1. for all $n \in \omega$, $\beta_n \notin E$; and
2. $A_{\delta+1}$ is generated mod A_δ by a set of elements of the form

$$(1) \quad \frac{t(\bar{y}_\delta) - a}{d}$$

where $t(\bar{y}_\delta)$ is a linear combination of the elements of \bar{y}_δ , $d \in \mathbb{Z}$, and $a \in \bigoplus_{n \in \omega} K_{\beta_n, \beta_{n+1}}$.

Moreover, given $\mu < \delta$, we can choose η_δ such that $\eta_\delta(0) > \mu$.

Proof. Let Y_δ and η_δ be as in the definition of standard form above. Let $\bar{y}_\delta = \{y_{\delta,i} : i \in I\}$ be a maximal linearly independent subset of Y_δ . By the remark preceding the lemma we can (by replacing $y_{\delta,i}$ by $y_{\delta,i} - \pi_{\mu+1}(y_{\delta,i})$) assume that $\eta_\delta(0) > \mu$.

If d divides $t(\bar{y}_\delta) \bmod A_{\nu+1}$ for some integer d and linear combination $t(\bar{y}_\delta)$, then d divides $t(\bar{y}_\delta) - a$ where $a = \pi_{\nu+1}(t(\bar{y}_\delta)) = \sum_{\beta \in S} \pi_{\beta, \beta+1}(t(\bar{y}_\delta))$ for some finite subset $S \subseteq \text{rg}(\eta_\delta)$. \square

Proposition 5. *Let G and G' be \aleph_1 -separable groups such that G is in standard form. Suppose that they have filtrations $\{G_\nu : \nu \in \omega_1\}$ and $\{G'_\nu : \nu \in \omega_1\}$ respectively such that the filtration of G attests that G is in standard form and $E = \{\nu \in \omega_1 : G_\nu \text{ is not a summand of } G\} = \{\nu \in \omega_1 : G'_\nu \text{ is not a summand of } G'\}$. Suppose also that for all limit ordinals δ , given a ladder η_δ on δ , there is an isomorphism $\theta_\delta : G_{\delta+1} \rightarrow G'_{\delta+1}$ such that for all $n \in \omega$, $\theta_\delta[G_{\eta_\delta(n)}] = G'_{\eta_\delta(n)}$ and $\theta_\delta[G_{\eta_\delta(n)+1}] = G'_{\eta_\delta(n)+1}$. Then G and G' are filtration-equivalent.*

Proof. We can assume that the filtration of G is as in Lemma 4. We prove by induction on ν the following:

if $\mu < \nu$ and $\mu, \nu \in \omega_1 \setminus E$ and $f : G_\mu \rightarrow G'_\mu$ is a level-preserving isomorphism, then f extends to a level-preserving isomorphism $g : G_\nu \rightarrow G'_\nu$.

If $\nu = \tau + 1$ where $\tau \notin E$, then the result follows easily by induction and the fact that G_ν/G_τ and G'_ν/G'_τ are free. If ν is a limit ordinal, choose a ladder ζ_ν on ν such that $\zeta_\nu(0) > \mu$ and for all n , $\zeta_\nu(n) \notin E$, and extend f successively, by induction, to $g_n : G_{\zeta_\nu(n)} \rightarrow G'_{\zeta_\nu(n)}$, and let $g = \bigcup_n g_n$.

The crucial case is when $\nu = \delta + 1$ where $\delta \in E$. Let η_δ be as in Lemma 4 with $\eta_\delta(0) > \mu$, and let θ_δ be the corresponding isomorphism given by the hypothesis of this proposition. Let $C_{\delta,n} = K_{\beta_n, \beta_{n+1}}$. By induction, extend f to a level-preserving isomorphism $f'_0 : G_{\eta_\delta(0)} \rightarrow G'_{\eta_\delta(0)}$ and then extend it to $g_0 : G_{\eta_\delta(0)+1} \rightarrow G'_{\eta_\delta(0)+1}$ by letting $g_0 \upharpoonright C_{\delta,0} = \theta_\delta \upharpoonright C_{\delta,0}$. Clearly g_0 is level-preserving. By induction extend g_0 to a level-preserving $f_1 : G_{\eta_\delta(1)} \rightarrow G'_{\eta_\delta(1)}$ and then to $g_1 : G_{\eta_\delta(1)+1} \rightarrow G'_{\eta_\delta(1)+1}$ by letting $g_1 \upharpoonright C_{\delta,1} = \theta_\delta \upharpoonright C_{\delta,1}$. Continuing in this way we obtain level-preserving isomorphisms $g_n : G_{\eta_\delta(n)+1} \rightarrow G'_{\eta_\delta(n)+1}$ for each n . Let $\tilde{g} = \bigcup_n g_n : G_\delta \rightarrow G'_\delta$.

By Lemma 4, $G_{\delta+1}$ is generated mod G_δ by a set of elements of the form

$$\frac{t(\bar{y}_\delta) - a}{d}$$

where $a \in \bigoplus_{n \in \omega} C_{\delta, n}$; hence $G'_{\delta+1}$ is generated mod G'_δ by elements

$$\frac{t(\theta_\delta(\bar{y}_\delta)) - \theta_\delta(a)}{d}.$$

But then since $\tilde{g}(a) = \theta_\delta(a)$ for each such a by construction, we can extend \tilde{g} to $g : G_{\delta+1} \rightarrow G'_{\delta+1}$ by sending each $y_{\delta, i}$ in \bar{y}_δ to $\theta_\delta(y_{\delta, i})$. Since \bar{y}_δ is linearly independent over G_δ , this is a well-defined homomorphism. \square

Theorem 6. *Suppose A and A' are \aleph_1 -separable groups of cardinality \aleph_1 and at least one of them is in standard form. If A and A' are $\omega^2 + \omega$ -equivalent, then they are filtration-equivalent.*

Proof. We can suppose that A is in standard form, and that we have chosen a filtration, $\{A_\nu : \nu \in \omega_1\}$, which attests to the fact. Moreover, we can assume that if $\delta \in E = \{\delta : A_\delta \text{ is not a direct summand of } A\}$, then $(A_{\delta+1}/A_\delta)^* = 0$. (Use Stein's Lemma [2, Exercise 3, p. 112], and replace $A_{\delta+1}$ by a direct summand, if necessary.)

Since A is quotient-equivalent to A' by Proposition 3, we can assume that there is a filtration $\{A'_\nu : \nu \in \omega_1\}$ of A' such that $E = \{\delta : A'_\delta \text{ is not a direct summand of } A'\}$ and for $\delta \in E$, $A'_{\delta+1}/A'_\delta \cong A_{\delta+1}/A_\delta = 0$, so in particular $(A'_{\delta+1}/A'_\delta)^* = 0$.

Fix a bijection $\psi_{\alpha\beta} : \omega \rightarrow (A_\beta \setminus A_\alpha) \cup (A'_\beta \setminus A'_\alpha)$ for each $\alpha < \beta$. Let $\psi = \{\psi_{\alpha\beta} : \alpha < \beta < \omega_1\}$.

When we talk about moves in a game, we refer to the game $\text{EF}_{\omega^2 + \omega}(A, A')$. Given a strictly increasing finite sequence of countable ordinals $\alpha_1 < \alpha_2 < \dots < \alpha_n$, we will say that \forall plays according to ψ and $\langle \alpha_1, \alpha_2, \dots, \alpha_n \rangle$ for the first ωn moves if the $\omega k + l$ move of player \forall is $\psi_{\alpha_k \alpha_{k+1}}(l)$ for $k = 0, \dots, n - 1$ and $l \in \omega$ (where $\alpha_0 = 0$).

Suppose that τ is a w.s. for \exists in the game $\text{EF}_{\omega^2 + \omega}(A, A')$. Let C be the set of all $\delta < \omega_1$ such that, for any integers $n > 0$ and $m \geq 0$ and any ordinals $\alpha_1 < \alpha_2 < \dots < \alpha_n < \delta$, if \forall plays according to ψ and $\langle \alpha_1, \alpha_2, \dots, \alpha_n \rangle$ for the first ωn moves and then plays any elements of A_δ for the next m moves, then the responses of \exists using τ are all in $A_\delta \cup A'_\delta$.

Then C is a cub: for the proof of unboundedness, note that there are only countably many possibilities that one has to close under: choice of n and m , choice of $\alpha_1 < \alpha_2 < \dots < \alpha_n$, and choice of moves $\omega n, \omega n + 1, \dots, \omega n + m - 1$. (The earlier moves are determined by the $\psi_{\alpha_k \alpha_{k+1}}$ and by τ .)

There is a continuous strictly increasing function $\tilde{h} : \omega_1 \rightarrow \omega_1$ whose range is C . Define $h : \omega_1 \rightarrow \omega_1$ by

$$h(\beta) = \begin{cases} \tilde{h}(\beta) + 1 & \text{if } \beta \text{ is a successor and } \tilde{h}(\beta) \in E, \\ \tilde{h}(\beta) & \text{otherwise.} \end{cases}$$

Let $G_\alpha = A_{h(\alpha)}$ and $G'_\alpha = A'_{h(\alpha)}$. Then for successor β , G_β is a summand of A and, for limit δ , $G_\delta = A_{\tilde{h}(\delta)} + \bigcup_{\beta < \delta} A_{h(\beta)} = \bigcup_{\beta < \delta} G_\beta$, so $\{G_\alpha : \alpha \in \omega_1\}$ (resp. $\{G'_\alpha : \alpha \in \omega_1\}$) is a filtration of A (resp. A'). Given a limit ordinal δ and a ladder η_δ on δ , it follows—from Lemma 1 and the definition of C —that there is an isomorphism $\theta_\delta : G_{\delta+1} \rightarrow G'_{\delta+1}$ such that, for all $n \in \omega$, $\theta_\delta[G_{\eta_\delta(n)}] = G'_{\eta_\delta(n)}$ and $\theta_\delta[G_{\eta_\delta(n)+1}] = G'_{\eta_\delta(n)+1}$. In fact, θ_δ is the partial isomorphism which

results because \exists wins the game where the $\omega k + l$ move of \forall is

$$\Psi_{h(\eta_\delta(n)), h(\eta_\delta(n)+1)}(l)$$

when $k = 2n$, and is

$$\Psi_{h(\eta_\delta(n+1)), h(\eta_\delta(n+1))}(l)$$

when $k = 2n + 1$, and the $\omega^2 + m$ move of \forall is $\Psi_{h(\delta), h(\delta+1)}(m)$.

Thus we have satisfied the hypotheses of Proposition 5 so we conclude that A and A' are filtration-equivalent. \square

Now we can prove Theorem 2. PFA implies that every strongly \aleph_1 -free abelian group of cardinality \aleph_1 is \aleph_1 -separable and in standard form. Moreover, assuming PFA, filtration-equivalent \aleph_1 -separable groups of cardinality \aleph_1 are isomorphic. Thus the result follows from Theorem 6.

2. A DIAMOND CONSTRUCTION: ONE TREE

The result to be proved in this section is the following:

Theorem 7. *Assume \diamond . For any bounded tree T_1 there exist non-isomorphic \aleph_1 -separable groups G^0 and G^1 of cardinality \aleph_1 which are T_1 -equivalent (and filtration-equivalent) are both in standard form.*

Proof. We will present the proof in layers of increasing detail.

(I) Fix a stationary subset E of ω_1 consisting of limit ordinals and such that E is the disjoint union of two uncountable subsets E_0 and E_1 such that $\diamond(E_1)$ holds.

Given a bounded tree T (which in practice will be determined by, but not equal to, T_1), we shall identify its nodes with countable ordinals in such a way that if $\nu <_T \mu$ (in the tree ordering), then $\nu < \mu$ (as ordinals).

By induction on $\alpha < \omega_1$ we will define the following data:

1. continuous chains $\{G_\nu^l : \nu < \alpha\}$ of countable free groups (for $l = 0, 1$) such that, for all $\nu < \mu < \alpha$, G_μ^l/G_ν^l is free if $\nu \notin E_1$, and if $\nu \in E_1$, then $G_{\nu+1}^l/G_\nu^l$ has rank at most 1;
2. homomorphisms $\pi_{\nu, \mu}^l : G_\mu^l \rightarrow G_\nu^l$ for $\nu \leq \mu < \alpha$ and $\nu \notin E_1 \cup \{\sigma + 1 : \sigma \in E_0\}$ such that: $\pi_{\nu, \mu}^l$ is the identity on G_ν^l ; for $\nu \leq \mu < \rho$, $\pi_{\nu, \mu}^l \subseteq \pi_{\nu, \rho}^l$; and for $\tau < \nu \leq \mu$, $\pi_{\tau, \nu}^l \circ \pi_{\nu, \mu}^l = \pi_{\tau, \mu}^l$ (i.e., $\pi_{\nu, \mu}^l$ is a projection and the system of projections is coherent);
3. for each ν with $\nu + 1 < \alpha$ an isomorphism $f_\nu^0 : G_{\nu+1}^0 \rightarrow G_{\nu+1}^1$ satisfying:

$$\text{if } \nu_1 <_T \nu_2, \text{ then } f_{\nu_2}^0 \upharpoonright G_{\nu_1+1}^0 = f_{\nu_1}^0.$$

(These partial isomorphisms will give \exists her winning strategy.)

For convenience we will use f_ν^1 to denote $(f_\nu^0)^{-1} : G_{\nu+1}^1 \rightarrow G_{\nu+1}^0$.

Define $G^l = \bigcup_{\nu < \omega_1} G_\nu^l$. (It depends on T , but we suppress that in the notation.) Now we will indicate how we choose T so that G^0 and G^1 are T_1 -equivalent.

Let $T_2 = {}^{<\omega_1} \omega_1 \setminus \emptyset$, i.e., the tree of non-empty countable sequences of countable ordinals, partially ordered by inclusion (so it has \aleph_1 nodes of height 0).

Let T be the product $T_1 \otimes T_2$, i.e., the (bounded) tree whose nodes are elements $(s, \sigma) \in T_1 \times T_2$, where s and σ have the same height, and the partial ordering is defined coordinate-wise. (As above, we identify the nodes of T with ordinals.)

Suppose we are able to carry out the construction outlined above for this T . Then since the G_ν^l are free, G^l is \aleph_1 -free. Moreover, for $\nu \notin E_1$, $\bigcup_{\mu < \omega_1} \pi_{\nu, \mu}^l : G^l \rightarrow G_\nu^l$ is a projection which shows that G_ν^l is a direct summand of G^l ; so G^l is \aleph_1 -separable (and has a coherent system of projections; the fact that it is in standard form will follow from the details of the construction—see part (V)).

We claim that G^0 and G^1 are T_1 -equivalent. In fact, here is \exists 's winning strategy in the T_1 -game. If in his first move \forall plays $s_0 \in T_1$ (which we may assume has height 0), and $y_0 \in G_{\gamma_0}^{l_0}$, \exists chooses α_0 such that $(s_0, \langle \alpha_0 \rangle) \in T$ is the element ν_0 of the enumeration of T , where $\nu_0 \geq \gamma_0$; and she plays $f_{\nu_0}^{l_0}(y_0) \in G_{\nu_0+1}^{l_0}$. (Note that the domain of $f_{\nu_0}^{l_0}$ is $G_{\nu_0+1}^{l_0} \supseteq G_{\gamma_0}^{l_0}$.) Suppose that after β moves \forall has chosen $s_0 <_{T_1} s_1 <_{T_1} \dots <_{T_1} s_i <_{T_1} \dots$ in the tree and $y_0, y_1, \dots, y_i, \dots$ in the groups where $y_i \in G^i$ ($i < \beta$), and \exists has responded to the i th move with $f_{\nu_i}^i(y_i)$ where $\nu_i = (s_i, \langle \alpha_0, \dots, \alpha_i \rangle)$. Now if \forall plays $s_\beta >_{T_1} s_i$ ($i < \beta$)—which we can assume has height β —and $y_\beta \in G_{\gamma_\beta}^{l_\beta}$, then \exists chooses α_β such that $(s_\beta, \langle \alpha_0, \dots, \alpha_\beta \rangle)$ is $\nu_\beta \geq \gamma_\beta$, and plays $f_{\nu_\beta}^{l_\beta}(y_\beta)$. Notice that $\nu_\beta >_T \nu_i$, so $f_{\nu_\beta}^{l_\beta}$ extends $f_{\nu_i}^i$ for $l = 0, 1$. Therefore the sequence of moves determines a partial isomorphism, so \exists will win.

(II) Of course, we also want to do the construction so that G^0 and G^1 are not isomorphic. This will be achieved by our construction of $G_{\delta+1}^l$ for $\delta \in E_1$ (plus the requirement 4 below); when $\delta \in E_1$ we will make use of the “guess” provided by $\diamond(E_1)$ of an isomorphism: $G_\delta^0 \rightarrow G_\delta^1$.

Our construction will be such that when $\alpha = \mu + 1$ where $\mu \notin E$, then

$$G_\alpha^l = G_\mu^l \oplus \mathbb{Z}x_{\mu,0}^l \oplus \mathbb{Z}x_{\mu,1}^l.$$

When $\alpha = \sigma + 1$ where $\sigma \in E_0$, then

$$G_\alpha^l = G_\sigma^l \oplus \bigoplus_{n \in \omega} \mathbb{Z}u_{\sigma,n}^l \oplus \mathbb{Z}v_{\sigma,n}^l.$$

We define

$$w_{\sigma,n} = 2u_{\sigma,n+1}^0 - u_{\sigma,n}^0.$$

Notice that $\{w_{\sigma,n} : n \in \omega\}$ generates a pure subgroup of $\bigoplus_{n \in \omega} \mathbb{Z}u_{\sigma,n}^0$ which is not a direct summand. Hence there is no isomorphism of $\bigoplus_{n \in \omega} \mathbb{Z}u_{\sigma,n}^0 \oplus \mathbb{Z}v_{\sigma,n}^0$ with $\bigoplus_{n \in \omega} \mathbb{Z}u_{\sigma,n}^1 \oplus \mathbb{Z}v_{\sigma,n}^1$ which takes each $w_{\sigma,n}$ to $v_{\sigma,n}^1$. In order to carry out the inductive construction we will define in addition:

4. subsets $W_\alpha[\Theta]$ of G_α^0 for every non-empty finite subset Θ of α which is an antichain in T , satisfying:
 - (a) for all $\alpha < \beta$, $W_\alpha[\Theta] \subseteq W_\beta[\Theta]$;
 - (b) every element of $W_\alpha[\Theta]$ is of the form $w_{\sigma,n}$ for some $\sigma \in E_0$, and $n \in \omega$.

The functions f_α^0 will be required to satisfy:

- (c) for all $\mu \leq \alpha$, $j \in \{0, 1\}$, $f_\alpha^0(x_{\mu,j}^0) = x_{\mu,j}^1$; moreover, if $w_{\sigma,n} \in W_{\alpha+1}[\Theta]$ and $\Theta \cap \{\nu : \nu \leq_T \alpha\} \neq \emptyset$, then $f_\alpha^0(w_{\sigma,n}) = v_{\sigma,n}^1$.

For any finite antichain Θ in T , let $W[\Theta] = \bigcup_\alpha W_\alpha[\Theta]$.

Now we will outline how we do the construction so that G^0 and G^1 are not isomorphic. Before we start, we choose a function Υ with domain E_0 which maps onto the set of all ω -sequences $\langle \Theta_n : n \in \omega \rangle$ of finite subsets of T such that $\bigcup_{n \in \omega} \Theta_n$ is an antichain; we also require that $\Upsilon(\sigma) = \langle \Theta_n : n \in \omega \rangle$ for uncountably many σ .

Suppose now that we have defined G_ν^l for $\nu \leq \alpha$. If $\alpha = \sigma \in E_0$, then $G_{\sigma+1}^l$ will be defined as indicated above and is such that (as we will prove)

$$(II.1) \text{ for all } e \in \{1, -1\}, \text{ there is no isomorphism of } G_{\sigma+1}^0 \text{ with } \bigoplus_{n \in \omega} \mathbb{Z}v_{\sigma,n}^1 \oplus C \text{ for any } C, \text{ which for all } n \in \omega \text{ takes } w_{\sigma,n} \text{ to } ev_{\sigma,n}^1.$$

Moreover $w_{\sigma,n}$ will be put into $W_{\sigma+1}[\Theta_n^\sigma]$. (This is the only way that an element becomes a member of a $W_\alpha[\Theta]$.)

If $\alpha = \delta \in E_1$ and $\beta < \delta$, we introduce the notation $A_{\beta,\delta} = \{t : t \text{ is } <_T\text{-minimal in } \delta \setminus \beta\}$ —so $A_{\beta,\delta}$ is an antichain. We fix finite subsets $\Theta_n^{\beta,\delta}$ of $A_{\beta,\delta}$ which form a chain such that $\bigcup_{n \in \omega} \Theta_n^{\beta,\delta} = A_{\beta,\delta}$. We consider the prediction given by $\diamond(E_1)$ of an isomorphism $h : G_\delta^0 \rightarrow G_\delta^1$ and we ask whether the following holds:

$$(II.2) \quad \exists \beta < \delta \forall e \in \{1, -1\} \forall n \in \omega \forall \mu < \delta \exists w_{\sigma,m} \in W_\delta[\Theta_n^{\beta,\delta}] \text{ such that } \mu < \sigma < \delta \text{ and } h(w_{\sigma,m}) \neq ev_{\sigma,m}^1.$$

We will do the construction of $G_{\delta+1}^l$ so that:

$$(II.3) \text{ If (II.2) holds, then } G_{\delta+1}^l/G_\delta^l \text{ is non-free rank 1 and } h \text{ does not extend to a homomorphism: } G_{\delta+1}^0 \rightarrow G_{\delta+1}^1.$$

Assuming we can do all of this, let us see why G^0 is not isomorphic to G^1 . Suppose, to the contrary, that there is an isomorphism $H : G^0 \rightarrow G^1$. Then there is a stationary set, S , of $\delta \in E_1$ where $\diamond(E_1)$ guesses $h = H \upharpoonright G_\delta^0$ and $H : G_\delta^0 \rightarrow G_\delta^1$. Note that Lemma 1 implies that H must map $G_{\delta+1}^0$ into $G_{\delta+1}^1$ because $G_{\delta+1}^0/G_\delta^0$ is non-free rank 1 but $G^1/G_{\delta+1}^1$ is \aleph_1 -free by construction. If for any such δ (II.2) holds, then $H \upharpoonright G_{\delta+1}^0$ would extend $h = H \upharpoonright G_\delta^0$, contradicting (II.3).

Since (II.2) fails, for all $\delta \in S$ and all $\beta < \delta$ there exists $e \in \{1, -1\}$ and a finite subset Θ of $A_{\beta,\delta}$ such that $H(w_{\sigma,n}) = ev_{\sigma,n}^1$ for all $w_{\sigma,n} \in W_\delta[\Theta]$ for sufficiently large $\sigma < \delta$. Now there is a cub C such that for all $\delta \in C$, all $e \in \{1, -1\}$, all $\beta < \delta$, and all finite subsets Θ of $A_{\beta,\delta}$, if $H(w_{\sigma,n}) \neq ev_{\sigma,n}^1$ for uncountably many $w_{\sigma,n} \in W[\Theta]$, then $H(w_{\sigma,n}) \neq ev_{\sigma,n}^1$ for $w_{\sigma,n} \in W_\delta[\Theta]$ with arbitrarily large $\sigma < \delta$. Thus for all $\delta \in C \cap S$ and all $\beta < \delta$, there exist $e \in \{1, -1\}$, and a finite subset Θ of $A_{\beta,\delta}$ such that $H(w_{\sigma,n}) = ev_{\sigma,n}^1$ for all $w_{\sigma,n} \in W[\Theta]$ with $\sigma > \alpha_\Theta$. Since $C \cap S$ is uncountable, it follows easily that there exist $e \in \{1, -1\}$ and an uncountable set $\{\Theta_\nu : \nu < \omega_1\}$ of pairwise disjoint finite antichains such that $H(w_{\sigma,n}) = ev_{\sigma,n}^1$ for all $w_{\sigma,n} \in W[\Theta_\nu]$

with $\sigma > \alpha_\nu$ for all $\nu < \omega_1$. Since T has no uncountable branches, by a standard argument (see, for example, [5, Lemma 24.2, p. 245]), there is a countably infinite subset $\{\nu_n : n \in \omega\}$ of ω_1 such that $\bigcup\{\Theta_{\nu_n} : n \in \omega\}$ is an antichain. There exists $\sigma \in E_0$ such that $\Upsilon(\sigma) = \langle \Theta_{\nu_n} : n \in \omega \rangle$ and $\sigma > \alpha_{\nu_n}$ for all n . Now $H \upharpoonright G_{\sigma+1}^0$ is such that, for all $n \in \omega$, $H(w_{\sigma,n}) = ev_{\sigma,n}^1$ since $w_{\sigma,n} \in W_{\sigma+1}[\Theta_{\nu_n}]$; this contradicts (II.1), since $\bigoplus_{n \in \omega} \mathbb{Z}v_{\sigma,n}^1$ is a direct summand of $G_{\sigma+1}^1$, and hence of G^1 (by 2).

(III) The next step is to describe in detail the recursive construction of the data satisfying the properties 1, 2, 3 and 4, as well as (II.1) and (II.3). So assume that we have defined G_ν^l and $W_\nu[\Theta]$ for $\nu < \alpha$ and f_ν^l for $\nu + 1 < \alpha$.

There are several cases to consider.

Case 1: α is a limit ordinal. We let $G_\alpha^l = \bigcup_{\nu < \alpha} G_\nu^l$, $W_\alpha[\Theta] = \bigcup_{\nu < \alpha} W_\nu[\Theta]$. Clearly the desired properties are satisfied.

If α is a successor, $\alpha = \mu + 1$, we will define G_α^l so that

(III.1) if $B = \{t : t <_T \mu\}$ and we define $g_B = \bigcup\{f_t^0 : t \in B\}$, then g_B (which is a function by 3) extends to an isomorphism, f_μ^0 , of G_μ^0 onto G_α^1 which satisfies 4(c); i.e. for all $\nu \leq \mu$, $j \in \{0, 1\}$, $f_\mu^0(x_{\nu,j}^0) = x_{\nu,j}^1$ and if $w_{\sigma,n} \in W_\alpha[\Theta]$ and $\Theta \cap \{\nu : \nu \leq_T \mu\} \neq \emptyset$, then $f_\mu^0(w_{\sigma,n}) = v_{\sigma,n}^1$.

Leaving the verification of (III.1) to the next part, we will show how to define the data at α (except for the definition of the $\pi_{\sigma,\alpha}^l$ which we defer to part (V)).

Case 2: $\alpha = \mu + 1$ for some $\mu \notin E$. As described above, define

$$G_\alpha^l = G_\mu^l \oplus \mathbb{Z}x_{\mu,0}^l \oplus \mathbb{Z}x_{\mu,1}^l.$$

Let $W_\alpha[\Theta] = W_\mu[\Theta]$ for every $\Theta \subseteq \mu$ ($= \emptyset$ if Θ is not a subset of μ). Assuming (III.1), we have f_μ^0 as desired.

Case 3: $\alpha = \sigma + 1$, where $\sigma \in E_0$. In this case, as stated before,

$$G_\alpha^l = G_\sigma^l \oplus \bigoplus_{n \in \omega} \mathbb{Z}u_{\sigma,n}^l \oplus \mathbb{Z}v_{\sigma,n}^l$$

and recall that $w_{\sigma,n}$ is defined to be $2u_{\sigma,n+1}^0 - u_{\sigma,n}^0$. Say $\Upsilon(\sigma) = \langle \Theta_n^\sigma : n \in \omega \rangle$. Define

$$W_\alpha[\Theta] = \begin{cases} W_\sigma[\Theta] \cup \{w_{\sigma,n}\} & \text{if } \Theta = \Theta_n^\sigma, \\ W_\sigma[\Theta] & \text{otherwise.} \end{cases}$$

Assuming (III.1) (with $\mu = \sigma$), we can define f_σ^0 . Now let us see why (II.1) holds. Suppose to the contrary that there is an isomorphism $H : G^0 \rightarrow G^1$ contradicting (II.1). Now $\bigoplus_{n \in \omega} \mathbb{Z}v_{\sigma,n}^1$ is a direct summand of G_α^1 and hence (by 2) a direct summand of G^1 . Thus $H^{-1}[\bigoplus_{n \in \omega} \mathbb{Z}v_{\sigma,n}^1]$ is a direct summand of G^0 . But by assumption on H , $H^{-1}[\bigoplus_{n \in \omega} \mathbb{Z}v_{\sigma,n}^1] = \bigoplus_{n \in \omega} \mathbb{Z}w_{\sigma,n}$ and the latter is *not* a direct summand of G^0 because the coset of $u_{\sigma,0}^0$ is a non-zero element of $G^0 / \bigoplus_{n \in \omega} \mathbb{Z}w_{\sigma,n}$ which is divisible by all power of 2 by definition of the $w_{\sigma,n}$.

Case 4: $\alpha = \delta + 1$, where $\delta \in E_1$. If (II.2) fails, let $G_{\delta+1}^l = G_\delta^l$. Otherwise, let β be as in (II.2). We introduce some ad hoc notation. For any finite subset

Θ of $A_{\beta, \delta}$, let f_Θ be the function whose domain is the subgroup generated by $\{x_{\mu, j}^0 : \mu \notin E, \mu < \delta, j \in \{0, 1\}\} \cup W_\delta[\Theta]$ such that $f_\Theta(x_{\mu, j}^0) = x_{\mu, j}^1$ and $f_\Theta(w_{\sigma, n}) = v_{\sigma, n}^1$. Notice that, for all $u \in \text{dom}(f_\Theta)$ and all $\nu \in \Theta$, if $\nu \leq_T \rho$ and $u \in \text{dom}(f_\rho^0)$, then $f_\Theta(u) = f_\rho^0(u)$ by 4(c). Let $\Theta_n^{\beta, \delta}$ be as before (finite subsets forming a chain whose union is $A_{\beta, \delta}$); for short, let $\Theta_n = \Theta_n^{\beta, \delta}$. We claim that:

(III.2) given $m, m' \in \mathbb{Z} \setminus \{0\}$, $n \in \omega$, $\gamma \in G_\delta^1$, for arbitrarily large $\gamma < \delta$ there exists $k^0 \in \text{dom}(f_{\Theta_n}) \cap G_{\gamma+2}^0$ such that k^0 is pure-independent mod $G_{\gamma+1}^0$ and is such that $mh(k^0) \neq m'f_{\Theta_n}(k^0) + \gamma$. Moreover, $f_{\Theta_n}(k^0)$ is pure-independent mod $G_{\gamma+1}^1$.

Supposing this is true—we will prove it in part (IV)—let us define $G_{\delta+1}^1$. Fix a ladder η_δ on δ . Also, enumerate in an ω -sequence all triples $\langle r, d, v \rangle$ where $r \in \omega$, $d \in \mathbb{Z} \setminus \{0\}$, and $g \in G_\delta^1$ so that the n th triple $\langle r, d, g \rangle$ satisfies $n > r$. By (III.2) we can inductively define primes p_n , ordinals $\gamma_n \geq \eta_\delta(n)$, and elements $k_{\delta, n}^0 \in \text{dom}(f_{\Theta_n}) \cap G_{\gamma_n+2}^0$ pure-independent over $G_{\gamma_n+1}^0$ such that (if the n th triple is $\langle r, d, g \rangle$) p_n does not divide $mh(k_{\delta, n}^0) - m'f_{\Theta_n}(k_{\delta, n}^0) - \gamma$ where

$$m = \prod_{i=0}^{n-1} p_i,$$

$$m' = d \prod_{i=r}^{n-1} p_i,$$

$$\gamma = \sum_{j=0}^{n-1} \left(\prod_{i=0}^{j-1} p_i \right) h(k_{\delta, j}^0) + g - d \sum_{j=r}^{n-1} \left(\prod_{i=r}^{j-1} p_i \right) f_{\Theta_n}(k_{\delta, j}^0).$$

(Note that since G_δ^1 is free, every non-zero element is divisible by only finitely many primes, so we can take p_n to be any sufficiently large prime.) Then we let $G_{\delta+1}^0$ be generated by $G_\delta^0 \cup \{z_{\delta, n}^0 : n \in \omega\}$ modulo the relations

$$p_n z_{\delta, n+1}^0 = z_{\delta, n}^0 + k_{\delta, n}^0$$

and $G_{\delta+1}^1$ is defined similarly, except that we impose the relations

$$p_n z_{\delta, n+1}^1 = z_{\delta, n}^1 + f_{\Theta_n}(k_{\delta, n}^0).$$

We need to show that h does not extend to a homomorphism: $G_{\delta+1}^0 \rightarrow G_{\delta+1}^1$. If it does, then $h(z_{\delta, 0}^0) = dz_{\delta, r}^1 + g$ for some $r \in \omega$, $d \in \mathbb{Z} \setminus \{0\}$, and $g \in G_\delta^1$. Let n be such that $\langle r, d, g \rangle$ is the n th triple in the list. Now, in $G_{\delta+1}^0$ we have

$$\left(\prod_{i=0}^n p_i \right) z_{\delta, n+1}^0 = z_{\delta, 0}^0 + \sum_{j=0}^n \left(\prod_{i=0}^{j-1} p_i \right) k_{\delta, j}^0$$

so, applying h , we conclude that p_n divides

$$dz_{\delta, r}^1 + g + \sum_{j=0}^n \left(\prod_{i=0}^{j-1} p_i \right) h(k_{\delta, j}^0).$$

On the other hand, in $G_{\delta+1}^1$ we have p_n divides

$$dz_{\delta,r}^1 + d \sum_{j=r}^n \left(\prod_{i=r}^{j-1} p_i \right) f_{\Theta_j}(k_{\delta,j}^0)$$

so, subtracting, we obtain a contradiction since p_n divides

$$mh(k_{\delta,n}^0) - m' f_{\Theta_n}^0(k_{\delta,n}^0) - y,$$

where m , m' , and y are as above.

We let $W_{\delta+1}[\Theta] = W_{\delta}[\Theta]$ for any subset Θ of δ (and $= \emptyset$ if $\Theta \not\subseteq \delta$). By (III.1) we can define f_{δ}^0 .

(IV) In this layer we will prove (III.1) and (III.2).

First let us prove (III.2) since for the purposes of proving (III.1) we will need more information about the nature of the elements $k_{\delta,n}^0$. Fix m , m' , n , y , γ as in (III.2); there are several cases. In the first two cases we can use any $\gamma < \delta$.

Case (i): $y \neq 0$. If neither $x_{\gamma+1,0}^0$ nor $x_{\gamma+1,1}^0$ will serve for k^0 , then $x_{\gamma+1,0}^0 - x_{\gamma+1,1}^0$ will.

Case (ii): $y = 0$, $m \neq \pm m'$. Let $k^0 = x_{\gamma+1,0}^0$. Then by construction, k^0 generates a cyclic summand of G_{δ}^0 ; hence $f_{\Theta_n}(k^0)$ and $h(k^0)$ both generate cyclic summands of G_{δ}^1 . Hence $mh(k^0) \neq m' f_{\Theta_n}^0(k^0)$.

Case (iii): $y = 0$, $m = m'$. Pick $\gamma = \sigma$ sufficiently large so that there exists $w_{\sigma,j} \in G_{\gamma+1}^0 \cap W_{\delta}[\Theta_n]$ such that $f_{\Theta_n}(w_{\sigma,j}) \neq h(w_{\sigma,j})$. If $x_{\gamma+1,0}^0$ will not serve for k^0 (i.e., $h(x_{\gamma+1,0}^0) = x_{\gamma+1,0}^1$), then we can take k^0 to be $x_{\gamma+1,0}^0 + w_{\sigma,j}$.

Case (iv): $y = 0$, $m = -m'$. Similarly k^0 can be taken to be of the form $x_{\gamma+1,0}^0$ or $x_{\gamma+1,0}^0 - w_{\sigma,j}$ where $f_{\Theta_n}(w_{\sigma,j}) \neq -h(w_{\sigma,j})$.

Now if we examine the construction in Case 4 of (III) and the proof above we see that

(IV.1) each $k_{\delta,n}^0$ can be (and will be) taken to be of the form $x_{\mu_n,j_n}^0 \pm \xi_{\delta,n}$ where $\xi_{\delta,n}$ is 0, $x_{\sigma,j}^0$ or $w_{\sigma,j}$ for some σ, j .

We will say that $w_{\sigma,j}$ is a part of $k_{\delta,n}^0$ in case $\xi_{\delta,n}$ is $w_{\sigma,j}$.

Before beginning the proof of (III.1), let us observe the following facts:

(IV.2) Given $\sigma \in E_0$ and $N \in \omega$, there is an isomorphism $g' : \bigoplus_{n \in \omega} \mathbb{Z}u_{\sigma,n}^0 \oplus \mathbb{Z}v_{\sigma,n}^0 \rightarrow \bigoplus_{n \in \omega} \mathbb{Z}u_{\sigma,n}^1 \oplus \mathbb{Z}v_{\sigma,n}^1$ such that $g'(w_{\sigma,n}) = v_{\sigma,n}^1$ for $n \leq N$ and $g'(u_{\sigma,n}^0) = u_{\sigma,n}^1$ for $n \geq N+1$.

Indeed, we can define $g'(u_{\sigma,n}^0) = 2g'(u_{\sigma,n+1}^0) - v_{\sigma,n}^1$ for $n \leq N$ (and the other values appropriately).

(IV.3) Given an isomorphism $g : G_{\delta}^0 \rightarrow G_{\delta}^1$ where $\delta \in E_1$, we can extend g to an isomorphism $g' : G_{\delta+1}^0 \rightarrow G_{\delta+1}^1$ provided that (using the notation of Case 4) $g(k_{\delta,n}^0) = f_{\Theta_n}(k_{\delta,n}^0)$ for almost all $n \in \omega$.

Indeed, if $g(k_{\delta,n}^0) = f_{\Theta_n}(k_{\delta,n}^0)$ for all $n \geq N$, we can define $g'(z_{\delta,n}^0) = z_{\delta,n}^1$ for $n \geq N$ and $g'(z_{\delta,n}^0) = p_n g'(z_{\delta,n+1}^0) - g(k_{\delta,n}^0)$ for $n < N$ by “downward induction”. We will apply (IV.3) to the situation of (III.1), with $g = g_B$, $\delta = \mu$, $\delta + 1 = \alpha$; if we are in Case 4, then the hypothesis on g in (IV.3) will hold if there exists $t \in B$ such that $t \geq \beta$ (where β is as in Case 4).

We return to the notation of (III.1). Let $\tau = \sup\{t + 1 : t \in B\}$; then $\text{dom } g_B = G_\tau^0$. Assume first that $\tau = \mu$. In case $G_{\mu+1}^l/G_\mu^l$ is free there is no problem extending g_B ; in the other case $\mu = \delta \in E_1$ and by the remarks above we can extend g_B since there exists $t \in B$ such that $t \geq \beta$ (since $\sup B = \delta$).

We are left with the case when $\tau < \mu$. We will first define an extension of g_B to a partial isomorphism \tilde{g}_B whose domain is

$$\text{dom}(g_B) + \left\langle \begin{array}{l} \{x_{\nu,j}^0 : \nu \leq \mu, j = 0, 1\} \\ \cup \{u_{\sigma,n}^0 : \sigma \in E_0 \cap \mu + 1, n \in \omega\} \\ \cup \{v_{\sigma,n}^0 : \sigma \in E_0 \cap \mu + 1, n \in \omega\} \end{array} \right\rangle.$$

Notice that every $k_{\delta,n}^0$ for $\delta \leq \mu$, $n \in \omega$ belongs to the domain of \tilde{g}_B . We let $\tilde{g}_B(x_{\nu,j}^0) = x_{\nu,j}^1$ for all ν, j . By enumerating in an ω -sequence the set $(E_0 \cup E_1) \cap (\mu + 1)$ we can define by recursion the values $\tilde{g}_B(u_{\sigma,n}^0)$ and $\tilde{g}_B(v_{\sigma,n}^0)$ so that:

- $\tilde{g}_B(w_{\sigma,n}) = v_{\sigma,n}^1$ whenever $w_{\sigma,n} \in W_{\mu+1}[\Theta]$ for some Θ with $B \cap \Theta \neq \emptyset$;
- for all $\sigma \in E_0$ with $\tau \leq \sigma \leq \mu$, for almost all $n \in \omega$, $\tilde{g}_B(u_{\sigma,n}^0) = u_{\sigma,n}^1$; and
- for all $\delta \in E_1$ with $\tau \leq \delta \leq \mu$, for almost all $n \in \omega$, if (for some σ, m) $w_{\sigma,m}$ is a part of $k_{\delta,n}^0$, then $\tilde{g}_B(w_{\sigma,m}) = v_{\sigma,m}^1$.

The first condition is required by 4(c). In view of (IV.2), there is no conflict between the first two conditions because, for any $\sigma \in E_0$, $\bigcup_{n \in \omega} \Theta_n^\sigma$ is an antichain, so there is at most one n such that $\Theta_n^\sigma \cap B \neq \emptyset$.

To be sure that the third condition can indeed be satisfied, we need to consider the case that, for some $\delta \in E_1$, there are infinitely many n such that there exists w_{σ_n, m_n} which is a part of $k_{\delta,n}^0$ and belongs to the domain of g_B . Say this is the case for n belonging to the (infinite) set $Y \subseteq \omega$ (for a fixed δ). Then for each $n \in Y \exists t_n \in B$ such that $t_n \geq \sigma_n$. Suppose that the construction of $G_{\delta+1}^l$ uses $A_{\beta, \delta} = \bigcup_{n \in \omega} \Theta_n^{\beta, \delta}$. Selecting one $n_* \in Y$, we see that, since $\Theta_{n_*}^{\beta, \delta} \subseteq \sigma_{n_*}$, $\sigma_{n_*} > \beta$ and hence $t_{n_*} \in A_{\beta, \delta}$. Therefore there exists M such that, for all $n \geq M$, $t_{n_*} \in \Theta_n^{\beta, \delta}$. But then, for $n \in Y$ with $n \geq M$, $t_n \geq \sigma_n \supseteq \Theta_n^{\beta, \delta}$, so $t_{n_*} \leq t_n$ and thus $t_{n_*} \leq_T t_n$. By the construction in Case 4 and by 4(c), $g_B(w_{\sigma_n, m_n}) = v_{\sigma_n, m_n}^1$ for $n \in Y$, $n \geq M$. Moreover, there is no conflict between the last two conditions because, by construction, if $\delta \in E_1$ and $\sigma \in E_0$, then $w_{\sigma, m} \in W_\delta[\Theta_n^{\beta, \delta}]$ if and only if $\Theta_n^{\beta, \delta} = \Theta_m^\sigma$, but the elements of $\{\Theta_m^\sigma : m \in \omega\}$ are disjoint and the $\Theta_n^{\beta, \delta}$ form a chain under \subseteq .

It remains to extend \tilde{g}_B to f_μ^0 by defining $f_\mu^0(z_{\delta,n}^0)$ for $\tau \leq \delta \leq \mu$, $n \in \omega$. This is possible by observation (IV.3) because of the construction of \tilde{g}_B .

(V) We will define the projections $\pi_{\nu, \mu}^l$ by induction on μ and then verify the conditions to be in standard form (see Section 1 or [2, Definition 1.9(ii), p.

257]). We refer to the cases of the construction in part (III). In Case 1, we take unions. In Case 2, for $\nu < \mu + 1$ we let $\pi_{\nu, \mu+1}^l$ be the extension of $\pi_{\nu, \mu}^l$ which sends each $x_{\mu, j}^l$ to 0. (Here, $\pi_{\mu, \mu}^l$ is the identity.) In Case 3, for $\nu \leq \sigma$ we let $\pi_{\nu, \sigma+1}^l$ be the extension of $\pi_{\nu, \sigma}^l$ which sends each $u_{\sigma, n}^l$ and each $v_{\sigma, n}^l$ to 0.

Finally, for Case 4, we use the notation of that case. We define $\pi_{\nu, \delta+1}^0(z_{\delta, n}^0) = -\sum_{j=n}^m d_{n, j} k_{\delta, j}^0$ where m is maximal such that $\gamma_m + 2 \leq \nu$ and $d_{n, j} = \prod_{i=n}^{j-1} p_i$ (and $d_{n, 0} = 1$). (Compare [2, pp. 249f].) The definition of $\pi_{\nu, \delta+1}^1$ is similar, replacing $k_{\delta, j}^0$ by $f_{\Theta_j}(k_{\delta, j}^0)$. Let $Y_{\delta}^l = \{z_{\delta, n}^l : n \in \omega\}$. Then we can easily verify the conditions of [2, Definition 1.9(ii), p. 257] using the information in the proof of (III.2) about the form of $k_{\delta, j}^0$.

This completes the proof of Theorem 7.

3 A NON-STRUCTURE THEOREM

Our goal is to generalize the construction in the previous section to prove:

Theorem 8. *Assume \diamond . There exists an \aleph_1 -separable group G^0 and for each bounded tree T_1 an \aleph_1 -separable group G^{T_1} which is T_1 -equivalent to G^0 but not isomorphic to G^0 . Moreover, all the groups are of cardinality \aleph_1 and in standard form.*

Proof. We assume familiarity with the previous proof and outline the modifications, in layers of increasing detail.

(VI) Fix a stationary subset E of ω_1 consisting of limit ordinals (> 0) and such that E is the disjoint union of two subsets E_0 and E_1 such that cardinality $\diamond(E_0)$ and $\diamond(E_1)$ hold. ($\diamond(E_0)$ is not essential, but convenient.)

We need only consider bounded trees T on ω_1 such that if $\nu <_T \mu$ (in the tree ordering), then $\nu < \mu$ (as ordinals). For each $\delta \in E_1$ (resp. $\sigma \in E_0$), diamond will give us a “prediction” $T_{\delta} = \langle \delta, <_{\delta} \rangle$ (resp. T_{σ}) of the restriction of a bounded tree to δ (resp. σ). If $\mu < \delta$, we write $T_{\delta} \upharpoonright \mu$ for $\langle \mu, <_{\delta} \cap (\mu \times \mu) \rangle$.

By induction on $\delta \in \{0\} \cup E$ we will define the following data:

1. continuous chains $\{G_{\nu}^{\delta} : \nu \leq \delta + 1\}$ of countable free groups such that, for all $\nu < \mu \leq \delta + 1$, $G_{\mu}^{\delta}/G_{\nu}^{\delta}$ is free if $\nu \notin E_1$, and if $\nu \in E_1$, then $G_{\nu+1}^{\delta}/G_{\nu}^{\delta}$ has rank at most 1.
2. projections $\pi_{\nu, \mu}^{\delta} : G_{\mu}^{\delta} \rightarrow G_{\nu}^{\delta}$ for $\nu \leq \mu \leq \delta + 1$ and $\nu \notin E_1 \cup \{\sigma + 1 : \sigma \in E_0\}$ such that: for $\nu \leq \mu < \rho$, $\pi_{\nu, \mu}^{\delta} \subseteq \pi_{\nu, \rho}^{\delta}$; and for $\tau < \nu \leq \mu$, $\pi_{\tau, \nu}^{\delta} \circ \pi_{\nu, \mu}^{\delta} = \pi_{\tau, \mu}^{\delta}$;
3. for each $\delta \in E$ and each $\nu \leq \delta$ an isomorphism $f_{\nu}^{\delta} : G_{\nu+1}^0 \rightarrow G_{\nu+1}^{\delta}$ satisfying:

$$\text{if } \nu_1 <_{\delta} \nu_2, \text{ then } f_{\nu_2}^{\delta} \upharpoonright G_{\nu_1+1}^0 = f_{\nu_1}^{\delta}.$$

Moreover, we require that if $\delta < \delta'$ are elements of E such that $T_{\delta} = T_{\delta'} \upharpoonright \delta$, then $G_{\nu}^{\delta} = G_{\nu}^{\delta'}$ for $\nu \leq \delta + 1$; $\pi_{\nu, \mu}^{\delta} = \pi_{\nu, \mu}^{\delta'}$ for $\nu \leq \mu \leq \delta + 1$; and $f_{\nu}^{\delta'} = f_{\nu}^{\delta}$ for $\nu \leq \delta$.

Define $G^0 = \bigcup_{\nu < \omega_1} G_{\nu}^0$ and for each bounded tree T on ω_1 let $G^T = \bigcup \{G_{\nu}^{\delta} : T_{\delta} = T \upharpoonright \delta, \nu \leq \delta + 1\}$. As before, given T_1 we can choose T so that G^0 and G^T are T_1 -equivalent.

We indicate how to modify the previous construction so that G^0 and G^T are not isomorphic. Our construction will be such that when $\alpha = \mu + 1$ where $\mu \notin E$, then

$$(*) \quad G_\alpha^0 = G_\mu^0 \oplus \mathbb{Z}x_{\mu,0}^0 \oplus \mathbb{Z}x_{\mu,1}^0$$

and

$$(**) \quad G_\alpha^\delta = G_\mu^\delta \oplus \mathbb{Z}x_{\mu,0}^1 \oplus \mathbb{Z}x_{\mu,1}^1$$

for $\delta \in E$, $\alpha < \delta$. When $\alpha = \sigma + 1$ where $\sigma \in E_0$, then

$$(***) \quad G_\alpha^0 = G_\sigma^0 \oplus \bigoplus_{n \in \omega} \mathbb{Z}u_{\sigma,n}^0 \oplus \mathbb{Z}v_{\sigma,n}^0$$

and

$$(***) \quad G_\alpha^\delta = G_\sigma^\delta \oplus \bigoplus_{n \in \omega} \mathbb{Z}u_{\sigma,n}^1 \oplus \mathbb{Z}v_{\sigma,n}^1$$

for $\delta \in E$, $\sigma < \delta$.

We define

$$w_{\sigma,n} = 2u_{\sigma,n+1}^0 - u_{\sigma,n}^0.$$

In order to carry out the inductive construction we will define in addition:

4. for $\delta \in E$ and $\alpha \leq \delta + 1$, subsets $W_\alpha^\delta[\Theta]$ of G_α^0 for every nonempty finite subset Θ of α which is an antichain in T_δ , satisfying:

(a) for all $\alpha < \beta$, $W_\alpha^\delta[\Theta] \subseteq W_\beta^\delta[\Theta]$;

(b) every element of $W_\alpha^\delta[\Theta]$ is of the form $w_{\sigma,n}$ for some $n \in \omega$ and some $\sigma \in E_0$ such that $T_\delta \upharpoonright \sigma = T_\sigma$.

The function f_α^δ will be required to satisfy (as before):

(c) for all $\mu \leq \alpha$, $j \in \{0, 1\}$, $f_\alpha^\delta(x_{\mu,j}^0) = x_{\mu,j}^1$; moreover if $w_{\sigma,n} \in W_{\alpha+1}^\delta[\Theta]$ and $\Theta \cap \{\nu : \nu \leq_\delta \alpha\} \neq \emptyset$, then $f_\alpha^\delta(w_{\sigma,n}) = v_{\sigma,n}^1$.

Moreover, in order to carry out the inductive construction we will also require the following for all $\delta \in E$, $\alpha \leq \delta$:

(d) if $\sigma \in E_0$ with $\sigma \leq \alpha + 1$ and $T_\delta \upharpoonright \sigma \neq T_\sigma$, then $f_\alpha^\delta(u_{\sigma,n}^0) = u_{\sigma,n}^1$ for all $n \in \omega$;

(e) for all pairs β_1, β_2 with $\sup\{t : t <_\delta \alpha\} \leq \beta_1 < \beta_2 \leq \alpha$, it is the case for almost all $n \in \omega$ that for all $w_{\sigma,m} \in W_{\alpha+1}^\delta[\Theta_n^{\beta_1, \beta_2}]$ we have $f_\alpha^\delta(w_{\sigma,m}) = v_{\sigma,m}^1$.

(The notation $\Theta_n^{\beta_1, \beta_2}$ is defined before (II.2).)

$\diamond(E_0)$ gives us for each $\sigma \in E_0$ a "prediction" $\Upsilon(\sigma) = \langle \Theta_n^\sigma : n \in \omega \rangle$ of an ω -sequence of finite subsets of T_σ such that $\bigcup_{n \in \omega} \Theta_n^\sigma$ is an antichain in T_σ . The proof that G^0 and G^T are not isomorphic will then work as before.

(VII) The next step is to describe in detail the inductive construction of the data satisfying the properties given above. Our construction is by induction on the elements of E . At stage $\delta \in E$ we will define G_α^0 and G_α^δ for any $\alpha \leq \delta + 1$ for which they are not already defined. We will have already defined G_ν^0 for $\nu \leq \sup\{\delta' + 1 : \delta' \in E, \delta' < \delta\}$. By following the prescriptions in (*) and (***) , we can assume that G_ν^0 is defined for all $\nu \leq \delta$.

Let $\gamma = \sup\{\delta' + 1 : \delta' \in E \cap \delta, T_\delta \upharpoonright \delta' = T_{\delta'}\}$. Then we need to define G_α^δ for $\gamma < \alpha \leq \delta + 1$. We need to do this in such a way that we are able to define the partial isomorphisms f_α^δ . We shall leave the details of the latter to the next section and describe the construction of the groups here. There are two cases to consider.

Case 1: $\gamma = \delta \in E$. Then G_δ^δ is already defined. If $\delta \in E_0$, follow the prescription in (***) and (****). If $\delta \in E_1$, $\diamond(E_1)$ gives us an isomorphism $h : G_\delta^0 \rightarrow G_\delta^\delta$; the construction of $G_{\delta+1}^0$ and G_δ^δ is essentially the same as in the previous theorem (Case 4 of (III)); in particular, if (II.2) holds, we use an antichain $A_{\beta, \delta}^\delta = \{t : t \text{ is } <_\delta\text{-minimal in } \delta \setminus \beta\}$; $G_{\delta+1}^0$ is generated by $G_\delta^0 \cup \{z_{\delta, n}^0 : n \in \omega\}$ subject to relations $p_n z_{\delta, n+1}^0 = z_{\delta, n}^0 + k_{\delta, n}^0$ (which keep h from extending) and $G_{\delta+1}^\delta$ is generated by $G_\delta^\delta \cup \{z_{\delta, n}^\delta : n \in \omega\}$ subject to relations $p_n z_{\delta, n+1}^\delta = z_{\delta, n}^\delta + k_{\delta, n}^\delta$ (where $k_{\delta, n}^\delta = f_{\Theta_n}^\delta(k_{\delta, n}^0)$).

For the purpose of later stages of the construction we also define, for any $\delta_1 > \delta$ such that $\delta_1 \in E$ and $T_{\delta_1} \upharpoonright \delta \neq T_\delta$, elements $k_{\delta, n}^{\delta_1} \in G_{\delta_1}^{\delta_1}$. We know that $k_{\delta, n}^0$ has the form $x_{\mu_n, j_n}^0 \pm \xi_{\delta, n}$ where $\xi_{\delta, n}$ is either 0, $x_{\sigma, j}^0$, or $w_{\sigma, j}$ for some σ, j (cf. (IV.1)). In case $\xi_{\delta, n}$ is 0, let $k_{\delta, n}^{\delta_1} = x_{\mu_n, j_n}^1$; in case $\xi_{\delta, n} = x_{\sigma, j}^0$, let $k_{\delta, n}^{\delta_1} = x_{\mu_n, j_n}^1 \pm x_{\sigma, j}^1$. Finally, if $\xi_{\delta, n} = w_{\sigma, j}$, let $k_{\delta, n}^{\delta_1} = x_{\mu_n, j_n}^1 \pm \xi'_{\delta, n}$ where

$$\xi'_{\delta, n} = \begin{cases} w_{\sigma, j}^1 & \text{if } T_{\delta_1} \upharpoonright \sigma \neq T_\sigma, \\ v_{\sigma, j}^1 & \text{if } T_{\delta_1} \upharpoonright \sigma = T_\sigma, \end{cases}$$

and $w_{\sigma, j}^1 = 2u_{\sigma, j+1}^1 - u_{\sigma, j}^1$. We will be able to show (in the next section) the following:

$$(VII.1) \text{ for any branch } B \text{ in } T_{\delta_1} \upharpoonright \delta \text{ with } \delta = \sup\{t+1 : t \in B\}, \\ g_B = \bigcup \{f_\alpha^{\delta_1} : \alpha \in B\} \text{ is such that, for almost all } n, g_B(k_{\delta, n}^0) = k_{\delta, n}^{\delta_1}.$$

(This is evidence of what, in view of (IV.3), will enable us to extend functions.)

Case 2: $\gamma < \delta$. We need to define G_α^δ for $\gamma + 1 \leq \alpha \leq \delta + 1$ by induction on α . If we have defined G_α^δ for $\alpha \leq \rho < \delta$, and ρ does not belong to E_1 , we follow the prescription in (**) or (****). If $\rho \in E_1$, then $T_\delta \upharpoonright \rho \neq T_\rho$ (by definition of γ). By induction $G_{\rho+1}^0$ is constructed as in Case 1 and we have $k_{\rho, n}^\delta$ as there (with δ playing the role of δ_1 and ρ playing the role of δ). In particular, $G_{\rho+1}^0$ is generated by $G_\rho^0 \cup \{z_{\rho, n}^0 : n \in \omega\}$ subject to relations $p_n z_{\rho, n+1}^0 = z_{\rho, n}^0 + k_{\rho, n}^0$. We define $G_{\rho+1}^\delta$ to be generated by $G_\rho^\delta \cup \{z_{\rho, n}^\delta : n \in \omega\}$ subject to relations $p_n z_{\rho, n+1}^\delta = z_{\rho, n}^\delta + k_{\rho, n}^\delta$. Finally, we define $G_{\delta+1}^\delta$ as in Case 1.

The definition of the $W_\alpha^\delta[\Theta]$ will be as in (III); specifically, $W_{\alpha+1}^\delta[\Theta] = W_\alpha^\delta[\Theta]$ unless $\alpha = \sigma \in E_0$, $T_\delta \upharpoonright \sigma = T_\sigma$ and $\Theta = \Theta_n^\sigma$ for some n , in which case $W_{\sigma+1}^\delta[\Theta] = W_\sigma^\delta[\Theta] \cup \{w_{\sigma, n}\}$.

(VIII) We have defined the groups and the sets $W_\alpha^\delta[\Theta]$; the last step is to show that the partial isomorphisms f_ν^δ can be defined satisfying the conditions in 4.

First let us verify (VII.1). Let δ and δ_1 be as in Case 1 of (VII) and suppose B is a branch in $T_{\delta_1} \upharpoonright \delta$ with $\delta = \sup\{t + 1 : t \in B\}$. Then g_B is an

isomorphism: $G_\delta^0 \rightarrow G_\delta^{\delta_1}$ and we want to show that $g_B(k_{\delta,n}^0) = k_{\delta,n}^{\delta_1}$ for almost all n . Recall that $k_{\delta,n}^0$ has the form $x_{\mu_n, j_n}^0 \pm \xi_{\delta,n}$ where $\xi_{\delta,n}$ is either 0, $x_{\sigma,j}^0$, or $w_{\sigma,j}$ for some σ, j ; the only case we need to worry about is when $\xi_{\delta,n} = w_{\sigma,j}$.

Let $\mu = \sup\{\alpha < \delta : T_\delta \upharpoonright \alpha = T_{\delta_1} \upharpoonright \alpha\}$; so $\mu < \delta$ and $G_\alpha^{\delta_1} = G_\alpha^\delta$ for $\alpha \leq \mu$. Suppose that $G_{\delta+1}^0$ and $G_{\delta+1}^{\delta_1}$ are defined using $A_{\beta,\delta}^\delta = \bigcup_{n \in \omega} \Theta_n^{\beta,\delta}$ as in Case 1 of (VII) and Case 4 of (III). We consider several cases. First, suppose that there exists $t \in A_{\beta,\delta}^\delta$ with $t \geq \mu$. Then for almost all n , $t \in \Theta_n^{\beta,\delta}$ and thus if $w_{\sigma,j} \in W_\delta^\delta[\Theta_n^{\beta,\delta}]$, then $\sigma > t \geq \mu$; hence $T_{\delta_1} \upharpoonright \sigma \neq T_\delta \upharpoonright \sigma$ and it follows from 4(d) that $g_B(k_{\delta,n}^0) = k_{\delta,n}^{\delta_1}$. If this case does not hold, then $A_{\beta,\delta}^\delta \subseteq \mu$ so $A_{\beta,\delta}^\delta = A_{\beta,\mu}^\delta$ is an antichain in $T_{\delta_1} \upharpoonright \mu = T_\delta \upharpoonright \mu$. If there exists $t \in B$ with $\beta \leq t < \mu$, then there exists $t \in B$ with $t \in A_{\beta,\delta}^\delta$ and hence $t \in \Theta_n^{\beta,\delta}$ for almost all n ; it follows easily that, for almost all n , $g_B(k_{\delta,n}^0) = k_{\delta,n}^{\delta_1}$ (considering separately the cases when $\sigma \leq \mu$ and $\sigma > \mu$). In the remaining case, if $\alpha = \inf\{t \in B : t \geq \beta\}$, then $\alpha \geq \mu$ so we have $\sup\{t : t <_{\delta_1} \alpha\} \leq \beta < \mu \leq \alpha$ and we have the desired conclusion by 4(e)—again distinguishing between the cases when $\sigma \leq \mu$ and $\sigma > \mu$. This completes the proof of (VII.1).

Now we need to verify the analog of (III.1). Letting δ and γ be as in (VII), we need to define f_α^δ for $\gamma \leq \alpha \leq \delta$. Fix α and let $B = \{t < \gamma : t <_\delta \alpha\}$ and $g_B = \bigcup\{f_t^\delta : t \in B\}$. We can suppose that α is $<_\delta$ -minimal among elements of $\{\beta : \gamma \leq \beta \leq \alpha\}$.

We will first define an extension of g_B to a partial isomorphism \tilde{g}_B whose domain is

$$\text{dom}(g_B) + \left\langle \begin{array}{l} \{x_{\nu,j}^0 : \nu \leq \alpha, j = 0, 1\} \\ \cup \{u_{\sigma,n}^0 : \sigma \in E_0 \cap (\alpha + 1), n \in \omega\} \\ \cup \{v_{\sigma,n}^0 : \sigma \in E_0 \cap (\alpha + 1), n \in \omega\} \end{array} \right\rangle.$$

Using an enumeration in an ω -sequence of $Y_0 \cup Y_1$ where

$$Y_0 = \{\sigma \in E_0 : \sup B \leq \sigma < \gamma \text{ and } T_\delta \upharpoonright \sigma = T_\sigma\}$$

and

$$Y_1 = \{(\beta_1, \beta_2) : \sup B \leq \beta_1 < \beta_2 \leq \alpha\}$$

we can define \tilde{g}_B such that

- (c') for all $\nu \leq \alpha, j \in \{0, 1\}$, $\tilde{g}_B(x_{\nu,j}^0) = x_{\nu,j}^1$; moreover, if $w_{\sigma,n} \in W_{\gamma+1}^\delta[\Theta]$ and $\Theta \cap B \neq \emptyset$, then $\tilde{g}_B(w_{\sigma,n}) = v_{\sigma,n}^1$;
- (d') if $\sigma \in E_0 \cap \alpha + 2$, then $\tilde{g}_B(u_{\sigma,n}^0) = u_{\sigma,n}^1$ for almost all n , and if $T_\delta \upharpoonright \sigma \neq T_\sigma$, then $\tilde{g}_B(u_{\sigma,n}^0) = u_{\sigma,n}^1$ for all n ; and
- (e') for all pairs β_1, β_2 with $\sup B \leq \beta_1 < \beta_2 \leq \alpha$, it is the case for almost all $n \in \omega$ that for all $w_{\sigma,m} \in W_{\alpha+1}^\delta[\Theta_n^{\beta_1, \beta_2}]$ we have $\tilde{g}_B(w_{\sigma,m}) = v_{\sigma,m}^1$.

Now $\tilde{g}_B(k_{\rho,n}^0)$ is defined for all $\rho \in E_1$ with $\rho \leq \alpha$. We need to define $f_\alpha^\delta(z_{\rho,n}^\delta)$ for all such $\rho \geq \sup B$. In view of (IV.3), we can do this provided that $\tilde{g}_B(k_{\rho,n}^0) = k_{\rho,n}^\delta$ for almost all $n \in \omega$. We consider separately the cases:

$T_\delta \upharpoonright \rho = T_\rho$; and $T_\delta \upharpoonright \rho \neq T_\rho$. The first case is as in (IV); the last is as in the proof of (VII.1) (with δ playing the role of δ_1 , ρ playing the role of δ and using (d') and (e')).

This completes the proof of Theorem 8.

REFERENCES

1. P. Eklof, *The structure of ω_1 -separable groups*, Trans. Amer. Math. Soc. **279** (1983), 497–523.
2. P. C. Eklof and A. H. Mekler, *Almost free modules*, North-Holland, 1990.
3. T. Hyttinen, *Model theory for infinite quantifier languages*, Fund. Math. **134** (1990), 125–142.
4. T. Hyttinen and H. Tuuri, *Constructing strongly equivalent nonisomorphic models for unstable theories*, Ann. Pure Appl. Logic **52** (1991), 203–248.
5. T. Jech, *Set theory*, Academic Press, 1978.
6. M. Karttunen, *Model theory for infinitely deep languages*, Ann. Acad. Sci. Fenn. Ser. A I Math. Dissertationes **50** (1984), 897–908.
7. A. H. Mekler, *Proper forcing and abelian groups*, Abelian Group Theory, Lecture Notes in Math., vol. 1006, Springer-Verlag, 1983, pp. 285–303.
8. ———, *The structure of groups that are almost the direct sum of countable abelian groups*, Trans. Amer. Math. Soc. **303** (1987), 145–160.
9. A. Mekler, S. Shelah and J. Väänänen, *The Enrenfeucht-Fraïssé game of length ω_1* , Trans. Amer. Math. Soc. **339** (1993), 567–580.
10. A. Mekler and J. Oikkonen, *Abelian p -groups with no invariants*, J. Pure Appl. Algebra **87** (1993), 51–59.
11. J. Oikkonen, *Enrenfeucht-Fraïssé-games and nonstructure theorems*, preprint.
12. J. Väänänen, *Games and trees in infinitary logic: a survey*, Quantifiers (M. Krynicki, M. Mostowski and L. Szczerba, eds.), Kluwer Acad. Publ. (to appear).

(P. Eklof and H. Foreman) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, IRVINE, CALIFORNIA 92717

E-mail address: peklof@math.uci.edu

E-mail address: mforeman@math.uci.edu

(S. Shelah) DEPARTMENT OF MATHEMATICS, HEBREW UNIVERSITY, JERUSALEM, ISRAEL

E-mail address: shelah@math.huji.ac.il

(S. Shelah) DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW BRUNSWICK, NEW JERSEY 08903