

SUCCESSOR OF SINGULARS: COMBINATORICS AND NOT COLLAPSING CARDINALS $\leq \kappa$ IN $(< \kappa)$ -SUPPORT ITERATIONS

BY

SAHARON SHELAH*

*Institute of Mathematics, The Hebrew University of Jerusalem
Givat Ram, 91904 Jerusalem, Israel*

and

*Department of Mathematics, Rutgers University
New Brunswick, NJ 08854, USA*

and

*Mathematics Department, University of Wisconsin – Madison
Madison, WI 53706, USA*

e-mail: shelah@math.huji.ac.il

ABSTRACT

On the one hand, we deal with $(< \kappa)$ -supported iterated forcing notions which are $(\mathcal{E}_0, \mathcal{E}_1)$ -complete, bearing in mind problems on Whitehead groups, uniformizations and the general problem. We deal mainly with the case of a successor of the singular cardinal. This continues [Sh 587]. On the other hand, we deal with complimentary ZFC combinatorial results.

ANNOTATED CONTENTS

- §1. GCH implies for successor of singular no stationary S has uniformization 128
 [For λ strong limit singular, for stationary $S \subseteq S_{cf(\lambda)}^{\lambda^+}$ we prove strong negation of uniformization for some S -ladder system and even weak versions of diamond. E.g., if λ is singular strong limit and $2^\lambda = \lambda^+$, then there are $\gamma_i^\delta < \delta$ increasing in $i < cf(\lambda)$ with limit δ for each $\delta \in S$ such that for every $f: \lambda^+ \rightarrow \alpha^* < \lambda$ for stationarily many $\delta \in S$, for every i we have $f(\gamma_{2i}^\delta) = f(\gamma_{2i+1}^\delta)$.]
- §2. Case C: Forcing for successor of singulars 131
 [Let λ be strong limit singular $\kappa = \lambda^+ = 2^\lambda$, $S \subseteq S_{cf(\lambda)}^\kappa$ stationary not reflecting. We present the consistency of a forcing axiom implying, e.g.: if h_δ is a function from A_δ to θ , $A_\delta \subseteq \delta = \sup(A_\delta)$,

* I would like to thank Alice Leonhardt for the beautiful typing.
 This research was supported by The Israel Science Foundation founded by the Israel Academy of Sciences and Humanities. Publication 667.
 Received November 16, 1998 and in revised form March 5, 2001

$\text{otp}(A_\delta) = \text{cf}(\lambda), \theta < \lambda$, then for some $h: \kappa \rightarrow \theta$ for every $\delta \in S$ we have $h_\delta \subseteq^* h$.]

- §3. κ^+ -c.c. and κ^+ -pic 141
 [In the forcing axioms we would like to allow forcing notions of cardinality $> \kappa$; for this we use a suitable chain condition (allowed here and in [Sh 587]). This sheds more light on the strongly inaccessible case and we comment on this (and forcing against cases of diamonds).]
- §4. Existence of non-free Whitehead (and $\text{Ext}(G, \mathbb{Z}) = 0$) abelian groups in successor of singulars 150
 [We use the information on the existence of weak version of the diamond for $S \subseteq S_{\text{cf}(\lambda)}^{\lambda^+}$, λ strong limit singular with $2^\lambda = \lambda^+$, to prove that there are some abelian groups with special properties (from reasonable assumptions). We also get more combinatorial principles on $\lambda = \mu^+$, $\mu > \text{cf}(\mu)$ (even if just $\lambda = \lambda^{2^\sigma}$).]
- References 155

§1. GCH implies for successor of singular no stationary S has uniformization

We show that a major improvement in [Sh 587] over [Sh 186] for inaccessible (every ladder on S has uniformization rather than some ladder on S) cannot be done for successor of singulars. This is continued in §4.

1.1 FACT: Assume

- (a) λ is strong limit singular with $2^\lambda = \lambda^+$, let $\text{cf}(\lambda) = \sigma$
- (b) $S \subseteq \{\delta < \lambda^+ : \text{cf}(\delta) = \sigma\}$ is stationary.

Then we can find $\langle \gamma_i^\delta : i < \sigma > : \delta \in S \rangle$ such that

- (α) γ_i^δ is increasing (with i) with limit δ
- (β) if $\mu < \lambda$ and $f: \lambda^+ \rightarrow \mu$ then the following set is stationary:
 $\{\delta \in S : f(\gamma_{2i}^\delta) = f(\gamma_{2i+1}^\delta) \text{ for every } i < \sigma\}$.

Moreover

- (β)⁺ if $f_i: \lambda^+ \rightarrow \mu_i, \mu_i < \lambda$ for $i < \sigma$ then the following set is stationary:
 $\{\delta \in S : f_i(\gamma_{2i}^\delta) = f_i(\gamma_{2i+1}^\delta) \text{ for every } i < \sigma\}$.

Proof: This will prove 1.2, too. We first concentrate on (α) + (β) only.

Let $\lambda = \sum_{i < \sigma} \lambda_i$, λ_i a cardinal increasing continuous with i , $\lambda_{i+1} > 2^{\lambda_i}$, $\lambda_0 > 2^\sigma$. For $\alpha < \lambda^+$, let $\alpha = \bigcup_{i < \sigma} a_{\alpha, i}$ such that $|a_{\alpha, i}| \leq \lambda_i$. Without loss of generality $\delta \in S \Rightarrow \delta$ divisible by λ^ω (ordinal exponentiation). For $\delta \in S$ let $\langle \beta_i^\delta : i < \sigma \rangle$ be increasing continuous with limit δ , β_i^δ divisible by λ and > 0 . For $\delta \in S$ let $\langle b_i^\delta : i < \sigma \rangle$ be such that: $b_i^\delta \subseteq \beta_i^\delta$, $|b_i^\delta| \leq \lambda_i$, b_i^δ is increasing

continuous with i and $\delta = \bigcup_{i < \sigma} b_i^\delta$ (e.g., we can let $b_i^\delta = \bigcup_{j_1, j_2 < i} a_{\beta_{j_1, j_2}^\delta} \cup \lambda_i$). We further demand $\lambda_i \subseteq b_i^\delta \cap \lambda$. Let $\langle f_\alpha^* : \alpha < \lambda^+ \rangle$ list the two-place functions with domain an ordinal $< \lambda^+$ and range $\subseteq \lambda^+$. Let $S = \bigcup_{\mu < \lambda} S_\mu$, with each S_μ stationary and $\langle S_\mu : \mu < \lambda \rangle$ pairwise disjoint. We now fix $\mu < \lambda$ and will choose $\bar{\gamma}^\delta = \langle \gamma_i^\delta : i < \sigma \rangle$ for $\delta \in S_\mu$ such that clause (α) holds and clause (β) holds (that is, for every $f: \lambda^+ \rightarrow \mu$ for stationary many $\delta \in S_\mu$ the conclusion of clause (β) holds); this clearly suffices.

Now for $\delta \in S_\mu$ and $i < j < \sigma$ we can choose $\zeta_{i,j,\varepsilon}^\delta$ (for $\varepsilon < \lambda_j$) (really here we use just $\varepsilon = 0, 1$) such that:

- (A) $\langle \zeta_{i,j,\varepsilon}^\delta : \varepsilon < \lambda_j \rangle$ is a strictly increasing sequence of ordinals,
- (B) $\beta_i^\delta < \zeta_{i,j,\varepsilon}^\delta < \beta_{i+1}^\delta$ (can even demand $\zeta_{i,j,\varepsilon}^\delta < \beta_i^\delta + \lambda$),
- (C) $\zeta_{i,j,\varepsilon}^\delta \notin \{ \zeta_{i_1, j_1, \varepsilon_1}^\delta : j_1 < j, \varepsilon_1 < \lambda_{j_1} \text{ (and } i_1 < \sigma, \text{ really only } i_1 = i \text{ matters)} \}$,
- (D) for every $\alpha_1, \alpha_2 \in b_j^\delta$, the sequence $\langle \text{Min}\{\lambda_j, f_{\alpha_1}^*(\alpha_2, \zeta_{i,j,\varepsilon}^\delta)\} : \varepsilon < \lambda_j \rangle$ is constant, i.e., one of the following occurs:

- (α) $\varepsilon < \lambda_j \Rightarrow (\alpha_2, \zeta_{i,j,\varepsilon}^\delta) \notin \text{Dom}(f_{\alpha_1}^*)$,
- (β) $\varepsilon < \lambda_j \Rightarrow f_{\alpha_1}^*(\alpha_2, \zeta_{i,j,\varepsilon}^\delta) = f_{\alpha_1}^*(\alpha_2, \zeta_{i,j,0}^\delta)$, well defined,
- (γ) $\varepsilon < \lambda_j \Rightarrow f_{\alpha_1}^*(\alpha_2, \zeta_{i,j,\varepsilon}^\delta) \geq \lambda_j$, well defined.

For each $i < j < \sigma$ we use “ λ is strong limit $> \lambda_j \geq \sum_{j_1 < j} \lambda_{j_1} + \sigma$ ”.

Let $G = \{g : g \text{ a function from } \sigma \text{ to } \sigma \text{ such that } (\forall i < \sigma)(i < g(i))\}$. For each function $g \in G$ we try $\bar{\gamma}^{g,\delta} = \langle \zeta_{i,g(i),0}^\delta, \zeta_{i,g(i),1}^\delta : i < \sigma \rangle$, i.e., $\langle \zeta_{2i}^{g,\delta}, \zeta_{2i+1}^{g,\delta} \rangle = \langle \gamma_{i,g(i),0}^\delta, \gamma_{i,g(i),1}^\delta \rangle$.

Now we ask for each $g \in G$:

Question $_g^\mu$: Does $\langle \bar{\gamma}^{g,\delta} : \delta \in S_\mu \rangle$ satisfy

$$(\forall f \in {}^{\lambda^+}\mu)(\exists^{\text{stat}} \delta \in S_\mu) \left(\bigwedge_{i < \sigma} f(\gamma_{2i}^{g,\delta}) = f(\gamma_{2i+1}^{g,\delta}) \right)?$$

If for some $g \in G$ the answer is yes, we are done. Assume not; so for each $g \in G$ we can find $f_g: \lambda^+ \rightarrow \mu$ and a club E_g of λ^+ such that

$$\delta \in S_\mu \cap E_g \Rightarrow (\exists i < \sigma) (f_g(\gamma_{2i}^{g,\delta}) \neq f_g(\gamma_{2i+1}^{g,\delta})),$$

which means

$$\delta \in S_\mu \cap E_g \Rightarrow (\exists i < \sigma) [f_g(\zeta_{i,g(i),0}^\delta) \neq f_g(\zeta_{i,g(i),1}^\delta)].$$

Let $G = \{g_\varepsilon : \varepsilon < 2^\sigma\}$, so we can find a 2-place function f^* from λ^+ to μ satisfying $f^*(\varepsilon, \alpha) = f_{g_\varepsilon}(\alpha)$ when $\varepsilon < 2^\sigma$, $\alpha < \lambda^+$. Hence for each $\alpha < \lambda^+$ there is $\gamma[\alpha] < \lambda^+$ such that $f^* \upharpoonright \alpha = f_{\gamma[\alpha]}^*$.

Let $E^* = \bigcap_{\varepsilon < 2^\sigma} E_{g_\varepsilon} \cap \{\delta < \lambda^+ : \text{for every } \alpha < \delta \text{ we have } \gamma[\alpha] < \delta\}$. Clearly it is a club of λ^+ , hence we can find $\delta \in S_\mu \cap E^*$. Now $\beta_{i+1}^\delta < \delta$ hence $\gamma[\beta_{i+1}^\delta] < \delta$ (as $\delta \in E^*$), but $\delta = \bigcup_{i < \sigma} b_i^\delta$ hence for some $j < \sigma$, $\gamma[\beta_{i+1}^\delta] \in b_j^\delta$; as b_j^δ increases with j we can define a function $h: \sigma \rightarrow \sigma$ by $h(i) = \text{Min}\{j : j > i + 1 \text{ and } \mu < \lambda_j \text{ and } \gamma[\beta_{i+1}^\delta] \in b_j^\delta\}$. So $h \in G$, hence for some $\varepsilon(*) < 2^\sigma$ we have $h = g_{\varepsilon(*)}$. Now looking at the choice of $\zeta_{i,h(i),0}^\delta, \zeta_{i,h(i),1}^\delta$ we know (remember $2^\sigma < \lambda_0 \subseteq b_j^\delta$ and $\mu < \lambda_{h(i)}$)

$$\begin{aligned} (\forall \varepsilon < 2^\sigma)(\forall \alpha \in b_{h(i)}^\delta)[\text{Rang}(f_\alpha^*) \subseteq \mu \ \& \ \text{Dom}(f_\alpha^*) \supseteq \beta_{i+1}^\delta \rightarrow f_\alpha^*(\varepsilon, \zeta_{i,h(i),0}^\delta) \\ &= f_\alpha^*(\varepsilon, \zeta_{i,h(i),1}^\delta)]. \end{aligned}$$

In particular this holds for $\varepsilon = \varepsilon(*), \alpha = \gamma[\beta_{i+1}^\delta]$, so we get

$$f_{\gamma[\beta_{i+1}^\delta]}^*(\varepsilon(*), \zeta_{i,h(i),0}^\delta) = f_{\gamma[\beta_{i+1}^\delta]}^*(\varepsilon(*), \zeta_{i,h(i),1}^\delta).$$

By the choice of f^* and of $\gamma[\beta_{i+1}^\delta]$ this means

$$f_{g_{\varepsilon(*)}}(\zeta_{i,h(i),0}^\delta) = f_{g_{\varepsilon(*)}}(\zeta_{i,h(i),1}^\delta);$$

but $h = g_{\varepsilon(*)}$, and the above equality means $f_{g_{\varepsilon(*)}}^*(\gamma_{2i}^{g_{\varepsilon(*)},\delta}) = f_{g_{\varepsilon(*)}}^*(\gamma_{2i+1}^{g_{\varepsilon(*)},\delta})$, and this holds for every $i < \sigma$, and $\delta \in E^* \Rightarrow \delta \in E_{g_{\varepsilon(*)}}$, so we get a contradiction to the choice of $(f_{g_{\varepsilon(*)}}, E_{\varepsilon(*)})$. So we have finished proving $(\alpha) + (\beta)$.

How do we get $(\beta)^+$ of 1.1, too? The first difference is in phrasing the question. Now, for $g \in G$; it is

Question $^\mu$: Does $\langle \bar{\gamma}^{g,\delta} : \delta \in S_\mu \rangle$ satisfy:

$$\begin{aligned} \left((\forall f_0 \in \lambda^+ \mu_0)(\forall f_1 \in \lambda^+ \mu_1) \cdots (\forall f_i \in \lambda^+ \mu_i) \cdots \right)_{i < \sigma} \\ (\exists^{\text{stat}} \delta \in S_\mu) \left(\bigwedge_{i < \sigma} f_i(\gamma_{2i}^{g,\delta}) = f_i(\gamma_{2i+1}^{g,\delta}) \right). \end{aligned}$$

If for some g the answer is yes, we are done; so assume not. Therefore we have $f_{g,i} \in \lambda^+(\mu_i)$ for $g \in G, i < \sigma$ and club E_g of λ^+ such that

$$\delta \in S_\mu \cap E_g \Rightarrow (\exists i < \sigma)(f_{g,i}(\gamma_{2i}^{g,\delta}) \neq f_{g,i}(\gamma_{2i+1}^{g,\delta})).$$

A second difference is the choice of f^* as $f^*(\sigma\varepsilon + i, \alpha) = f_{g_\varepsilon,i}(\alpha)$ for $\varepsilon < 2^\sigma, i < \sigma, \alpha < \lambda^+$.

Lastly, the equations later change slightly. ■_{1.1}

1.2 Fact (1) Under the assumptions (a)+(b) of 1.1, letting $\bar{\lambda} = \langle \lambda_i : i < \sigma \rangle$ be increasingly continuous with limit λ such that $2^\sigma < \lambda_0, 2^{\lambda_i} < \lambda_{i+1}$ we have

$(*)_1 + (*)_2$ where

$(*)_1$ we can find $\langle \langle \gamma_\zeta^\delta : \zeta < \lambda \rangle : \delta \in S \rangle$ such that

(α) γ_ζ^δ is increasing in ζ with limit δ ,

(β)⁺ if $f_i: \lambda^+ \rightarrow \lambda_{i+1}$, for $i < \sigma$, then the following set is stationary

$\{\delta \in S : f_i(\gamma_\zeta^\delta) = f_i(\gamma_\xi^\delta) \text{ when } \zeta, \xi \in [\lambda_i, \lambda_{i+1}] \text{ for every } i < \sigma\}$;

$(*)_2$ moreover, if $F_i: [\lambda^+]^{<\lambda} \rightarrow [\lambda^+]^{\lambda^+}$ for $i < \sigma$ (or just $F_i: [\lambda^+]^{<\lambda} \rightarrow [\lambda^+]^\lambda$) and $\sup(w) < \min(F_i(w))$ for $w \in [\lambda^+]^{<\lambda}$, for each $i < \sigma$, then in addition we can demand

(i) $\{\gamma_\zeta^\delta : \zeta \in [\lambda_i, \lambda_{i+1}]\} \subseteq F_i(\{\gamma_\zeta^\delta : \zeta < \lambda_i\})$,

(ii) $|\{\langle \gamma_\zeta^\delta : \zeta < \zeta^* \rangle : \gamma_{\zeta^*}^\delta = \gamma\}| \leq \lambda$ for each $\gamma < \lambda^+$ and $\zeta^* < \sigma$.

(2) Assume $\lambda, \langle \lambda_i : i < \sigma \rangle$ are as in part (1) and $\langle C_\delta : \delta \in S \rangle$ is given; it guesses clubs (for λ^+ , which means that for every club E of λ^+ the set $\{\delta \in S : C_\delta \subseteq E\}$ is a stationary subset of λ^+) and $C_\delta = \{\alpha[\delta, i] : i < \sigma\}$, $\alpha[\delta, i]$ divisible by λ^ω increasing in i with limit δ ; $\langle \text{cf}(\alpha[\delta, i+1]) : i < \sigma \rangle$ is increasing with limit λ and let $\beta(\delta, i) = \sum_{j < i} \lambda_j \times \text{cf}(\alpha[\delta, j])$. Then

$(*)$ we can find $\langle \langle \gamma_\zeta^\delta : \zeta < \lambda \rangle : \delta \in S \rangle$ such that

(α) $\langle \gamma_\zeta^\delta : \zeta < \lambda \rangle$ is increasing with limit δ , (for $\delta \in S$),

(β) $\sup\{\gamma_\zeta^\delta : \gamma_\zeta^\delta < \beta[\delta, j+1]\} = \alpha[\delta, j]$,

(γ) for every $f_i \in {}^{(\lambda^+)}(\mu_i)$ for $i < \sigma$ where $\mu_i < \lambda$ and club E of λ^+ , for stationarily many $\delta \in S$ we have $\{\gamma_i^\delta : i < \lambda\} \subseteq E$ and $f_i(\gamma_\zeta^\delta) = f_i(\gamma_\varepsilon^\delta)$, when $\zeta, \varepsilon \in [\beta[\delta, i] + \lambda_i \xi, \beta[\delta, i] + \lambda_i \xi + \lambda_i)$ and $\xi < \text{cf}(\alpha[\delta, i])$.

Proof: (1) The same proof as in 1.1 for $(*)_1$, but see a proof after the proof of 4.2.

(2) Should be clear, too. $\blacksquare_{1.2}$

§2. Case C: Forcing for successor of singulars

We continue [Sh 587].

2.1 Hypothesis: (1) λ strong limit singular $\sigma = \text{cf}(\lambda) < \lambda$, $\kappa = \lambda^+$, $\mu^* \geq \kappa$, $2^\lambda = \lambda^+$.

2.2 Definition: (1) Let $\mathfrak{C}_{<\kappa}(\mu^*)$ be the family of $\hat{E}_0 \subseteq \{\bar{a} : \bar{a} = \langle a_i : i \leq \alpha \rangle \text{ where } \alpha < \kappa, a_i \in [\mu^*]^{<\kappa} \text{ increasing continuous, and } a_i \cap \kappa \in \kappa\}$ such that: for every $\theta = \text{cf}(\theta) < \lambda$, χ large enough and $x \in \mathcal{H}(\chi)$ we can find $\langle N_i : i \leq \theta \rangle$ obeying $\bar{a} \in \hat{E}_0$ (with error some n , see [Sh 587, B.5.1(1)]) and such that $x \in N_0$; this repeats [Sh 587, B.5.1(2)]; formally we should say that \bar{N} obeys \bar{a} for μ^* .

(2) $\mathfrak{C}_{<\kappa}^1(\mu^*)$ is the family of $\hat{\mathcal{E}}_1 \subseteq \{\bar{a} = \langle a_i : i \leq \sigma \rangle, a_i \text{ increasing continuous, } i < \sigma \Rightarrow |a_i| < \lambda \text{ and } \lambda + 1 \subseteq \bigcup_{i < \sigma} a_i\}$.

2.3 Definition: (1) We say $\bar{M} = \langle M_i : i \leq \sigma \rangle$ is ruled by $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$ if, for some $\chi > \mu^*$:

- (a) $\hat{\mathcal{E}}_0 \in \mathfrak{C}_{<\kappa}(\mu^*), \hat{\mathcal{E}}_1 \in \mathfrak{C}_{<\kappa}^1(\mu^*)$,
- (b) for* some $\langle \bar{M}^i : -1 \leq i < \sigma \rangle$ and $\langle \bar{N}^i : -1 \leq i < \sigma \rangle$ we have:
 - (α) $M_i \prec (\mathcal{H}(\chi), \epsilon, <_\chi^*)$,
 - (β) \bar{M} obeys some $\bar{a} \in \hat{\mathcal{E}}_1$ for some finite error (so for some n , for every $i, a_i \subseteq M_i \cap \mu^* \subseteq a_{i+n}$) and $\bar{M} \upharpoonright (i+1) \in M_{i+1}$ and $j < i \Rightarrow M_j \prec M_i$ and M_i is increasing continuous,
 - (γ) $[M_{i+1}]^{2^{\|M_i\|}} \subseteq M_{i+1}$ for i a limit ordinal $< \sigma$,
 - (δ) $\bar{M}^i = \langle M_\alpha^i : \alpha \leq \delta_i \rangle, \bar{N}^i = \langle N_\alpha^i : \alpha \leq \delta_i \rangle$ and $M_\alpha^i \prec N_\alpha^i \prec (\mathcal{H}(\chi), \epsilon, <_\chi^*)$ and $\lambda + 1 \subseteq N_\alpha^i$ and $\|M_\alpha^i\| = \|M_\alpha^i\|^{\|M_i\|}$ for $\alpha < \delta_i$ non-limit, $[M_\beta^i]^{\|M_i\|} \subseteq M_{\beta+1}^i$, for $\beta < \delta_i$,
 - (ϵ) $\langle N_\alpha^i : \alpha \leq \delta_i \rangle = \bar{N}^i$ obeys some $\bar{b}_i \in \hat{\mathcal{E}}_0$ for some finite error and \bar{M}^i, \bar{N}^i are increasing continuous,
 - (ζ) $M_{i+1} = M_{\delta_i}^i \subseteq N_{\delta_i}^i$ and $\langle (\bar{M}^j, \bar{N}^j) : j < i \rangle \in M_0^i$,
 - (η) $\delta_i \subseteq M_{i+1}$ (hence $\delta_i < \lambda$) and $\lambda \subseteq N_\alpha^i$,
 - (θ) $\text{cf}(\delta_i) > 2^{\|M_i\|}$ for i limit,
 - (ι) $\bar{N}^i \upharpoonright (\alpha + 1), \bar{M}^i \upharpoonright (\alpha + 1) \in M_{\alpha+1}^i$ for $\alpha < \delta_i, i < \sigma$, hence $N_\beta^i = \text{Sk}_{(\mathcal{H}(\chi), \epsilon, <_\chi^*)}(M_\beta^i \cup \lambda)$ when $i < \sigma$ and $\beta \leq \delta_i$ is a limit ordinal,
 - (κ) $N_{\delta_i}^i \prec N_0^j$ for $i < j$,
 - (λ) $M_i \prec M_0^i, M_i \in M_0^i$.

(2) We say above that $(\langle \bar{M}^i : i < \sigma \rangle, \langle \bar{N}^i : i < \sigma \rangle)$ is an $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$ -approximation to \bar{M} .

(3) Let $\mathfrak{C}_{<\kappa}^\spadesuit(\mu^*)$ be the family of $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$ such that:

- (a) $\hat{\mathcal{E}}_0 \in \mathfrak{C}_{<\kappa}(\mu^*)$ and $\hat{\mathcal{E}}_1 \in \mathfrak{C}_{<\kappa}^1(\mu^*)$,
- (b) for χ large enough and $x \in \mathcal{H}(\chi)$ we can find \bar{M} which is ruled by $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$ and $x \in M_0$,
- (c) $\hat{\mathcal{E}}_0$ is closed (see below).

(4) $\hat{\mathcal{E}}_0$ is closed if $\langle a_i : i \leq \alpha \rangle \in \hat{\mathcal{E}}_0, \gamma \leq \beta \leq \alpha$ implies $\langle a_i : i \in [\beta, \gamma] \rangle \in \hat{\mathcal{E}}_0$.

Remark: (1) In Definition 2.3(1), letting $\bar{N} = \bar{N}^0 \hat{\ } \bar{N}^1 \dots$, i.e., $\bar{N} = \langle N_i : i < \lambda \rangle, N_\epsilon =: N_\alpha^i$ if $\epsilon = \sum_{j < i} \delta_j + \alpha$; hence $\text{lg}(\bar{N}) = \lambda$ and $\bar{N} \upharpoonright (i_0 + 1) \in N_{i_0+1}$ so \bar{N} is \prec -increasingly continuous, and $\gamma < \lambda \Rightarrow \bar{N} \upharpoonright \gamma \in N_{\gamma+1}$.

* We may later ignore the $i = -1$ in our notation.

2.4 CLAIM: (1) Assume $\hat{\mathcal{E}}_0 \in \mathfrak{C}_{<\kappa}(\mu^*)$ and $\bar{\mathbb{Q}} = \langle \mathbb{P}_\alpha, \mathbb{Q}_i : i < \gamma \rangle$ is a ($< \kappa$)-support iteration such that $\Vdash_{\mathbb{P}_i} \text{“}\mathbb{Q}_i \text{ is strongly } \hat{\mathcal{E}}_0\text{-complete”}$ for each $i < \gamma$; see [Sh 587, B.5.3(3)]. Then \mathbb{P}_γ is strongly $\hat{\mathcal{E}}_0$ -complete (hence $\mathbb{P}_\gamma/\mathbb{P}_\beta$).

(2) If \mathbb{Q} is $\hat{\mathcal{E}}_0$ -complete, then $\mathbf{V}^{\mathbb{Q}} \models \hat{\mathcal{E}}_0$ non-trivial.

Proof: By [Sh 587, B.5.6] (here the choice “for any regular cardinal $\theta < \kappa$ ” rather than “for any cardinal $\theta < \kappa$ ” in [Sh 587, B.5.1(2)] is important). $\blacksquare_{2.4}$

2.5 Definition: Let $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1) \in \mathfrak{C}_{<\kappa}^\clubsuit(\mu^*)$ and let \mathbb{Q} be a forcing notion.

(1) For a sequence $\bar{M} = \langle M_i : i \leq \sigma \rangle$ ruled by $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$ with an $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$ -approximation $(\langle \bar{M}^i : i < \sigma \rangle, \langle \bar{N}^i : i < \sigma \rangle)$ and a condition $r \in \mathbb{Q}$ we define a game $\mathfrak{G}_{\bar{M}, \langle \bar{M}^i : i < \sigma \rangle, \langle \bar{N}^i : i < \sigma \rangle}^\clubsuit(\mathbb{Q}, r)$ between two players COM and INC.

The play lasts σ moves during which the players construct a sequence $\langle i_0, p, \langle p_i, \bar{q}_i : i_0 - 1 \leq i < \sigma \rangle \rangle$ such that $i_0 < \sigma$ is non-limit, $p \in M_{i_0} \cap \mathbb{Q}$, $p_i \in M_{i+1} \cap \mathbb{Q}$, $\bar{q}_i = \langle q_{i,\varepsilon} : \varepsilon < \delta_i \rangle \subseteq \mathbb{Q}$ (where $\delta_i + 1 = \text{lg}(\bar{N}^i)$).

The player INC first decides what is $i_0 < \delta$ and then it chooses a condition $p \in \mathbb{Q} \cap M_{i_0}$ stronger than r . Next, at the stage $i \in [i_0 - 1, \delta)$ of the game, COM chooses $p_i \in \hat{\mathbb{Q}} \cap M_{i+1}$ such that:

- (i) $p \leq_{\mathbb{Q}} p_i$,
- (ii) $(\forall j < i)(\forall \varepsilon < \delta_j)(q_{j,\varepsilon} \leq_{\mathbb{Q}} p_i)$,
- (iii) if i is a non-limit ordinal, then $p_i \in \hat{\mathbb{Q}}$ is minimal satisfying (i)+(ii),
- (iv) if i is a limit ordinal, then $p_i \in \mathbb{Q}$.

Now the player INC answers, choosing an increasing sequence $\bar{q}_i = \langle q_{i,\varepsilon} : \varepsilon < \delta_i \rangle$ such that $p_i \leq_{\mathbb{Q}} q_{i,0}$ and \bar{q}_i is $(\bar{N}^i \upharpoonright [\alpha, \delta_i], \mathbb{Q})^*$ -generic for some $\alpha < \delta_i$ (see [Sh 587, B.5.3.1]) and $\beta < \delta_i \Rightarrow \bar{q}_i \upharpoonright (\beta + 1) \in M_{i,\beta+1}$. The player COM wins if it has always legal moves and the sequence $\langle p_i : i < \sigma \rangle$ has an upper bound in \mathbb{Q} .

(2) We say that the forcing notion \mathbb{Q} is complete for $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$ or $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$ -complete if

- (a) \mathbb{Q} is strongly complete for $\hat{\mathcal{E}}_0$ and
- (b) for a large enough regular χ , for some $x \in \mathcal{H}(\chi)$, for every sequence \bar{M} ruled by $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$ with an $\hat{\mathcal{E}}_0$ -approximation $(\langle \bar{M}^i : i < \sigma \rangle, \langle \bar{N}^i : i < \sigma \rangle)$ and such that $x \in M_0$ and for any condition $r \in \mathbb{Q} \cap M_0$, the player INC does not have a winning strategy in the game $\mathfrak{G}_{\bar{M}, \langle \bar{M}^i : i < \sigma \rangle, \langle \bar{N}^i : i < \sigma \rangle}^\clubsuit(\mathbb{Q}, r)$.

2.6 PROPOSITION: Assume

- (a) $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1) \in \mathfrak{C}_{<\kappa}^\clubsuit(\mu^*)$,
- (b) \mathbb{Q} is a forcing notion for $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$.

Then $\Vdash_{\mathbb{Q}} \text{“}(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1) \in \mathfrak{C}_{<\kappa}^\clubsuit(\mu^*)\text{”}$.

Proof: Straightforward (and not used in this paper).

2.7 PROPOSITION: Assume that $\hat{\mathcal{E}} \in \mathfrak{C}_{<\kappa}(\mu^*)$ is closed and $\bar{\mathbb{Q}} = \langle \mathbb{P}_\alpha, \mathbb{Q}_\alpha : \alpha < \gamma \rangle$ is a $(< \kappa)$ -support iteration of forcing notions which are strongly complete for $\hat{\mathcal{E}}$. Let $\mathcal{T} = (T, <^\pm, rk)$ be a standard $(w, \alpha_0)^\gamma$ -tree (see [Sh 587, A.3.3]), $\|\mathcal{T}\| < \lambda, w \subseteq \gamma, \alpha_0$ an ordinal, and let $\bar{p} = \langle p_t : t \in T \rangle \in FTr'(\bar{\mathbb{Q}})$; see [Sh 587, A.3.2]. Suppose that \mathcal{I} is an open dense subset of \mathbb{P}_γ . Then there is $\bar{q} = \langle q_t : t \in T \rangle \in FTr'(\bar{\mathbb{Q}})$ such that $\bar{p} \leq \bar{q}$ and, for each $t \in T$,

- (a) $q_t \in \{q \upharpoonright rk(t) : q \in \mathcal{I}\}$, and
- (b) for each $\alpha \in \text{Dom}(q_t)$, one of the following occurs:
 - (i) $q_t(\alpha) = p_t(\alpha)$,
 - (ii) $\Vdash_{\mathbb{P}_\alpha} \text{“} q_t(\alpha) \in \mathbb{Q}_\alpha \text{”}$ (not just in the completion $\hat{\mathbb{Q}}_\alpha$),
 - (iii) $\Vdash_{\mathbb{P}_\alpha} \text{“} \text{there is } r \in \mathbb{Q}_\alpha \text{ such that } \hat{\mathbb{Q}}_\alpha \models p_t(\alpha) \leq r \leq q_t(\alpha) \text{”}$ (not really needed).

Proof: Just like the proof of [Sh 587, B.7.1].

Our next proposition corresponds to [Sh 587, B.7.2], which corresponds to [Sh 587, A.3.6]. The difference with [Sh 587, B.7.2] is the appearance of the \bar{M}, \bar{M}^i .

2.8 PROPOSITION: Assume that $\hat{\mathcal{E}} \in \mathfrak{C}_{<\kappa}(\mu^*)$ is closed and $\bar{\mathbb{Q}} = \langle \mathbb{P}_\alpha, \mathbb{Q}_\alpha : \alpha < \gamma \rangle$ is a $(< \kappa)$ -support iteration and $x = \langle x_\alpha : \alpha < \gamma \rangle$ is such that

$$\Vdash_{\mathbb{P}_\alpha} \text{“} \mathbb{Q}_\alpha \text{ is strongly complete for } \hat{\mathcal{E}} \text{ with witness } x_\alpha \text{”}$$

(for $\alpha < \gamma$). Further suppose that

- (α) (\bar{N}, \bar{a}) is an $\hat{\mathcal{E}}$ -complementary pair (see [Sh 587, B.5.1]), $\bar{N} = \langle N_i : i \leq \delta \rangle$ and $x, \hat{\mathcal{E}}, \bar{\mathbb{Q}} \in N_0$,
- (β) $\mathcal{T} = (T, <^\pm, rk) \in N_0$ is a standard $(w, \alpha_0)^\gamma$ -tree, $w \subseteq \gamma \cap N_0, \|w\| < \text{cf}(\delta), \alpha_0$ is an ordinal, $\alpha_1 = \alpha_0 + 1$ and $0 \in w$,
- (γ) $\bar{p} = \langle p_t : t \in T \rangle \in FTr'(\bar{\mathbb{Q}}) \cap N_0, w \in N_0$, (of course $\alpha_0 \in N_0$, on FTr' see [Sh 587, A.3.2]),
- (δ) $\bar{M} = \langle M_i : i \leq \delta \rangle, M_i \prec (H(\chi), \in, <_\chi^*), M_i$ is increasing continuous, $[M_i]^{\|w\|+|T|} \subseteq M_{i+1}$ and the pair $(\bar{M} \upharpoonright (i+1), \bar{N} \upharpoonright (i+1))$ belongs to $M_{i+1}, M_i \prec N_i$ and $w \cup \{x, \hat{\mathcal{E}}_0, \bar{\mathbb{Q}}\} \in M_0$,
- (ε) for $i \leq \delta, \mathcal{T}_i = (T_i, <_i, rk_i)$ is such that T_i consists of all sequences $t = \langle t_\zeta : \zeta \in \text{dom}(t) \rangle$ such that $\text{dom}(t)$ is an initial segment of w , and
 - (i) each t_ζ is a sequence of length α_1 ,
 - (ii) $\langle t_\zeta \upharpoonright \alpha_0 : \zeta \in \text{dom}(t) \rangle \in T$,

- (iii) for each $\zeta \in \text{dom}(t)$, either $t_\zeta(\alpha_0) = *$ or $t_\zeta(\alpha_0) \in M_i$ is a \mathbb{P}_ζ -name for an element of \mathbb{Q}_ζ and
 if $t_\zeta(\alpha) \neq *$ for some $\alpha < \alpha_0$, then $t_\zeta(\alpha_0) \neq *$,
- (iv) $\text{rk}_i(t) = \min(w \cup \{\zeta\} \setminus \text{dom}(t))$ and $<_i$ is the extension relation.

Then

- (a) each \mathcal{T}_i is a standard $(w, \alpha_1)^\gamma$ -tree, $\|\mathcal{T}_i\| \leq \|T\| \cdot \|M_i\|^{\|w\|}$, and if $i < \delta$ then $\mathcal{T}_i \in N_{i+1}$,
- (b) \mathcal{T} is the projection of each \mathcal{T}_i onto (w, α_0) and \mathcal{T}_i is increasing with i ,
- (c) there is $\bar{q} = \langle q_t : t \in T_\delta \rangle \in FTr'(\bar{\mathbb{Q}})$ such that
- (i) $\bar{p} \leq_{\text{proj}_T^{\mathcal{T}_\delta}} \bar{q}$,
 - (ii) if $t \in T_\delta \setminus \{<>\}$ then the condition $q_t \in \mathbb{P}'_{\text{rk}_\delta(t)}$ is an upper bound of an $(\bar{N} \upharpoonright [i_0, \delta], \mathbb{P}_{\text{rk}_\delta(t)})^*$ -generic sequence (where $i_0 < \delta$ is such that $t \in T_{i_0}$) and for every $\beta \in \text{dom}(q_t) = N_\delta \cap \text{rk}(t)$, $q_t(\beta)$ is a name for the least upper bound in \mathbb{Q}_β of an $(\bar{N}[G_\beta] \upharpoonright [\xi, \delta], \mathbb{Q}_\beta)^*$ -generic sequence (for some $\xi < \delta$),
 [Note that by [Sh 587, B.5.5], the first part of the demand on q_t implies that, if $i_0 \leq \xi$, then $q_t \upharpoonright \beta$ forces that $(\bar{N}[G_\beta] \upharpoonright [\xi, \delta], \bar{a} \upharpoonright [\xi, \delta])$ is an $\hat{\mathcal{S}}$ -complementary pair.]
 - (iii) if $t \in T_\delta, t' = \text{proj}_T^{\mathcal{T}_\delta}(t) \in T, \zeta \in \text{dom}(t)$ and $t_\zeta(\alpha_0) \neq *$, then
 $q_t \upharpoonright \zeta \Vdash_{\mathbb{P}_\zeta} "p_{t'}(\zeta) \leq_{\mathbb{Q}_\zeta} t_\zeta(\alpha_0) \Rightarrow t_\zeta(\alpha_0) \leq_{\mathbb{Q}_\zeta} q_t(\zeta)",$
 - (iv) $q_{<>} = p_{<>}$.

Proof: Clauses (a) and (b) should be clear. Clause (c) is proved as in [Sh 587, B.7.2]. $\blacksquare_{2.8}$

Remark: In 2.9 below it is proved as in the inaccessible case, i.e., the proofs of ([Sh 587, B.7.3]) with $\bar{M}, \langle \bar{N}^i : i < \sigma \rangle$ as in Definition 2.5. We define the trees point: in stage i using trees \mathcal{T}_i with set of levels $w_i = M_i \cap \gamma$ and looking at all possible moves of COM, i.e., $p_i \in M_{i+1} \cap \mathbb{P}_\gamma$, so constructing this tree of conditions in δ_i stages, in stage $\varepsilon < \delta_i$, has $|N_\varepsilon^i \cap M_{i+1}|^{2^{\|M_i\|}}$ nodes.

Now

$$p \in \mathbb{P}_\gamma \cap M_{i+1} \Rightarrow \text{Dom}(p) \subseteq M_{i+1}$$

but

$$p \in \mathbb{P}_\gamma \cap M_{i+1} \Rightarrow \text{Dom}(p) \subseteq M_\sigma = \bigcup_{i < \omega \sigma} N_{\delta_i}^i,$$

$$p \in \mathbb{P}_\gamma \cap N_\varepsilon^i \Rightarrow \text{Dom}(p) \subseteq N_\varepsilon^i.$$

So in limit cases $i < \sigma$: the existence of limit is by the clause (μ) of Definition 2.3. In the end we use the winning of the play and then need to find a branch in the tree of conditions of level σ : like Case A using $\hat{\mathcal{E}}_0$.

2.9 THEOREM: Suppose that $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1) \in \mathfrak{C}_{<\kappa}^\spadesuit(\mu^*)$ (so $\hat{\mathcal{E}}_0 \in \mathfrak{C}_{<\kappa}(\mu^*)$) and $\bar{\mathbb{Q}} = \langle \mathbb{P}_\alpha, \bar{\mathbb{Q}}_\alpha : \alpha < \gamma \rangle$ is a $(< \kappa)$ -support iteration such that for each $\alpha < \gamma$

$$\Vdash_{\mathbb{P}_\alpha} \text{“}\bar{\mathbb{Q}}_\alpha \text{ is complete for } (\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)\text{”}.$$

Then

- (a) $\Vdash_{\mathbb{P}_\gamma} (\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1) \in \mathfrak{C}_{<\kappa}^\spadesuit(\mu^*)$; moreover
- (b) \mathbb{P}_γ is complete for $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$.

Proof: We need only part (a) of the conclusion, so we concentrate on it. Let χ be a regular large enough regular cardinal, \bar{x} be a name for an element of $\mathcal{H}(\chi)$ and $p \in \mathbb{P}_\gamma$. Let $x_\alpha \in \mathcal{H}(\chi)$ be a \mathbb{P}_α -name for the witness that $\bar{\mathbb{Q}}_\alpha$ is (forced to be) complete for $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$ and let $\bar{x} = \langle x_\alpha : \alpha < \gamma \rangle$. Since $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1) \in \mathfrak{C}_{<\kappa}^\spadesuit(\mu^*)$, we find $\bar{M} = \langle M_i : i \leq \sigma \rangle$ which is ruled by $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$ with an $\hat{\mathcal{E}}_0$ -approximation $\langle \bar{M}^i, \bar{N}^i : -1 \leq i < \sigma \rangle$ and such that $p, \bar{\mathbb{Q}}, \bar{x}, \hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1 \in M_0$ (see 2.3). Let $\bar{N}^i = \langle N_\varepsilon^i : \varepsilon \leq \delta_i \rangle$ and let $\bar{a}^i \in \hat{\mathcal{E}}_0$ be such that (\bar{N}^i, \bar{a}^i) is an $\hat{\mathcal{E}}_0$ -complementary pair and let $\bar{M}^i = \langle M_\varepsilon^i : \varepsilon \leq \delta_i \rangle$. Let $w_i = \{0\} \cup \bigcup_{w_j \leq i} (\gamma \cap M_{w_j})$ (for $i \leq \delta$). By the demands of 2.3 we know that $\|w_i\| < \text{cf}(\delta_i)$, $w_i \in M_0^i$.

By induction on $i \leq \sigma$ we define standard $(w_i, i)^\gamma$ -trees $\mathcal{T}_i \in M_{i+1}$ and $\bar{p}^i = \langle p_t^i : t \in T_i \rangle \in \text{FTr}'(\bar{\mathbb{Q}}) \cap M_{i+1}$ such that $\|T_i\| \leq \|M_i\|^{\|w_i\|} \leq \|M_{i+1}\|$ if i is limit or 0, $w_{i+1} = w_i$ hence $\mathcal{T}_{i+1} = \mathcal{T}_i$, and if $j < i \leq \delta$ then $\mathcal{T}_j = \text{proj}_{(w_j, j+1)}^{(w_i, i+1)}(\mathcal{T}_i)$ and $\bar{p}^j \leq_{\text{proj}_{\mathcal{T}_j}^{\mathcal{T}_i}} \bar{p}^i$.

CASE 1: $i = 0$.

Let T_0^* consist of all sequences $\langle t_\zeta : \zeta \in \text{dom}(t) \rangle$ such that $\text{dom}(t)$ is an initial segment of w_0 and $t_\zeta = \langle \rangle$ for $\zeta \in \text{dom}(t)$. Thus T_0^* is a standard $(w_0, 0)^\gamma$ -tree, $\|T_0^*\| = \|w_0\| + 1$. For $t \in T_0^*$ let $p_t^{*0} = p \upharpoonright \text{rk}_0^*(t)$. Clearly the sequence $\bar{p}^{*0} = \langle p_t^{*0} : t \in T_0^* \rangle$ is in $\text{FTr}'(\bar{\mathbb{Q}}) \cap N_0^{-1}$. Apply 2.8 to $\hat{\mathcal{E}}_0, \bar{\mathbb{Q}}, \bar{N}^{-1}, T_0^*, w_0$ and \bar{p}^{*0} (note that $\|M_\varepsilon^{-1}\|^{\|w_0\|} \subseteq M_\varepsilon^{-1}$ for $\varepsilon < \delta_0$). As a result we get a $(w_0, 1)^\gamma$ -tree \mathcal{T}_0 (the one called \mathcal{T}_{δ_0} there) and $\bar{p}^0 = \langle p_t^0 : t \in T_0 \rangle \in \text{FTr}'(\bar{\mathbb{Q}}) \cap M_1$ (the one called \bar{q} there) satisfying clauses $(\varepsilon), (c)(i)$ –(iv) of 2.8 and such that $\|T_0\| \leq \|N_{\delta_0}^{-1}\|^{\|w_0\|} = \|M_0\|^{\|w_0\|} = \|M_0\|$ (remember $\text{cf}(\delta_0) > 2^{\|M_0\|}$). So, in particular, if $t \in T_0, \zeta \in \text{dom}(t)$ then $t_\zeta(0) \in M_1$ is either $*$ of a \mathbb{P}_ζ -name for an element of $\bar{\mathbb{Q}}_\zeta$.

Moreover, we additionally require that $(\mathcal{T}_0, \bar{p}^0)$ is the $<_\chi^*$ -first with all these properties, so $\mathcal{T}_0, \bar{p}^0 \in M_1$.

CASE 2: $i = i_0 + 1$.

We proceed similarly to the previous case. Suppose we have defined \mathcal{T}_{i_0} and \bar{p}^{i_0} such that $\mathcal{T}_{i_0}, \bar{p}^{i_0} \in M_{i_0+1}$, $\|T_{i_0}\| \leq \|M_{i_0+1}\|$. Let \mathcal{T}_i^* be a standard $(w_i, i_0)^\gamma$ -tree such that

T_i^* consists of all sequences $\langle t_\zeta : \zeta \in \text{dom}(t) \rangle$ such that $\text{dom}(t)$ is an initial segment of w_i and

$$\langle t_\zeta : \zeta \in \text{dom}(t) \cap w_{i_0} \rangle \in T_{i_0} \text{ and } (\forall \zeta \in \text{dom}(t) \setminus w_{i_0})(\forall j < i_0)(t_\zeta(j) = *).$$

Thus, $T_{i_0} = \text{proj}_{(w_{i_0}, i_0)}^{(w_i, i)}(T_i^*)$ and $\|T_i^*\| \leq \|M_i\|$. Let $p_t^{*i} = p_{t'}^{i_0} \upharpoonright \text{rk}_i^*(t)$ for $t \in T_i^*$, $t' = \text{proj}_{T_{i_0}}^{T_i}(t)$. Now apply 2.8 to $\hat{\mathcal{E}}_0, \hat{\mathcal{Q}}, \bar{N}^{i_0}, T_i^*, w_i$ and \bar{p}^{*i} (check that the assumptions are satisfied). So we get a standard $(w_i, i_0 + 1)^\gamma$ -tree T_i and a sequence \bar{p}^i satisfying (ε) , (c)(i)–(iv) of 2.8, and we take the $\langle \chi \rangle$ -pair (T_i, \bar{p}^i) with these properties. In particular, we will have $\|T_i\| \leq \|M_{i_0}\| \cdot \|N_{\delta_1}^{i_0}\|^{M_{i_0}} = \|M_{i_0+1}\|$ and $\bar{p}^i, T_i \in M_{i+1}$.

CASE 3: i is a limit ordinal.

Suppose we have defined T_j, \bar{p}^j for $j < i$ and we know that $\langle (T_j, \bar{p}^j) : j < i \rangle \in M_{i+1}$ (this is the consequence of taking “the $\langle \chi \rangle$ -first such that ...”). Let $T_i^* = \lim(\langle T_j : j < i \rangle)$. Now, for $t \in T_i^*$ we would like to define p_t^{*i} as the limit of $p_{\text{proj}_{T_j}^{T_i^*}(t)}^j$. However, our problem is that we do not know if the limit exists.

Therefore, we restrict ourselves to these t for which the respective sequence has an upper bound. To be more precise, for $t \in T_i^*$ we apply the following procedure.

⊗ Let $t^j = \text{proj}_{T_j}^{T_i^*}(t)$ for $j < i$. Try to define inductively a condition $p_t^{*i} \in \mathbb{P}_{\text{rk}_i^*(t)}$ such that $\text{dom}(p_t^{*i}) = \cup\{\text{dom}(p_{t^j}^j) \cap \text{rk}_i^*(t) : j < i\}$. Suppose we have successfully defined $p_t^{*i} \upharpoonright \alpha$ for $\alpha \in \text{dom}(p_t^{*i})$, in such a way that $p_t^{*i} \upharpoonright \alpha \geq p_{t^j}^j \upharpoonright \alpha$ for all $j < i$. We know that

$$p_t^{*i} \upharpoonright \alpha \Vdash_{\mathbb{P}_\alpha} \text{“the sequence } \langle p_{t^j}^j(\alpha) : j < i \rangle \text{ is } \leq_{\mathbb{Q}_\alpha} \text{-increasing”}.$$

So now, if there is a \mathbb{P}_α -name τ for an element of \mathbb{Q}_α such that

$$p_t^{*i} \upharpoonright \alpha \Vdash_{\mathbb{P}_\alpha} \text{“}(\forall j < i)(p_{t^j}^j(\alpha) \leq_{\mathbb{Q}_\alpha} \tau)\text{”},$$

then we take the \mathbb{P}_α -name of the lub of $\langle p_{t^j}^j(\alpha) : j < i, p_{t^j}^j(\alpha) \neq * \rangle$ in $\hat{\mathbb{Q}}$, and we continue. If there is no such τ , then we decide that $t \notin T_i^+$ and we stop the procedure.*

Now, let T_i^+ consist of those $t \in T_i^*$ for which the above procedure resulted in a successful definition of $p_t^{*i} \in \mathbb{P}_{\text{rk}_i^*(t)}$. It might not be clear at the moment if T_i^+ contains anything more than $\langle \rangle$, but we will see that this is the case. Note that

$$\|T_i^+\| \leq \|T_i^*\| \leq \prod_{j < i} \|T_j\| \leq \prod_{j < i} \|M_j\| \leq 2^{\|M_i\|} \leq \|M_0^i\|.$$

* Generally in such situation we can act as in 2.7 to get a real decision, i.e., if $p_t^{*i} \upharpoonright (\alpha + 1)$ is not well defined while $p_t^{*i} \upharpoonright \alpha$ is well defined then $p_t^{*i} \upharpoonright \alpha \Vdash$ “the sequence $\langle p_{t^j}^j(\alpha) : j < i \rangle$ has no $\leq_{\mathbb{Q}_\alpha}$ -upper bound”. But the need has not arisen here.

Moreover, for nonlimit $\varepsilon > 2$ we have $\|M_\varepsilon^i\|^{w_i} + \|T_\varepsilon^+\| \leq \|M_\varepsilon^i\|^{M_\varepsilon} \subseteq M_{\varepsilon+1}^i$ and $\mathcal{T}_i^+, \bar{p}^{*i} \in M_{i+1}$. Let $\mathcal{T}_i = \mathcal{T}_i^*, \bar{p}^i = \bar{p}^{*i}$ (this time there is no need to take the $\langle \cdot \rangle_\chi^*$ -first pair as the process leaves no freedom). So we have finished case 3.

After the construction is carried out we continue in a similar manner as in [Sh 587, A.3.7] (but note a slightly different meaning of the \ast 's here).

So we let $\mathcal{T}_\sigma = \lim(\langle \mathcal{T}_i : i < \sigma \rangle)$. It is a standard $(\sigma, \sigma)^\gamma$ -tree. By induction on $\alpha \in w_\sigma \cup \{\gamma\}$ we choose $q_\alpha \in \mathbb{P}'_\alpha$ and a \mathbb{P}_α -name t_α such that:

- (a) $\Vdash_{\mathbb{P}_\alpha} "t_\alpha \in T_\sigma \ \& \ \text{rk}_\delta(t_\alpha) = \alpha"$ and let $i_0^\alpha = \min\{i < \delta : \alpha \in M_i\} < \sigma$,
- (b) $\Vdash_{\mathbb{P}_\alpha} "t_\beta = t_\alpha \upharpoonright \beta"$ for $\beta < \alpha$,
- (c) $\text{dom}(q_\alpha) = w_\delta \cap \alpha$,
- (d) if $\beta < \alpha$ then $q_\beta = q_\alpha \upharpoonright \beta$,
- (e) $p_{\text{proj}_{\mathcal{T}_i^\delta}(t_\alpha)}^i$ is well defined and $p_{\text{proj}_{\mathcal{T}_i^\delta}(t_\alpha)}^i \upharpoonright \alpha \leq q_\alpha$ for each $i < \sigma$,
- (f) for each $\beta < \alpha$

$q_\alpha \Vdash_{\mathbb{P}_\alpha} "(\forall i < \delta)((t_{\beta+1})_\beta(i) = \ast \Leftrightarrow i < i_0^\beta)"$ and the sequence

$$\langle i_0^\beta, p_{\text{proj}_{\mathcal{T}_i^\delta}(t_{\beta+1})}^{i_0^\beta}(\beta), \langle (t_{\beta+1})_\beta(i), p_{\text{proj}_{\mathcal{T}_i^\delta}(t_{\beta+1})}^i(\beta) : i_0^\beta \leq i < \delta \rangle \rangle$$

is a result of a play of the game $\mathfrak{G}_{\bar{M}[G_\beta], \langle \bar{N}^i[G_\beta] : i < \delta \rangle}^\spadesuit(\mathbb{Q}_\beta, 0_{\mathbb{Q}_\beta})$,
won by player COM",

- (g) the condition q_α forces (in \mathbb{P}_α) that
"the sequence $\bar{M}[G_{\mathbb{P}_\alpha}] \upharpoonright [i_\alpha, \delta]$ is ruled by $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$ and
 $\langle \bar{N}^i[G_{\mathbb{P}_\alpha}] : i_0^\alpha \leq i < \sigma \rangle$ is its $\hat{\mathcal{E}}_0$ -approximation".

(Remember: $\hat{\mathcal{E}}_1$ is closed under end segments.) This is done completely in parallel to the last part of the proof of [Sh 587, A.3.7].

Finally, look at the condition q_γ and the clause (g) above. ■_{2.9}

2.10 Generalization (1) $\hat{\mathcal{E}}_1$ is a set of triples $\langle \bar{a}, \langle \bar{b}^i, \bar{a}^i : i < \sigma \rangle, \bar{\lambda} \rangle$, $\bar{a} = \langle a_i : i \leq \sigma \rangle$, $\bar{a}^i = \langle a_\alpha^i : \alpha \leq \delta_i \rangle$, $\bar{b}^i = \langle b_\alpha^i : \alpha \leq \delta_i \rangle \in \hat{\mathcal{E}}_0$, $a_{\delta_i}^i = a_{i+1}$, $a_i \subseteq b_0^i$, $\lambda = \langle \lambda_i : i < \sigma \rangle$ an increasing sequence of cardinals $< \lambda$, $\sum \lambda_i = \lambda$.

(2) We say $(\bar{M}, \langle \bar{M}^i : i < \sigma \rangle, \langle \bar{N}^i : i < \sigma \rangle)$ obeys $(\bar{a}, \langle \bar{b}^i : i < \bar{\lambda} \rangle)$ **iff**: $M_i \cap \mu^* = a_i$, \bar{N}^i obeys \bar{b}^i all things in 2.3 but $\lambda_i \geq \|M_i\|$, $\lambda_i \geq \prod_{j \leq i} \|M_j\|$, $[M_\alpha^i]^{\lambda_i} \subseteq M_{\alpha+1}^i$ for $\alpha < \delta_i$ (so earlier $\lambda_i = 2^{\|M_i\|}$).

2.11 Conclusion (1) Assume

- (a) $S \subseteq \{\delta < \kappa : \text{cf}(\delta) = \sigma\}$ is stationary not reflecting,
- (b) $\bar{a} = \langle \bar{a}_\delta : \delta \in S \rangle$, $\bar{a}_\delta = \langle a_{\delta,i} : i \leq \sigma \rangle$, $\delta = a_{\delta,\sigma}$ and $a_{\delta,i}$ increasing with i and $i < \sigma \Rightarrow |a_{\delta,i}| < \lambda$ and $\sup(a_{\delta,i}) < \delta$
[variant: $\bar{\lambda}^\delta = \langle \lambda_i^\delta : i < \sigma \rangle$ increasing with limit λ],

- (c) we let $\mu^* = \kappa$, $\hat{\mathcal{E}}_0 = \hat{\mathcal{E}}_0[S] = \{\bar{a} : \bar{a} = \langle a_i : i \leq \alpha \rangle, \alpha < \kappa, a_i \in \kappa \setminus S \text{ increasing continuous}\}$,
- (d) $\hat{\mathcal{E}}_1 = \{\bar{a}_\delta : \delta \in S\}$
(or $\{\langle \bar{a}_\delta, \langle \bar{a}^{\delta,i}, \bar{b}^{i,\delta} : i < \sigma \rangle, \bar{\lambda}^\delta \rangle : \delta \in S\}$ appropriate for (2.10)),
- (e) we assume the pair $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1) \in \mathfrak{C}_{<\kappa}^\blacklozenge(\mu^*)$,
- (f) $\mu = \mu^\kappa, \kappa < \tau = \text{cf}(\tau) < \mu$.

Then for some $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$ -complete forcing notion \mathbb{P} of cardinality μ we have

$\Vdash_{\mathbb{P}}$ “forcing axiom for $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$ -complete forcing notion
of cardinality $\leq \kappa$ and $< \tau$ of open dense sets”

and in $\mathbf{V}^{\mathbb{P}}$ the set S is still stationary (by preservation of $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$ -nontrivial).

(2) If clauses (a),(c) holds and \diamond_S , then for some \bar{a} , if we define $\hat{\mathcal{E}}_1$ as in clause (d) then clauses (b),(d),(e) hold.

Proof: (1) See more at the end of §3.

(2) Easy. ■_{2.11}

2.12 Application: In $\mathbf{V}^{\mathbb{P}}$ of 2.11:

(a) if

- (i) $\theta < \lambda, A_\delta \subseteq \delta = \sup(A_\delta)$ for $\delta \in S$,
- (ii) $|A_\delta| < \theta$,
- (iii) $\bar{h} = \langle h_\delta : \delta \in S \rangle, h_\delta: A \rightarrow \theta$,
- (iv) $A_\delta \subseteq \bigcup \{a_{\delta,i+1} \setminus a_{\delta,i} : i < \sigma\}$,

then for some $h: \kappa \rightarrow \theta$ and club E of κ we have $(\forall \delta \in S \cap E)[h_\delta \subseteq^* h]$ where $h' \subseteq^* h''$ means that $\sup(\text{Dom}(h')) > \sup\{\alpha : \alpha \in \text{Dom}(h') \text{ and } \alpha \notin \text{Dom}(h'')\}$ or

$\alpha \in \text{Dom}(h'') \ \& \ h'(\alpha) \neq h''(\alpha)$,

(b) if we add: “ h_δ constant”, then we can omit the assumption (iii),

(c) we can weaken $|A_\delta| < \theta$ to $|A_\delta \cap a_{\delta,i+1}| \leq |a_{\delta,i}|$,

(d) in (c) we can weaken $|A_\delta| \leq \theta \vee |A_\delta \cap a_{\delta,i+1}| \leq |a_{\delta,i}|$ to $h_\delta \upharpoonright a_{\delta,i+1}$ belongs to $M_{i+1} \cap N_\alpha^i$ for some $\alpha < \delta_i$

(remember $\text{cf}(\sup a_{\delta,i+1}) > \lambda_i^\delta$).

2.13 Remark: (1) Compared to [Sh 186] the new point in the application is (b).

(2) You may complain why not having the best of (a)+(b), i.e., combine their good points. The reason is that this is impossible by §1, §4; the situation is different in the inaccessible case.

Proof: Should be clear. Still, we say something in case h_δ constant, that is (b).

Let

$$\mathbb{Q} = \{(h, C) : h \text{ is a function with domain an ordinal} \\ \alpha < \kappa = \lambda^+, \\ C \text{ a closed subset of } \alpha + 1, \alpha \in C \\ \text{and } (\forall \delta \in C \cap S \cap (\alpha + 1))(h_\delta \subseteq^* h)\}$$

with the partial order being inclusion.

For $p \in \mathbb{Q}$ let $p = (h^p, C^p)$.

So clearly if $(h, C) \in \mathbb{Q}$ and $\alpha = \text{Dom}(h) < \beta \in \kappa$ then for some h_1 we have $h \subseteq h_1 \in \mathbb{Q}_1$, $\text{Dom}(h_1) = \beta$; moreover, if $\gamma < \theta$ & $\beta \notin S$ then $(h, C) \leq (h \cup \gamma_{[\alpha, \beta]}, C \cup \{\beta\}) \in \mathbb{Q}$.

The main point is proving \mathbb{Q} is complete for $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$. Now “ \mathbb{Q} is strongly complete for $\hat{\mathcal{E}}_0$ ” is proved as in [Sh 587, B.6.5.1, B.6.5.2] (or 3.14 below which is somewhat less similar). The main point is clause (b) of 2.5(2); that is, let $\bar{M}, \langle \bar{M}^i : i < \omega\sigma \rangle, \langle \bar{N}^i : i < \omega\sigma \rangle$ be as there. In the game $\mathfrak{G}_{\bar{M}, \langle N_i : i < \omega\sigma \rangle}(r, \mathbb{Q})$ from 2.5(1), we can even prove that the player COM has a winning strategy: in stage i (non-trivial): if h_δ is constantly $\gamma < \theta$ or just $h_\delta \upharpoonright (A_\delta \cap a_{\delta, i+1} \setminus a_{\delta, i})$ is constantly $\gamma < \theta$ then we let

$$p_i = \left(\cup \{h^{\alpha_\zeta^j} : j < i \text{ and } \zeta < \delta_i\} \cup \gamma_{[N_{\delta_i}^i \cap \kappa, \beta_i]}, \right. \\ \left. \text{closure}(\cup \{C^{\alpha_\zeta^j} : j < i \text{ and } \zeta < \delta_i\} \cup \{\beta_i\}) \right)$$

for some $\beta_i \in M_{i+1} \cap \kappa \setminus M_i$ large enough such that $A_\delta \cap M_{i+1} \cap \kappa \subseteq \beta_i$. ■2.12

Remark: In the example of uniformizing (see [Sh 587]), if we use this forcing, the density is less problematic.

2.14 CLAIM: (1) In 2.12's conclusion we can omit the club E , that is, let $E = \kappa$ and demand $(\forall \delta \in S)(h_\delta \subseteq^* h)$ provided that we add in 2.12, recalling $S \subseteq \kappa$ does not reflect is a set of limit ordinals and

$$\bar{A} = \langle A_\delta : \delta \in S \rangle, A_\delta \subseteq \delta = \sup(A_\delta)$$

satisfies

(*) $\delta_1 \neq \delta_2$ in $S \Rightarrow \sup(A_{\delta_1} \cap A_{\delta_2}) < \delta_1 \cap \delta_2$.

(2) If $(\forall \delta \in S) \text{otp}(A_\delta) = \theta$ this always holds.

Proof: We define $\mathbb{Q} = \{h : \text{Dom}(h) \text{ is an ordinal } < \kappa \text{ and } h(\beta) \neq 0 \wedge \beta \in \text{Dom}(h) \rightarrow (\exists \delta \in S)[h_\delta(\beta) = h(\beta)] \text{ and } \delta \in (\text{Dom}(h) + 1) \cap S \text{ implies } h_\delta \subseteq^* h\}$ ordered by \subseteq . Now we should prove the parallel of the fact:

☒' if $p \in \mathbb{Q}$, $\alpha = \text{Dom}(p) < \beta < \kappa$ then there is q such that $p \leq q \in \mathbb{Q}$ and $\text{Dom}(q) = \beta$.

Why does this hold? We can find $\langle A'_\delta : \delta \in S \cap (\beta + 1) \rangle$ such that $A'_\delta \subseteq A_\delta$, $\sup(A_\delta \setminus A'_\delta) < \delta$ and $\bar{A}' = \langle A'_\delta : \delta \in S \cap (\beta + 1) \rangle$ is pairwise disjoint.

Now choose q as follows:

$$\text{Dom}(q) = \beta,$$

$$q(j) = \begin{cases} p(j) & \text{if } j < \alpha, \\ h_\delta(j) & \text{if } j \in A'_\delta \setminus \alpha \text{ and } \delta \in S \cap (\beta + 1) \setminus (\alpha + 1), \\ 0 & \text{if otherwise.} \end{cases}$$

Why does \bar{A}' exist? Prove by induction on β that for any $\bar{A}^1, \langle A'_\delta : \delta \in S \cap (\alpha + 1) \rangle$ as above and β satisfying $\alpha < \beta < \kappa$, we can extend \bar{A}^1 to $\langle A'_\delta : \delta \in S \cap (\beta + 1) \rangle$ which is as above. ■_{2.14}

2.15 Remark: Note: concerning κ inaccessible we could imitate what is here: having $M_{i+1} \not\prec_{\neq} N_{\delta_i}^i, \bigcup_{i < \delta} M_i = \bigcup_{i < \delta} N_{\delta_i}^i$.

As long as we are looking for a proof that no sequences of length $< \kappa$ are added, the gain is meagre (restricting the \bar{q} 's by $\bar{q} \upharpoonright \alpha \in N'_{\alpha+1}$). Still, if you want to make the uniformization and some diamond we may consider this.

2.16 Comment: We can weaken further the demand, by letting COM have more influence. E.g., we have (in 2.3) $\delta_i = \lambda_i = \text{cf}(\lambda_i) = \|M_{i+1}\|, D_i$ a $|a_i|^+$ -complete filter on λ_i , the choice of \bar{q}^i in the result of a game in which INC should have chosen a set of players $\in D_i$ and \diamond_{D_i} holds (as in the treatment of case E^* here).

The changes are obvious, but I do not see an application at the moment.

§3. κ^+ -c.c. and κ^+ -pic

We intend to generalize pic of [Sh f, Ch.VIII, §1]. The intended use is for iteration with each forcing $> \kappa$ — see use in [Sh f]. In [Sh 587, B.7.4] we assume each \mathbb{Q}_i of cardinality $\leq \kappa$. Usually $\mu = \kappa^+$.

Note: $\hat{\mathcal{E}}_0$ is as in the accessible case, in [Sh 587], but this part works in the other cases. In particular, in Cases A, B (in [Sh 587]'s context) if the length of $\bar{a} \in \hat{\mathcal{E}}_0$ is $< \lambda$ (remember $\kappa = \lambda^+$), then we have $(< \lambda)$ -completeness implies $\hat{\mathcal{E}}_0$ -completeness AND in 3.7 even $\bar{a} \in \hat{\mathcal{E}}_0 \Rightarrow \ell g(\bar{a}) = \omega$ is O.K.

In Case A on the $S_0 \subseteq S_\lambda^\kappa$ if $\ell g(\bar{a}) = \lambda, a_\lambda \in S_0$ is O.K., too. STILL one can start with other variants of completeness which is preserved.

3.1 Context: We continue [Sh 587, B.5.1–B.5.7(1)] (except the remark [Sh 587, B.5.2(3)]) under the weaker assumption $\kappa = \kappa^{<\kappa} > \aleph_0$, so κ is not necessarily

strongly inaccessible; also in our $\hat{\mathcal{E}}$'s we allow \bar{a} such that $|a_\delta| = |\delta|$ is strongly inaccessible.

3.2 Definition: Assume:

- ☒(a) $\mu = \text{cf}(\mu) > |\alpha|^{<\kappa}$ for $\alpha < \mu$,
- (b) the triple $(\kappa, \mu^*, \hat{\mathcal{E}}_0)$ satisfies: $\kappa = \text{cf}(\kappa) > \aleph_0, \mu^* \geq \kappa, \hat{\mathcal{E}}_0 \subseteq \{\bar{a} : \bar{a} \text{ an increasing continuous sequence of members of } [\mu^*]^{<\kappa} \text{ of limit length } < \kappa \text{ with } a_i \cap \kappa \in \kappa\}$, and
- (c) $S^\square \subseteq \{\delta < \mu : \text{cf}(\delta) \geq \kappa\}$ stationary.

For $\ell = 1, 2$ we say \mathbb{Q} satisfies $(\mu, S^\square, \hat{\mathcal{E}}_0)$ -pic $_\ell$ if, for some $x \in \mathcal{H}(\chi)$ (can be omitted, essentially, i.e., replaced by \mathbb{Q}), we have

(*) if

- (α) $S \subseteq S^\square$ is stationary and $\langle \mu, S, \hat{\mathcal{E}}_0, x \rangle \in N_0^\alpha$,
- (β) for $\alpha \in S, \delta_\alpha < \kappa$, and
 - (i) if $\ell = 1, \bar{N}^\alpha = \langle N_i^\alpha : i \leq \delta_\alpha \rangle$ and $c_\alpha = \delta_\alpha$ and $\bar{N}^{\alpha,*} = \bar{N}^\alpha$,
 - (ii) if $\ell = 2$ then $\bar{N}^{\alpha,*} = \langle N_i^\alpha : i \leq \delta_\alpha \rangle, \bar{N}^\alpha = \langle N_i^\alpha : i \in c_\alpha^+ \rangle$ where $c_\alpha \subseteq \delta_\alpha = \sup(c_\alpha), c_\alpha^+ = c_\alpha \cup \{\delta_\alpha\}, c_\alpha$ is closed, $\gamma < \beta \in c_\alpha \Rightarrow c_\alpha \cap \gamma \in N_\beta^\alpha$,
- (γ) $(\bar{N}^\alpha, \bar{a}^\alpha)$ is $\hat{\mathcal{E}}_0$ -complementary (see [Sh 587, B.5.3]); so \bar{N}^α obeys $\bar{a}^\alpha \in \hat{\mathcal{E}}_0$ (with some error n_α) (so here we have $\|N_{\delta_\alpha}^\alpha\| < \kappa, \delta_\alpha < \kappa$),
- (δ) \bar{p}^α is $(\bar{N}^\alpha, \mathbb{Q})^1$ -generic (see [Sh 587, Definition B.5.3.1]),
- (ε) $\alpha \in N_0^\alpha$ and
 - (i) if $\ell = 1$, then for some club C of μ for every $\alpha \in S$ we have $\langle (\bar{N}^\beta, \bar{p}^\beta) : \beta \in S \cap C \cap \alpha \rangle$ belongs to N_0^α ,
 - (ii) if $\ell = 2$, then for some club C of μ for every $\alpha \in S \cap C$ and $i < \delta_\alpha$ we have $\langle (\bar{N}^{\beta,*} \upharpoonright (i+1), \bar{p}^\beta \upharpoonright (i+1)) : \beta \in S \cap C \rangle$ belongs to N_{i+1}^α ,
- (ζ) we define a function g with domain S as follows: $g(\alpha) = (g_0(\alpha), g_1(\alpha))$ where

$$g_0(\alpha) = N_{\delta_\alpha}^\alpha \cap \left(\bigcup_{\beta < \alpha} N_{\delta_\beta}^\beta \right) \text{ and } g_1(\alpha) = (N_{\delta_\alpha}^\alpha, N_i^\alpha, c)_{i < \delta_1, c \in g_0(\alpha)} / \cong,$$

then we can find a club C of μ such that:

if $\alpha < \beta$ & $g(\alpha) = g(\beta)$ & $\alpha \in C \cap S$ & $\beta \in C \cap S$ then $\delta_\alpha = \delta_\beta, g(\alpha) = g(\beta)$, for some $h, N_{\delta_\alpha}^\alpha \cong_h N_{\delta_\beta}^\beta$ (really unique), and for each $i < \delta_\alpha$ the function h maps N_i^α to N_i^β, p_i^α to p_i^β and $\{p_i^\alpha : i < \delta_\alpha\} \cup \{p_i^\beta : i < \delta_\beta\}$ has an upper bound.

3.3 CLAIM: Assume \boxtimes , i.e., (a), (b), (c) of 3.2 and

(d) $\hat{\mathcal{E}}_0$ is non-trivial, which means:

for every χ large enough and $x \in \mathcal{H}(\chi)$ there is $\bar{N} = \langle N_i : i \leq \delta \rangle$ increasingly continuous, $N_i \prec (\mathcal{H}(\chi), \in)$, $x \in N_i$, $\|N_i\| < \kappa$, $\bar{N} \upharpoonright (i+1) \in N_{i+1}$ and \bar{N} obeys some $\bar{a} \in \hat{\mathcal{E}}_0$ with some finite error n ,

(e) \mathbb{Q} is a strongly $cl(\hat{\mathcal{E}}_0)$ -complete forcing notion (hence adding no new bounded subsets of κ) where $cl(\hat{\mathcal{E}}_0) =: \{\bar{a} \upharpoonright [\alpha, \beta] : \bar{a} \in \hat{\mathcal{E}}_0 \text{ and } \alpha \leq \beta \leq lg(\bar{a})\}$,

(f) \mathbb{Q} satisfies $(\mu, S^\square, \hat{\mathcal{E}}_0)$ -pic $_\ell$ where $\ell \in \{1, 2\}$.

Then \mathbb{Q} satisfies the μ -c.c. provided that

(*) $\ell = 1$ or $\ell = 2$ and $\hat{\mathcal{E}}_0$ is fat; see below.

3.4 Definition: We say $\hat{\mathcal{E}}_0 \in \mathfrak{C}_{<\kappa}^-(\mu^*)$ is fat, if in the following game $\mathcal{D}_{\kappa, \mu^*}(\hat{\mathcal{E}}_0)$ between fat and lean, the fat player has a winning strategy.

A play last, κ moves; in the α -th move:

Case 1: α nonlimit.

The player lean chooses a club $Y_\alpha \subseteq [\mu^*]^{<\kappa}$, the fat player chooses $a_\alpha \in Y_\alpha$ and $\mathcal{P}_\alpha \subseteq \{c : c \subseteq \alpha \text{ is closed}\}$ of cardinality $< \kappa$.

Case 2: α limit.

We let $Y_\alpha = [\mu_0]^{<\kappa}$ and $a_\alpha = \cup\{a_\beta : \beta < \alpha\}$ and the player fat chooses $\mathcal{P}_\alpha \subseteq \{C : C \subseteq \alpha \text{ is closed}\}$ of cardinality $< \kappa$.

In a play, fat wins iff for some limit ordinal α and $c \in \mathcal{P}_\alpha$ we have:

(*) (i) $\beta \in c \Rightarrow c \cap \beta \in \mathcal{P}_\beta$,

(ii) $\alpha = \sup(c)$,

(iii) $\langle a_\beta : \beta \in c \cup \{\alpha\} \rangle \in \hat{\mathcal{E}}_0$.

3.5 Remark: (0) With more care in the game (Definition 3.4) we incorporate choosing the $p^\alpha - s$. In 3.2(*) (ε) (ii) we can add $\langle N_{i+1}^\beta : \beta \in \alpha \cap c \rangle$ belong to N_{i+1}^α .

(1) In Definition 3.4, without loss of generality $c \in \mathcal{P}_\alpha \& \beta \in c \Rightarrow c \cap \beta \in \mathcal{P}_\beta$.

(2) If κ is strongly inaccessible, without loss of generality we have $\mathcal{P}_\alpha = \mathcal{P}(\alpha)$, so fat has a winning strategy.

(3) In general being fat is a weak demand, e.g., if $\hat{\mathcal{E}}_0 \supseteq \{\bar{a} : \bar{a} = \langle a_i : i \leq \omega \rangle, a_\omega = \bigcup_n a_n, a_i \in [\mu^*]^{<\kappa} \text{ is increasing}\}$.

Proof of 3.3: Case 1: $\ell = 1$.

Assume $p_\alpha \in \mathbb{Q}$ for $\alpha < \mu$ and let χ be large enough and x as in Definition 3.2. We choose $(\bar{N}^\alpha, \bar{p}^\alpha)$ by induction on $\alpha < \mu$ as follows. If $\langle (\bar{N}^\beta, \bar{p}^\beta) : \beta < \alpha \rangle$ is already defined, as $\hat{\mathcal{E}}_0$ is non-trivial there is a pair $(\bar{N}^\alpha, \bar{a}^\alpha)$ which is $\hat{\mathcal{E}}_0$ -complementary and $\langle (\bar{N}^\beta, \bar{p}^\beta) : \beta < \alpha \rangle, \mathbb{Q}, \langle p_\beta : \beta < \mu \rangle, p_\alpha, \alpha, x$ belong to N_0^α

and let $\bar{N}^\alpha = \langle N_i^\alpha : i \leq \delta_i \rangle$. So $p_\alpha \in N_0^\alpha$ and we can choose $p_{\alpha,i} \in N_{i+1}^\alpha$ such that $p_\alpha = p_{\alpha,0}$ and $\langle p_{\alpha,i} : i < \delta_\alpha \rangle$ is $(\bar{N}^\alpha, \mathbb{Q})^1$ -generic.

[Why? By the proof of [Sh 587, B.5.6.4].] Now by “ \mathbb{Q} is $(\mu, S^\square, \hat{\mathcal{E}}_0)$ -pic $_\ell$ ”, for some $\alpha < \beta$ in S^\square , $\{p_i^\alpha : i < \delta_\alpha\} \cup \{p_i^\beta : i < \delta_\beta\}$ has a common upper bound hence, in particular, p_α, p_β are compatible.

Case 2: $\ell = 2$.

Assume $p_\alpha \in \mathbb{Q}$ for $\alpha < \mu$ and let χ be large enough. Let **St** be a winning strategy for the player fat in the game $\partial_{\kappa, \mu^*}(\hat{\mathcal{E}}_0)$. Now we choose by induction on $i < \kappa$, the tuple $(N_i^\alpha, \mathcal{P}_i^\alpha, Y_i^\alpha, \bar{p}_i^\alpha)$ where $\bar{p}_i^\alpha = \langle p_{i,c}^\alpha : c \in \mathcal{P}_i^\alpha \rangle$ for $\alpha < \mu$ such that:

- ⊠(a) $M_i^\alpha \prec (\mathcal{H}(\chi), \in, <_\chi^*)$,
- (b) M_i^α increasing continuous in i ,
- (c) $\|M_i^\alpha\| < \kappa$ and $\langle M_j^\alpha : j \leq i \rangle \in M_{i+1}^\alpha$ and $M_i^\alpha \cap \kappa \in \kappa$, and $p_\alpha \in M_i^\alpha$,
- (d) $\langle Y_j^\alpha, M_j^\alpha \cap \mu^*, \mathcal{P}_j^\alpha : j \leq i \rangle$ is an initial segment of a play of $\partial_{\kappa, \mu^*}(\hat{\mathcal{E}}_0)$ in which the player fat uses his winning strategy **St**,
- (e) $\langle (M_j^\beta, \mathcal{P}_j^\beta, Y_j^\beta, \bar{p}_j^\beta) : j \leq i, \beta \in S \rangle$ belongs to N_{i+1}^α (hence $\mathcal{P}_j^\alpha \subseteq M_{j+1}^\alpha$, etc.),
- (f) $p_{i,c}^\alpha \in \mathbb{Q} \cap N_{i+1}^\alpha$,
- (g) if $c \in \mathcal{P}_i^\alpha$ and $\langle p_{j,c \cap j}^\alpha : j \in c \rangle$ has an upper bound then $p_{i,c}^\alpha$ is such a bound,
- (h) $p_{i,c}^\alpha \in \cap \{ \mathcal{I} : \mathcal{I} \in M_i^\alpha \text{ is a dense open subset of } \mathbb{Q} \}$.

Can we carry the induction?

For i limit let $M_i^\alpha = \cup \{M_j^\alpha : j < i\}$ and choose $Y_i^\alpha, \mathcal{P}_i^\alpha$ by clause (d), i.e., by the rules of the game $\partial_{\kappa, \mu^*}(\hat{\mathcal{E}}_0)$ and p_i^α by clause (g)+(h) (possible as forcing by \mathbb{Q} adds no new sequences of length $< \kappa$ of members of \mathbf{V}). For i non-limit, let $x_i = \langle (M_j^\beta, \mathcal{P}_j^\beta, Y_j^\beta, \bar{p}_j^\beta) : j \leq i, \beta \in S \rangle$, let $Y_i^\alpha = \{a : a \in [\mu^*]^{<\kappa} \text{ and } \alpha \in a \text{ and } a = \mu^* \cap \text{Sk}_{(\mathcal{H}(\chi), \in, <_\chi^*)}^{<\kappa}(\{x_i \times \mathbb{Q}, \mathbf{St}, \alpha\})\}$ ($\text{Sk}^{<\kappa}$ means $a \in Y_i^\alpha \Rightarrow a \cap \kappa \in \kappa$) and let $(a_i^\alpha, \mathcal{P}_i^\alpha)$ be the move which the strategy **St** dictates to the player fat if the i -th move of lean is Y_i^α (and the play so far is $\langle (Y_j^\alpha, M_j^\alpha \cap \mu^*, \mathcal{P}_{\alpha,j}) : j < i \rangle$). Now we choose $M_i^\alpha = \text{Sk}_{(\mathcal{H}(\chi), \in, <_\chi^*)}^{<\kappa}(\{x_i, \mathbb{Q}, \mathbf{St}, \alpha\})$ and \mathcal{P}_i^α has already been chosen and $\bar{p}_i^\alpha = \langle p_{i,c}^\alpha : c \in \mathcal{P}_i^\alpha \rangle$ as in the limit case.

Having carried out the induction, for each $\alpha \in S$ in the play $\langle (Y_i^\alpha, M_i^\alpha \cap \mu^*, \mathcal{P}_i^\alpha) : i < \kappa \rangle$ the player fat wins the game having used the strategy **St**, hence there are a limit ordinal $i_\alpha < \kappa$ and closed $c_\alpha \in \mathcal{P}_{i_\alpha}$ such that $i_\alpha = \sup(c_\alpha)$ and $\langle M_j^\alpha : j \in c_\alpha \cup \{i_\alpha\} \rangle$ obeys some member \bar{a}_α of $\hat{\mathcal{E}}_0$. As \mathbb{Q} is $cl(\hat{\mathcal{E}}_0)$ -complete we can prove by induction on $j \in c_\alpha \cup \{i_\alpha\}$ that $\varepsilon < j$ & $\varepsilon \in C_\alpha \Rightarrow \mathbb{Q} \models p_{\varepsilon, c_\alpha \cap \varepsilon}^\alpha \leq p_{j, c_\alpha \cap j}^\alpha$.

Let $\delta_\alpha = i_\alpha, N_i^\alpha = M_i^\alpha$ for $i \leq \delta_\alpha$ and $\bar{p}^\alpha = \langle p_i^\alpha : i \in c_\alpha \rangle$. Now continue as in Case 1. ■_{3.3}

3.6 CLAIM: If $(*)$ of Definition 3.2, we can allow $\text{Dom}(g)$ to be a subset of $S \cap C$, $\langle A_i : i < \mu \rangle$ be an increasingly continuous sequence of sets, $|A_i| < \mu$, $N_{\delta_\alpha}^\alpha \subseteq A_{\alpha+1}$ replacing the definition of g, g_0 and by $g_0(\alpha) = N_{\delta_\alpha}^\alpha \cap A_\alpha$ and g_1 by $g_1(\alpha) = (N_{\delta_\alpha}^\alpha, N_i^\alpha, c)_{i < \delta_\alpha, c \in g_0(c)} / \cong$ (and get an equivalent definition).

Remark: If $\text{Dom}(g) \cap S^\square$ is not stationary, the definition says nothing.

Proof: Straightforward.

3.7 CLAIM: Assume clauses \boxtimes , i.e., (a), (b), (c) of 3.2 and (d) of 3.3.

For $(< \kappa)$ -support iteration $\bar{\mathbb{Q}} = \langle \mathbb{P}_i, \mathbb{Q}_i : i < \alpha \rangle$, if we have $\Vdash_{\mathbb{P}_i}$ “ \mathbb{Q}_i is $(\mu, S^\square, \hat{\mathcal{E}}_0)$ -pic $_\ell$ ” for each $i < \alpha$ and forcing with $\text{Lim}(\bar{\mathbb{Q}})$ add no bounded subsets of κ , then \mathbb{P}_γ and $\mathbb{P}_\gamma/\mathbb{P}_\beta$, for $\beta \leq \gamma \leq \text{lg}(\bar{\mathbb{Q}})$, are $\hat{\mathcal{E}}_0$ -complete $(\mu, S^\square, \hat{\mathcal{E}}_0)$ -pic $_\ell$.

3.8 Remark: We can omit the assumption “ $\text{Lim}(\bar{\mathbb{Q}})$ add no bounded subsets of κ ” if we add the assumption $\text{cl}(\hat{\mathcal{E}}_0) \in \mathfrak{C}_{< \kappa}(\mu^*)$, see [Sh 587, Def. B.5.1(2)], because with the latter assumption the former follows by [Sh 587, B.5.6].

Proof: Similar to [Sh f, Ch. VIII]. We first concentrate on

Case 1: $\ell = 1$.

It is enough to prove for \mathbb{P}_α .

We prove this by induction on α . Let $\Vdash_{\mathbb{P}_i}$ “ \mathbb{Q}_i is $(\mu, S^\square, \hat{\mathcal{E}}_0)$ -pic $_\ell$ as witnessed by x_i and let $\chi_i = \text{Min}\{\chi : x_i \in \mathcal{H}(\chi)\}$ ”.

Let $x = (\mu^*, \kappa, \mu, S^\square, \hat{\mathcal{E}}_0, \langle \langle \chi_i, x_i \rangle : i < \text{lg}(\bar{\mathbb{Q}}) \rangle)$ and assume χ is large enough such that $x \in \mathcal{H}(\chi)$ and let $\langle \langle \bar{N}^\alpha, \bar{p}^\alpha \rangle : \alpha \in S \rangle$ be as in Definition 3.2, so $S \subseteq S^\square$ is stationary and $\bar{N}^\alpha = \langle N_i^\alpha : i \leq \delta_\alpha \rangle$. We define a g by

\boxtimes_1 g is a function with domain S ,

\boxtimes_2 $g(\alpha) = \langle g_\ell(\alpha) : \ell < 2 \rangle$ where

$$g_0(\alpha) = (N_{\delta_\alpha}^\alpha) \cap \left(\bigcup_{\beta < \alpha} N_{\delta_\beta}^\beta \right),$$

$$g_1(\alpha) = \text{the isomorphic type of } (N_{\delta_\alpha}^\alpha, N_i^\alpha, p_i^\alpha, c)_{c \in g_0(\alpha)}.$$

Let C be a club of μ such that $\alpha \in S \cap C \Rightarrow \langle \langle \bar{N}^\beta, \bar{p}^\beta \rangle : \beta < \alpha \rangle \in N_\alpha^\alpha$ (recall $\ell = 1$).

Fix y such that $S_y = \{\alpha \in S : g(\alpha) = y \text{ and } \alpha \in C\}$ is stationary.

Let $w_\alpha = \bigcup_{i < \delta_\alpha} \text{Dom}(p_i^\alpha)$, $w_y^* = w_\alpha \cap g_0(\alpha)$ for $\alpha \in S_y$ (as $\alpha \in S_y$, clearly the set does not depend on the α). For each $\zeta \in w_y^*$ we define a \mathbb{P}_ζ -name, $\underline{S}_{y,\zeta}$, as follows:

$$\underline{S}_{y,\zeta} = \{\alpha \in S_y : (\forall i < \delta_\alpha)(p_i^\alpha \upharpoonright \zeta \in G_{\mathbb{P}_\zeta})\}.$$

Now we try to apply Definition 3.2 in $\mathbf{V}^{\mathbb{P}^\zeta}$ to

$$\langle \langle \langle N_i^\alpha[G_{\mathbb{P}_\zeta}] : i \leq \delta_\alpha \rangle, \langle p_i^\alpha(\zeta)[G_{\mathbb{P}_\zeta}] : i < \delta_\alpha \rangle \rangle : \alpha \in \underline{S}_{y,\zeta}[G_{\mathbb{P}_\zeta}] \rangle.$$

Clearly, if $\mathcal{S}_{y,\zeta}[G_{\mathbb{P}_\zeta}]$ is a stationary subset of μ , we can apply it and $g_{y,\zeta}$ is the \mathbb{P}_ζ -name of a function with domain $\mathcal{S}_{y,\zeta}$ defined like g in (*) of Definition 3.2. Now $g_{y,\zeta}$ is well defined, and actually can be computed if we use $A_\beta = \cup\{N_{\delta_\alpha}^\alpha[G_{\mathbb{P}_\zeta}] : \alpha < \beta\}$. So by an induction hypothesis on α there is a suitable \mathbb{P}_ζ -name \mathcal{C}_ζ of a club of μ st in addition, if $\mathcal{S}_{y,\zeta}[G_{\mathbb{P}_\zeta}]$ is not a stationary subset of μ , let \mathcal{C}_ζ be a club of μ disjoint to it. But as \mathbb{P}_ζ satisfies the μ -c.c., without loss of generality $\mathcal{C}_\zeta = C_\zeta$ so $C' = C \cap \bigcap_{\zeta \in w_y^*} C_\zeta$ is a club of μ . Now choose $\alpha_1 < \alpha_2$ from $S_y \cap C'$ and we choose by induction on $\varepsilon \in w' = w_y^* \cup \{0, \ell g(\bar{Q})\}$ a condition $q_\varepsilon \in \mathbb{P}_\varepsilon$ such that:

▣₃(i) $\varepsilon_1 < \varepsilon \Rightarrow q_{\varepsilon_1} = q_\varepsilon \upharpoonright \varepsilon_1$,

(ii) q_ε is a bound to $\{p_u^{\alpha_1} \upharpoonright \varepsilon : i < \delta_{\alpha_1}\} \cup \{p_i^{\alpha_2} \upharpoonright \varepsilon : i < \delta_{\alpha_2}\}$.

For $\varepsilon = 0$ let $q_0 = \emptyset$. We have nothing to do really if ε is with no immediate predecessor in w ; we let q_ε be $\cup\{q_{\varepsilon_1} : \varepsilon_1 < \varepsilon, \varepsilon_1 \in w'\}$. So let $\varepsilon = \varepsilon_1 + 1, \varepsilon_1 \in w'$; now if $q_\varepsilon \in G \subseteq \mathbb{P}_{\varepsilon_1}$, G generic over V , then $\alpha_1, \alpha_2 \in \mathcal{S}_{y,\varepsilon_1}[G]$, hence $\mathcal{S}_{y,\zeta}[G] \cap C_{\varepsilon_1}$ is non-empty, hence is stationary, and we use Definition 3.2.

Case 2: $p = 2$.

Similar proof. ▣_{3.7}

3.9 CLAIM: Assume $\mu = \text{cf}(\mu) > \kappa, (\forall \alpha < \mu)(|\alpha|^{<\kappa} < \mu)$,

$$S \subseteq \{\delta < \mu : \text{cf}(\delta) \geq \kappa\}$$

is stationary. If $|\mathbb{Q}| \leq \kappa$ or just $< \mu, \mathcal{E}_0 \in \mathcal{C}_{<\kappa}^<(\mu^*)$, that is $\subseteq \{\bar{a} : \bar{a} \text{ increasingly continuous of length } < \kappa, a_i \in [\mu^*]^{<\kappa} \text{ and } a_i \cap \kappa \in \kappa\}$ non-trivial, possibly just for one cofinality, say \aleph_0 , then \mathbb{Q} satisfies κ^+ -pic $_\ell$.

Proof: Trivial, we get same sequence of condition or just see the proof of [Sh 587, B.7.4]. ▣_{3.9}

3.10 Discussion: (1) What is the use of pic?

In the forcing axioms instead of “ $|\mathbb{Q}| \leq \kappa$ ” we can write “ \mathbb{Q} satisfies the $(\mu, S^\square, \hat{\mathcal{E}}_0)$ -pic”. This strengthens the axioms.

In [Sh f], in some cases the length of the forcing is bounded (there ω_2) but here there is no need (as in [Sh f, Ch. VII, §1]).

This section applies to all cases in [Sh 587] and its branches.

(2) Note that we can demand that the p_i^α satisfies some additional requirements (in Definition 3.2), say $p_{2i}^\alpha = F_{\mathbb{Q}}(\bar{N} \upharpoonright (2i+1), \bar{p}^\alpha \upharpoonright (2i+1))$.

Let us see how this improves somewhat the results of [Sh 587, B.8] on $\mathcal{C}_{<\kappa}^\clubsuit(\mu^*)$, see [Sh 587, B.5.7.3].

3.11 Definition: Assume

- ⊗ $\kappa > \aleph_0$ is strongly inaccessible and $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1) \in \mathfrak{C}_{<\kappa}^\spadesuit(\mu^*)$ and θ_0, θ_1 are regular cardinals $> \kappa, \theta_2$ a cardinal $> \kappa$ (let $\bar{\theta} = (\theta_0, \theta_1, \theta_2)$, the usual case is $\theta_0 = \kappa^+$) and $\hat{\mathcal{E}} \subseteq \hat{\mathcal{E}}_1$ is nontrivial (see Definition 3.3, clause (d)) and $\ell \in \{1, 2\}$.

Let $Ax_{\theta_1, \theta_2}^\kappa(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1, \mathcal{E})$, the forcing axiom for $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1, \mathcal{E})$, and $\bar{\theta} = (\theta_0, \theta_1, \theta_2)$ be the following statement:

⊠ if

- (i) \mathbb{Q} is a forcing notion of cardinality $< \theta_1$,
- (ii) \mathbb{Q} is complete for $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$, see Definition [Sh 587, B.5.9(3)],
- (iii) \mathbb{Q} satisfies $(\theta_0, S^\square, \hat{\mathcal{E}})$ -pic $_\ell$,
- (iv) \mathcal{I}_i is a dense subset of \mathbb{Q} for $i < i^* < \theta_2$,

then there is a directed $H \subseteq \mathbb{Q}$ such that $(\forall i < i^*)(H \cap \mathcal{I}_i \neq \emptyset)$.

3.12 THEOREM: Assume ⊗ of Definition 3.11 and $\mu = \mu^{<\theta_1} = \mu^{<\theta_0} \geq \theta_0 + \theta_2$.

Then there is a forcing notion \mathbb{P} such that:

- (α) \mathbb{P} is complete for $\hat{\mathcal{E}}_0$,
- (β) \mathbb{P} has cardinality μ ,
- (γ) \mathbb{P} satisfies the θ_0 -c.c. and even the $(\kappa, \theta_0, \hat{\mathcal{E}})$ -pic $_\ell$,
- (δ) \mathbb{P} is complete for $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$, hence $\Vdash_{\mathbb{P}} “(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1) \in \mathfrak{C}_{<\kappa}^\spadesuit(\mu^*)”$ and more,
- (ε) $\Vdash_{\mathbb{P}} “Ax_{\bar{\theta}}^\kappa(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1, \mathcal{E})”$.

Proof: Like the proof of [Sh 587, B.8.2], using 3.7 instead of [Sh 587, B.7.4].

■_{3.12}

We may wonder how large can a stationary $S \subseteq \kappa$ be?

3.13 CLAIM: (1) Assume

- ⊗(a) κ is strongly inaccessible $> \aleph_0$,
- (b) $S \subseteq \kappa$ is stationary,
- (c) for letting $\mu^* = \kappa$ and $\hat{\mathcal{E}}_0 = \hat{\mathcal{E}}_0[S] = \{\bar{a} \in \mathfrak{C}_{<\kappa}(\mu^*) : \text{for every } i \leq \text{lg}(\bar{a}) \text{ we have } a_i \notin S\}$ we have $\hat{\mathcal{E}}_0 \in \mathfrak{C}_{<\kappa}(\mu^*)$,
- (d) we let $\hat{\mathcal{E}}_1 = \hat{\mathcal{E}}_1[S] = \{\bar{a} \in \mathfrak{C}_{<\kappa}(\mu^*) : \text{for every nonlimit } i \leq \text{lg}(\bar{a}) \text{ we have } a_i \notin S\}$.

Then

- (α) $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1) \in \mathfrak{C}_{<\kappa}^\spadesuit(\mu^*)$, see [Sh 587, B.5.7(3)].
- (2) The parallel of 2.11.

We now deal with forcing the failure of diamond on the set of inaccessibles.

3.14 CLAIM: Assume

- (a) $\kappa, S, \hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1$ are as in 3.13,
- (b) if $S_{bd} =: \{\theta < \kappa : \theta \text{ strongly inaccessible, } S \cap \theta \text{ is stationary in } \theta \text{ and } \diamond_{S \cap \theta}\}$ is not a stationary subset of κ ,
- (c) $\bar{A} = \langle A_\alpha : \alpha \in S \rangle, A_\alpha \subseteq \alpha$,
- (d) $\mathbb{Q} = \mathbb{Q}_{\bar{A}_1}$ is as in Definition 3.15 below,
- (e) $\hat{\mathcal{E}} \subseteq \hat{\mathcal{E}}_0$ is nontrivial.

Then

- (α) \mathbb{Q} is complete for $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$,
- (β) \mathbb{Q} satisfies the $(\kappa, \kappa^+, \hat{\mathcal{E}})$ -pic $_{\ell}$,
- (γ) \mathbb{Q} satisfies the κ^+ -c.c.

3.15 Definition: For $\kappa = \text{cf}(\kappa), S \subseteq \kappa = \text{sup}(S), \bar{A} = \langle A_\alpha : \alpha \in S \rangle$, with $A_\alpha \subseteq \alpha$ we define the forcing notions $\mathbb{Q} = \mathbb{Q}_{\bar{A}}^{\text{ad}}$ as follows:

- (a) $p \in \mathbb{Q}$ iff
 - (i) $p = (c, A) = (c^p, A^p)$,
 - (ii) c is \emptyset or a closed bounded subset of κ , hence has a last element,
 - (iii) $A \subseteq \text{sup}(c)$ such that,
 - (iv) if $\alpha \in C \cap S$ then $A \cap \alpha \neq A_\alpha$;
- (b) $p \leq q$ iff
 - (i) c^p is an initial segment of c^q ,
 - (ii) $A^p = A^q \cap \text{sup}(c^p)$.

Proof of 3.14: We concentrate on part (1), part (2)'s proof is similar. Now

- (*)₁ for every $\alpha < \kappa, \mathcal{I}_\alpha = \{p \in \mathbb{Q} : \alpha < \text{sup}(c^p)\}$ is dense open.
 [Why? If $p \in \mathbb{Q}$, let $\beta = \text{sup}(c^p) + 1 + \alpha$ and $q = (c^p \cup \{\beta\}, A^p)$, so $p \leq q \in \mathcal{I}_\alpha$.]
- (*)₂ If $\delta < \kappa$ is a limit ordinal, $\langle p_i : i < \delta \rangle$ is $\leq_{\mathbb{Q}}$ -increasing and $\text{sup}(c^{p_i}) \leq \alpha_{i+1} < \text{sup}(c^{p_{i+1}})$ for $i < \delta$, and for limit $i, \alpha_i = \cup\{\alpha_j : j < i\}$ and $\{\alpha_{1+i} : i < \delta\}$ is disjoint to S , then $p = (\bigcup_{i < \delta} c_i^{p_i}, \bigcup_{i < \delta} A^{p_i})$ is a $\leq_{\mathbb{Q}}$ -lub of $\langle p_i : i < \delta \rangle$.
 [Why? Just think.]
- (*)₃ Forcing with \mathbb{Q} adds no new sequences of length $< \kappa$ of ordinals (or members of \mathbf{V}).
 [Why? By (*)₂ + the assumption \oplus , clause (c) of Claim 3.13 as in [Sh 587, B.6].]
- (*)₄ \mathbb{Q} is complete for $\hat{\mathcal{E}}_0$
 [Why? Just think.]

(*)₅ \mathbb{Q} is complete for $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$; see [Sh 587, Def. B.5.9(3)].

[Why? Let χ be large enough and let $\langle M_i : i < \delta \rangle$ be ruled by $(\hat{\mathcal{E}}_0, \hat{\mathcal{E}}_1)$, with $\hat{\mathcal{E}}_0$ -approximation $\langle (\bar{N}^i, \bar{a}^i) : i < \delta \rangle$, see [Sh 587, Def. B.5.9(1)] and $r \in \mathbb{Q} \cap M_0$ and $S, \kappa, \bar{A} \in M_0$ and we have to prove that the player COM has a winning strategy in the game $\mathfrak{D}_{\bar{M}, \langle \bar{N}^i : i < \delta \rangle}(\mathbb{Q}, r)$.]

For this we proved by induction on $\delta < \kappa$ (a limit ordinal) the statement

\boxtimes_δ if $\langle M_i : i \leq \delta \rangle, \langle \bar{N}^i : i < \delta \rangle, r$ are as above (but α may be a nonlimit ordinal) $\bar{b} = \langle b_i : i < \delta \rangle, b_i \in [M_{i+1} \cap \kappa \setminus M_i]^{\leq \|M_i\|}$ and $B \subseteq M_\delta \cap \kappa$ (or just $B \subseteq \cup \{b_i : i < \delta\}$); then we can find p such that $r \leq p \in \mathbb{Q}$ and $A^p \cap b_i = B \cap b_i$ for every $i < \delta$ and $\sup(c^p) = M_\delta \cap \kappa$.

Case 1: α nonlimit. Trivial.

Case 2: α limit and for some $i < \alpha$, we have $\text{cf}(\delta) \leq \|M_i\|$.

Let $\theta = \text{cf}(\delta)$ and let $\langle \delta_\varepsilon : \varepsilon \leq \theta \rangle$ be increasing continuous, $\delta_0 = 0, \|M_{\delta_1}\| > \theta$ and $\delta_\theta = \delta$.

Choose $b \subseteq M_{\delta_1+1} \cap \kappa \setminus M_{\delta_1} \setminus b_{\delta_1}$ of cardinality θ and choose $b' \subseteq b$ such that $\zeta \in (\varepsilon, \delta] \Rightarrow A_{M_{\delta_\zeta} \cap \kappa} \cap b \neq b'$. By the induction hypothesis, we can find $r_{\delta_1} \in M_{\delta_1+1}$ such that $\sup(c^{r_1}) = M_{\delta_1} \cap \kappa, r \leq r_{\delta_0}, \beta < \delta_1 \Rightarrow A^{r_1} \cap b_\beta = B \cap b_\beta$ and r_1 is (M_β, \mathbb{Q}) -generic for every $\beta \leq \delta_1$. Let r_1^+ be such that $r_{\delta_1} \leq r_1^+ \in \mathbb{Q} \cap M_{\delta_1+1}$ and $\sup(b_{\delta_1} \cup b) < \sup(r_{\delta_1}^+)$ and $A^{r_1^+} \cap b_{\delta_1} = B \cap b_{\delta_1}$ and $A^{r_1^+} \cap b = b'$. Now we choose, by induction on $\varepsilon \in [2, \delta]$, a condition r_ε such that $r_\varepsilon \in M_{\delta_\varepsilon+1}$, $\sup(c^{r_\varepsilon}) = M_{\delta_\varepsilon} \cap \kappa, r_1^+ \leq r_\varepsilon, [\zeta \in [2, \varepsilon] \Rightarrow r_\zeta \leq r_\varepsilon]$ and $\beta < \delta_\varepsilon \Rightarrow A^{r_\varepsilon} \cap b_\beta = B \cap b_\beta$ and r_ε is (M_γ, \mathbb{Q}) -generic for $\gamma \leq \delta_\varepsilon$. For limit $\varepsilon, r_\varepsilon$ is uniquely determined and is $\in \mathbb{Q}$ by the choice of r_1^+ . For ε nonlimit use the induction hypothesis for $\langle M_\beta : \beta \in [\delta_\varepsilon + 1, \delta_{\varepsilon+1}] \rangle$.

Case 3: Neither Case 1 nor Case 2.

So α is strongly inaccessible, call it θ and $\theta = M_\theta \cap \kappa$; so as $\{\kappa, S\} \in M_\theta \prec (\mathcal{H}(\chi), \in, <^*)$, necessarily $\delta = \sup(S), \delta \in S_{bd}$ and $\neg \diamond_{\theta \cap S}$ (e.g., $\theta \cap S$ is not stationary in S). Choose for each $\beta < \theta$ an ordinal $\gamma_\beta \in M_{\beta+1} \cap \kappa \setminus M_\beta \setminus b_\beta$ and let $A'_i = \{j < i : \gamma_j \in A_{M_\beta \cap \kappa}\}$ for $i \in S \cap \theta$.

Now $\langle A'_i : i \in S \cap \theta \rangle$ cannot be a diamond sequence for θ , hence we can find $X \subseteq \theta$ and club C^- of θ such that $\delta \in X \cap S \Rightarrow A_\delta^- \neq X \cap \delta$. Let $C = \{i < \theta : i \text{ limit}, (\forall j < i)(\alpha_j < i) \text{ and } i \in C^- \text{ and } M_i \cap \kappa = i\}$, clearly C is a club of θ . Let $b_\beta^+ = a_\beta \cup \{\gamma_\beta\}, B^+ = B \cup \{\gamma_\beta : \beta \in X\}$, and proceed naturally.

■_{3.14}

3.16 Remark: So we can iterate and get that (GCH and) diamond fail for “most” stationary subsets of any strongly inaccessibles. We shall return to this elsewhere.

§4. Existence of non-free Whitehead (and $\text{Ext}(G, \mathbb{Z}) = \{0\}$) abelian groups in successor of singulars

In [Sh 587], the consistency with GCH of the following is proved for some regular uncountable κ : there is a κ -free nonfree abelian group of cardinality κ , and all such groups are Whitehead. We use κ inaccessible, here we ask: is this assumption necessary for the first such κ ?

The following claim seems to support the hope for a positive answer.

4.1 CLAIM: Assume

- (a) λ is strong limit singular, $\sigma = \text{cf}(\lambda) < \lambda, \kappa = \lambda^+ = 2^\lambda$,
- (b) $S \subseteq \{\delta < \kappa : \text{cf}(\delta) = \sigma\}$ is stationary,
- (c) S does not reflect or at least,
- (c)⁻ $\bar{A} = \langle A_\delta : \delta \in S \rangle, \text{otp}(A_\delta) = \sigma, \text{sup}(A_\delta) = \delta$, and \bar{A} is λ -free, i.e., for every $\alpha^* < \kappa$ we can find $\langle \alpha_\delta : \delta \in \alpha^* \cap S \rangle, \alpha_\delta < \delta$ such that $\langle A_\delta \setminus \alpha_\delta : \delta \in S \cap \alpha^* \rangle$ is a sequence of pairwise disjoint sets,
- (d) $\langle G_i : i \leq \sigma \rangle$ is a sequence of abelian groups such that:
 - (α) $\delta < \sigma$ limit $\Rightarrow G_\delta = \bigcup_{i < \delta} G_i$,
 - (β) $i < j \leq \sigma \Rightarrow G_j/G_i$ free and $G_i \subseteq G_j$,
 - (γ) $G_\sigma / \bigcup_{i < \sigma} G_i$ is not Whitehead,
 - (δ) $|G_\sigma| < \lambda$,
 - (ε) $G_0 = \{0\}$.

Then

(1) There is a strongly κ -free abelian group of cardinality κ which is not Whitehead, in fact $\Gamma(G) \subseteq S$.

(2) There is a strongly κ -free abelian group G^* of cardinality κ satisfying $\text{HOM}(G^*, \mathbb{Z}) = \{0\}$, in fact $\Gamma(G^*) \subseteq S$ (in fact, the same abelian group can serve).

(3) We can rephrase clause (d)(γ) of the assumption, i.e., " $G_\sigma / \bigcup_{i < \sigma} G_i$ is not Whitehead", by:

(d)(γ)⁻ some $f^* \in \text{HOM}(\bigcup_{i < \sigma} G_i, \mathbb{Z})$ cannot be extended to $f' \in \text{HOM}(G_\sigma, \mathbb{Z})$.

We first note:

4.2 CLAIM: Assume

- (a) λ strong limit singular, $\sigma = \text{cf}(\lambda) < \lambda, \kappa = 2^\lambda = \lambda^+$,
- (b) $S \subseteq \{\delta < \kappa : \text{cf}(\delta) = \sigma \text{ and } \lambda^\omega \text{ divides } \delta \text{ for simplicity}\}$ is stationary,
- (c) $A_\delta \subseteq \delta = \text{sup}(A_\delta), \text{otp}(A_\delta) = \sigma, A_\delta = \{\alpha_{\delta, \zeta} : \zeta < \sigma\}$ increasing with ζ ,
- (d) $h_0: \kappa \rightarrow \kappa$ and $h_1: \kappa \rightarrow \sigma$ are such that

$$(\forall \alpha < \kappa)(\forall \zeta < \sigma)(\forall \gamma \in (\alpha, \kappa))(\exists^\lambda \beta \in [\gamma, \gamma + \lambda])(h_0(\beta) = \alpha \text{ and } h_1(\beta) = \zeta),$$

and $(\forall \alpha < \kappa) h_0(\alpha) \leq \alpha$,

- (e) $\bar{\lambda} = \langle \lambda_\zeta : \zeta < \sigma \rangle$ is increasing continuous with limit λ such that $\lambda_0 = 0$ and $\zeta < \sigma \Rightarrow \lambda_{\zeta+1} = \text{cf}(\lambda_{\zeta+1}) > \sigma$.

Then we can choose $\langle (g_\delta, \langle \gamma_\zeta^\delta : \zeta < \lambda \rangle) : \delta \in S \rangle$ such that

- \odot_1 (i) $\langle \gamma_\zeta^\delta : \zeta < \lambda \rangle$ is strictly increasing with limit δ ,
 (ii) if $\lambda_\zeta \leq \xi < \lambda_{\zeta+1}$ then $h_0(\gamma_\xi^\delta) = h_0(\gamma_{\lambda_\zeta}^\delta) = \alpha_{\delta, \zeta}$ and $h_1(\gamma_\xi^\delta) = h_1(\gamma_{\lambda_\zeta}^\delta) = \zeta$,
 (iii) h_δ^* a partial function from κ to κ , $\text{sup}(\text{Dom}(h_\delta^*)) < \gamma_\zeta^\delta$ for $\delta \in S$;
 \odot_2 for every $f: \kappa \rightarrow \kappa, B \in [\kappa]^{<\lambda}$ and $g_\zeta^2: \kappa \rightarrow \lambda_{\zeta+1}$ for $\zeta < \sigma$ there are stationarily many $\delta \in S$ such that:
 (i) $h_\delta^* = f \upharpoonright B$,
 (ii) if $\lambda_\zeta \leq \xi < \lambda_{\zeta+1}$ then $g_\zeta^2(\gamma_\xi^\delta) = g_\zeta^2(\gamma_{\lambda_\zeta}^\delta)$.

Remark: Note that when subtraction or division* is meaningful, \odot_2 is quite strong.

Proof: By the proofs of 1.1 and 1.2 (one can use guessing clubs by $\alpha_{\delta, \zeta}$'s, and can demand that $\beta_{2\zeta}^\delta, \beta_{2\zeta+1}^\delta \in [\alpha_{\delta, \zeta}, \alpha_{\delta, \zeta} + \lambda)$).

But to help the reader, we give a proof.

Let $\lambda = \sum_{i < \sigma} \lambda_i$, λ_i increasing continuous, $\lambda_{i+1} > 2^{\lambda_i}, \lambda_0 = 0, \lambda_1 > 2^\sigma$. Let $M_i \prec (\mathcal{H}((2^\kappa)^+), \in, <^*)$ be increasing continuous, $\|M_i\| = \lambda, \langle M_j : j \leq i \rangle \in M_{i+1}, \lambda + 1 \subseteq M_i$ and $\langle \bar{A}, h_0, h_1, \bar{\lambda} \rangle \in M_0$. For $\alpha < \lambda^+$, let $\alpha = \bigcup_{i < \sigma} \alpha_{\alpha, i}$ such that $|a_{\alpha, i}| \leq \lambda_i$ and $a_{\alpha, i} \in M_{\alpha+1}$ and even $\langle a_{\beta, i} : i < \sigma \rangle: \beta \leq \alpha \in M_{\alpha+1}$. Without loss of generality $\delta \in S \Rightarrow \delta$ divisible by λ^ω (ordinal exponentiation). For $\delta \in S$ let $\bar{\beta}^\delta = \langle \beta_i^\delta : i < \sigma \rangle$ be increasing continuous with limit δ, β_i^δ divisible by λ and > 0 . For $\delta \in S$ let $\langle b_i^\delta : i < \sigma \rangle$ be such that $b_i^\delta \subseteq \beta_i^\delta, |b_i^\delta| \leq \lambda_i, b_i^\delta$ is increasing continuous in i and $\delta = \bigcup_{i < \sigma} b_i^\delta$ (e.g., $b_i^\delta = \bigcup_{j_1, j_2 < i} \alpha_{\beta_{j_1, j_2}^\delta} \cup \lambda_i$). We further demand $\lambda_i \subseteq b_i^\delta \cap \lambda$. Let $\langle f_\alpha^* : \alpha < \lambda^+ \rangle$ list the two-place functions with domain an ordinal $< \lambda^+$ and range $\subseteq \lambda^+$. Let be the set of functions h , $\text{Dom}(h) \in [\kappa]^{<\lambda}, \text{Rang}(h) \subseteq \kappa$, so $|H| = \kappa$. Let $S = \cup \{S_h : h \in H\}$, with each S_h stationary and $\langle S_h : h \in H \rangle$ pairwise disjoint. Without loss of generality $\delta \in S_h \Rightarrow \text{sup}(\text{Dom}(h)) < \beta_0^\delta$. Let h_δ^* be h when $\delta \in S_h$. We now fix $h \in H$ and choose $\bar{\gamma}^\delta = \langle \gamma_i^\delta : i < \lambda \rangle$ for $\delta \in S_h$ such that clauses $\odot_1 + \odot_2$ for our fixed h (and $\delta \in S_h$ ignoring h in \odot_2) hold; this clearly suffices.

Now for $\delta \in S_h$ and $i < \sigma$ and $g \in {}^\sigma \sigma$ we can choose $\zeta_{i, g, \varepsilon}^\delta$ (for $\varepsilon < \lambda_{i+1}$) such that:

* Namely, x_β belongs to some additive group G^* for $\beta < \kappa, \hat{g} \in \text{Hom}(G^*, H^*), g(\beta) = \hat{g}(x_\beta)$, then for some δ as in \odot_2 , we have $g(x_{\beta_\xi}^\delta - x_{\beta_{\lambda_\zeta}^\delta})$ is 0_{H^*} ; similarly for multiplicative groups.

- (A) $\langle \zeta_{i,g,\varepsilon}^\delta : \varepsilon < \lambda_{i+1} \rangle$ is a strictly increasing sequence of ordinals,
 (B) $\beta_i^\delta < \zeta_{i,g,\varepsilon}^\delta < \beta_{i+1}^\delta$ (we can even demand $\zeta_{i,j,\varepsilon}^\delta < \beta_i^\delta + \lambda$),
 (C) $h_0(\zeta_{i,g,\varepsilon}^\delta) = \alpha_{\delta,i}$ and $h_1(\zeta_{i,g,\varepsilon}^\delta) = i$,
 (D) for** every $\alpha_1, \alpha_2 \in b_{g(i)}^\delta$, the sequence $\langle \text{Min}\{\lambda_{g(i)}, f_{\alpha_1}^*(\alpha_2, \zeta_{i,g,\varepsilon}^\delta) : \varepsilon < \lambda_{i+1}\} \rangle$ is constant, i.e., one of the following occurs:
 (α) $\varepsilon < \lambda_{i+1} \Rightarrow (\alpha_2, \zeta_{i,g,\varepsilon}^\delta) \notin \text{Dom}(f_{\alpha_1}^*)$,
 (β) $\varepsilon < \lambda_{i+1} \Rightarrow f_{\alpha_1}^*(\alpha_2, \zeta_{i,g,\varepsilon}^\delta) = f_{\alpha_1}^*(\alpha_2, \zeta_{i,j,0}^\delta)$ well defined,
 (γ) $\varepsilon < \lambda_j, f_{\alpha_1}^*(\alpha_2, \zeta_{i,g,\varepsilon}^\delta) \geq \lambda_j$, well defined. We can add $\langle f_{\alpha_1}^*(\alpha_2, \zeta_{i,g,\varepsilon}^\delta) : \varepsilon < \lambda_i \rangle$ is constant or strictly increasing,
 (E) for some $j < \sigma$, we have $(\forall \varepsilon < \lambda_{i+1})[\zeta_{i,g,\varepsilon}^\delta \in a_{\alpha,j}]$ where
 $\alpha = \sup\{\zeta_{i,g,\varepsilon}^\delta : \varepsilon < \lambda_{i+1}\}$ (remember $\sigma \neq \lambda_{i+1}$ are regular).

For each function $g \in {}^\sigma \sigma$ we try $\bar{\gamma}^{g,\delta} = \langle \gamma_{\varepsilon}^{\delta,g} : \varepsilon < \lambda \rangle$ if $\lambda_i \leq \varepsilon < \lambda_{i+1}$ then $\gamma_{\alpha}^{\delta,g} = \zeta_{i,g,\varepsilon}^\delta$. Now for some g it works. ■_{4.2}

Proof of 1.2(1): Let $M = \cup\{M_\alpha : \alpha < \kappa\}$, $M_\alpha \prec (\mathcal{H}(2^\kappa)^+, \in)$ has cardinality λ , M_α is increasing continuous, $\langle M_\beta : \beta \leq \alpha \rangle \in M_\alpha$ and $\langle F_i : i < \sigma \rangle$ belongs to M_0 . Let $E_0 = \{\delta < \kappa : M_\delta \cap \kappa = \delta\}$ and $E = \text{acc}(E)$. The proof is like the proof of 4.2 with the following changes:

- (i) $\beta_i^\delta \in E_0$ for $\delta \in S \cap E$,
 (ii) in clause (A) we demand $\langle \zeta_{i,g,\varepsilon}^\delta : g \in G, \varepsilon < \lambda_{i+1} \rangle$ belongs to $M_{\beta_{i+1}^\delta}$ (hence also $\langle \zeta_{j,g,\varepsilon}^\delta : g \in G, \varepsilon < \lambda_{j+1} : j \leq i \rangle$ belongs to $M_{\beta_{i+1}^\delta}$),
 (iii) clause (c) is replaced by: $\zeta_{i,g,\varepsilon}^\delta \in F_i(\{\zeta_{j,g \upharpoonright (j+1), \varepsilon}^\delta : \varepsilon < \lambda_{j+1} \text{ and } j < i\})$.

■_{1.2}

Proof of 4.1: (1) We apply 4.2 to the $\langle A_\delta : \delta \in S \rangle$ from 4.1, and any h_0, h_1 as in clause (d) of 4.2.

Let $\{t_\gamma^{i,j} + G_i : \gamma < \theta^{i,j}\}$ be a free basis of G^j/G^i for $i < j \leq \sigma$. If $i = 0, j = \sigma$ we may omit the i, j , i.e., $t_\zeta = t_\zeta^{0,\sigma}$ and $\theta = \theta^{0,\sigma}$. Let $\theta + \aleph_0 = |G_\sigma| < \lambda$; actually $\theta^{\zeta, \zeta+1} < \lambda_\zeta$ is enough; without loss of generality $\theta < \lambda_1$ in 4.2. Let $\beta_{\zeta,i}^\delta = \gamma_{\xi(\zeta,i)}^\delta$ where $\xi(\zeta, i) = \bigcup_{\varepsilon < \zeta} \lambda_\varepsilon + 1 + i$ for $\delta \in S, \zeta < \sigma, i < \theta$.

Let $\beta_\delta(*) = \text{Min}\{\beta : \beta \in \text{Dom}(h_\delta^*), h_\delta^*(\beta) \neq 0\}$, if well defined where h_δ^* is from 4.2.

Clearly (see \odot_1 (iii) of 4.2) we have $\beta_\delta(*) \notin \{\beta_{\zeta,i}^\delta : \zeta < \sigma, i < \theta\}$ (or omit $\lambda_\zeta, \beta_{\zeta,i}^\delta$ for ζ too small). We define an abelian group G^* : it is generated by $\{x_\alpha : \alpha < \kappa\} \cup \{y_\gamma^\delta : \gamma < \theta \text{ and } \delta \in S\}$ freely except for the relations:

$$(*)_1 \quad \sum_{\gamma < \theta} a_\gamma y_\gamma^\delta = \sum \{b_{\zeta,\gamma} (x_{\beta_{\zeta,\gamma}^\delta} - x_{\gamma_{\lambda_\zeta}^\delta}) : \zeta < \sigma \text{ and } \gamma < \theta^{\zeta, \zeta+1}\}$$

when $G_\sigma \models \sum_{\gamma < \theta^{0,\sigma}} a_\gamma t_\gamma = \sum \{b_{\zeta,\gamma} t_\gamma^{\zeta, \zeta+1} : \zeta < \sigma \text{ and } \gamma < \theta^{\zeta, \zeta+1}\}$ where

** We can use a colouring which uses, e.g., $\langle \zeta_{j,g,\varepsilon}^\delta : j < i, \varepsilon < \lambda_{j+1} \rangle$ as a parameter.

$a_\gamma, b_{\zeta, \gamma} \in \mathbb{Z}$ but all except finitely many are zero.

There is a (unique) homomorphism \mathbf{g}_δ from G_σ into G^* induced by $\mathbf{g}_\delta(t_\gamma) = y_\gamma^\delta$. As usual it is an embedding. Let $\text{Rang}(\mathbf{g}_\delta) = G^{<\delta>}$.

For $\beta < \kappa$ let G_β^* be the subgroup of G^* generated by

$$\{x_\alpha : \alpha < \beta\} \cup \{y_\gamma^\delta : \gamma < \theta^{0, \sigma} \text{ and } \delta \in \beta \cap S\}.$$

It can be described similarly to G^* .

Fact A: G^* is strongly λ -free.

Proof: For $\alpha^* < \beta^* < \kappa$, we can find $\langle \alpha_\delta : \delta \in S \cap (\alpha^*, \beta^*] \rangle$ such that $\langle A_\delta \setminus \alpha_\delta : \delta \in S \cap (\alpha^*, \beta^*] \rangle$ are pairwise disjoint and disjoint to α^* hence the sequence $\langle \{\beta_{\zeta, i}^\delta : i < \theta, \zeta \in (\text{Min}\{\xi < \sigma : \beta_{\zeta, 0}^\delta > \alpha_\delta\}, \sigma)\} : \delta \in S \cap (\alpha^*, \beta^*] \rangle$ is a sequence of pairwise disjoint sets.

For $\delta \in S \cap (\alpha^*, \beta^*]$, let $\zeta_\delta = \text{Min}\{\zeta : \beta_{\zeta, 0}^\delta > \alpha_\delta\} < \sigma$. Now easily $G_{\beta^*+1}^*$ is generated as an extension of $G_{\alpha^*+1}^*$ by $\{\mathbf{g}_\delta(t_\gamma^{\zeta_\delta, \sigma}) : \gamma < \theta^{\zeta_\delta, \sigma} \text{ and } \delta \in S \cap (\alpha^*, \beta^*] \cup \{x_\alpha : \alpha \in (\alpha^*, \beta^*] \text{ and for no } \delta \in S \cap (\alpha^*, \beta^*] \text{ do we have } \alpha \in \{\beta_{\zeta, i}^\delta : i < \theta^{\zeta, \sigma} \text{ and } \zeta < \zeta_\delta\}\}$; moreover, $G_{\beta^*+1}^*$ is freely generated (as an extension of $G_{\alpha^*+1}^*$). So $G_{\beta^*+1}^*/G_{\alpha^*+1}^*$ is free; as also G_1^* is free, we have shown Fact A.

Fact B: G^* is not Whitehead.

Proof: We choose, by induction on $\alpha \leq \kappa$, an abelian group H_α and a homomorphism $\mathbf{h}_\alpha : H_\alpha \rightarrow G_\alpha^* = \langle \{x_\beta : \beta < \alpha\} \cup \{y_\gamma^\delta : \gamma < \theta, \delta \in S \cap \alpha\} \rangle_{G^*}$ increasing continuous in α , with kernel \mathbb{Z} , $\mathbf{h}_0 = \text{zero}$ and $\mathbf{k}_\alpha : G_\alpha^* \rightarrow H_\alpha$ is a not necessarily linear mapping such that $\mathbf{h}_\alpha \circ \mathbf{k}_\alpha = \text{id}_{G_\alpha^*}$. We identify the set of members of $H_\alpha, G_\alpha, \mathbb{Z}$ with subsets of $\lambda \times (1 + \alpha)$ such that $O_{H_\alpha} = O_{\mathbb{Z}} = 0$.

Usually we have no freedom or no interesting freedom. But we have for $\alpha = \delta + 1, \delta \in S$. What we demand is $(G^{<\delta>})$ — see before Fact A):

(*)₂ letting $H^{<\delta>} = \{x \in H_{\delta+1} : \mathbf{h}_{\delta+1}(x) \in G^{<\delta>}\}$, if $s^* = g_\delta(x_{\beta_\delta^*}) \in \mathbb{Z} \setminus \{0\}$ (g_δ from 4.2), then there is no homomorphism $f_\delta : G^{<\delta>} \rightarrow H^{<\delta>}$ such that

$$(\alpha) f_\delta(x_{\beta_{\zeta, i}^\delta}) - \mathbf{k}_\delta(x_{\beta_{\zeta, i}^\delta}) \in \mathbb{Z} \text{ is the same for all } i \in (\bigcup_{\varepsilon < \zeta} \lambda_\varepsilon, \lambda_\zeta]$$

$$(\beta) \mathbf{h}_{\delta+1} \circ f_\delta = \text{id}_{G^{<\delta>}}.$$

[Why is this possible? By non-Whiteheadness of $G^\sigma / \bigcup_{i < \sigma} G^i$, that is, see (d)(γ)⁻ in 4.1.]

The rest should be clear.

Proof of 4.1(2): Of course, similar to that of 4.1(1) but with some changes.

Step A: Without loss of generality there is a homomorphism f^* from $\bigcup_{i < \sigma} G^i$ to \mathbb{Z} which cannot be extended to a homomorphism from G_σ to \mathbb{Z} .

[Why? Standard, see [Fu].]

Step B: During the construction of G^* , we choose G_α^* by induction on $\alpha \leq \kappa$, but if $h_\delta^*(0)$ from 4.2 is a member of G_δ^* in $(*)_1$ we replace $(x_{\beta_{\zeta,\gamma}^\delta} - x_{\gamma_{\lambda_\zeta}^\delta})$ by $(x_{\beta_{\zeta,\gamma}^\delta} - x_{\beta_{\lambda_\zeta}^\delta} + f^*(t_\gamma^{\zeta,\zeta+1})h_\delta^*(0))$; note that $f^*(t_\gamma^{\zeta,\zeta+1}) \in \mathbb{Z}$ and $h_\delta^*(0) \in G_\delta^*$.

So if in the end $f: G^* \rightarrow \mathbb{Z}$ is a non-zero homomorphism, let $x^* \in G^*$ be such that $f(x^*) \neq 0$ and $|f^*(x^*)|$ is minimal under this, so without loss of generality it is 1. Hence for some $\delta \in S$ we have:

$$(*)_3 \quad f(g_\delta(0)) = 1_{\mathbb{Z}},$$

$$(*)_4 \quad f(x_{\gamma_{\lambda_{\zeta+1+\gamma}}^\delta}) = f(x_{\gamma_{\lambda_\zeta}^\delta}) \text{ for } \gamma \in \lambda_{\zeta+1} \setminus \lambda_\zeta,$$

that is, $f(x_{\beta_{\zeta,\gamma}^\delta}) = f(x_{\gamma_{\lambda_\zeta}^\delta})$

(in fact, this holds for stationarily many ordinals $\delta \in S$).

So we get an easy contradiction.

(3) The proof is included in the proof of part (2). ■_{4.1}

We also note the following consequence of a conclusion of an instance of GCH.

4.3 CLAIM: Assume

$$(a) \quad \lambda = \mu^+ \text{ and } \mu > \sigma = \text{cf}(\mu),$$

$$(b) \quad \lambda = \lambda^\theta \text{ where } \theta = 2^\sigma$$

$$(\text{equivalently } \mu^\theta = \mu^+ > 2^\theta),$$

$$(c) \quad S \subseteq \{\delta < \lambda : \text{cf}(\delta) = \sigma\} \text{ is stationary,}$$

$$(d) \quad \bar{\eta} = \langle \eta_\delta : \delta \in S \rangle \text{ with } \eta_\delta \text{ an increasing sequence of length } \sigma \text{ with limit } \delta.$$

Then we can find $\langle \bar{A}^\delta : \delta \in S \rangle$ such that:

$$(\alpha) \quad \bar{A}^\delta = \langle A_i^\delta : i < \sigma \rangle,$$

$$(\beta) \quad A_i^\delta \in [\delta]^{< \mu} \text{ and } \sup(A_i^\delta) < \delta,$$

$$(\beta)^+ \text{ for some } \langle \lambda_i^* : i < \sigma \rangle \text{ increasing with limit } \lambda, |A_i^\delta| < \lambda_i^*,$$

$$(\gamma) \text{ for every } h: \lambda \rightarrow \lambda, \text{ for stationarily many } \delta \in S \text{ we have } (\forall i < \sigma) [h(\eta_\delta(i)) \in A_i^\delta].$$

4.4 Remark: (1) We can restrict ourselves to $h: \lambda \rightarrow \mu$ in clause (γ) , and then, of course, we can use $\langle \langle A_i^\delta : i < \sigma \rangle : \delta \in S \rangle$ with $A_i^\delta \subseteq \mu$.

(2) We can add to the conclusion " $A_i^\delta \subseteq \eta_\delta(i+1)$ " if $\bar{\eta}$ guess clubs.

Proof: Let $\langle \lambda_i : i < \sigma \rangle$ be increasing continuous with limit μ . Let $\langle \bar{\alpha}_\gamma : \gamma < \lambda \rangle$ list ${}^\theta \lambda$, so $\bar{\alpha}_\gamma = \langle \alpha_{\gamma,\varepsilon} : \varepsilon < \theta \rangle$ and, without loss of generality, $\alpha_{\gamma,\varepsilon} \leq \gamma$. For each $\delta \in S$ let $\langle b_i^\delta : i < \sigma \rangle$ be an increasing continuous sequence of subsets of δ

* What does this mean? $f^*(x^*)$ is an integer, so its absolute value is well defined.

with union δ such that $|b_i^\delta| < \mu$ and $\sup(b_i^\delta) < \delta$; for $(\beta)^+$, moreover, $|b_i^\delta| \leq \lambda_i$; this is possible as $\text{cf}(\delta) = \sigma = \text{cf}(\mu) < \mu$. Let $\langle g_\varepsilon : \varepsilon < \theta \rangle$ list ${}^\sigma\sigma$ and define $A_i^{\varepsilon, \delta} = \{\alpha_{\gamma, \varepsilon} : \gamma \in b_{g_\varepsilon(i)}^\delta\}$. Now $A_i^{\varepsilon, \delta}$ is a set of cardinality $\leq |b_{g_\varepsilon(i)}^\delta| < \mu$ and $\sup(A_i^{\varepsilon, \delta}) \leq \sup(b_{g_\varepsilon(i)}^\delta)$ (as we have demanded that $\alpha_{\gamma, \varepsilon} \leq \gamma$), but $\sup(b_{g_\varepsilon(i)}^\delta) < \delta$ by the choice of the b_j^δ 's, hence $\sup(A_i^{\varepsilon, \delta}) < \delta$. So for each $\varepsilon < \theta$ the sequence $\bar{\mathbf{A}}^\varepsilon = \langle \bar{A}^{\varepsilon, \delta} : \delta \in S \rangle$, where $\bar{A}^{\varepsilon, \delta} = \langle A_i^{\varepsilon, \delta} : i < \sigma \rangle$ satisfies clauses $(\alpha) + (\beta)$ and $(\beta)^+$ when relevant. Hence it suffices to prove that for some $\varepsilon < \theta$ the sequence $\bar{\mathbf{A}}^\varepsilon$ satisfies clause (γ) , too. Assume toward a contradiction that for every $\varepsilon < \theta$ the sequence $\bar{\mathbf{A}}^\varepsilon$ fails clause (γ) , hence there is $h_\varepsilon : \lambda \rightarrow \lambda$ which exemplifies this, that is, for some club E_ε of λ , $\delta \in E_\varepsilon \cap S \Rightarrow (\exists i < \sigma)[h_\varepsilon(\eta_\delta(i)) \notin A_i^{\varepsilon, \delta}]$. So for every $\beta < \lambda$ the sequence $\langle h_\varepsilon(\beta) : \varepsilon < \theta \rangle$ belongs to ${}^\theta\lambda$, hence is equal to $\bar{\alpha}_{h(\beta)}$ for some $h(\beta) < \lambda$. Clearly $E = \{\delta < \lambda : \delta \text{ a limit ordinal and } (\forall \beta < \delta) h(\beta) < \delta\}$ is a club of λ (recall $\theta < \lambda$), hence we can find $\delta(*) \in E \cap S$. We define $g^* : \sigma \rightarrow \sigma$ by $g^*(i) = \text{Min}\{j < \sigma : h(\eta_{\delta(*)}(j)) \in b_j^\delta\}$, now g^* is well defined as, for $i < \sigma$, the ordinal $h(\eta_{\delta(*)}(i))$ is $< \delta(*)$ (as $\delta(*) \in E$) and $\eta_{\delta(*)}(i) < \delta(*)$ and $\delta = \bigcup_{j < \sigma} b_j^\delta$. As $g^* \in {}^\sigma\sigma$, clearly for some $\varepsilon(*) < \theta$ we have $g_{\varepsilon(*)} = g^*$.

So, for any $i < \sigma$, let $\gamma_i = h(\eta_{\delta(*)}(i))$; now $h(\eta_{\delta(*)}(i)) \in b_{g^*(i)}^\delta$ (by the choice of g^*) and $g^*(i) = g_{\varepsilon(*)}(i)$ by the choice of $\varepsilon(*)$, together with $\gamma_i \in b_{g_{\varepsilon(*)}(i)}^\delta$. But $A_i^{\varepsilon(*), \delta(*)} = \{\alpha_{\gamma, \varepsilon(*)} : \gamma \in b_{g_{\varepsilon(*)}(i)}^\delta\}$ by the choice of $A_i^{\varepsilon(*), \delta(*)}$, hence $\alpha_{\gamma_i, \varepsilon(*)} \in A_i^{\varepsilon(*), \delta(*)}$; but as $\gamma_i = h(\eta_{\delta(*)}(i))$, by the choice of h we have $h_{\varepsilon(*)}(\eta_{\delta(*)}(i)) = \alpha_{\gamma_i, \varepsilon(*)} \in A_i^{\varepsilon(*), \delta(*)}$.

So $(\forall i < \sigma)(h_{\varepsilon(*)}(\eta_{\delta(*)}(i)) \in A_i^{\varepsilon(*), \delta(*)})$ which, by the choice of $h_{\varepsilon(*)}$, implies $\delta(*) \notin E_{\varepsilon(*)}$, but $\delta(*) \in E \subseteq \bigcap_{\varepsilon < \theta} E_\varepsilon$, a contradiction. $\blacksquare_{4.3}$

References*

- [Fu] L. Fuchs, *Infinite Abelian Groups*, I, II, Academic Press, New York, 1970, 1973.
- [Sh 587] S. Shelah, *Not collapsing cardinals $\leq \kappa$ in $(< \kappa)$ -support iterations*, Israel Journal of Mathematics, to appear, math.LO/9707225.
- [Sh 186] S. Shelah, *Diamonds, uniformization*, The Journal of Symbolic Logic **49** (1984), 1022–1033.
- [Sh f] S. Shelah, *Proper and Improper Forcing*, Perspectives in Mathematical Logic, Springer, New York, 1998.

* References of the form math.XX/... refer to the xxx.lanl.gov archive.