

LARGE CARDINALS IMPLY THAT EVERY REASONABLY DEFINABLE SET OF REALS IS LEBESGUE MEASURABLE

BY

SAHARON SHELAH^{†,a,c} AND HUGH WOODIN^{††,b,c}^a*Institute of Mathematics, The Hebrew University of Jerusalem, Jerusalem, Israel;*^b*Department of Mathematics, California Institute of Technology, Pasadena, California, USA;**and* ^c*Department of Mathematics, Rutgers University, New Brunswick, New Jersey, USA*

ABSTRACT

We prove that if there is a supercompact cardinal or much smaller large cardinals, then every set of reals from $L(\mathbf{R})$ is Lebesgue measurable, and similar results. We also introduce some large cardinals.

§1. Introduction

In the old days Solovay proved the, by now classical, result:

1.1. THEOREM. *If $V^1 = V^{\text{Levi}(\aleph_\omega < \kappa)}$ (see below), κ strongly inaccessible, then V^1 satisfies:*

- (a) *every set of reals from $L(\mathbf{R})$ is Lebesgue measurable,*
- (b) *every set of reals from $L(\mathbf{R})$ has the property of Baire,*
- (c) *in $L(\mathbf{R})$ there is no well ordering of the reals,*
- (d) *in $L(\mathbf{R}) \models \omega \rightarrow (\omega)_2^\omega$ (added by Mathias [Mt])*

(and on classical results see Solovay [So]).

For so long some of the central set theorists hoped to prove measurability, etc., of definable sets from large cardinals, and then also the stronger conclusion — *AD*. See Moschovakis [M1] for classical results.

[†] The first author thanks the United States–Israel Binational Science Foundation for partially supporting this research, M. Gitik, J. I. Ihoda and Y. Koppelvich for corrections and Alice Leonhardt for the beautiful typing. Publication no. 241.

^{††} The second author thanks the National Science Foundation for supporting this research.

Received December 21, 1986 and in final revised form September 12, 1989

Magidor [Mg] proved: if there is a measurable cardinal and a precipitous ideal on ω_1 , then (a), (b), (c) of 1.1 hold for Σ_3^1 and, from stronger assumptions, Σ_4^1 (precipitous ideals on $\mathcal{P}_{<\aleph_1}(\aleph_1)$, $\mathcal{P}_{<\aleph_1}(\aleph_3)$). Woodin [W1] proved that “in $L(\mathbf{R})$ ” every set of reals is Lebesgue measurable, has the Baire property and is not a well ordering of the reals (1.3 (a), (b), (c) for short) by proving $L(\mathbf{R}) \models$ “ AD ” assuming there is an elementary embedding $j: L(V_{\lambda+1}) \rightarrow L(V_{\lambda+1})$ with critical point $< \lambda$. This work starts from the thesis that Martin’s proof of Π_2^1 determinacy is the right proof.

It did not affect much the theology of the Los Angeles school of descriptive set theorists, by which:

- (A) Determinacy has nothing to do with large cardinals [inconsistency of supercompact? “I have no understanding/interest/stake in them”].
- (B) Determinacy is much stronger than all known large cardinals.
It also fits well with the theology of general set theorists by which:
- (C) Every interesting statement is equi-consistent with some large cardinal.
- (D) A consistency result means one from a large cardinal, as those are at least half way to being axioms, not mere statements, hence the experimental observation that they are linearly ordered.

The large majority look at consistency results from AD as implications.

But by the L.A. school

- (E) AD is much more reliable, having inner models.

Anyhow everybody agrees with:

- (F) Elementary embeddings come only from large cardinals.

By this theology “there is an \aleph_2 -saturated normal ideal on ω_1 ” should be equi-consistent with almost huge — one side was known, the other waits for the inner model.

There was no change by results of Foreman [F], which got results on measurability, etc. from “potent axioms” (saying “large filter” exists on small cardinals), as we do not know much on their consistency strength. This picture was shattered when in Spring 1984 Foreman, Magidor and Shelah [FMS1, §1] was written. From this paper’s point of view the important thing was that (relying on [Sh1]):

1.2. THEOREM. *Suppose κ is a supercompact; we can force, without collapsing \aleph_1 , “ D_{ω_1} is \aleph_2 -saturated”; there is κ -c.c., semi-proper forcing notion P forcing this.*

So some generic elementary embeddings in small cardinals are novel creatures, not extending some old one for large cardinals in an inner model.

In [FMS1]

1.3. THEOREM. *If $MA^+(\aleph_1\text{-complete})$ (see 1.12 below) then any forcing notion preserving stationarity of a subset of ω_1 is semi-proper.*

Using the sealing forcing of maximal antichains the following theorem was proved:

1.4. THEOREM. *If $MA^+(\aleph_1\text{-complete})$ then D_{ω_1} (the club filter on ω_1) is precipitous and even semi-proper, subject to the following definition:*

1.5. DEFINITION. Let D be a fine normal filter on $I \subseteq \mathcal{P}(\lambda)$ (our main interest in $I = \omega_1$ or $I \subseteq \{a \subseteq \lambda : |a| < \aleph_1\}$, so we assume it in (2) below).

(1) We say $\{A_i : i < i(*)\}$ (or $\bar{A} = \langle A_i : i < i(*) \rangle$) is a maximal antichain (for D) if $A_i \subseteq I$, $A_i \neq \emptyset \pmod{D}$, $[i \neq j \Rightarrow A_i \cap A_j = \emptyset \pmod{D}]$, and $[A \subseteq I \wedge \bigwedge_{i < i(*)} A \cap A_i = \emptyset \pmod{D} \Rightarrow A = \emptyset \pmod{D}]$.

(2) \bar{A} is semi-proper if: for every large enough cardinal χ and countable $N < (H(\chi), \in)$, such that $\bar{A} \in N$, $N \cap \lambda \in I$ there is a countable N_1 , $N < N_1 < (H(\chi), \in)$, $N \cap \lambda = N_1 \cap \lambda$ such that $(\exists i < \text{lg}(\bar{A})) [N_1 \cap \lambda \in A_i \wedge i \in N_1]$ (equivalently $\text{seal}(\bar{A})$ is semi-proper forcing notion; see proof of 2.4).

(3) D is semi-proper if every maximal antichain is.

1.6. REMARK. When we are interested in uncountable models, the natural restrictions on N , N_1 are more complicated; see [FMS1].

Much parallel activity follows. Foreman, Magidor and Shelah [FMS1], [FMS2] contain some further related theorems, e.g. non- (λ, λ^+) -regular uniform ultrafilters on λ^+ ($\lambda \geq \aleph_0$ regular) were constructed. [This raises a theological question — who first proved its consistency? See (D), (E) above.] See also the present paper and [Sh2], [Sh3] and [Sh4].

Our result here is

1.7. THEOREM. (1) *If there is a supercompact cardinal, then $L(\mathbf{R})$ satisfies (a), (b), (c) and (d) of 1.1 holds (in fact \bigoplus_κ holds — see below).*

(2) *In fact much lower cardinals suffice.*

This work was continued by Woodin [W2] who proved (under similar assumptions) that every set in $L(\mathbf{R})$ is represented by a weakly homogeneous tree.

Then Martin and Steel [MS] proved that every set of reals represented by sufficiently weakly homogeneous trees (with measures κ^+ -complete) is determined if κ satisfies Definition 4.3, thus deducing $L(\mathbf{R}) \models AD$ from much less

than supercompact (relying on the result of [W2] mentioned above for sets in $L(\mathbf{R})$).

This paper is written in the generic way — repeating the steps of its conception, as we feel this has interest *per se*, and as anyhow “everybody” knows the proofs. Naturally there is some overlapping with other works of Spring 1984 in 2.4 with [FMS1], and in 3.4 with [Sh3].

We define here two large cardinals: $\text{Pr}_a(\lambda, f)$, $\text{Pr}_a(\lambda)$ by Shelah (Definition 3.5) and $\text{Pr}_b(\lambda)$ by Woodin — now called a Woodin cardinal.

We also used freely facts which are well known.

Let us mention

1.8. DEFINITION. $\text{Levi}(\lambda, < \mu) = \{f: f \text{ a two-place function from ordinals to ordinals, } \text{Dom}(f) \text{ has cardinality } < \lambda \text{ and } [f(\alpha, \beta) = \gamma \Rightarrow \alpha < \mu \wedge \beta < \lambda \wedge \gamma < \alpha]\}$.

1.9. DEFINITION. \bigoplus_κ (for a universe V) will be the statement: in $V^1 = V^{\text{Levi}(\aleph_\nu, < \kappa)}$ there is an elementary embedding $J: L(\mathbf{R}^V) \rightarrow L(\mathbf{R}^{V^1})$ (you can add “with a class of ordinals $\alpha, j(\alpha) = \alpha$ ”, for this paper).

A rephrasing (essentially) of 1.1 is

1.10. COROLLARY. *If $\bigoplus_\kappa, \kappa > \aleph_1$ then (a) + (b) + (c) + (d) of 1.1 holds (i.e. for $L(\mathbf{R}^V)$).*

1.11. DEFINITION. (1) $MA_\alpha(\text{Pr})$ means: *if P is a forcing notion satisfying Pr, D_i a dense subset of P for $i < \omega_1, \mathcal{S}_\beta$ a P -name of a stationary subset of ω_1 for $\beta < \alpha$, then there is a directed $G \subseteq P$ such that: $G \cap D_i \neq \emptyset$ for $i < \omega_1$, and $\{\zeta < \omega_1: (\exists p \in G)p \Vdash “i \in \mathcal{S}_\beta”\}$ is stationary for $\beta < \alpha$.*

(2) Let $MA^+(\text{Pr})$ be $MA_1(\text{Pr})$.

$P \triangleleft Q$ means complete subforcing, i.e. $P \subseteq Q$ and every maximal antichain of P is a maximal antichain of Q .

§2. The dawn

A conclusion to 1.2, using weakly homogeneous trees, was

2.1. THEOREM. *If κ is supercompact, then every projective set is Lebesgue measurable and with the property of Baire.*

The second step was

2.2. CONCLUSION. If κ is supercompact then \bigoplus_κ (hence, by 1.11, every set in $L(\mathbf{R})$ is Lebesgue measurable, etc.).

This was obtained by putting together the following simultaneously obtained results:

2.3. THEOREM. *If κ is weakly compact, and*
 $\textcircled{a}_\kappa^{\text{id}}$: *for some κ -c.c. forcing notion P of power κ not adding reals, \Vdash_P "there is a normal \aleph_2 -saturated ideal I on ω_1 "*
then \bigoplus_κ holds.

2.4. LEMMA. *For κ -supercompact and $S \subseteq \omega_1$ stationary, co-stationary:*

\textcircled{a}^S : *there is a κ -c.c. forcing notion P of power, κ not adding reals \Vdash_P " $D_{\omega_1} + S$ is \aleph_2 -saturated".*

PROOF OF 2.3. In $V^1 = V^P$ let $Q = \{A \subseteq \omega_1 : A \notin I\}$ ordered by inverse inclusion, so Q is \aleph_2 -c.c., i.e. κ -c.c., hence in V , $R \stackrel{\text{def}}{=} P * Q$ is κ -c.c. Let $R = \bigcup_{i < \kappa} R_i$, R_i increasing continuous, $|R_i| < \kappa$. As κ is weakly compact and R is κ -c.c., $W \stackrel{\text{def}}{=} \{\lambda < \kappa : \lambda \text{ regular, } R_\lambda \triangleleft R, R_\lambda \text{ satisfies } \lambda\text{-c.c.}\}$ is stationary, so every real in V^R is in V^{R_λ} for some λ . Now we shall show: In V^R we can rearrange R as $\text{Levi}(\aleph_0, < \kappa)$: let $S = \{F : F \text{ an isomorphism from } R_\lambda (\lambda \in W) \text{ onto } \text{Levi}(\aleph_0, < \lambda)\}$, ordered by inclusion. S as a forcing notion adds no real as W is stationary, so $\mathbf{R}(V^R) = \mathbf{R}(V^{(R \ast S)})$, but the latter is $\mathbf{R}(V^{\text{Levi}(\aleph_0, < \kappa)})$. Now in $V^R = V^{P * Q}$, \mathcal{G}_Q is a normal (generic) ultrafilter on $\mathcal{P}(\omega_1)^{V^P} = \mathcal{P}(\omega_1)^{V^1}$ so the usual ultrapower $(V^1)^{\omega_1 / \mathcal{G}_Q} = M$ (by Mostowski collapse) is defined, as well as the embedding $j : V^1 \rightarrow M$. As I is \aleph_2 -saturated (in V^P) every Q -name \check{r} of a real (also ordinal) can be represented by $f : \omega_1 \rightarrow \omega : \check{r}(n) = l \Leftrightarrow f^{-1}(\{l\}) \in \mathcal{G}_Q$; hence in V^R , $\mathbf{R}(V^R) = \mathbf{R}(M)$. So $j \upharpoonright L(\mathbf{R}^V)$ exemplify \bigoplus_κ remembering $\mathbf{R}(V^P) = \mathbf{R}(V)$.

PROOF OF 2.4. We use RCS iteration $\bar{Q} = \langle P_i, \check{Q}_j : i \leq \kappa, j < \kappa \rangle$. For each i , if there is in V^{P_i} a maximal antichain $A = \langle A_\alpha^i : \alpha < \alpha^*(*) \rangle$, $|\alpha^*(*)| > \aleph_1$, of $D_{\omega_1} + S$ (i.e. $\omega_1 - S \subseteq A_\alpha^i \neq \emptyset \pmod{D_{\omega_1} + S}$, $A_\alpha^i \cap A_\beta^i = \emptyset \pmod{D_{\omega_1} + S}$) and

$$\text{seal}(\bar{A}) = \{(f, C) : f \text{ a function from some countable ordinal } \alpha + 1 \text{ to } \alpha^*(*), C \text{ a closed subset of } \alpha + 1, (\forall \delta \in C \cap S) \delta \in \bigcup_{i < \delta} A_{f(i)}\},$$

order is defined by:

$$(f_1, C_1) \leq (f_2, C_2) \Leftrightarrow f_1 \subseteq f_2 \wedge (C_1 = C_2 \cap \text{Dom } f_1)$$

is semi-proper and Laver's diamond ([L]) gives for i a P_i -name of \bar{A} , then choose Q_i as such a seal(\bar{A}). Otherwise

$$Q_i = \text{Levi}(\aleph_1, (2^{\aleph_1})^{V^{P_i}}).$$

By the $(\omega_1 - S)$ -completeness of the Q_i 's, reals are not added (by [Sh1], Ch. V).

By the semi-proper iteration lemma (see [Sh1], Ch. X) S is stationary in V^P and also we know P_κ satisfies the κ -c.c. Now suppose $p \Vdash_P \langle A_i : i < \kappa \rangle$ is a maximal antichain for $D_{\omega_1} + S$; as we have used Laver's diamond we have $V^P \models MA^+$ (\aleph_1 -complete forcing), hence $V^P \models$ "a forcing is semi-proper iff it does not destroy stationary subsets of ω_1 ", but $Q_{\bar{A}}$ destroys the stationary of no subset of ω_1 . By the reflection properties of κ , we could have chosen Q_i as $Q_{\bar{A}, i}$ for many i 's; as \bar{Q} was chosen by Laver's diamond we have done it, and then seal(\bar{A}) makes the diagonal union of A_j ($j < i$) contain a club, contradicting $A_i \neq \emptyset \pmod{D_{\omega_1} + S}$.

2.5. REMARK. Everybody knows that we can replace in 2.4 P by $\text{Levi}(\aleph_1, < \kappa)$ and get an \aleph_2 -saturated normal ideal on ω_1 (as $\text{Levi}(\aleph_1, < \kappa)$ is embeddable into P by a \triangleleft -embedding; equivalently, in V^P we can find a directed subset of $\text{Levi}(\aleph_1, < \kappa)^V$ generic over V).

§3. The dusty road

We may wonder what is the right cardinal in 2.2. Analyzing the proof of $MA^+(\aleph_1\text{-complete})$ in the case needed in 2.4:

3.1. THEOREM. *In 2.4 and 2.2 we may replace supercompactness by superstrong:*

there is an elementary embedding $j : V \rightarrow M$, κ the critical ordinal of j , and $H(j(\kappa)) \subseteq M$ (M a transitive class).

3.1A. CLAIM. In 2.3 we can replace the use of an \aleph_2 -saturated ideal by the use of a presaturated one.

In the next stage one may wonder whether we can economize in 2.1, and hopefully find a proof using n cardinals for sets of reals defined by $\Sigma_{n+n_0}^1$ formulas.

3.2. LEMMA. (1) *Suppose $P_0 \triangleleft P_1 \triangleleft \dots \triangleleft P_n$, P_l a forcing notion adding no reals, P_l not destroying stationary subsets of ω_1 from V or V^{P_m} for $m < l$, and $S \subseteq \omega_1$ stationary, $\Vdash_{P_l} "D_{\omega_1} + S \text{ is } \aleph_2\text{-saturated}"$, $\Vdash_{P_l} "\kappa_l = \aleph_2"$, $|P_l| < \aleph_{l+1}$, and κ_l is strongly inaccessible.*

Then every Σ_{n+2}^1 set of reals is Lebesgue measurable, Ramsey and has the property of Baire.

(2) Suppose we have such $S, P_l (l \leq \omega)$ then $L(\mathbf{R})$ satisfies (a), (b), (c) and (d) of 1.1.

But when does this occur?

3.3. DEFINITION. (1) Let $(*)_{\kappa}^+$ mean:

$$\| \text{Levi}(\aleph_1, < \kappa) \| \text{ “} D_{\omega_1} \text{ is semi-proper”}.$$

(2) $(*)_{\kappa}$ means: letting $P = \text{Levi}(\aleph_0, < \kappa)$, if $\bar{A} = \langle A_{\alpha}; \alpha < \kappa \rangle$ is a maximal antichain in V^P , then for $\alpha < \kappa$

$$\| \text{Levi}(\aleph_1, < \kappa) \| \text{ “} [\text{Levi}(\aleph_1, < \kappa) / \text{Levi}(\aleph_1, < \alpha)] * \text{seal}(\bar{A}) \text{ is semi-proper”}.$$

3.3A. REMARK. Clearly $(*)_{\kappa}^+ \Rightarrow (*)_{\kappa}$.

3.4. LEMMA. (1) If κ is measurable, $(*)_{\kappa}^+$ holds even after forcing by any Q , $|Q| < \kappa$ (semi-proper if you want), $S \subseteq \omega_1$ stationary, costationary, then for some forcing notion, P , κ -c.c. of power κ not adding reals, $|P| = \kappa$, $\| \text{Levi}(\aleph_1, < \kappa) \|_P \text{ “} D_{\omega_1} + S \text{ is } \aleph_2\text{-saturated”}.$

(2) So if $\aleph_0 < \kappa_0 < \dots < \kappa_n$, each κ_l as in 3.4(1), then there are P_l (for $l = 0, n$) as required in 3.2.

But the assumption in 3.4 is not “a large cardinal axiom”; the problem is not only esthetical — the large cardinal axioms are easily ordered on a scale. So

3.5. DEFINITION. $\text{Pr}_a(\kappa)$ means: $\text{Pr}_a(\kappa, f)$ for every $f: \kappa \rightarrow \kappa$, which means: there is $j: V \rightarrow M$ (elementary embedding into a transitive class) with critical point κ such that $H(j(f)(\kappa)) \subset M$ and $M^{<\kappa} \subseteq M$.

This property arises from the further analysis of the proof of $MA^+(\aleph_1\text{-complete})$. (Note $\| \text{Levi}(\aleph_1, < \kappa) \| \text{ “} MA^+(\aleph_1\text{-complete}) \text{”} \Rightarrow (*)_{\kappa}^+.$)

Certainly in the large cardinal scale, its use of embedding somewhat deviates.

3.6. LEMMA. If $\text{Pr}_a(\kappa)$ then κ is measurable, $(*)_{\kappa}^+$; moreover, if $|P| < \kappa$ then $\| \text{Levi}(\aleph_1, < \kappa) \|_P \text{ “} \text{Pr}_a(\kappa) \text{”}$ hence $\| \text{Levi}(\aleph_1, < \kappa) \|_P \text{ “} (*)_{\kappa}^+ \text{”}$

3.7. CONCLUSION. If there are $\omega + 1$ many cardinals κ , satisfying $\text{Pr}_a(\kappa)$,

then the conclusions of 1.1 (measurability, Baire and non-well orderings of \mathbf{R}) in $L(\mathbf{R})$ hold.

PROOF OF 3.2. (1) Let $Q_j = \{A : A \in V^{P_j}, A \subseteq \omega_1, A \neq \emptyset \text{ mod } D_{\omega_1} + S\}$ ordered by inverse inclusion. Note that Q_j is a P_j -name.

Clearly $Q_j \in V^{P_j}$ and by a hypothesis (forcing with P_l preserves stationarity of subsets of ω_1 from V and even from V^{P_m} , $m < l$) we know $Q_j \subseteq Q_{j+1}$.

More still: $P_j * Q_j \leq P_{j+1} * Q_{j+1}$ [in V^{P_j} , $D_{\omega_1} + S$ is \aleph_2 -saturated, so Q_j satisfies the \aleph_2 -c.c., hence every maximal antichain of Q_j (from V^{P_j}) has cardinality $\leq \aleph_1$, hence its diagonal union include $\omega_1 - S$ on a club, hence is also a maximal antichain of Q_{j+1}]. Let $G_{P_n} \subseteq P_n$ be generic over V , thus $G_{P_j} \stackrel{\text{def}}{=} G_{P_n} \cap P_j$ is generic over V . Let $G_{Q_n} \subseteq Q_n$ be generic over $V[G_{P_n}]$, then $G_{Q_j} \stackrel{\text{def}}{=} G_{Q_n} \cap Q_j$ is a generic subset of Q_j over $V[G_{P_j}]$. Now in $V[G_{P_j}, G_{Q_j}]$ we can compute the ultrapower $V_j \stackrel{\text{def}}{=} V^{\omega_1/G_{Q_j}}$, where we consider only functions $h : \omega_1 \rightarrow V$ which belong to $V[G_{P_j}]$. Set $V_{-1} = V$, and by Los theorem $V_j < V_{j+1}$ (before the Moskowski collapse). As in $V[G_{P_j}]$, $D_{\omega_1} + S$ is \aleph_2 -saturated, it is well known that V_j is well-founded, and we can identify it with a well-founded class in $V[G_{P_j}][G_{Q_j}]$. Also, by \aleph_2 -saturation of $D_{\omega_1} + S$, we know that every real number which belongs to $V[G_{P_j}][G_{Q_j}]$, in $V[G_{P_j}]$, has a Q_j -name τ , hence for some $\langle f_n^r : n < \omega \rangle \in V[G_{P_j}]$, $f_n^r : \omega_1 \rightarrow \{0, 1\}$, $\tau[G_{Q_j}](n) = 0 \Leftrightarrow (f_n^r)^{-1}(\{0\}) \in G_{Q_j}$ hence $\tau[G_{Q_j}] \in V_j$. So $\mathbf{R}^{V_j} = \mathbf{R}^{V[G_{P_j}, G_{Q_j}]}$.

We conclude that

$$\mathbf{R}(V) < \mathbf{R}(V[G_{P_0}, G_{Q_0}]) < \dots < \mathbf{R}(V[G_{P_n}, G_{Q_n}]).$$

But for each $i < n$, $P_i * Q_i \in V_{\kappa_{i+1}}$, κ_{i+1} is strongly inaccessible in V and all cardinals $< \kappa_{i+1}$ are collapsed to ω in $V[G_{P_{i+1}}, G_{Q_{i+1}}]$. Hence for $i < n$ we can find cardinals $\gamma_{i+1} \in (\kappa_i, \kappa_{i+1})$ and mutual generics $g_{i+1} \subset \text{Levi}(\omega, \gamma_{i+1})$ such that:

$$\begin{aligned} \mathbf{R}(V[G_{P_0}, G_{Q_0}]) &\subset \mathbf{R}(V[g_1]) \subset \mathbf{R}(V[G_{P_1}, G_{Q_1}]) \subset \mathbf{R}(V[g_1, g_2]) \\ &\subset \dots \subset \mathbf{R}(V[g_1, \dots, g_n]) \subset \mathbf{R}(V[G_{P_n}, G_{Q_n}]). \end{aligned}$$

Suppose

$$\varphi(\bar{y}) = \exists x_1 \forall x_2 \dots Qx_n \psi(\bar{x}, \bar{y})$$

is a Σ_{n+2}^1 formula (so ψ is Σ_2^1 or Π_2^1 depending on the parity of n). Then for all \bar{b} from $\mathbf{R}(V[G_{P_0}, G_{Q_0}])$,

$$V[G_{P_n}, G_{Q_n}] \models \varphi[\bar{b}]$$

iff

$$\exists a_1 \in \mathbf{R}(V[g_1]) \forall a_2 \in \mathbf{R}(V[g_1, g_2]) \cdots \forall a_n \in \mathbf{R}(V[g_1, \dots, g_n]) \psi[\bar{a}, \bar{b}]$$

(ψ is either Σ_2^1 or Π_2^1 and so $\psi[a, b]$ is absolute).

Suppose $A \subset \mathbf{R}(V[G_{P_n}, G_{Q_n}])$ is Σ_{n+2}^1 in parameters from $\mathbf{R}(V)$. Then by the homogeneity of the Levy collapses and since (c^+) is collapsed to ω in $V[G_{P_0}, G_{Q_0}]$ it follows that $A \cap V[G_{P_0}, G_{Q_0}]$ is Lebesgue measurable, etc. in $V[G_{P_0}, G_{Q_0}]$. But $\mathbf{R}(V[G_{P_0}, G_{Q_0}]) < \mathbf{R}(V[G_{P_n}, G_{Q_n}])$ hence $A \cap V[G_{P_0}, G_{Q_0}]$ is the Σ_{n+2}^1 set as computed in $V[G_{P_0}, G_{Q_0}]$. Finally $\mathbf{R}(V) < \mathbf{R}(V[G_{P_0}, G_{Q_0}])$ and so this Σ_{n+2}^1 set computed in V is Lebesgue measurable, etc. in V .

(2) For $i \leq \omega$ choose G_{P_i}, G_{Q_i} as in the proof of (1). For each $j \leq \omega$ set $V_j = V^{\omega_i}/G_{Q_i}$ (where we use all functions $h: \omega_1 \rightarrow V$ from $V[G_{P_j}]$ and we do not pass to the Mostowski collapse). $V_i < V_j$ for $i < j \leq \omega$.

Let $V = \bigcup_{i < \omega} V_i$. Thus $V < V_\omega$. V_ω is well founded and so V is well founded. Let $N = \text{collapse}(V)$ and let $\kappa = \sup\{\kappa_i \mid i < \omega\}$. Then there exists an elementary embedding

$$J: V \rightarrow N.$$

Notice $\mathbf{R}(N) = \bigcup_{i < \omega} \mathbf{R}(N_i) = \bigcup_{i < \omega} \mathbf{R}(V[G_{P_i}, G_{Q_i}])$ where, for $i < \omega$, $N_i = \text{collapse}(V_i)$.

$P_i * Q_i \in V_{\kappa_{i+1}}$ and $P_i * G_i$ collapses all cardinals $< \kappa_i$ to ω . Hence we can force over $V[G_{P_\omega}, G_{Q_\omega}]$ to find $G \subset \text{Levi}(\omega, < \kappa)$ such that G is V -generic and

$$\mathbf{R}(N) = \bigcup \mathbf{R}(V[G_i])$$

where for $i < \omega$, $G_i = G \cap \text{Levi}(\omega, < \kappa_i)$. Thus $L(\mathbf{R})^N$ can be viewed as a Solovay model obtained from collapsing below κ , which is a singular strong limit. Therefore

$$L(\mathbf{R})^N \models \text{"Every set is Lebesgue measurable . . ."}.$$

But there is an elementary embedding

$$J: V \rightarrow N,$$

hence

$$L(\mathbf{R}) \models \text{"Every set is Lebesgue measurable . . ."}.$$

PROOF OF 3.6. The measurability and preservation by small forcing are obvious.

Let us prove $(*)_\kappa^+$. Let $\langle A_i : i < \kappa \rangle$ be a $\text{Levi}(\aleph_1, < \kappa)$ -name of a maximal antichain for D_ω . Define $f_0: \kappa \rightarrow \kappa$ by: $f_0(i)$ is the minimal cardinal such that

for every stationary subset A of ω_1 from $V^{\text{Levi}(\aleph_1, < \aleph_\alpha(i))}$, for some $\alpha < f_0(i)$, $A \cap A_\alpha$ is stationary (remember $\text{Levi}(\aleph_1, < \kappa)$ satisfies the κ -c.c.); $f_1(i)$ is the minimal cardinal such that $\langle A_j : j < f_0(i) \rangle$ is a name in $\text{Levi}(\aleph_1, < f_1(i))$; $f_2(i) = \beth_{18}(f_1(i))^+$. Let $j : V \rightarrow M$ be as in 3.5 for the function f_2 . Let $G_{j(\kappa)} \subseteq \text{Levi}(\aleph_1, < j(\kappa))$ be generic over M , so $G_i \stackrel{\text{def}}{=} G_{j(\kappa)} \cap \text{Levi}(\aleph_1, < i)$ is generic over V whenever $i < j(f_2)(\kappa)$. In $V[G_\kappa]$, $\langle A_i[G_\kappa] : i < \kappa \rangle$ is a maximal anti-chain of D_{ω_1} . Let $<^*$ be a well ordering of $H(2^\kappa)^{V[G_\kappa]}$,

$$S = \{N : N < (H(2^\kappa)^{V[G_\kappa]})^\infty, \in, <^*\}, N \text{ countable, and there is no } N_1, N < N_1 < (H(2^{2^\kappa})^\infty, \in, <^*), N_1 \text{ countable, } N_1 \cap \omega_1 = N \cap \omega_1 \in \bigcup_{i \in N_1} A_i[G_\kappa]\}.$$

Clearly in V there exists a $\text{Levi}(\aleph_1, \leq 2^{2^\kappa})$ -name $\langle \dot{a}_i : i < \omega_1 \rangle$ such that

$$\Vdash_{\text{Levi}(\aleph_1, < 2^{2^\kappa})} \text{“}\langle \dot{a}_i : i < \omega_1 \rangle \text{ is increasing, continuous, each } \dot{a}_i \text{ countable, } \bigcup_{i < \omega_1} \dot{a}_i = H(2^\kappa)^{V[G_\kappa]} \text{”}.$$

Let $A = \{i : (\exists N \in S)(\omega_1 \cap \dot{a}_i \subseteq |N| \subseteq \dot{a}_i)\}$. As $\text{Levi}(\aleph_1, \leq 2^{2^\kappa})$ preserve stationary sets of $\{b : b \subseteq H(2^\kappa) \text{ is countable}\}$, we have that

$$\Vdash_{\text{Levi}(\aleph_1, \leq 2^{2^\kappa})} \text{“}A \text{ is a stationary subset of } \omega_1 \text{”}.$$

This holds also in M , so there exists $i < j(f_0)(\kappa)$ such that, in M ,

$$\Vdash_{\text{Levi}(\aleph_1, < j(f_0)(\kappa))} \text{“}A \cap A_i \text{ is a stationary subset of } \omega_1 \text{”}.$$

Therefore (see choice of f_2) this last assertion holds in V too (remember that $H(j(f_2)(\kappa)) \subseteq M$ and note that $S, A \in M$), so in $M[G_\kappa]$ letting $\kappa(0) = (j(f_1))(\kappa)$, $\kappa(1) = 2^{2^{\kappa(0)}}$, $P_\kappa = \text{Levi}(\aleph_1, < \kappa)$.

$$S_1 = \{N < (H(\kappa(1))^{M[G_\kappa]})^\infty, \in, <^*\} : N \text{ countable and for some } p \in P_{\kappa(0)}/G_\kappa, p \text{ is } (N, P_{\kappa(0)}/G_\kappa)\text{-semi-generic, } p \Vdash \text{“}N \cap \omega_1 \in A \cap A_i \text{”}\}$$

is a stationary subset of $[H(2^{2^{\kappa(0)}})^{M[G_\kappa]}]^\omega$. But

$$H(2^{2^{\kappa(1)}})^{M[G_\kappa]} = H(2^{2^{\kappa(0)}})^{V[G_\kappa]}$$

hence the same holds with $V[G_\kappa]$ instead of $M[G_\kappa]$.

Now in $V[G_\kappa]$,

$$S_2 = \{N \cap H(\kappa(1)): N < H(\aleph_3(j(\kappa))^+)^{M[G_\kappa]}, \in, <^*\}, N \text{ countable} \\ \langle q_i : i < \omega_1 \rangle, j \upharpoonright \aleph_3(\kappa)^+, A, A_i, \text{ belongs to } N\}$$

is a club of $S_{\leq \aleph_0}(H(\kappa(1)))$; remember $M^{<\kappa} \subseteq M$. So there is $N_2 \in S_2 \cap S_1$, and let N_1 be the Skolem hull of $|N_2|$ in $(H(\aleph_3(j(\kappa))^+)^{M[G_\kappa]})^+$. And let $p \in P_{\kappa(0)}/G_\kappa$ be as in the definition of S_1 . So clearly (in $M[G_\kappa]$) p is also $(N_2, P_{f(j(\kappa))})$ -semi-generic. So let $\delta \stackrel{\text{def}}{=} N_1 \cap \omega_1$; as $p \Vdash \delta \in A$, for some $q, p \leq q \in P_{\kappa(0)}/G_\kappa$ and $b \in S$,

$$q \Vdash \text{“}\delta \subseteq b \subseteq a_\delta\text{”}.$$

Also, as q is $(N_1, P_{\kappa(0)}/G_\kappa)$ -semi-generic, $\langle q_i : i < \omega_1 \rangle \in N_1$ we have that

$$q \Vdash \text{“}N_1 \cap H(2^\kappa)^{V[G_\kappa]} = a_\delta\text{”}.$$

Therefore

$$q \Vdash \text{“}b \subseteq N_1 \cap H(2^{2^\kappa})^{V[G_\kappa]}\text{”}$$

so really $b \subseteq N_1$, and thus (see definition of S_2) $j(b) = j''(b) = \{j(x) : x \in b\} \subseteq N_1$, and therefore

$$M[G_\kappa] \vDash j(b) \in j(S).$$

But N_2, i, q contradict this. This concludes the proof of 3.6.

We can skip 3.4.

3.8. CLAIM. If $\aleph_0 < \kappa_1 < \kappa_2 < \dots < \kappa_n$ are strongly inaccessible and $(*)_{\kappa_i}$ then the conclusion of 3.2 holds.

PROOF. Let $P_l = \text{Levi}(\aleph_1, \kappa_l)$ for $l = 1, \dots, n+1$ where we stipulate $\kappa_{n+1} = \aleph_8(\kappa_n)$. Let S_l be the P_{l+1} -name of the diagonal intersection of the diagonal union of every $\bar{A} \in V^{P_l}$, a maximal antichain of D_{ω_1} in V^{P_l} . The union and intersection are well defined modulo D_{ω_1} ; as in $V^{P_{l+1}}$ they are of \aleph_1 subsets of ω_1 . As $(*)_{\kappa_i}$, S_l is a stationary subset of κ_l , with stationary intersection with any stationary $S \in V^{P_l}$ ($S \subseteq \omega_1$). So we can show inductively $S^* = \bigcap_{l \leq n} S_l \neq \emptyset \pmod{D_{\omega_1}}$ in $V^{P_{n+1}}$. Now let $Q = \{A \subseteq S^* : A \text{ stationary}\}$ (in $V^{P_{n+1}}$), ordered by inverse inclusion, and proceed as in the proof of 3.2.

3.9. REMARK. (1) However, 3.8 is not a big saving, as from the large cardinals for which we can use 3.8 we could also use 3.2 (i.e. get \aleph_2 -saturated ideals $D_{\omega_1} + S$) (by [Sh 4]).

(2) Here, we can require on j only “ $M^\omega \subseteq M$ ” instead of “ $M^{<\kappa} \subseteq M$ ”.

§4. Happy end

A close look at 3.5 and 3.6 shows their lousiness — $\text{Pr}_a(\kappa)$ implies that there are many smaller cardinals with the desired properties. Further analysis leads to 4.3 below, which is further deviating but totally in the scale of large cardinals.

4.1. DEFINITION. $\text{Pr}_b(\kappa)$, now usually called “ κ is a Woodin cardinal”, means:

For every $f: \kappa \rightarrow \kappa$, there is an elementary embedding $j: V \rightarrow M$ with critical point $\lambda < \kappa$, such that $H(j(f)(\lambda)) \subseteq M$ and $f''\lambda \subset \lambda$.

So κ is a Mahlo cardinal, but not necessarily a weakly compact cardinal.

4.2. THEOREM. *If $\text{Pr}_b(\kappa)$ then $(*)_\kappa$.*

PROOF. Let $P \stackrel{\text{def}}{=} \text{Levi}(\aleph_1, < \kappa)$, and $\langle A_i : i < \kappa \rangle$ be a P -name of a maximal antichain for D_{ω_1} (in V^P). Define $f_l: \kappa \rightarrow \kappa$ for $l = 0, 1, 2$, as in the proof of 3.6. Let

$$C = \{\delta < \kappa : \delta \text{ a limit cardinal and } i < \delta \Rightarrow f_1(i) < \delta\};$$

it is closed unbounded. So for some stationary set of $\lambda \in C$, there is an elementary embedding $j: V \rightarrow M$, with critical point λ , $H(j(f)(\lambda)) \subseteq M$. We continue as in 3.6.

As in 3.6 for each such λ , in $V^{\text{Levi}(\aleph_1, < \lambda)}$, $\langle A_i : i < \lambda \rangle$ is semi-proper. Now if $N < (H(\aleph_3(\kappa), \in))$ is countable, $\kappa \in N$, $\langle A_i : i < \lambda \rangle \in N$ and $(p, q) \in \text{Levi}(\aleph_1, < \kappa) * \text{seal}(\bar{A})$, then for some λ

- (i) $\lambda \in C \cap N$,
- (ii) $\Vdash_{\text{Levi}(\aleph_1, < \lambda)} \langle A_i : i < \lambda \rangle \text{ is semi-proper}$,
- (iii) $p \in \text{Levi}(\aleph_1, < \lambda)$.

We can find $p_1, p \leq p_1 \in \text{Levi}(\aleph_1, < \lambda)$ and p_1 is $(N, \text{Levi}(\aleph_1, \lambda))$ -generic. Let $p_1 \in G_\lambda \subseteq \text{Levi}(\aleph_1, < \lambda)$, G_λ generic over V . In $V[G_\lambda]$ we can find $N_1, N < N_1 < (H(\aleph_3(\kappa))[G_\lambda], \in)$ and for some $i \in N_1 \cap \lambda$, $\delta \stackrel{\text{def}}{=} N \cap \delta = N_1 \cap \delta \in A_i[G_\lambda]$. We can find $G \subseteq \text{Levi}(\aleph_1, < \kappa)$ generic over V , $G_\lambda \subseteq G$, and then find $q_1, q \leq q_1 \in \text{seal}(\bar{A}[G])$, $q_1(N_1, \text{seal}(\bar{A}))$ -generic. So there is a $\text{Levi}(\aleph_1, < \kappa)$ name \check{q}_1 for such q_1 and (p_1, \check{q}_1) is a required $(N, \text{Levi}(\aleph_1, < \lambda)) * \text{seal}(\bar{A})$ -semi-generic.

4.3. CONCLUSION. If there are $\omega + 1$ many cardinals as in 4.1, then $L(\mathbf{R})$ satisfies (a), (b), (c), (d) of 1.1.

PROOF. By 4.2, 3.8.

Why is this a happy end? Because Martin and Steel seem to be proving that finitely many cardinals do not suffice for 1.1(c) (hence (a), (b)).

4.4. THEOREM. *If κ satisfies Definition 4.1 in $L(V_\kappa)$ and $L(V_\kappa)^*$ exists, then all Σ_3^1 sets are Lebesgue measurable.*

The proof is combining what was done above with classical proofs. Steel, hearing about the previous work but not 4.4, proved this independently. (Of course, for Σ_{n+3}^1 we need n cardinals, λ below κ satisfying $\text{Pr}_b(\lambda)$.)

The proof is different than for 3.2; one proves by induction on n that Σ_{n+4}^1 -truth is absolute between forcing extensions by partial orders of size $<$ the least λ with $\text{Pr}_b(\lambda)$. Here $L(V_\kappa)^*$ is used. Without it we get only that Σ_{n+3}^1 -truth is absolute.

Note

4.5. CLAIM. Really the inaccessibility can be replaced by $2^{2^{\kappa_l}} < \kappa_{l+1}$ (with $\kappa_{-1} = \aleph_2$).

4.6. CLAIM. We can prove our theorems also from a (strongly) compact cardinal.

4.7. REMARK. For more on $(*)_\kappa$ see Woodin [W3], where in an equivalent form (towers) they are used to produce strange forcing notions, which were previously produced only in models of AD .

REFERENCES

- [F] M. Foreman, *Potent axioms*, Trans. Am. Math. Soc. **294** (1986), 1–28.
- [FMS1] M. Foreman, M. Magidor and S. Shelah, *Martin maximum, saturated ideals and non-regular ultrafilters I*, Ann. of Math. **127** (1988), 1–47.
- [FMS2] M. Foreman, M. Magidor and S. Shelah, *Martin maximum, saturated ideals and non-regular ultrafilters II*, Ann. of Math. **127** (1988), 521–545.
- [L] R. Laver, *Making the supercompactness of κ indestructible under κ -directed closed forcing*, Isr. J. Math. **29** (1978), 385–388.
- [Mg] M. Magidor, *Precipitous ideal Σ_4^1 sets*, Isr. J. Math. **35** (1980), 109–134.
- [M1] Y. Moschovakis, *Descriptive Set Theory*, North-Holland, Amsterdam, 1980.
- [MS] D. A. Martin and J. R. Steel, *Determinacy in $L(\mathbb{R})$ follows from large cardinals*, Proc. Natl. Acad. Sci. U.S.A., submitted.
- [Mt] A. D. R. Mathias, *On sequences generic in the sense of Prikry*, J. Aust. Math. Soc. **15** (1973), 409–416.
- [Sh1] S. Shelah, *Proper Forcing*, Lecture Notes in Math. **940**, Springer-Verlag, Berlin, 1982.
- [Sh2] S. Shelah, *Notes, part (A): On normal ideals and Boolean algebras*, in *Around Classification Theory*, Lecture Notes in Math. **1182**, Springer-Verlag, Berlin, 1986.
- [Sh3] S. Shelah, *Iterated forcing and normal ideals on ω_1* , Isr. J. Math. **60** (1987), 345–380.
- [Sh4] S. Shelah, *Notes 3/85*, to appear in [Sh 1], new edition.

[So1] R. M. Solovay, *The cardinality Σ_2^1 sets of reals*, in *Foundation of Mathematics*, Symposium paper commemorating the sixtieth birthday of Kurt Godel, Springer-Verlag, Berlin, 1969, pp. 59–73.

[So2] R. M. Solovay, *A model of set theory in which every set of reals is Lebesgue measurable*, *Ann. of Math.* **92** (1970), 1–56.

[W1] H. Woodin, *Embedding $j: L(V_{\lambda+1}) \rightarrow L(V_{\lambda+1})$ and determinacy*, Lectures L.A., Cabal seminar, Spring 1984.

[W2] H. Woodin, *Every set of $L(\mathbb{R})$ has a weakly homogeneous tree resp. from a supercompact*, L.A., Cabal seminar, Spring 1985.

[W3] H. Woodin, *Σ_1^2 absoluteness and supercompact cardinals*, Notes, May 1985.