

CARDINAL INVARIANTS \mathfrak{b}_κ AND \mathfrak{t}_κ

Saharon Shelah and Zoran Spasojević

Communicated by Žarko Mijajlović

ABSTRACT. We study cardinal invariants \mathfrak{b}_κ and \mathfrak{t}_κ , the natural generalizations of the invariants \mathfrak{b} and \mathfrak{t} to a regular cardinal κ .

0. Introduction

The cardinal invariants \mathfrak{b} and \mathfrak{t} were introduced by Rothberger [Ro39], [Ro48]. They are cardinals between \aleph_1 and 2^{\aleph_0} and have been extensively studied over the years. The survey paper [Bsxx] contains much information about these two invariants as well as many other cardinal invariants of the continuum.

The goal of this paper is to study the natural generalizations of \mathfrak{b} and \mathfrak{t} to higher regular cardinals, namely \mathfrak{b}_κ and \mathfrak{t}_κ respectively, where κ is a regular cardinal. The results presented here are that the relationship $\mathfrak{t} \leq \mathfrak{b}$ (shown by Rothberger [Ro48]) also holds for \mathfrak{b}_κ and \mathfrak{t}_κ and that, assuming, e.g., that $\kappa = \kappa^{<\kappa} \geq \beth_\omega$, if $\kappa \leq \mu < \mathfrak{t}_\kappa$ then $2^\mu = 2^\kappa$. These results are then used as constraints in the forcing construction of models in which \mathfrak{b}_κ and \mathfrak{t}_κ can take on essentially any preassigned regular value.

The cardinal \mathfrak{b}_κ was studied in [CuSh:541] where it was shown that the value of \mathfrak{b}_κ does not have any influence on the value of 2^μ for $\kappa \leq \mu < \mathfrak{b}_\kappa$ even if *GCH* is assumed to hold below κ . However, the same does not hold for \mathfrak{t}_κ as it is shown in Section 2.

In an earlier version, a wrong “improvement” of Section 1 was used and we thank the referee for detecting this.

1. Conventions and elementary facts

1.1. NOTATION. 1) For cardinals λ and κ let $[\kappa]^\lambda = \{X \subseteq \kappa : |X| = \lambda\}$ and ${}^\lambda\kappa$ is the set of functions from λ to κ . The symbol κ^λ is used to denote the cardinality of the set $\{f : f : \lambda \rightarrow \kappa\}$.

2000 *Mathematics Subject Classification*. Primary 03E17; Secondary 03E35.

Research of the first author was partially supported by the NSF and Israel Science Foundation. Publication 643.

2) For $A, B \in [\kappa]^\kappa$ let $A \subseteq^* B$ iff $|A \setminus B| < \kappa$ and $A \subset^* B$ iff $|A \setminus B| < \kappa \wedge |B \setminus A| = \kappa$. Let “ A is an almost subset of B ” mean $A \subseteq^* B$. For $f, g \in {}^\kappa\kappa$ let $f <^* g$ iff $\exists \beta < \kappa \forall \alpha > \beta (f(\alpha) < g(\alpha))$. Then $\mathcal{F} \subseteq {}^\kappa\kappa$ is unbounded (or \leq^* -unbounded) in $({}^\kappa\kappa, <^*)$ mean that $\forall f \in {}^\kappa\kappa \exists g \in \mathcal{F} (g \not<^* f)$.

3) For a filter D on a set A let $A = \text{Dom}(D)$.

1.2. DEFINITION. For regular cardinal κ let

$$\begin{aligned} \mathfrak{b}_\kappa &= \min\{\mathcal{F} : \mathcal{F} \subseteq {}^\kappa\kappa \text{ and } \mathcal{F} \text{ is } <^* \text{-unbounded in } {}^\kappa\kappa\}, \\ \mathfrak{t}_\kappa &= \min\{\mathcal{T} : \mathcal{T} \subseteq [\kappa]^\kappa, |\mathcal{T}| \geq \kappa, \mathcal{T} \text{ is well ordered by } \subset^*, \\ &\quad (\forall C \in [\kappa]^\kappa) (|\kappa \setminus C| = \kappa \Rightarrow \exists A \in \mathcal{T} (|A \setminus C| = \kappa)) \\ &\quad \mathcal{T} \text{ is with no } \subseteq^* \text{-last element} \\ &\quad \text{and has no } \subseteq^* \text{-unbounded subset of cardinality } < \kappa\}. \end{aligned}$$

(Observe that some of the requirements in the definition of \mathfrak{t}_κ follow from the others.) In this notation $\mathfrak{b} = \mathfrak{b}_\omega$ and $\mathfrak{t} = \mathfrak{t}_\omega$.

An equivalent formulation of \mathfrak{t}_κ is obtained if \supseteq^* is used instead of \subseteq^* . Standard arguments show that

1.3. FACT. For any regular cardinal κ , $\kappa^+ \leq \mathfrak{b}_\kappa$, $\mathfrak{t}_\kappa \leq 2^\kappa$ and in the definition of \mathfrak{b}_κ , \mathcal{F} may be assumed to be well ordered by $<^*$ and consisting only of strictly increasing functions. Thus, both \mathfrak{b}_κ and \mathfrak{t}_κ are regular cardinals.

If $\kappa > \omega$ then there exists a countable \subseteq^* -decreasing sequence of elements in $[\kappa]^\kappa$, which does not have a lower bound with respect to \subseteq^* .

1.4. FACT. $\mathfrak{t}_\kappa \leq \mathfrak{b}_\kappa$.

PROOF. On the case $\kappa = \omega$ see, e.g., [Bsxx]. So assume $\kappa > \omega$ and by way of contradiction assume $\mathfrak{b}_\kappa < \mathfrak{t}_\kappa$. Let $\{f_\alpha : \alpha < \mathfrak{b}_\kappa\} \subseteq {}^\kappa\kappa$ be $<^*$ -unbounded in $({}^\kappa\kappa, <^*)$ and without loss of generality by 1.3 such that $\alpha < \beta \rightarrow f_\alpha <^* f_\beta$. For each $\alpha < \mathfrak{b}_\kappa$ let $C_\alpha = \{\xi < \kappa : \forall \zeta < \xi (f_\alpha(\zeta) < \xi)\}$. Then each C_α is closed unbounded in κ and $\alpha \leq \beta \rightarrow C_\beta \subseteq^* C_\alpha$. Easily, $(\forall \alpha < \mathfrak{b}_\kappa) (\exists \beta) (\alpha < \beta < \mathfrak{b}_\kappa \wedge C_\beta \subset^* C_\alpha)$, so without loss of generality $\alpha < \beta \rightarrow C_\beta \subset^* C_\alpha$. Since $\kappa < \mathfrak{b}_\kappa$ and we are assuming that $\mathfrak{b}_\kappa < \mathfrak{t}_\kappa$, there is $A \in [\kappa]^\kappa$ such that $\forall \alpha < \mathfrak{b}_\kappa (A \subset^* C_\alpha)$. Let $f : \kappa \rightarrow A$ be such that $\forall \xi < \kappa (\xi < f(\xi))$. Fix $\alpha < \mathfrak{b}_\kappa$ and let $i_\alpha \in [\kappa]^{<\kappa}$ be such that $A \setminus i_\alpha \subseteq C_\alpha \setminus i_\alpha$. Let $\xi \in \kappa \setminus i_\alpha$ and $\zeta = \min(C_\alpha \setminus (\xi + 1))$ and note that $f_\alpha(\xi) < \zeta$ by the definition of C_α . However, $A \setminus \xi \subseteq C_\alpha \setminus \xi$, and $\xi < f(\xi)$, so $\zeta \leq f(\xi)$. In other words, $(\forall \xi \in \kappa \setminus i_\alpha) (f_\alpha(\xi) < f(\xi))$; hence f is a $<^*$ -bound for $\{f_\alpha : \alpha < \mathfrak{b}_\kappa\}$. This is a contradiction and the lemma is proved. $\square_{1.4}$

2. Combinatorics

The goal of this section is to show that if, e.g., $\kappa^{<\kappa} = \kappa \geq \beth_\omega$ then $2^\mu = 2^\kappa$ for any μ with $\kappa \leq \mu < \mathfrak{t}_\kappa$. Naturally we start from the scheme of the proof of $\omega \leq \mu < \mathfrak{t} \rightarrow 2^\mu = 2^\omega$, namely to use $\mu < \mathfrak{t}_\kappa$ to construct a binary tree in $(\mathcal{P}(\kappa), \subset^*)$ of height μ . However, when κ is uncountable a difficulty arises in the construction

at limit stages of cofinality less than κ , a case which does not occur when $\kappa = \aleph_0$. The difficulty comes from the fact that a \subseteq^* -decreasing sequence in $[\kappa]^\kappa$ of limit length less than κ may not have a lower bound. To deal with this difficulty, a notion of a closed subset of κ with respect to a certain parameter is introduced next.

From where comes the condition $\kappa \geq \mathfrak{J}_\omega$? From using a result from pcf theory [Sh:460], so called “revised GCH”.

2.1. MAIN LEMMA. *If κ is regular, then for every $\mu < \mathfrak{t}_\kappa$ we have $2^\mu = 2^\kappa$ provided that at least one of the following conditions holds:*

CND_1 : $\kappa \geq \mathfrak{J}_\omega$ and $\kappa = \kappa^{<\kappa}$.

CND_2 : diamond on κ or at least $D\ell_\kappa$ (see below, see more e.g., in [Sh:460]).

CND_3 : there is a sequence $\bar{D} = \langle D_{\delta,\gamma} : \delta \text{ limit} < \kappa \text{ and } \gamma < \gamma_\delta \rangle$ such that:

- (a) $\gamma_\delta < \kappa$ (for each limit $\delta < \kappa$)
- (b) $D_{\delta,\gamma}$ is a filter on δ to which all cobounded subsets of δ belongs
- (c) for every unbounded subset A of κ , for stationarily many ordinals $\delta < \kappa$ we have: for some $\gamma < \gamma_\delta$, $A \cap \delta \in D_{\delta,\gamma}$
- (d) moreover, if $\tau < \kappa$ is regular and A_i an unbounded subset of κ for $i < \tau$ and for $i < j < \tau$, A_j is an almost subset of A_i (i.e., $A_j \subseteq^* A_i$) then for stationarily many $\delta < \kappa$ some $\gamma < \gamma_\delta$ satisfies: for every $i < \tau$ we have $A_i \cap \delta \in D_{\delta,\gamma}$.

2.2. DEFINITION. Let $D\ell_\kappa$ mean: κ is regular uncountable and there is a sequence $\langle \mathcal{P}_\alpha : \alpha < \kappa \rangle$ such that:

- (a) \mathcal{P}_α is a family of subsets of α
- (b) \mathcal{P}_α has cardinality $< \kappa$
- (c) for every $A \subseteq \kappa$ the set $\{\delta < \kappa : A \cap \delta \in \mathcal{P}_\delta\}$ is a stationary subset of κ .

PROOF. By [Sh:460], CND_1 implies CND_2 . Easily CND_2 implies CND_3 , so we shall assume the latter. Let E be a club of κ such that each member is a limit ordinal and $\delta < \alpha \in E$ implies $(\delta + \gamma_\delta + \omega) < \alpha$; here δ always denotes a limit ordinal. The proof is preceded by a definition and some facts.

Below A, B denote subsets of κ , a fix regular uncountable cardinal; unbounded means unbounded in κ .

2.3. DEFINITION. 1) A subset A of κ is called (E, \bar{D}) -closed when: for every $\delta \in E$, if $A \cap \delta \in D_{\delta,\gamma}$ then $\delta + \gamma \in A$.

2) The atomic (E, \bar{D}) -closure $\text{atcl}(A)$ of A , a subset of κ is $A \cup \{\delta + \gamma : \delta \in E, \gamma < \gamma_\delta \text{ and } A \cap \delta \in D_{\delta,\gamma}\}$.

3) We define $\text{cl}^\alpha(A)$ for A a subset of κ and α an ordinal, by induction on α : $\text{cl}^\alpha(A) = A \cup \bigcup \{\text{atcl}(\text{cl}^\beta(A)) : \beta < \alpha\}$.

4) We define the (E, \bar{D}) -closure of A , $\text{cl}(A)$ as $\text{cl}^\alpha(A)$ for every α large enough; (see 2.4(3)).

2.4. FACT. (1) κ is (E, \bar{D}) -closed, unbounded in κ .

(2) For $\alpha < \beta$ we have $A \subseteq \text{cl}^\alpha(A) \subseteq \text{cl}^\beta(A)$.

(3) $\text{cl}^\alpha(A) \subseteq \text{cl}^\kappa(A) = \text{cl}^\beta(A)$ if $\alpha < \kappa \leq \beta$.

(4) If $\delta \in E$ and $A \subseteq \kappa$ then $\text{cl}(A \cap (\delta + \gamma_\delta)) = \text{cl}(A) \cap (\delta + \gamma_\delta)$.

2.5. FACT. For A a subset of κ

(a) $\text{cl}(A)$ is a (E, \bar{D}) -closed set

(b) $\text{cl}(A)$ is the minimal (E, \bar{D}) -closed set which includes A

(c) $\text{cl}(A)$ is bounded (in κ) iff A is bounded (in fact if A is a subset of $\delta \in E$ then $\text{cl}(A)$ is a subset of $\delta + \gamma_\delta$).

2.6. FACT. If A is (E, \bar{D}) -closed and unbounded, and B is an unbounded, almost a subset of A then $\text{cl}(B)$ is an unbounded, (E, \bar{D}) -closed almost subset of A .

PROOF. By 2.4(4).

2.7. FACT. $\text{cl}(A)$ is the increasing union of $\text{cl}(A \cap \alpha)$ for $\alpha < \kappa$.

2.8. FACT. If A is (E, \bar{D}) -closed and unbounded (subset of κ) then we can find two disjoint (E, \bar{D}) -closed unbounded subsets of it.

PROOF. We choose by induction on $i < \kappa$, ordinals α_i, β_i such that:

(*) α_i, β_i are distinct members of A and larger than the supremum of the (E, \bar{D}) -closure of $\{\alpha_j, \beta_j : j < i\}$.

There is no problem to do it and $\text{cl}(\{\alpha_i : i < \kappa\}), \text{cl}(\{\beta_i : i < \kappa\})$ are two sets as required. $\square_{2.8}$

2.9. FACT. If A_i is a (E, \bar{D}) -closed unbounded subset of κ for $i < \tau$, τ a regular cardinal $< \kappa$ and for $i < j < \tau$ the set A_j is an almost subset of A_i then their intersection is a (E, \bar{D}) -closed unbounded subset of κ .

PROOF. By the last demand in CND_3 , i.e., clause (d). $\square_{2.9}$

CONTINUATION OF THE PROOF OF 2.1. Now let $\mu < \mathfrak{t}_\kappa$. We choose by induction on $\zeta \leq \mu$ for every sequence η of zeroes and ones of length ζ , a set A_η such that:

(A) A_η is a subset of κ of cardinality κ

(B) A_η is (E, \bar{D}) -closed

(C) if ρ is an initial segment of η then $A_\eta \setminus A_\rho$ has cardinality $< \kappa$

(D) if $\rho \in {}^\varepsilon 2$, $\varepsilon < \zeta$ then $A_{\rho \frown \langle 0 \rangle}, A_{\rho \frown \langle 1 \rangle}$ are disjoint.

If we succeed, clearly $\{A_\rho : \rho \text{ a sequence of zeroes and ones of length } \mu\}$ is a family of 2^μ pairwise almost disjoint subsets of κ , so $2^\mu \leq 2^\kappa$ thus finishing.

In stage $\zeta = 0$ use fact 2.4(1).

In limit stages of cofinality $\geq \kappa$, we use the hypothesis $\mu < \mathfrak{t}_\kappa$ to get an unbounded A_ρ^0 , almost included in each $A_{\rho \upharpoonright \varepsilon}$ for $\varepsilon < \zeta$. Let A_ρ be $\text{cl}(A_\rho^0)$, it is (E, \bar{D}) -closed (by fact 2.5, clause (a)) is unbounded (by fact 2.5, clause (c)) and is almost a subset of $A_{\rho \upharpoonright \varepsilon}$ for each $\varepsilon < \zeta$ by Fact 2.6.

In limit stages ζ of cofinality $< \kappa$, we choose an increasing sequence $\langle \varepsilon_i : i < \text{cf}(\zeta) \rangle$ of ordinals $< \zeta$ converging to ζ . We let $A_\rho^0 = \bigcap \{A_{\rho \upharpoonright \varepsilon_i} : i < \text{cf}(\zeta)\}$. By the Fact 2.9, A_ρ^0 has cardinality κ . Let A_ρ be the (E, \bar{D}) -closure of A_ρ^0 ; now A_ρ is an unbounded (E, \bar{D}) -closed subset of κ (see Fact 2.5, clauses (c), (a)), it is almost subset of each $A_{\rho \upharpoonright \varepsilon_i}$ by Fact 2.6, hence is as required.

Lastly, for successor stages use Fact 2.8. $\square_{2.1}$

2.10. REMARK. In Lemma 2.1 it suffices to assume the following variant.

CND_4 : we can find a set \mathcal{D} such that:

- (a) \mathcal{D} is a family of $\leq \kappa$ filters
- (b) each filter D from \mathcal{D} , is a filter on some $\alpha = \alpha[D] < \kappa$
- (c) if $A \in [\kappa]^\kappa$ then for some $D \in \mathcal{D}$ we have¹ $A \cap \alpha[D] \in D$
- (d) if $\theta = \text{cf}(\theta) < \kappa$ and $A_i \in [\kappa]^\kappa$ for $i < \kappa$ and $i < j < \theta \Rightarrow A_j \subseteq A_i$ then for some $D \in \mathcal{D}$ we have $j < \theta \Rightarrow A_j \in D$
- (e) for each $\beta < \kappa$, the set $\{D : \beta \cap \alpha[D] \in D\}$ has cardinality $< \kappa$.

Why? First note that for some \bar{D}

- (*) $\bar{D} = \langle D_i : i \in S^* \rangle$ list \mathcal{D} with no repetitions where $\alpha[D_i] \leq i$ and $S^* \subseteq \kappa$ is unbounded.

[Why? Note that $|\mathcal{D}| \leq \kappa$ by clause (a) but if $|\mathcal{D}| < \kappa$ then $\alpha^* = \bigcup \{\alpha[D] : D \in \mathcal{D}\}$ is $< \kappa$ so $A =: (\alpha, \kappa) \in [\kappa]^\kappa$ but by clause (c) there is $D \in \mathcal{D}$, such that $A \cap \alpha[D] \in D$ easy contradiction. So together $|\mathcal{D}| = \kappa$, let $\langle D_i^0 : i < \kappa \rangle$ list \mathcal{D} , and let us define $\zeta(i) < \kappa$ strictly increasing such that $\alpha[D_i^0] < \zeta(i)$. Now let $S^* = \{\zeta(i) : i < \lambda\}$, $D_{\zeta(i)} = D_i^0$.]

Also let E be $\{\delta < \lambda : \delta = \sup(S^* \cap \delta), \delta \text{ limit ordinal and for no } \alpha < \delta \leq \zeta \in S^* \text{ do we have } \alpha \cap \text{Dom}(D_\zeta) \in D_\zeta\}$, so E is a club of λ , and continue as above.

3. Forcing

For regular κ we know that $\kappa < \mathfrak{t}_\kappa \leq \mathfrak{b}_\kappa$, both regular, so we may wonder are there additional restrictions.

We use the previous section by which if $\diamond_{\kappa, 2^\kappa} < 2^\lambda$ then $\mathfrak{t} \leq \lambda$, so making \mathfrak{b}_κ larger than some such λ guarantees this.

Let κ be a regular uncountable cardinal and let λ, μ, θ be cardinals such that $\kappa < \lambda \leq \mu \leq \theta$ with λ, μ regular and $\text{cf}(\theta) \geq \lambda$. This section deals with the construction of a model for $\mathfrak{t}_\kappa = \lambda$, $\mathfrak{b}_\kappa = \mu$ and $2^\kappa = \theta$. The idea behind the construction is as follows: Start with a countable transitive model (c.t.m.) N for $ZFC + GCH$. Expand N to a model M by forcing with the standard partial order for adding θ^+ many subsets of λ (see below). Then

$$M \models \text{“}\forall \xi < \lambda (2^\xi = \xi^+ \wedge 2^\lambda = \theta^+) \text{”}.$$

In M , perform an iterated forcing construction with $< \kappa$ -supports of length $\theta \cdot \mu$ (ordinal product) with κ -closed and κ^+ -c.c. partial orders as follows: At stages which are not of the form $\theta \cdot \xi$ ($\xi < \mu$) towers in $(\mathcal{P}(\kappa), \subseteq^*)$ of height η are destroyed for $\kappa < \eta < \lambda$. At stages of the form $\theta \cdot \xi$ a function from κ to κ is added to eventually dominate all the functions from κ to κ constructed by that stage. The bookkeeping is arranged in such a way that by the end of the construction all

¹we can let \mathcal{D} be over $\text{Dom}(D) \subseteq \alpha$, no real difference

towers of height η for $\kappa < \eta < \lambda$ are considered so that in the final model $\mathfrak{t}_\kappa \geq \lambda$. However, in the final model

$$\forall \xi ((\xi < \kappa \rightarrow 2^\xi = \xi^+) \wedge (\kappa \leq \xi < \lambda \rightarrow 2^\xi = \theta)) \wedge 2^\lambda = \theta^+$$

so that, by the previous section, $\mathfrak{t}_\kappa = \lambda$. By virtue of adding dominating functions at stages of the form $\theta \cdot \xi$, the final model has a scale in $(\kappa^\kappa, <^*)$ of order type μ so that $\mathfrak{b}_\kappa = \mu$.

The rest of this section deals with the details of the construction. In showing that the final model has the desired properties it is important to know that cardinals are not collapsed. A standard way of proving this is to show that the final partial order obtained by the iteration is κ -closed and has the κ^+ -cc this follows (see [Sh:80]) but a self-contained proof is given. And to show that the final partial order has the two properties, the names for the partial orders used in the iteration must be carefully selected. The discussion here will be analogous to the discussion in the final section of [Ku83] which deals with countable support iterations. Also many proofs are omitted here since they are analogous to the proofs of the corresponding facts in [Ku83].

3.1. DEFINITION. Let \mathbb{P} be a partial order and π a \mathbb{P} -name for a partial order. π is full for $< \kappa$ -sequences iff whenever $\alpha < \kappa$, $p \in \mathbb{P}$, $\rho_\xi \in \text{dom}(\pi)$ ($\xi < \alpha$) and for each $\xi < \zeta < \alpha$

$$p \Vdash \text{“}\rho_\zeta, \rho_\xi \in \pi \wedge \rho_\zeta \leq \rho_\xi\text{”}$$

then there is a $\sigma \in \text{dom}(\pi)$ such that $p \Vdash \text{“}\sigma \in \pi\text{”}$ and $p \Vdash \text{“}\sigma \leq \rho_\xi\text{”}$ for all $\xi < \alpha$.

The reason for using names which are full for $< \kappa$ -sequences is because of the following

3.2. LEMMA. *Let M be a c.t.m. for ZFC and in M let*

$$\langle \langle \mathbb{P}_\xi : \xi \leq \alpha \rangle, \langle \pi_\xi : \xi < \alpha \rangle \rangle$$

be a $< \kappa$ -support iterated forcing construction and suppose that for each ξ , the \mathbb{P}_ξ -name π_ξ is full for $< \kappa$ -sequences. Then \mathbb{P}_α is κ -closed in M .

The next few paragraphs show how to select names for partial orders in the construction so that they are full for $< \kappa$ -sequences. First consider the partial order which destroys a tower in $(\mathcal{P}(\kappa), C^*)$. Let ϵ be a regular cardinal with $\kappa < \epsilon < \lambda$ and $a = \langle a_\xi : \xi < \epsilon \rangle$ a tower in $(\mathcal{P}(\kappa), C^*)$. In the following subsets of κ are identified with their characteristic functions.

3.3. DEFINITION. $\mathbb{T}_a = \{(s, x) : s \text{ is a function} \wedge \text{dom}(s) \in \kappa \wedge \text{ran}(s) \subseteq 2 \wedge x \in [\epsilon]^{< \kappa}\}$ with $(s_2, x_2) \leq (s_1, x_1)$ iff

$$(1) s_1 \subseteq s_2 \wedge x_1 \subseteq x_2, \quad (2) \forall \xi \in x_1 \forall \eta \in \text{dom}(s_2) \setminus \text{dom}(s_1) (a_\xi(\eta) \leq s_2(\eta)).$$

Then \mathbb{T}_a is a partial order and it is κ -closed and κ^+ -c.c. (assuming $\kappa^{< \kappa} = \kappa$). Let G be \mathbb{T}_a -generic over M and $b = \bigcup \{s : \exists x ((s, x) \in G)\}$. Since G intersects suitably chosen dense subsets of \mathbb{T}_a in M , then $b \subseteq \kappa$, $|b| = |\kappa \setminus b| = \kappa$ and $\forall \xi < \epsilon (a_\xi \subseteq^* b)$ so that a ceases to be a tower in $M[G]$.

Since the $< \kappa$ -support iteration is sensitive to the particular names used for the partial orders, a suitable name for \mathbb{T}_a is formulated next.

3.4. DEFINITION. Assume that $\mathbb{P} \in M$, (\mathbb{P} is κ -closed) and

$$\mathbf{1} \Vdash \text{“}\tau \text{ is an } \check{\epsilon}\text{-tower in } (\mathcal{P}(\kappa), \subset^*)\text{”}.$$

A standard name for \mathbb{T}_τ is $\langle \sigma, \leq_\sigma, \mathbf{1}_\sigma \rangle$, where

$$\begin{aligned} \sigma &= \{ \langle \text{op}(\check{s}, \rho), \mathbf{1}_\mathbb{P} \rangle : s \text{ is a function } \wedge \text{dom}(s) \in \kappa \wedge \text{ran}(s) \subseteq 2^\kappa \\ &\quad \mathbf{1} \Vdash \text{“}\rho \subseteq \tau \wedge |\rho| < \kappa\text{”} \wedge \rho \text{ is a nice name for a subset of } \tau \} \end{aligned}$$

and $\mathbf{1}_\sigma = \text{op}(\check{0}, \check{0})$.

Here op is the invariant name for the ordered pair and ρ is a nice name for a subset of τ if $\rho = \bigcup \{ \{ \pi \} \times A_\pi : \pi \in \text{dom}(\tau) \}$ and each A_π is an antichain in \mathbb{P} . It is irrelevant what type of name we use for \leq_σ as long as it is forced by $\mathbf{1}_\mathbb{P}$ to be the correct partial order on \mathbb{T}_τ .

In M , let \mathbb{P} , τ , and σ be as in the definition above. Let G be \mathbb{P} -generic over M and $a = \tau_G$. Then in $M[G]$, $\sigma_G = \mathbb{T}_a$. In addition, σ is full for $< \kappa$ -sequences.

The dominating function partial order is considered next. Let $F \subseteq \kappa^\kappa$. In the final construction F will be equal to κ^κ , but for the general discussion F is any subset of κ^κ .

3.5. DEFINITION. $\mathbb{D}_F = \{ (s, x) : s \text{ is a function } \wedge \text{dom}(s) \in \kappa \wedge \text{ran}(s) \subseteq \kappa \wedge x \in [F]^{<\kappa} \}$ where $(s_2, x_2) \leq (s_1, x_1)$ if

- (1) $s_1 \subseteq s_2 \wedge x_1 \subseteq x_2$,
- (2) $\forall f \in x_1 \forall \alpha \in \text{dom}(s_2) \setminus \text{dom}(s_1) (f(\alpha) < s_2(\alpha))$.

Then \mathbb{D}_F is a partial order and is κ -closed and κ^+ -c.c. (assuming $\kappa^{<\kappa} = \kappa$). Let G be \mathbb{D}_F -generic over M and $g = \bigcup \{ s : \exists x ((s, x) \in G) \}$. Then since G intersects suitably chosen dense subsets of \mathbb{D}_F in M , g is a function from κ to κ which eventually dominates every function in F , i.e., $\forall f \in F (f <^* g)$.

3.6. DEFINITION. Assume that $\mathbb{P} \in M$, (\mathbb{P} is κ -closed) M , and $\mathbf{1} \Vdash \text{“}\varphi \subseteq \kappa^\kappa\text{”}$. The standard \mathbb{P} -name for \mathbb{D}_φ is $\langle \psi, \leq_\psi, \mathbf{1}_\psi \rangle$, where

$$\begin{aligned} \psi &= \{ \langle \text{op}(\check{s}, \phi), \mathbf{1}_\mathbb{P} \rangle : s \text{ is a function } \wedge \text{dom}(s) \in \kappa \wedge \text{ran}(s) \subseteq \kappa \wedge \\ &\quad \mathbf{1}_\mathbb{P} \Vdash \text{“}\phi \subseteq \varphi \wedge |\phi| < \kappa\text{”} \wedge \\ &\quad \phi \text{ is a nice name for a subset of } \varphi \} \end{aligned}$$

and $\mathbf{1}_\psi = \text{op}(\check{0}, \check{0})$.

The choice of the \mathbb{P} -name \leq_ψ is, once again, irrelevant as long as it is forced by $\mathbf{1}_\mathbb{P}$ to be the correct partial order on \mathbb{D}_φ .

In M , let \mathbb{P} , φ , ψ , be as above. Let G be \mathbb{P} -generic over M and $F = \varphi_G$. Then, in $M[G]$, $\psi_G = \mathbb{D}_F$. In addition, ψ is full for $< \kappa$ -sequences. The use of full names for $< \kappa$ -sequences will guarantee, as indicated earlier, that the iteration is κ -closed. The use of standard names will imply that the iteration also satisfies the κ^+ -cc so that all the cardinals are preserved in the final model.

Now follows the main result of this section.

3.7. THEOREM. *Let N be a c.t.m. for $ZFC + GCH$ and, in N , let $\kappa < \lambda \leq \mu \leq \theta$ be cardinals such that κ, λ, μ are regular and $\text{cf}(\theta) \geq \lambda$. Then there is a cardinal preserving extension $M[G]$ of N such that*

$$M[G] \models \text{“}\mathfrak{t}_\kappa = \lambda \wedge \mathfrak{b}_\kappa = \mu \wedge 2^\kappa = \theta\text{”}.$$

PROOF. Let α, β be cardinals with α regular, $\alpha < \beta$, and $\text{cf}(\beta) > \alpha$. Then $\mathbb{F}n(\beta \times \alpha, 2, \alpha)$ is the standard partial order for adding β -many subsets of α (see [Ku83]). It is α -closed and α^+ -c.c. (assuming $\alpha^{<\alpha} = \alpha$), so it preserves cardinals.

Let N be a c.t.m. for $ZFC + GCH$. In N , let $\kappa < \lambda \leq \mu \leq \theta$ be cardinals such that κ, λ, μ are regular and $\text{cf}(\theta) > \kappa$. The goal is to produce an extension of N in which $\mathfrak{t}_\kappa = \lambda$, $\mathfrak{b}_\kappa = \mu$ and $2^\kappa = \theta$. Let H be $\mathbb{F}n(\theta^+ \times \lambda, 2, \lambda)$ -generic over N and let $N[H] = M$. Then

$$M \models \text{“}ZFC + \forall \xi < \lambda (2^\xi = \xi^+) + 2^\lambda = \theta^+\text{”}$$

κ, λ, μ are still regular and all the cardinals are preserved. Now, in M , perform an iterated forcing construction of length $\theta \cdot \mu$ (ordinal product) with $< \kappa$ -supports, i.e., build an iterated forcing construction

$$\langle \langle \mathbb{P}_\xi : \xi \leq \theta \cdot \mu \rangle, \langle \pi_\xi : \xi < \theta \cdot \mu \rangle \rangle$$

with supports of size less than κ each \mathbb{P}_ξ having cardinality $\leq \lambda$.

Given \mathbb{P}_ξ , if ξ is not of the form $\theta \cdot \xi$, list all the \mathbb{P}_ξ -names for towers in $(\mathcal{P}(\kappa), \subset^*)$ of size η for all $\kappa < \eta < \lambda$; for example, let $\langle \sigma_\gamma^\xi : \gamma < \theta \rangle$ enumerate all \mathbb{P}_ξ -names σ such that for some η , with $\kappa < \eta < \lambda$, σ is a nice \mathbb{P}_ξ -name for a subset of $(\eta \times \kappa)$ with the property that there is a name τ_γ^ξ such that

$$\mathbf{1} \Vdash \text{“}\tau_\gamma^\xi = \{x \subseteq \kappa : \exists \zeta < \eta (x = \{\nu : (\zeta, \nu) \in \sigma_\gamma^\xi\})\} \\ \text{is a tower in } (\mathcal{P}(\kappa), \subset^*) \text{ of size } \eta\text{”}.$$

Let $\Theta = (\theta \cdot \mu) \setminus \{\theta \cdot \xi : \xi < \mu\}$ and let $f : \Theta \rightarrow (\theta \cdot \mu) \times \theta$ be a bookkeeping function such that f is onto and $\forall \xi, \beta, \gamma (f(\xi) = (\beta, \gamma) \rightarrow \beta < \xi)$. If $f(\xi) = (\beta, \gamma)$, let τ_ξ be a \mathbb{P}_ξ -name for the same object for which τ_γ^β is a \mathbb{P}_β -name. Let π_ξ be the standard \mathbb{P}_ξ -name for \mathbb{T}_{τ_ξ} . And if ξ is of the form $\theta \cdot \zeta$, let φ_ξ be a \mathbb{P}_ξ -name for κ^κ and let π_ξ be the standard \mathbb{P}_ξ -name for \mathbb{D}_{φ_ξ} . This finishes the iteration.

By Lemma 9 $\mathbb{P}_{\theta \cdot \mu}$ is κ -closed in M . In fact, $\mathbb{P}_{\theta \cdot \mu}$ has the property that each decreasing sequence of length $< \kappa$ has a greatest lower bound so that the set \mathbb{P}' of elements $p \in \mathbb{P}_{\theta \cdot \mu}$ with the property that the first coordinate of $p(\gamma)$, for $\gamma \in \text{dom}(p)$, is a real object and not just a \mathbb{P}_γ -name, is dense in $\mathbb{P}_{\theta \cdot \mu}$. Therefore, to show that $\mathbb{P}_{\theta \cdot \mu}$ also has the κ^+ -cc in M it suffices to show that \mathbb{P}' has the κ^+ -cc in M . So, in M , let $p^\gamma \in \mathbb{P}'$ for $\gamma < \kappa^+$. By $\kappa^{<\kappa} = \kappa$, the Δ -system lemma (see Theorem II,1.6 in [Ku83]) implies that there is an $X \in [\kappa^+]^{\kappa^+}$ such that $\{\text{support}(p^\gamma) : \gamma < \kappa^+\}$ for a Δ -system with root r . Let $p^\gamma = \langle \rho_\xi^\gamma : \xi < \theta \cdot \mu \rangle$,

and let $\rho_\xi^\gamma = \text{op}(\check{s}_\xi^\gamma, \sigma_\xi^\gamma)$. By $\kappa^{<\kappa} = \kappa$, there is a $Y \in [X]^{\kappa^+}$ such that for all $\xi \in r$, the s_ξ^γ for $\gamma \in Y$ are all the same; say $s_\xi^\gamma = s_\xi$ for $\xi \in r$ and $\gamma \in Y$. But then the p^γ for $\gamma \in Y$ are pairwise compatible; to see this observe that if $\gamma, \delta \in Y$, then p^γ, p^δ have as a common extension $\langle p_\xi : \xi < \theta \cdot \mu \rangle$, where ρ_ξ is

(a) ρ_ξ^γ if $\xi \notin \text{support}(p^\delta)$, (b) ρ_ξ^δ if $\xi \notin \text{support}(p^\gamma)$, (c) $\text{op}(\check{s}_\xi, \sigma_\xi)$ if $\xi \in r$,

where σ_ξ is a nice name which satisfies $\mathbf{1}_\xi \Vdash \sigma_\xi = \sigma_\xi^\gamma \cup \sigma_\xi^\delta$. So $\mathbb{P}_{\theta \cdot \mu}$ has the κ^+ -c.c. and together with being κ -closed preserves all the cardinal numbers. Let G be $\mathbb{P}_{\theta \cdot \mu}$ -generic over M . Since at each stage of the form $\theta \cdot \xi$, a function from κ to κ is added which eventually dominates all the functions in κ^κ constructed by that stage, it follows that, in $M[G]$, there is a scale in $(\kappa^\kappa, <^*)$ of order type μ so that $\mathfrak{b}_\kappa = \mu$. In addition, since at each stage of the iteration a new element to κ^κ or $\mathcal{P}(\kappa)$ is added, it follows that $M[G] \models "2^\kappa = |\theta \cdot \mu| = \theta"$. Finally, $M[G]$ contains no towers in $(\mathcal{P}(\kappa), \subseteq^*)$ of order type η for $\kappa < \eta < \lambda$ since by the bookkeeping device all such towers are considered and eventually destroyed at some stage of the iteration, so that $\mathfrak{t}_\kappa \geq \lambda$. However, $M[G] \models "\forall \xi (\kappa \leq \xi < \lambda \rightarrow 2^\xi = \theta)"$ and $M[G] \models "2^\lambda = \theta^+"$ since $M \models "2^\lambda = \theta^+"$ and clearly \diamond_κ holds (e.g., without loss of generality $M[G] \models \diamond_\kappa$ and κ -closed forcing preserve it so that by the previous section $\mathfrak{t}_\kappa = \lambda$). This finishes the proof of this theorem. \square

References

- Bsxx Andreas Blass, *Combinatorial Cardinal Characteristics of the Continuum*, in M. Foreman, A. Kanamori, and M. Magidor, editors, *Handbook of Set Theory*, Kluwer,
 CuSh:541 James Cummings and Saharon Shelah, *Cardinal invariants above the continuum*, Ann. Pure Appl. Logic **75** (1995), 251–268.
 Ku83 Kenneth Kunen, *Set Theory: An Introduction to Independence Proofs*, volume 102 of Studies in Logic and the Foundations of Mathematics, North-Holland, Amsterdam, 1983.
 Ro39 Fritz Rothberger, *Sur un ensemble toujours de première catégorie qui est dépourvu de la propriété λ* , Fundam. Math. **32** (1939), 294–300.
 Ro48 Fritz Rothberger, *On some problems of Hausdorff and Sierpiński*, Fundam. Math. **35** (1948), 29–46.
 Sh:80 Saharon Shelah, *A weak generalization of MA to higher cardinals*, Israel J. Math. **30** (1978), 297–306.
 Sh:460 Saharon Shelah, *The Generalized Continuum Hypothesis revisited*, Israel J. Math. **116** (2000), 285–321.

Institute of Mathematics
 The Hebrew University
 Jerusalem, Israel

(Received 06 03 2000)
 (Revised 10 02 2003)

Rutgers University
 Mathematics Department
 New Brunswick, NJ USA

Department of Mathematics
 Massachusetts Institute of Technology
 Cambridge, MA 02139