

Universal graphs and functions on ω_1

Saharon Shelah^{a,1}, Juris Steprāns^{b,2}

^a*Department of Mathematics, Rutgers University, Hill Center, Piscataway, New Jersey, U.S.A. 08854-8019*
and

*Institute of Mathematics, Hebrew University
Givat Ram, Jerusalem 91904, Israel*

^b*Department of Mathematics, York University, 4700 Keele Street, Toronto, Ontario, Canada M3J 1P3*

Abstract

It is shown to be consistent with various values of \mathfrak{b} and \mathfrak{d} that there is a universal graph on ω_1 . Moreover, it is also shown that it is consistent that there is a universal graph on ω_1 — in other words, a universal symmetric function from ω_1^2 to 2 — but no such function from ω_1^2 to ω . The method used relies on iterating well known reals, such as Miller and Laver reals, and alternating this with the PID forcing which adds no new reals.

1. Introduction

1.1. Background to the problem

It is well known that countably saturated models are universal for models of cardinality \aleph_1 . However, in the case of graphs, the existence of a saturated model is equivalent to $2^{\aleph_0} = \aleph_1$. These two observations immediately raise the question of whether it is possible to have a universal graph of cardinality \aleph_1 in the absence of the Continuum Hypothesis. This provided the motivation for the articles [1], [2] and [3] which solved not only this question, but also others about the universality of different structures.

While the results to be presented here have their roots in this work, they are also motivated by considerations that are not model theoretic. A function $U : X \times X \rightarrow X$ is said to be (*Sierpiński*) *universal* if for any $G : X \times X \rightarrow X$ there is $e : X \rightarrow X$ such that $G(x, y) = U(e(x), e(y))$ for all x and y in X . The function e will be called an *embedding* of G into U . An early reference to this notion can be found in Problem 132 of the Scottish Book [4] in which Sierpiński asked if there is a Borel function which is universal in this sense, when X is the real line. He had already shown in [5] that there is a Borel universal function assuming the Continuum Hypothesis. This notion of universal function is also studied in Rado [6]. More recently, this notion and various generalizations of it were studied in [7], in which a restricted form of the following definition appears as Definition 7.5.

Definition 1.1. A function $U : \kappa \times \kappa \rightarrow \lambda$ is *weakly universal* if for every $f : \kappa \times \kappa \rightarrow \lambda$ there exist one-to-one functions $h : \kappa \rightarrow \kappa$ and $k : \lambda \rightarrow \lambda$ such that $k(f(\alpha, \beta)) = U(h(\alpha), h(\beta))$ for all α and β in κ . The pair (h, k) will be called a weak embedding.

Definition 7.4 of [7] defines a function $U : \kappa \times \kappa \rightarrow \kappa$ to be *model theoretically universal* if it is weakly universal, as in Definition 1.1, but with $h = k$. Note that U is Sierpiński universal if it is weakly universal with k being the identity. Remark 7.7 of [7] claims that all three notions — Sierpiński universal, weakly universal, and model theoretically universal — are equivalent for maps into 2.

The following is Theorem 5.9 of [7] showing that there is no difference between asking about the existence of universal graphs — in other words, symmetric, irreflexive functions from ω_1^2 to 2 — and non-symmetric functions from ω_1^2 to 2.

Theorem 1.2. *For any infinite cardinal κ the following are equivalent:*

1. *For each $n \in \mathbb{N}$ there is a universal function from $\kappa \times \kappa$ to n .*

Email addresses: shelah@math.rutgers.edu (Saharon Shelah), steprans@yorku.ca (Juris Steprāns)

¹The first author's research for this paper was partially supported by the United States-Israel Binational Science Foundation (grant no. 2006108), and by the National Science Foundation (grant no. NSF-DMS 0600940). This article has been assigned number 1088 in the first author's list of publications that can be found at <http://shelah.logic.at>

²The second author's research for this paper was partially supported by NSERC of Canada.

2. For some $n \in \mathbb{N}$ with $n \geq 2$ there is a universal function from $\kappa \times \kappa$ to n .
3. There is a symmetric, irreflexive function from $\kappa \times \kappa$ to 2 universal for all symmetric, irreflexive functions from $\kappa \times \kappa$ to 2
4. There is a universal graph on κ .

Given Theorem 1.2 it is reasonable to focus attention only on symmetric functions from ω_1^2 to ω and this will be done from now on. Moreover, given that there are only two possible values for k when $\lambda = 2$, the validity of Remark 7.7 of [7] should now be clear and, of course, assuming the Continuum Hypothesis there is a Sierpiński universal function from \mathbb{R}^2 to \mathbb{R} , and hence there is also a weakly universal function from ω_1^2 to 2. Problem 7.8 of [7] asks if the existence of one type of universal function implies the existence of the others in general. To provide a negative answer to the question, it is therefore necessary to consider models where the Continuum Hypothesis fails. This provides an other path to the questions considered in [1], [2] and [3].

When $\kappa = \omega_1$ and $\lambda = 2$ it was shown in [1] and [2] that it is consistent with the failure of the Continuum Hypothesis that there is a symmetric universal function $U : \omega_1^2 \rightarrow 2$ which satisfies all three universality properties for symmetric functions. The methods used in the generalization of this result by Mekler to other theories in [3] provide examples of Sierpinski universal functions $U : \omega_1^2 \rightarrow \lambda$ for λ equal to 2, ω or ω_1 . In the models of [1] and [2] the cardinal invariants \mathfrak{b} and \mathfrak{d} have the following values respectively: \aleph_1 and \aleph_2 . One might, therefore ask, whether these values are needed for the existence of universal functions from $U : \omega_1^2 \rightarrow \lambda$ with the failure of the Continuum Hypothesis. This question becomes even more interesting in light of the positive results to be presented in Lemma 6.1 and Lemma 7.8 which rely on small values of \mathfrak{b} and \mathfrak{d} respectively. It will be shown in Corollary 3.10 that it is consistent with the existence of universal functions from $U : \omega_1^2 \rightarrow 2$ that $\mathfrak{b} = \mathfrak{d} = \aleph_2$ and in Corollary 4.14 that it is consistent with the existence of universal functions from $U : \omega_1^2 \rightarrow 2$ that $\mathfrak{b} = \mathfrak{d} = \aleph_1$.

However, the main goal of this paper will be to answer Problem 5.10 of [7]. This is done in Corollary 6.2 which establishes that there is a Sierpiński universal function from ω_1^2 to 2 but no such function from ω_1^2 to ω . It is worth recalling that Theorem 5.9 of [7] asserts that if $2 \leq n < \omega$ then there is a Sierpiński universal function from ω_1^2 to n if and only if there is a Sierpiński universal function from ω_1^2 to n . A solution to Problem 7.8 of [7] will also be presented under the assumption that \mathfrak{d} is small. This and related questions are discussed in §7. That section also contains some arguments extending the methods of [1] to some weak versions of embedding. Finally, if one is only interested in obtaining a model of set theory in which the Continuum Hypothesis fails, yet there is a universal graph of cardinality \aleph_1 , then §2 presents an easier argument than the original of [1].

1.2. Notation and terminology

Since trees will play a central role in the following discussion, it may be worthwhile reviewing some notation and terminology, even though most of this is standard and almost all of the notation used will follow that of Sections 1.1.D and 7.3.D of [8]. By a tree T will be meant a subset $T \subseteq \omega^{<\omega} = \bigcup_{n \in \omega} \omega^n$ that is closed under initial segments — in other words, if $t \in T$ and $k \leq |t|$ then $t \upharpoonright k \in T$. If T is a tree and $t \in T$ then $T[t]$ will denote the tree defined by

$$T[t] = \{s \in T \mid s \subseteq t \text{ or } t \subseteq s\}$$

and $\text{succ}_T(t)$ will denote the set $\{s \in T \mid s \supseteq t \text{ and } |s| = |t| + 1\}$.

A tree T will be called *infinite splitting* if $|\text{succ}_T(t)| \in \{1, \aleph_0\}$ for each $t \in T$. Define $\text{split}(T) = \{t \in T \mid |\text{succ}_T(t)| = \aleph_0\}$ and define

$$\text{split}_n(T) = \{t \in \text{split}(T) \mid |\{k \in |t| \mid t \upharpoonright k \in \text{split}(T)\}| = n\}.$$

If T is infinite splitting then let $\Psi_T : \omega^{<\omega} \rightarrow \text{split}(T)$ be the unique bijection from $\omega^{<\omega}$ to $\text{split}(T)$ preserving the lexicographic ordering. For $t \in \omega^{<\omega}$ let $T\langle t \rangle = T[\Psi_T(t)]$ — the reader is warned that this notation differs from [7]. Hence $\text{stem}(T)$ can be defined to be $\Psi_T(\emptyset)$ for infinite splitting trees T .

Let $\{u_i\}_{i \in \omega}$ enumerate $\omega^{<\omega}$ in such a way that if $k < |u_i|$ then there is $j \in i$ such that $u_i \upharpoonright k = u_j$. Then for infinite splitting trees T and S the ordering \leq_n is defined by $T \leq_n S$ if $T \subseteq S$ and $\Psi_S(u_j) = \Psi_T(u_j)$ for all $j \in n$.

Finally, recall that Miller forcing, or rational perfect set forcing, is denoted by \mathbf{PT} in [8] and consists of all infinite branching trees ordered by inclusion. Laver forcing, on the other hand, is denoted by \mathbf{LT} in [8] and consists of all infinite branching trees such that $T \setminus \text{split}(T)$ is finite, also ordered by inclusion. In the case of Laver forcing the notion of a *front* is useful: If $T \in \mathbf{LT}$ then $W \subseteq T$ is a front if it consists of incomparable elements of T and every maximal branch of T contains an element of W .

1.3. The universal graphs

The graphs that will be shown to be universal in the arguments to follow will all come from some initial model of set theory in which the Continuum Hypothesis holds.

Definition 1.3. Given any function $G : \omega_1^2 \rightarrow \omega$ and $\eta \in \omega_1$ define $G^\eta : \eta \rightarrow \omega$ by $G^\eta(\zeta) = G(\zeta, \eta)$ and then define $S^\eta(G) = \{G^\mu \upharpoonright \eta\}_{\mu \geq \eta}$. A function $G : \omega_1^2 \rightarrow \omega$ such that $S^\eta(G)$ is everywhere non-meagre for each $\eta \geq \omega$ will be called *category saturated*.

If one does not wish to restrict to irreflexive functions then one can define

$$S^{\eta,k}(G) = \{G^\mu \upharpoonright \eta \mid \mu \geq \eta \text{ and } G(\mu, \mu) = k\}$$

and then ask that $S^{\eta,k}(G)$ is everywhere non-meagre for each $\eta \geq \omega$ for each $k \in \omega$. Since this added complication currently provides no new insights, this work will focus on irreflexive functions only.

Definition 1.4. Let ν be an atomic probability measure on ω and let ν^η be the Fubini product of this measure on ω^η for any $\eta \in \omega_1$. A function $G : \omega_1^2 \rightarrow \omega$ such that $S^\eta(G)$ has outer measure 1 for each $\eta \geq \omega$ will be called *ν -saturated*. The notion defined here will not be needed in this full generality. For the purposes of this article a function $G : \omega_1^2 \rightarrow \omega$ will be called *measure saturated* if it is ν -saturated where ν is the measure on 2 giving each point equal measure.

Lemma 1.5. *Assuming the Continuum Hypothesis, there is a symmetric function from ω_1^2 to ω that is category saturated and ν -saturated for every atomic probability measure ν .*

Indeed, using the Continuum Hypothesis it is easy to construct a symmetric function $G : \omega_1^2 \rightarrow \omega$ such that $S^\eta(G) = \omega^\eta$ for each $\eta \in \omega_1$. Note, however, that adding a real will destroy this stronger property; nevertheless, in certain generic extensions the weaker properties of being category saturated or ν -saturated may persist.

2. Category saturated graphs are universal after adding Miller reals

2.1. A P -ideal from a class of names

This section will describe an ideal that arises in generic extensions in which reals are added. It can easily, but inaccurately, be described as the collection of all sets in the extension that have a name that is a continuous function whose range is almost disjoint.

Definition 2.1. If $G \subseteq \mathbf{PT}$ is generic over V define $\mathcal{S}(\mathbf{PT})$ to be the set of all $S \in [\omega_1]^{\aleph_0}$ such that there is $T \in G$ and $\psi : T \rightarrow [\omega_1]^{<\aleph_0}$ such that

1. $\psi \in V$
2. $\psi(s) \cap \psi(t) = \emptyset$ unless $s = t$
3. $T \Vdash_{\mathbf{PT}} "S = \bigcup_{j=0}^{\infty} \psi(r_G \upharpoonright j)"$ where $r_G : \omega \rightarrow \omega$ is the generic real obtained from the generic set G .

Functions satisfying (2) will be said to have *disjoint range*. Given any $\psi : T \rightarrow [\omega_1]^{<\aleph_0}$ let S_ψ be a \mathbf{PT} -name for $\bigcup_{j=0}^{\infty} \psi(r_G \upharpoonright j)$ where $r_G = \bigcup_{T \in G} \mathbf{stem}(T)$ and G is the generic set.

Lemma 2.2. *If $S \in \mathcal{S}(\mathbf{PT})$ then there is $T \in G$ and $\psi : T \rightarrow [\omega_1]^{<\aleph_0}$ such that $\psi(t) = \emptyset$ unless $t \in \mathbf{split}(T)$ and $T \Vdash_{\mathbf{PT}} "S = S_\psi"$.*

PROOF. Given $S \in \mathcal{S}(\mathbf{PT})$ it suffices to show that for $T \in \mathbf{PT}$ there is $\bar{T} \subseteq T$ and $\psi : \bar{T} \rightarrow [\omega_1]^{<\aleph_0}$ such that $\psi(t) = \emptyset$ unless $t \in \mathbf{split}(\bar{T})$ and $\bar{T} \Vdash_{\mathbf{PT}} "S = S_\psi"$. To this end use the definition of $\mathcal{S}(\mathbf{PT})$ to find T^* , \hat{T} and $\hat{\psi} : \hat{T} \rightarrow [\omega_1]^{<\aleph_0}$ such that $T^* \Vdash_{\mathbf{PT}} "S = S_{\hat{\psi}}"$ and $\hat{T} \in G$. Observe that $\bar{T} = T^* \cap \hat{T} \in \mathbf{PT}$ because otherwise it would follow that $T^* \Vdash_{\mathbf{PT}} "\hat{T} \notin G"$.

Then define $\psi : \bar{T} \rightarrow [\omega_1]^{<\aleph_0}$ by

$$\psi(t) = \begin{cases} \bigcup \left\{ \hat{\psi}(t \upharpoonright k) \mid (\forall j \in [k, |t|]) t \upharpoonright j \notin \mathbf{split}(\bar{T}) \right\} & \text{if } t \in \mathbf{split}(\bar{T}) \\ \emptyset & \text{otherwise.} \end{cases}$$

It should be obvious that $\psi : \bar{T} \rightarrow [\omega_1]^{<\aleph_0}$ has disjoint range and that $\bar{T} \Vdash_{\mathbf{PT}} "S_\psi = S_{\hat{\psi}} = S"$.

Lemma 2.3. *Let $G \subseteq \mathbf{PT}$ be generic over V and $S \subseteq \omega_1$ in $V[G]$. Then the following are equivalent:*

1. $S \in \mathcal{S}(\mathbf{PT})$
2. for all $F : \omega \times \omega \rightarrow \omega_1$ in V such that the mapping $j \mapsto F(n, j)$ is one-to-one for all $n \in \omega$, there is $g : \omega \rightarrow \omega$ in V such that for all $n \in \omega$ there is $k \leq g(n)$ such that $F(n, k) \notin S$
3. for all $F : \omega \times \omega \rightarrow \omega_1$ in V such that the mapping $j \mapsto F(n, j)$ is one-to-one for all $n \in \omega$, there is a one-to-one function $g : \omega \times \omega \rightarrow \omega$ in V such that $F(n, g(n, m)) \notin S$ for all $(n, m) \in \omega \times \omega$.

PROOF. To see that (1) implies (3) suppose that $S \in \mathcal{S}(\mathbf{PT})$. Using Lemma 2.2 find

$$\psi : \mathbf{split}(T) \rightarrow [\omega_1]^{<\aleph_0}$$

such that $T \Vdash_{\mathbf{PT}} "S = S_\psi"$ and $\psi(t) = \emptyset$ unless $t \in \mathbf{split}(T)$. Let $F : \omega \times \omega \rightarrow \omega_1$ belong to V such that the mapping $j \mapsto F(n, j)$ is one-to-one for all $n \in \omega$. Let $\{(i_n, j_n)\}_{n \in \omega}$ enumerate $\omega \times \omega$ and construct trees T_n and distinct integers $g(i_n, j_n)$ such that

1. $T_0 = T$
2. $T_{n+1} \leq_n T_n$
3. $F(\ell, g(i_\ell, j_\ell)) \notin \bigcup_{i \in n} \bigcup_{j \leq |\Psi_{T_n}(u_i)|} \psi(\Psi_{T_n}(u_i) \upharpoonright j)$ for all $\ell \in n$.

To construct T_{n+1} first choose $g(i_n, j_n)$ such that $g(i_n, j_n) \neq g(i_\ell, j_\ell)$ for $\ell \in n$ and such that

$$F(i_n, g(i_n, j_n)) \notin \bigcup_{i \in n} \bigcup_{j \leq |\Psi_{T_n}(u_i)|} \psi(\Psi_{T_n}(u_i) \upharpoonright j).$$

Then note that for each $k \leq n$ there is at most one $s_k \in T_n$ such that $F(i_k, g(i_k, j_k)) \in \psi(s_k)$ and none of these s_k is below any $\Psi_{T_n}(u_\ell)$ for $\ell \in n$. It is therefore easy to find $T_{n+1} \subseteq T_n$ such that $\Psi_{T_{n+1}}(u_n) \neq s_i$ for all $i \leq n$ and, hence, the induction hypotheses are satisfied. Let $T^* = \bigcap_n T_n$ and observe that the function $g : \omega \times \omega \rightarrow \omega$ defined by the induction belongs to V and that

$$T^* \Vdash_{\mathbf{PT}} "(\forall n \in \omega) F(n, g(i_n, j_n)) \notin S".$$

To see that (2) implies (1) suppose that S satisfies (2). Let $S = \{\sigma_j\}_{j \in \omega}$ be an enumeration of S and let $T \in G$ be such that there is $\psi : \mathbf{split}_n(T) \rightarrow \omega_1$ such that $T_t \Vdash_{\mathbf{PT}} "\sigma_n = \psi(t)"$ for each $t \in \mathbf{split}_n(T)$. Now let B be the set of $t \in \mathbf{split}(T)$ such that there is a family $\{w_i^t\}_{i \in \omega} \subseteq \mathbf{split}(T)$ such that

1. $w_i^t \supseteq t$
2. $w_i(|t|) \neq w_j(|t|)$ unless $i = j$
3. there is $\theta_n \in \prod_{i=n}^\infty (|t|, |w_i^t|)$ and ρ_n for each $n \in \omega$ such that if $n \leq i$ then $\psi(w_i^t \upharpoonright \theta_n(i)) = \rho_n$
4. $\rho_n \neq \rho_m$ unless $n = m$.

It will first be shown that B cannot be dense in T .

For if B were dense in T then it would be possible to find $\bar{T} \subseteq T$ such that $\bar{T} \in \mathbf{PT}$, $\mathbf{split}(\bar{T}) \subseteq B$ and if $t \in \mathbf{split}_m(\bar{T})$ and $\{w_i^t\}_{i \in \omega}$ enumerates $\{s \in \mathbf{split}_{m+1}(\bar{T}) \mid s \supseteq t\}$ then there are $\theta_n^t \in \prod_{i=n}^\infty (|t|, |w_i^t|)$ and ρ_n^t for each $n \in \omega$ witnessing that $t \in B$. Let $\{t_i\}_{i \in \omega}$ enumerate $\mathbf{split}(\bar{T})$ and define $F : \omega \times \omega \rightarrow \omega_1$ by $F(m, n) = \rho_n^{t_m}$.

Now given any $T^* \subseteq \bar{T}$ such that $T^* \in \mathbf{PT}$ and any $g \in \omega^\omega \cap V$ it follows that $\mathbf{stem}(T^*) = t_m$ for some m . There is then some $n \in \omega$ and some t^* such that $t^* \supseteq w_n^{t_m} \supseteq t_m$ and $n > g(m)$. Then $T_{t^*}^* \Vdash_{\mathbf{PT}} "\{\rho_i^{t_m}\}_{i=0}^{g(m)+1} \subseteq S"$ yielding a contradiction to the hypothesis on S .

Hence it will be assumed that $B = \emptyset$. Construct a sequence $T = T_0 \geq_0 T_1 \geq_1 T_2 \dots$ such that for each $k \in n$ there is a family $\{w_i^k\}_{i \in \omega}$ satisfying:

1. $w_i^k \supseteq \Psi_{T_n}(u_k)$
2. $w_i^k(|\Psi_{T_n}(u_k)|) \neq w_m^k(|\Psi_{T_n}(u_k)|)$ unless $i = m$

3. there is some $J_k \in \omega$ and $\theta_j \in \prod_{i=j}^{\infty} (|\Psi_{T_n}(u_k)|, |w_i^k|)$ and distinct ρ_j^k for each $j \in J_k$ such that if $j \leq i$ then $\psi(w_i^k \upharpoonright \theta_j(i)) = \rho_j^k$ and, moreover, the set $\{\rho_j^k\}_{j \in J_k}$ is maximal with this property.

Completing this induction is easy using the fact that $T_n \cap B = \emptyset$. Let $T^* = \bigcap_n T_n$ and let $\bar{\psi}(\Psi_{T^*}(u_k)) = \{\rho_j^k\}_{j \in J_k}$.

Next observe that there is $T^{**} \subseteq T^*$ such that $\mathbf{split}(T^{**}) = \mathbf{split}(T^*) \cap T^{**}$ and

$$(\forall s \in T_n \setminus \{\Psi_{T_n}(u_k) \mid k \in n\}) \psi(s) \notin \bigcup_{k \in n} \bar{\psi}(\Psi_{T_n}(u_k)) \quad (2.1)$$

$$(\forall k < m < n) \bar{\psi}(\Psi_{T_n}(u_k)) \cap \bar{\psi}(\Psi_{T_n}(u_m)) = \emptyset \quad (2.2)$$

To see this construct a sequence $T = T_0 \geq_0 T_1 \geq_1 T_2 \dots$ such that $\bar{\psi}(\Psi_{T_n}(u_i)) \cap \bar{\psi}(\Psi_{T_n}(u_j)) = \emptyset$ for $i < j < n$. Given T_n , let j be such that u_j is maximal from the set $\{u_i \mid i < n \text{ and } u_i \subseteq u_n\}$. There cannot be infinitely many $s \in \mathbf{succ}_{T_n}(\Psi_{T_n}(u_j))$ such that there is $\bar{s} \in T_n[s]$ such that $\psi(\bar{s}) \in \bigcup_{k \in n} \bar{\psi}(\Psi_{T_n}(u_k)) \neq \emptyset$ because this would violate the maximality of $\bar{\psi}(\Psi_{T_n}(u_j))$. This makes it easy to find T_{n+1} . Then $\bar{\psi} : T^{**} \rightarrow [\omega_1]^{<\aleph_0}$ witness that $S \in \mathcal{S}(\mathbf{PT})$.

To see that (3) implies (2) let $g(n) = g(n, 0) + 1$.

Corollary 2.4. $\mathcal{S}(\mathbf{PT})$ is closed under subsets and finite unions. Moreover if $S \in \mathcal{S}(\mathbf{PT})$ then S contains no infinite set from V .

PROOF. The fact that $\mathcal{S}(\mathbf{PT})$ is closed under subsets follows from (3) of Lemma 2.3. To see that $\mathcal{S}(\mathbf{PT})$ is closed under finite unions, let S_0 and S_1 belong to $\mathcal{S}(\mathbf{PT})$. Given $F : \omega \times \omega \rightarrow \omega_1$ let $g_0 : \omega \times \omega \rightarrow \omega$ witness (3) for S_0 . Then, define $\tilde{F}(i, j) = F(i, g_0(i, j))$ and apply (3) of Lemma 2.3 to \tilde{F} and S_1 to get $g_1 : \omega \times \omega \rightarrow \omega$. Then g_1 also witnesses (3) for $S_0 \cup S_1$ for the given function F .

To see the final claim let $X \in [\omega_1]^{\aleph_0}$ belong to V and let $S \in \mathcal{S}(\mathbf{PT})$. Then let $F : \omega \times \omega \rightarrow \omega_1$ be such that $\{F(i, j)\}_{j \in \omega}$ enumerates X on each $i \in \omega$. It follows from (2) that $X \not\subseteq S$.

It will be useful to observe that it is possible to isolate part of the proof of Lemma 2.3 to show the following.

Corollary 2.5. If $T \in \mathbf{PT}$ and $\psi_i : T \rightarrow [\omega_1]^{<\aleph_0}$ have disjoint range for $i \in k$ then there is $T^* \subseteq T$ such that $\mathbf{split}(T^*) = \mathbf{split}(T) \cap T^*$ and the function ψ defined by $\psi(t) = \bigcup_{i \in k} \psi_i(t)$ has disjoint range.

PROOF. The key point is that at each stage of a fusion argument only finitely many nodes of the tree need to be eliminated.

Question 2.6. If G is \mathbf{PT} generic over V does it follow that $\mathcal{S}(\mathbf{PT})$ is equal to the

$$\{X \in [\omega_1]^{\aleph_0} \mid \mathcal{P}(X) \cap V = [X]^{<\aleph_0}\}$$

in $V[G]$?

Lemma 2.7. If $\psi_i : T_i \rightarrow [\omega_1]^{<\aleph_0}$ have disjoint range for $i \in \omega$ then there are $\bar{T}_i \subseteq T_i$ such that

$$\mathbf{split}(\bar{T}_i) = \mathbf{split}(T_i) \cap \bar{T}_i \quad (2.3)$$

$$(\forall i < j < \omega)(\forall t \in \bar{T}_i)(\forall s \in \bar{T}_j) \text{ if } t \supseteq \mathbf{stem}(T_i) \text{ and } s \supseteq \mathbf{stem}(T_j) \text{ then } \psi_j(s) \cap \psi_i(t) = \emptyset \quad (2.4)$$

PROOF. This is a routine fusion argument. Using Lemma 2.2 it may be assumed that $\psi_i(t) = \emptyset$ unless $t \in \mathbf{split}(T_i)$. The key point is that given $n \in \omega$ and $T_{n,i} \subseteq T_i$ for $i \in \omega$ there are $T_{n+1,j}$ for $j \in \omega$ such that

- $T_{n+1,j} \leq_n T_{n,j}$ for $j \in n$
- $T_{n+1,j} \subseteq T_{n,j}$ for $j \in \omega$
- $\mathbf{split}(T_{n+1,j}) = \mathbf{split}(T_{n,j}) \cap T_{n+1,j}$ for $j \in \omega$
- $T_{n,j} \langle u_i \rangle = T_{n+1,j}[\Psi_{T_{n,j}}(u_i)]$ for $j \leq n$ and $i \in n$
- $\psi_i(\Psi_{T_{n,i}}(u_m)) \cap \psi_j(\Psi_{T_{n,j}}(u_k)) = \emptyset$ if $i < j \leq n$ and $k, m \in n$.

To find $\Psi_{T_{n+1,j}}(u_n)$ for $j \leq n$ use the fact that each ψ_i has disjoint range and each $T_{n,j}$ is infinite branching to eliminate the finitely many nodes violating this requirement — only $\{\mathbf{stem}(T_i)\}_{i \in \omega}$ cannot be omitted.

Lemma 2.8. *If S is \mathbf{PT} -name such that $T \Vdash_{\mathbf{PT}} "S \in \mathcal{S}(\mathbf{PT})"$ then there is $\bar{T} \subseteq T$ and $\psi : \bar{T} \rightarrow [\omega_1]^{<\aleph_0}$ such that*

- $\text{stem}(\bar{T}) = \text{stem}(T) = t^*$
- ψ has disjoint range
- $\psi(t^* \upharpoonright j) = \emptyset$ for $j \leq |t^*|$
- $\bar{T} \Vdash_{\mathbf{PT}} "S \equiv^* S_\psi"$.

PROOF. For each $t \in \text{succ}_T(t^*)$ use Lemma 2.2 to find $T_t \leq T[t]$ and ψ_t such that

- $\psi_t : T_t \rightarrow [\omega_1]^{<\aleph_0}$ has disjoint range
- $\psi_t(t \upharpoonright j) = \emptyset$ for $j \leq |t|$
- $T_t \Vdash_{\mathbf{PT}} "S \equiv^* S_{\psi_t}"$.

Now apply Lemma 2.7 to the family $\{\psi_t\}_{t \in \text{succ}_T(t^*)}$ to find $\tilde{T}_t \subseteq T_t$ such that ψ defined by

$$\psi(s) = \begin{cases} \psi_t(s) & \text{if } (\exists t \in \text{succ}_T(t^*)) s \supseteq t \text{ and } s \in \tilde{T}_t \\ \emptyset & \text{if } (\exists t \in \text{succ}_T(t^*)) s \subseteq t \end{cases}$$

has disjoint range. Let $\bar{T} = \bigcup_{t \in \text{succ}_T(t^*)} \tilde{T}_t$. It is immediate that ψ and \bar{T} satisfy the lemma.

Lemma 2.9. *If G is \mathbf{PT} generic over V then $\mathcal{S}(\mathbf{PT})$ is a P -ideal in $V[G]$.*

PROOF. Suppose that $\{S_n\}_{n \in \omega}$ are \mathbf{PT} -names such that $T \Vdash_{\mathbf{PT}} "S_n \in \mathcal{S}(\mathbf{PT})"$ for each n . If it is possible to construct T_n and ψ_n such that

1. $T_0 = T$
2. $T_{n+1} \leq_n T_n$
3. $\psi_n : T_n \rightarrow [\omega_1]^{<\aleph_0}$ has disjoint range
4. $\psi_n(\Psi_{T_n}(u_i) \upharpoonright j) = \psi_{n+1}(\Psi_{T_n}(u_i) \upharpoonright j)$ if $i \in n$ and $j \leq |\Psi_{T_n}(u_i) \upharpoonright j|$
5. if $t \in T_{n+1}$ then $\psi_n(t) \subseteq \psi_{n+1}(t)$
6. $(\forall i \leq n) T_n \Vdash_{\mathbf{PT}} "S_i \subseteq^* S_{\psi_n}"$.

Then let $\bar{T} = \bigcap_{k \in \omega} T_k$ and let ψ be defined by $\psi(t) = \bigcup_k \psi_k(t)$ noting that this is finite by (4). Then

$$\bar{T} \Vdash_{\mathbf{PT}} "S_\psi \in \mathcal{S}(\mathbf{PT}) \text{ and } (\forall j) S_\psi \supseteq^* S_j".$$

To complete the construction suppose that T_n and ψ_n are given. For $i \in n$ let

$$U_i = \{t \in T \langle u_i \rangle \mid (\forall j \in n) \text{ if } u_j \subseteq t \text{ then } u_j \subseteq u_i\}.$$

Use Lemma 2.8 to find $T_i^* \subseteq U_i$ and $\psi_i^* : T_i^* \rightarrow [\omega_1]^{<\aleph_0}$ such that

- $\text{stem}(T_i^*) = \Psi_{T_n}(u_i)$
- ψ_i^* has disjoint range
- $\psi_i^*(\Psi_{T_n}(u_i) \upharpoonright j) = \emptyset$ for $j \leq |\Psi_{T_n}(u_i)|$
- $T_i^* \Vdash_{\mathbf{PT}} "S_n \equiv^* S_{\psi_i^*}"$.

Use Corollary 2.5 for each $i \in n$ to find $T_i^{**} \subseteq T_i^*$ such that $\mathbf{stem}(T_i^{**}) = \Psi_{T_n}(u_i)$ and the function ψ_i^{**} defined by $\psi_i^{**}(s) = \psi_i^*(s) \cup \psi_n(s)$ has disjoint range. Then apply Lemma 2.7 to find $\tilde{T}_i \subseteq T_i^{**}$ such that $\mathbf{stem}(\tilde{T}_i) = \Psi_{T_n}(u_i)$ and

$$(\forall i < j < n)(\forall t \in \tilde{T}_i)(\forall s \in \tilde{T}_j) \text{ if } t \supseteq \Psi_{T_n}(u_i) \text{ and } s \supseteq \Psi_{T_n}(u_j) \text{ then } \psi_j^{**}(s) \cap \psi_i^{**}(t) = \emptyset.$$

Since this only requires a finite amount of further pruning, it may be assumed that

$$\psi_i^{**}(t) \cap \left(\bigcup_{j \in n} \left(\bigcup_{k \leq |\Psi_{T_n}(u_j)|} \psi_n(\Psi_{T_n}(u_j) \upharpoonright k) \right) \right) = \emptyset$$

for each $i \in n$ and $t \in \tilde{T}_i$. Now let $T_{n+1} = \bigcup_{i \in n} \tilde{T}_i$ and define ψ_{n+1} by

$$\psi_{n+1}(t) = \begin{cases} \psi_n(t) & \text{if } (\exists i \in n) t \subseteq \Psi_{T_n}(u_i) \\ \psi_n(t) \cup \psi_i^{**}(t) & \text{if } (\exists i \in n) t \supseteq \Psi_{T_n}(u_i) \end{cases}$$

and check that this satisfies the induction hypotheses.

Lemma 2.10. *If G is \mathbf{PT} generic over V and $S \in \mathcal{S}(\mathbf{PT})$ and $f : S \rightarrow 2$ is a function in $V[G]$ then there is $T \in G$ and $\psi \in V$ defined on T with disjoint range and $f^* \in V$ such that*

1. $T \Vdash_{\mathbf{PT}} "S = S_\psi"$
2. $f^* : \bigcup_{t \in T} \psi(t) \rightarrow 2$
3. if $t \in T$ then $T[t] \Vdash_{\mathbf{PT}} "f \upharpoonright \psi(t) = f^* \upharpoonright \psi(t)"$ and, hence, $T \Vdash_{\mathbf{PT}} "f = f^* \upharpoonright S"$.

PROOF. It suffices to show that the set of $T \in \mathbf{PT}$ satisfying Conditions (1) to (3) is dense. Given $T \in \mathbf{PT}$ use Lemma 2.2 to find $\bar{T} \subseteq T$ and ψ such that $\bar{T} \Vdash_{\mathbf{PT}} "S = S_\psi"$ and such that $\psi(t) = \emptyset$ unless $t \in \mathbf{split}(\bar{T})$. Now construct T_n and f_n^* such that:

- $T_0 = \bar{T}$
- $T_{n+1} \leq_n T_n$
- the domain of f_n^* is $\bigcup_{j \in n} \psi(\Psi_{T_n}(u_j))$
- if $j \in n$ then $T_n \langle u_j \rangle \Vdash_{\mathbf{PT}} "f_n^* \upharpoonright \psi(\Psi_{T_n}(u_j)) \subseteq f"$.

Then if $T_\omega = \bigcap_{n \in \omega} T_n$ it is clear that $\psi \upharpoonright T_\omega$ and $f^* = \bigcup_{n \in \omega} f_n^*$ witness that T_ω satisfies Conditions (1) to (3) of the lemma.

To complete the induction it suffices to note that $T_n \langle u_n \rangle \Vdash_{\mathbf{PT}} "\psi(\Psi_{T_n}(u_n)) \subseteq S"$ and hence for each $s \in \mathbf{succ}_{T_n}(\Psi_{T_n}(u_n))$ there is an extension $T^s \subseteq T_n[s]$ and f_s such that $T^s \Vdash_{\mathbf{PT}} "f \upharpoonright \psi(\Psi_{T_n}(u_n)) = f_s"$. Now choose T_{n+1} such that $T_{n+1} \leq_n T_n$ and there is $\bar{f} : \psi(\Psi_{T_n}(u_n)) \rightarrow 2$ such that $f_s = \bar{f}$ for all $s \in \mathbf{succ}_{T_{n+1}}(\Psi_{T_n}(u_n))$. Let $f_{n+1}^* = f_n^* \cup \bar{f}$.

Lemma 2.11. *If G is \mathbf{PT} generic over V and $S \in \mathcal{S}(\mathbf{PT})$, $S \subseteq \xi \in \omega_1$, $f : S \rightarrow 2$ is a function in $V[G]$ and $Z \subseteq 2^\xi$ is nowhere meagre, then there is $z \in Z$ such that $f \subseteq z$.*

PROOF. Given f find T , ψ and f^* satisfying Conditions (1) to (3) of Lemma 2.10. Using Lemma 2.2 it may be assumed that $\psi(t) = \emptyset$ if $t \notin \mathbf{split}(T)$. Let $\bar{f} = f^* \upharpoonright \psi(\mathbf{stem}(T))$ and let \mathcal{O} be the open set $\{h \in 2^\xi \mid h \supseteq \bar{f}\}$. Then $Z \cap \mathcal{O}$ is not meagre in \mathcal{O} . Note that

$$\mathcal{O} \cap \{h \in 2^\xi \mid (\forall k \in \omega)(\forall t \in \mathbf{split}_k(T)) \mid \{s \in \mathbf{split}_{k+1}(T) \mid s \supseteq t \text{ and } f^* \upharpoonright \psi(s) \subseteq h\} = \aleph_0\}$$

is a dense G_δ in \mathcal{O} and hence there is some $z \in Z$ such that $\bar{f} \subseteq z$ and

$$(\forall k \in \omega)(\forall t \in \mathbf{split}_k(T)) \mid \{s \in \mathbf{split}_{k+1}(T) \mid s \supseteq t \text{ and } f^* \upharpoonright \psi(s) \subseteq h\} = \aleph_0$$

It follows that there is $T^* \subseteq T$ such that $T^* \in \mathbf{PT}$ and such that if $t \in \mathbf{split}(T^*)$ then $f^* \upharpoonright \psi(t) \subseteq z$ and hence $f^* \subseteq z$. It follows that $T^* \Vdash_{\mathbf{PT}} "f \subseteq z"$.

2.2. Applying the P-ideal dichotomy

This section will use some results from [9]. (The strengthened results of [10] will not be needed here). Recall that if \mathcal{I} is an ideal on a set X then $Y \subseteq X$ is said to be orthogonal to \mathcal{I} if $[Y]^{\aleph_0} \cap \mathcal{I} = \emptyset$.

Lemma 2.12. *If G is \mathbf{PT} generic over V then no uncountable subset of ω_1 in $V[G]$ is orthogonal to $\mathcal{S}(\mathbf{PT})$.*

PROOF. Suppose that Z is a \mathbf{PT} -name such that $T \Vdash_{\mathbf{PT}} "Z \in [\omega_1]^{\aleph_1}"$. It suffices to construct a sequence of conditions $T_n \in \mathbf{PT}$ and ordinals ζ_n such that:

- $T_0 = T$,
- $T_{n+1} \leq_n T_n$ for each n
- $T_n \langle u_j \rangle \Vdash_{\mathbf{PT}} "\zeta_j \in Z"$ for each $j \in n$
- the mapping $j \mapsto \zeta_j$ is one-to-one.

To carry out the construction let T_n be given and let $\eta = \max_{j \in n} \zeta_{u_j}$. Find $T^* \subseteq T_n \langle u_n \rangle$ and $\zeta_n > \eta$ such that $T^* \Vdash_{\mathbf{PT}} "\zeta_n \in Z"$. Let $T_{n+1} = (T_n \setminus T_n \langle u_n \rangle) \cup T^*$.

The applications of the P-ideal dichotomy in [9] and [10] all rely on simply finding an uncountable set all of whose countable subsets belong to a given ideal. The arguments to be presented here rely on a stronger version of the axiom, but one which is, nevertheless, true in the model constructed in [9]. The following theorem is implicit in Lemma 3.1 of [9]; the following argument simply verifies this assertion.

Theorem 2.13 (Abraham & Todorcevic). *If \mathcal{I} is a P-ideal on ω_1 then there is a partial order $\mathbb{P}_{\mathcal{I}}$, that adds no reals, even when iterated with countable support, such that $\mathbb{P}_{\mathcal{I}}$ adds a set $Z \subseteq \omega_1$ such that for any $W \subseteq \omega_1$ in V which is not the union of countably many sets orthogonal to \mathcal{I}*

$$1 \Vdash_{\mathbb{P}_{\mathcal{I}}} "Z \cap W \neq \emptyset" \quad (2.5)$$

$$1 \Vdash_{\mathbb{P}_{\mathcal{I}}} "(\forall \eta \in \omega_1) Z \cap \eta \in \mathcal{I}." \quad (2.6)$$

PROOF. The proof is the same as that in [9]. To begin, given a non-principal P-ideal \mathcal{I} on ω_1 , fix $A_\xi \subseteq \xi$ for each $\xi \in \omega_1$ such that if $\xi \in \eta$ then $A_\xi \subseteq^* A_\eta$ and such that every member of \mathcal{I} is almost included in some A_ξ . The partial order $\mathbb{P}_{\mathcal{I}}$ is defined to consist of pairs $p = (x_p, \mathfrak{X}_p)$ such that:

- $x_p \in \mathcal{I}$ (The reader should not be confused by the claim in [9] that x_p can be any countable subset of ω_1 .)
- $|\mathfrak{X}_p| \leq \aleph_0$
- $\mathfrak{X}_p \subseteq [\omega_1]^{\aleph_1}$.

The ordering on $\mathbb{P}_{\mathcal{I}}$ is defined by defining $p \leq q$ if:

- $x_p \cap \sup(x_q) = x_q$
- $\mathfrak{X}_p \supseteq \mathfrak{X}_q$
- $\{\xi \in X \mid x_p \setminus x_q \subseteq A_\xi\} \in \mathfrak{X}_p$ for every $X \in \mathfrak{X}_q$.

Lemma 3.1 of [9] establishes that if ω_1 cannot be decomposed into countably many sets orthogonal to \mathcal{I} , then for each $\gamma \in \omega_1$ the set of $p \in \mathbb{P}_{\mathcal{I}}$ such that $x_p \setminus \gamma \neq \emptyset$ is a dense subset of $\mathbb{P}_{\mathcal{I}}$. It will now be verified that the same argument establishes that if $W \subseteq \omega_1$ is not the union of countably many sets orthogonal to \mathcal{I} then $D(W) = \{p \in \mathbb{P}_{\mathcal{I}} \mid x_p \cap W \neq \emptyset\}$ is a dense subset of $\mathbb{P}_{\mathcal{I}}$.

To see this suppose that there is no member of $D(W)$ extending p . Then for each $\mu \in W$ there is some $X \in \mathfrak{X}_p$ such that

$$X(\mu) = \{\xi \in X \mid \mu \in A_\xi\}$$

is countable. For $X \in \mathfrak{X}_p$ let

$$B(X) = \{\mu \in W \mid |X(\mu)| = \aleph_0\}$$

and note that the hypothesis on W implies that $\bigcup_{X \in \mathfrak{X}_p} B(X) \supseteq W$. It therefore suffices to show that each $B(X)$ is orthogonal to \mathcal{I} . To see that this is so, suppose that b is an infinite subset of $B(X)$ and that $b \in \mathcal{I}$. Then $\{\xi \in X \mid b \subseteq^* A_\xi\}$ is clearly a co-countable subset of X because of the cofinality of $\{A_\xi\}_{\xi \in \omega_1}$. Moreover

$$\{\xi \in X \mid b \subseteq^* A_\xi\} = \bigcup_{F \in [b]^{<\aleph_0}} \{\xi \in X \mid (b \setminus F) \subseteq A_\xi\}$$

and hence there is some $F \in [b]^{<\aleph_0}$ such that $\{\xi \in X \mid (b \setminus F) \subseteq A_\xi\}$ is uncountable. Then if $\mu \in b \setminus F$ this contradicts that $X(\mu) \supseteq \{\xi \in X \mid (b \setminus F) \subseteq A_\xi\}$ and $|X(\mu)| = \aleph_0$ because $\mu \in B(X)$.

Theorem 2.14. *Let V be a model of set theory and suppose that $U : \omega_1^2 \rightarrow 2$ is a symmetric, category saturated function in V and that $G \subseteq \mathbf{PT}$ is generic over V . In $V[G]$ let $H \subseteq \mathbb{P}_{\mathcal{S}(\mathbf{PT})}$ be generic over $V[G]$. Then in $V[G][H]$ the function U is universal.*

PROOF. Using Theorem 2.13 in $V[G]$ there is $R \subseteq \omega_1$ such that $[R]^{\aleph_0} \subseteq \mathcal{S}(\mathbf{PT})$ and $R \cap Y \neq \emptyset$ for each uncountable $Y \in V[G]$. Given $W : \omega_1^2 \rightarrow 2$ which is symmetric, construct by induction embeddings $e_\eta : \eta \rightarrow R$ of $W \upharpoonright \eta^2$ into U such that $e_\eta \subseteq e_\zeta$ if $\eta \leq \zeta$.

Since limit stages of the induction are trivial, it suffices to show that given e_η there is $e_{\eta+1}$ as required. Let S be the range of e_η and suppose that $S \subseteq \xi$. Then $S \in [R]^{\aleph_0} \subseteq \mathcal{S}(\mathbf{PT})$. Let $f : S \rightarrow 2$ be defined by $f(\sigma) = W(e_\eta^{-1}(\sigma), \eta)$ and note that $f \in V[G]$ since $V[G]$ and $V[G][H]$ have the same reals. Recall that \mathbf{PT} preserves non-meagre sets by Theorems 6.3.20 and 7.3.46 in [8]. By Lemma 2.11 it therefore follows that, using the notation of Definition 1.3, $Y = \{\gamma \in \omega_1 \mid f \subseteq U^\gamma\}$ is an uncountable set in $V[G]$. By Theorem 2.13 it is possible to find $\gamma \in R \setminus \xi$ such that $f \subseteq U^\gamma$ and, hence, $W(e_\eta^{-1}(\sigma), \eta) = f(\sigma) = U(\sigma, \gamma)$ for all $\sigma \in \mu$. Let $e_{\eta+1} = e_\eta \cup \{(\eta, \gamma)\}$.

Corollary 2.15. *It is consistent with $\mathfrak{b} = \aleph_1$ and $\mathfrak{d} = \aleph_2$ that there is a universal graph on ω_1 .*

PROOF. The required model is the one obtained by starting with a model of the Continuum Hypothesis (in order to use Lemma 1.5) and iterating, with countable support, ω_2 Miller reals at even coordinates and forcing with $\mathbb{P}_{\mathcal{S}(\mathbf{PT})}$ at odd coordinates. Any category saturated graph in the initial model — and, in particular, any saturated graph — will be universal in the final extension. To see this, begin by observing that by Theorem 7.3.46 of [8] it follows that \mathbf{PT} preserves $\sqsubseteq^{\text{Cohen}}$ as defined in Definition 6.3.15 of [8]. Since $\mathbb{P}_{\mathcal{S}(\mathbf{PT})}$ is proper and adds no new reals it is immediate that it also preserves $\sqsubseteq^{\text{Cohen}}$. By Theorem 6.3.20 of [8] it follows that the entire countable support iteration preserves non-meagre sets and, hence, any category saturated graph in the initial model remains category saturated.

To see that all of these graphs are universal use Lemma 3.4 and Lemma 3.6 of [9] to conclude that each partial order in the ω_2 length iteration is proper and has the \aleph_2 -pic of Definition 2.1 on page 409 of [11]. By Lemma 2.4 on page 410 of [11] it follows that the iteration has the \aleph_2 chain condition and, hence, that any graph on ω_1 appears at some stage. It is then routine to apply Theorem 2.14.

It may be worth observing that Theorem 2.14 actually yields that if V is any model of set theory in which there is a category saturated graph and if G is \mathbf{PT} generic over V and H is $\mathbb{P}_{\mathcal{S}(\mathbf{PT})}$ generic over $V[G]$ then already in the model $V[G][H]$ the category saturated graph is universal. In fact, any proper forcing that preserves non-meagreness can be added to the iteration. As a consequence, one can obtain a model with a universal graph in which PFA for non-meagreness preserving partial orders holds; in particular, it is consistent that there are no Suslin trees and there is a universal graph. However the following question seems to be open.

Question 2.16. Does PFA imply there is no universal graph of cardinality \aleph_1 ? What about MA_{\aleph_1} ?

One might also wonder whether one could start with a model in which $2^{\aleph_0} > \aleph_1$ and force with \mathbf{PT} and then $\mathbb{P}_{\mathcal{S}(\mathbf{PT})}$ to obtain a model with a universal graph where $2^{\aleph_0} > \aleph_1$. However, Osvaldo Guzman has pointed out that Peter Simon's argument [12] proving that Sacks forcing collapses the continuum to \mathfrak{b} also works for \mathbf{PT} .

3. Measure saturated graphs are universal after adding Laver reals

Definition 3.1. If $G \subseteq \mathbf{LT}$ is generic over V define $\mathcal{S}(\mathbf{LT})$ to be the set of all $S \subseteq \omega_1$ such that there is $T \in G$ and $\psi : T \rightarrow [\omega_1]^{<\aleph_0}$ such that

1. $\psi \in V$

2. if $s \neq t$ then $\psi(s) \cap \psi(t) = \emptyset$
3. for each $t \in T$ there is $K \in \omega$ such that $|\psi(s)| < K$ for all $s \in \mathbf{succ}_T(t)$.
4. $T \Vdash_{\mathbf{LT}} "S = S_\psi"$

with S_ψ defined as in Definition 2.1. Using the analogue of Lemma 2.2 it can and will be assumed that $\psi(t) = \emptyset$ if $t \not\subseteq \mathbf{stem}(T)$.

Notation 3.2. For a tree $W \subseteq \omega^{<\omega}$ let $\max(W)$ denote the maximal elements of W .

Lemma 3.3. *If S is an \mathbf{LT} -name such that $T \Vdash_{\mathbf{LT}} "S \in \mathcal{S}(\mathbf{LT})"$ then there is $\bar{T} \subseteq T$ and $\psi : \bar{T} \rightarrow [\omega_1]^{<\aleph_0}$ such that*

- $\mathbf{stem}(T) = \mathbf{stem}(\bar{T})$
- ψ satisfies the requirements of Definition 3.1
- $\bar{T} \Vdash_{\mathbf{LT}} "S \equiv^* S_\psi"$.

PROOF. For each $w \in \mathbf{succ}_T(\mathbf{stem}(T))$ let $T_w \subseteq T[w]$ be such that $\mathbf{stem}(T_w) = w$ and there is a well founded tree $W_w \subseteq T_w$ such that $\max(W_w)$ is a front and for each $t \in \max(W_w)$ there is $\psi_t : T_w[t] \rightarrow [\omega_1]^{<\aleph_0}$ that satisfies the requirements of Definition 3.1 and such that

$$T_w[t] \Vdash_{\mathbf{LT}} "S_{\psi_t} = S".$$

The existence of W_w follows from a standard rank argument using the fact that for every $U \in \mathbf{LT}$ such that $U \Vdash_{\mathbf{LT}} "S \in \mathcal{S}(\mathbf{LT})"$ there is $\bar{U} \subseteq U$ such that $\bar{U} \Vdash_{\mathbf{LT}} "S = S_\psi"$ for some $\psi : \bar{U} \rightarrow [\omega_1]^{<\aleph_0}$ in V with disjoint range.

Note that Lemma 2.7 remains true for Laver forcing and apply it to the family

$$\{\psi_t \upharpoonright T_w[t] \mid w \in \mathbf{succ}_T(\mathbf{stem}(T)) \text{ and } t \in \max(W_w)\}$$

to find $\tilde{T}_t \subseteq T_w[t]$ such that $\tilde{T}_t \Vdash_{\mathbf{LT}} "S \equiv^* S_{\tilde{\psi}_t}"$ where $\tilde{\psi}_t$ agrees with ψ_t except that $\tilde{\psi}_t(t) = \emptyset$ and observe that

$$\psi = \bigcup_{w \in W} \left(\bigcup_{t \in \max(W_w)} \tilde{\psi}_t \upharpoonright \tilde{T}_t \right)$$

satisfies the requirements of Definition 3.1. Let $\bar{T} = \bigcup_{w \in W} \bigcup_{t \in \max(W_w)} \tilde{T}_t$. Then ψ and \bar{T} satisfy the lemma.

Lemma 3.4. *If $G \subseteq \mathbf{LT}$ is generic over V and in $V[G]$ and $S \in \mathcal{S}(\mathbf{LT})$ and $f : S \rightarrow 2$ then there is $T \in G$, $\psi : T \rightarrow [\omega_1]^{<\aleph_0}$ with disjoint range and $f^* : \bigcup_{t \in T} \psi(t) \rightarrow 2$, both ψ and f^* in V , such that $T \Vdash_{\mathbf{LT}} "f^* \upharpoonright S = f"$.*

PROOF. The proof is similar to that of Lemma 2.10. Note, however, that by (3) of Definition 3.1 it can be assumed that $\{\psi(s) \mid s \in \mathbf{succ}_T(t)\}$ forms a Δ -system with root R_t for each t . For each $s \in \mathbf{succ}_T(t)$ use the following fact: If W is a well founded tree such that $|\mathbf{succ}_W(w)| = \aleph_0$ for each $w \in W \setminus \max(W)$ whose maximal nodes are coloured in finitely many colours then there is $\bar{W} \subseteq W$ such that

- $|\mathbf{succ}_{\bar{W}}(w)| = \aleph_0$ for each $w \in \bar{W} \setminus \max(\bar{W})$
- $\mathbf{stem}(\bar{W}) = \mathbf{stem}(W)$
- $\mathbf{split}(\bar{W}) = \mathbf{split}(W) \cap (\bar{W} \setminus \max(\bar{W}))$
- all the nodes in $\max(\bar{W})$ are coloured the same colour.

to conclude that there is $T_s \leq_0 T[s]$ and f_s such that $T_s \Vdash_{\mathbf{LT}} "f \upharpoonright \psi(s) = f_s"$. There is then an infinite set of $s \in \mathbf{succ}_T(t)$ such that $f_s \upharpoonright R_t$ all agree. From this it follows that there is $T_t^* \leq_0 T[t]$ and f_t^* such that $T_t^* \Vdash_{\mathbf{LT}} "f \upharpoonright \psi(s) = f_t^*" for all $s \in \mathbf{succ}_{T_t^*}(t)$.$

Lemma 3.5. *If $G \subseteq \mathbf{LT}$ is generic over V then $\mathcal{S}(\mathbf{LT})$ is a P -ideal.*

PROOF. The fact $\mathcal{S}(\mathbf{LT})$ is closed under finite unions and subsets follows from an argument similar to that of Corollary 2.4. The proof of Lemma 2.9 using Lemma 3.3 instead of Lemma 2.8 can now be applied to show that $\mathcal{S}(\mathbf{LT})$ is a P-ideal. The only question the reader may have is as to why Condition (3) of Definition 3.1 is satisfied. This is easily dealt with by restricting the domain of ψ_n so that if $t \in \mathbf{domain}(\psi_n)$ then $|t| \geq n$.

Lemma 3.6. *If G is \mathbf{LT} generic over V and $S \in \mathcal{S}(\mathbf{LT})$, $S \subseteq \xi \in \omega_1$, $f : S \rightarrow 2$ is a function in $V[G]$ and $Z \subseteq 2^\xi$ has full outer measure, then there is $z \in Z$ such that $f \subseteq z$.*

PROOF. Given f find T , ψ and f^* satisfying the conclusion of Lemma 3.4. Let $\bar{f} = f^* \upharpoonright \psi(\mathbf{stem}(T))$, recalling that $\psi(t) = \emptyset$ if $t \subsetneq \mathbf{stem}(T)$, and let \mathcal{O} be the open set $\{h \in 2^\xi \mid h \supseteq \bar{f}\}$. Note that for a given $t \in T$ Condition (3) of the definition of $\mathcal{S}(\mathbf{LT})$ implies that there is a positive lower bound for the measure of $\{h \in 2^\xi \mid f^* \upharpoonright \psi(s) \subseteq h\}$ as s ranges over $\mathbf{succ}_T(t)$. Therefore

$$\{h \in 2^\xi \mid |\{s \in \mathbf{succ}_T(t) \mid f^* \upharpoonright \psi(s) \subseteq h\}| = \aleph_0\}$$

has measure one and hence, since it is independent from \mathcal{O} , it follows that

$$\mathcal{O} \cap \{h \in 2^\xi \mid (\forall t \in \mathbf{split}(T)) |\{s \in \mathbf{succ}_T(t) \mid f^* \upharpoonright \psi(s) \subseteq h\}| = \aleph_0\}$$

has positive measure and so there is some $z \in Z$ such that $\bar{f} \subseteq z$ and

$$(\forall t \in \mathbf{split}(T)) |\{s \in \mathbf{succ}_T(t) \mid f^* \upharpoonright \psi(s) \subseteq z\}| = \aleph_0.$$

It follows that there is $T^* \subseteq T$ such that $T^* \in \mathbf{LT}$ and such that $\mathbf{stem}(T^*) = \mathbf{stem}(T)$ and $f^* \upharpoonright \psi(s) \subseteq z$ for all $s \in \mathbf{split}(T^*)$. Then $T^* \Vdash_{\mathbf{LT}} "f \subseteq z"$. It follows that there is a dense set of conditions forcing the conclusion of the lemma.

Lemma 3.7. *If W is a well founded subtree of $\omega^{<\omega}$ such that $|\mathbf{succ}_W(t)| = \aleph_0$ for every $t \in W \setminus \max(W)$ and if $\theta : \max(W) \rightarrow \omega_1$ then there is a subtree $W^* \subseteq W$ and a one-to-one function $\theta^* : \max(W^*) \rightarrow \omega_1$ such that*

- $|\mathbf{succ}_{W^*}(t)| = \aleph_0$ for every $t \in W^* \setminus \max(W^*)$
- if $w \in \max(W) \cap \max(W^*)$ then $\theta^*(w) = \theta(w)$
- if $w \in \max(W^*) \setminus \max(W)$ then there is a subtree $W_w \subseteq W[w]$ such that $\theta(\bar{w}) = \theta^*(w)$ for each $\bar{w} \in \max(W_w)$ and such that $|\mathbf{succ}_{W_w}(t)| = \aleph_0$ for every $t \in W_w \setminus \max(W_w)$ such that $w \subseteq t$.

PROOF. This is standard argument by induction on the rank of the tree W .

Lemma 3.8. *If G is \mathbf{LT} generic over V then no uncountable subset of ω_1 is orthogonal to $\mathcal{S}(\mathbf{LT})$ in $V[G]$.*

PROOF. This is similar to the proof of Lemma 2.12. Given $T \in \mathbf{LT}$ such that $T \Vdash_{\mathbf{LT}} "Z \in [\omega_1]^{\aleph_1}"$ proceed by induction on n to find $T_{n+1} \leq_n T_n$ (with $T_0 = T$) such that for each $i \leq n$ there is a well founded tree $W_i \subseteq T_{n+1} \langle u_i \rangle$ such that $\max(W_i)$ is a front in $T_{n+1} \langle u_i \rangle$ and a one-to-one function $\psi_i : \max(W_i) \rightarrow \omega_1$ such that $T_{n+1} \langle u_i \rangle \Vdash_{\mathbf{LT}} "\psi_i(w) \in Z"$ for each $w \in \max(W_i)$. To carry out the induction, given T_n , let σ be a name for the least element of Z greater than all elements of

$$\bigcup_{i \in n} \left(\bigcup_{w \in \max(W_i)} \psi_i(w) \right)$$

and then find $T^* \subseteq T_n \langle u_n \rangle$ such that $\mathbf{stem}(T^*) = \Psi_{T_n}(u_n)$ and there is a well founded tree $W^* \subseteq T^*$ such that $\max(W^*)$ is a front in T^* and $\theta : \max(W^*) \rightarrow \omega_1$ such that $T^* \langle u_n \rangle \Vdash_{\mathbf{LT}} "\sigma = \theta(w)"$ for each $w \in \max(W^*)$. Then apply Lemma 3.7 to yield $W_{n+1} \subseteq W^*$ and θ^* and let $T_{n+1} = \bigcup_{w \in \max(W_{n+1})} T^* \langle u_n \rangle [w]$ and then let $\psi_{n+1} = \theta^*$. Then let $\bar{T} = \bigcap_{n \in \omega} T_n$ and define ψ by $\psi(t) = \{\psi_j(t) \mid t \in \max(W_j)\}$ and note that

$$|\psi(t)| \leq |\{j \in \omega \mid t \in \max(W_j)\}| \leq |t|$$

as required by Condition (3) of the definition of $\mathcal{S}(\mathbf{LT})$. It follows that $\bar{T} \Vdash_{\mathbf{LT}} "S_\psi \in [Z]^{\aleph_0}"$.

Theorem 3.9. *Let V be a model of set theory and suppose that $U : \omega_1^2 \rightarrow 2$ is a symmetric, measure saturated function in V and that $G \subseteq \mathbf{LT}$ is generic over V . In $V[G]$ let $H \subseteq \mathbb{P}_{\mathcal{S}(\mathbf{LT})}$ be generic over $V[G]$. Then in $V[G][H]$ the function U is universal.*

PROOF. This is the same as the proof of Theorem 2.14 using measure in the place of category and $\sqsubseteq^{\mathbf{Random}}$ in the place of $\sqsubseteq^{\mathbf{Cohen}}$.

Corollary 3.10. *It is consistent with $\mathfrak{b} = \mathfrak{d} = \aleph_2$ that there is a universal graph on ω_1 .*

PROOF. This the same as the proof of Corollary 2.15.

4. Measure saturated graphs are universal after adding ω^ω -bounding reals

This section contains the key consistency result needed to establish the main result of this article, namely that the existence of a universal graph on ω_1 does not entail the existence of a universal function from ω_1^2 to ω . The key concepts are already contained in §3 the only difference being that the partial order $\mathbf{PT}_{f,g}$ of Definition 7.3.3 of [8] will be used in the place of \mathbf{LT} . As can be seen in the exposition of $\mathbf{PT}_{f,g}$ in [8], there are a great many analogies between this partial order and \mathbf{LT} and these explain the similarities between §3 and this section. The following definition introduces the key technical notion.

Definition 4.1. The partial order $\mathbf{PT}_{f,g}$ will be used with the functions f and g defined as follows. First let $a_n > 0$ be such that $\sum_{n=0}^{\infty} a_n < 1$. Let $g(0, 0) = 1$. If $g(n, n)$ has been defined let $f(n) = \max(g(n, n), 2^n)$. Then let $g(n+1, 0) = 1$ and then choose $g(n+1, k+1)$ be so large that if

- $[X_{i,j}]_{i \in g(n+1, k+1), j \in n+1}$ is a matrix of independent 2-valued random variables
- the probability that $X_{i,j} = 1$ is $1/2$
- $\varphi : g(n+1, k+1) \times (n+1) \rightarrow 2$

then the probability that

$$|\{i \in g(n+1, k+1) \mid (\forall j \in n+1) X_{i,j} = \varphi(j)\}| \geq g(n+1, k) \quad (4.1)$$

is greater than $1 - a_n / \prod_{m=0}^n f(m)$. It will also be required that the following inequalities hold:

$$g(n, j+1) > g(n, j)^2 \quad (4.2)$$

$$g(n+1, j+1) > g(n+1, j)f(n) \quad (4.3)$$

$$g(n, j+1) > g(n, j) \left(\prod_{m \in n+1} f(m) \right)^2 \quad (4.4)$$

$$g(n, j+1) > g(n, j) + \sum_{j \leq n} j \left(\prod_{m \in j+1} f(m) \right)^2 \quad (4.5)$$

$$g(n, j+1) > g(n, j) \prod_{m \in n+1} 2^m \quad (4.6)$$

all of which are easily obtained. The functions f and g are fixed throughout this section.

Recall that $\mathbf{PT}_{f,g}$ consists of trees $T \subseteq \bigcup_{n \in \omega} \prod_{i \in n} f(i)$ such that there is a function $r : \omega \rightarrow \omega$ satisfying that $\lim_{n \rightarrow \infty} r(n) = \infty$ and such that

$$|\mathbf{succ}_T(t)| > g(|t|, r(|t|))$$

for all $t \in T$. For any $T \in \mathbf{PT}_{f,g}$ fix $r_T : \omega \rightarrow \omega$ witnessing that $T \in \mathbf{PT}_{f,g}$. The ordering on $\mathbf{PT}_{f,g}$ is inclusion.

Definition 4.2. If $G \subseteq \mathbf{PT}_{f,g}$ is generic over V define $\mathcal{S}(\mathbf{PT}_{f,g})$ to be the set of all $S \subseteq \omega_1$ such that there is $T \in G$ and $\psi : T \rightarrow [\omega_1]^{< \aleph_0}$ such that

1. $\psi \in V$
2. if $s \neq t$ then $\psi(s) \cap \psi(t) = \emptyset$
3. $\lim_{t \in T} |\psi(t)|/|t| = 0$

4. $T \Vdash_{\mathbf{PT}_{f,g}} "S = S_\psi"$.

recalling that S_ψ is defined in Definition 2.1. As in Definition 3.1 it will be assumed that $\psi(s) = \emptyset$ if $s \not\subseteq \mathbf{stem}(T)$.

Notation 4.3. In this section it will be convenient, for $k \in \omega$ and any tree T , to use the notation $T(k) = \{t \in T \mid |t| = k\}$ and to note that $|T(k)| \leq \prod_{j \in k} f(j)$ if $T \in \mathbf{PT}_{f,g}$.

The following is the lemma that corresponds to Lemma 2.7 in the context of the partial order $\mathbf{PT}_{f,g}$.

Lemma 4.4. *If $\{T_i\}_{i \in m}$ and $\{\psi_i\}_{i \in m}$ satisfy the following*

- $m \leq \prod_{j \in n+1} f(j)$
- $T_i \in \mathbf{PT}_{f,g}$
- $|\mathbf{stem}(T_i)| = n$ for each i
- $r_{T_i}(\ell) > 2$ for each $i \in m$ and $\ell \geq n$
- each $\psi_i : T_i \rightarrow [\omega_1]^{<\aleph_0}$ satisfies the requirements of Definition 4.2

then there are $T_i^* \subseteq T_i$ such that

$$(\forall t \in T_i^*) |\mathbf{succ}_{T_i^*}(t)| \geq g(|t|, r_{T_i}(|t|) - 2) \quad (4.7)$$

$$(\forall i < j < m)(\forall t \in T_i^*)(\forall s \in T_j^*) \text{ if } |t| > n \text{ and } |s| > n \text{ then } \psi_i(t) \cap \psi_j(s) = \emptyset. \quad (4.8)$$

PROOF. Begin by noting the following simple fact: If $\mathcal{F}_i \in [\omega_1]^{<k}$ is a pairwise disjoint family for each $i \in u$ then there are $\mathcal{G}_i \subseteq \mathcal{F}_i$ such that $\bigcup_{i \in u} \mathcal{G}_i$ is a pairwise disjoint family and $|\mathcal{G}_i| \geq |\mathcal{F}_i|/uk$. Given $k \geq n$, define $\mathcal{F}_{t,i}^k = \{\psi_i(s) \mid s \in \mathbf{succ}_{T_i}(t)\}$ for each $i \in m$ and $t \in T_i(k)$. Using Inequality (4.4), for each $k \geq n$ choose $\mathcal{G}_{t,i}^k \subseteq \mathcal{F}_{t,i}^k$ such that $|\mathcal{G}_{t,i}^k| \geq g(k, r_{T_i}(k) - 1)$ and

$$\mathcal{Z}_k = \bigcup_{i \in m} \left(\bigcup_{t \in T_i(k)} \mathcal{G}_{t,i}^k \right)$$

is a pairwise disjoint family for each $k \geq n$. Using the fact that $|\bigcup \mathcal{Z}_k| < k(\prod_{j \in k} f(j))^2$ and Inequality (4.5) it is possible to find $\mathcal{H}_{t,i} \subseteq \mathcal{G}_{t,i}^{|t|}$ such that

$$\bigcup_{k \in \omega} \bigcup_{i \in m} \left(\bigcup_{t \in T_i(k)} \mathcal{H}_{t,i} \right)$$

is a pairwise disjoint family and such that $|\mathcal{H}_{t,i}| \geq g(|t|, r_{T_i}(|t|) - 2)$ if $t \in T_i$. It is then immediate that there are $T_i^* \subseteq T_i$ such that $\mathbf{stem}(T_i^*) = \mathbf{stem}(T_i)$ and such that $\mathbf{succ}_{T_i^*}(t) = \mathcal{H}_{t,i}$ if $t \supseteq \mathbf{stem}(T_i^*)$ and $t \in T_i^*$. These T_i^* satisfy the lemma.

Corollary 4.5. $\mathcal{S}(\mathbf{PT}_{f,g})$ is closed under finite unions.

PROOF. This follows from an argument similar to that of Lemma 4.4 using the following fact: If

$$\{a_{i,j}\}_{i \in u, j \in n} \subseteq [\omega_1]^{<k}$$

is such that for each $i \in u$ the family $\{a_{i,j}\}_{j \in n}$ is pairwise disjoint then there is $Y \subseteq n$ such that $|Y| \geq n/uk$ and the family $\{\bigcup_{i \in u} a_{i,j}\}_{j \in Y}$ is pairwise disjoint. Of course, it is also necessary to note that the limit in Condition (3) of Definition 4.2 is preserved by finite sums.

Lemma 4.6. *If $T \in \mathbf{PT}_{f,g}$ is such that*

$$T \Vdash_{\mathbf{PT}_{f,g}} "S \in \mathcal{S}(\mathbf{PT}_{f,g})" \quad (4.9)$$

$$M \in \omega \text{ is such that } (\forall \ell \geq |\mathbf{stem}(T)|) r_T(\ell) > M \quad (4.10)$$

then there is $T^* \subseteq T$ and $\psi : T^* \rightarrow [\omega_1]^{<\aleph_0}$ such that

- ψ has disjoint range and satisfies Condition (3) of Definition 4.2
- $\mathbf{stem}(T) = \mathbf{stem}(T^*)$
- $T^* \Vdash_{\mathbf{PT}_{f,g}} "S \equiv^* S_\psi"$
- $|\mathbf{succ}_{T^*}(t)| \geq g(|t|, M)$ for all $t \in T^*$ such that $t \supseteq \mathbf{stem}(T)$.

PROOF. Begin by observing that for all $T \in \mathbf{PT}_{f,g}$ there exists $T' \subseteq T$ such that

$$(\exists \psi : T \rightarrow [\omega_1]^{<\aleph_0}) \psi \text{ satisfies Condition (3) of Definition 4.2 and}$$

$$T' \Vdash_{\mathbf{PT}_{f,g}} "S \equiv^* S_\psi" \text{ and } \psi(s) = \emptyset \text{ if } s \subseteq \mathbf{stem}(T)$$

$$\text{and } (\forall s \in T') \text{ if } s \supseteq \mathbf{stem}(T') \text{ then } |\mathbf{succ}_{T'}(s)| \geq g(|s|, M+2). \quad (4.11)$$

To see this simply find $\bar{T} \subseteq T$ and \tilde{T} and $\bar{\psi} : \tilde{T} \rightarrow [\omega_1]^{<\aleph_0}$ such that $\bar{T} \Vdash_{\mathbf{PT}_{f,g}} "\tilde{T} \in G \text{ and } S \equiv^* S_{\bar{\psi}}"$. Note that $\bar{T} \cap \tilde{T} \in \mathbf{PT}_{f,g}$ because $\bar{T} \Vdash_{\mathbf{PT}_{f,g}} "\tilde{T} \in G"$. Choose ℓ such that $r_{\bar{T} \cap \tilde{T}}(i) > M+2$ for all $i \geq \ell$ and let $t \in \bar{T} \cap \tilde{T}(\ell)$. Let $T' = (\bar{T} \cap \tilde{T})[t]$ and define ψ with domain T' by

$$\psi(s) = \begin{cases} \emptyset & \text{if } s \subseteq t \\ \bar{\psi}(s) & \text{otherwise.} \end{cases}$$

Now define $\rho(t) = 0$ if there is $T' \subseteq T[t]$ such that $T \in \mathbf{PT}_{f,g}$, $\mathbf{stem}(T) = t$ and (4.11) holds. Define

$$\rho(t) = \min(\{\alpha + 1 \mid |\{s \in \mathbf{succ}_T(t) \mid \rho(s) \leq \alpha\}| \geq g(|t|, r(|t|) - 1)\})$$

and note that (4.11) and a standard argument show that $\rho(\mathbf{stem}(T))$ is defined. There is no harm in assuming that $r_T(|\mathbf{stem}(T)|) > 1$.

It follows that there is some $\hat{T} \subseteq T$ and $k \in \omega$ such that

$$(\forall s \in \hat{T}) \text{ if } |s| < k \text{ then } |\mathbf{succ}_{\hat{T}}(s)| \geq g(|s|, r_T(|s|) - 1)$$

and, moreover, for each $t \in \hat{T}(k)$ there is $\psi_t : \hat{T}[t] \rightarrow [\omega_1]^{<\aleph_0}$ with disjoint range such that ψ_t satisfies Condition (3) of Definition 4.2 and

$$\hat{T}[t] \Vdash_{\mathbf{PT}_{f,g}} "S \equiv^* S_{\psi_t}" \quad (4.12)$$

$$(\forall j \leq k) \psi_t(t \upharpoonright j) = \emptyset \quad (4.13)$$

$$(\forall s \in \hat{T}[t]) \text{ if } s \supseteq t \text{ then } |\mathbf{succ}_{\hat{T}}(s)| \geq g(|s|, M+2). \quad (4.14)$$

Now use Lemma 4.4 to find $T^t \subseteq \hat{T}[t]$ such that

$$\bigcup_{t \in T(k)} \psi_t \upharpoonright T^t$$

is a function with disjoint range and $|\mathbf{succ}_{T^t}(s)| \geq g(|s|, M)$ for all $s \in T^t$ such that $s \supseteq t$. Let $T^* = \bigcup_{t \in T(k)} \psi_t \upharpoonright T^t$.

Corollary 4.7. *If $T \in \mathbf{PT}_{f,g}$ and*

$$T \Vdash_{\mathbf{PT}_{f,g}} "(\forall n \in \omega) S_n \in \mathcal{S}(\mathbf{PT}_{f,g})"$$

then there is $T^ \subseteq T$ and $\psi : T^* \rightarrow [\omega_1]^{<\aleph_0}$ with disjoint range satisfying Condition (3) of Definition 4.2 such that*

$$T^* \Vdash_{\mathbf{PT}_{f,g}} "(\forall n \in \omega) S_n \subseteq^* S_\psi"$$

and, hence, $1 \Vdash_{\mathbf{PT}_{f,g}} "\mathcal{S}(\mathbf{PT}_{f,g}) \text{ is a } P\text{-ideal}"$.

PROOF. The fact that $\mathcal{S}(\mathbf{PT}_{f,g})$ is closed under finite unions was established in Corollary 4.5. Use Lemma 4.11 to prove that it is closed under subsets in the same way that Lemma 3.4 was used in Lemma 3.5 to prove the corresponding fact about $\mathcal{S}(\mathbf{LT})$.

Inductively find T_n, K_n and ψ_n such that

- $T_{n+1} \subseteq T_n$
- $T_{n+1}(K_n) = T_n(K_n)$
- $\psi_n : \{t \in T_n \mid |t| > K_n\} \rightarrow [\omega_1]^{<\aleph_0}$ has disjoint range
- $|\psi_n(t)|/|t| < 1/n$ if $t \in T_n$ and $|t| > K_n$
- $\psi_{n+1}(t) \supseteq \psi_n(t)$ for all t in the domain of ψ_{n+1}
- $|\text{succ}_{T_n}(t)| \geq g(|t|, n)$ for $t \in T_n$ if $|t| > K_n$
- $K_{n+1} > K_n \geq n$
- $T_n \Vdash_{\mathbf{PT}_{f,g}} "S_n \equiv^* S_{\psi_n}"$.

and then let $T^* = \bigcap_{n \in \omega} T_n$ and define $\psi(t) = \bigcup_{n \in \omega} \psi_n(t)$. Observe that t is in the domain of only finitely many ψ_n and, moreover, if k is maximal such that $t \in \text{domain}(\psi_k)$ then $|\psi_k(t)|/|t| < 1/k$ and $\psi_k(t) \supseteq \psi_j(t)$ for any j such that $t \in \text{domain}(\psi_j)$. Hence $\lim_{t \in T^*} |\psi(t)|/|t| = 0$.

At stage n choose K_{n+1} so large that $r_{T_n}(j) \geq n+4$ for all $j \geq K_{n+1}$. Then for all $t \in T_n(K_{n+1})$ use Lemma 4.6 to find $T'_t \subseteq T_n[t]$ such that $|\text{succ}_{T'_t}(s)| \geq g(|s|, n+3)$ for all $s \in T'_t$ such that $s \supseteq t$ and $\psi_t : T'_t \rightarrow [\omega_1]^{<\aleph_0}$ with disjoint range such that

$$T'_t \Vdash_{\mathbf{PT}_{f,g}} "S_{n+1} \equiv^* S_{\psi_t}"$$

and such that $\psi_t(s) = \emptyset$ if $|s| \leq K_{n+1}$. Then use Lemma 4.4 to find $T_t^{**} \subseteq T'_t$ satisfying Conditions 4.7 and 4.8. In particular, $r_{T_t^{**}}(j) \geq n+1$ if $j \geq |t|$. Then use the argument of Corollary 4.5 to find $T'_t \subseteq T_t^{**}$ such that $r_{T'_t}(j) \geq n$ if $j \geq |t|$ and such that the function ψ_{n+1} defined by $\psi_{n+1}(s) = \psi_t(s) \cup \psi_n(s)$ if $s \in T'_t$ and $t \in T_n(K_{n+1})$ has disjoint range. Let

$$T_{n+1} = \bigcup_{t \in T_n(K_{n+1})} T'_t$$

and observe that all of the induction hypotheses are satisfied.

The following lemma is the counterpart to Lemma 3.7 in the context of $\mathbf{PT}_{f,g}$.

Lemma 4.8. *If W is a finite subtree of $\bigcup_{n \in \omega} \prod_{i \in n} f(i)$ such that $|\text{succ}_W(t)| \geq g(|t|, r(|t|))$ for every $t \in W \setminus \max(W)$ and if $\theta : \max(W) \rightarrow \omega_1$ then there is a subtree $W^* \subseteq W$ and a one-to-one function $\theta^* : \max(W^*) \rightarrow \omega_1$ such that*

- $|\text{succ}_{W^*}(t)| \geq g(|t|, r(|t|) - 2)$ for every $t \in W^* \setminus \max(W^*)$
- if $w \in \max(W) \cap \max(W^*)$ then $\theta^*(w) = \theta(w)$
- if $w \in \max(W^*) \setminus \max(W)$ then there is a subtree $W_w \subseteq W[w]$ such that $\theta(\bar{w}) = \theta^*(w)$ for each $\bar{w} \in \max(W_w)$ and such that $|\text{succ}_{W_w}(t)| \geq g(|t|, r(|t|) - 2)$ for every $t \in W_w \setminus \max(W_w)$ such that $w \subseteq t$.

PROOF. Proceed by induction on the difference between the height of W and the cardinality of $\text{stem}(W)$, the case when $\text{stem}(W)$ is the maximal element of W being trivial. Given the result for n let W be a tree such that the difference between the height of W and the cardinality of $s_W = \text{stem}(W)$ is $n+1$. For each $s \in \text{succ}_W(s_W)$ the difference between the height of $W[s]$ and the cardinality of $s = \text{stem}(W[s])$ is n and hence there are W_s^* and θ_s^* witnessing that the lemma holds.

Let $A = \{s \in \text{succ}_W(s_W) \mid \max(W_s^*) = \{s\}\}$. The first case to consider is that

$$|A| \geq g(|s_W|, r(s_W) - 1).$$

Suppose first that there is $\alpha \in \omega_1$ such that if $B_\alpha = \{s \in \text{succ}_W(s_W) \mid \theta_s^*(s) = \alpha\}$ then

$$|B_\alpha| \geq g(|s_W|, r(s_W) - 2).$$

Then let W^* be the tree with maximal element s_W and let $\theta^*(s_W) = \alpha$. Let $W_{s_W} = \bigcup_{s \in B_\alpha} W_s$.

On the other hand, if $|B_\alpha| < g(|s_W|, r(s_W) - 1)$ for each $\alpha \in \omega_1$ then use Inequality (4.2) to conclude that there is $C \subseteq A$ such that $|C| \geq g(|s_W|, r(s_W) - 2)$ and the mapping $s \mapsto \theta_s^*(s)$ is one-to-one on C . In this case let W^* be the tree whose set of maximal elements is precisely C . For each $s \in C$ the set W_s is already defined from the induction hypothesis.

Now suppose that $|A| < g(|s_W|, r(s_W) - 1)$. Observe that if $s \in \text{succ}_W(s_W) \setminus A$ and $w \in W_s^*$ is such that $\text{succ}_{W_s^*}(w) \subseteq \max(W_s^*)$ then $|w| > |s_W|$ and hence by Inequality (4.3) it follows that

$$g(|w|, r(|w|)) > g(|w|, r(|w|) - 1)f(|s_W|) \geq g(|w|, r(|w|) - 1)|\text{succ}_W(s_W)|.$$

From this it is easy to find for each $w \in W_s^*$ such that $\text{succ}_{W_s^*}(w) \subseteq \max(W_s^*)$ a set $Z_w \subseteq \text{succ}_{W_s^*}(w)$ such that $|Z_w| > g(|w|, r(|w|) - 1)$ and such that

$$\theta^* = \bigcup_{s \in \text{succ}_W(s_W) \setminus A} \bigcup \{ \theta_s^* \upharpoonright Z_w \mid w \in W_s^* \text{ and } \text{succ}_{W_s^*}(w) \subseteq \max(W_s^*) \}$$

is a one-to-one function. Let W^* be the tree whose set of maximal elements is

$$\bigcup_{s \in \text{succ}_W(s_W) \setminus A} \bigcup \{ Z_w \mid w \in W_s^* \text{ and } \text{succ}_{W_s^*}(w) \subseteq \max(W_s^*) \}.$$

The trees W_t for $t \in Z_w \subseteq \max(W^*)$ are already defined from the induction hypothesis.

Lemma 4.9. *If G is $\mathbf{PT}_{f,g}$ generic over V then every infinite subset of ω_1 is orthogonal to $\mathcal{S}(\mathbf{PT}_{f,g})$ in $V[G]$.*

PROOF. This is similar to the proof of Lemma 3.8 but using Lemma 4.8.

The stronger conclusion of Lemma 4.9 that all infinite sets are orthogonal to $\mathcal{S}(\mathbf{PT}_{f,g})$, as opposed to uncountable sets in the conclusion of Lemma 3.8, will not be used here, but seems worthy of note in any case.

Lemma 4.10. *Let $T \in \mathbf{PT}_{f,g}$, $K = |\text{stem}(T)|$ and $k = \prod_{m \in K} 2^m$. Then for any $J \geq K$ and $\theta : T(J) \rightarrow k$ there is $W \subseteq T$ such that $\max(W) \subseteq T(J)$ and $\theta \upharpoonright \max(W)$ is constant and such that $|\text{succ}_W(t)| \geq g(|t|, r_T(|t|) - 1)$ for every $t \in W \setminus \max(W)$.*

PROOF. Proceed by induction on $J - K$ noting that if $J = K$ the result is trivial. If the result is true for n let T and J be given such that $J - K = n + 1$. Since $k < \prod_{m \in K+1} 2^m$ it follows from the induction hypothesis that there is $W_s \subseteq T[s]$ such that $\max(W_s) \subseteq T[s](J)$ and such that $\theta \upharpoonright \max(W_s)$ is constant with value v_s and such that $|\text{succ}_{W_s}(t)| \geq g(|t|, r(|t|) - 1)$ for each $s \in \text{succ}_T(\text{stem}(T))$ and $t \in W_s \setminus \max(W_s)$. Now use Inequality (4.6) of Definition 4.1 to find $Z \subseteq \text{succ}_T(\text{stem}(T))$ and v such that $v_s = v$ for $s \in Z$ and such that $|Z| \geq g(K, r(K) - 1)$. Then $W = \bigcup_{z \in Z} W_z$ is the required tree.

Lemma 4.11. *If $T \in \mathbf{PT}_{f,g}$ and*

$$T \Vdash_{\mathbf{PT}_{f,g}} \text{“} S \in \mathcal{S}(\mathbf{PT}_{f,g}) \text{ and } f : S \rightarrow 2 \text{”}$$

then there are $T^ \subseteq T$ and $\psi : T^* \rightarrow [\omega_1]^{<\aleph_0}$ such that $T^* \Vdash_{\mathbf{PT}_{f,g}} \text{“} S \equiv^* S_\psi \text{”}$ and there is*

$$f^* : \bigcup_{t \in T^*} \psi(t) \rightarrow 2$$

such that $T^ \Vdash_{\mathbf{PT}_{f,g}} \text{“} f \equiv^* f^* \upharpoonright S \text{”}$.*

PROOF. Begin by using Lemma 4.6 to find $\bar{T} \subseteq T$ and $\psi : \bar{T} \rightarrow [\omega_1]^{<\aleph_0}$ such that $\bar{T} \Vdash_{\mathbf{PT}_{f,g}} \text{“} S \equiv^* S_\psi \text{”}$. A standard rank argument can then be used to find $T^{**} \subseteq \bar{T}$ and an increasing sequence of positive integers $\{L_i\}_{i \in \omega}$ such that $T^{**} \in \mathbf{PT}_{f,g}$ and for each $t \in T^{**}(L_{i+1})$ there is $f_t : \bigcup_{m \in L_i} \psi(t \upharpoonright m) \rightarrow 2$ such that

$$T^{**}[t] \Vdash_{\mathbf{PT}_{f,g}} \text{“} f_t = f \upharpoonright \left(\bigcup_{m \in L_i} \psi(t \upharpoonright m) \right) \text{”}.$$

Then use Lemma 4.10 to find T^* such that $r_{T^*}(n) \geq r_{T^{**}}(n) - 1$ and such that $f_t = f_s$ if $|s| = |t| = L_{i+1}$ and $s \upharpoonright L_i = t \upharpoonright L_i$.

Lemma 4.12. *If G is $\mathbf{PT}_{f,g}$ generic over V and $S \in \mathcal{S}(\mathbf{PT}_{f,g})$, $S \subseteq \xi \in \omega_1$, $f : S \rightarrow 2$ is a function in $V[G]$ and $Z \subseteq \omega^\xi$ has full outer measure, then there is $z \in Z$ such that $f \subseteq z$.*

PROOF. Given f find T , ψ and f^* satisfying the conclusion of Lemma 4.11. Let \bar{f} be a finite partial function such that $\bar{f} \supseteq \psi(\text{stem}(T))$ and

$$(f^* \upharpoonright S \setminus \text{domain}(\bar{f})) \cup \bar{f} = f$$

and let \mathcal{O} be the open set $\{h \in 2^\xi \mid h \supseteq \bar{f}\}$. Then $Z \cap \mathcal{O}$ has positive outer measure. Note that Condition 4.1 of Definition 4.1 implies that for any $t \in T$ the probability that

$$\{h \in 2^\xi \mid |\{s \in \text{succ}_T(t) \mid f^* \upharpoonright \psi(s) \subseteq h\}| \geq g(|t|, r_T(|t|) - 1)\}$$

is greater than $1 - a_{|t|} / \prod_{m=0}^{|t|} f(m)$ and hence, for any n the probability that

$$\{h \in 2^\xi \mid (\forall t \in T(n)) \mid \{s \in \text{succ}_T(t) \mid f^* \upharpoonright \psi(s) \subseteq h\}| \geq g(|t|, r_T(|t|) - 1)\}$$

is greater than $1 - a_{|t|}$. It follows from the choice of the a_n that the probability that

$$\{h \in 2^\xi \mid (\forall t \in T) \mid \{s \in \text{succ}_T(t) \mid f^* \upharpoonright \psi(s) \subseteq h\}| \geq g(|t|, r_T(|t|) - 1)\}$$

is positive and, moreover, this set is independent from \mathcal{O} . It follows that there is some $z \in Z$ such that

$$(\forall t \in T) \mid \{s \in \text{succ}_T(t) \mid f^* \upharpoonright \psi(s) \subseteq z\}| \geq g(|t|, r(|t|) - 1)$$

and $\bar{f} \subseteq z$. It follows that there is $T^* \subseteq T$ such that $T^* \in \mathbf{PT}_{f,g}$ and such that if $t \in T$ then $f^* \upharpoonright \psi(t) \subseteq z$. Then $T^* \Vdash_{\mathbf{PT}_{f,g}} \text{“} f \subseteq z \text{”}$.

Theorem 4.13. *Let V be a model of set theory and suppose that $U : \omega_1^2 \rightarrow 2$ is a symmetric, measure saturated function in V and that $G \subseteq \mathbf{PT}_{f,g}$ is generic over V . In $V[G]$ let $H \subseteq \mathbb{P}_{S(\mathbf{PT}_{f,g})}$ be generic over $V[G]$. Then in $V[G][H]$ the graph U is universal.*

PROOF. This is the same as the proof of Theorem 3.9.

Corollary 4.14. *It is consistent that $\mathfrak{b} = \mathfrak{d} = \aleph_1$ and there is a universal graph on ω_1 .*

PROOF. This the same as the proof of Corollary 2.15 but using that $\mathbf{PT}_{f,g}$ is ω^ω bounding.

Finally, a result will be stated without proof that establishes a weak form of universal function from ω_1^2 to ω in the models of this section and the previous one.

Definition 4.15. A function $F : \omega_1^2 \rightarrow \omega$ will be called *bounded* if there is function $B : \omega_1 \rightarrow \omega$ such that $F(\eta, \zeta) < \min(B(\eta), B(\zeta))$ for all η and ζ .

If ν is some measure on ω giving each singleton positive measure and $U : \omega_1^2 \rightarrow \omega$ is ν -saturated then a slight modification of the methods of §3 and §4 yields a model where U is universal for all bounded function from ω_1^2 to ω .

5. Tree partial orders with additional structure

5.1. The general framework

The results of the preceding sections were originally obtained by a more complicated argument that was eventually replaced by the simpler arguments described in §2, §3 and §4. However, there is at least one result for which the simplified argument does not seem to be sufficient; this is the question of finding a universal function from ω_1^2 to ω rather than a universal function from ω_1^2 to 2. This section will describe an argument showing that it is consistent with $\mathfrak{b} < \mathfrak{d}$ that there is a universal function from ω_1^2 to ω . The argument has wider applicability though, which motivated readers will be able to find on their own. Note, however, that in changing the range of the universal object from 2 to ω the argument in the last paragraph of Lemma 2.10 is no longer valid. The remedy for this is Lemma 5.6.

Definition 5.1. Let $G_0 : \omega_1^2 \rightarrow \omega$ and $G_1 : \omega_1^2 \rightarrow \omega$ be symmetric functions and let $\mathcal{E}(G_0, G_1)$ denote the set of all finite, one-to-one functions e that are isomorphisms between $G_1 \upharpoonright \text{domain}(e)^2$ and $G_0 \upharpoonright \text{range}(e)^2$; in other words, $G_1(\eta, \zeta) = G_0(e(\eta), e(\zeta))$ for all distinct η and ζ in the domain of e .

Definition 5.2. If $G_0 : \omega_1^2 \rightarrow \omega$ and $G_1 : \omega_1^2 \rightarrow \omega$ are symmetric functions and $T \subseteq \omega^{<\omega}$ is a tree then a function $E : T \rightarrow \mathcal{E}(G_0, G_1)$ will be called *good* if:

- (a) if s and t belong to T and $s \subseteq t$ then $E(s) \subseteq E(t)$
- (b) if s and t belong to T then $\mathbf{range}(E(t)) \cap \mathbf{range}(E(s)) = \mathbf{range}(E(s \cap t))$.

The following definitions will be used only in the context of $\mathbb{P} = \mathbf{PT}$ but it seems worth providing the more general context since the definitions are applicable for any partial order consisting of trees ordered by inclusion.

Definition 5.3. Let \mathbb{P} be a tree partial order. If $G_0 : \omega_1^2 \rightarrow \omega$ and $G_1 : \omega_1^2 \rightarrow \omega$ are symmetric functions define \mathbb{P}_{G_0, G_1} to consist of triples (T, E, η) such that

1. $T \in \mathbb{P}$
2. $E : T \rightarrow \mathcal{E}(G_0, G_1)$ is good
3. $\eta \in \omega_1$.

If $p = (T, E, \eta) \in \mathbb{P}_{G_0, G_1}$ the notation (T^p, E^p, η^p) will be used to denote (T, E, η) . Define $p \leq q$ if and only if

4. $T^p \subseteq T^q$
5. $E^p(t) = E^q(t)$ for each $t \in T^p$ such that $t \subsetneq \mathbf{stem}(T^p)$
6. $E^p(t) \supseteq E^q(t)$ for each $t \in T^p$ such that $t \supseteq \mathbf{stem}(T^p)$
7. $(\mathbf{range}(E^p(t)) \setminus \mathbf{range}(E^q(t))) \cap \eta^q = \emptyset$ for all $t \in T^p$
8. $\eta^p \geq \eta^q$.

Definition 5.4. If $G \subseteq \mathbb{P}_{G_0, G_1}$ is generic define $E_G : \omega_1 \rightarrow \omega_1$ by $E_G = \bigcup_{p \in G} E(\mathbf{stem}(T^p))$.

It is immediate that E_G is a partial embedding of G_1 into G_0 . However, some extra requirements will be needed to guarantee that E_G is a total embedding. The following arguments restrict attention to Miller forcing.

5.2. Modifying Miller forcing

The first thing to check is that \mathbf{PT}_{G_0, G_1} is proper, indeed, that it satisfies Axiom A. This will rely on the partial orders \prec_n defined in Definition 7.3.44 of [8]. However, the proof is not immediate as this is the place at which η^p , the third component of $p \in \mathbf{PT}_{G_0, G_1}$, is used.

Lemma 5.5. *If $G_0 : \omega_1^2 \rightarrow \omega$ and $G_1 : \omega_1^2 \rightarrow \omega$ are symmetric functions then \mathbf{PT}_{G_0, G_1} satisfies Axiom A.*

PROOF. For p and q in \mathbf{PT}_{G_0, G_1} define $p \prec_n q$ if

1. $p \leq q$
2. $\mathbf{split}_n(T^p) = \mathbf{split}_n(T^q)$
3. $E^p(t \upharpoonright k) = E^q(t \upharpoonright k)$ for all $t \in \mathbf{split}_n(T^p)$ and $k \leq |t|$.

It needs to be verified that these partial orders witness that \mathbf{PT}_{G_0, G_1} satisfies Axiom A. The only point that needs an argument is that given a dense $D \subseteq \mathbf{PT}_{G_0, G_1}$, $p \in \mathbf{PT}_{G_0, G_1}$ and $n \in \omega$ there is $q \prec_n p$ such that for each $t \in \mathbf{split}_n(T^p) = \mathbf{split}_n(T^q)$ the condition $(T^q[t], E^q \upharpoonright T^q[t], \eta^q) \in D$.

To see that this is so, let $\{t_k\}_{k \in \omega}$ enumerate $\mathbf{split}_n(T^p)$. Construct inductively $p_k \in \mathbf{PT}_{G_0, G_1}$ and $\eta_k \in \omega_1$ such that:

- $\eta_0 = \eta^p$
- $p_k \leq (T^p[t_k], E^p \upharpoonright T^p[t_k], \eta_k)$
- $p_k \in D$
- $\eta_k \supseteq \bigcup_{i \in k} \bigcup_{s \in T^{p_i}} \mathbf{range}(E^{p_i}(s))$

- $\eta_k \geq \eta^{p_i}$ for $i \in k$.

Then let $T^q = \bigcup_{k \in \omega} T^{p_k}$ and $E^q = \bigcup_{k \in \omega} E^{p_k}$ and $\eta^q = \bigcup_{k \in \omega} \eta^{p_k}$ and set $q = (T^q, E^q, \eta^q)$. Note that Condition (3) holds because of (5) in Definition 5.3.

Lemma 5.6. *If $G_0 : \omega_1^2 \rightarrow \omega$ is category saturated and $\xi \in \omega_1$ then*

$$\{p \in \mathbf{PT}_{G_0, G_1} \mid \xi \in \mathbf{domain}(E^p(\mathbf{stem}(T^p)))\}$$

is dense in \mathbf{PT}_{G_0, G_1} .

PROOF. Let $p \in \mathbf{PT}_{G_0, G_1}$. For $s \in \mathbf{split}_m(T^p)$ let

$$\mathcal{F}(s) = \{E^p(t) \setminus E^p(s) \mid t \in \mathbf{split}_{m+1}(T) \text{ and } t \supseteq s\}$$

and note that any two distinct functions in $\mathcal{F}(s)$ have disjoint ranges by (b) of Definition 5.2. For $f \in \mathcal{F}(s)$ define $F_{\xi, f} : \mathbf{range}(f) \rightarrow 2$ by $F_{\xi, f}(f(\zeta)) = G_1(\zeta, \xi)$.

Let $\mu \in \omega_1$ be so large that the range of each $E^p(t)$ is contained in μ . Note that if

$$\mathcal{D}(s) = \{h \in 2^\mu \mid h \supseteq F_{\xi, f} \text{ for infinitely many } f \in \mathcal{F}(s)\}$$

then $\mathcal{D}(s)$ is a dense G_δ subset of 2^μ with the product topology. Since G_0 is category saturated it is possible to find $\xi^* > \eta^p$ such that $G_0^{\xi^*} \upharpoonright \mu \in \mathcal{D}(s)$ for all $s \in \mathbf{split}(T^p)$ where $G_0^{\xi^*}$ is defined in Definition 1.3. Moreover, it can be assumed that $G_0(E^p(\mathbf{stem}(T^p))(\zeta), \xi^*) = G_1(\zeta, \xi)$ for each $\zeta \in \mathbf{domain}(\mathbf{stem}(T^p))$.

Now let T be set of all initial segments of elements of

$$\left\{ t \in T^p \mid t \supseteq \mathbf{stem}(T^p) \text{ and } F_{\xi, E^p(t)} \subseteq G_0^{\xi^*} \right\}$$

and note that $T \in \mathbf{PT}$ and, hence, that $(T, E^p \upharpoonright T, \eta^p) \in \mathbf{PT}_{G_0, G_1}$. Moreover if E is defined by $E(t) = E^p(t) \cup \{(\xi, \xi^*)\}$ for all $t \in T$ such that $t \supseteq \mathbf{stem}(T)$ then E is good and hence $(T, E, \eta^p) \in \mathbf{PT}_{G_0, G_1}$.

The following is the version of Theorem 7.3.46 required for \mathbf{PT}_{G_0, G_1} .

Theorem 5.7. \mathbf{PT}_{G_0, G_1} *preserves $\sqsubseteq^{\mathbf{Cohen}}$ as defined in Definition 6.3.15 on page 295 of [8].*

PROOF. Given $p \in \mathbf{PT}_{G_0, G_1}$ and a countable elementary submodel \mathfrak{N} such that

$$p \Vdash_{\mathbf{PT}_{G_0, G_1}} \text{“}c \text{ is a Cohen real over } \mathfrak{N}\text{”}$$

let $T' \in \mathbf{PT}$ be the tree constructed in the proof of Theorem 7.3.46 in [8]. Observe that this proof does not require extending E^p or η^p . Hence $q = (T', E^p, \eta^p) \in \mathbf{PT}_{G_0, G_1}$ and it is easy to check that

$$q \Vdash_{\mathbf{PT}_{G_0, G_1}} \text{“}c \text{ is a Cohen real over } \mathfrak{N}[G]\text{”}.$$

Corollary 5.8. *The countable support iteration of partial orders of the form \mathbf{PT}_{G_0, G_1} preserves the non-meagreness of non-meagre sets from the ground model.*

PROOF. This follows from Theorem 6.3.20 in [8].

Theorem 5.9. *It is consistent with set theory that $\mathfrak{b} = \aleph_1$, $\mathfrak{d} = \aleph_2 = \mathfrak{c}$ and there is a universal function from ω_1^2 to ω .*

PROOF. Let V be a model of the Continuum Hypothesis and, in V , let $\{G_\xi\}_{\xi \in \omega_2}$ enumerate all \mathbb{P} names for functions from ω_1^2 to ω where \mathbb{P} is a partial order of cardinality \aleph_1 . Also in V , using Lemma 1.5 let G be a category saturated function from ω_1^2 to ω . Let \mathbb{P}_ξ be defined inductively so that if ξ is a limit ordinal then \mathbb{P}_ξ is the countable support limit of $\{\mathbb{P}_\eta\}_{\eta \in \xi}$ and, if G_ξ is a \mathbb{P}_ξ name then $\mathbb{P}_{\xi+1} = \mathbb{P}_\xi * \mathbf{PT}_{G, G_\xi}$ and let $\mathbb{P}_{\xi+1} = \mathbb{P}_\xi * \mathbf{PT}$ otherwise.

If $H \subseteq \mathbb{P}_{\omega_2}$ is generic over V and $\bar{G} : \omega_1^2 \rightarrow \omega$ belongs to $V[H]$ then, since \mathbb{P}_{ω_2} is proper by Lemma 5.5, it follows that $\bar{G} \in V[H \cap \mathbb{P}_\xi]$ for some $\xi \in \omega_2$. Hence there is some $\eta \in \omega_2$ such that G_η is a \mathbb{P}_η name for \bar{G} .

By Lemma 6.3.19 of [8] and Corollary 5.8 it follows that

$$1 \Vdash_{\mathbb{P}_\eta} \text{“}G \text{ is category saturated”}$$

and hence by Lemma 5.6 it follows that the domain of $E_{H \cap \mathbf{PT}_{G, G_\eta}}$ as defined in Definition 5.4 is all of ω_1 and so $E_{H \cap \mathbf{PT}_{G, G_\eta}}$ is an embedding of G_η into G as required.

Note that, unlike the corresponding arguments in §2, §3 and §4, the argument of Theorem 5.9 deals with only one function at each stage of the iteration. While in the earlier sections the saturated graph constructed using the Continuum Hypothesis in the ground model remains universal at all stages of the iteration, in Theorem 5.9 this is only achieved at the end of the iteration.

6. Universal graphs are not the same as universal functions

This section describes the combinatorial reason why the existence of a universal graph does not imply the existence of a universal function with range ω .

Lemma 6.1. *If $\mathfrak{b} = \aleph_1$ and there is a sequences of pairs of natural numbers $\{(m_i, n_i)\}_{i \in \omega}$ such that $m_i < n_i < m_{i+1}$ for each $i \in \omega$ then*

$$\left(\forall \mathcal{F} \subseteq \left[\prod_{i \in \omega} [n_i]^{m_i} \right]^{\aleph_1} \right) \left(\exists g \in \prod_{i \in \omega} n_i \right) (\forall f \in \mathcal{F}) (\exists m \in \omega) (\forall k \geq m) g(k) \notin f(k) \quad (6.1)$$

then there is no Sierpiński universal $c : \omega_1^2 \rightarrow \omega$.

PROOF. Let $B_\eta : \eta \rightarrow \omega$ be a bijection for each $\eta \in \omega_1$. Suppose that $c : \omega_1^2 \rightarrow \omega$ is a Sierpiński universal function. If $\eta \in \xi \in \omega_1$ and $j \in \omega$ let

$$f_{\eta, \xi}(j) = \{c(B_\eta^{-1}(k), \xi) \in n_j \mid k \in m_j\}$$

and use the hypothesis of the lemma to find a function $g_\eta \in \prod_{i \in \omega} n_i$ such that $g_\eta(j) \notin f_{\eta, \xi}(j)$ for every $\xi \in \omega_1 \setminus \eta$ and for all but finitely many $j \in \omega$.

Let \mathcal{U} be a family of increasing functions from ω to ω that is \leq^* unbounded and such that $|\mathcal{U}| = \aleph_1$. Let $\psi : \mathcal{U} \times \omega_1 \rightarrow \omega_1$ be a bijection and define

$$b : \omega \times \omega_1 \rightarrow \omega$$

by $b(j, \psi(u, \eta)) = g_\eta(u(j))$.

Now suppose that $e : \omega_1 \rightarrow \omega_1$ is an embedding of the partial function b into c . Let η be such that $e(j) \in \eta$ for all $j \in \omega$ and let $u \in \mathcal{U}$ be such that there are infinitely many k such that $B_\eta(e(k)) \in m_{u(k)}$. Choose j so large that $g_\eta(u(j)) \notin f_{\eta, e(\psi(u, \eta))}(u(j))$ and such that $B_\eta(e(j)) \in m_{u(j)}$. Then

$$b(j, \psi(u, \eta)) = g_\eta(u(j)) \neq c(B_\eta^{-1}(B_\eta(e(j))), e(\psi(u, \eta))) = c(e(j), e(\psi(u, \eta)))$$

contradicting that e is an embedding.

Corollary 6.2. *The existence of a universal graph on ω_1 does not imply the existence of a Sierpiński universal function from ω_1^2 to ω .*

PROOF. Recall that the model establishing Corollary 4.14 is obtained from the ω_2 length iteration of the partial orders $\mathbf{PT}_{f,g}$ of Theorem 4.13. Using $n_i = f(i)$ and $m_i = g(i, 0)$ it follows that given \mathcal{F} in the intermediate model iterating up to ζ the generic function in $\prod_{i \in \omega} n_i$ added at stage ζ witnesses that Condition (6.1) holds in this model. Now apply Lemma 6.1.

7. Other types of embeddings

There are various way of generalizing the notions of embeddings discussed in the introduction. The following two are singled out because something can be said about them.

Definition 7.1. A function $U : \kappa \times \kappa \rightarrow \lambda$ will be called (ρ_0, ρ_1) -weakly universal if for every $f : \kappa \times \kappa \rightarrow \lambda$ there exists a one-to-one function $h : \kappa \rightarrow \kappa$ and functions $e_0 : \lambda \rightarrow \lambda$ and $e_1 : \lambda \rightarrow \lambda$ such that

$$(\forall \alpha \in \kappa) (\forall \beta \in \kappa) e_0(f(\alpha, \beta)) = e_1(U(h(\alpha), h(\beta))) \quad (7.1)$$

$$\text{if } \rho_i > 1 \text{ then } (\forall \xi \in \lambda) |e_i^{-1}(\xi)| < \rho_i \quad (7.2)$$

$$\text{if } \rho_i = 1 \text{ then } (\forall \xi \in \lambda) e_i(\xi) = \xi \quad (7.3)$$

The triple (h, e_0, e_1) will be called a (ρ_0, ρ_1) -weak embedding and, of course, in the case that $\rho_i = 1$ the e_i will usually be ignored. Note that a $(1, 1)$ weak embedding is just an embedding.

While asking for the function $U : \kappa \times \kappa \rightarrow \lambda$ to be $(\lambda^+, 1)$ -weakly universal is trivial (just let U be constant) this is not so clear for the notion of $(1, \lambda^+)$ -weakly universal.

Proposition 7.2. *A function $U : \kappa \times \kappa \rightarrow \lambda$ is $(1, \lambda^+)$ -weakly universal if and only if for every $f : \kappa \times \kappa \rightarrow \lambda$ there exists a one-to-one function $h : \kappa \rightarrow \kappa$ such that*

$$\text{if } f(\alpha, \beta) \neq f(\alpha^*, \beta^*) \text{ then } U(h(\alpha), h(\beta)) \neq U(h(\alpha^*), h(\beta^*))$$

for all α, β, α^* and β^* in κ .

PROOF. Given the property define $e_1 : \lambda \rightarrow \lambda$ by $e_1(U(h(\alpha), h(\beta))) = f(\alpha, \beta)$. The other implication is even more trivial.

Trivial as it may seem, it is worth noting that Proposition 7.2 implies that it is easy to find a $(1, \kappa^+)$ -weakly universal function $U : \kappa \times \kappa \rightarrow \kappa$ (simply make U one-to-one). It is shown in unpublished notes of Tanmay Inamdar [13] that Martin's Axiom for partial orders with Knaster's Property implies that there is a $(1, \aleph_1)$ -weakly universal function from ω_1^2 to ω . That argument is reproduced in the following. Recall that a partial order has Knaster's Property (Property K) if every uncountable subset contains an uncountable subset, any two of whose elements are compatible.

Proposition 7.3. *Martin's Axiom for partial orders with Property K implies that there is a $(1, \aleph_1)$ -weakly universal function from ω_1^2 to ω*

PROOF. Begin by constructing $\Psi_\alpha : \alpha \rightarrow \omega$ by induction on $\alpha \in \omega_1$ such that for each α and $Z \in [\omega_1 \setminus \alpha]^{<\aleph_0}$ there is $S \in [\alpha]^{<\aleph_0}$ such that for all distinct ξ and η in $\alpha \setminus S$ and ζ and μ in Z

$$\Psi_\zeta(\xi) \neq \Psi_\mu(\eta).$$

If $\beta \in \alpha$ let $U(\alpha, \beta) = \Psi_\alpha(\beta)$.

Given $F : \omega_1^2 \rightarrow \omega$ let $\{\mathfrak{M}_\xi\}_{\xi \in \omega_1}$ be a continuous, increasing sequence of countable elementary submodels of $(H(\omega_2), U, F, \in)$ and let \mathbb{P} consist of all finite functions e such that

- $e : \Gamma \rightarrow \omega_1$ is a strictly increasing function with $\Gamma \in [\omega_1]^{<\aleph_0}$
- if $\{\alpha, \beta\}, \{\alpha', \beta'\} \in [\Gamma]^2$ and $F(\alpha, \beta) \neq F(\alpha', \beta')$ then $U(\alpha, \beta) \neq U(\alpha', \beta')$
- if $\gamma \in \Gamma \cap \mathfrak{M}_\xi$ then $e(\gamma) \in \mathfrak{M}_\xi$.

Now if $\{e_\xi\}_{\xi \in \omega_1} \subseteq \mathbb{P}$ then let $A \subseteq \omega_1$ be uncountable such that that $|e_\alpha| = k$ for each $\alpha \in A$ and for each α and β in A if $\psi : \text{domain}(e_\alpha) \rightarrow \text{domain}(e_\beta)$ is the order preserving bijection

$$U(e_\alpha(\xi), e_\alpha(\eta)) = U(e_\beta(\psi(\xi)), e_\beta(\psi(\eta)))$$

$$F(\xi, \eta) = F(\psi(\xi), \psi(\eta))$$

for all ξ and η in the domain of e_α . Moreover, it can be assumed that $\{\text{range}(e_\alpha)\}_{\alpha \in A}$ form a Δ -system with root E and that if $\alpha \in \beta$ then

$$\max(E) < \min(\text{range}(e_\alpha) \setminus E) < \max(\text{range}(e_\alpha) \setminus E) < \min(\text{range}(e_\beta) \setminus E).$$

Now suppose that $\{\alpha_n\}_{n \leq \omega}$ is a subset of A enumerated in increasing order. Let $\alpha = \min(\text{range}(e_{\alpha_\omega}) \setminus E)$ and let $Z = \text{range}(e_{\alpha_\omega}) \setminus E$. Then apply the property of U to find $S \in [\alpha]^{<\aleph_0}$ such that for all distinct ξ and η in $\alpha \setminus S$ and ζ and μ in Z

$$\Psi_\zeta(\xi) \neq \Psi_\mu(\eta).$$

Then let $n \in \omega$ be so large that $S \cap \text{range}(e_{\alpha_n}) = \emptyset$. It is then easy to check that $e_{\alpha_n} \cap e_{\alpha_\omega} \in \mathbb{P}$. An application of the Dushnik–Miller partition relation $\omega_1 \rightarrow (\omega_1, \omega + 1)$ then yields that \mathbb{P} has Property K. The necessary density can be proved by an argument similar to that of the chain condition.

On the other hand, it will follow from Proposition 7.19 that it is consistent with Martin's Axiom for partial orders with Property K of cardinality \aleph_1 that there is no universal function from ω_1^2 to ω ; Corollary 5.22 of [7] establishes that it is consistent with Martin's Axiom for ccc partial orders of cardinality \aleph_1 that there is not even a universal function ω_1^2 to 2.

However, even the following question seems open.

Question 7.4. Is there a $(1, \aleph_1)$ -weakly universal function from ω_1^2 to ω ?

A key motivating question from the introduction was Problem 7.8 of [7].

Question 7.5. Does the existence of a $(2, 1)$ -weakly universal function from ω_1^2 to ω imply the existence of a universal function from ω_1^2 to ω ?

As will be seen by the next result, this question has an elementary solution not requiring the forcing methods developed in earlier sections if an assumption about the cardinal invariant \mathfrak{d} is made.

Proposition 7.6. *If $\mathfrak{d} = \aleph_1$ then the existence of a $(2, 1)$ -weakly universal function from ω_1^2 to ω implies the existence of a universal function from ω_1^2 to ω .*

PROOF. Let $U : \omega_1^2 \rightarrow \omega$ be a $(2, 1)$ -weakly universal function. It will be shown that there is some $D : \omega \rightarrow \omega$ such that for every $f : \omega_1^2 \rightarrow \omega$ there is an embedding $h : \omega_1 \rightarrow \omega_1$ and a bijection $e : \omega \rightarrow \omega$ such that

1. $f(\alpha, \beta) = e(U(h(\alpha), h(\beta)))$ for all α and β
2. $e(n) < D(n)$ and $e^{-1}(n) < D(n)$ for all n .

If this fails then let \mathfrak{D} be a dominating family in ω^ω of cardinality \aleph_1 and for each $D \in \mathfrak{D}$ let f_D be a function be such that there are no h and e satisfying Conditions 1 and 2. There is no harm in assuming that the domain of f_D is X_D^2 and that the $\{X_D\}_{D \in \mathfrak{D}}$ are pairwise disjoint. Then let $f = \bigcup_{D \in \mathfrak{D}} f_D$.

Using that U is a $(2, 1)$ -weakly universal function let $h : \omega_1 \rightarrow \omega_1$ and $e : \omega \rightarrow \omega$ be such such that Condition 1 is satisfied for f . Since \mathfrak{D} is dominating there must be some $D \in \mathfrak{D}$ such that Condition 2 is satisfied for e and D . Restricting h to X_D^2 yields a contradiction.

Now fix D satisfying Conditions 1 and 2. Then $\{J_i\}_{i \in \omega} \subseteq \omega$ be an increasing sequence such that if $n \in J_i$ then $D(n) \in J_{i+1}$. Then define $U^*(\alpha, \beta) = i$ if $J_{2i-1} \leq U(\alpha, \beta) < J_{2i}$. To see that U^* is a universal function from ω_1^2 to ω let $f^* : \omega_1^2 \rightarrow \omega$. Define $f(\alpha, \beta) = J_{f^*(\alpha, \beta)}$. Using the choice of D , let $h : \omega_1 \rightarrow \omega_1$ and $e : \omega \rightarrow \omega$ be such that Conditions 1 and 2 are satisfied. Then $U^*(h(\alpha), h(\beta)) = f^*(\alpha, \beta)$ if $J_{2f^*(\alpha, \beta)-1} \leq U(\alpha, \beta) < J_{2f^*(\alpha, \beta)}$. From Condition 1 it follows that $e(U(\alpha, \beta)) = f(\alpha, \beta) = J_{f^*(\alpha, \beta)}$ and hence $U(\alpha, \beta) = e^{-1}(J_{f^*(\alpha, \beta)}) \in [J_{2f^*(\alpha, \beta)-1}, J_{2f^*(\alpha, \beta)})$ as required.

Corollary 7.7. *If $\mathfrak{d} = \aleph_1$ then the existence of a $(1, 2)$ -weakly universal function from ω_1^2 to ω implies the existence of a universal function from ω_1^2 to ω .*

PROOF. The existence of a $(1, 2)$ -weakly universal function from ω_1^2 to ω is equivalent to the existence of a $(2, 1)$ -weakly universal function from ω_1^2 to ω .

Proposition 7.8. *If $\mathfrak{d} = \aleph_1$ and there is an (\aleph_0, k) -weakly universal function from ω_1^2 to ω then there is a $(2, k)$ -weakly universal function from ω_1^2 to ω .*

PROOF. Suppose there is an (\aleph_0, k) -weakly universal function $U : \omega_1^2 \rightarrow \omega$. Let $\mathfrak{D} \subseteq \omega^\omega$ be a dominating family of cardinality \mathfrak{d} consisting of strictly increasing functions. For $d \in \mathfrak{D}$ let

$$K_d^n = \underbrace{d \circ d \circ \dots \circ d}_{n \text{ iterations}}(0)$$

and let $F_d : \omega \rightarrow \omega$ be defined by $F_d(i) = K_d^{4j}$ for all i such that $K_d^{4j-2} \leq i < K_d^{4j+2}$. Let k_d be the function defined by $k_d(j) = K_d^{4j}$ and note that k_d is one-to-one.

Then $F_d \circ U : \omega_1^2 \rightarrow \{K_d^{4j}\}_{j \in \omega}$ is not $(2, k)$ -weakly universal function and there is a witness to this, namely a function W_d that is not a $(2, k)$ -weakly embeddable into $F_d \circ U$. There is no harm in assuming that there is a partition $\{P_d\}_{d \in \mathfrak{D}}$ of ω_1 such that the domain of W_d is P_d^2 so that $W_d : P_d^2 \rightarrow \omega$. Let

$$W = \bigcup_{d \in \mathfrak{D}} k_d \circ W_d.$$

Using that U is (\aleph_0, k) -weakly universal let $h : \omega_1 \rightarrow \omega_1$ be one-to-one and let $e_0 : \omega \rightarrow \omega$ be finite-to-one and $e_1 : \omega \rightarrow \omega$ be $< k$ -to-one such that $e_0(W(\alpha, \beta)) = e_1(U(h(\alpha), h(\beta)))$. Now let $d \in \mathfrak{D}$ be such that

$$d(i) \supseteq \bigcup_{j \leq i} \bigcup_{a=0}^1 \bigcup_{b=0}^1 e_a^{-1}(e_b(j)) \tag{7.4}$$

and note that $(h \upharpoonright P_d, e_0, e_1)$ is an (\aleph_0, k) -weak embedding of $k_d \circ W_d$ to U . It will be shown that $(h \upharpoonright P_d, k_d, e_1)$ is actually a $(2, k)$ -weak embedding of W_d into $F_d \circ U$; in other words, that $k_d \circ W_d(\alpha, \beta) = e_1 \circ F_d \circ U(h(\alpha), h(\beta))$ for α and β in P_d .

To see that this is the case, let $(\alpha, \beta) \in P_d^2$. Then $k_d(W_d(\alpha, \beta)) = K_d^{2j}$ for $j = W_d(\alpha, \beta)$ and hence, using that $e_1(K_d^{4j-1}, K_d^{4j+1}) \subseteq (K_d^{4j-2}, K_d^{4j+2})$, it suffices to show that $K_d^{4j-1} \leq U(h(\alpha), h(\beta)) < K_d^{4j+1}$. But

$$e_0(K_d^{4j}) = e_0(k_d(W_d(\alpha, \beta))) = e_0(W(\alpha, \beta)) = e_1(U(h(\alpha), h(\beta)))$$

and so it suffices to show that $K_d^{4j-1} \leq e_1^{-1}e_0(K_d^{4j}) < K_d^{4j+1} = d(K_d^{4j})$. This follows immediately from (7.4).

Corollary 7.9. *If $\mathfrak{d} = \aleph_1$ and there is an $(\aleph_0, 1)$ -weakly universal function from ω_1^2 to ω then there is a universal function from ω_1^2 to ω .*

PROOF. Use Proposition 7.8 and Proposition 7.6.

The following definition provides an other way of generalizing the notions of embeddings and universality.

Definition 7.10. Given a family $\mathcal{A} \subseteq \mathcal{P}(\lambda)$, a function $U : \kappa \times \kappa \rightarrow \mathcal{A}$ is \mathcal{A} -weakly universal if for every $f : \kappa \times \kappa \rightarrow \lambda$ there exists a one-to-one function $h : \kappa \rightarrow \kappa$ such that $f(\alpha, \beta) \in U(h(\alpha), h(\beta))$ for all α and β in κ .

Definition 7.11. Considered for the following assertions about the existence of an \mathcal{A} -weakly universal function:

1. There is an \mathcal{A} -weakly universal function with $\mathcal{A} = [\omega]^{<\aleph_0}$.
2. There is an \mathcal{A} -weakly universal function with $\mathcal{A} = [m]^k$ for positive integers m and k .
3. There is an \mathcal{A} -weakly universal function with $\mathcal{A} = \{A \subseteq \omega \mid |\omega \setminus A| = \aleph_0\}$.
4. There is an \mathcal{A} -weakly universal function with $\mathcal{A} = \{\omega \setminus \{j\} \mid j \in \omega\}$.
5. There is an \mathcal{A} -weakly universal function with $\mathcal{A} = [\omega]^k$ for some positive integer k .

Proposition 7.12. *Assertion (3) of Definition 7.11 is equivalent to Assertion (4).*

PROOF. It is immediate that Assertion (3) implies Assertion (4). To see the converse let $\{P_i\}_{i \in \omega}$ be a partition of ω into infinite sets. Given U witnessing that Assertion (4) holds, define $U^*(\alpha, \beta) = \bigcup_{j \in U(\alpha, \beta)} P_j$. Now if $F : [\omega_1]^2 \rightarrow \omega$ let $F^*(\alpha, \beta)$ satisfy that $F(\alpha, \beta) \in P_{F^*(\alpha, \beta)}$. If $e : \omega_1 \rightarrow \omega_1$ is a weak embedding of F^* into U then it is also an embedding into U^* .

The following monotonicity is immediate.

Proposition 7.13. *If $a \leq b \leq c \leq d$ and Assertion (2) holds for $[d]^a$ then it also holds for $[c]^b$.*

Proposition 7.14. *If $k < m$ and Assertion (2) holds for $[k+1]^k$ then it also holds for $[m]^k$.*

PROOF. Let $J = \binom{m}{k+1}$. It is then possible to choose functions $t_i : J \rightarrow k+1$ for $i \in m$ such that for each $x \in [m]^{k+1}$ there is $j \in J$ such that $\{t_i(j) \mid i \in x\} = k+1$. It then follows that if $\psi : (k+1)^J \rightarrow \mathcal{P}(J)$ is defined by

$$\psi(s) = \{i \in m \mid (\forall j \in J) s(j) \neq t_i(j)\}$$

then $|\psi(s)| \leq k$ for each $s \in (k+1)^J$.

Now, given a function U with domain $[\omega_1]^2$ that is $[k+1]^k$ -weakly universal let the domain of U^* be $[\omega_1^J]^2$ and let $U^*(f, g) = \psi(f * g)$ where the function $f * g$ is defined by $(k+1) \setminus \{f * g(j)\} = U(f(j), g(j))$.

To see that U^* is $[m]^k$ -weakly universal let $F : [\omega_1^J]^2 \rightarrow m$. Let $\{g_\xi\}_{\xi \in \omega_1}$ enumerate ω_1^J and define $F^* : [\omega_1 \times J]^2 \rightarrow k+1$ to be any function such that $F^*((\zeta, j), (\eta, j)) = t_{F(g_\zeta, g_\eta)}(j)$. Let $e : \omega_1 \times \omega_1$ be a $[k+1]^k$ -weak embedding of F^* into U . Then define $\bar{e} : \omega_1^J \rightarrow \omega_1^J$ by $\bar{e}(g_\xi)(j) = e(\xi, j)$.

To see that \bar{e} is a $[m]^k$ -weak embedding of F into U^* let $(\eta, \zeta) \in [\omega_1^J]^2$. It must be shown that $F(g_\eta, g_\zeta) \in U^*(\bar{e}(g_\eta), \bar{e}(g_\zeta))$ or, in other words, that $F(g_\eta, g_\zeta) \in \psi(\bar{e}(g_\eta) * \bar{e}(g_\zeta))$. So it needs to be shown that

$$(\forall j \in J) \bar{e}(g_\eta) * \bar{e}(g_\zeta) \neq t_{F(g_\eta, g_\zeta)}(j)$$

and this is equivalent to

$$(\forall j \in J) t_{F(g_\eta, g_\zeta)}(j) \in U(\bar{e}(g_\eta)(j), \bar{e}(g_\zeta)(j)) = U(e(\eta, j), e(\zeta, j)).$$

But $U(e(\eta, j), e(\zeta, j))$ contains $F^*((\eta, j), (\zeta, j)) = t_{F(g_\eta, g_\zeta)}(j)$.

The following is an immediate consequence of Propositions 7.13 and 7.14.

Corollary 7.15. *If (2) of Definition 7.11 holds then there is a unique k such that*

- *there is a universal $[k + 1]^k$ -weakly universal function,*
- *there are $[m]^n$ -weakly universal functions for all $m \geq n \geq k$*
- *there is no $[m]^n$ -weakly universal function if $n < k$ and $m > n$.*

Definition 7.16. Define

$$\mathfrak{K} = \sup(\{k \in \omega \mid \text{there is no } [k + 1]^k\text{-weakly universal function}\})$$

and define $\mathfrak{K}_\omega = \sup(\{k \in \omega \mid \text{there is no } [\omega]^k\text{-weakly universal function}\})$.

It now follows that of the assertions listed in Definition 7.11 only the following need be considered:

1. $\mathfrak{K} < \omega$.
2. There is a $[\omega]^{<\aleph_0}$ -weakly universal function.
3. There is a $\{\omega \setminus \{j\} \mid j \in \omega\}$ -weakly universal function.
4. $\mathfrak{K}_\omega < \omega$.

Observe that Assertion (1) implies Assertion (2) which, in turn, implies Assertion (3). To see that none of these arrows can be reversed two propositions will be useful.

Proposition 7.17. *If $\mathfrak{b} > \aleph_1$ then there is no $[\omega]^{<\aleph_0}$ -weakly universal function.*

PROOF. Suppose that $U : \omega_1^2 \rightarrow [\omega]^{<\aleph_0}$ is an $[\omega]^{<\aleph_0}$ -weakly universal function. For $\xi \in \eta \in \omega_1$ let $F_{\xi,\eta} : \xi \rightarrow \omega$ be defined by $F_{\xi,\eta}(\zeta) = \max(U(\zeta, \eta))$. Using the hypothesis that $\mathfrak{b} > \aleph_1$ let $F_\xi : \xi \rightarrow \omega$ be such that $F_\xi(\zeta) > F_{\xi,\eta}(\zeta)$ for all $\eta > \xi$ and for all but finitely many $\zeta \in \xi$. Let $F : \omega_1^2 \rightarrow \omega$ be defined by $F(\xi, \zeta) = F_\xi(\zeta)$ for $\zeta < \xi$.

If $e : \omega_1 \rightarrow \omega_1$ were a weak embedding of F into U then note that $F_{e(\omega), e(\omega+1)}(e(n)) \leq \max(U(e(\omega + 1), e(n)))$ for all $n \in \omega$ and hence $F(e(\omega + 1), e(n)) \notin U(e(\omega + 1), e(n))$ for all but finitely many $n \in \omega$.

Proposition 7.18. *If there is a universal graph on ω_1 then Assertion 3 holds.*

PROOF. Given a universal graph $U : \omega_1^2 \rightarrow 2$ let $U^*(\eta, \zeta) = \omega \setminus \{U(\eta, \zeta)\}$. Now if $F : \omega_1^2 \rightarrow \omega$ define $F^* : \omega_1^2 \rightarrow 2$ by

$$F^*(\eta, \zeta) = \begin{cases} 0 & \text{if } F(\eta, \zeta) \neq 0 \\ 1 & \text{if } F(\eta, \zeta) = 0 \end{cases}$$

An embedding of F^* into U witnesses that F embeds weakly into U^* .

By Corollary 3.10 it follows from Proposition 7.17 that Assertion (2) fails in the iterated Laver model yet, by Proposition 7.18, Assertion (3) holds there. On the other hand, Assertion (2) does not imply Assertion (1) as can be seen in the model obtained by adding \aleph_2 random reals by Corollary 7.22. In the iterated $\mathbb{P}_{f,g}$ model $\mathfrak{K} = 1$ but $\mathfrak{K}_\omega = \omega$, so Assertion (4) does imply Assertion (1). However, Assertion (1) easily implies Assertion (4). Even though Assertion (3) is the weakest, even this weak form of embedding can fail as the next proposition shows.

Proposition 7.19. *If $U : [\omega_1]^2 \rightarrow \omega$ and $U^*(\eta, \zeta) = \omega \setminus \{U(\eta, \zeta)\}$ and \mathbb{C} is Cohen forcing and \mathbb{P} has Property K and G is $\mathbb{C} * \mathbb{P}$ generic over V then there is $F : [\omega_1]^2 \rightarrow 2$ in $V[G]$ such that for any one-to-one $h : \omega_1 \rightarrow \omega_1$ there are η and ζ such that $F(\eta, \zeta) \notin U^*(h(\eta), h(\zeta))$.*

PROOF. Let $T : [\omega_1]^2 \rightarrow \omega$ witness that $\aleph_1 \not\rightarrow [\aleph_1]_{\aleph_0}^2$. (See [14] for Todorcevic's construction of such a function.) Let $c : \omega \rightarrow \omega$ be the Cohen real added by G — in other words, $c = \bigcup(G \cap \mathbb{C})$ — and define $F = c \circ T$. If $h : \omega_1 \rightarrow \omega_1$ is one-to-one let $g_\eta * p_\eta$ be a $\mathbb{C} * \mathbb{P}$ such that $g_\eta * p_\eta \Vdash_{\mathbb{C} * \mathbb{P}} "h(\eta) = \eta^*"$. There is then some Cohen condition $g \in \mathbb{C}$ and an uncountable $\Gamma \subseteq \omega_1$ such that $g_\eta = g$ for all $\eta \in \Gamma$ and such that $g * p_\eta$ and $g * p_\zeta$ are compatible if η and ζ are in Γ . Let $k \in \omega \setminus \text{domain}(g)$. There are then η and ζ in Γ such that $T(\eta, \zeta) = k$. Extend g to g^* such that $g^*(k) = U(\eta^*, \zeta^*)$ to yield a contradiction.

Corollary 7.20. *In any model obtained by adding \aleph_2 Cohen reals with finite support Assertion (3) fails.*

Proposition 7.21. *If $2 \leq k < \omega$ and $U : [\omega_1]^2 \rightarrow [k]^{k-1}$ and r is a random real then there is $F : [\omega_1]^2 \rightarrow k$ in $V[r]$ such that for any one-to-one $h : \omega_1 \rightarrow \omega_1$ there are η and ζ such that $F(\eta, \zeta) \notin U(\eta, \zeta)$.*

PROOF. Let $T : [\omega_1]^2 \rightarrow \omega$ witness that $\aleph_1 \not\rightarrow [\aleph_1]_{\aleph_0}^2$. Let $r : \omega \rightarrow k$ be a random real and define $F = r \circ T$. If $h : \omega_1 \rightarrow \omega_1$ is one-to-one let B_η be a Borel set of positive measure forcing that $h(\eta) = \eta^*$.

Let Γ be an uncountable subset of ω_1 and $\mu > 0$ be such that B_η has measure greater than μ for each $\eta \in \Gamma$. Then, for each $\eta \in \Gamma$ let C_η be a clopen set in k^ω such that

$$\lambda(C_\eta \Delta B_\eta) < \mu/2(k+1).$$

Let Γ^* be an uncountable subset of Γ for which there is a clopen set C such that $C_\eta = C$ for every $\eta \in \Gamma^*$. By the definition of random forcing there is then some $j \in \omega$ such that if $S(j, k)$ is the Boolean value of the statement $r(j) = k$ then $\lambda(S(j, i) \cap C) = \lambda(C)/k$ for each $i \in k$. There are then η and ζ in Γ^* such that $T(\eta, \zeta) = j$. But $\lambda(C \setminus B_\eta \cap B_\zeta) < 1/(k+1)$ and hence $S(j, i) \cap B_\eta \cap B_\zeta$ has positive measure for each $i \in k$. In particular, if $j \notin U(\eta^*, \zeta^*)$ then $S(j, i) \cap B_\eta \cap B_\zeta$ forces that $F(\eta, \zeta) \notin U(h(\eta), h(\zeta))$.

Corollary 7.22. *In any model obtained by adding \aleph_2 random reals $\mathfrak{R} = \omega$.*

However, it follows from the next result that Assertion (2) holds in the \aleph_2 random real model.

Proposition 7.23. *If $\mathfrak{d} = \aleph_1$ then Assertion (2) holds.*

PROOF. For each $\eta \in \omega_1$ let \mathcal{D}_η be a dominating family on ω^η . Let $\{S_\eta\}_{\eta \in \omega_1}$ be a partition of ω_1 into uncountable sets. Then let \mathcal{D}_η be enumerated as $\{d_{\eta, \zeta}\}_{\zeta \in S_\eta \setminus \eta}$. Then for $\xi < \eta < \zeta < \omega_1$ and $\zeta \in S_\eta$ define $U(\xi, \zeta) = d_{\eta, \zeta}(\xi)$ (thinking of $d_{\eta, \zeta}(\xi)$ as an initial segment of natural numbers) noting that there is no ambiguity in this definition.

Now given $F : [\omega_1]^2 \rightarrow \omega$ construct the embedding h by induction. If $h \upharpoonright \alpha$ has been defined let η be so large that it contains the image of α under h . Let $f : \eta \rightarrow \omega$ be any function such that $f(h(\beta)) = F(\beta, \alpha)$ for all $\alpha \in \beta$. There is then $\zeta \in S_\eta$ such that $d_{\eta, \zeta}(\xi) > f(\xi)$ for all $\xi \in \eta$. Let $h(\alpha) = \zeta$.

Proposition 7.24. *If $2^{\aleph_0} > \aleph_1$ and \clubsuit holds then $\mathfrak{R} = \omega$.*

PROOF. Given $U : [\omega_1]^2 \rightarrow [k]^{k-1}$ let $\psi : \omega_1 \rightarrow \omega_1 \times \omega_1$ be a bijection. Let $\{s_\eta : \omega \rightarrow \eta\}_{\eta \in \omega_1}$ witness that \clubsuit holds. Now construct $F : [\omega_1]^2 \rightarrow k$ by induction.

Suppose that $F \upharpoonright \xi$ has been constructed and the range of $F \upharpoonright \xi$ is cofinal in η . Suppose also that $\psi(s_\eta(j)) = (\psi_0(j), \psi_1(j))$ and that the mapping $j \mapsto \psi_1(j)$ is one-to-one. For each $\zeta \geq \eta$ let $f_{\eta, \zeta} : \omega \rightarrow k$ be the function defined by $k \setminus \{f_{\eta, \zeta}(j)\} = U(\zeta, \psi_1(j))$. Since $2^{\aleph_0} > \aleph_1$ there is some function $g : \omega \rightarrow k$ such that for each $\zeta \geq \eta$ there are infinitely many j such that $g(j) = f_{\eta, \zeta}(j)$. Define $F(\psi_1(j), \xi) = g(j)$ and $F(\mu, \xi)$ arbitrarily if μ is not one of the $\psi_1(j)$.

Now suppose that $e : \omega_1 \rightarrow \omega_1$ is an embedding of F into U . There is then some $\eta \in \omega$ such that $s_\eta(j) \in \psi^{-1}(e)$ for all but finitely many $j \in \omega$ and, moreover, if $\psi(s_\eta(j)) = (\psi_0(j), \psi_1(j))$ then $\lim_{j \rightarrow \infty} \psi_0(j) = \lim_{j \rightarrow \infty} \psi_1(j) = \eta$. Then $e(\eta) \geq \eta$ and there are infinitely many j such that $f_{\eta, e(\eta)} = U(\psi_1(j), e(\eta))$ contradicting that e is a $[k]^{k-1}$ -weak embedding.

Definition 7.25. Let the hypothesis \clubsuit^- assert that there are $s_\sigma : \omega \rightarrow \sigma \times \sigma$ such that for any one-to-one $e : \omega_1 \rightarrow \omega_1$ there are is a stationary set S such that for each $\sigma \in S$ there are infinitely many j such that $s_\sigma(j) \in e$.

Proposition 7.26. *If $\mathfrak{b} > \aleph_1$ and \clubsuit^- holds then there is no $[\omega]^{< \aleph_0}$ -weakly universal function on $[\omega_1]^2$.*

PROOF. The proof is the same as that for Proposition 7.24.

Theorem 7.27. *For any positive integer $m > 1$ it is consistent that $\mathfrak{R}_\omega = m$.*

Note that Theorem 7.27 for the case $k = 1$ yields a model of set theory in which there is no universal function from $[\omega_1]^2$ to ω yet there is a $[\omega]^2$ -weakly universal function. The following theorem provides a companion result by showing that there is a model of set theory in which there is an $[\omega]^{< \aleph_0}$ -weakly universal function but there is not even a $(2, 2)$ -universal function.

Theorem 7.28. *It is consistent that there is a there is an $[\omega]^{<\aleph_0}$ -weakly universal function but there is no $(2, 2)$ -universal function.*

Only the proof of both Theorem 7.27 will be presented. However, it should be noted that the use of a measurable cardinal is not essential, but it does reduce the technical complexities needed in the arguments based on [1] or [3]. Since the key idea does not benefit from the large cardinal hypothesis, the reader interested in eliminating this hypothesis will be able to combine the proofs to be presented here with the details of [1] or [3] to produce a ZFC argument. The details, along with the proof of Theorem 7.28 are presented in an appendix.

Let κ be a measurable cardinal and let \mathcal{F} be a normal, κ -complete ultrafilter on κ . Let \mathbb{L} be the Levy collapse of κ thought of as the partial order for adding a generic function from $L : [\kappa]^2 \rightarrow \omega$ with finite conditions such that for each $\alpha \in \kappa$ the mapping $\beta \mapsto L(\{\beta, \alpha\})$ is one-to-one on α . For a countable set x let \mathbb{C}_x be Cohen forcing thought of as the partial order of finite functions from ω to x .

Definition 7.29. Let R be graph on κ and $U : [\kappa]^2 \rightarrow [\omega]^{<\aleph_0}$. Suppose that $Z \in \mathcal{F}$ is enumerated in increasing order as $\{z_\xi\}_{\xi \in \kappa}$ and $Y \in \mathcal{F}^*$ is also enumerated as

$$Y = \{y_{\xi,n}\}_{\xi \in \kappa, n \in \omega} \text{ and } (\forall n \in \omega) z_{\xi+1} < y_{\xi,n} < y_{\xi,n+1} < z_{\xi+2}. \quad (7.5)$$

Define $\mathbb{E}(U, R, Z, Y)$ to the set of all pairs (e, b) such that:

1. e is a finite, partial function from κ to J
2. if α and β are in $\mathbf{domain}(e)$ then $R(\alpha, \beta) \in U(e(\alpha), e(\beta))$
3. if $\alpha \in \mathbf{domain}(e)$ then $e(\alpha) = y_{\alpha,n}$ for some $n \in \omega$
4. $b \in [\kappa]^{<\aleph_0}$

and define $(e, b) \leq (e', b')$ if $e \supseteq e'$, $b \supseteq b'$ and the range of $e \setminus e'$ is disjoint from b' .

Definition 7.30. Suppose that \mathbb{Q} is a finite support iteration of $\{\mathbb{Q}_\xi\}_{\xi \in \kappa^+}$ such that $\mathbb{Q}_0 = \mathbb{L} \times \mathbb{C}_m$. If $G : [\kappa]^2 \rightarrow \omega$ is a generic function for \mathbb{L} and $H : \omega \rightarrow m$ is generic for \mathbb{C}_m so that $G \times H$ is generic for $\mathbb{Q}_0 * \mathbb{Q}_1 = \mathbb{L} * \mathbb{C}_m$ then let $U_m = H \circ G$. Suppose further that for every $\xi > 1$

$$\mathbb{Q}_{\xi+1} = \mathbb{Q}_\xi * \mathbb{C}_m \text{ if } \xi \text{ is a limit ordinal} \quad (7.6)$$

and otherwise

$$\mathbb{Q}_{\xi+1} = \mathbb{Q}_\xi * \mathbb{E}(U_m, R_{\xi+1}, Z_{\xi+1}, J_{\xi+1}). \quad (7.7)$$

As usual, a condition q will be called *determined* if for each non-limit ξ there are $e_{q,\xi}$ and $b_{q,\xi}$ such that

$$q \upharpoonright \xi \Vdash_{\mathbb{Q}_\xi} \text{“} q(\xi) = (e_{q,\xi}, b_{q,\xi}) \text{”}$$

and if ξ is a limit then

$$q \upharpoonright \xi \Vdash_{\mathbb{Q}_\xi} \text{“} q(\xi) = f_{q,\xi} \text{”}.$$

To simplify notation somewhat, and since m will be fixed, let $\mathbb{E}(U_m, R_{\xi+1}, Z_{\xi+1}, J_{\xi+1})$ be denoted by $\mathbb{E}_m, (R_{\xi+1}, Z_{\xi+1}, J_{\xi+1})$. In a further attempt to simplify notation $e_{q,\xi}$ will simply be denoted as $e_{q,\xi}$ and the same will hold for $b_{q,\xi}$ and $f_{q,\xi}$. While the determined conditions are dense, the following weaker form of being determined will play a key role: If $q \in \mathbb{Q}_\nu$, α is a limit, $q \upharpoonright (\alpha + 1)$ is determined and $J \notin \mathbf{domain}(f_{q,\xi})$ then q will be said to be *weakly determined by α and J* if for each $j \in m$ there is a determined condition $q[j]$ such that:

- $q \upharpoonright \alpha = q[j] \upharpoonright \alpha$
- $q \upharpoonright \alpha \Vdash_{\mathbb{Q}_\alpha} \text{“} (f \cup \{(J, j)\}) * q \upharpoonright (\nu \setminus (\alpha + 1)) = q[j] \upharpoonright (\nu \setminus \alpha) \text{”}.$

Notation 7.31. For $X \subseteq V_\kappa$ define $\mathbf{Reflect}(X) = \left\{ \zeta \in \kappa \mid (V_\zeta, X \cap V_\zeta, \in) \prec_{\Pi_1^1} (V_\kappa, X, \in) \right\}$.

Definition 7.32. For $\nu \in \kappa^+$ let \leq_ν be an ordering of κ in order type ν . Then $\leq_\nu \subseteq V_\kappa$ and it is possible, using the ordering \leq_ν , to find a copy of $\{\mathbb{Q}_\xi\}_{\xi \in \nu}$ in V_κ . With this copy it makes sense to consider $\mathbf{Reflect}(\mathbb{Q}_\nu)$. Now, for $\eta \in \mathbf{Reflect}(\mathbb{Q}_\nu)$ and $q \in \mathbb{Q}_\nu$ that is determined, define $\mathbf{proj}_\eta(q)$ by:

- $\mathbf{proj}_\eta(q)(\xi) = (e_{q,\xi} \upharpoonright \eta, b_{q,\xi} \cap \eta)$ if ξ is not a limit
- $\mathbf{proj}_\eta(q)(\xi) = f_{q,\xi}$ if ξ is a limit.

provided that $\xi \in \nu \cap V_\eta$ or, more precisely, if there is $\xi^* \in V_\eta$ such that ξ^* is in the range of \leq_ν and $\{\alpha \in \mathbf{domain}(\leq_\nu) \mid \alpha \leq_\nu \xi^*\}$ has order type ξ in the ordering \leq_ν . Then, if $\alpha \in \nu \cap V_\eta$ and q is weakly determined by α and J define $\mathbf{proj}_\eta(q)$ so that it is also weakly determined by α and J and satisfies that $\mathbf{proj}_\eta(q)[j] = \mathbf{proj}_\eta(q[j])$.

Note that it is possible that $\mathbf{proj}_\eta(q) \notin \mathbb{Q}_\nu$ and even that $\mathbf{proj}_\eta(q) \notin \mathbb{Q}_\nu \cap V_\eta$. However, the following lemma often suffices.

Lemma 7.33. *If $\eta \in \mathbf{Reflect}(\mathbb{Q}_\nu)$ and $q \in \mathbb{Q}_\nu$ then $\mathbf{proj}_\eta(q) \in \mathbb{Q}_\nu \cap V_\eta$.*

PROOF. It suffices to show that for all ξ such that $(V_\eta, V_\eta \cap \nu) \models \xi \in \nu$

$$\mathbf{proj}_\eta(q \upharpoonright \xi) \Vdash_{\mathbb{Q}_\xi} “(\forall \alpha \in \mathbf{domain}(e_{q,\xi}))(\forall \beta \in \mathbf{domain}(e_{q,\xi})) U_m(e_{q,\xi}(\alpha), e_{q,\xi}(\beta)) = R_\xi(\alpha, \beta)”$$

but this easily follows by induction and the fact that $V_\eta \cap \mathbb{Q}_\xi$ contains a maximal antichain deciding $R_\xi(\alpha, \beta)$ for all α and β in $\mathbf{domain}(e_{q,\xi}) \cap \eta$.

Lemma 7.34. *Suppose that R_ξ is a \mathbb{Q}_ξ -name for a graph on κ for each $\xi \in \kappa^+$. It is then possible to choose inductively Z_ξ and Y_ξ for $\xi \in \kappa^+$ so that if $\{\mathbb{Q}_\xi\}_{\xi \in \kappa^+}$ is as in Definition 7.30 then there are monotone $\Psi_\xi : [\xi]^{<\aleph_0} \rightarrow \kappa$ for each $\xi \in \kappa^+$ and such that*

1. if $q \in \mathbb{Q}_\nu$
2. $\eta \in Z = Z_\nu \setminus \Psi_\nu(\mathbf{domain}(q))$
3. $\bar{\alpha} \in \nu \cap V_\eta$
4. q is weakly determined by $\bar{\alpha}$ and J

then there is q^* such that

5. $q^* \leq \mathbf{proj}_\eta(q)$
6. q^* is weakly determined by $\bar{\alpha}$ and J
7. $q^* \in V_\eta$

with the property that if

8. $\bar{q} \leq q^*$ is in V_η and is weakly determined by $\bar{\alpha}$ and J
9. $u \in \mathbb{L}$ is such that $\mathbf{domain}(u) \subseteq \bigcup_{\alpha \in Z \setminus \eta} \{\alpha\} \times \alpha$
10. $q(0) \cup u \in \mathbb{L}$

then there is r such that $r \leq \bar{q}$, $r \leq u$, $r \leq q$ and r is weakly determined by $\bar{\alpha}$ and J . (Observe that even if $\bar{\alpha}$ and J are not mentioned, the result still has content and this form of the lemma will also be used.)

PROOF. Proceed by induction on ν and suppose the result holds for all $\xi \in \nu$. Consider first the case that ν is a limit and that $\mathbf{cof}(\nu) < \kappa$. Let $C \subseteq [\nu]^{\mathbf{cof}(\nu)}$ be a cofinal set and define $\Psi_\nu(a) = \bigcup_{\gamma \in C} \Psi_\gamma(a \setminus \{\nu\})$. It is easily checked that this works since the iteration is with finite support.

If $\mathbf{cof}(\nu) = \kappa$. Let $\{\gamma_\xi\}_{\xi \in \kappa}$ be an increasing sequence cofinal in ν . Let $Z_\nu = \left\{ \rho \in \kappa \mid \rho \in \bigcap_{\xi \in \rho} Z_{\gamma_\xi} \right\}$. For each $\xi \in \kappa$ let γ_ξ^* be such that $Z_{\gamma_\xi} \setminus \gamma_\xi^* \subseteq Z_\nu$ and then define $\Psi_\nu(a)$ by choosing ξ such that $a \subseteq \gamma_\xi$ and letting $\bar{\Psi}_\nu(a) = \max(\Psi_{\gamma_\xi}(a), \gamma_\xi^*)$ and, to achieve monotonicity, let $\Psi_\nu(a) = \max_{\bar{a} \subseteq a} (\bar{\Psi}_\nu(\bar{a}))$.

Now let $\nu = \xi + 1$. Moreover, assume that ξ is not a limit since otherwise defining $\Psi_\nu(a \cup \{\xi\}) = \Psi_\xi(a)$ is easily verified to satisfy the lemma. Let $Y_\nu \in \mathcal{F}^*$ be a cofinal subset of $Z_\xi \cap \mathbf{Reflect}(\mathbb{Q}_\nu)$ so that if $Z_\nu = (Z \cap R(\mathbb{Q}_\nu)) \setminus Y_\nu$ then (7.5) holds. Let $\mathbb{Q}_\nu = \mathbb{Q}_\xi * \mathbb{E}_m(R_\nu, Z_\nu, Y_\nu)$

Now suppose that $q \in \mathbb{Q}_\nu$ is weakly determined by $\bar{\alpha}$ and J , $\eta \in Z_\nu \setminus \Psi_\nu(\mathbf{domain}(q))$ and $\bar{\alpha} \in \nu \cap V_\eta$. For any condition \tilde{q} that is weakly determined by $\bar{\alpha}$ and J define

$$\mu(\tilde{q})^* = \max(\{e_{\tilde{q}[j],\xi}(\zeta) \mid j \in m \text{ and } \zeta \in \mathbf{domain}(e_{\tilde{q}[j],\xi})\})$$

with the convention that $\max(\emptyset) = -1$. Proceed by induction on $\mu^* \in \kappa$ to show that the Lemma holds at η for all \tilde{q} that are weakly determined by $\bar{\alpha}$ and J and such that $\mu(\tilde{q})^* \leq \mu^*$. Since the case when $\mu(q)^* = -1$ is easy, it can be assumed that $\mu(q)^* = \mu^* \geq 0$. There are now two cases to consider, either $\mu^* < \eta$ or $\mu^* > \eta$. It is not possible that $\mu^* = \eta$ because $e_{q[j],\xi}(\zeta) \in Y_\nu$ and $\eta \in Z_\nu$ and $Y_\nu \cap Z_\nu = \emptyset$.

In the first case let q^* be the condition weakly determined by $\bar{\alpha}$ and J obtained by setting $q^*[j] = \tilde{q}^*[j]^*(e_{q[j],\xi}, b_{q[j],\xi})$. Now suppose that

$$\mathbf{domain}(u) \subseteq \bigcup_{\alpha \in Z_\nu \setminus \eta} \{\alpha\} \times \alpha \quad (7.8)$$

and that $\bar{q} \leq q^*$ is weakly determined by $\bar{\alpha}$ and J . Then $\eta \in Z_\nu \subseteq Z_\xi$ and $\eta > \Psi_\xi(\mathbf{domain}(q^*))$ and by the induction hypothesis there is $\bar{r} \in \mathbb{Q}_\xi$ that is weakly determined by $\bar{\alpha}$ and J and such that $\bar{r} \leq \bar{q} \upharpoonright \xi$, $\bar{r} \leq u$ and $\bar{r} \leq q \upharpoonright \xi$. Then define $r^* \in \mathbb{Q}_\nu$ such that it is weakly determined by $\bar{\alpha}$ and J and $r^*[j] \upharpoonright \xi = \bar{r}[j] \upharpoonright \xi$ and $r^*[j](\xi) = \bar{q}[j](\xi)$. This satisfies the requirements.

On the other hand, suppose that $\mu^* > \eta$ and that the Lemma holds at η for all \tilde{q} that are weakly determined by $\bar{\alpha}$ and J and such that $\mu(\tilde{q})^* < \mu^*$. Note that

$$\mu^* \in Z_\nu \setminus \Psi_\nu(\mathbf{domain}(q)) \subseteq Z_\xi \setminus \Psi_\xi(\mathbf{domain}(q \upharpoonright \xi)) \quad (7.9)$$

and so the induction hypothesis that holds at ξ can be applied to μ^* and $q \upharpoonright \xi$ to find $q_1^* \leq \mathbf{proj}_{\mu^*}(q \upharpoonright \xi) \in \mathbb{Q}_\xi$ satisfying the conclusion of Lemma 7.34. Let

$$B = \{\zeta \in \mu^* \mid (\mu^*, \zeta) \in \mathbf{domain}(q(0))\}. \quad (7.10)$$

Then define $q_2^* \in \mathbb{Q}_\nu$ to be a condition that is weakly determined by $\bar{\alpha}$ and J and defined by setting

$$q_2^*[j] = q_1^*[j]^*(e_{q[j],\xi} \upharpoonright (e_{q[j],\xi}^{-1}(\mu^*)), (b_{q[j],\xi} \cap \mu^*) \cup B)$$

for each $j \in m$. Note that $\mu^*(q_2^*) < \mu^*$ and so the induction hypothesis on μ^* can be applied to q_2^* and η to find a condition q^* that is weakly determined by $\bar{\alpha}$ and J and such that

$$q^* \leq \mathbf{proj}_\eta(q_2^*) \leq \mathbf{proj}_\eta(q)$$

and the conclusion of Lemma 7.34 is satisfied for $\mathbf{proj}_\eta(q_2^*)$.

To see that q^* satisfies the conclusion of Lemma 7.34 for q suppose that u satisfies (9) and (10) and that $\bar{q} \leq q^*$ is weakly determined by $\bar{\alpha}$ and J and that $\bar{q} \in V_\eta$. It follows that there is $r^* \in \mathbb{Q}_\nu \cap V_{\mu^*}$ that is weakly determined by $\bar{\alpha}$ and J such that $r^* \leq \bar{q}$, $r^* \leq u \upharpoonright V_{\mu^*}$ and $r^* \leq q_2^*$.

Let $A = \{j \in m \mid (\exists \mu_j) e_{r^*[j],\xi}(\mu_j) = \mu^*\}$. Now construct by induction $j \in m$ determined conditions $r_j \in \mathbb{Q}_\xi \cap V_{\mu^*}$ such that:

1. $r_0 \upharpoonright \bar{\alpha} \leq r^* \upharpoonright \bar{\alpha}$
2. if $j > 0$ then $r_j \upharpoonright \bar{\alpha} \leq r_{j-1} \upharpoonright \bar{\alpha}$
3. $f_{r_j, \bar{\alpha}}(J) = j$
4. if $j \in A$ then for each $\zeta \in \mathbf{domain}(e_{r^*[j],\xi})$ there is $v_{j,\zeta}$ such that $r_j \Vdash_{\mathbb{Q}_\xi} "R_\nu(\zeta, \mu_j) = v_{j,\zeta}"$.

There is then a condition \bar{r} that is weakly determined by $\bar{\alpha}$ and J and such that $\bar{r} \upharpoonright \bar{\alpha} = r_{m-1} \upharpoonright \bar{\alpha}$ and $\bar{r}[j] \leq r_j$ for each $j \in m$. Moreover, $\bar{r} \in \mathbb{Q}_\xi \cap V_{\mu^*}$ and $\bar{r} \leq \bar{q} \upharpoonright \xi$.

Now observe that if

$$E = \zeta \in \left(\bigcup_{j \in m} \mathbf{domain}(e_{r^*[j],\xi}) \right) \setminus \left(\bigcup_{j \in m} \mathbf{domain}(e_{q[j],\xi}) \right)$$

then $(\mu^*, \zeta) \notin \mathbf{domain}(q(0))$ and, since $\bar{r} \in V_{\mu^*}$ and $\bar{r} \leq \mathbf{proj}_\eta(q)$, it follows that $(\mu^*, \zeta) \notin \mathbf{domain}(\bar{r}(0))$. Hence it is possible to find $u^* : \{\mu^*\} \times E \rightarrow \mu^*$ and $f^* \supseteq \bar{r}(1)$ such that $(\bar{r}(0) \cup u^*, f^*) \leq (\bar{r}(0), \bar{r}(1))$ and such that $f^*(u^*(\mu^*, \zeta)) = \{v_{j,\zeta} \mid j \in m\}$ for each $\zeta \in E$. Then define $\bar{u} = (u \setminus V_{\mu^*}) \cup u^*$ and observe that

$$\mathbf{domain}(\bar{u}) \subseteq \bigcup_{\alpha \in Z_\xi \setminus (\Psi_\xi(\mathbf{domain}(q)) \cup \mu^*)} \{\alpha\} \times \alpha$$

and, moreover, the choice of B in (7.10) guarantees that $q(0) \cup \bar{u} \in \mathbb{L}$. By the induction hypothesis on ξ it is then possible to find a condition \tilde{r} that is weakly determined by $\bar{\alpha}$ and J and such that $\tilde{r} \leq \bar{r}$, $\tilde{r} \leq q \upharpoonright \xi$ and $\tilde{r} \leq \bar{u}$ and, hence, $\tilde{r} \leq u$.

Finally, let $r \in \mathbb{Q}_\nu$ be a condition that is weakly determined by $\bar{\alpha}$ and J and such that $r[j] = \tilde{r}[j] * (e_{r^*[j],\xi} \cup \{(\mu_j, \mu^*)\}, b_{r^*[j],\xi})$ if $j \in A$ and $r[j] = \tilde{r}[j] * (e_{r^*[j],\xi}, b_{r^*[j],\xi})$ otherwise. Observe that

$$r[j] \Vdash_{\mathbb{Q}_\nu} \text{“} R_\nu(\zeta, \mu_j) \in U_m(e_{r[j],\xi}(\zeta), e_{r[j],\xi}(\mu_j)) \text{”}$$

for all j and $\zeta \in \mathbf{domain}(e_{r[j],\xi})$ and so it follows that $r \leq q$. Moreover, $r \upharpoonright \xi \leq \tilde{r} \leq \bar{r} \leq r^* \upharpoonright \xi \leq \bar{q} \upharpoonright \xi$. Then (4) guarantees that $r \leq r^* \leq \bar{q}$ and, hence, is the desired condition.

Corollary 7.35. *If $\zeta \in \nu$ and W is a \mathbb{Q}_ζ name such that*

$$1 \Vdash_{\mathbb{Q}_\zeta} \text{“} W : [\kappa]^2 \rightarrow [\omega]^{m-1} \text{”}$$

*and $\zeta < \bar{\alpha} < \nu$ so that $\mathbb{Q}_{\bar{\alpha}+1} = \mathbb{Q}_{\bar{\alpha}} * \mathbb{C}_m$ and let $T : [\omega_1]^2 \rightarrow \omega$ be any function such that the image of $[X]^2$ under T is infinite for each uncountable $X \subseteq \omega_1$. (See [14] for such a function.) Then*

$$1 \Vdash_{\mathbb{Q}_\zeta} \text{“} \textit{there is no weak embedding of } c_{\bar{\alpha}} \circ T \textit{ into } W \text{”}.$$

PROOF. Suppose the lemma fails and that

$$1 \Vdash_{\mathbb{Q}_\zeta} \text{“} w \textit{ is a weak embedding of } c_{\bar{\alpha}} \circ T \textit{ into } W \text{”}.$$

For each $\xi \in \kappa$ find $q_\xi \in \mathbb{Q}_\zeta$ and ξ^* such that $q_\xi \Vdash_{\mathbb{Q}_\zeta} \text{“} w(\xi) = \xi^* \text{”}$. Using the fact that \mathcal{F} is countably complete, find $J \in \omega$ such that $Z = \{\xi \in \kappa \mid J \notin \mathbf{domain}(q_\xi(\bar{\alpha}))\} \in \mathcal{F}$ so that q_ξ is weakly determined by $\bar{\alpha}$ and J for each $\xi \in Z$. Using Lemma 7.34 find $q_\xi^* \in V_\alpha$ for each $\xi \in Z$ such that $q_\xi^* \leq \mathbf{proj}_\xi(q_\xi)$ and each q_ξ^* is weakly determined by $\bar{\alpha}$ and J and such that the conclusion of Lemma 7.34 is satisfied. Using the normality of \mathcal{F} find p such that $Z' = \{\xi \in \kappa \mid q_\xi^* = p\} \in \mathcal{F}$. Finally, using the choice of T let α and β be in Z' such that

- $\Psi_\nu(\mathbf{domain}(p)) < \alpha < \beta$
- $q_\alpha \in V_\beta$
- $T(\alpha, \beta) = J$
- α and β are in Z_ν .

Using the conclusion of Lemma 7.34 for q_α with $\bar{q} = p$ (and ignoring u since it does not play a role at this point) it is possible to find $r' \in V_\beta$ such that $r' \leq q_\alpha$ and r' is weakly determined by $\bar{\alpha}$ and J . Then using the conclusion of Lemma 7.34 for q_β with $\bar{q} = r'$ it is possible to find r such that $r \leq q_\alpha$, $r \leq q_\beta$ and r is weakly determined by $\bar{\alpha}$ and J . Then let $r^* \leq r \upharpoonright \bar{\alpha}$ be such that $r^* \Vdash_{\mathbb{Q}_{\bar{\alpha}}} \text{“} W(\alpha^*, \beta^*) = a \text{”}$ for some $a \in [\omega]^{m-1}$. Then let $j \in m \setminus a$ and extend r^* so that $r^*(\bar{\alpha}) = j$. It follows that $r^* \Vdash_{\mathbb{Q}_{\bar{\alpha}}} \text{“} c_{\bar{\alpha}}(T(\alpha, \beta)) = c_{\bar{\alpha}}(J) = j \notin a \text{”}$.

An examination of the proof of Corollary 7.35 will reveal that it contains a proof of the following.

Corollary 7.36. *If R_ξ is a \mathbb{Q}_ξ -name for a graph on κ for each $\xi \in \nu$ and Z_ξ and J_ξ are chosen as in Lemma 7.34 so that $\mathbb{Q}_{\xi+1} = \mathbb{Q}_\xi * \mathbb{E}(R_\xi, Z_\xi, J_\xi)$ then \mathbb{Q}_ν has the κ -cc.*

The proof of the chain condition is often sufficient to establish the required density, but that is not the case here. The following result uses the full conclusion of Lemma 7.34.

Corollary 7.37. *If $\xi \in \kappa$ and (7.7) holds at ν then $\{q \in \mathbb{Q}_{\nu+1} \mid \xi \in \mathbf{domain}(q(\nu))\}$ is dense in $\mathbb{Q}_{\nu+1}$.*

PROOF. Let ξ and q be given and suppose that $q \upharpoonright \nu \Vdash_{\mathbb{Q}_\nu} "q((\nu) = (e, b))"$. Using the notation of Definition 7.29 note that $z_{\xi+1} > \xi > e(\eta) > z_{\eta+1} > \eta$ if $\eta \in \mathbf{domain}(e) \cap \xi$. Furthermore, $z_{\xi+1} \in Z_\nu$. Using Lemma 7.34 (and ignoring $\bar{\alpha}$ and J) let $q^* \leq \mathbf{proj}_{z_\xi}(q)$ satisfy the conclusion of that lemma. Since $z_{\xi+1} \in \mathbf{Reflect}(\mathbb{Q}_\nu)$ it is possible to extend q^* to $\bar{q} \in V_{z_{\xi+1}}$ deciding $R_\nu(\eta, \xi)$ for all $\eta \in \mathbf{domain}(e) \cap \xi$. Let n be such that $\{y_{\xi, n}\} \times y_{\xi, n} \cap \mathbf{domain}(q(0)) = \emptyset$ and such that $y_{\xi, n} \notin b$. Then let $u(y_{\xi, n}, e(\eta)) \in \omega$ be such that $\bar{q} \Vdash_{\mathbb{Q}_\nu} "R_\nu(\eta, \xi) = u(y_{\xi, n}, e(\eta))"$ for $\eta \in \mathbf{domain}(e) \cap \xi$. Then let r be such that $r \leq \bar{q}$, $r \leq q$ and if $(r(0), r(1)) = (\bar{u}, f)$ then $f(\bar{u}(y_{\xi, n}, e(\eta))) = \{u(y_{\xi, n}, e(\eta))\}$ for $\eta \in \mathbf{domain}(e) \cap \xi$. If $r(\nu) = (e^*, b^*)$ and $\xi \in \mathbf{domain}(e^*)$ there is nothing to do. Otherwise let $\bar{e} = e \cup \{(\xi, y_{\xi, n})\}$ and let $r^* \upharpoonright \nu = r$ and $r^*(\nu) = (\bar{e}, b)$.

PROOF (OF THEOREM 7.27). Consider the model obtained by forcing with \mathbb{Q}_{κ^+} as defined in Definition 7.30. Since the first coordinate of \mathbb{Q}_{κ^+} is \mathbb{L} , Corollary 7.36 implies that in that model $\kappa^+ = \aleph_2$. It then follows from Corollary 7.37 that $\aleph_\omega \leq m$ in that model. Finally, Corollary 7.35 together with Corollary 7.36 implies that $\aleph_\omega = m$.

As already mentioned, the proof of Theorem 7.27 does not require a large cardinal and the technical details of the ZFC proof can be found in the appendix. The other advantage of the proof presented here, is that the ideas may have further applications. For example, the original hope for this argument was that it would solve Question 2.16.

8. References

- [1] S. Shelah, On universal graphs without instances of CH, *Ann. Pure Appl. Logic* 26 (1) (1984) 75–87. doi:10.1016/0168-0072(84)90042-3.
URL [http://dx.doi.org.ezproxy.library.yorku.ca/10.1016/0168-0072\(84\)90042-3](http://dx.doi.org.ezproxy.library.yorku.ca/10.1016/0168-0072(84)90042-3)
- [2] S. Shelah, Universal graphs without instances of CH: revisited, *Israel J. Math.* 70 (1) (1990) 69–81. doi:10.1007/BF02807219.
URL <http://dx.doi.org.ezproxy.library.yorku.ca/10.1007/BF02807219>
- [3] A. H. Mekler, Universal structures in power \aleph_1 , *J. Symbolic Logic* 55 (2) (1990) 466–477. doi:10.2307/2274640.
URL <http://dx.doi.org.ezproxy.library.yorku.ca/10.2307/2274640>
- [4] R. D. Mauldin (Ed.), *The Scottish Book*, Birkhäuser, Boston, Mass., 1981, mathematics from the Scottish Café, Including selected papers presented at the Scottish Book Conference held at North Texas State University, Denton, Tex., May 1979.
- [5] W. Sierpiński, Sur une fonction universelle de deux variables réelles., *Bull. Int. Acad. Polon. Sci. A* 1936, 8-12 (1936). (1936).
- [6] R. Rado, Universal graphs and universal functions, *Acta Arith.* 9 (1964) 331–340.
- [7] P. B. Larson, A. W. Miller, J. Steprāns, W. A. R. Weiss, Universal functions, *Fund. Math.* 227 (3) (2014) 197–246. doi:10.4064/fm227-3-1.
URL <http://dx.doi.org/10.4064/fm227-3-1>
- [8] T. Bartoszyński, H. Judah, *Set theory*, A K Peters, Ltd., Wellesley, MA, 1995, on the structure of the real line.
- [9] U. Abraham, S. Todorčević, Partition properties of ω_1 compatible with CH, *Fund. Math.* 152 (2) (1997) 165–181.
- [10] S. Todorčević, A dichotomy for P-ideals of countable sets, *Fund. Math.* 166 (3) (2000) 251–267.
- [11] S. Shelah, *Proper and improper forcing*, 2nd Edition, *Perspectives in Mathematical Logic*, Springer-Verlag, Berlin, 1998. doi:10.1007/978-3-662-12831-2.
URL <http://dx.doi.org.ezproxy.library.yorku.ca/10.1007/978-3-662-12831-2>
- [12] P. Simon, Sacks forcing collapses \mathfrak{c} to \mathfrak{b} , *Comment. Math. Univ. Carolin.* 34 (4) (1993) 707–710.
- [13] T. Inamdar, Universal functions, notes written during HIF Semester Cambridge (October 2015).
- [14] S. Todorčević, Partitioning pairs of countable ordinals, *Acta Math.* 159 (3-4) (1987) 261–294. doi:10.1007/BF02392561.
URL <http://dx.doi.org.ezproxy.library.yorku.ca/10.1007/BF02392561>

9. Appendix

The proofs of Theorem 7.27 and Theorem 7.28 will be based on arguments in [1] and [3]. However, while some familiarity with those articles may be useful, the main argument will be presented in detail here since various modifications will need to be made to both [1] and [3]. Before presenting the arguments, some common definitions will be presented.

Definition 9.1. Given $U : [\omega_1]^2 \rightarrow [\omega]^{<\aleph_0}$ and $R : [\omega_1]^2 \rightarrow \omega$ and $A \subseteq \omega_1$ and $\mathfrak{M} = \{\mathfrak{N}_\xi\}_{\xi \in \omega_1}$ define $\mathbb{E}(U, R, A, \mathfrak{M})$ to consist of pairs (e, X) such that

1. $X \in [\omega_1]^{<\aleph_0}$
2. $\mathbf{domain}(e) \in [\omega_1]^{<\aleph_0}$
3. $\mathbf{range}(e) \subseteq A \setminus X$
4. if $\{\eta, \zeta\} \in [\mathbf{domain}(e)]^2$ then $R(\eta, \zeta) \in U(e(\eta), e(\zeta))$
5. for all $\xi \in \mathbf{domain}(e)$ if $\xi \in \mathfrak{N}_{\eta+1} \setminus \mathfrak{N}_\eta$ then $e(\xi) \in \mathfrak{N}_{\tau+\eta+1} \setminus \mathfrak{N}_{\tau+\eta}$ where $\tau = \omega_1 \cap \mathfrak{N}_\eta$.

The partial order \leq on $\mathbb{E}(U, R, A, \mathfrak{M})$ is coordinatewise inclusion. Note that (5) implies that e is strictly increasing.

Finite support iterations of length κ where $\kappa \in \{\omega_2, \omega_3\}$ will be constructed such that $\{R_\alpha\}_{\alpha \in \kappa}$ will be an enumeration, each element occurring cofinally often, of all the potential forcing names for functions from $[\omega_1]^2$ to ω in partial orders that are hereditarily of cardinality no greater than κ :

- If $\kappa = \omega_2$ there is, as in [1], a family $\{A_\xi\}_{\xi \in \omega_2}$ of uncountable subsets of ω_1 any two of which have finite intersection and there are increasing families of countable elementary submodels $\mathfrak{M}_\alpha = \{\mathfrak{N}_{\alpha, \xi}\}_{\xi \in \omega_1}$ such that

$$\{\alpha\} \cup \{\mathfrak{M}_\beta\}_{\beta \in \alpha} \cup \{(A_\xi, R_\xi)\}_{\xi \in \omega_2} \in \mathfrak{N}_{\alpha, 0} \quad (9.1)$$

for each $\alpha \in \omega_2$ and if α is a successor ordinal and R_α is a \mathbb{P}_α name and $1 \Vdash_{\mathbb{P}_\alpha} "R_\alpha : [\omega_1]^2 \rightarrow \omega"$ then $\mathbb{P}_{\alpha+1} = \mathbb{P}_\alpha * \mathbb{Q}_\alpha$ where \mathbb{Q}_α is a \mathbb{P}_α name such that

$$1 \Vdash_{\mathbb{P}_\alpha} "Q_\alpha = \mathbb{E}(R_0, R_\alpha, A_\alpha, \mathfrak{M}_\alpha)".$$

- If $\kappa = \omega_3$ then $2^{\aleph_1} = \aleph_2$ in the ground model and there are increasing families of countable elementary submodels $\mathfrak{M}_\alpha = \{\mathfrak{N}_{\alpha, \xi}\}_{\xi \in \omega_1}$ such that

$$\{\alpha\} \cup \{\mathfrak{M}_\beta\}_{\beta \in \alpha} \cup \{R_\xi\}_{\xi \in \omega_3} \in \mathfrak{N}_{\alpha, 0} \quad (9.2)$$

for each $\alpha \in \omega_3$ and if α is a successor ordinal and R_α is a \mathbb{P}_α name and $1 \Vdash_{\mathbb{P}_\alpha} "R_\alpha : [\omega_1]^2 \rightarrow \omega"$ then $\mathbb{P}_{\alpha+1} = \mathbb{P}_\alpha * \mathbb{Q}_\alpha$ where \mathbb{Q}_α is a \mathbb{P}_α name such that

$$1 \Vdash_{\mathbb{P}_\alpha} "Q_\alpha = \mathbb{E}(R_0, R_\alpha, \omega_1, \mathfrak{M}_\alpha)".$$

In either case, if α is a limit or R_α is not a \mathbb{P}_α name let $\mathbb{P}_{\alpha+1} = \mathbb{P}_\alpha * \mathbb{C}$ where \mathbb{C} is Cohen forcing, which will thought of as adding a generic function $c_\alpha : \omega \rightarrow k+1$ if $\kappa = \omega_2$ and $c_\alpha : \omega \rightarrow \omega$ if $\kappa = \omega_3$. Let

$$W = \{\xi \in \kappa \mid \xi \text{ is not a limit and } 1 \Vdash_{\mathbb{P}_\alpha} "R_\alpha : [\omega_1]^2 \rightarrow \omega" \}.$$

Note that the definition of \mathbb{P}_α implies that $\mathbb{P}_\alpha \in \mathfrak{N}_{\alpha, 0}$.

Definition 9.2. As usual, define a condition $p \in \mathbb{P}_\kappa$ to be determined if:

$$(\forall \sigma \in \mathbf{domain}(p) \cap W)(\exists (e_{p(\sigma)}, X_{p(\sigma)})) p \restriction \sigma \Vdash_{\mathbb{P}_\sigma} "p(\sigma) = (e_{p(\sigma)}, X_{p(\sigma)})" \quad (9.3)$$

$$(\forall \sigma \in \mathbf{domain}(p) \setminus W)(\exists f_{p(\sigma)}) p \restriction \sigma \Vdash_{\mathbb{P}_\sigma} "p(\sigma) = f_{p(\sigma)}" \quad (9.4)$$

$$(\forall \sigma \in \mathbf{domain}(p))(\forall \tau \in \omega_1)(\forall \{\xi, \xi^*\} \in [\mathbf{domain}(e_{p(\sigma)})]^2 \cap \mathfrak{N}_{\sigma, \tau}) \\ (\exists r = r_{p(\sigma)}(\xi, \xi^*))(\exists q = q_{p(\sigma)}(\tau, \xi, \xi^*) \in \mathfrak{N}_{\sigma, \tau} \cap \mathbb{P}_\sigma) p \restriction \sigma \leq q \text{ and } q \Vdash_{\mathbb{P}_\sigma} "R_\sigma(\xi, \xi^*) = r" \quad (9.5)$$

$$(\forall \sigma \in \mathbf{domain}(p) \setminus W) \mathbf{range}(e_{p(\sigma)}) \subseteq \mathbf{domain}(p(0)) \quad (9.6)$$

The use of checks to denote canonical names for elements of the ground model will be omitted in the sequel.

Claim 9.3. *If p and q are determined conditions that belong to \mathbb{P}_κ and for each $\zeta \in \mathbf{domain}(p) \cap \mathbf{domain}(q)$ either $q(\zeta) \leq p(\zeta)$ or $p(\zeta) \leq q(\zeta)$ and $p \vee q$ is defined by*

$$p \vee q(\zeta) = \begin{cases} \min(q(\zeta), p(\zeta)) & \text{if } \zeta \in \mathbf{domain}(q) \cap \mathbf{domain}(p) \\ q(\zeta) & \text{if } \zeta \in \mathbf{domain}(q) \setminus \mathbf{domain}(p) \\ p(\zeta) & \text{if } \zeta \in \mathbf{domain}(p) \setminus \mathbf{domain}(q) \end{cases}$$

then $p \vee q \in \mathbb{P}_{\omega_2}$ and $p \vee q \leq p$ and $p \vee q \leq q$.

PROOF. Proceed by induction on $\zeta \in \mathbf{domain}(p \vee q)$.

Definition 9.4. If p and q are determined conditions in \mathbb{P}_κ and $\rho \in \omega_1$ define $p \sqsupset_\rho q$ to mean that the following conditions hold:

$$p(0) \supseteq q(0) \upharpoonright [\rho]^2 \tag{9.7}$$

$$(\forall \zeta \in \mathbf{domain}(p)) e_{p(\zeta)} \supseteq e_{q(\zeta)} \upharpoonright \rho \tag{9.8}$$

$$(\forall \zeta \in \mathbf{domain}(p)) X_{p(\zeta)} \supseteq X_{q(\zeta)} \cap \rho \tag{9.9}$$

If $p \in \mathbb{P}_{\omega_2}$ is a determined condition define $\mathbf{height}(p)$ to be the least ordinal $\rho \in \omega_1$ such that:

$$\mathbf{domain}(p(0)) \subseteq [\rho]^2 \tag{9.10}$$

$$(\forall \zeta \in \mathbf{domain}(p)) \mathbf{range}(e_{p(\zeta)}) \subseteq \rho \tag{9.11}$$

$$(\forall \zeta \in \mathbf{domain}(p)) X_{p(\zeta)} \subseteq \rho \tag{9.12}$$

PROOF (PROOF OF THEOREM 7.27). To begin, a key change is that \mathbb{Q}_0 defined in §1.4 of [1] will be defined to consist of functions from $[\Gamma]^2$ to $[\omega]^{k+1}$ where $\Gamma \in [\omega_1]^{<\aleph_0}$. In keeping with the notation of [1], if $G \subseteq \mathbb{Q}_0$ is generic then $\bigcup_{f \in G} f$ will be denoted by R_0 .

With these minor modifications it is left to the reader to verify that the arguments of [1] establish that \mathbb{P}_{ω_2} has the countable chain condition and that in any generic extension by \mathbb{P}_{ω_2} the function R_0 is $[\omega]^{k+1}$ -weakly universal. (However, the fact that \mathbb{P}_{ω_2} has the countable chain condition is also easily established using the arguments to be presented in the following.) In particular, it is easy to check that R_0 has the following stronger version of finite saturation: For every $B \in [\omega_1]^{<\aleph_0}$ and $f : B \rightarrow 2$ and $\eta \in \omega_1$ there is ζ such that $\eta < \zeta < \eta + \omega$ such that $R_0(\beta, \zeta) = f(\beta)$ for each $\beta \in B$. Note that R_0 has the stronger finite saturation property in any extension of the ground model. Given $G \subseteq \mathbb{P}_\alpha$ generic over V and any $(e, X) \in \mathbb{Q}_\alpha$ in $V[G]$ and $\xi \in \omega_1$ let η such that $\xi \in \mathfrak{N}_{\alpha, \eta+1} \setminus \mathfrak{N}_{\alpha, \eta}$. There is then some $\zeta \in \mathfrak{N}_{\alpha, \tau+\zeta+1} \setminus \mathfrak{N}_{\alpha, \tau+\zeta}$ where $\tau = \omega_1 \cap \mathfrak{N}_{\alpha, \eta}$ such that $R_0(e(\mu), \zeta) = R_\alpha(\mu, \xi)$ for each $\mu \in \mathbf{domain}(e)$. Then $e \cup \{(\xi, \zeta)\}$ is strictly increasing and so $(e \cup \{(\xi, \zeta)\}, X) \leq_{\mathbb{Q}_\alpha} (e, X)$. In other words,

$$1 \Vdash_{\mathbb{P}_\alpha} \text{“}\{(e, X) \in \mathbb{Q}_\alpha \mid \xi \in \mathbf{domain}(e)\} \text{ is dense.”}$$

To see that there is no $[k+1]^k$ -weakly universal function $U : [\omega_1]^2 \rightarrow [k+1]^k$ suppose that U is a counterexample in $V[G]$ where $G \subseteq \mathbb{P}_{\omega_2}$ is generic. There is then some $\alpha \in \omega_2$ such that $U \in V[G \cap \mathbb{P}_\alpha]$. Let $\bar{\alpha} > \alpha$ be an ordinal that adds a Cohen function $c_{\bar{\alpha}} : \omega \rightarrow k+1$ and let $T : [\omega_1]^2 \rightarrow \omega$ be any function such that the image of $[X]^2$ under T is infinite for each uncountable $X \subseteq \omega_1$. (See [14] for such a function.) Then suppose that $E : \omega_1 \rightarrow \omega_1$ is an embedding of $c_{\bar{\alpha}} \circ T$ into U such that $c_{\bar{\alpha}}(T(\eta, \zeta)) \in U(E(\eta), E(\zeta))$ for all η and ζ in ω_1 .

Then using standard arguments it is possible to find an uncountable subset $\Gamma \subseteq \omega_1$ and $f \in \mathbb{C}$ such that the following hold:

5. for each ξ there is some $\xi^* \in \omega_1$ such that $p_\xi \Vdash_{\mathbb{P}_{\omega_2}} \text{“}E(\xi) = \xi^*\text{”}$
6. p_ξ is determined for each $\xi \in \Gamma$
7. $\{\mathbf{domain}(p_\xi) \mid \xi \in \Gamma\}$ form a Δ -system with root S
8. $\{\mathbf{domain}(p_\xi(\delta)) \mid \xi \in \Gamma\}$ form a Δ -system with root S_δ for each $\delta \in S$
9. there is $\beta \in \omega_1$ such that if $\xi \in \eta$ are in Γ and $\delta \in \mathbf{domain}(p_\xi)$ and $\zeta \in \mathbf{domain}(p_\eta)$ then $\min(\mathbf{domain}(p_\eta(\zeta)) \setminus \beta > \max(\mathbf{range}(p_\xi(\delta)))$

10. $p_\xi(\bar{\alpha}) = f$ for each $\xi \in \Gamma$
11. if $\sigma \in S \cap W$ then there are e^σ such that $e_{p_\xi(\sigma)} \upharpoonright S_\sigma = e^\sigma$ for all $\xi \in \Gamma$
12. $\{X_{p_\xi(\delta)} \mid \xi \in \Gamma\}$ form a Δ -system with root X_δ for each $\delta \in S \cap W$.

Let μ be so large that $\mu > \beta$ and $\mu > \max(\bigcup_{\delta \in S} S_\delta \cup X_\delta)$ and such that if $\sigma < \delta$ and $\{\sigma, \delta\} \subseteq S$ then $A_\sigma \cap A_\delta \subseteq \mu$. Now let ξ and η be in Γ such that $\mu < \xi < \eta$ and such that $J = T(\xi, \eta) \notin \mathbf{domain}(f)$.

As in [1] let $\{\rho_i\}_{i=1}^m$ enumerate

$$\{e_{p_\eta(\zeta)}(\theta) \mid \zeta \in \mathbf{domain}(p_\eta) \cap W \text{ and } \theta \in \mathbf{domain}(e_{p_\eta(\zeta)}) \setminus \mu\}$$

in increasing order. If $1 \leq i \leq m$ let

$$W_i = \{\zeta \in \mathbf{domain}(p_\eta) \cap W \mid (\exists \theta \in \mathbf{domain}(e_{p_\eta(\zeta)}) \setminus \mu) e_{p_\eta(\zeta)}(\theta) = \rho_i\}.$$

For each $\zeta \in W_i$ let $\nu(\zeta, i)$ be such that there is $\theta(\zeta, i) \in \mathfrak{N}_{\zeta, \nu(\zeta, i)}$ and $\kappa(\zeta, i)$ such that

$$\omega_1 \cap \mathfrak{N}_{\zeta, \nu(\zeta, i)} < \kappa(\zeta, i) < (\omega_1 \cap \mathfrak{N}_{\zeta, \nu(\zeta, i)}) \cdot 2$$

such that

$$e_{p_\eta(\zeta)}(\theta(\zeta, i)) \in \mathfrak{N}_{\zeta, \kappa(\zeta, i)+1} \setminus \mathfrak{N}_{\zeta, \kappa(\zeta, i)} \quad (9.13)$$

and then let $Y_i = \{\omega_1 \cap \mathfrak{N}_{\zeta, \kappa(\zeta, i)} \mid \zeta \in W_i\}$.

Families of conditions $\{q_\ell^j\}_{j \in k+1}$ will be constructed by induction on ℓ for $1 \leq \ell \leq m$ such that the following are satisfied:

13. $q_{\ell+1}^j \leq q_\ell^j \leq p_\xi$
14. $\mathbf{height}(q_\ell^j) \leq \rho_\ell + 1$
15. $q_\ell^j \sqsupset_{\rho_\ell+1} p_\eta$
16. $q_\ell^0 \upharpoonright \alpha = q_\ell^j \upharpoonright \alpha$ for all j
17. $q_0^j(\bar{\alpha})(J) = j$.

If this can be accomplished then it follows from (13) that $q_m^j \leq p_\xi$ and from (15) that $q_m^j \leq p_\eta$ for each j . Then note that from (16) it follows that $q_m^0 \upharpoonright \alpha = q_m^j \upharpoonright \alpha$ for all j .

Then let $\bar{q} \in \mathbb{P}_\alpha$ be such that $\bar{q} \leq q_m^0 \upharpoonright \alpha$ and $\bar{q} \Vdash_{\mathbb{P}_\alpha} "U(\xi^*, \eta^*) = k+1 \setminus \{j\}"$ for some $j \in k+1$ and define q by

$$q(\zeta) = \begin{cases} \bar{q}(\zeta) & \text{if } \zeta \in \alpha \\ q_m^j(\zeta) & \text{if } \zeta \geq \alpha \end{cases}$$

and note that $q \leq q_m^j$ and that (17) implies that

$$q \Vdash_{\mathbb{P}_{\omega_2}} "c_{\bar{\alpha}} \circ T(\xi, \eta) = j \notin U(\xi^*, \eta^*) = U(E(\xi), E(\eta))"$$

contradicting that U is $[k+1]^k$ -weakly universal.

The family $\{q_\ell^j\}_{j \in k+1}$ will also be constructed by induction on j . To start, let

$$q_0^j(\zeta) = \begin{cases} p_\xi(\zeta) & \text{if } \zeta \neq \bar{\alpha} \\ f \cup \{(J, j)\} & \text{if } \zeta = \bar{\alpha} \end{cases}$$

for all j . The following additional induction hypothesis will be used: for every $\ell \leq m$ and $j < k+1$ and $\zeta \in \mathbf{domain}(p_\eta)$

$$X_{q_\ell^j(\zeta)} \supseteq \left(\bigcup_{\{\delta, \bar{\delta}\} \in [\mathbf{domain}(p_\eta)]^2} A_\delta \cap A_{\bar{\delta}} \right) \cap (\rho_\ell + 1) \setminus \mathbf{range}(e_{p_\eta(\zeta)}) \quad (9.14)$$

$$X_{q_\ell^j(\zeta)} \supseteq \bigcup_{u \in k+1 \setminus \{j\}} \mathbf{range}(e_{q_{\ell-1}^u(\zeta)}) \setminus \mathbf{range}(e_{p_\eta(\zeta)}) \quad (9.15)$$

$$X_{q_\ell^j(\zeta)} \supseteq \bigcup_{u \in j} \mathbf{range}(e_{q_\ell^u(\zeta)}) \setminus \mathbf{range}(e_{p_\eta(\zeta)}) \quad (9.16)$$

$$X_{q_\ell^j(\zeta)} \supseteq \{\beta \in \rho_\ell \mid (\exists v \geq \ell) \{\beta, \rho_v\} \in \mathbf{domain}(p_\eta(0))\} \setminus \mathbf{range}(e_{p_\eta(\zeta)}). \quad (9.17)$$

To construct the family $\{q_\ell^j\}_{j \in k+1}$ define \hat{q}_ℓ^j for $\ell \geq 1$ by

$$\hat{q}_\ell^j(\zeta) = \begin{cases} q_{\ell-1}^k(\zeta) & \text{if } j = 0 \text{ and } \zeta < \alpha \\ q_\ell^{j-1}(\zeta) & \text{if } j > 1 \text{ and } \zeta < \alpha \\ (e_{p_\varepsilon}(\zeta), X(\zeta, j, \ell)) & \text{if } \zeta \in S \text{ and } \ell = 1 \\ (e_{q_{\ell-1}^j(\zeta)}, X(\zeta, j, \ell)) & \text{if } \zeta \in \mathbf{domain}(p_\eta) \text{ and } \ell > 1 \\ f \cup \{(J, j)\} & \text{if } \zeta = \bar{\alpha} \text{ and } \ell = 1 \\ q_{\ell-1}^j(\zeta) & \text{otherwise} \end{cases} \quad (9.18)$$

where $X(\zeta, j, \ell)$ is chosen to satisfy conditions (9.14) to (9.17). Observe that $\hat{q}_\ell^j \in \mathbb{P}_{\omega_2}$ and $\hat{q}_\ell^j \leq q_{\ell-1}^j$.

Let $\{\nu_i\}_{i=1}^n$ enumerate Y_ℓ in increasing order. Conditions $q_{\ell,i}^j$ will be constructed by induction on i such that:

18. $q_{\ell,i+1}^j \leq q_{\ell,i}^j \leq \hat{q}_\ell^j$
19. $\mathbf{height}(q_{\ell,i}^j) \leq \nu_i$
20. $q_{\ell,i}^j(0) \supseteq p_\eta(0) \upharpoonright [\nu_i]^2$
21. $q_{\ell,i}^j \sqsupset_{\nu_i} p_\eta$
22. letting $Z_u = \{\zeta \in W_\ell \mid \omega_1 \cap \mathfrak{N}_{\zeta, \kappa(\zeta, \ell)} = \nu_u\}$ there is

$$r_{\zeta,i}^j : \mathbf{domain}(e_{q_{\ell,i}^j(\zeta)}) \cap \nu_i \rightarrow k+1$$

for each $\zeta \in \bigcup_{u=1}^i Z_u$ such that

$$q_{\ell,i}^j \upharpoonright \zeta \Vdash_{\mathbb{P}_\zeta} \text{“} R_\zeta(\delta, \theta(\zeta, i)) = r_{\zeta,i}^j(\delta)\text{”}$$

for all $\delta \in \mathbf{domain}(e_{q_{\ell,i}^j(\zeta)}) \cap \nu_i$

23. if $\zeta \in \bigcup_{u=1}^i Z_u$ and $\delta \in \mathbf{domain}(p_\eta(\zeta))$ then $r_{\zeta,i}^j(\delta) = r_{p_\eta(\zeta)}(\delta, \theta(\zeta, i))$ as in (9.5).

If this can be accomplished for each $j \in k+1$ then define a function H by

$$H(e_{q_{\ell,n}^j(\zeta)}(\delta), \rho_\ell) = \{r_{\zeta,n}^j(\delta) \mid j \in k+1\}$$

for $\zeta \in W_\ell$ and $\delta \in \mathbf{domain}(e_{q_{\ell,n}^j(\zeta)}) \setminus \mathbf{domain}(e_{p_\eta(\zeta)})$. To see that H is a well defined function observe that if

$$e_{q_{\ell,n}^j(\zeta)}(\delta) = e_{q_{\ell,n}^{j'}(\zeta')}(\delta') \quad (9.19)$$

and $\delta \notin \mathbf{domain}(e_{p_\eta(\zeta)})$ and $\delta' \notin \mathbf{domain}(e_{p_\eta(\zeta)})$ then from (9.14) and the fact that $W_\ell \subseteq \mathbf{domain}(p_\eta)$ it follows that if $\zeta \neq \zeta'$ then $e_{q_{\ell,n}^j(\zeta)}(\delta) \in A_\zeta \setminus A_{\zeta'}$ and $e_{q_{\ell,n}^{j'}(\zeta')}(\delta') \in A_{\zeta'} \setminus A_\zeta$ contradicting (9.19). Hence it must be that $\zeta = \zeta'$.

Appealing to (9.16) it follows that $j = j'$. Since $e_{q_{\ell,n}^j(\zeta)}$ is strictly increasing it follows that $\delta = \delta'$. It is immediate that

$H(\xi, \xi') \in [\omega]^{k+1}$ for all (ξ, ξ') in the domain of H .

Then define q_ℓ^j by

$$q_\ell^j(\zeta) = \begin{cases} q_{\ell,n}^j(0) \cup H \cup (p_\eta(0) \upharpoonright [\rho_\ell + 1]^2) & \text{if } \zeta = 0 \\ q_{\ell,n}^j(\zeta) & \text{if } \zeta \notin W_\ell \\ q_{\ell,n}^j(\zeta) \cup \{(\theta(\zeta, \ell), \rho_\ell)\} & \text{if } \zeta \in W_\ell. \end{cases}$$

Observe that $q_{\ell,n}^j(0) \cup (p_\eta(0) \upharpoonright [\nu_i]^2)$ is a function by (19) and (20). Next note that if $(e_{q_{\ell,n}^j(\zeta)}(\delta), \rho_\ell) \in \mathbf{domain}(H)$ then $\delta \notin \mathbf{domain}(p_\eta(\zeta))$ and hence $(e_{q_{\ell,n}^j(\zeta)}(\delta), \rho_\ell) \notin \mathbf{domain}(p_\eta(0))$ by (9.17). In other words, $q_{\ell,n}^j(0) \cup H \cup (p_\eta(0) \upharpoonright [\rho_\ell + 1]^2)$ is a function.

Hence, all that needs to be verified, therefore, is that for each $\zeta \in W_\ell$, $j \in k + 1$ and $\delta \in \mathbf{domain}(e_{q_\ell^j(\zeta)})$

$$q_\ell^j \Vdash_{\mathbb{P}_{\omega_2}} \text{“} R_\zeta(\delta, \theta(\zeta, \ell)) \in R_0(e_{q_\ell^j(\zeta)}(\delta), \rho_\ell)\text{”}.$$

But if $\zeta \in W_\ell$ then $\zeta \in \bigcup_{u=1}^m Z_u$ and so from (22) if

$$\delta \in \mathbf{domain}(e_{q_\ell^j(\zeta)}) \setminus \mathbf{domain}(e_{p_\eta(\zeta)} \upharpoonright \mu) = \mathbf{domain}(e_{q_{\ell,n}^j(\zeta)}) \setminus \mathbf{domain}(e_{p_\eta(\zeta)} \upharpoonright \mu)$$

it follows immediately from the definition of H that

$$q_\ell^j \Vdash_{\mathbb{P}_{\omega_1}} \text{“} R_\zeta(\delta, \theta(\zeta, \ell)) = r_{\zeta,n}^j(\delta) \in H(e_{q_\ell^j(\zeta)}(\delta), e_{q_\ell^j(\zeta)}(\theta(\zeta, \ell))) = R_0(e_{q_\ell^j(\zeta)}(\delta), \rho_\ell)\text{”}.$$

On the other hand, if $\delta \in \mathbf{domain}(e_{p_\eta(\zeta)})$ then by (23)

$$q_\ell^j \Vdash_{\mathbb{P}_{\omega_1}} \text{“} R_\zeta(\delta, \theta(\zeta, \ell)) = r_{p_\eta(\zeta)}(\delta, \theta(\zeta, \ell)) \in R_0(e_{p_\eta(\zeta)}(\delta), \rho_\ell) = R_0(e_{q_\ell^j(\zeta)}(\delta), \rho_\ell)\text{”}$$

with the inclusion holding since $p_\eta \upharpoonright \zeta \Vdash_{\mathbb{P}_\zeta} \text{“} p_\eta^j(\zeta) \in \mathbb{Q}_\zeta = \mathbb{E}(R_0, R_\zeta, A_\zeta, \mathfrak{M}_\zeta)\text{”}$ and the last equality holding by (21).

To see that (18) implies (16) use the first two cases of (9.18). To see that (21) implies (15) note that $\nu_i < \rho_\ell$ and $\theta(\zeta, \ell)$ is the maximum element of the domain of $e_{p_\eta(\zeta)} \upharpoonright \rho_\ell$ for each $\zeta \in Z_i$ because each $e_{p_\eta(\zeta)} \upharpoonright \rho_\ell$ is strictly increasing.

To carry out the construction of the $q_{\ell,i}^j$ let $q_{\ell,0}^j = \hat{q}_\ell^j$. Then, given $q_{\ell,i-1}^j$, let $\{\zeta_v\}_{v=1}^t$ be a decreasing enumeration of Z_i from (22) and construct conditions $\{z_v^j\}_{v=1}^t$ such that:

$$24. z_0^j = q_{\ell,i-1}^j$$

$$25. z_{v+1}^j \leq z_v^j$$

$$26. \mathbf{height}(z_v^j) \leq \nu_i \text{ for each } j$$

$$27. q_{\ell,u}^j(\zeta) = z_v^j(\zeta) \text{ if } u < i \text{ and } \zeta \in Z_u$$

$$28. z_u^j(\zeta) = z_v^j(\zeta) \text{ if } u < v \text{ and } \zeta \geq \zeta_u$$

$$29. z_v^j \sqsupset_{\nu_i} p_\eta \text{ for each } j$$

$$30. \text{ there is } r_v^j : \mathbf{domain}(z_v^j(\zeta_v)) \cap \nu_i \rightarrow k + 1 \text{ such that}$$

$$(\forall \delta \in \mathbf{domain}(e_{z_v^j(\zeta_v)} \cap \nu_i)) z_v^j \upharpoonright \zeta_v \Vdash_{\mathbb{P}_{\zeta_v}} \text{“} R_{\zeta_v}(\delta, \theta(\zeta_v, i)) = r_v^j(\delta)\text{”}.$$

$$31. \text{ if } \delta \in \mathbf{domain}(e_{p_\eta(\zeta_v)}) \text{ then } r_v^j(\delta) = r_{p_\eta(\zeta_v)}(\delta, \theta(\zeta_v, i)).$$

If this can be accomplished then let $q_{\ell,i}^j = z_i^j$ and let

$$r_{\zeta,i}^j = \begin{cases} r_v^j & \text{if } \zeta = \zeta_v \\ r_{\zeta,i-1}^j & \text{otherwise.} \end{cases}$$

To see that (30) implies (22) let $\zeta \in \bigcup_{u=1}^i Z_u$. Then if $\zeta \in Z_i$ there is some $v \leq t$ such that $\zeta = \zeta_v$. By (30) it follows that

$$z_v^j \upharpoonright \zeta_v \Vdash_{\mathbb{P}_{\zeta_v}} \text{“} R_{\zeta_v}(\delta, \theta(\zeta_v, i)) = r_v^j(\delta) = r_{\zeta,i}^j(\delta)\text{”}$$

and hence by (25) that

$$q_{\ell,i}^j \upharpoonright \zeta_v \Vdash_{\mathbb{P}_{\zeta_v}} \text{“} R_{\zeta_v}(\delta, \theta(\zeta_v, i)) = r_{\zeta,i}^j(\delta)\text{”}$$

for all $\delta \in \mathbf{domain}(e_{z_v^j(\zeta_v)}) \cap \nu_i$. But by (28) and the fact that the ζ_u are decreasing with u it follows that $\mathbf{domain}(e_{z_v^j(\zeta_v)}) \cap \nu_i = \mathbf{domain}(e_{q_{\ell,i}^j(\zeta_v)}) \cap \nu_i$ as required by (22). On the other hand, if $\zeta \in Z_u$ for some $u < i$ then by (27) and (28) it follows that $q_{\ell,i}^j(\zeta) = q_{\ell,u}^j(\zeta)$ and hence

$$q_{\ell,i}^j \upharpoonright \zeta \Vdash_{\mathbb{P}_\zeta} \text{“} R_\zeta(\delta, \theta(\zeta, i)) = r_{\zeta,i}^j(\delta)\text{”}$$

for all $\delta \in \mathbf{domain}(e_{q_{\ell,u}^j(\zeta)}) \cap \nu_i$ by the induction hypothesis.

To see that (31) implies that (23) holds the following claim will be used.

Claim 9.5. *If $1 \leq u \leq v \leq t$ and $\zeta \in Z_v$ then $Z_u \cap \mathfrak{M}_{\zeta, \kappa(\zeta, \ell)} \cap \zeta = \emptyset$.*

PROOF. Let $\bar{\zeta} \in Z_u \cap \zeta$. Then $\omega_1 \cap \mathfrak{M}_{\bar{\zeta}, \kappa(\bar{\zeta}, \ell)} = \nu_u$ and so $e_{p_\eta(\bar{\zeta})}(\theta(\bar{\zeta}, \ell)) = \rho_\ell$ and $\theta(\bar{\zeta}, \ell) \in \nu_u$. Since

$$\rho_\ell = e_{p_\eta(\zeta)}(\theta(\zeta, \ell)) \notin \nu_v = \omega_1 \cap \mathfrak{M}_{\zeta, \kappa(\zeta, \ell)}$$

it follows that $\mathfrak{M}_{\bar{\zeta}} \notin \mathfrak{M}_{\zeta, \kappa(\zeta, \ell)}$. Since (9.2) holds, it follows that $\bar{\zeta} \notin \mathfrak{M}_{\zeta, \kappa(\zeta, \ell)}$.

If $\zeta = \zeta_v$ for some $v \leq t$ then it is immediate that (31) implies (23). On the other hand, Claim 9.5 implies that there are no other instances to consider except those that have been dealt with at previous stages of the induction by (27).

It will now be shown how to construct z_v^j given z_{v-1}^j . Using the notation established in Definition 9.2, let

$$\bar{q} = \bigvee_{\delta \in \mathbf{domain}(e_{p_\eta(\zeta_v)})} q_{p_\eta(\zeta_v)}(\kappa(\zeta_v, i), \delta, \theta(\zeta_v, \ell)) \quad (9.20)$$

and note that $\bar{q} \in \mathbb{P}_{\zeta_v}$ because $\bar{q}(\zeta) \geq p_\eta(\zeta)$ for all $\zeta \in \zeta_v$ by (9.5). Moreover, $\bar{q} \in \mathfrak{M}_{\zeta_v, \kappa(\zeta_v, i)}$ because $\zeta_v \in \mathfrak{M}_{\zeta_v, 0}$. Observe that Claim 9.3 implies that $\bar{z} = z_{v-1}^j \vee \bar{q} \in \mathbb{P}_{\omega_2}$ because of (26), (29) and the fact that $\nu_i = \omega_1 \cap \mathfrak{M}_{\zeta_v, \kappa(\zeta_v, i)}$. Begin by defining the function z with domain $\mathbf{domain}(\bar{z}) \cap \mathfrak{M}_{\zeta_v, \kappa(\zeta_v, i)}$ such that $z(\zeta) = \bar{z}(\zeta)$. It may not be the case that $z \in \mathbb{P}_{\omega_2}$ but, nevertheless, the function z does belong to $\mathfrak{M}_{\zeta_v, \kappa(\zeta_v, i)}$. Since \bar{z} witnesses that the following statement holds:

$$\begin{aligned} (\exists q \in \mathbb{P}_{\omega_2})(\forall \zeta \in \mathbf{domain}(z)) e_{q(\zeta)} \supseteq e_{z(\zeta)} \text{ and } X_{q(\zeta)} \supseteq X_{z(\zeta)} \text{ and} \\ (\forall \theta \in \mathbf{domain}(e_{p_\eta(\zeta_v)})) q \upharpoonright \zeta \Vdash_{\mathbb{P}_\zeta} \text{“} R_\zeta(\theta, \theta(\zeta_v, \ell)) = r_{p_\eta(\zeta_v)}(\theta, \theta(\zeta_v, \ell))\text{”} \end{aligned} \quad (9.21)$$

it follows that there is $\bar{z} \in \mathfrak{M}_{\zeta_v, \kappa(\zeta_v, \ell)} \cap \mathbb{P}_{\omega_2}$ satisfying (9.21). By elementarity there is then $\bar{z}_v^j \leq \bar{z}$ in $\mathfrak{M}_{\zeta_v, \kappa(\zeta_v, \ell)} \cap \mathbb{P}_{\zeta_v}$ and r_v^j such that (30) holds. Then use Claim 9.3 to let $z_v^j = \bar{z}_v^j \vee z_{v-1}^j$. Note that $z_v^j \leq \bar{z}_v^j$, $z_v^j \leq z_{v-1}^j$ and (28) holds since $\bar{z}_v^j \in \mathbb{P}_{\zeta_v}$. Observe that (27) holds since if $\zeta \in \zeta_v \cap Z_u$ for some $u < i$ then $\zeta \notin \mathbf{domain}(\bar{z}_v^j)$ by Claim 9.5. Hence $z_v^j(\zeta) = z_{v-1}^j(\zeta)$ in this case.

PROOF (PROOF OF THEOREM 7.28). The model is obtained by forcing with \mathbb{P}_{ω_3} over a model V such that $V \models 2^{\aleph_1} = \aleph_2$. A key change from the proof of Theorem 7.27 and §1.4 of [1] is that \mathbb{Q}_0 will be defined to consist of functions from $[\Gamma]^2$ to $[\omega]^{< \aleph_0}$ where $\Gamma \in [\omega_1]^{< \aleph_0}$. In keeping with the notation of [1], if $G \subseteq \mathbb{Q}_0$ is generic then $\bigcup_{f \in G} f$ will be denoted by R_0 .

It is easy to see that in order to show that there is no $(2, 2)$ -universal function it suffices to show that there is no $(2, 1)$ -universal function. So suppose that $U : [\omega_1]^2 \rightarrow \omega$ is such a function. As in the proof of Theorem 7.27 let $\alpha \in \omega_3$ be such that U has a \mathbb{P}_α name. Also as in the proof of Theorem 7.27, for every limit $\bar{\alpha} > \alpha$ let $E^{\bar{\alpha}}$ be a name for an embedding of $c_{\bar{\alpha}} \circ T$ into F . Then let $\Gamma^{\bar{\alpha}}$ satisfy requirements (5) to (??) of Theorem 7.27 with $E^{\bar{\alpha}}$ in the place of E and $p_{\xi}^{\bar{\alpha}}$ in place of p_ξ .

Letting $D^{\bar{\alpha}} = \bigcup_{\xi \in \Gamma^{\bar{\alpha}}} \mathbf{domain}(p_{\xi}^{\bar{\alpha}})$ and using that $2^{\aleph_1} = \aleph_2$ it is possible to find $A \subseteq \omega_3$ of cardinality \aleph_3 and Γ such that:

1. $\Gamma^{\bar{\alpha}} = \Gamma$ for all $\bar{\alpha} \in A$
2. $\{D^{\bar{\alpha}} \mid \bar{\alpha} \in A\}$ form a Δ -system with root D^*
3. the structures $\{p_{\xi}^{\bar{\alpha}}\}_{\xi \in \Gamma}$ are all isomorphic and, indeed, if α_1 and α_2 are in A then there is a bijection $\Psi_{\alpha_1, \alpha_2} : D^{\alpha_1} \rightarrow D^{\alpha_2}$ such that $\Psi_{\alpha_1, \alpha_2} \upharpoonright D^*$ is the identity and $p_{\xi}^{\alpha_1}(\eta) = p_{\xi}^{\alpha_2}(\Psi_{\alpha_1, \alpha_2}(\eta))$ for all $\xi \in \Gamma$ and $\eta \in \mathbf{domain}(p_{\xi}^{\alpha_1})$.

Now let α_1 and α_2 be distinct elements of A . Observe that if $\xi \in \Gamma$ then because the $p_\xi^{\alpha_i}$ agree on D^* it is possible to define p_ξ to be $p_\xi^{\alpha_1} \vee p_\xi^{\alpha_2}$ and have that $p_\xi \in \mathbb{P}_{\omega_3}$ and $p_\xi \leq p_\xi^{\alpha_1}$ and $p_\xi \leq p_\xi^{\alpha_2}$.

Let $f \in \mathbb{C}$ be such that $p_\xi(\alpha_1) = p_\xi(\alpha_2) = f$ for all $\xi \in \Gamma$ and let $J \in \omega$ be such that $\mathbf{domain}(f) \subseteq J$. Now let $\xi_1 \in \Gamma$ be such that there is $n_1 > J$ such that $B = \{\xi \in \Gamma \mid T(\xi_1, \xi) = n_1\}$ is uncountable. Then let $\xi_2 < \xi_3$ be in B such that $T(\xi_2, \xi_3) = n_2 > n_1$. If it can be shown that there is $q \in \mathbb{P}_{\omega_3}$ such that $q \leq p_{\xi_i}$ for $i \leq 3$ and such that $q(\alpha_1)(n_1) = q(\alpha_1)(n_2)$ and $q(\alpha_2)(n_1) \neq q(\alpha_2)(n_2)$ then it would follow that

$$q \Vdash_{\mathbb{P}_{\omega_3}} \text{“} |\{c_{\alpha_1} \circ T(\xi_i, \xi_j) \mid 1 \leq i < j \leq 3\}| = 1 < |\{c_{\alpha_2} \circ T(\xi_i, \xi_j) \mid 1 \leq i < j \leq 3\}| \text{”}$$

$$(\forall i \leq 3) q \Vdash_{\mathbb{P}_{\omega_3}} \text{“} E^{\alpha_1}(\xi_i) = E^{\alpha_2}(\xi_i) \text{”}$$

implying, since the functions from ω to ω witnessing (2,1)-universality are one-to-one, that

$$q \Vdash_{\mathbb{P}_{\omega_3}} \text{“} |\{U(E^{\alpha_1}(\xi_i), E^{\alpha_1}(\xi_j)) \mid 1 \leq i < j \leq 3\}| < |\{U(E^{\alpha_2}(\xi_i), E^{\alpha_2}(\xi_j)) \mid 1 \leq i < j \leq 3\}| \text{”} \quad (9.22)$$

which is a contradiction since

$$\{U(E^{\alpha_1}(\xi_i), E^{\alpha_1}(\xi_j)) \mid 1 \leq i < j \leq 3\} = \{U(E^{\alpha_2}(\xi_i), E^{\alpha_2}(\xi_j)) \mid 1 \leq i < j \leq 3\}.$$

Applying the following claim first to p_{ξ_1} and p_{ξ_2} to yield \bar{q} and then to \bar{q} and p_{ξ_3} will furnish the q of (9.22).

Claim 9.6. *If q_1 and q_2 are in \mathbb{P}_{ω_3} and there are μ_1 and μ_2 such that*

- $\mathbf{height}(q_1) \leq \mu_1$
- $\mathbf{height}(q_2) \leq \mu_2$
- $q_1 \sqsubseteq_{\mu_1} q_2$

then there is q such that $q \leq q_1$, $q \leq q_2$ and $\mathbf{height}(q) \leq \mu_2$.

PROOF. As in the proof of Theorem 7.27 let $\{\rho_i\}_{i=1}^m$ enumerate

$$\{e_{q_2(\zeta)}(\theta) \mid \zeta \in \mathbf{domain}(q_2) \text{ and } \theta \in \mathbf{domain}(e_{q_2(\zeta)}) \setminus \mu_1\}$$

in increasing order. If $1 \leq i \leq m$ let

$$W_i = \{\zeta \in \mathbf{domain}(q_2) \cap W \mid (\exists \theta \in \mathbf{domain}(e_{q_2(\zeta)}) \setminus \mu_1) e_{q_2(\zeta)}(\theta) = \rho_i\}.$$

For each $\zeta \in W_i$ let $\nu(\zeta, i)$ be such that there is $\theta(\zeta, i) \in \mathfrak{N}_{\zeta, \nu(\zeta, i)}$ and $\kappa(\zeta, i)$ such that

$$\omega_1 \cap \mathfrak{N}_{\zeta, \nu(\zeta, i)} < \kappa(\zeta, i) < (\omega_1 \cap \mathfrak{N}_{\zeta, \nu(\zeta, i)}) \cdot 2$$

such that

$$e_{q_2(\zeta)}(\theta(\zeta, i)) \in \mathfrak{N}_{\zeta, \kappa(\zeta, i)+1} \setminus \mathfrak{N}_{\zeta, \kappa(\zeta, i)} \quad (9.23)$$

and then let $Y_i = \{\omega_1 \cap \mathfrak{N}_{\zeta, \kappa(\zeta, i)} \mid \zeta \in W_i\}$.

Conditions $\{\bar{q}_\ell\}_{1 \leq \ell \leq m}$ will be constructed by induction on ℓ such that the following are satisfied:

4. $\bar{q}_{\ell+1} \leq \bar{q}_\ell \leq q_1$
5. $\mathbf{height}(\bar{q}_\ell) \leq \rho_\ell + 1$
6. $\bar{q}_\ell \sqsupseteq_{\rho_\ell} q_2$

If this can be accomplished then it follows from (4) that $\bar{q}_m \leq q_1$ and from (6) that $\bar{q}_m \leq q_2$ and that $\mathbf{height}(\bar{q}_m) \leq \rho_m \leq \mu_2$ so that \bar{q}_m satisfies the conclusion of the claim.

To start, let $\bar{q}_0 = q_1$. The following additional induction hypothesis will be used: for every $\ell \leq m$ and $\zeta \in \mathbf{domain}(q_2)$

$$X_{\bar{q}_\ell(\zeta)} \supseteq \{\beta \in \rho_\ell \mid (\exists v \geq \ell) \{\beta, \rho_v\} \in \mathbf{domain}(q_2(0)) \setminus \mathbf{range}(e_{q_2(\zeta)})\}. \quad (9.24)$$

Given $\bar{q}_{\ell-1}$ let $\{\nu_i\}_{i=1}^n$ enumerate Y_ℓ in increasing order. Conditions $q_{\ell, i}$ will be constructed by induction on i such that:

7. $q_{\ell, i+1} \leq q_{\ell, i} \leq \bar{q}_{\ell-1}$
8. $\mathbf{height}(q_{\ell, i}) \leq \nu_i$
9. $q_{\ell, i} \sqsupseteq_{\nu_i} q_2$ for each $\eta \in \mathbf{domain}(q_{\ell, i})$
10. letting $Z_i = \{\zeta \in W_\ell \mid \omega_1 \cap \mathfrak{M}_{\zeta, \kappa(\zeta, \ell)} = \nu_i\}$ there is

$$r_{\zeta, i} : \mathbf{domain}(e_{q_{\ell, i}(\zeta)}) \cap \nu_i \rightarrow k+1$$

for each $\zeta \in \bigcup_{u=1}^{\ell} Z_u$ such that

$$q_{\ell, i} \upharpoonright \zeta \Vdash_{\mathbb{P}_\zeta} \text{“} R_\zeta(\delta, \theta(\zeta, i)) = r_{\zeta, i}(\delta)\text{”}$$

for all $\delta \in \mathbf{domain}(e_{q_{\ell, i}(\zeta)}) \cap \nu_i$

11. if $\zeta \in \bigcup_{u=1}^{\ell} Z_u$ and $\delta \in \mathbf{domain}(e_{q_2(\zeta)})$ then $r_{\zeta, i}(\delta) = r_{q_2(\zeta)}(\delta, \theta(\zeta, i))$ as in (9.5).

If this can be accomplished then define H to be the function with domain

$$\{(\zeta, \delta) \mid \zeta \in W_\ell \text{ and } \delta \in \mathbf{domain}(e_{q_{\ell, n}(\zeta)}) \setminus \mathbf{domain}(e_{q_2(\zeta)})\}$$

and constant value

$$\{r_{\zeta, n}(\delta) \mid (\zeta, \delta) \in \mathbf{domain}(H)\}$$

for all $(\zeta, \delta) \in \mathbf{domain}(H)$.

Then define \bar{q}_ℓ by

$$\bar{q}_\ell(\zeta) = \begin{cases} q_{\ell, n}(0) \cup H \cup q_2(0) \upharpoonright [\rho_\ell]^2 & \text{if } \zeta = 0 \\ q_{\ell, n}(\zeta) & \text{if } \zeta \notin W_\ell \\ q_{\ell, n}(\zeta) \cup \{(\theta(\zeta, \ell), \rho_\ell)\} & \text{if } \zeta \in W_\ell. \end{cases}$$

Observe that $q_{\ell, n}(0) \cup (q_2(0) \upharpoonright [\rho_\ell]^2)$ is a function by (8) and (9). Next note that if

$$(e_{q_{\ell, n}(\zeta)}(\delta), \rho_\ell) \in \mathbf{domain}(H)$$

then $\delta \notin \mathbf{domain}(q_2(\zeta))$ and hence $(e_{q_{\ell, n}(\zeta)}(\delta), \rho_\ell) \notin \mathbf{domain}(q_2(0))$ by (9.24). In other words,

$$q_{\ell, n}(0) \cup H \cup (q_2(0) \upharpoonright [\rho_\ell + 1]^2)$$

is a function.

All that needs to be verified, therefore, is that for each $\zeta \in W_\ell$, and $\delta \in \mathbf{domain}(e_{\bar{q}_\ell(\zeta)})$

$$\bar{q}_\ell \Vdash_{\mathbb{P}_{\omega_2}} \text{“} R_\zeta(\delta, \theta(\zeta, \ell)) \in R_0(e_{q_\ell(\zeta)}(\delta), \rho_\ell)\text{”}.$$

But if $\zeta \in W_\ell$ then $\zeta \in \bigcup_{u=1}^n Z_u$ and so from (10) if

$$\delta \in \mathbf{domain}(e_{\bar{q}_\ell(\zeta)}) \setminus \mathbf{domain}(e_{q_2(\zeta)} \upharpoonright \mu) = \mathbf{domain}(e_{q_{\ell, n}(\zeta)}) \setminus \mathbf{domain}(e_{q_2(\zeta)} \upharpoonright \mu)$$

it follows immediately from the definition of H that

$$\bar{q}_\ell \Vdash_{\mathbb{P}_{\omega_1}} \text{“} R_\zeta(\delta, \theta(\zeta, \ell)) = r_{\zeta, n}(\delta) \in H(e_{\bar{q}_\ell(\zeta)}(\delta), e_{\bar{q}_\ell(\zeta)}(\theta(\zeta, \ell))) = R_0(e_{\bar{q}_\ell(\zeta)}(\delta), \rho_\ell)\text{”}.$$

On the other hand, if $\delta \in \mathbf{domain}(e_{q_2(\zeta)})$ then by (11)

$$\bar{q}_\ell \Vdash_{\mathbb{P}_{\omega_1}} \text{“} R_\zeta(\delta, \theta(\zeta, \ell)) = r_{q_2(\zeta)}(\delta, \theta(\zeta, \ell)) \in R_0(e_{q_2(\zeta)}(\delta), \rho_\ell) = R_0(e_{\bar{q}_\ell(\zeta)}(\delta), \rho_\ell)\text{”}$$

with the inclusion holding since $q_2 \upharpoonright \zeta \Vdash_{\mathbb{P}_\zeta} \text{“} q_2(\zeta) \in \mathbb{Q}_\zeta = \mathbb{E}(R_0, R_\zeta, \omega_1, \mathfrak{M}_\zeta)\text{”}$ and the last equality holding by (9). To see that (9) implies (6) note that $\nu_i < \rho_\ell$ and $\theta(\zeta, \ell)$ is the maximum element of the domain of $e_{q_2(\zeta)} \upharpoonright \rho_\ell$ for each $\zeta \in Z_i$ because each $e_{q_2(\zeta)} \upharpoonright \rho_\ell$ is strictly increasing.

To carry out the construction of the $q_{\ell, i}$ let $q_{\ell, 0} = \bar{q}_{\ell-1}$. Then, given $q_{\ell, i}$, let $\{\zeta_v\}_{v=1}^t$ be a decreasing enumeration of Z_i and construct conditions $\{z_v\}_{v=1}^t$ such that:

12. $z_0 = q_{\ell, i}$
13. $z_{v+1} \leq z_v$
14. $\mathbf{height}(z_v) \leq \nu_i$
15. $z_v^j(\zeta) \sqsubseteq_{\nu_i} e_{q_2}$
16. there is $r_v : \mathbf{domain}(z_v(\zeta_v)) \cap \nu_i \rightarrow k + 1$ such that

$$(\forall \delta \in \mathbf{domain}(z_v(\zeta_v)) \cap \nu_i) z_v \upharpoonright \zeta_v \Vdash_{\mathbb{P}_{\zeta_v}} "R_{\zeta_v}(\delta, \theta(\zeta_v, i)) = r_v(\delta)"$$

$$(\forall \delta \in \mathbf{domain} e_{\bar{q}_{\ell-1}(\zeta_v)}) r_v(\delta) = r_{\bar{q}_{\ell-1}(\zeta_v)}(\delta, \theta(\zeta, i)).$$

If this can be accomplished then let $q_{\ell, i+1} = z_t$ and let $r_{\zeta_v, i+1} = r_v$. Note that (16) implies that (10) and (11) hold. To construct z_v given z_{v-1} begin by defining \bar{q} as in (9.20) and then proceed exactly as in the proof of Theorem 7.27.