

ON INCOMPACTNESS FOR CHROMATIC NUMBER OF
GRAPHS
SH1006

SAHARON SHELAH

ABSTRACT. We deal with compactness. Assume the existence of non-reflecting stationary subset of the regular cardinal λ of cofinality κ . We prove that one can define a graph G whose chromatic number is $> \kappa$, while the chromatic number of every subgraph $G' \subseteq G, |G'| < \lambda$ is $\leq \kappa$. The main case is $\kappa = \aleph_0$.

Date: January 31, 2014.

2010 Mathematics Subject Classification. Primary: 03E05; Secondary: 05C15.

Key words and phrases. set theory, graphs, chromatic number, compactness, non-reflecting stationary sets.

The author thanks Alice Leonhardt for the beautiful typing. First typed: July 3, 2011. Pf for J.: Sept 2012.

Anotated Content

§0 Introduction, pg.3

§(0A) The questions and results, pg.3

§(0AB) Preliminaries, pg.3

§1 From a non-reflecting stationary set, pg.5

[We show that “ $S \subseteq S_\kappa^\lambda$ is stationary not reflecting” implies compactness for length λ for “chromatic number = κ ”.]

§2 From almost free, pg.8

[Here we weaken the assumption in §1 to “ $\mathcal{A} \subseteq {}^\kappa \text{Ord}$ is almost free”.]

§ 0. INTRODUCTION

§ 0(A). **The questions and results.** During the Hajnal conference (June 2011) Magidor asked me on incompactness of “having chromatic number \aleph_0 ”; that is, there is a graph G with λ nodes, chromatic number $> \aleph_0$ but every subgraph with $< \lambda$ nodes has chromatic number \aleph_0 when:

- (*)₁ λ is regular $> \aleph_1$ with a non-reflecting stationary $S \subseteq S_{\aleph_0}^\lambda$, possibly though better not, assuming some version of GCH.

Subsequently also when:

- (*)₂ $\lambda = \aleph_{\omega+1}$.

Such problems were first asked by Erdős-Hajnal, see [?]; we continue [?].

First answer was using BB, see [?, 3.24] so assuming

- ⊕ (a) $\lambda = \mu^+$
 (b) $\mu^{\aleph_0} = \mu$
 (c) $S \subseteq \{\delta < \lambda : \text{cf}(\delta) = \aleph_0\}$ is stationary not reflecting

or just

- ⊕' (a) $\lambda = \text{cf}(\lambda)$
 (b) $\alpha < \lambda \Rightarrow |\alpha|^{\aleph_0} < \lambda$
 (c) as above.

However, eventually we get more: if $\lambda = \aleph^{\aleph_0} = \text{cf}(\lambda)$ and $S \subseteq S_{\aleph_0}^\lambda$ is stationary non-reflective then we have λ -incompactness for \aleph_0 -chromatic. In fact, we replace \aleph_0 by $\kappa = \text{cf}(\kappa) < \lambda$ using a suitable hypothesis.

Moreover, if $\lambda^\kappa > \lambda$ we still get $(\lambda^\kappa, \lambda)$ -incompactness for κ -chromatic number. In §2 we use quite free family of countable sequences.

In subsequent work we shall solve also the parallel of the second question of Magidor, i.e.

- (*)₂ for regular $\kappa \geq \aleph_0$ and $n < \omega$ there is a graph G of chromatic number $> \kappa$ but every sub-graph with $< \aleph_{\kappa \cdot n+1}$ nodes has chromatic number $\leq \kappa$.

In fact, considerably is proved, see [?]. We thank Menachem Magidor for asking, Peter Komjath for stimulating discussion and Paul Larson, Shimoni Garti and the referee for some comments.

§ 0(B). Preliminaries.

Definition 0.1. For a graph G , let $\text{ch}(G)$, the chromatic number of G be the minimal cardinal χ such that there is colouring \mathbf{c} of G with χ colours, that is \mathbf{c} is a function from the set of nodes of G into χ or just a set of of cardinality $\leq \chi$ such that $\mathbf{c}(x) = \mathbf{c}(y) \Rightarrow \{x, y\} \notin \text{edge}(G)$.

Definition 0.2. 1) We say “we have λ -incompactness for the $(< \chi)$ -chromatic number” or $\text{INC}_{\text{chr}}(\lambda, < \chi)$ when: there is a graph G with λ nodes, chromatic number $\geq \chi$ but every subgraph with $< \lambda$ nodes has chromatic number $< \chi$.
 2) If $\chi = \mu^+$ we may replace “ $< \chi$ ” by μ ; similarly in 0.3.

We also consider

Definition 0.3. 1) We say “we have (μ, λ) -incompactness for $(< \chi)$ -chromatic number” or $\text{INC}_{\text{chr}}(\mu, \lambda, < \chi)$ when there is an increasing continuous sequence $\langle G_i : i \leq \lambda \rangle$ of graphs each with $\leq \mu$ nodes, G_i an induced subgraph of G_λ with $\text{ch}(G_\lambda) \geq \chi$ but $i < \lambda \Rightarrow \text{ch}(G_i) < \chi$.
 2) Replacing (in part (1)) χ by $\bar{\chi} = \langle \chi_0, \chi_1 \rangle$ means $\text{ch}(G_\lambda) \geq \chi_1$ and $i < \lambda \rightarrow \text{ch}(G_i) < \chi_0$; similarly in 0.2 and parts 3),4) below.
 3) We say we have incompactness for length λ for $(< \chi)$ -chromatic (or $\bar{\chi}$ -chromatic) number when we fail to have (μ, λ) -compactness for $(< \chi)$ -chromatic (or $\bar{\chi}$ -chromatic) number for some μ .
 4) We say we have $[\mu, \lambda]$ -incompactness for $(< \chi)$ -chromatic number or $\text{INC}_{\text{chr}}[\mu, \lambda, < \chi]$ when there is a graph G with μ nodes, $\text{ch}(G) \geq \chi$ but $G^1 \subseteq G \wedge |G^1| < \lambda \Rightarrow \text{ch}(G^1) < \chi$.
 5) Let $\text{INC}_{\text{chr}}^+(\mu, \lambda, < \chi)$ be as in part (1) but we add that even the $\text{cl}(G_i)$, the colouring number of G_i is $< \chi$ for $i < \lambda$, see below.
 6) Let $\text{INC}_{\text{chr}}^+[\mu, \lambda, < \chi]$ be as in part (4) but we add $G^1 \subseteq G \wedge |G^1| < \lambda \Rightarrow \text{cl}(G^1) < \chi$.
 7) If $\chi = \kappa^+$ we may write κ instead of “ $< \chi$ ”.

Definition 0.4. 1) For regular $\lambda > \kappa$ let $S_\kappa^\lambda = \{\delta < \lambda : \text{cf}(\delta) = \kappa\}$.
 2) We say C is a $(\geq \theta)$ -closed subset of a set B of ordinals when: if $\delta = \sup(\delta \cap B) \in B$, $\text{cf}(\delta) \geq \theta$ and $\delta = \sup(C \cap \delta)$ then $\delta \in C$.

Definition 0.5. For a graph G , the colouring number $\text{cl}(G)$ is the minimal κ such that there is a list $\langle a_\alpha : \alpha < \alpha(*) \rangle$ of the nodes of G such that $\alpha < \alpha(*) \Rightarrow \kappa > |\{\beta < \alpha : \{a_\beta, a_\alpha\} \in \text{edge}(G)\}|$.

§ 1. FROM NON-REFLECTING STATIONARY IN COFINALITY \aleph_0

Claim 1.1. *There is a graph G with λ nodes and chromatic number $> \kappa$ but every subgraph with $< \lambda$ nodes have chromatic number $\leq \kappa$ when:*

- ⊕ (a) λ, κ are regular cardinals
- (b) $\kappa < \lambda = \lambda^\kappa$
- (c) $S \subseteq S_\kappa^\lambda$ is stationary, not reflecting.

Proof. Stage A: Let $\bar{X} = \langle X_i : i < \lambda \rangle$ be a partition of λ to sets such that $|X_i| = \bar{\lambda}$ or just $|X_i| = |i + 2|^\kappa$ and $\min(X_i) \geq i$ and let $X_{<i} = \cup\{X_j : j < i\}$ and $X_{\leq i} = X_{<(i+1)}$. For $\alpha < \lambda$ let $\mathbf{i}(\alpha)$ be the unique ordinal $i < \lambda$ such that $\alpha \in X_i$. We choose the set of points = nodes of G as $Y = \{(\alpha, \beta) : \alpha < \beta < \lambda, \mathbf{i}(\beta) \in S \text{ and } \alpha < \mathbf{i}(\beta)\}$ and let $Y_{<i} = \{(\alpha, \beta) \in Y : \mathbf{i}(\beta) < i\}$.

Stage B: Note that if $\lambda = \kappa^+$, the complete graph with λ nodes is an example (no use of the further information in ⊕). So without loss of generality $\lambda > \kappa^+$.

Now choose a sequence satisfying the following properties, exists by [?, Ch.III]:

- ⊕ (a) $\bar{C} = \langle C_\delta : \delta \in S \rangle$
- (b) $C_\delta \subseteq \delta = \sup(C_\delta)$
- (c) $\text{otp}(C_\delta) = \kappa$ such that $(\forall \beta \in C_\delta)(\beta + 1, \beta + 2 \notin C_\delta)$
- (d) \bar{C} guesses¹clubs.

Let $\langle \alpha_{\delta, \varepsilon}^* : \varepsilon < \kappa \rangle$ list C_δ in increasing order.

For $\delta \in S$ let Γ_δ be the set of sequence $\bar{\beta}$ such that:

- ⊕ _{$\bar{\beta}$} (a) $\bar{\beta}$ has the form $\langle \beta_\varepsilon : \varepsilon < \kappa \rangle$
- (b) $\bar{\beta}$ is increasing with limit δ
- (c) $\alpha_{\delta, \varepsilon}^* < \beta_{2\varepsilon+i} < \alpha_{\delta, \varepsilon+1}^*$ for $i < 2, \varepsilon < \kappa$
- (d) $\beta_{2\varepsilon+i} \in X_{<\alpha_{\delta, \varepsilon+1}^*} \setminus X_{\leq \alpha_{\delta, \varepsilon}^*}$ for $i < 2, \varepsilon < \kappa$
- (e) $(\beta_{2\varepsilon}, \beta_{2\varepsilon+1}) \in Y$ hence $\in Y_{<\alpha_{\delta, \varepsilon+1}^*} \subseteq Y_{<\delta}$ for each $\varepsilon < \kappa$

(can ask less).

So $|\Gamma_\delta| \leq |\delta|^\kappa \leq |X_\delta| \leq \lambda$ hence we can choose a sequence $\langle \bar{\beta}_\gamma : \gamma \in X'_\delta \subseteq X_\delta \rangle$ listing Γ_δ .

Now we define the set of edges of G : $\text{edge}(G) = \{(\alpha_1, \alpha_2), (\min(C_\delta), \gamma)\} : \delta \in S, \gamma \in X'_\delta$ hence the sequence $\bar{\beta}_\gamma = \langle \beta_{\gamma, \varepsilon} : \varepsilon < \kappa \rangle$ is well defined and we demand $(\alpha_1, \alpha_2) \in \{(\beta_{\gamma, 2\varepsilon}, \beta_{\gamma, 2\varepsilon+1}) : \varepsilon < \kappa\}$.

Stage C: Every subgraph of G of cardinality $< \lambda$ has chromatic number $\leq \kappa$.

For this we shall prove that:

$$\oplus_1 \text{ch}(G|Y_{<i}) \leq \kappa \text{ for every } i < \lambda.$$

This suffice as λ is regular, hence every subgraph with $< \lambda$ nodes is included in $Y_{<i}$ for some $i < \lambda$.

For this we shall prove more by induction on $j < \lambda$:

¹the guessing clubs are used only in Stage D.

$\oplus_{2,j}$ if $i < j, i \notin S, \mathbf{c}_1$ a colouring of $G \upharpoonright Y_{<i}, \text{Rang}(\mathbf{c}_1) \subseteq \kappa$ and $u \in [\kappa]^\kappa$ then there is a colouring \mathbf{c}_2 of $G \upharpoonright Y_{<j}$ extending \mathbf{c}_1 such that $\text{Rang}(\mathbf{c}_2 \upharpoonright (Y_{<j} \setminus Y_{<i})) \subseteq u$.

Case 1: $j = 0$
Trivial.

Case 2: j successor, $j - 1 \notin S$

Let i be such that $j = i + 1$, but then every node from $Y_j \setminus Y_i$ is an isolated node in $G \upharpoonright Y_{<j}$, because if $\{(\alpha, \beta), (\alpha', \beta')\}$ is an edge of $G \upharpoonright Y_j$ then $\mathbf{i}(\beta), \mathbf{i}(\beta') \in S$ hence necessarily $\mathbf{i}(\beta) \neq j - 1 = i, \mathbf{i}(\beta') \neq j - 1 = i$ hence both $(\alpha, \beta), (\alpha', \beta')$ are from Y_i .

Case 3: j successor, $j - 1 \in S$

Let $j - 1$ be called δ so $\delta \in S$. But $i \notin S$ by the assumption in $\oplus_{2,j}$ hence $i < \delta$. Let $\varepsilon(*) < \kappa$ be such that $\alpha_{\delta, \varepsilon(*)}^* > i$.

Let $\langle u_\varepsilon : \varepsilon \leq \kappa \rangle$ be a sequence of subsets of u , a partition of u to sets each of cardinality κ ; actually the only disjointness used is that $u_\kappa \cap (\bigcup_{\varepsilon < \kappa} u_\varepsilon) = \emptyset$.

We let $i_0 = i, i_{1+\varepsilon} = \cup\{\alpha_{\delta, \varepsilon(*)+1+\zeta}^* + 1 : \zeta < 1 + \varepsilon\}$ for $\varepsilon < \kappa, i_\kappa = \delta$ and $i_{\kappa+1} = \delta + 1 = j$.

Note that:

- $\varepsilon < \kappa \Rightarrow i_\varepsilon \notin S_j$.

[Why? For $\varepsilon = 0$ by the assumption on i , for ε successor i_ε is a successor ordinal and for i limit clearly $\text{cf}(i_\varepsilon) = \text{cf}(\varepsilon) < \kappa$ and $S \subseteq S_\kappa^\lambda$.]

We now choose $\mathbf{c}_{2,\zeta}$ by induction on $\zeta \leq \kappa + 1$ such that:

- $\mathbf{c}_{2,0} = \mathbf{c}_1$
- $\mathbf{c}_{2,\zeta}$ is a colouring of $G \upharpoonright Y_{<i_\zeta}$
- $\mathbf{c}_{2,\zeta}$ is increasing with ζ
- $\text{Rang}(\mathbf{c}_{2,\zeta} \upharpoonright (Y_{<i_{\zeta+1}} \setminus Y_{<i_\zeta})) \subseteq u_\xi$ for every $\xi < \zeta$.

For $\zeta = 0, \mathbf{c}_{2,0}$ is \mathbf{c}_1 so is given.

For $\zeta = \varepsilon + 1 < \kappa$: use the induction hypothesis, possible as necessarily $i_\varepsilon \notin S$.

For $\zeta \leq \kappa$ limit: take union.

For $\zeta = \kappa + 1$, note that each node b of $Y_{<i_\zeta} \setminus Y_{<i_\kappa}$ is not connected to any other such node and if the node b is connected to a node from $Y_{<i_\kappa}$ then the node b necessarily has the form $(\min(C_\delta), \gamma), \gamma \in X'_\delta$, hence $\bar{\beta}_\gamma$ is well defined, so the node $b = (\min(C_\delta), \gamma)$ is connected in G , more exactly in $G \upharpoonright Y_{\leq \delta}$ exactly to the κ nodes $\{(\beta_{\gamma, 2\varepsilon}, \beta_{\gamma, 2\varepsilon+1}) : \varepsilon < \kappa\}$, but for every $\varepsilon < \kappa$ large enough, $\mathbf{c}_{2,\kappa}((\beta_{\gamma, 2\varepsilon}, \beta_{\gamma, 2\varepsilon+1})) \in u_\varepsilon$ hence $\notin u_\kappa$ and $|u_\kappa| = \kappa$ so we can choose a colour.

Case 4: j limit

By the assumption of the claim there is a club e of j disjoint to S and without loss of generality $\min(e) = i$. Now choose $\mathbf{c}_{2,\xi}$ a colouring of $Y_{<\xi}$ by induction on $\xi \in e \cup \{j\}$, increasing with ξ such that $\text{Rang}(\mathbf{c}_{2,\xi} \upharpoonright (Y_{<\xi} \setminus Y_{<i})) \subseteq u$ and $\mathbf{c}_{2,0} = \mathbf{c}_1$

- For $\xi = \min(e) = i$ the colouring $\mathbf{c}_{2,\xi} = \mathbf{c}_{2,i} = \mathbf{c}_1$ is given,

- for ξ successor in e , i.e. $\in \text{nacc}(e) \setminus \{i\}$, use the induction hypothesis with ξ , $\max(e \cap \xi)$ here playing the role of j, i there recalling $\max(e \cap \xi) \in e, e \cap S = \emptyset$
- for $\xi = \sup(e \cap \xi)$ take union.

Lastly, for $\xi = j$ we are done.

Stage D: $\text{ch}(G) > \kappa$.

Why? Toward a contradiction, assume \mathbf{c} is a colouring of G with set of colours $\subseteq \kappa$. For each $\gamma < \lambda$ let $u_\gamma = \{\mathbf{c}((\alpha, \beta)) : \gamma < \alpha < \beta < \lambda \text{ and } (\alpha, \beta) \in Y\}$. So $\langle u_\gamma : \gamma < \lambda \rangle$ is \subseteq -decreasing sequence of subsets of κ and $\kappa < \lambda = \text{cf}(\lambda)$, hence for some $\gamma(*) < \lambda$ and $u_* \subseteq \kappa$ we have $\gamma \in (\gamma(*), \lambda) \Rightarrow u_\gamma = u_*$.

Hence $E = \{\delta < \lambda : \delta \text{ is a limit ordinal } > \gamma(*) \text{ and } (\forall \alpha < \delta)((\mathbf{i}(\alpha) < \delta) \text{ and for every } \gamma < \delta \text{ and } i \in u_* \text{ there are } \alpha < \beta \text{ from } (\gamma, \delta) \text{ such that } (\alpha, \beta) \in Y \text{ and } \mathbf{c}((\alpha, \beta)) = i)\}$ is a club of λ .

Now recall that \bar{C} guesses clubs hence for some $\delta \in S$ we have $C_\delta \subseteq E$, so for every $\varepsilon < \kappa$ we can choose $\beta_{2\varepsilon} < \beta_{2\varepsilon+1}$ from $(\alpha_{\delta, \varepsilon}^*, \alpha_{\delta, \varepsilon+1}^*)$ such that $(\beta_{2\varepsilon}, \beta_{2\varepsilon+1}) \in Y$ and $\varepsilon \in u_* \Rightarrow \mathbf{c}((\beta_{2\varepsilon}, \beta_{2\varepsilon+1})) = \varepsilon$. So $\langle \beta_\varepsilon : \varepsilon < \kappa \rangle$ is well defined, increasing and belongs to Γ_δ , hence $\bar{\beta}_\gamma = \langle \beta_\varepsilon : \varepsilon < \kappa \rangle$ for some $\gamma \in X_\delta$, hence $(\alpha_{\delta, 0}^*, \gamma)$ belongs to Y and is connected in the graph to $(\beta_{2\varepsilon}, \beta_{2\varepsilon+1})$ for $\varepsilon < \kappa$. Now if $\varepsilon \in u_*$ then $\mathbf{c}((\beta_{2\varepsilon}, \beta_{2\varepsilon+1})) = \varepsilon$ hence $\mathbf{c}((\alpha_{\delta, 0}^*, \gamma)) \neq \varepsilon$ for every $\varepsilon \in u_*$, so $\mathbf{c}((\alpha_{\delta, 0}^*, \gamma)) \in \kappa \setminus u_*$. But $u_* = u_{\alpha_{\delta, 0}^*}$ and $\mathbf{c}((\alpha_{\delta, 0}^*, \gamma)) \in \kappa \setminus u_*$, so we get contradiction to the definition of $u_{\alpha_{\delta, 0}^*}$. $\square_{1.1}$

Similarly

Claim 1.2. *There is an increasing continuous sequence $\langle G_i : i \leq \lambda \rangle$ of graphs each of cardinality λ^κ such that $\text{ch}(G_\lambda) > \kappa$ and $i < \lambda$ implies $\text{ch}(G_i) \leq \kappa$ and even $\text{cl}(G_i) \leq \kappa$ when:*

- \boxplus (a) $\lambda = \text{cf}(\lambda)$
- (b) $S \subseteq \{\delta < \lambda : \text{cf}(\delta) = \kappa\}$ is stationary not reflecting.

Proof. Like 1.1 but the X_i are not necessarily $\subseteq \lambda$ or use 2.2. $\square_{1.2}$

§ 2. FROM ALMOST FREE

Definition 2.1. Suppose $\eta_\beta \in {}^\kappa \text{Ord}$ for every $\beta < \alpha(*)$ and $u \subseteq \alpha(*)$, and $\alpha < \beta < \alpha(*) \Rightarrow \eta_\alpha \neq \eta_\beta$.

1) We say $\{\eta_\alpha : \alpha \in u\}$ is free when there exists a function $h : u \rightarrow \kappa$ such that $\langle \{\eta_\alpha(\varepsilon) : \varepsilon \in [h(\alpha), \kappa)\} : \alpha \in u \rangle$ is a sequence of pairwise disjoint sets.

2) We say $\{\eta_\alpha : \alpha \in u\}$ is weakly free when there exists a sequence $\langle u_{\varepsilon, \zeta} : \varepsilon, \zeta < \kappa \rangle$ of subsets of u with union u , such that the function $\eta_\alpha \mapsto \eta_\alpha(\varepsilon)$ is a one-to-one function on $u_{\varepsilon, \zeta}$, for each $\varepsilon, \zeta < \kappa$.

Claim 2.2. 1) We have $\text{INC}_{\text{chr}}(\mu, \lambda, \kappa)$ and even $\text{INC}_{\text{chr}}^+(\mu, \lambda, \kappa)$, see Definition 0.3(1), (5) when:

- ⊞ (a) $\alpha(*) \in [\mu, \mu^+)$ and λ is regular $\leq \mu$ and $\mu = \mu^\kappa$
- (b) $\bar{\eta} = \langle \eta_\alpha : \alpha < \alpha(*) \rangle$
- (c) $\eta_\alpha \in {}^\kappa \mu$
- (d) $\langle u_i : i \leq \lambda \rangle$ is a \subseteq -increasing continuous sequence of subsets of $\alpha(*)$ with $u_\lambda = \alpha(*)$
- (e) $\bar{\eta} \upharpoonright u_\alpha$ is free iff $\alpha < \lambda$ iff $\bar{\eta} \upharpoonright u_\alpha$ is weakly free.

2) We have $\text{INC}_{\text{chr}}[\mu, \lambda, \kappa]$ and even $\text{INC}_{\text{chr}}^+[\mu, \lambda, \kappa]$, see Definition 0.3(4) when:

- ⊞₂ (a), (b), (c) as in ⊞ from 2.2
- (d) $\bar{\eta}$ is not free
- (e) $\bar{\eta} \upharpoonright u$ is free when $u \in [\alpha(*)]^{< \lambda}$.

Proof. We concentrate on proving part (1) the chromatic number case; the proof of part (2) and of the colouring number are similar. For $\mathcal{A} \subseteq {}^\kappa \text{Ord}$, we define $\tau_{\mathcal{A}}$ as the vocabulary $\{P_\eta : \eta \in \mathcal{A}\} \cup \{F_\varepsilon : \varepsilon < \kappa\}$ where P_η is a unary predicate, F_ε a unary function (will be interpreted as possibly partial).

Without loss of generality for each $i < \lambda$, u_i is an initial segment of $\alpha(*)$ and let $\mathcal{A} = \{\eta_\alpha : \alpha < \alpha(*)\}$ and let $<_{\mathcal{A}}$ be the well ordering $\{(\eta_\alpha, \eta_\beta) : \alpha < \beta < \alpha(*)\}$ of \mathcal{A} .

We further let $K_{\mathcal{A}}$ be the class of structures M such that (pedantically, $K_{\mathcal{A}}$ depend also on the sequence $\langle \eta_\alpha : \alpha < \alpha(*) \rangle$):

- ⊞₁ (a) $M = (|M|, F_\varepsilon^M, P_\eta^M)_{\varepsilon < \kappa, \eta \in \mathcal{A}}$
- (b) $\langle P_\eta^M : \eta \in \mathcal{A} \rangle$ is a partition of $|M|$, so for $a \in M$ let $\eta_a = \eta_a^M$ be the unique $\eta \in \mathcal{A}$ such that $a \in P_\eta^M$
- (c) if $a_\ell \in P_{\eta_\ell}^M$ for $\ell = 1, 2$ and $F_\varepsilon^M(a_2) = a_1$ then $\eta_1(\varepsilon) = \eta_2(\varepsilon)$ and $\eta_1 <_{\mathcal{A}} \eta_2$.

Let $K_{\mathcal{A}}^*$ be the class of M such that

- ⊞₂ (a) $M \in K_{\mathcal{A}}$
- (b) $\|M\| = \mu$
- (c) if $\eta \in \mathcal{A}$, $u \subseteq \kappa$ and $\eta_\varepsilon <_{\mathcal{A}} \eta$, $\eta_\varepsilon(\varepsilon) = \eta(\varepsilon)$ and $a_\varepsilon \in P_{\eta_\varepsilon}^M$ for $\varepsilon \in u$ then for some $a \in P_\eta^M$ we have $\varepsilon \in u \Rightarrow F_\varepsilon^M(a) = a_\varepsilon$ and $\varepsilon \in \kappa \setminus u \Rightarrow F_\varepsilon^M(a)$ not defined.

Clearly

\boxplus_3 there is $M \in K_{\mathcal{A}}^*$.

[Why? As $\mu = \mu^\kappa$ and $|\mathcal{A}| = \mu$.]

\boxplus_4 for $M \in K_{\mathcal{A}}$ let G_M be the graph with:

- set of nodes $|M|$
- set of edges $\{\{a, F_\varepsilon^M(a)\} : a \in |M|, \varepsilon < \kappa \text{ when } F_\varepsilon^M(a) \text{ is defined}\}$.

Now

\boxplus_5 if $u \subseteq \alpha(*)$, $\mathcal{A}_u = \{\eta_\alpha : \alpha \in u\} \subseteq \mathcal{A}$ and $\bar{\eta} \upharpoonright u$ is free, and $M \in K_{\mathcal{A}}$ then $G_{M, \mathcal{A}_u} := G_M \upharpoonright (\cup\{P_\eta^M : \eta \in \mathcal{A}_u\})$ has chromatic number $\leq \kappa$; moreover has colouring number $\leq \kappa$.

[Why? Let $h : u \rightarrow \kappa$ witness that $\bar{\eta} \upharpoonright u$ is free and for $\varepsilon < \kappa$ let $\mathcal{B}_\varepsilon := \{\eta_\alpha : \alpha \in u \text{ and } h(\alpha) = \varepsilon\}$, so $\mathcal{B} = \cup\{\mathcal{B}_\varepsilon : \varepsilon < \kappa\}$, hence it is enough to prove for each $\varepsilon < \kappa$ that $G_{\mu, \mathcal{B}_\varepsilon}$ has chromatic number $\leq \kappa$. To prove this, by induction on $\alpha \leq \alpha(*)$ we choose $\mathbf{c}_\alpha^\varepsilon$ such that:

- $\boxplus_{5.1}$ (a) $\mathbf{c}_\alpha^\varepsilon$ is a function
 (b) $\langle \mathbf{c}_\beta : \beta \leq \alpha \rangle$ is increasing continuous
 (c) $\text{Dom}(\mathbf{c}_\alpha^\varepsilon) = B_\alpha^\varepsilon := \cup\{P_{\eta_\beta}^M : \beta < \alpha \text{ and } \eta_\beta \in \mathcal{B}_\varepsilon\}$
 (d) $\text{Rang}(\mathbf{c}_\alpha^\varepsilon) \subseteq \kappa$
 (e) if $a, b \in \text{Dom}(\mathbf{c}_\alpha)$ and $\{a, b\} \in \text{edge}(G_M)$ then $\mathbf{c}_\alpha(a) \neq \mathbf{c}_\alpha(b)$.

Clearly this suffices. Why is this possible?

If $\alpha = 0$ let $\mathbf{c}_\alpha^\varepsilon$ be empty, if α is a limit ordinal let $\mathbf{c}_\alpha^\varepsilon = \cup\{\mathbf{c}_\beta^\varepsilon : \beta < \alpha\}$ and if $\alpha = \beta + 1 \wedge \alpha(\beta) \neq \varepsilon$ let $\mathbf{c}_\alpha = \mathbf{c}_\beta$.

Lastly, if $\alpha = \beta + 1 \wedge h(\beta) = \varepsilon$ we define $\mathbf{c}_\alpha^\varepsilon$ as follows for $a \in \text{Dom}(\mathbf{c}_\alpha^\varepsilon)$, $\mathbf{c}_\alpha^\varepsilon(a)$ is:

Case 1: $a \in B_\beta^\varepsilon$.

Then $\mathbf{c}_\alpha^\varepsilon(a) = \mathbf{c}_\beta^\varepsilon(a)$.

Case 2: $a \in B_\alpha^\varepsilon \setminus B_\beta^\varepsilon$.

Then $\mathbf{c}_\alpha^\varepsilon(a) = \min(\kappa \setminus \{\mathbf{c}_\beta^\varepsilon(F_\zeta^M(a)) : \zeta < \varepsilon \text{ and } F_\zeta^M(a) \in \text{Dom}(\mathbf{c}_\beta^\varepsilon)\})$.

This is well defined as:

- $\boxplus_{5.2}$ (a) $B_\alpha^\varepsilon = B_\beta^\varepsilon \cup P_{\eta_\beta}^M$
 (b) if $a \in B_\beta^\varepsilon$ then $\mathbf{c}_\beta^\varepsilon(a)$ is well defined (so case 1 is O.K.)
 (c) if $\{a, b\} \in \text{edge}(G_M)$, $a \in P_{\eta_\beta}^M$ and $b \in B_\alpha^\varepsilon$ then $b \in B_\beta^\varepsilon$ and $b \in \{F_\zeta^M(a) : \zeta < \varepsilon\}$
 (d) $\mathbf{c}_\alpha^\varepsilon(a)$ is well defined in Case 2, too
 (e) $\mathbf{c}_\alpha^\varepsilon$ is a function from B_α^ε to κ
 (f) $\mathbf{c}_\alpha^\varepsilon$ is a colouring.

[Why? Clause (a) by $\boxplus_{5.1}(c)$, clause (b) by the induction hypothesis and clause (c) by $\boxplus_1(c) + \boxplus_4$. Next, clause (d) holds as $\{\mathbf{c}_\beta^\varepsilon(F_\zeta^M(a)) : \zeta < \varepsilon \text{ and } F_\zeta^M(a) \in B_\beta^\varepsilon = \text{Dom}(\mathbf{c}_\beta^\varepsilon)\}$ is a set of cardinality $\leq |\varepsilon| < \kappa$. Clause (e) holds by the choices of the $\mathbf{c}_\alpha^\varepsilon(a)$'s. Lastly, to check that clause (f) holds assume $\{a, b\}$ is an edge of $G_M \upharpoonright B_\alpha^\varepsilon$, without loss of generality for some $\zeta < \kappa$ we have $b = F_\zeta^M(a)$, hence $\eta_a^M <_{\mathcal{A}} \eta_b^M$. If $a, b \in B_\beta^\varepsilon$ use the induction hypothesis. Otherwise, $\zeta < \varepsilon$ by the definition of “ h witnesses $\bar{\eta} \upharpoonright u$ is free” and the choice of B_α^ε in $\boxplus_{5.1}(c)$. Now use the choice of $\mathbf{c}_\alpha^\varepsilon(a)$ in Case 2 above.]

So indeed \boxplus_5 holds.]

\boxplus_6 $\text{chr}(G_M) > \kappa$ if $M \in K_{\mathcal{A}}^*$.

Why? Toward contradiction assume $\mathbf{c} : G_M \rightarrow \kappa$ is a colouring. For each $\eta \in \mathcal{A}$ and $\varepsilon < \kappa$ let $\Lambda_{\eta, \varepsilon} = \{\nu : \nu \in \mathcal{A}, \nu <_{\mathcal{A}} \eta, \nu(\varepsilon) = \eta(\varepsilon) \text{ and for some } a \in P_\nu^M \text{ we have } \mathbf{c}(a) = \varepsilon\}$.

Let $\mathcal{B}_\varepsilon = \{\eta \in \mathcal{A} : |\Lambda_{\eta, \varepsilon}| < \kappa\}$. Now if $\mathcal{A} \neq \cup\{\mathcal{B}_\varepsilon : \varepsilon < \kappa\}$ then pick any $\eta \in \mathcal{A} \setminus \cup\{\mathcal{B}_\varepsilon : \varepsilon < \kappa\}$ and by induction on $\varepsilon < \kappa$ choose $\nu_\varepsilon \in \Lambda_{\eta, \varepsilon} \setminus \{\nu_\zeta : \zeta < \varepsilon\}$, possible as $\eta \notin \mathcal{B}_\varepsilon$ by the definition of \mathcal{B}_ε . By the definition of $\Lambda_{\eta, \varepsilon}$ there is $a_\varepsilon \in P_{\nu_\varepsilon}^M$ such that $\mathbf{c}(a_\varepsilon) = \varepsilon$. So as $M \in K_{\mathcal{A}}^*$ there is $a \in P_\eta^M$ such that $\varepsilon < \kappa \Rightarrow F_\varepsilon^M(a) = a_\varepsilon$, but $\{a, a_\varepsilon\} \in \text{edge}(G_M)$ hence $\mathbf{c}(a) \neq \mathbf{c}(a_\varepsilon) = \varepsilon$ for every $\varepsilon < \kappa$, contradiction. So $\mathcal{A} = \cup\{\mathcal{B}_\varepsilon : \varepsilon < \kappa\}$.

For each $\varepsilon < \kappa$ we choose $\zeta_\eta < \kappa$ for $\eta \in \mathcal{B}_\varepsilon$ by induction on $<_{\mathcal{A}}$ such that $\zeta_\eta \notin \{\zeta_\nu : \nu \in \Lambda_{\eta, \varepsilon} \cap \mathcal{B}_\varepsilon\}$. Let $\mathcal{B}_{\varepsilon, \zeta} = \{\eta \in \mathcal{B}_\varepsilon : \zeta_\eta = \zeta\}$ for $\varepsilon, \zeta < \kappa$ so $\mathcal{A} = \cup\{\mathcal{B}_{\varepsilon, \zeta} : \varepsilon, \zeta < \kappa\}$ and clearly $\eta \mapsto \eta(\varepsilon)$ is a one-to-one function with domain $\mathcal{B}_{\varepsilon, \zeta}$, contradiction to “ $\bar{\eta} = \bar{\eta} \upharpoonright u_\lambda$ is not weakly free”. $\square_{2.2}$

Observation 2.3. 1) If $\mathcal{A} \subseteq \kappa^\mu$ and $\eta \neq \nu \in \mathcal{A} \Rightarrow (\forall^\infty \varepsilon < \kappa)(\eta(\varepsilon) \neq \nu(\varepsilon))$ then \mathcal{A} is free iff \mathcal{A} is weakly free.

2) The assumptions of 2.2(2) hold when: $\mu \geq \lambda > \kappa$ are regular, $S \subseteq S_\kappa^\mu$ stationary, $\bar{\eta} = \langle \eta_\delta : \delta \in S \rangle$, η_δ an increasing sequence of ordinals of length κ with limit δ such that $u \subseteq [\lambda]^{<\lambda} \Rightarrow \langle \text{Rang}(\eta_\delta) : \eta \in u \rangle$ has a one-to-one choice function.

Conclusion 2.4. Assume that for every graph G , if $H \subseteq G \wedge |H| < \lambda \Rightarrow \text{chr}(H) \leq \kappa$ then $\text{chr}(G) \leq \kappa$.

Then:

- (A) if $\mu > \kappa = \text{cf}(\mu)$ and $\mu \geq \lambda$ then $\text{pp}(\mu) = \mu^+$
- (B) if $\mu > \text{cf}(\mu) \geq \kappa$ and $\mu \geq \lambda$ then $\text{pp}(\mu) = \mu^+$, i.e. the strong hypothesis
- (C) if $\kappa = \aleph_0$ then above λ the SCH holds.

Proof. Clause (A): By 2.2 and [?, Ch.II], [?, Ch.IX, §1].

Clause (B): Follows from (A) by [?, Ch.VIII, §1].

Clause (C): Follows from (B) by [?, Ch.IX, §1]. $\square_{2.4}$

EINSTEIN INSTITUTE OF MATHEMATICS, EDMOND J. SAFRA CAMPUS, GIVAT RAM, THE HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM, 91904, ISRAEL, AND, DEPARTMENT OF MATHEMATICS, HILL CENTER - BUSCH CAMPUS, RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY, 110 FRELINGHUYSEN ROAD, PISCATAWAY, NJ 08854-8019 USA

Email address: shelah@math.huji.ac.il

URL: <http://shelah.logic.at>