

CICHOŃ'S MAXIMUM

MARTIN GOLDSTERN, JAKOB KELLNER, AND SAHARON SHELAH

ABSTRACT. Assuming four strongly compact cardinals, it is consistent that all entries in Cichoń's diagram (apart from $\text{add}(\mathcal{M})$ and $\text{cof}(\mathcal{M})$, whose values are determined by the others) are pairwise different; more specifically that

$$\aleph_1 < \text{add}(\mathcal{N}) < \text{cov}(\mathcal{N}) < \mathfrak{b} < \text{non}(\mathcal{M}) < \text{cov}(\mathcal{M}) < \mathfrak{d} < \text{non}(\mathcal{N}) < \text{cof}(\mathcal{N}) < 2^{\aleph_0}.$$

INTRODUCTION

Independence. How many Lebesgue null sets are required to cover the real line? Obviously countably many are not enough, as the countable union of null sets is null; and obviously continuum many are enough, as $\bigcup_{r \in \mathbb{R}} \{r\} = \mathbb{R}$.

The answer to our question is a cardinal number called $\text{cov}(\mathcal{N})$. As we have just seen,

$$\aleph_0 = |\mathbb{N}| < \text{cov}(\mathcal{N}) \leq |\mathbb{R}| = 2^{\aleph_0}.$$

In particular, if the Continuum Hypothesis (CH) holds (i.e., if there are no cardinalities strictly between $|\mathbb{N}|$ and $|\mathbb{R}|$, or equivalently: if $\aleph_1 = 2^{\aleph_0}$), then $\text{cov}(\mathcal{N}) = 2^{\aleph_0}$; but without CH, the answer could also be some cardinal less than 2^{\aleph_0} . According to Cohen's famous result [Coh63], CH is independent of the usual axiomatization of mathematics, the set theoretic axiom system ZFC. I.e., we can prove that the ZFC axioms neither imply CH nor imply $\neg\text{CH}$. For this result, Cohen introduced the method of forcing, which has been continuously expanded and refined ever since. Forcing also proves that the value of $\text{cov}(\mathcal{N})$ is independent. For example, $\text{cov}(\mathcal{N}) = \aleph_1 < 2^{\aleph_0}$ is consistent, as is $\aleph_1 < \text{cov}(\mathcal{N}) = 2^{\aleph_0}$.

Cichoń's diagram. $\text{cov}(\mathcal{N})$ is a so-called cardinal characteristic of the continuum. Other well-studied characteristics include the following:

- $\text{add}(\mathcal{N})$ is the smallest number of Lebesgue null sets whose union is not null.
- $\text{non}(\mathcal{N})$ is the smallest cardinality of a non-null set.
- $\text{cof}(\mathcal{N})$ is the smallest size of a cofinal family of null sets, i.e., a family that contains for each null set N a superset of N .
- Replacing “null” with “meager”, we can analogously define $\text{add}(\mathcal{M})$, $\text{non}(\mathcal{M})$, $\text{cov}(\mathcal{M})$, and $\text{cof}(\mathcal{M})$.
- In addition, we define \mathfrak{b} as the smallest size of an unbounded family, i.e., a family \mathcal{H} of functions from \mathbb{N} to \mathbb{N} such that for every $f : \mathbb{N} \rightarrow \mathbb{N}$ there is some $h \in \mathcal{H}$ which is not almost everywhere bounded by f .

Equivalently, $\mathfrak{b} = \text{add}(\mathcal{K}) = \text{non}(\mathcal{K})$, where \mathcal{K} is the σ -ideal generated by the compact subsets of the irrationals.

Date: October 5, 2020.


2010 Mathematics Subject Classification. 03E17.

Key words and phrases. Set theory of the reals, Cichoń's diagram, Forcing, Compact cardinals.

Supported by Austrian Science Fund (FWF): P29575 & I3081 and National Science Foundation NSF DMS-1362974 (first author), Austrian Science Fund (FWF): P26737 & P30666 (second author) European Research Council grant ERC-2013-ADG 338821 (third author). This is publication number 1122 of the third author.

- And \mathfrak{d} is the smallest size of a dominating family, i.e., a family \mathcal{H} such that for every $f : \mathbb{N} \rightarrow \mathbb{N}$ there is some $h \in \mathcal{H}$ such that $(\exists n \in \mathbb{N})(\forall m > n) h(m) > f(m)$.
Equivalently, $\mathfrak{d} = \text{cov}(\mathcal{K}) = \text{cof}(\mathcal{K})$.
- For the ideal ctbl of countable sets, we trivially get $\text{add}(\text{ctbl}) = \text{non}(\text{ctbl}) = \aleph_1$ and $\text{cov}(\text{ctbl}) = \text{cof}(\text{ctbl}) = 2^{\aleph_0}$.

The characteristics we have mentioned so far,¹ and the basic relations between them, can be summarized in Cichoń’s diagram:



An arrow from \mathfrak{x} to \mathfrak{y} indicates that ZFC proves $\mathfrak{x} \leq \mathfrak{y}$. Moreover, $\max(\mathfrak{d}, \text{non}(\mathcal{M})) = \text{cof}(\mathcal{M})$ and $\min(\mathfrak{b}, \text{cov}(\mathcal{M})) = \text{add}(\mathcal{M})$. A (by now) classical series of theorems [Bar84, BJS93, CKP85, JS90, Kam89, Mil81, Mil84, RS83, RS85] proves these (in)equalities in ZFC and shows that they are the only ones provable. More precisely, all assignments of the values \aleph_1 and \aleph_2 to the characteristics in Cichoń’s Diagram are consistent with ZFC, provided they do not contradict the above (in)equalities. (A complete proof can be found in [BJ95, ch. 7].)

Note that Cichoń’s diagram shows a fundamental asymmetry between the ideals of Lebesgue null sets and of meager sets (we will mention another one in the context of large cardinals). Any such asymmetry is hidden if we assume CH, as under CH not only all the characteristics are \aleph_1 , but even the Erdős-Sierpiński Duality Theorem holds [Oxt80, ch. 19]: There is an involution $f : \mathbb{R} \rightarrow \mathbb{R}$ (i.e., a bijection such that $f \circ f = \text{Id}$) such that $A \subseteq \mathbb{R}$ is meager iff $f''A$ is null.

So it is settled which assignments of \aleph_1 and \aleph_2 to Cichoń’s diagram are consistent. It is more challenging to show that the diagram can contain more than two different cardinal values. For recent progress in this direction see, e.g., [Mej13, GMS16, FGKS17, KTT18].

The result of this paper is in some respect the strongest possible, as we show that consistently *all* the entries are pairwise different (apart from the two equalities provable in ZFC mentioned above). Of course one can ask more; see the questions in Section 4. In particular, we use large cardinals in the proof.

Large cardinals. As mentioned, ZFC is an axiom system for the whole of mathematics. A much “weaker” axiom system (for the natural numbers) is PA (Peano arithmetic).

Gödel’s Incompleteness Theorem shows that a theory such as PA or ZFC can never prove its own consistency. On the other hand, it is trivial to show in ZFC that PA is consistent (as in ZFC we can construct \mathbb{N} and prove that it satisfies PA). We can say that ZFC has a higher consistency strength than PA.

One axiom of ZFC is INF, the statement “there is an infinite cardinal”. If we remove INF from ZFC, we end up with a theory ZFC^0 that can still describe concrete hereditarily finite objects and can be interpreted (admittedly in a not very natural way) as a weak version of

¹There are many other cardinal characteristics, see for example [Bla10], but the ones in Cichoń’s diagram seem to be considered to be the most important ones.

PA which has the same consistency strength as PA.² So we can say that adding an infinite cardinal to ZFC⁰ increases the consistency strength.

There are notions of cardinal numbers much “stronger” than just “infinite”. Often, such large cardinal assumptions (abbreviated LC in the following) have the following form:

There is a cardinal $\kappa > \aleph_0$ that behaves towards the smaller cardinals in a similar way as \aleph_0 behaves to finite numbers.

A forcing proof shows, e.g.,

If ZFC is consistent, then ZFC+¬CH is consistent,

and this implication can be proved in a very weak system such as PA. However, we cannot prove (not even in ZFC) for any large cardinal

“if ZFC is consistent, then ZFC+LC is consistent”;

because in ZFC+LC we can prove the consistency of ZFC. We say: LC has a higher consistency strength than ZFC.

An instance of a large cardinal (in fact a very weak one, a so-called inaccessible cardinal), appears in another striking example of the asymmetry between measure and category: The following statement is equiconsistent with an inaccessible cardinal [Sol70, She84]:

All projective³ sets of reals are Lebesgue measurable.

In contrast, according to [She84] no large cardinal assumption is required to show the consistency of

All projective sets of reals have the property of Baire.

So we can assume “for free” that all (reasonable) sets have the Baire property, whereas we have to provide additional consistency strength for Lebesgue measurability.

In the case of our paper, we require (the consistency of) the existence of four compact cardinals to prove our main result. It seems unlikely that any large cardinals are actually required; but a proof without them would probably be considerably more complicated. It is not unheard of that ZFC results first have (simpler) proofs using large cardinal assumptions; an example can be found in [She04].

Annotated Contents. From now on, we assume that the reader is familiar with some basic properties of the characteristics defined above, as well as with the associated forcing notions Cohen, amoeba, random, Hechler and eventually different, all of which can be found, e.g., in [BJ95].

This paper consists of three parts:

In Section 1, we present a finite support ccc iteration \mathbb{P}^5 forcing that $\aleph_1 < \text{add}(\mathcal{N}) < \text{cov}(\mathcal{N}) < \mathfrak{b} < \text{non}(\mathcal{M}) < \text{cov}(\mathcal{M}) = 2^{\aleph_0}$. This result is not new: Such a forcing was introduced in [GMS16], and we follow this construction quite closely. However, we need GCH in the ground model, whereas [GMS16] requires $2^\chi \gg \lambda$ for some $\chi < \lambda$. Also, we describe how the inequalities are “strongly witnessed”, see Definitions 1.8 and 1.15.

In Section 2, we show how to construct (under GCH) for κ strongly compact and $\theta > \kappa$ regular a “BUP-embedding” from κ to θ , i.e., an elementary embedding $j : V \rightarrow M$ with critical point κ and $\text{cf}(j(\kappa)) = |j(\kappa)| = \theta$ such that M is transitive and $<\kappa$ -closed and

²More concretely, $\text{ZF}_{\text{fin}} := \text{ZFC}^0 + \neg\text{INF}$ can be seen to be “equivalent” to PA (i.e., mutually interpretable); this goes back to Ackermann [Ack37], see the survey [KW07].

³This is the smallest family containing the Borel sets and closed under continuous images, complements, and countable unions. In practice, all sets used in mathematics that are defined without using AC are projective. Alternatively we could use the statement: “ZF (without the Axiom of choice) holds and all sets of reals are Lebesgue measurable.”




FIGURE 1. Our cardinal configuration (the λ_i are increasing).

such that $j''S$ is cofinal in $j(S)$ for every $\leq \kappa$ -directed partial order S . For a ccc forcing P we investigate $j(P)$ and show that $j(P)$ forces the same values to some characteristics in Cichoń’s diagram as P and different values to others, in a very controlled way; assuming that there were “strong witnesses” for P forcing the initial values, as described in Section 1.

Section 3 contains the main result of this paper: Assuming four strongly compact cardinals, we let k be the composition of four such BUP-embeddings, mapping \mathbb{P}^5 to a ccc forcing \mathbb{P}^9 . We then show that \mathbb{P}^9 forces

$$\aleph_1 < \text{add}(\mathcal{N}) < \text{cov}(\mathcal{N}) < \mathfrak{b} < \text{non}(\mathcal{M}) < \text{cov}(\mathcal{M}) < \mathfrak{d} < \text{non}(\mathcal{N}) < \text{cof}(\mathcal{N}) < 2^{\aleph_0},$$

i.e., we get for increasing cardinals λ_i the constellation of Figure 1.

Boolean ultrapowers as used in this paper were investigated by Mansfield [Man71] and recently applied e.g. by the third author with Malliaris [MS16] and with Raghavan [RS], where Boolean ultrapowers of forcing notions are used to force specific values to certain cardinal characteristics. Recently the third author developed a method of using Boolean ultrapowers to control characteristics in Cichoń’s diagram. A first (and simpler) application of these methods is given in [KTT18].

We mention some open questions in Section 4.

Acknowledgments. We thank three anonymous referees for pointing out several unclarities and typos; and Moti Gitik and Diego Mejía for suggestions to improve the presentation.

1. THE INITIAL FORCING

1.1. Good iterations and the LCU property. We want to show that some forcing \mathbb{P}^5 results in $\mathfrak{x} = \lambda_i$ for certain characteristics \mathfrak{x} . So we have to show two “directions”, $\mathfrak{x} \leq \lambda_i$ and $\mathfrak{x} \geq \lambda_i$. For most of the characteristics, one direction will use the fact that \mathbb{P}^5 is “good”; a notion introduced by Judah and the third author [JS90] and Brendle [Bre91]. We now recall the basic facts of good iterations, and specify the instances of the relations we use.

Assumption 1.1. *We will consider binary relations R on $X = \omega^\omega$ (or on $X = 2^\omega$) that satisfy the following: There are relations R^k such that $R = \bigcup_{k \in \omega} R^k$, each R^k is a closed subset (and in fact absolutely defined) of $X \times X$, and for $g \in X$ and $k \in \omega$, the set $\{f \in X : f R^k g\}$ is nowhere dense (and of course closed). Also, for all $g \in X$ there is some $f \in X$ with $f R g$.*

We will actually use another space as well, the space C of strictly positive rational sequences $(q_n)_{n \in \omega}$ such that $\sum_{n \in \omega} q_n \leq 1$. It is easy to see that C is homeomorphic to ω^ω , when we equip the rationals with the discrete topology and use the product topology. Let us fix one such (absolutely defined) homeomorphism.

We use the following instances of relations R on X ; it is easy to see that they all satisfy the assumption (for $X_1 = C$ we use the homeomorphism mentioned above):

Definition 1.2. 1. $X_1 = C$: $f R_1 g$ if $(\forall^* n \in \omega) f(n) \leq g(n)$.

(We use $\forall^* n$ as abbreviation for $(\exists n_0)(\forall n > n_0)$.)

2. Fix a partition $(I_n)_{n \in \omega}$ of ω with $|I_n| = 2^{n+1}$.
 $X_2 = 2^\omega$: $f R_2 g$ if $(\forall^* n \in \omega) f \upharpoonright I_n \neq g \upharpoonright I_n$.
3. $X_3 = \omega^\omega$: $f R_3 g$ if $(\forall^* n \in \omega) f(n) \leq g(n)$.
4. $X_4 = \omega^\omega$: $f R_4 g$ if $(\forall^* n \in \omega) f(n) \neq g(n)$.

Note that Assumption 1.1 is satisfied, witnessed by the relations R_i^k defined by replacing $(\forall^* n \in \omega)$ with $(\forall n \geq k)$.

We say “ f is bounded by g ” if $f R g$; and, for $\mathcal{Y} \subseteq \omega^\omega$, “ f is bounded by \mathcal{Y} ” if $(\exists y \in \mathcal{Y}) f R y$. We say “unbounded” for “not bounded”. (I.e., f is unbounded by \mathcal{Y} if $(\forall y \in \mathcal{Y}) \neg f R y$.) We call \mathcal{X} an R-unbounded family, if $\neg(\exists g)(\forall x \in \mathcal{X}) x R g$, and an R-dominating family if $(\forall f)(\exists x \in \mathcal{X}) f R x$.

- Let \mathfrak{b}_i be the minimal size of an R_i -unbounded family,
- and let \mathfrak{d}_i be the minimal size of an R_i -dominating family.

We only need the following connections between R_i and the cardinal characteristics:

- Lemma 1.3.** 1. $\text{add}(\mathcal{N}) = \mathfrak{b}_1$ and $\text{cof}(\mathcal{N}) = \mathfrak{d}_1$.
 2. $\text{cov}(\mathcal{N}) \leq \mathfrak{b}_2$ and $\text{non}(\mathcal{N}) \geq \mathfrak{d}_2$.
 3. $\mathfrak{b} = \mathfrak{b}_3$ and $\mathfrak{d} = \mathfrak{d}_3$.
 4. $\text{non}(\mathcal{M}) = \mathfrak{b}_4$ and $\text{cov}(\mathcal{M}) = \mathfrak{d}_4$.

Proof. (3) holds by definition. (1) can be found in [BJ95, 6.5.B]. (4) is a result of [Mil82, Bar87], cf. [BJ95, 2.4.1 and 2.4.7].

To prove (2), note that for fixed $f \in 2^\omega$ the set $\{g \in 2^\omega : \neg f R_2 g\}$ is a null set, call it N_f . Let \mathcal{F} be an R_2 -unbounded family. Then $\{N_f : f \in \mathcal{F}\}$ covers 2^ω : Fix $g \in 2^\omega$. As g does not bound \mathcal{F} , there is some $f \in \mathcal{F}$ unbounded by g , i.e., $g \in N_f$. Let X be a non-null set. Then X is R_2 -dominating: For any $f \in 2^\omega$ there is some $x \in X \setminus N_f$, i.e., $f R_2 x$. \square

We will also use:

Lemma 1.4. [BJ95] Amoeba forcing \mathbb{A} adds a dominating element \bar{b} of C , i.e., $\mathbb{A} \Vdash \bar{q} R_1 \bar{b}$ for all $\bar{q} \in C \cap V$.

Proof. Let us define a slalom S to be a function $S : \omega \rightarrow [\omega]^{<\omega}$ such that $|S(n)| > 0$ and $\sum_{n=1}^{\infty} \frac{|S(n)|}{n^2} < \infty$.

Amoeba forcing will add a null set covering all old null sets, and therefore (according to [BJ95, 2.3.3]) a slalom S covering all old slaloms. Set $a_n := \frac{|S(n)|}{n^2}$, $M := \sum_{n=1}^{\infty} a_n$, set M' the smallest natural number $\geq M$, and set $b_n := \frac{a_{n+1}}{M'}$. Then it is easy to see that $(b_n)_{n \in \omega} \in C$ dominates every old sequence $(q_n)_{n \in \omega}$ in C . \square

Definition 1.5. [JS90] Let P be a ccc forcing, λ an uncountable regular cardinal, and R as above. P is (R, λ) -good, if for each P -name $r \in \omega^\omega$ there is (in V) a nonempty set $\mathcal{Y} \subseteq \omega^\omega$ of size $< \lambda$ such that every f (in V) that is R -unbounded by \mathcal{Y} is forced to be R -unbounded by r as well.

Note that λ -good trivially implies μ -good if $\mu \geq \lambda$ are regular.

How do we get good forcings? Let us just note the following results:

Lemma 1.6. A finite support (henceforth abbreviated FS) iteration of Cohen forcing is good for any (R, λ) , and the composition of two (R, λ) -good forcings is (R, λ) -good.

Assume that $(P_\alpha, Q_\alpha)_{\alpha < \delta}$ is a FS ccc iteration. Then P_δ is (R, λ) -good, if each Q_α is forced to satisfy the following:

1. For $R = R_1$: $|Q_\alpha| < \lambda$, or Q_α is σ -centered, or Q_α is a sub-Boolean-algebra of the random algebra.
2. For $R = R_2$: $|Q_\alpha| < \lambda$, or Q_α is σ -centered.
4. For $R = R_4$: $|Q_\alpha| < \lambda$.

(Remark: For R_3 the same holds as for R_4 , which however is of no use for our construction.)

Proof. (R, λ) -goodness is preserved by FS ccc iterations (in particular compositions), as proved in [JS90], cf. [BJ95, 6.4.11–12]. Also, ccc forcings of size $< \lambda$ are (R, λ) -good [BJ95, 6.4.7]; which takes care of the case $|Q_\alpha| < \lambda$ (and in particular of Cohen forcing). So it remains to show that (for $i = 1, 2$) the “large” iterands in the list are (R_i, λ) -good.

For R_1 this follows from [JS90] and [Kam89], cf. [BJ95, 6.5.17–18].

For R_2 , this is proven in [Bre91], and as the proof is very short, we give it here: Write Q_α as union $\bigcup_{k \in \omega} Q^k$ of centered sets. Given the Q_α -name r , pick a countable elementary submodel N containing r and Q_α , and set $\mathcal{Y} = N \cap 2^\omega$. Assume towards a contradiction that f is unbounded by \mathcal{Y} , but is forced by p_0 to be bounded by r , i.e., p_0 forces $(\forall n > n_0) f \upharpoonright I_n \neq r \upharpoonright I_n$. Now p_0 may not be in N , but there is some $k_0 \in \omega$ such that $p_0 \in Q^{k_0}$. In N , we can pick for each $n \in \omega$ some $s_n \in 2^{I_n}$ such that no $q \in Q^{k_0}$ forces $r \upharpoonright I_n \neq s_n$. (There are only finitely many $s \in 2^{I_n}$; if each s is forbidden by some q , then the common stronger element would prevent all possibilities for $r \upharpoonright I_n$.) So in N , we get some $g \in 2^\omega$ such that $g \upharpoonright I_n = s_n$. As f is unbounded by \mathcal{Y} (or equivalently: by N), there is some $n > n_0$ such that $f \upharpoonright I_n = g \upharpoonright I_n = s_n$, which implies that p_0 (as an element of Q^{k_0}) does not force $r \upharpoonright I_n \neq f \upharpoonright I_n$, a contradiction. \square

Lemma 1.7. *Let $\lambda \leq \kappa \leq \mu$ be uncountable regular cardinals. After forcing with μ many Cohen reals $(c_\alpha)_{\alpha \in \mu}$, followed by an (R, λ) -good forcing, we get: For every real r in the final extension, the set $\{\alpha \in \kappa : c_\alpha \text{ is unbounded by } r\}$ is cobounded in κ . I.e., $(\exists \alpha \in \kappa) (\forall \beta \in \kappa \setminus \alpha) \neg c_\beta R r$.*

(The Cohen real c_β can be interpreted both as Cohen generic element of 2^ω and as Cohen generic element of ω^ω ; we use the interpretation suitable for the relation R .)

Proof. Work in the intermediate extension after κ many Cohen reals; let us call it V_κ . The remaining forcing (i.e., $\mu \setminus \kappa$ many Cohens composed with the good forcing) is good; so applying the definition we get (in V_κ) a set \mathcal{Y} of size $< \lambda$.

As the initial Cohen extension is ccc, and $\kappa \geq \lambda$ is regular, we get some $\alpha \in \kappa$ such that each element y of \mathcal{Y} already exists in the extension by the first α many Cohens, call it V_α . The set of reals M_y bounded by y is meager (and absolute). Any c_β for $\beta \in \kappa \setminus \alpha$ is Cohen over V_α , and therefore not in M_y , i.e., not bounded by y , i.e., not by \mathcal{Y} . So according to the definition of good, each such c_β is unbounded by r as well, for the given r . \square

In the light of this result, let us revisit Lemma 1.3 with some new notation, the “linearly cofinally unbounded” property LCU:

Definition 1.8. For $i = 1, 2, 3, 4$, γ a limit ordinal, and P a ccc forcing notion, let $\text{LCU}_i(P, \gamma)$ stand for:

There is a sequence $(x_\alpha)_{\alpha \in \gamma}$ of P -names of elements of X_i (the domain of the relation R_i) such that for every such P -name y
 $(\exists \alpha \in \gamma) (\forall \beta \in \gamma \setminus \alpha) P \Vdash \neg x_\beta R_i y$.

Lemma 1.9. \bullet $\text{LCU}_i(P, \delta)$ is equivalent to $\text{LCU}_i(P, \text{cf}(\delta))$.

- If λ is regular, then $\text{LCU}_i(P, \lambda)$ implies $\mathfrak{b}_i \leq \lambda$ and $\mathfrak{d}_i \geq \lambda$.

In particular:

1. $\text{LCU}_1(P, \lambda)$ implies $P \Vdash (\text{add}(\mathcal{N}) \leq \lambda \ \& \ \text{cof}(\mathcal{N}) \geq \lambda)$.
2. $\text{LCU}_2(P, \lambda)$ implies $P \Vdash (\text{cov}(\mathcal{N}) \leq \lambda \ \& \ \text{non}(\mathcal{N}) \geq \lambda)$.
3. $\text{LCU}_3(P, \lambda)$ implies $P \Vdash (\mathfrak{b} \leq \lambda \ \& \ \mathfrak{d} \geq \lambda)$.
4. $\text{LCU}_4(P, \lambda)$ implies $P \Vdash (\text{non}(\mathcal{M}) \leq \lambda \ \& \ \text{cov}(\mathcal{M}) \geq \lambda)$.

Proof. Assume that $(\alpha_\beta)_{\beta \in \text{cf}(\delta)}$ is increasing continuous and cofinal in δ . If $(x_\alpha)_{\alpha \in \delta}$ witnesses $\text{LCU}_i(P, \delta)$, then $(x_{\alpha_\beta})_{\beta \in \text{cf}(\delta)}$ witnesses $\text{LCU}_i(P, \text{cf}(\delta))$. And if $(x_\beta)_{\beta \in \text{cf}(\delta)}$ witnesses $\text{LCU}_i(P, \text{cf}(\delta))$, then $(y_\alpha)_{\alpha \in \delta}$ witnesses $\text{LCU}_i(P, \delta)$, where $y_\alpha := x_\beta$ for $\alpha \in [\alpha_\beta, \alpha_{\beta+1})$.

The set $\{x_\alpha : \alpha \in \lambda\}$ is certainly forced to be \mathbb{R}_i -unbounded; and given a set $Y = \{y_j : j < \theta\}$ of $\theta < \lambda$ many P -names, each has a bound $\alpha_j \in \lambda$ so that $(\forall \beta \in \lambda \setminus \alpha_j) P \Vdash \neg x_\beta \mathbb{R}_i y_j$, so for any $\beta \in \lambda$ above all α_j we get $P \Vdash \neg x_\beta \mathbb{R}_i y_j$ for all j ; i.e., Y cannot be dominating. \square

Remark 1.10. $\mathfrak{b}_i \leq \lambda$ is equivalent to the existence of a sequence $(x_\alpha : \alpha \in \lambda)$ with the property $(\forall y) (\exists \alpha) \neg (x_\alpha \mathbb{R}_i y)$; such a sequence might be called a “witness” for $\mathfrak{b}_i \leq \lambda$. In LCU we demand a stronger property; a sequence $(x_\alpha : \alpha < \lambda)$ with this stronger property could informally be called a “strong witness” for $\mathfrak{b}_i \leq \lambda$. Similarly, the next subsection introduces a different notion, COB, corresponding to “strong witnesses” for $\mathfrak{d}_i \leq \mu$.

1.2. The initial forcing \mathbb{P}^5 : Partial forcings and the COB property. Assume we have a forcing iteration $(P_\beta, Q_\beta)_{\beta < \alpha}$ with limit P_α , where each Q_β is forced by P_β to be a set of reals such that the generic filter of Q_β is determined (in a Borel way)⁴ from some generic real η_β . Fix some $w \subseteq \alpha$. We define the P_α -name Q_α to consist of all random forcing conditions that can be Borel-calculated from generics at w alone.

More explicitly:

Definition 1.11. (1) q is in Q_α if there are in the ground model V a countable subset $u \subseteq w$ and a Borel function $B : \mathbb{R}^u \rightarrow \mathbb{R}$ such that $q = B((\eta_\beta)_{\beta \in u})$ is a random condition.

Being a random condition is a Borel property (if we fix some suitable representation of random forcing). Accordingly, we can restrict ourselves to the case that B is a Borel function whose image consists of random conditions only.

- (2) We call a pair (B, u) as above “a w -groundmodel-code” or just “code”. Note that this code is a ground model object. Q_α consists exactly of the evaluations of such codes.
- (3) We call a condition $(p, q) \in P_\alpha * Q_\alpha$ “determined at position α ”, if there is a code (B, u) such that p forces that (B, u) is a code for q . (Note that generally we only have a P_α -name for a code.) Given some (p, q) , we can obviously find $p' \leq p$ such that (p', q) is determined at α .
- (4) We will later also consider so-called “groundmodel-code-sequences” for elements of Q_α , that is (in V) a sequence $(B_n, u_n)_{n \in \omega}$ of codes, where u_n is in w_α . Of course not every ω -sequence of Q_α -conditions in the P_α -extension is described by a ground model sequence. (In particular, there will only be few ground model sequences, but many new ω -sequences in the extension.)

⁴More specifically, we require that the Borel function for Q_β is already fixed in the ground model. For example, assume Q_β is random forcing, defined as the set of all positive pruned trees T , i.e., trees $T \subseteq 2^{<\omega}$ without leaves such that $[T]$ has positive measure. Then the generic filter G for this forcing is determined by the generic real η (the random real), and G consists of those trees T such that $\eta \in [T]$, which is a Borel relation. See [KTT18, Sec. 1.2] for a formal definition and more details.

Clearly, in the P_α extension, Q_α is a subforcing (not necessarily a complete one) of the full random forcing, and if p, q in Q_α are incompatible in Q_α then they are incompatible in random forcing. (Two compatible conditions p, q have a canonical conjunction $p \wedge q$ (the intersection), and if p and q are both Borel-calculated from w , then so is the intersection.) In particular Q_α is ccc.

We call this forcing “partial random forcing defined from w ”. Analogously, we define the “partial Hechler”, “partial eventually different”⁵ and “partial amoeba” forcings (and the same argument shows that these forcings are also ccc).

Assume that λ is regular uncountable and that $\mu < \lambda$ implies $\mu^{\aleph_0} < \lambda$. Then $|w| < \lambda$ implies that the sizes of the partial forcings defined by w are $< \lambda$.

We will assume the following throughout the paper:

Assumption 1.12. $\aleph_1 < \lambda_1 < \lambda_2 < \lambda_3 < \lambda_4 < \lambda_5$ are regular cardinals such that $\mu < \lambda_i$ implies $\mu^{\aleph_0} < \lambda_i$. Furthermore, λ_3 is the successor of a regular cardinal χ with $\chi^{\aleph_0} = \chi$, and $\lambda_5^{< \lambda_4} = \lambda_5$.

We set $\delta_5 = \lambda_5 + \lambda_5$, and partition $\delta_5 \setminus \lambda_5$ into unbounded sets S^1, S^2, S^3 and S^4 . Fix for each $\alpha \in \delta_5 \setminus \lambda_5$ some $w_\alpha \subseteq \alpha$ such that each $\{w_\alpha : \alpha \in S^i\}$ is cofinal⁶ in $[\delta_5]^{< \lambda_i}$.

The reader can assume that $(\lambda_i)_{i=1, \dots, 5}, (S^i)_{i=1, \dots, 4}$ as well as $(w_\alpha)_{\alpha \in S^i}$ for $i = 1, 2, 3$ have been fixed once and for all (let us call them “fixed parameters”), whereas we will investigate various possibilities for $\bar{w} = (w_\alpha)_{\alpha \in S^4}$ in the following Subsections 1.3 and 1.4. (We will call such a \bar{w} that satisfies the assumption a “cofinal parameter”).

Definition 1.13. Let $\mathbb{P}^5 = (P_\alpha, Q_\alpha)_{\alpha \in \delta_5}$ be the FS iteration where Q_α is Cohen forcing for $\alpha \in \lambda_5$, and

$$Q_\alpha \text{ is the partial } \left\{ \begin{array}{c} \text{amoeba} \\ \text{random} \\ \text{Hechler} \\ \text{eventually different} \end{array} \right\} \text{ forcing defined from } w_\alpha \text{ if } \alpha \text{ is in } \left\{ \begin{array}{c} S^1 \\ S^2 \\ S^3 \\ S^4 \end{array} \right.$$

According to Lemma 1.6 \mathbb{P}^5 is (λ_i, R_i) -good for $i = 1, 2, 4$, so Lemmas 1.7 and 1.9 give us:

Lemma 1.14. $\text{LCU}_i(\mathbb{P}^5, \kappa)$ holds for $i = 1, 2, 4$ and each regular cardinal κ in $[\lambda_i, \lambda_5]$.

So in particular, \mathbb{P}^5 forces $\text{add}(\mathcal{N}) \leq \lambda_1$, $\text{cov}(\mathcal{N}) \leq \lambda_2$, $\text{non}(\mathcal{M}) \leq \lambda_4$ and $\text{cov}(\mathcal{M}) = \text{non}(\mathcal{N}) = \text{cof}(\mathcal{N}) = \lambda_5 = 2^{\aleph_0}$; i.e., the respective characteristics in the left half of Cichoń’s diagram are small enough. It is easy to see that they are also large enough:

For example, the partial amoebas and the fact that $(w_\alpha)_{\alpha \in S^1}$ is cofinal ensure that \mathbb{P}^5 forces $\text{add}(\mathcal{N}) \geq \lambda_1$: Let $(N_k)_{k \in \mu}$, $\aleph_1 \leq \mu < \lambda_1$ be a family of \mathbb{P}^5 -names of null sets. Each N_k is a Borel-code, i.e., a real, i.e., a sequence of natural numbers, each of which is decided by a maximal antichain (labeled with natural numbers). Each condition in such an antichain has finite support, hence only uses finitely many coordinates in δ_5 . So all in all we get a set w^* of size $\leq \mu$ that already decides all N_k . (I.e., for each $k \in \mu$ there are a Borel function B in V and a sequence $(\alpha_j)_{j \in \omega}$ in V of elements of w^* such that $N_k = B(\eta_{\alpha_0}, \eta_{\alpha_1}, \dots)$.) There is some $\beta \in S^1$ such that $w_\beta \supseteq w^*$, and the partial amoeba forcing at β sees all the null sets N_k and therefore covers their union.

⁵See 1.22 for the definition.

⁶i.e., if $\alpha \in S^i$ then $|w_\alpha| < \lambda_i$, and for all $u \subseteq \delta_5$, $|u| < \lambda_i$ there is some $\alpha \in S^i$ with $w_\alpha \supseteq u$.

We will reformulate this in a slightly cumbersome manner that can be conveniently used later on, using the “cone of bounds” property COB:

Definition 1.15. For a ccc forcing notion P , regular uncountable cardinals λ, μ and $i = 1, 3, 4$, let $\text{COB}_i(P, \lambda, \mu)$ stand for:

There are a $<\lambda$ -directed partial order $(S, <)$ of size μ and a sequence $(g_s)_{s \in S}$ of P -names for reals such that for each P -name f of a real $(\exists s \in S)(\forall t > s) P \Vdash f R_i g_t$.

So s is the tip of a cone that consists of elements bounding f .

Lemma 1.16. For $i = 1, 3, 4$, $\text{COB}_i(P, \lambda, \mu)$ implies $P \Vdash (\mathfrak{b}_i \geq \lambda \ \& \ \mathfrak{d}_i \leq \mu)$.

Proof. $\mathfrak{d}_i \leq \mu$, as the set $(g_s)_{s \in S}$ is a dominating family of size μ . To show $\mathfrak{b}_i \geq \lambda$, assume $(f_\alpha)_{\alpha \in \theta}$ is a sequence of P -names of length $\theta < \lambda$. For each f_α there is a cone of upper bounds with tip $s_\alpha \in S$, i.e., $(\forall t > s_\alpha) P \Vdash f_\alpha R_i g_t$. As S is $<\lambda$ -directed, there is some t above all tips s_α . Accordingly, $P \Vdash f_\alpha R_i g_t$ for all α , i.e., $\{f_\alpha : \alpha \in \theta\}$ is not unbounded. \square

So for example, $\text{COB}_1(P, \lambda, \mu)$ implies $\lambda_1 \leq \mathfrak{b}_1 = \text{add}(\mathcal{N})$, etc. The definition and lemma would work for $i = 2$ as well, but would not be useful⁷ as we do not have $\mathfrak{b}_2 \leq \text{cov}(\mathcal{N})$. So instead, we define COB_2 separately:

Definition 1.17. For P, λ and μ as above, let $\text{COB}_2(P, \lambda, \mu)$ stand for:

There are a $<\lambda$ -directed partial order $(S, <)$ of size μ and a sequence $(g_s)_{s \in S}$ of P -names for reals such that for each P -name f of a null set $(\exists s \in S)(\forall t > s) P \Vdash g_t \notin f$.

Lemma 1.18. 1. $\text{COB}_1(P, \lambda, \mu)$ implies $P \Vdash (\text{add}(\mathcal{N}) \geq \lambda \ \& \ \text{cof}(\mathcal{N}) \leq \mu)$.
 2. $\text{COB}_2(P, \lambda, \mu)$ implies $P \Vdash (\text{cov}(\mathcal{N}) \geq \lambda \ \& \ \text{non}(\mathcal{N}) \leq \mu)$.
 3. $\text{COB}_3(P, \lambda, \mu)$ implies $P \Vdash (\mathfrak{b} \geq \lambda \ \& \ \mathfrak{d} \leq \mu)$.
 4. $\text{COB}_4(P, \lambda, \mu)$ implies $P \Vdash (\text{non}(\mathcal{M}) \geq \lambda \ \& \ \text{cov}(\mathcal{M}) \leq \mu)$.

Proof. The cases $i \neq 2$ are direct consequences of Lemmas 1.3 and 1.16. The proof for $i = 2$ is analogous to the proof of Lemma 1.16. \square

Lemma 1.19. $\text{COB}_i(\mathbb{P}^5, \lambda_i, \lambda_5)$ holds (for $i = 1, 2, 3, 4$).

Proof. Set $S = S^i$ and $s < t$ if $w_s \subsetneq w_t$. As λ_i is regular, $(S, <)$ is $<\lambda_i$ -directed. Let g_s be the generic added at s (e.g., the partial random real in case of $i = 2$, etc). A \mathbb{P}^5 -name f depends (in a Borel way) on the subsequence of generics indexed by a countable set $w^* \subseteq \delta$. Fix some $s \in S^i$ such that $w_s \supseteq w^*$. Pick any $t > s$. Then $w_t \supseteq w_s$, so w_t contains all information to calculate f , so we can show that $P \Vdash f R_i g_t$. Let us list the possible cases: $i = 2$: A partial random real g_t will avoid the null set f . $i = 3$: A partial Hechler real g_t will dominate f . $i = 4$: A partial eventually different real g_t will be eventually different from f . As for $i = 1$, we use⁸ Lemma 1.4. \square

So to summarize what we know so far about \mathbb{P}^5 :

- COB_i holds for $i = 1, 2, 3, 4$. So the left hand characteristics are large.

⁷More specifically: this definition would give us the property $g_t \notin f$ only for the null sets of the specific form $f = \{h : \neg r R_2 h\} = N_r$, for some $r \in 2^\omega$; whereas we will define COB_2 to deal with all names f of null sets.

⁸Alternatively, we could use, instead of amoeba, some other Suslin ccc forcing that more directly adds an \mathfrak{R}_1 -dominating element of \mathcal{C} .

• LCU_i holds for $i = 1, 2, 4$. So the left hand characteristics other than \mathfrak{b} are small. However, LCU_3 (corresponding to “ \mathfrak{b} small”) is missing; and we cannot get it by a simple “preservation of $(\mathbb{R}_3, \lambda_3)$ -goodness” argument. Instead, we will argue in the following two sections that it is possible to choose the parameter $(w_\alpha)_{\alpha \in S^4}$ in such a way that LCU_3 holds as well.

1.3. Dealing with \mathfrak{b} without GCH. In this section, we follow (and slightly modify) the main construction of [GMS16].

In this section (and this section only) we will assume the following (in addition to Assumption 1.12, i.e., in particular to the assumption $\lambda_3 = \chi^+$):

Assumption 1.20. (*This section only.*) $2^\chi = |\delta_5| = \lambda_5$.

Set $S^0 = \lambda_5 \cup S^1 \cup S^2 \cup S^3$. So $\delta_5 = S^0 \cup S^4$, and \mathbb{P}^5 is a FS ccc iteration along δ_5 such that $\alpha \in S^0$ implies $|Q_\alpha| < \lambda_3$, i.e., $|Q_\alpha| \leq \chi$. Let us fix P_α -names

$$(1.21) \quad i_\alpha : Q_\alpha \rightarrow \chi \text{ injective}$$

(for $\alpha \in S^0$). Note that we can strengthen each $p \in \mathbb{P}^5$ to some q such that $\alpha \in \text{supp}(q) \cap S^0$ implies $q \restriction \alpha \Vdash i_\alpha(q(\alpha)) = \check{j}$ for some $j \in \chi$.

For $\alpha \in S^4$, Q_α is a partial eventually different forcing. At this point, we should specify which variant of this forcing we actually use:⁹

Definition 1.22.

- Eventually different forcing \mathbb{E} consists of all tuples (s, k, φ) , where $s \in \omega^{<\omega}$, $k \in \omega$, and $\varphi : \omega \rightarrow [\omega]^{<k}$ satisfies $s(i) \notin \varphi(i)$ for all $i \in \text{dom}(s)$.
- We define $(s', k', \varphi') \leq (s, k, \varphi)$ if $s \subseteq s'$, $k \leq k'$, and $\varphi(i) \subseteq \varphi'(i)$ for all i .
- The generic object $g^* = \bigcup_{(s,k,\varphi) \in G_{\mathbb{E}}} s$ is a function such that each condition (s, k, φ) forces that s is an initial segment of g^* , and $g^*(i) \notin \varphi(i)$ for all i .
- We call $s \in \omega^{<\omega}$ the “stem” of (s, k, φ) and $k \in \omega$ the “width”.

A density argument shows that g^* will be eventually different from all functions $f : \omega \rightarrow \omega$ from V .

The following is easy to see:

- If $p, q \in \mathbb{E}$ are compatible, then they have a greatest lower bound.
- Any finite set of conditions with the same stem has a lower bound (again with the same stem). So \mathbb{E} is σ -centered.
- If $q = (s', k', \varphi')$ and $p = (s, k, \varphi)$ and s' extends s , then p and q are compatible iff $s'(i) \notin \varphi(i)$ for all $i \in \text{dom}(s')$.
- If a condition $q^* = (s^*, k^*, \varphi^*)$ is compatible with each condition in a finite set $B \subseteq \mathbb{E}$, and s^* extends s for each $(s, k, \varphi) \in B$, then the set $B \cup \{q^*\}$ has a lower bound. (Use s^* as stem, and take the pointwise union of all φ that occur in $B \cup \{q^*\}$.)

We will not force with \mathbb{E} , but with a partial version of \mathbb{E} . In the P_α -extension (for $\alpha \in S^4$), this partial forcing $Q_\alpha = \mathbb{E}'$ is a (generally not complete) sub-forcing of \mathbb{E} which is easily seen to be closed under conjunctions (i.e., under the partial operation “greatest lower bound” of finite sets of conditions). Note that this implies that compatibility is absolute between \mathbb{E} and \mathbb{E}' , and that the previous items also hold for \mathbb{E}' . For later reference, let us explicitly state the last item:

Fact 1.23. *Assume $\mathbb{E}' \subseteq \mathbb{E}$ is closed under conjunctions. If a condition $q^* = (s^*, k^*, \varphi^*)$ in \mathbb{E}' is compatible with each condition in a finite set $B \subseteq \mathbb{E}'$, and s^* extends s for each $(s, k, \varphi) \in B$, then the set $B \cup \{q^*\}$ has a lower bound in \mathbb{E}' .*

⁹In the previous section it did not matter which variant we use.

Definition 1.24. Let D be a non-principal ultrafilter on ω , and let $\bar{p} = (p_n)_{n \in \omega} = (s, k, \varphi_n)_{n \in \omega}$ be a sequence of conditions in \mathbb{E} with the same stem and the same width. We define $\lim_D \bar{p}$ to be (s, k, φ_∞) , where for all i and all j we have $j \in \varphi_\infty(i) \Leftrightarrow \{n : j \in \varphi_n(i)\} \in D$.

The following is easy to see: $\lim_D \bar{p} \in \mathbb{E}$ and if $q \leq \lim_D \bar{p}$, then the set $B := \{n \in \omega : p_n \text{ compatible with } q\}$ is in D .

(Proof: $q = (s', k', \varphi') \leq \lim_D \bar{p} = (s, k, \varphi_\infty)$. So for each $i \in \text{dom}(s')$, $s'(i) \notin \varphi_\infty(i)$, and by the definition of the limit, $A^i := \{n : s'(i) \notin \varphi_n(i)\} \in D$. If $n \in \bigcap_{i \in \text{dom}(s')} A^i$, then p_n is compatible with q .)

As B is defined using only compatibility, the statement still holds for compatibility preserving subforcings. We state it for later reference in the following form:

Fact 1.25. Assume that \mathbb{E}' is a subforcing of \mathbb{E} closed under conjunctions, let \bar{p} be a sequence of \mathbb{E}' conditions with the same stem and width, and assume that $\lim_D(\bar{p}) \in \mathbb{E}'$ and that $q \leq_{\mathbb{E}'} \lim_D(\bar{p})$. Then $B := \{n \in \omega : p_n \text{ compatible with } q\}$ is in D .

Definition 1.26.

- A “partial guardrail” is a function h defined on a subset of δ_5 such that $h(\alpha) \in \chi$ for $\alpha \in S^0 \cap \text{dom}(h)$, and $h(\alpha) \in \omega^{<\omega} \times \omega$ for $\alpha \in S^4 \cap \text{dom}(h)$.
- A “countable guardrail” is a partial guardrail with countable domain. A “full guardrail” is a partial guardrail with domain δ_5 .

We will use the following lemma, which is a consequence of the Engelking-Karłowicz theorem [EK65] on the density of box products (cf. [GMS16, 5.1]):

Lemma 1.27. (As $|\delta_5| \leq 2^\chi$ and $\chi^{\aleph_0} = \chi$.) There is a family H^* of full guardrails with $|H^*| = \chi$, such that each countable guardrail is extended by some $h \in H^*$. We will fix such an H^* and enumerate it as $(h_\varepsilon)_{\varepsilon \in \chi}$.

Note that the notion of guardrail (and the density property required in Lemma 1.27) only depends on χ , δ_5 , S^0 and S^4 , i.e., on fixed parameters; so we can fix an H^* that will work for all cofinal parameters $\bar{w} = (w_\alpha)_{\alpha \in S^4}$.

Once we have decided on \bar{w} , and thus have defined \mathbb{P}^5 , we can define the following:

Definition 1.28. A condition $p \in \mathbb{P}^5$ follows the full guardrail h , if

- for all $\alpha \in S^0 \cap \text{dom}(p)$, the empty condition of P_α forces that $p(\alpha) \in Q_\alpha$ and $i_\alpha(p(\alpha)) = h(\alpha)$ (where i_α is defined in (1.21)), and
- for all $\alpha \in S^4 \cap \text{dom}(p)$:
 - $p \restriction \alpha$ forces that the pair of stem and width of $p(\alpha)$ is equal to $h(\alpha)$, and moreover
 - p is determined at α . (This was defined in 1.11(3): We already know in V a code (B, u) that evaluates to $p(\alpha)$.)

As we are dealing with a FS iteration, the set of conditions p determined at each position $\alpha \in \text{dom}(p)$ is easily seen to be dense (by induction). So note that

- the set of conditions p such that there is *some* guardrail h such that p follows h , is dense; while
- for each *fixed* guardrail h , the set of all conditions p following h , is *centered* (i.e., each finitely many such p are compatible).

Definition 1.29.

- A “ Δ -system with root ∇ following the full guardrail h ” is a family $\bar{p} = (p_i)_{i \in I}$ of conditions all following h , where $(\text{dom}(p_i) : i \in I)$ is a Δ -system with root ∇ in the usual sense (so $\nabla \subseteq \delta_5$ is finite).

- We will be particularly interested in countable Δ -systems. Let $(p_n : n \in \omega)$ be such a Δ -system with root ∇ following h , and assume that $\bar{D} = (D_\alpha : \alpha \in u)$ is

a sequence such that $u \supseteq \nabla \cap S^4$ and each D_α is a P_α -name of an ultrafilter on ω . Then we define the $\lim_{\bar{D}} \bar{p}$ to be the following function with domain ∇ :

- If $\beta \in \nabla \cap S^0$, then $\lim_{\bar{D}} \bar{p}(\beta)$ is the common value of all $p_n(\beta)$. (Recall that this value is already determined by the guardrail h .)
- If $\alpha \in \nabla \cap S^4$, then $\lim_{\bar{D}} \bar{p}(\alpha)$ is (forced by \mathbb{P}_α^5 to be) $\lim_{D_\alpha} (p_n(\alpha))_{n \in \omega}$.

Note that in general $\lim_{\bar{D}} \bar{p}$ will not be a condition in \mathbb{P}^5 : For $\alpha \in S^4 \cap \nabla$, the object $\lim_{\bar{D}} \bar{p}(\alpha)$ will be forced to be in the eventually different forcing \mathbb{E} , but not necessarily in the *partial* eventually different forcing $Q_\alpha \subseteq \mathbb{E}$.

Also note the following: If \bar{p} is a countable Δ -system, and $\alpha \in \nabla \cap S^4$, then $(p_n(\alpha))_{n \in \omega}$ is a ground-model-code-sequence (see Definition 1.11(4)). This follows trivially from the definition of “ p_n follows h ” and the fact that \bar{p} is in V .

Recall that we assume all of the parameters defining $\mathbb{P}^5 = (P_\alpha, Q_\alpha)_{\alpha \in \delta_5}$ to be fixed, apart from $(w_\alpha)_{\alpha \in S^4}$. Once we fix w_α for $\alpha \in S^4 \cap \beta$, we know P_β .

Lemma/Construction 1.30. *We can construct by induction on $\alpha \in \delta_5$ the sequences $(D_\alpha^\varepsilon)_{\varepsilon \in \chi}$ and, if $\alpha \in S^4$, also w_α , such that:*

- (a) Each D_α^ε is a P_α -name of a nonprincipal ultrafilter extending $\bigcup_{\beta < \alpha} D_\beta^\varepsilon$.
- (b) For each countable Δ -system \bar{p} in P_α which follows the guardrail $h_\varepsilon^* \in H^*$:
 $\lim_{(D_\beta^\varepsilon)_{\beta < \alpha}} \bar{p}$ is in $P_\alpha \dots$
- (c) ... and forces that $A_{\bar{p}} := \{n \in \omega : p_n \in G_\alpha\}$ is in D_α^ε .
- (d) (If $\alpha \in S^4$) $w_\alpha \subseteq \alpha$, $|w_\alpha| < \lambda_4$, and for all ground-model-code-sequences¹⁰ for elements of Q_α , the D_α^ε -limit is forced to be in Q_α as well (for all $\varepsilon \in \chi$).
 (Actually, the set of w_α satisfying this is an ω_1 -club set in $[\alpha]^{<\lambda_4}$.¹¹)

Proof. (b) for α limit: The root of a Δ -system is finite and therefore below some $\beta < \alpha$, so the limit exists (by induction) already in P_β .

(a+c) for α limit: It is enough to show, for each $\varepsilon \in \chi$, that P_α forces that the following generates a proper filter (i.e., any finite intersection of elements of this set is nonempty):

$$\bigcup_{\beta < \alpha} D_\beta^\varepsilon \cup \{A_{\bar{p}} : \bar{p} \text{ is a countable } \Delta\text{-system following } h_\varepsilon^* \text{ and } \lim_{(D_\beta^\varepsilon)_{\beta < \alpha}} \bar{p} \in G_\alpha\}.$$

(Then we let D_α^ε be any ultrafilter extending this set.)

So assume towards a contradiction that $q \in P_\alpha$ forces that $A \cap A_{\bar{p}^0} \cap \dots \cap A_{\bar{p}^{n-1}} = \emptyset$, where $A \in D_{\beta_0}^\varepsilon$ for some $\beta_0 < \alpha$ (we can assume β_0 is already decided in V) and \bar{p}^i as above with $q \leq \lim_{(D_\beta^\varepsilon)_{\beta < \alpha}} \bar{p}^i$ for $i < n$. Let $\beta_1 < \alpha$ be the maximum of the union of the roots of the \bar{p}^i , and set $\beta_2 := \max(\text{supp}(q))$ and $\gamma := \max(\beta_0, \beta_1, \beta_2) + 1$. By the induction hypothesis, q forces $A' := A \cap \bigcap_{i < n} A_{\bar{p}^i \upharpoonright \gamma} \in D_\gamma^\varepsilon$ (as $\lim_{(D_\beta^\varepsilon)_{\beta < \gamma}} \bar{p}^i \upharpoonright \gamma = \lim_{(D_\beta^\varepsilon)_{\beta < \alpha}} \bar{p}^i$, since the root lies below γ). As A' is a P_γ -name, we can find $q' \leq q$ in P_γ and $\ell \in \omega$ such that $q' \Vdash \ell \in A'$. We now find $q'' \leq q'$ in P_α by defining $q''(\beta)$ for each element β of the finite set $\bigcup_{i < n} \text{supp}(\bar{p}_\ell^i) \setminus \gamma$: For such β in S^0 , the guardrail gives a specific value $h_\varepsilon^*(\beta) \in Q_\beta$, which we use for $q''(\beta)$ as well. For $\beta \in S^4$, all conditions $p_\ell^i(\beta)$ (where defined) have the same stem and width $h_\varepsilon^*(\beta)$; hence there is a common extension $q''(\beta)$.

Clearly q'' forces that ℓ is in the allegedly empty set, the desired contradiction.

¹⁰see Definition 1.11(4).

¹¹I.e., for each $w^* \in [\alpha]^{<\lambda_4}$ there is a $w_\alpha \supseteq w^*$ satisfying (d), and if $(w^i)_{i \in \omega_1}$ is an increasing sequence of sets satisfying (d), then the limit $w_\alpha := \bigcup_{i \in \omega_1} w^i$ satisfies (d) as well.

(b) for $\alpha = \gamma + 1$ successor: Assume the nontrivial case, $\gamma \in S^4$: Write the Δ -system as $(p_i, q_i)_{i \in \omega}$ with $(p_i, q_i) \in P_\gamma * Q_\gamma$. As noted above, $(q_n)_{n \in \omega}$ is a ground-model-code-sequence, and by induction (d) holds for w_γ . So it is forced that the D_γ^ε -limit q^* of the q_n is in Q_γ . Again by induction, the limit p^* of the p_n exists as well; and (p^*, q^*) is the required limit.

(a+c) for $\alpha = \gamma + 1$ successor: We again have to show that P_α forces that the following is a filter base, for each $\varepsilon \in \chi$:

$$D_\gamma^\varepsilon \cup \{A_{\bar{p}} : \bar{p} \text{ is a countable } \Delta\text{-system following } h_\varepsilon^* \text{ and } \lim_{(D_\beta^\varepsilon)_{\beta < \alpha}} \bar{p} \in G_\alpha\}.$$

As above, assume that q forces $A \cap A_{\bar{p}^0} \cap \dots \cap A_{\bar{p}^{n-1}} = \emptyset$.

We can assume that $q \upharpoonright \gamma$ forces that $q(\gamma)$ is stronger than the limit of all $\bar{p}^i(\gamma)$ (for $i < n$). Thus, by Fact 1.25, each $B_i := \{\ell \in \omega : q(\gamma) \text{ compatible with } p_\ell^i(\gamma)\}$ is forced to be in D_γ^ε .

By induction, $q \upharpoonright \gamma$ forces that $A' := A \cap \bigcap_{i < n} A_{\bar{p}^i \upharpoonright \gamma} \in D_\gamma^\varepsilon$, and therefore also forces that $B' = A' \cap \bigcap_{i < n} B_i$ is in the ultrafilter and in particular nonempty. Work in the P_γ -extension by some generic filter containing $q \upharpoonright \gamma$. Fix some $\ell \in B'$. By the definition of B_i , $q(\gamma)$ is compatible with each $p_\ell^i(\gamma)$ for $i < n$. According to Fact 1.23 there is a common lower bound q'' .

$q \upharpoonright \gamma \Vdash_{P_\gamma} q'' \Vdash_{Q_\gamma} \ell \in A_{\bar{p}^i}$. I.e., $q \upharpoonright \gamma * q'' \leq q$ forces that ℓ is an element of the allegedly empty set.

(d) For any $w \subseteq \alpha$, let Q^w be the (P_α -name for) the partial eventually different forcing defined using w . Start with some $w^0 \subseteq \alpha$ of size $< \lambda_4$. There are $|w^0|^{\aleph_0}$ many ground-model sequences in Q^{w^0} . For any ε and any such sequence, the D_α^ε -limit is a real; so we can extend w^0 by a countable set to some w' such that $Q^{w'}$ contains the limit. We can do that for all $\varepsilon \in \chi$ and all sequences, resulting in some $w^1 \supseteq w^0$ still of size $< \lambda_4$. We iterate this construction and get w^i for $i \leq \omega_1$, taking the unions at limits. Then $w_\alpha := w^{\omega_1}$ is as required, as $Q_\alpha := Q^{w_\alpha} = \bigcup_{i < \omega_1} Q^{w^i}$.

So this proof actually shows that the set of w_α with the desired property is an ω_1 -club. \square

After carrying out the construction of this lemma, we get a forcing notion \mathbb{P}^5 satisfying the following:

Lemma 1.31. $\text{LCU}_3(\mathbb{P}^5, \kappa)$ for $\kappa \in [\lambda_3, \lambda_5]$, witnessed by the sequence $(c_\alpha)_{\alpha < \kappa}$ of the first κ many Cohen reals.

Proof. We want to show that for every \mathbb{P}^5 -name y there are coboundedly many $\alpha \in \kappa$ such that $\mathbb{P}^5 \Vdash \neg c_\alpha \leq^* y$.

Assume that p^* forces that there are unboundedly many $\alpha \in \kappa$ with $c_\alpha \leq^* y$, and enumerate them as $(\alpha_i)_{i \in \kappa}$ in increasing order (so in particular $\alpha_i \geq i$). Pick $p_i \leq p^*$ deciding α_i to be some β_i , and also deciding n_i such that $(\forall m \geq n_i) c_{\alpha_i}(m) \leq y(m)$. We can assume that $\beta_i \in \text{dom}(p_i)$. Note that β_i is a Cohen position (as $\beta_i < \kappa \leq \lambda_5$), and we can assume that $p_i(\beta_i)$ is a Cohen condition in V (and not just a P_{β_i} -name for such a condition). By thinning out, we may assume:

- All n_i are equal to some n^* .
- $(p_i)_{i \in \kappa}$ forms a Δ -system with root ∇ .
- $\beta_i \notin \nabla$, hence all β_i are distinct.
- (For any $\beta \in \kappa$, at most $|\beta|$ many p_i can force $\alpha_i = \beta$, as p_i forces that $\alpha_i \geq i$ for all i .)
- $p_i(\beta_i)$ is always the same Cohen condition s , without loss of generality of length $n^{**} \geq n^*$.

(Otherwise extend s .)

Pick the first ω many elements $(p_i)_{i \in \omega}$ of this Δ -system. Now extend each p_i to p'_i by extending the Cohen condition $p_i(\beta_i) = s$ to $s \frown i$ (i.e., forcing $c_{\alpha_i}(n^{**}) = i$). Note that $(p'_i)_{i \in \omega}$ is still a countable Δ -system, following some new countable guardrail and therefore some full guardrail $h_\varepsilon^* \in H^*$.

Accordingly, the limit $\lim_{(D_\alpha^\varepsilon)_{\alpha \in \delta_5}} \bar{p}'$ forces that infinitely many of the p'_i are in the generic filter. But each such p'_i forces that $c_{\alpha_i}(n^{**}) = i \leq y(n^{**})$, a contradiction. \square

1.4. Recovering GCH. For the rest of the paper we will assume the following for the ground model V (in addition to Assumption 1.12):

Assumption 1.32. *GCH holds.*

(Note that this is incompatible with Assumption 1.20.)

Recall that all parameters used to define \mathbb{P}^5 are fixed, apart from $\bar{w} = (w_\alpha)_{\alpha \in S^4}$.

Lemma 1.33. *We can choose \bar{w} such that $\text{LCU}_3(\mathbb{P}^5, \kappa)$ holds for all regular $\kappa \in [\lambda_3, \lambda_5]$.*

For the proof, we will use the following easy observation:

Lemma 1.34. *Assume χ is a cardinal and B a set and $X^0 \in [B]^\chi$, \mathbb{R} is a χ^+ -cc forcing notion, and C is an \mathbb{R} -name such that the empty condition forces that C is an ω_1 -club subset of $[B]^\chi$. Then there is a set $X \supseteq X^0$ (in the ground model) such that the empty condition forces $X \in C$.*

Proof. By induction, choose (in the ground model) sequences $X^\alpha, \tilde{X}^\alpha$ for $\alpha < \omega_1$ such that X^α is in $[B]^\chi$, the sequence of the X^α is increasing with α , \tilde{X}^α is an \mathbb{R} -name, and the empty condition forces: “ \tilde{X}^α is in C and is a superset of X^α ; and the sequence of the \tilde{X}^α is increasing (not necessarily continuous).” Moreover, the empty condition forces $\tilde{X}^\alpha \subseteq X^{\alpha+1}$. (In a limit step γ , we set $X^\gamma = \bigcup_{\alpha < \gamma} X^\alpha$, and in a successor step $\alpha + 1$ we use χ^+ -cc to cover the name \tilde{X}^α .) Then $X = \bigcup_{\alpha \in \omega_1} X^\alpha$ is as required. \square

Proof of Lemma 1.33. Let \mathbb{R} be a $< \chi$ -closed χ^+ -cc p.o. that forces $2^\chi = \lambda_5$.

In the \mathbb{R} -extension V^* , Assumption 1.20 holds; and Assumption 1.12 still holds for the fixed parameters.¹²

So in V^* , we can perform the inductive Construction 1.30, where now “ground model” refers to V^* , not V (e.g., when we talk about determined positions, or ground-model-code-sequences, etc). Actually, we can construct in V the following, by induction on $\alpha \in \delta_5$, and starting with some cofinal $\bar{w}^{\text{initial}} = (w_\alpha^{\text{initial}})_{\alpha \in S^4}$ in V :

- An \mathbb{R} -name $(D_\alpha^\varepsilon)_{\varepsilon \in \chi}$ (forced to be constructed) according to 1.30(a,b,c).
- If $\alpha \in S^4$, some $w_\alpha \supseteq w_\alpha^{\text{initial}}$ in V such that \mathbb{R} forces w_α satisfies 1.30(d).
(We can do this by Lemma 1.34, as the set of potential w_α 's is an ω_1 -clubset of $[\alpha]^{< \lambda_4}$.)

So we get in V a cofinal parameter \bar{w} satisfying the following: In the \mathbb{R} -extension V^* , the same parameters define a forcing (call it $\mathbb{P}^{*,5}$) satisfying $\text{LCU}_3(\mathbb{P}^{*,5}, \kappa)$ in V^* .

$\mathbb{P}^{*,5}$ is basically the same as \mathbb{P}^5 . More formally:

¹²In particular, $(w_\alpha)_{\alpha \in S^i}$ is still cofinal in $[\delta_5]^{< \lambda_i}$: For $i = 1, 2$, the forcing \mathbb{R} doesn't add any new elements of $[\delta_5]^{< \lambda_i}$ as \mathbb{R} is λ_i -closed; for $i = 3$ any new subset of δ_5 of size $\theta < \lambda_3$ is contained in a ground model set of size at most $\theta \times \chi < \lambda_3$, as \mathbb{R} is χ^+ -cc.

In the \mathbb{R} -extension V^* , $\mathbb{P}^5 = (P_\alpha, Q_\alpha)_{\alpha < \delta_5}$ (the iteration constructed in V) is canonically densely embedded into $\mathbb{P}^{*,5} = (P_\alpha^*, Q_\alpha^*)_{\alpha < \delta_5}$ (the iteration constructed in V^* using the same parameters).

Proof: By induction, we show (in the \mathbb{R} -extension) that P_α^* forces that Q_α^* (evaluated by the P_α^* -generic) is equal to Q_α (evaluated by the induced P_α -generic, as per induction hypothesis): Every element of Q_α^* is a Borel function (which already exists in V) applied to the generics at a countable sequence of indices in w_α (which also already exists in V).

This implies:

In V , $\text{LCU}_3(\mathbb{P}^5, \kappa)$ holds for all $\kappa \in [\lambda_3, \lambda_5]$, witnessed by the first κ many Cohen reals.

Proof: Let y be a \mathbb{P}^5 -name of a real. In V^* , we can interpret y as $\mathbb{P}^{*,5}$ -name, and as $\text{LCU}_3(\mathbb{P}^{*,5}, \kappa)$ holds, we get $(\exists \alpha \in \kappa) (\forall \beta \in \kappa \setminus \alpha) \mathbb{P}^{*,5} \Vdash c_\beta \not\leq^* y$, where c_β is the Cohen added at β . As $\chi < \kappa$, there is in V an upper bound $\alpha^* < \kappa$ for the possible values of α . For any $\beta \in \kappa \setminus \alpha^*$, we have (in V) $\mathbb{P}^5 \Vdash c_\beta \not\leq^* y$ (by absoluteness). \square

To summarize:

Theorem 1.35. *Assuming GCH and given λ_i as in Assumption 1.12, we can find parameters¹³ such that the FS ccc iteration \mathbb{P}^5 as defined in 1.13 satisfies, for $i = 1, 2, 3, 4$:*

- $\text{LCU}_i(\mathbb{P}^5, \kappa)$ holds for any regular cardinal κ in $[\lambda_i, \lambda_5]$.
- $\text{COB}_i(\mathbb{P}^5, \lambda_i, \lambda_5)$ holds.

So in particular \mathbb{P}^5 forces $\text{add}(\mathcal{N}) = \lambda_1$, $\text{cov}(\mathcal{N}) = \lambda_2$, $\mathfrak{b} = \lambda_3$, $\text{non}(\mathcal{M}) = \lambda_4$ and $\text{cov}(\mathcal{M}) = \mathfrak{d} = \text{non}(\mathcal{N}) = \text{cof}(\mathcal{N}) = \lambda_5 = 2^{\aleph_0}$.

For the rest of the paper we fix these parameters and thus the forcing \mathbb{P}^5 .

2. BOOLEAN ULTRAPOWERS

In Subsections 2.1 and 2.2 we describe how to get an elementary embedding (which we call a *BUP-embedding*) $j : V \rightarrow M$ with $\text{cr}(j) = \kappa$ and $\text{cf}(j(\kappa)) = |j(\kappa)| = \theta$, assuming κ is strongly compact and $\theta > \kappa$ is a regular cardinal with $\theta^\kappa = \theta$.

In Subsections 2.3 and 2.4 we show how to use such embeddings to transform a ccc forcing P to $j(P)$ while preserving some of the values forced to the entries of Cichoń's diagram (and changing others).

2.1. Boolean ultrapowers. Boolean ultrapowers generalize ordinary ultrapowers by using arbitrary Boolean algebras instead of the power set algebra.

We assume that κ is strongly compact and that B is a κ -distributive, κ^+ -cc, atomless complete Boolean algebra. Then every κ -complete filter in B can be extended to a κ -complete ultrafilter U .¹⁴ Also, there is a maximal antichain A_0 in B of size κ such that $A_0 \cap U = \emptyset$ (i.e., U is not κ^+ -complete).¹⁵

For now, fix some κ -complete ultrafilter U .

The Boolean algebra B can be used as forcing notion. As usual, V (or: the ground model) denotes the universe we “start with”. In the following, we will not actually force with B (and in this subsection and the following subsection, we will not force with anything, we always

¹³I.e., we set $\delta_5 = \lambda_5 + \lambda_5$, and find $(S^i)_{i=1,\dots,4}$ and $\bar{w} = (w_\alpha)_{\alpha \in \delta_5}$.

¹⁴For this, neither κ^+ -cc nor atomless is required, and κ -complete is sufficient. The proof is straightforward; the first proof that we are aware of has been published in [KT64].

¹⁵Proof: Let A be a maximal antichain in the open dense set $B \setminus U$; by κ^+ -cc $|A| \leq \kappa$. And A cannot have size $< \kappa$, as otherwise it would meet the κ -complete U .

remain in V); but we still use forcing notation. In particular, we call the usual B -names “forcing names”.

A *BUP-name* (or: labeled antichain) x is a function $A \rightarrow V$ whose domain is a maximal antichain of B . We may write $A(x)$ to denote A .

Each BUP-name corresponds to a forcing name¹⁶ for an element of V . We will identify the BUP-name and the corresponding forcing name. In turn, every forcing name τ for an element of V has a forcing-equivalent BUP-name. In particular there is a *standard BUP-name* \check{v} for each $v \in V$.

We can calculate, for two BUP-names x and y , the Boolean value $\llbracket x = y \rrbracket$. We call x and y *equivalent*, if $\llbracket x = y \rrbracket \in U$ (the κ -complete ultrafilter fixed above).

For example, any two standard BUP-names for the same $v \in V$ trivially are equivalent (as $\mathbb{1}_B \in U$). So we can speak (modulo equivalence) of *the* standard BUP-name for v .

The *Boolean ultrapower* M^- consists of the equivalence classes $[x]$ of BUP-names x ; and we define $[x] \in^- [y]$ by $\llbracket x \in y \rrbracket \in U$. We are interested in the \in -structure (M^-, \in^-) . We let $j^- : V \rightarrow M^-$ map v to $[\check{v}]$.

Given BUP-names x_1, \dots, x_n and an \in -formula φ , the truth value $\llbracket \varphi^V(x_1, \dots, x_n) \rrbracket$ is well defined (it is the weakest element of B forcing that in the ground model $\varphi(x_1, \dots, x_n)$ holds, which makes sense as x_1, \dots, x_n are guaranteed to be in the ground model).

A straightforward induction (which can be found in [KTT18, Sec. 2]) shows:

- Łoś’s theorem: $(M^-, \in^-) \models \varphi([x_1], \dots, [x_n])$ iff $\llbracket \varphi^V(x_1, \dots, x_n) \rrbracket \in U$.
- $j^- : (V, \in) \rightarrow (M^-, \in^-)$ is an elementary embedding.
- In particular, (M^-, \in^-) is a ZFC model.

As U is σ -complete, (M^-, \in^-) is wellfounded. So we let M be the transitive collapse of (M^-, \in^-) , and let $j : V \rightarrow M$ be the composition of j^- with the collapse. We denote the collapse of $[x]$ by x^U . So in particular $\check{v}^U = j(v)$.

- Facts 2.1.**
- $M \models \varphi(x_1^U, \dots, x_n^U)$ iff $\llbracket \varphi^V(x_1, \dots, x_n) \rrbracket \in U$. In particular, $j : V \rightarrow M$ is an elementary embedding.
 - If $|Y| < \kappa$, then $j(Y) = j''Y$. In particular, j restricted to κ is the identity. M is closed under $<\kappa$ -sequences.
 - $j(\kappa) \neq \kappa$, i.e., $\kappa = \text{cr}(j)$.

As we have already mentioned, an arbitrary forcing name for an element of V has a forcing-equivalent BUP-name, i.e., a maximal antichain labeled with elements of V . If τ is a forcing name for an element of Y ($Y \in V$), then without loss of generality τ corresponds to a maximal antichain labeled with elements of Y . We call such an object y a “BUP-name for an element of $j(Y)$ ” (and not “for an element of Y ”, for the obvious reason: unlike in the case of a forcing extension, y^U is generally not in Y , but, by definition of \in^- , it is in $j(Y)$).

Lemma 2.2. *If the partial order (S, \leq) is $\leq\kappa$ -directed, then $j''S$ is cofinal in $j(S)$.*

Proof. Let x^U be some element of $j(S)$; without loss of generality we can assume that x is a labeled antichain which only uses elements of S as labels. The size of the antichain is at most κ , so all labels have some common upper bound s_0 . Then $\llbracket x \leq s_0 \rrbracket$ is $\mathbb{1}_B$, and thus in U ; so $(M^-, \in^-) \models [x] \leq [s_0]$, i.e., $j(s_0) \geq x^U$ as required. \square

For later reference, let us summarize what we know about j in the form of a definition:

¹⁶more specifically, to the forcing name $\{(\check{x}(a), a) : a \in A(x)\}$.

Definition 2.3. A BUP-embedding is an elementary embedding $j : V \rightarrow M$ (M transitive) with critical point κ , such that M is $<\kappa$ -closed and such that $j''S$ is cofinal in $j(S)$ for every $\leq\kappa$ -directed partial order S .

So the embedding j defined as above for a κ -distributive, κ^+ -cc atomless complete Boolean algebra and a κ -complete ultrafilter U is a BUP-embedding.

Lemma 2.4. *Let j be a BUP-embedding with $\text{cr}(j) = \kappa$.*

- *If $|A| < \kappa$, then $j''A = j(A)$.*
- *If S is a $<\lambda$ -directed partial order for some regular $\lambda < \kappa$, then $j(S)$ is $<\lambda$ -directed.*
- *If $\text{cf}(\alpha) \neq \kappa$, then $j''\alpha$ is cofinal in $j(\alpha)$, so in particular $\text{cf}(j(\alpha)) = \text{cf}(\alpha)$.*

Proof. For the second item, use that M believes that $j(S)$ is $<\lambda$ -directed and that M is $<\kappa$ -closed. For the last item, assume $\text{cf}(\alpha) = \lambda \neq \kappa$, witnessed by some strictly increasing cofinal function $f : \lambda \rightarrow \alpha$. If $\lambda < \kappa$, then M thinks that $j(f)$ is strictly increasing cofinal from $j(\lambda) = \lambda$ to $j(\alpha)$, which is absolute. If $\lambda > \kappa$, then α is a $\leq\kappa$ -directed (linear) order, so $j''\alpha$ is cofinal in $j(\alpha)$. So $j''f$, i.e. $(j(\zeta), j(f(\zeta)))_{\zeta \in \lambda}$, witnesses that $\text{cf}(j''\lambda) = \text{cf}(j''\alpha) = \text{cf}(j(\alpha))$, and $\text{cf}(j''\lambda) = \text{cf}(\lambda) = \lambda$ (as these orders are isomorphic). \square

2.2. The algebra and the filter. For a strongly compact cardinal we can get large $\text{cf}(j(\kappa))$:

Lemma 2.5. *Let κ be strongly compact, $\theta > \kappa$ and $\text{cf}(\theta) > \kappa$. Then there is a BUP-embedding j with $\text{cr}(j) = \kappa$ such that*

- (1) $\text{cf}(j(\kappa)) = \text{cf}(\theta)$ and $j(\kappa) \geq \theta$.
- (2) $|j(\mu)| \leq \max(\mu, \theta)^\kappa$ for any μ .
- (3) In particular, if $\theta^\kappa = \theta$ and $\kappa \leq \mu \leq \theta$ then $|j(\mu)| = \theta$.

We will use this in the following form:

Definition 2.6. A “BUP-embedding from κ to θ ” is a BUP-embedding j with critical point κ such that $\text{cf}(j(\kappa)) = |j(\kappa)| = \theta$ (in particular κ and θ are regular).

The lemma immediately implies:

Corollary 2.7. *Assume κ is strongly compact and $\theta > \kappa$ is a regular cardinal such that $\theta^\kappa = \theta$. Then there is a BUP-embedding j from κ to θ . (And $|j(\mu)| = \theta$ whenever $\kappa \leq \mu \leq \theta$.)*

Proof of Lemma 2.5. Let B be the complete Boolean algebra generated by the forcing notion $P_{\kappa, \theta}$ consisting of partial functions from θ to κ with domain of size $<\kappa$, ordered by extension. Clearly B is $<\kappa$ -distributive (as $P_{\kappa, \theta}$ is even $<\kappa$ -closed) and κ^+ -cc.

The forcing adds a canonical generic function $f^* : \theta \rightarrow \kappa$. So for each $\delta \in \theta$, $f^*(\delta)$ is a forcing name for an element of κ , and thus a BUP-name for an element of $j(\kappa)$.

Let x be some other BUP-name for an element of $j(\kappa)$, i.e., an antichain A of size κ labeled with elements of κ . As $P_{\kappa, \theta}$ is dense in $B \setminus \{0_B\}$, we can assume that $A \subseteq P_{\kappa, \theta}$. Let $\delta \in \theta$ be bigger than the supremum of the domain of a for each $a \in A$. We call such a pair (x, δ) “suitable”, and set $b_{x, \delta} := \llbracket f^*(\delta) > x \rrbracket$. We claim that these elements generate a κ -complete filter. To see this, fix suitable pairs (x_i, δ_i) for $i < \mu < \kappa$; we have to show that $\bigwedge_{i \in \mu} b_{x_i, \delta_i} \neq 0$. Enumerate $\{\delta_i : i \in \mu\}$ increasing (and without repetitions) as δ^ℓ for $\ell \in \gamma \leq \mu$. Set $A_\ell = \{i : \delta_i = \delta^\ell\}$. Given q_ℓ , define $q_{\ell+1} \in P_{\kappa, \theta}$ as follows: $q_{\ell+1} \leq q_\ell$; $\delta^\ell \in \text{supp}(q_{\ell+1}) \subseteq \delta^\ell \cup \{\delta^\ell\}$; and $q_{\ell+1} \upharpoonright \delta^\ell$ decides for all $i \in A_\ell$ the values of x_i to be some α_i ; and $q_{\ell+1}(\delta^\ell) = \sup_{i \in A_\ell} (\alpha_i) + 1$. This ensures that $q_{\ell+1}$ is stronger than b_{x_i, δ_i} for

$i \in A_\ell$. For $\ell \leq \gamma$ limit, let q_ℓ be the union of $\{q_k : k < \ell\}$. Then q_γ is stronger than each b_{x_i, δ_i} .

As κ is strongly compact, we can extend the κ -complete filter generated by all b_{x_i, δ_i} to a κ -complete ultrafilter U . Then the sequence $f^*(\delta) \underset{\delta \in \theta}{U}$ is strictly increasing (as $(f^*(\delta), \delta')$ is suitable for all $\delta < \delta'$) and cofinal in $j(\kappa)$ (as we have just seen); so $\text{cf}(j(\kappa)) = \text{cf}(\theta)$ and $j(\kappa) \geq \theta$.

To get an upper bound for $j(\mu)$ for any cardinal μ , we count all possible BUP-names for elements of $j(\mu)$. As we can assume that the antichains are subsets of $P_{\kappa, \theta}$, which has size $\theta^{<\kappa}$, we get the upper bound $|j(\mu)| \leq [\theta^{<\kappa}]^\kappa \times \mu^\kappa = \max(\theta, \mu)^\kappa$. \square

2.3. The ultrapower of a forcing notion. We now investigate the relation of a forcing notion $P \in V$ and its image $j(P) \in M$, which we use as forcing notion over V . (Think of P as being one of the forcings of Section 1; it has no relation with the Boolean algebra B used to construct j .)

Note that as $j(P) \in M$ and M is transitive, every $j(P)$ -generic filter G over V is trivially generic over M as well, and we will use absoluteness between $M[G]$ and $V[G]$ to prove various properties of $j(P)$.

Lemma 2.8. *Let $j : V \rightarrow M$ be elementary, M transitive and $<\kappa$ -closed with $\text{cr}(j) = \kappa$. Assume that P is ν -cc for some $\nu < \kappa$.*

- (1) $j(P)$ is ν -cc.
- (2) If τ is (in V) a $j(P)$ -name for an element of $M[G]$, then there is a $j(P)$ -name σ in M such that the empty condition forces $\sigma = \tau$.
- (3) In particular, every $j(P)$ -name for a real, a Borel-code, a countable sequence of reals, etc., is in M (more formally: has an equivalent name in M).
- (4) $M[G]$ is $<\kappa$ -closed in $V[G]$.
- (5) If $\xi < \kappa$ and P forces $2^\xi = \lambda$, then $j(P)$ forces $2^\xi = |j(\lambda)|$.
- (6) $j''P$, which is isomorphic to P via j , is a complete subforcing of $j(P)$.

Proof. (1): If $A \subseteq j(P)$ has size ν , then $A \in M$, and by elementarity M thinks that A is not an antichain, which is absolute.

(2): τ corresponds to (A, f) where $A \subseteq j(P)$ is a maximal antichain and $f : A \rightarrow M$ maps a to a $j(P)$ -name in M . As $j(P)$ is ν -cc and M $<\kappa$ -closed, (A, f) is in M and we can interpret in M (A, f) as a $j(P)$ -name σ .

This immediately implies (3) and (4): Given a $j(P)$ -name τ for a ζ -sequence of elements of $M[G]$, $\zeta < \kappa$, we can interpret τ as a ζ -sequence of names $(\tau_i)_{i < \zeta}$, and find for each τ_i an equivalent $j(P)$ -name σ_i in M . As M is $<\kappa$ -closed, the sequence $(\sigma_i)_{i < \zeta}$ is in M and defines a $j(P)$ -name in M equivalent to τ .

(And if τ is a $j(P)$ -name for a $<\kappa$ -sequence in $M[G]$, we can use the fact that κ is regular and that $j(P)$ is κ -cc to get a bound $\zeta < \kappa$ for the length of τ .)

(5) $M[G]$ thinks that $|2^\xi| = j(\lambda)$, and $2^\xi \cap V[G] = 2^\xi \cap M[G]$.

(6): It is clear that $j''P$ is an incompatibility-preserving subforcing of $j(P)$: $j(p) \leq j(q)$ in $j''P$ iff $p \leq q$ in P (by definition) iff M thinks that $j(p) \leq j(q)$ in $j(P)$ (by elementarity) iff this holds in V (by absoluteness); and the same argument works for compatibility instead of \leq . Similarly, assume $A \subseteq j''P$ is a maximal antichain. By definition, $B := j^{-1}(A) \subseteq P$ is one as well, and in particular of size $< \nu$. Therefore $j(B) = A$, and by elementarity M thinks that $A \subseteq j(P)$ is maximal, which holds in V by absoluteness. \square

To round off the picture, let us mention the following fact (which is however, not required for the rest of the paper):

Lemma 2.9. *If $P = (P_\alpha, Q_\alpha)_{\alpha < \delta}$ is a finite support (FS) ccc iteration of length δ , then $j(P)$ is a FS ccc iteration of length $j(\delta)$ (more formally: it is canonically equivalent to one).*

Proof. M certainly thinks that $j(P) = (P_\alpha^*, Q_\alpha^*)_{\alpha < j(\delta)}$ is a FS iteration of length $j(\delta)$.

By induction on α we define the FS ccc iteration $(\tilde{P}_\alpha, \tilde{Q}_\alpha)_{\alpha < j(\delta)}$ and show that P_α^* is a subforcing of \tilde{P}_α : Assume this is already the case for P_α^* . M thinks that Q_α^* is a P_α^* -name, so we can interpret it as \tilde{P}_α -name and use it as \tilde{Q}_α . Assume that (p, q) is an element (in V) of $\tilde{P}_\alpha * \tilde{Q}_\alpha$. So p forces that q is a name in M ; we can strengthen p to some p' that decides q to be the name $q' \in M$. By induction we can further strengthen p' to $p'' \in P_\alpha^*$, then $(p'', q') \in P_{\alpha+1}^*$ is stronger than (p, q) . (At limits there is nothing to do, as we use FS iterations.)

$j(P)$ is ccc according to Lemma 2.8(1). \square

2.4. Preservation of values of characteristics. Recall Definition 1.8 of LCU_i ; and Definitions 1.15 and 1.17 of COB_i .

Lemma 2.10. *Assume¹⁷ that P is ccc and that j is a BUP-embedding with critical point κ .*

- (1) $\text{LCU}_i(P, \delta)$ implies $\text{LCU}_i(j(P), j(\delta))$.
So if $\lambda \neq \kappa$ regular, then $\text{LCU}_i(P, \lambda)$ implies $\text{LCU}_i(j(P), \lambda)$.
- (2) Assume $\text{COB}_i(P, \lambda, \mu)$. If $\kappa > \lambda$, then $\text{COB}_i(j(P), \lambda, |j(\mu)|)$; if $\kappa < \lambda$, then $\text{COB}_i(j(P), \lambda, \mu)$.

Proof. (1) Let $\bar{x} = (x_\alpha)_{\alpha < \delta}$ be the sequence of P -names witnessing $\text{LCU}_i(P, \delta)$. So M thinks: For every $j(P)$ -name y of a real $(\exists \alpha \in j(\delta)) (\forall \beta \in j(\delta) \setminus \alpha) \neg ((j(\bar{x}))_\beta R_i y)$. This is absolute, so $j(\bar{x})$ witnesses $\text{LCU}_i(j(P), j(\delta))$.

The second claim follows from the fact that $\text{LCU}_i(j(P), j(\delta))$ is equivalent to $\text{LCU}_i(j(P), \text{cf}(j(\delta)))$ and that $\text{cf}(j(\lambda)) = \lambda$ for regular $\lambda \neq \kappa$.

(2) Let $(S, <)$ and \bar{g} witness $\text{COB}_i(P, \lambda, \mu)$. M thinks that

(*) for each $j(P)$ -name f : $(\exists s \in j(S)) (\forall t \in j(S)) (t > s \rightarrow j(P) \Vdash f R_i j(\bar{g})_t)$

(or, in the case $i = 2$, $j(P) \Vdash j(\bar{g})_t \notin f$, where f is the name of a null set). This is true in V as well: If f is a $j(P)$ -name for a real, then we can assume $f \in M$, and so we can find $s \in j(S)$ such that for all $t > s$, $M[G] \Vdash f R_i j(\bar{g})_t$, which holds in $V[G]$ as well, as R_i is absolute.

If $\lambda < \kappa$, then $j(\lambda) = \lambda$, and $j(S)$ is λ -directed in M and therefore in V as well, so we get $\text{COB}_i(j(P), \lambda, |j(\mu)|)$.

So assume $\lambda > \kappa$. We claim that $j''(S)$ and $j''\bar{g}$ witness $\text{COB}_i(j(P), \lambda, \mu)$. $j''S$ is isomorphic to S , so directedness is trivial. Given a $j(P)$ -name f of a real, without loss of generality in M , there is in M a cone with tip $s \in j(S)$ as in (*). As $j''S$ is cofinal in $j(S)$ there is some $s' \in S$ such that $j(s') > s$. Then for all $t > s'$, i.e., $j(t) > j(s')$, we get $j(P) \Vdash f R_i j(\bar{g})_t$. (Or, in case $i = 2$, $j(P) \Vdash j(\bar{g})_t \notin f$). \square

We list the specific cases that we will use:

Corollary 2.11. *Let j be a BUP embedding from κ to θ .*

- (a) $\text{LCU}_i(P, \lambda)$ for a regular $\lambda \neq \kappa$ implies $\text{LCU}_i(j(P), \lambda)$.
- (b) $\text{LCU}_i(P, \kappa)$ implies $\text{LCU}_i(j(P), \theta)$.
- (c) $\text{COB}_i(P, \lambda, \mu)$ for $\kappa > \lambda$ and $\kappa \leq \mu \leq \theta$ implies $\text{COB}_i(j(P), \lambda, \theta)$.

¹⁷For most of the Lemma, the requirements of Lemma 2.8 are sufficient: We use ccc only to simplify notation as we do not have to indicate where we calculate cofinalities (in V or the $j(P)$ extensions $V[G]$); and we need BUP-embedding for the last part of (2) only.

(d) $\text{COB}_i(P, \lambda, \mu)$ for $\kappa < \lambda$ implies $\text{COB}_i(j(P), \lambda, \mu)$.

3. A FINITE ITERATION OF BUP EMBEDDINGS

We now have everything required for the main result:

Theorem 3.1. *Assume GCH and that $\aleph_1 < \kappa_9 < \lambda_1 < \kappa_8 < \lambda_2 < \kappa_7 < \lambda_3 < \kappa_6 < \lambda_4 < \lambda_5 < \lambda_6 < \lambda_7 < \lambda_8 < \lambda_9$ are regular, λ_3 a successor of a regular cardinal, λ_i not successor of a cardinal with countable cofinality for $i = 1, 2, 4, 5$, and κ_i strongly compact for $i = 6, 7, 8, 9$. Then there is a ccc forcing notion \mathbb{P}^9 resulting in:*

$$\begin{aligned} \text{add}(\mathcal{N}) = \lambda_1 < \text{cov}(\mathcal{N}) = \lambda_2 < \mathfrak{b} = \lambda_3 < \text{non}(\mathcal{M}) = \lambda_4 < \\ < \text{cov}(\mathcal{M}) = \lambda_5 < \mathfrak{d} = \lambda_6 < \text{non}(\mathcal{N}) = \lambda_7 < \text{cof}(\mathcal{N}) = \lambda_8 < 2^{\aleph_0} = \lambda_9. \end{aligned}$$

Proof. For $i = 6, \dots, 9$, let j_i be a BUP-embedding from κ_i to λ_i , i.e., $\text{cf}(j_i(\kappa_i)) = |j_i(\lambda_i)| = \lambda_i$. (Such an embedding exists according to Corollary 2.7.)

We use \mathbb{P}^5 of Theorem 1.35, and set $\mathbb{P}^{i+1} := j_{i+1}(\mathbb{P}^i)$ for $i = 5, 6, 7, 8$. In particular, $\mathbb{P}^9 = j_9(j_8(j_7(j_6(\mathbb{P}^5))))$.

We enumerate the relevant characteristics of Cichoń's diagram as $\mathfrak{x}_1, \dots, \mathfrak{x}_8$ in the desired increasing order as displayed in Figure 1. For $i = 1, \dots, 4$ (i.e., \mathfrak{x}_i in the left half) we set $i^* := 9 - i$ (so \mathfrak{x}_{i^*} is the dual of \mathfrak{x}_i in the right half).

Recall that according to Lemmas 1.9 and 1.18, $\text{LCU}_i(\lambda)$ implies $\mathfrak{x}_i \leq \lambda$ and $\mathfrak{x}_{i^*} \geq \lambda$; and $\text{COB}_i(\lambda, \mu)$ implies $\mathfrak{x}_i \geq \lambda$ and $\mathfrak{x}_{i^*} \leq \mu$.

Claim: \mathbb{P}^9 forces $2^{\aleph_0} = \lambda_9$.

Proof: By induction on $i = 5, \dots, 8$ each \mathbb{P}^{i+1} forces $2^{\aleph_0} = j_{i+1}(\lambda_i) = \lambda_{i+1}$ (according to Lem. 2.8(5) and Cor. 2.7).

Claim: $\text{LCU}_i(\mathbb{P}^9, \lambda_i)$ holds for $i = 1, \dots, 4$; as well as $\text{LCU}_4(\mathbb{P}^9, \lambda_5)$.

Proof: The statements hold for \mathbb{P}^5 by Thm. 1.35 and are preserved by Cor. 2.11(a).

This implies $\mathfrak{x}_i \leq \lambda_i$ for $i = 1, \dots, 4$; as well as $\mathfrak{x}_5 = \text{cov}(\mathcal{M}) \geq \lambda_5$.

Claim: $\text{LCU}_i(\mathbb{P}^9, \lambda_{i^*})$ holds for $i = 1, 2, 3$.

Proof: Note that $\kappa_{i^*+1} < \lambda_i < \kappa_{i^*} < \lambda_5$. So $\text{LCU}_i(\mathbb{P}^5, \kappa_{i^*})$ holds (Thm. 1.35). This implies $\text{LCU}_i(\mathbb{P}^\ell, \kappa_{i^*})$ for $\ell = 5, \dots, i^* - 1$ (Cor. 2.11(a)), then $\text{LCU}_i(\mathbb{P}^\ell, \lambda_{i^*})$ for $\ell = i^*$ (Cor. 2.11(b)), and then again $\text{LCU}_i(\mathbb{P}^\ell, \lambda_{i^*})$ for $\ell = i^* + 1, \dots, 9$ (again Cor. 2.11(a)).

This implies $\mathfrak{x}_\ell \geq \lambda_\ell$ for $\ell = 6, 7, 8$.

Claim: $\text{COB}_i(\mathbb{P}^9, \lambda_i, \lambda_{i^*})$ holds for $i = 1, 2, 3, 4$.

Proof: $\text{COB}_i(\mathbb{P}^5, \lambda_i, \lambda_5)$ holds by Theorem 1.35 and implies $\text{COB}_i(\mathbb{P}^\ell, \lambda_i, \lambda_\ell)$ for $\ell = 5, \dots, i^*$ (while $\kappa_\ell > \lambda_i$) (Cor. 2.11(c)), then $\text{COB}_i(\mathbb{P}^\ell, \lambda_i, \lambda_{i^*})$ for $\ell = i^* + 1, \dots, 9$ (Cor. 2.11(d)).

This implies $\mathfrak{x}_i \geq \lambda_i$ for $i = 1, \dots, 4$ as well as $\mathfrak{x}_\ell \leq \lambda_\ell$ for $\ell = 5, \dots, 8$. \square

4. QUESTIONS

The result poses some obvious questions. (Since the initial submission of the paper, some of the questions found *partial answers* which we mention in the following.)

(a) Can we prove the result without using large cardinals?

It would be quite surprising if compact cardinals are needed, but a proof without them will probably be a lot more complicated.

Partial answers:

- Gitik [Git19] points out that certain extender embeddings are BUP-embeddings, and that a variation of superstrongs is sufficient to construct the BUP-embeddings required in our construction.

- As mentioned, we think that the *result* does not require any large cardinals. The *proof* in this paper obviously does: Gitik (ibid.) notes that a measurable κ with Mitchell order $\geq \kappa^{++}$ is required to get a BUP-embedding from κ to some regular $\lambda > \kappa$. More generally, an easy argument given in [GKMS19, Sec. 3.1] (following a deeper observation pointed out by Mildenerger [Mil98]), shows that at least $0^\#$ is required to get a constellation of models of the type used in our proof.¹⁸
 - In [BCM18] (building on [Mej19a]), a construction for the left half of Cichón's diagram is introduced that additionally forces $\text{non}(\mathcal{M}) < 2^{\aleph_0}$. Accordingly, three strongly compact cardinals (or: subcompacts) are sufficient to get the ten different values.
- (b) Does the result still hold for other specific values of λ_i , such as $\lambda_i = \aleph_{i+1}$?

In our construction, the regular cardinals λ_i for $i = 4, \dots, 9$ can be chosen quite arbitrarily (above the compact κ_6 , that is). However, $\aleph_1, \lambda_1, \lambda_2$ and λ_3 each have to be separated by a compact cardinal (and furthermore λ_3 has to be a successor of a regular cardinal).

Partial answer: In [GKMS19], it is shown that we can choose the values quite freely. E.g., $\lambda_i = \aleph_{i+1}$ is possible; as is basically “any choice” of successor cardinals. We also show that we can replace any number of instances of $<$ by $=$.


- (c) Are other linear orders between the characteristics of Cichón's diagram consistent?

Note that in this paper, we use a FS ccc iteration of length δ with uncountable cofinality, cf. 2.9, which always results in $\text{non}(\mathcal{M}) \leq \text{cof}(\delta) \leq \text{cov}(\mathcal{M})$. Under these restrictions, there are only four possible assignments. Of course there are a lot more¹⁹ possibilities to assign $\lambda_1, \dots, \lambda_8$ to Cichón's diagram in a way that satisfies the known ZFC-provable (in)equalities. Figure 2.B is an example. Such orders require entirely different methods. (Even to get just the five different values $\aleph_1 = \lambda_1 = \lambda_2 = \lambda_3 = \lambda_4 = \lambda_5 < \lambda_6 < \lambda_7 < \lambda_8 < \lambda_9$ in this figure turned out to be rather involved [FGKS17, Sec. 11].)

Partial answer: Another of the orders compatible with FS ccc iterations, the one of Figure 2.A, is consistent [KST19]. See also [Mej19b]. (A different initial forcing gives the modified ordering of the left hand side; then the same construction and proof as in this paper gives us the whole diagram.)

¹⁸More specifically, for $j_6 : V \rightarrow M$ and G $j_6(\mathbb{P}^5)$ -generic over V , we know that a $M[G]$ is a κ -closed transitive subclass of $V[G]$, cf. Lemma 2.8(4). And we have $M[G] \models \text{non}(\mathcal{M}) = j_6(\lambda_4)$ and $V[G] \models \text{non}(\mathcal{M}) = \lambda_4 < |j_6(\lambda_4)| = \lambda_6$, which implies at least $0^\#$ (and probably a measurable).

¹⁹In fact, we counted 57 in addition to the 4 that are compatible with FS ccc.



(A) An ordering compatible with FS ccc. (B) Another one, incompatible with FS ccc.

FIGURE 2. Alternative orderings of the cardinal characteristics.

- (d) Is it consistent that other cardinal characteristics that have been studied,²⁰ in addition to the ones in Cichoń's diagram, have pairwise different values as well?

Partial answer: In [GKMS19], it is forced that additionally $\aleph_1 < \mathfrak{m} < \mathfrak{p} < \mathfrak{h} < \text{add}(\mathcal{N})$ holds.

REFERENCES

- [Ack37] Wilhelm Ackermann, *Die Widerspruchsfreiheit der allgemeinen Mengenlehre*, Math. Ann. **114** (1937), no. 1, 305–315. MR 1513141
- [Bar84] Tomek Bartoszyński, *Additivity of measure implies additivity of category*, Trans. Amer. Math. Soc. **281** (1984), no. 1, 209–213. MR 719666
- [Bar87] ———, *Combinatorial aspects of measure and category*, Fund. Math. **127** (1987), no. 3, 225–239. MR 917147
- [BCM18] Jörg Brendle, Miguel A Cardona, and Diego A Mejía, *Filter-linkedness and its effect on preservation of cardinal characteristics*, arXiv:1809.05004, 2018.
- [BJ95] Tomek Bartoszyński and Haim Judah, *Set theory*, A K Peters, Ltd., Wellesley, MA, 1995, On the structure of the real line. MR 1350295
- [BJS93] Tomek Bartoszyński, Haim Judah, and Saharon Shelah, *The Cichoń diagram*, J. Symbolic Logic **58** (1993), no. 2, 401–423. MR 1233917 (94m:03077)
- [Bla10] Andreas Blass, *Combinatorial cardinal characteristics of the continuum*, Handbook of set theory, vol. 1, Springer, Dordrecht, 2010, pp. 395–489. MR 2768685
- [Bre91] Jörg Brendle, *Larger cardinals in Cichoń's diagram*, J. Symbolic Logic **56** (1991), no. 3, 795–810. MR 1129144
- [CKP85] J. Cichoń, A. Kamburelis, and J. Pawlikowski, *On dense subsets of the measure algebra*, Proc. Amer. Math. Soc. **94** (1985), no. 1, 142–146. MR 781072 (86j:04001)
- [Coh63] Paul Cohen, *The independence of the continuum hypothesis*, Proc. Nat. Acad. Sci. U.S.A. **50** (1963), 1143–1148. MR 0157890
- [EK65] R. Engelking and M. Karłowicz, *Some theorems of set theory and their topological consequences*, Fund. Math. **57** (1965), 275–285. MR 0196693
- [FGKS17] Arthur Fischer, Martin Goldstern, Jakob Kellner, and Saharon Shelah, *Creature forcing and five cardinal characteristics in Cichoń's diagram*, Arch. Math. Logic **56** (2017), no. 7-8, 1045–1103. MR 3696076
- [Git19] Moti Gitik, *A remark on large cardinals in “cichon maximum”*, unpublished note, April 18th 2019, <http://www.math.tau.ac.il/~gitik/ssstr.pdf>.
- [GKMS19] Martin Goldstern, Jakob Kellner, Diego Mejía, and Saharon Shelah, *Controlling cardinal characteristics without adding reals*, arXiv:1904.02617v1, 2019.
- [GMS16] Martin Goldstern, Diego Alejandro Mejía, and Saharon Shelah, *The left side of Cichoń's diagram*, Proc. Amer. Math. Soc. **144** (2016), no. 9, 4025–4042. MR 3513558
- [JS90] Haim Judah and Saharon Shelah, *The Kunen-Miller chart (Lebesgue measure, the Baire property, Laver reals and preservation theorems for forcing)*, J. Symbolic Logic **55** (1990), no. 3, 909–927. MR 1071305
- [Kam89] Anastasis Kamburelis, *Iterations of Boolean algebras with measure*, Arch. Math. Logic **29** (1989), no. 1, 21–28. MR 1022984
- [KST19] Jakob Kellner, Saharon Shelah, and Anda Tănăsie, *Another ordering of the ten cardinal characteristics in cichoń's diagram*, Comment. Math. Univ. Carolin. **60** (2019), no. 1, 61–95.
- [KT64] H. J. Keisler and A. Tarski, *From accessible to inaccessible cardinals. Results holding for all accessible cardinal numbers and the problem of their extension to inaccessible ones*, Fund. Math. **53** (1964), 225–308. MR 0166107
- [KTT18] Jakob Kellner, Anda Ramona Tănăsie, and Fabio Elio Tonti, *Compact cardinals and eight values in Cichoń's diagram*, J. Symb. Log. **83** (2018), no. 2, 790–803. MR 3835089
- [KW07] Richard Kaye and Tin Lok Wong, *On interpretations of arithmetic and set theory*, Notre Dame J. Formal Logic **48** (2007), no. 4, 497–510. MR 2357524
- [Man71] Richard Mansfield, *The theory of Boolean ultrapowers*, Ann. Math. Logic **2** (1971), no. 3, 297–323. MR 0300887
- [Mej13] Diego Alejandro Mejía, *Matrix iterations and Cichon's diagram*, Arch. Math. Logic **52** (2013), no. 3-4, 261–278. MR 3047455

²⁰The most important ones are described in [Bla10].

- [Mej19a] Diego A. Mejía, *Matrix iterations with vertical support restrictions*, Proceedings of the 14th and 15th Asian Logic Conferences (Byunghan Kim, Jörg Brendle, Gyesik Lee, Fenrong Liu, R Ramanujam, Shashi M Srivastava, Akito Tsuboi, and Liang Yu, eds.), World Sci. Publ., 2019, pp. 213–248.
- [Mej19b] Diego A. Mejía, *A note on “Another ordering of the ten cardinal characteristics in Cichoń’s Diagram” and further remarks*, To appear in *Kyōto Daigaku Sūrikaiseki Kenkyūsho Kōkyūroku*, [arXiv:1904.00165](https://arxiv.org/abs/1904.00165), 2019.
- [Mil81] Arnold W. Miller, *Some properties of measure and category*, Trans. Amer. Math. Soc. **266** (1981), no. 1, 93–114. MR 613787 (84e:03058a)
- [Mil82] ———, *A characterization of the least cardinal for which the Baire category theorem fails*, Proc. Amer. Math. Soc. **86** (1982), no. 3, 498–502. MR 671224
- [Mil84] ———, *Additivity of measure implies dominating reals*, Proc. Amer. Math. Soc. **91** (1984), no. 1, 111–117. MR 735576 (85k:03032)
- [Mil98] Heike Mildnerberger, *Changing cardinal invariants of the reals without changing cardinals or the reals*, J. Symbolic Logic **63** (1998), no. 2, 593–599. MR 1625907
- [MS16] M. Malliaris and S. Shelah, *Existence of optimal ultrafilters and the fundamental complexity of simple theories*, Adv. Math. **290** (2016), 614–681. MR 3451934
- [Oxt80] John C. Oxtoby, *Measure and category*, second ed., Graduate Texts in Mathematics, vol. 2, Springer-Verlag, New York-Berlin, 1980. MR 584443
- [RS] D. Raghavan and S. Shelah, *Boolean ultrapowers and iterated forcing*, Preprint.
- [RS83] Jean Raisonniér and Jacques Stern, *Mesurabilité et propriété de Baire*, C. R. Acad. Sci. Paris Sér. I Math. **296** (1983), no. 7, 323–326. MR 697963 (84g:03077)
- [RS85] ———, *The strength of measurability hypotheses*, Israel J. Math. **50** (1985), no. 4, 337–349. MR 800191
- [She84] Saharon Shelah, *Can you take Solovay’s inaccessible away?*, Israel J. Math. **48** (1984), no. 1, 1–47. MR 768264
- [She04] ———, *Two cardinal invariants of the continuum ($\mathfrak{d} < \mathfrak{a}$) and FS linearly ordered iterated forcing*, Acta Math. **192** (2004), no. 2, 187–223. MR 2096454
- [Sol70] Robert M. Solovay, *A model of set-theory in which every set of reals is Lebesgue measurable*, Ann. of Math. (2) **92** (1970), 1–56. MR 0265151

INSTITUTE OF DISCRETE MATHEMATICS AND GEOMETRY, TECHNISCHE UNIVERSITÄT WIEN (TU WIEN).
Email address: martin.goldstern@tuwien.ac.at
URL: <http://www.tuwien.ac.at/goldstern/>

INSTITUTE OF DISCRETE MATHEMATICS AND GEOMETRY, TECHNISCHE UNIVERSITÄT WIEN (TU WIEN).
Email address: jakob.kellner@tuwien.ac.at
URL: <http://dmg.tuwien.ac.at/kellner/>

THE HEBREW UNIVERSITY OF JERUSALEM AND RUTGERS UNIVERSITY.
Email address: shlhetal@mat.huji.ac.il
URL: <http://shelah.logic.at/>