

On some variants of the club principle

Ashutosh Kumar*, Saharon Shelah†

Abstract

We study some asymptotic variants of the club principle. Along the way, we construct some forcings and use them to separate several of these principles.

1 Introduction

For a regular uncountable cardinal κ and a stationary $S \subseteq \bullet Lim(\kappa)$, the club principle \clubsuit_S says the following: There exists $\bar{A} = \langle A_\delta : \delta \in S \rangle$ where each A_δ is an unbounded subset of δ of order type $cf(\delta)$ such that for every $A \in [\kappa]^\kappa$, there exists some (equivalently, stationary many) $\delta \in S$ such that $A_\delta \subseteq A$. We say that \bar{A} is a \clubsuit_S witnessing sequence. If $\kappa = \omega_1$ and $S = \bullet Lim(\omega_1)$ is the set of all countable limit ordinals, we drop the S and write \clubsuit .

In [?], it was shown that \clubsuit^1 does not imply \clubsuit where \clubsuit^1 is the following statement: There exists $\bar{A} = \langle A_\delta : \delta \in \bullet Lim(\omega_1) \rangle$ where each A_δ is an unbounded subset of δ of order type ω such that for every $A \in [\omega_1]^{\aleph_1}$, there exists δ such that $A_\delta \setminus A$ is finite. For some other variants of the club principle, see [?, ?, ?].

Definition 1.1. For $a \in (0, 1]$ and a stationary set $S \subseteq \bullet Lim(\omega_1)$, the principle $\clubsuit_S^{\text{inf} \geq a}$ says the following: There exists $\bar{A} = \langle A_\delta : \delta \in S \rangle$ such that

- (a) each $A_\delta = \{\alpha_{\delta,n} : n < \omega\}$ and $\alpha_{\delta,n}$'s are increasing cofinal in δ and
- (b) for every $A \in [\omega_1]^{\aleph_1}$, there exists $\delta \in S$ such that

$$\liminf_n \frac{|\{k < n : \alpha_{\delta,k} \in A\}|}{n} \geq a$$

If $S = \bullet Lim(\omega_1)$, we write $\clubsuit^{\text{inf} \geq a}$. By \clubsuit^{lim} , we mean $\clubsuit^{\text{inf} \geq 1}$.

*Einstein Institute of Mathematics, The Hebrew University of Jerusalem, Edmond J Safra Campus, Givat Ram, Jerusalem 91904, Israel; email: akumar@math.huji.ac.il; Supported by a Postdoctoral Fellowship at the Einstein Institute of Mathematics funded by European Research Council grant 338821

†Einstein Institute of Mathematics, The Hebrew University of Jerusalem, Edmond J Safra Campus, Givat Ram, Jerusalem 91904, Israel and Department of Mathematics, Rutgers, The State University of New Jersey, Hill Center-Busch Campus, 110 Frelinghuysen Road, Piscataway, NJ 08854-8019, USA; email: shelah@math.huji.ac.il; Partially supported by European Research Council grant 338821; Publication no. 1136

It is clear that \clubsuit^1 implies \clubsuit^{lim} and for $0 < a < b \leq 1$, $\clubsuit^{\text{inf} \geq b}$ implies $\clubsuit^{\text{inf} \geq a}$. At the end of Section 1, we show that under CH, all of these principles are equivalent to diamond.

Theorem 1.2. *Assume CH. Then for every $a \in (0, 1]$, $\clubsuit^{\text{inf} \geq a}$ implies \diamond .*

The bulk of the work in this paper is to show the following.

Theorem 1.3.

- (1) $\clubsuit^{\text{lim}} \wedge \neg \clubsuit^1$ is consistent.
- (2) For every $a \in (0, 1]$, $\clubsuit^{\text{inf} \geq a} \wedge (\forall b > a) \neg \clubsuit^{\text{inf} \geq b}$ is consistent.
- (3) For every $a \in (0, 1]$, $\neg \clubsuit^{\text{inf} \geq a} \wedge (\forall b < a) \clubsuit^{\text{inf} \geq b}$ is consistent.

In Sections 2-5 we prove Theorem ??(1). In Section 6, we supply the necessary modifications to get parts (2) and (3). The forcing used is quite flexible and can be useful for separating many similar principles.

In Section 7, we introduce $\clubsuit^{\text{sup} \geq a}$ (defined analogously) and prove the following in ZFC.

Theorem 1.4. *For every $a, b \in (0, 1)$, $\clubsuit^{\text{sup} \geq a}$ is equivalent to $\clubsuit^{\text{sup} \geq b}$.*

Finally, in Section 8, we prove that

Theorem 1.5. $\clubsuit^{\text{sup} \geq 0.5} \wedge \neg \clubsuit^{\text{sup} \geq 1}$ is consistent.

On notation: $\bullet \text{Lim}(\kappa)$ denotes the set of all limit ordinals below κ . $\text{cf}(\alpha)$ is the cofinality of α . $S_\delta^\kappa = \{\alpha < \kappa : \text{cf}(\alpha) = \text{cf}(\delta)\}$. For $k \leq \omega$, ω^k is the k th ordinal power of ω with under ordinal exponentiation. For a, b sets of ordinals, then we write $a < b$ to denote $(\forall \alpha \in a)(\forall \beta \in b)(\alpha < \beta)$. In forcing, we use the convention that a larger condition is the stronger one - $p \geq q$ means p extends q .

1.1 CH and \clubsuit^{inf}

Recall that \diamond says the following: There exists $\langle A_\delta : \delta \in \bullet \text{Lim}(\omega_1) \rangle$ where each $A_\delta \subseteq \delta$ such that for every $A \in [\omega_1]^{\aleph_1}$, $\{\delta \in \bullet \text{Lim}(\omega_1) : A_\delta = A \cap \delta\}$ is stationary. An equivalent formulation (see [?]) is the following: There exists $\langle \mathcal{A}_\delta : \delta \in \bullet \text{Lim}(\omega_1) \rangle$ where each \mathcal{A}_δ is a countable family of subsets of δ such that for every $A \in [\omega_1]^{\aleph_1}$, $\{\delta \in \bullet \text{Lim}(\omega_1) : A \cap \delta \in \mathcal{A}_\delta\}$ is stationary.

Proof of Theorem ??: Assume CH. Suppose $a \in (0, 1]$ and $\clubsuit^{\text{inf} \geq a}$ holds as witnessed by $\bar{A} = \langle A_\delta : \delta \in \bullet \text{Lim}(\omega_1) \rangle$. Let $A_\delta = \{\alpha_{\delta, n} : n < \omega_1\}$ list A_δ in increasing order. Using CH, fix $\langle B_i : i < \omega_1 \rangle$ such that each $B_i \subseteq i$ and for every $B \in [\omega_1]^{\leq \aleph_0}$, there are uncountably many $i < \omega_1$ for which $B = B_i$.

For $\delta \in \bullet \text{Lim}(\omega_1)$, define \mathcal{A}_δ as follows. $A \in \mathcal{A}_\delta$ iff for some $u \subseteq \omega$ the following hold.

- (a) $\liminf_n |u \cap n|/n \geq a$.

(b) For every $m < n$ in u , $B_{\alpha_{\delta,m}} = B_{\alpha_{\delta,n}} \cap \alpha_{\delta,m}$ and $A = \bigcup_{n \in u} B_{\alpha_{\delta,n}}$.

We claim that each \mathcal{A}_δ is finite. In fact, $|\mathcal{A}_\delta| \leq 1/a$. To see this assume otherwise and let $\{A_k : k < K\}$ be pairwise distinct members of \mathcal{A}_δ where $Ka > 1$. Choose $\langle u_k : k < K \rangle$ witnessing $A_k \in \mathcal{A}_k$. Choose $N_1 < N_2$ such that the following hold.

(i) $\langle A_k \cap \alpha_{\delta, N_1} : k < K \rangle$ has pairwise distinct members

(ii) $|u_k \cap [N_1, N_2]| > (N_2 - N_1)/K$ for each $k < K$

By (ii), it follows that for some $j < k < K$, $[N_1, N_2] \cap u_j \cap u_k \neq \emptyset$. But if $n \in [N_1, N_2] \cap u_j \cap u_k$, then $B_{\alpha_{\delta,n}} = A_j \cap \alpha_{\delta,n} = A_k \cap \alpha_{\delta,n}$ which is impossible by (i).

To complete the proof it is enough to show the following.

Claim 1.6. *For every $X \in [\omega_1]^{\aleph_1}$, for every club $E \subseteq \omega_1$, there exists $\delta \in E$ such that $C \cap \delta \in \mathcal{A}_\delta$.*

Proof of Claim ???: Construct $\langle \alpha_i : i < \omega_1 \rangle$ such that α_i 's are increasing and for every $i < \omega_1$, $X \cap \sup_{j < i} \alpha_j = B_{\alpha_i}$. Choose $\delta \in E$ and $u \subseteq \omega$ such that $\liminf_n |u \cap n|/n \geq a$ and $\{\alpha_{\delta,n} : n \in u\} \subseteq \{\alpha_i : i < \omega_1\}$. It follows that $X \cap \delta = \bigcup_{n \in u} B_{\alpha_{\delta,n}} \in \mathcal{A}_\delta$. \square

2 Creatures

Fix a family $\{S_k : k < \omega\}$ of pairwise disjoint stationary subsets of ω_1 consisting of limit ordinals. We describe a ccc forcing which is somewhat intermediate between adding \aleph_1 Cohen reals and adding a Cohen subset of ω_1 .

Definition 2.1. *We say that (CR, Σ) is an \aleph_1 -CP (creating pair) if the following hold.*

(A) *We call members of CR creatures. For each $\mathfrak{c} \in CR$,*

(i) $\mathfrak{c} = (\text{dom}(\mathfrak{c}), \text{pos}(\mathfrak{c}), f_{\mathfrak{c}})$.

(ii) $\text{dom}(\mathfrak{c})$ is a non empty subset of ω_1 of order type $< \omega^\omega$.

(iii) *For every limit $\delta < \omega_1$, if $\text{dom}(\mathfrak{c}) \cap \delta$ is unbounded in δ , then for some $k \geq 1$, $\delta \in S_k$ and $\text{otp}(\text{dom}(\mathfrak{c}) \cap \delta) = \varepsilon + \omega^j$ for some $\varepsilon < \omega^\omega$ and $1 \leq j \leq k$ - In particular, for every $\delta \in S_0$, $\text{dom}(\mathfrak{c}) \cap \delta$ is bounded below δ .*

(iv) $\text{pos}(\mathfrak{c})$ (possibilities for \mathfrak{c}) is a countable set of functions from $\text{dom}(\mathfrak{c})$ to $\{0, 1\}$ and $f_{\mathfrak{c}} \in \text{pos}(\mathfrak{c})$.

(v) *If $\text{dom}(\mathfrak{c})$ is finite, then $\text{pos}(\mathfrak{c}) = \{f_{\mathfrak{c}}\}$ - We call such \mathfrak{c} finite creature.*

(B) *For every finite $u \subseteq \omega_1$, and $f : u \rightarrow \{0, 1\}$, there exists $\mathfrak{c} \in CR$ such that $\text{dom}(\mathfrak{c}) = u$ and $f_{\mathfrak{c}} = f$.*

(C) *For every $\delta < \omega_1$, $|\{\mathfrak{c} \in CR : \text{dom}(\mathfrak{c}) \subseteq \delta\}| \leq \aleph_0$.*

(D) Σ is a function with domain CR that satisfies the following.

(i) $\Sigma(\mathbf{c})$ is a countable set of finite tuples $\bar{\mathfrak{d}} = \langle \mathfrak{d}_k : k < n \rangle$ where

(a) $\mathfrak{d}_k \in CR$,

(b) $\text{dom}(\mathbf{c}) = \bigcup_{k < n} \text{dom}(\mathfrak{d}_k)$,

(c) $\text{dom}(\mathfrak{d}_k) < \text{dom}(\mathfrak{d}_{k+1})$ and

(d) whenever $f_k \in \text{pos}(\mathfrak{d}_k)$ for $k < n$, $\bigcup_{k < n} f_k \in \text{pos}(\mathbf{c})$.

(ii) Cuts: If $\mathbf{c} \in CR$ and $\alpha \in \text{dom}(\mathbf{c})$ then for some $\bar{\mathfrak{d}} = \langle \mathfrak{d}_k : k < n \rangle \in \Sigma(\mathbf{c})$, there exists $k < n$ such that $\min(\text{dom}(\mathfrak{d}_k)) = \alpha$.

(iii) $\langle \mathbf{c} \rangle \in \Sigma(\mathbf{c})$.

(iv) Transitivity: If $\langle \mathbf{c}_k : k < n \rangle \in \Sigma(\mathbf{c})$ and $\langle \mathfrak{d}_{k,l} : l < n_k \rangle \in \Sigma(\mathbf{c}_k)$ for $k < n$, then $\langle \mathfrak{d}_{k,l} : k < n, l < n_k \rangle \in \Sigma(\mathbf{c})$.

(E) Finite joins: If $\{\mathfrak{d}_k : k < n\} \subseteq CR$ and $\text{dom}(\mathfrak{d}_k) < \text{dom}(\mathfrak{d}_{k+1})$, then there exists $\mathbf{c} \in CR$ such that

(i) $\text{dom}(\mathbf{c}) = \bigcup_{k < n} \text{dom}(\mathfrak{d}_k)$,

(ii) $\text{pos}(\mathbf{c}) = \{\bigcup_{k < n} f_k : (\forall k < n)(f_k \in \text{pos}(\mathfrak{d}_k))\}$,

(iii) $f_{\mathbf{c}} = \bigcup_{k < n} f_{\mathfrak{d}_k}$ and

(iv) $\Sigma(\mathbf{c}) = \{\bigcup_{i < n} \bar{f}_i : (\forall i < n)(\bar{f}_i \in \Sigma(\mathfrak{d}_i))\}$.

Definition 2.2. Suppose (CR, Σ) is an \aleph_1 -CP. Define $\mathbb{Q} = \mathbb{Q}_{CR, \Sigma}$ to be the forcing whose conditions are $p = \{\mathbf{c}_k : k < n\}$ where $\mathbf{c}_k \in CR$ and $\text{dom}(\mathbf{c}_k) < \text{dom}(\mathbf{c}_{k+1})$. We write $\text{dom}(p)$ for $\bigcup_{\mathbf{c} \in p} \text{dom}(\mathbf{c})$. For $p, q \in \mathbb{Q}$, define $p \leq q$ iff for every $\mathbf{c} \in p$, there exists $\bar{\mathfrak{d}} = \langle \mathfrak{d}_k : k < n \rangle \in \Sigma(\mathbf{c})$ such that $\{\mathfrak{d}_k : k < n\} \subseteq q$. Define $\mathbb{Q} \upharpoonright \alpha = \{p \in \mathbb{Q} : \text{dom}(p) \subseteq \alpha\}$. Let

$$\mathring{f}_{\mathbb{Q}} = \bigcup \{f_{\mathfrak{d}} : (\exists p \in G_{\mathbb{Q}})(\mathfrak{d} \in p \text{ is a finite creature})\}$$

Note that $\Vdash_{\mathbb{Q}} \mathring{f}_{\mathbb{Q}} : \omega_1 \rightarrow \{0, 1\}$

Example: Let CR be the set of all finite creatures $\mathbf{c} = (F, \{f\}, f)$ - So $F \subseteq \omega_1$ is finite and $f : F \rightarrow \{0, 1\}$. Let $\Sigma(\mathbf{c})$ be the set of all $\bar{\mathfrak{d}}$ such that the join of the members of $\bar{\mathfrak{d}}$ is \mathbf{c} . Then forcing with $\mathbb{Q} = \mathbb{Q}_{CR, \Sigma}$ is same as adding \aleph_1 Cohen reals. Note that this destroys all old witnesses to \clubsuit^{lim} . We would later add more creatures to CR in such a way that while some old \clubsuit^{lim} witnessing sequences are preserved, all old \clubsuit^1 witnessing sequences are destroyed.

Recall that a forcing notion \mathbb{Q} has \aleph_1 as a precaliber if whenever $\{p_i : i < \omega_1\} \subseteq \mathbb{Q}$, there exists $X \in [\omega_1]^{\aleph_1}$ such that $\{p_i : i \in X\}$ is centered - i.e., for every finite $F \subseteq X$, there exists $p \in \mathbb{Q}$ such that $(\forall i \in F)(p_i \leq p)$.

Claim 2.3. Suppose (CR, Σ) is an \aleph_1 -CP. Let $\mathbb{Q} = \mathbb{Q}_{CR, \Sigma}$. Then \mathbb{Q} has \aleph_1 as a precaliber.

Proof of Claim ???: Suppose $\{p_i : i < \omega_1\} \in [\mathbb{Q}]^{\aleph_1}$. The map $i \mapsto k(i) = \sup(\bigcup_{\mathfrak{c} \in p_i} \text{dom}(\mathfrak{c}) \cap i)$ is regressive on S_0 . Choose $X_1 \in [S_0]^{\aleph_1}$ and $k(\star) < \omega_1$ such that for every $i \in X_1$, $k(i) = k(\star)$ and for every $i < j$ in X_1 , $\text{dom}(p_i) \cap \text{dom}(p_j) \subseteq k(\star)$. Using Definition ??(D)(ii), by possibly extending each p_i , we can assume that for every $\mathfrak{c} \in p_i$, either $\text{dom}(\mathfrak{c}) \subseteq k(\star)$ or $\text{inf}(\text{dom}(\mathfrak{c})) \geq k(\star)$. Since $\{\mathfrak{c} \in \text{CR} : \text{dom}(\mathfrak{c}) \subseteq k(\star)\}$ is countable, we can find $X \in [X_1]^{\aleph_1}$ such that for every $i \in X$, $\{\mathfrak{c} \in p_i : \text{dom}(\mathfrak{c}) \subseteq k(\star)\}$ does not depend on $i \in X$. Now for any finite $F \subseteq X$, $\bigcup_{i \in F} p_i$ is a common extension of $\{p_i : i \in F\}$. \square

Claim 2.4. *Suppose (CR, Σ) is an \aleph_1 -CP. Let $\mathbb{Q} = \mathbb{Q}_{\text{CR}, \Sigma}$. Let $\langle p_i : i < \omega_1 \rangle$ be a sequence of conditions in \mathbb{Q} such that for every $i < j < \omega_1$, $\sup(\text{dom}(p_i)) < \sup(\text{dom}(p_j))$. Then there exist $X \in [\omega_1]^{\aleph_1}$, $\langle q_i : i \in X \rangle$, $m < n < \omega$ such that for every $i \in X$*

- (a) $q_i \in \mathbb{Q}$, $q_i \geq p_i$ and $\text{dom}(q_i) = \text{dom}(p_i)$,
- (b) $q_i = \{\mathfrak{c}_{i,k} : k < n\}$ and for every $k < n - 1$, $\text{dom}(\mathfrak{c}_{i,k}) < \text{dom}(\mathfrak{c}_{i,k+1})$,
- (c) for $k < m$, $\mathfrak{c}_{i,k} = \mathfrak{c}_k$ does not depend on $i \in X$.
- (d) for every $j < j'$ in X , $\text{dom}(\mathfrak{c}_{j,n-1}) < \text{dom}(\mathfrak{c}_{j',m})$ and
- (e) $\text{otp}(\text{dom}(\mathfrak{c}_{i,k}))$ does not depend on $i \in X$.

Proof of Claim ???: Just follow the argument in the proof of Claim ?? noting that $\text{dom}(p_i)$'s are unbounded in ω_1 . \square

3 Countable joins

In the course of club preservation arguments, we would like to be able to form new creatures out of old ones in the following way. Suppose $\langle q_i : i \geq 1 \rangle$ is a sequence of conditions in $\mathbb{Q} = \mathbb{Q}_{\text{CR}, \Sigma}$ which forms a Δ -system of an appropriate kind - It satisfies clauses (b)-(e) in Claim ??. We'd like to construct a new condition $q \in \mathbb{Q}$ such that $q \Vdash_{\mathbb{Q}} \text{"}\lim_n |\{i < n : q_i \in G_{\mathbb{Q}}\}|/n = 1 \text{"}$ and $\{i < \omega : q_i \notin G_{\mathbb{Q}}\}$ is infinite". This will require us to add "countable joins" of certain sequences of creatures to CR . This section introduces the countable join construction.

Definition 3.1. *For $\alpha < \omega_1$, we say that (CR_p, Σ_p) is a partial \aleph_1 -CP at α if for some \aleph_1 -CP (CR, Σ) ,*

- (1) $\text{CR}_p = \text{CR} \upharpoonright \alpha = \{\mathfrak{c} \in \text{CR} : \sup(\text{dom}(\mathfrak{c})) < \alpha\}$ and
- (2) $\Sigma_p = \Sigma \upharpoonright \text{CR}_p$.

Definition 3.2. *Suppose $k_{\star} \geq 1$, $\delta \in S_{k_{\star}}$ and (CR_p, Σ_p) is a partial \aleph_1 -CP at δ . Suppose $m < n < \omega$, $\delta \in S_{k_{\star}}$ for $k_{\star} \geq 1$ and $\bar{\mathfrak{d}}_i = \langle \mathfrak{d}_{i,k} : k < n \rangle$ satisfy the following for $1 \leq i < \omega$.*

- (a) $\mathfrak{d}_{i,k} \in \text{CR}_p$.
- (b) $\mathfrak{d}_{i,j} = \mathfrak{d}_j$ does not depend on i for $j < m$.

- (c) $\text{dom}(\mathfrak{d}_{i,k}) < \text{dom}(\mathfrak{d}_{i,k+1})$.
- (d) $\text{dom}(\mathfrak{d}_{i,n-1}) < \text{dom}(\mathfrak{d}_{i+1,m})$.
- (e) $\text{otp}(\text{dom}(\mathfrak{d}_{i,k}))$ only depends on k .
- (f) $W = \bigcup \{ \text{dom}(\mathfrak{d}_{i,k}) : 1 \leq i < \omega, k < n \}$ is unbounded in δ and has order type $\varepsilon + \omega^{j_\star}$ for some $\varepsilon < \omega_1$ and $1 \leq j_\star \leq k_\star$.

We say that $\langle \bar{\mathfrak{d}}_i : i \geq 1 \rangle$ is a joinable candidate for (CR_p, Σ_p) at δ .

For each $N \geq 1$ where N is a power of 2, we define new creatures $\mathfrak{c}_N^\star = (\text{dom}(\mathfrak{c}_N^\star), \text{pos}(\mathfrak{c}_N^\star), f_{\mathfrak{c}_N^\star})$ and $\Sigma_\star(\mathfrak{c}_N^\star)$, as follows.

- (1) $\text{dom}(\mathfrak{c}_1^\star) = W$ and $\text{dom}(\mathfrak{c}_N^\star) = \bigcup \{ \text{dom}(\mathfrak{d}_{i,k}) : N \leq i < \omega, m \leq k < n \}$ for $N \geq 2$.
- (2) $f_{\mathfrak{c}_1^\star} = \bigcup \{ f_{\mathfrak{d}_{i,k}} : 1 \leq i < \omega, k < n \}$ and $f_{\mathfrak{c}_N^\star} = \bigcup \{ f_{\mathfrak{d}_{i,k}} : N \leq i < \omega, m \leq k < n \}$ for $N \geq 2$.
- (3) $\Sigma_\star(\mathfrak{c}_1^\star)$ is the smallest family satisfying the following.
 - (i) $\langle \mathfrak{c}_1^\star \rangle \in \Sigma_\star(\mathfrak{c}_1^\star)$.
 - (ii) Whenever $j > 1$ is a power of 2 and $\langle \mathfrak{d}'_{i,k} : i < j, m \leq k < n \rangle$, $\langle \bar{\mathfrak{f}}_{i,k} : i < j, m \leq k < n \rangle$ and $\langle \bar{\mathfrak{g}}_k : k < m \rangle$ satisfy (a)-(d) below, we have, under appropriate order

$$\bigcup \{ \bar{\mathfrak{g}}_k : k < m \} \cup \bigcup \{ \bar{\mathfrak{f}}_{i,k} : i < j, m \leq k < n \} \cup \{ \mathfrak{c}_j^\star \} \in \Sigma_\star(\mathfrak{c}_1^\star)$$

- (a) $\mathfrak{d}'_{i,k} \in CR_p$ and $\text{dom}(\mathfrak{d}'_{i,k}) = \text{dom}(\mathfrak{d}_{i,k})$.
- (b) $|\{ i \in [j_1, j_2] : (\exists k \in [m, n]) (\mathfrak{d}'_{i,k} \neq \mathfrak{d}_{i,k}) \}| \leq (j_2 - j_1) / \log_2(j_1)$ for every $2 \leq j_1 < j_2 \leq j$ where j_1, j_2 are powers of 2.
- (c) $\bar{\mathfrak{f}}_{i,k} \in \Sigma(\mathfrak{d}'_{i,k})$.
- (d) $\bar{\mathfrak{g}}_k \in \Sigma(\mathfrak{d}_k)$.

- (4) For $N \geq 2$, $\Sigma_\star(\mathfrak{c}_N^\star)$ is the smallest family satisfying the following.

- (i) $\langle \mathfrak{c}_N^\star \rangle \in \Sigma_\star(\mathfrak{c}_N^\star)$.
- (ii) Whenever $j > N$ is a power of 2 and $\langle \mathfrak{d}'_{i,k} : N \leq i < j, m \leq k < n \rangle$ and $\langle \bar{\mathfrak{f}}_{i,k} : N \leq i < j, m \leq k < n \rangle$ satisfy (a)-(c) below, we have, under appropriate order

$$\bigcup \{ \bar{\mathfrak{f}}_{i,k} : i < j, m \leq k < n \} \cup \{ \mathfrak{c}_j^\star \} \in \Sigma_\star(\mathfrak{c}_N^\star)$$

- (a) $\mathfrak{d}'_{i,k} \in CR_p$ and $\text{dom}(\mathfrak{d}'_{i,k}) = \text{dom}(\mathfrak{d}_{i,k})$.
- (b) $|\{ i \in [j_1, j_2] : (\exists k \in [m, n]) (\mathfrak{d}'_{i,k} \neq \mathfrak{d}_{i,k}) \}| \leq (j_2 - j_1) / \log_2(j_1)$ for every $N \leq j_1 < j_2 \leq j$ where j_1, j_2 are powers of 2.
- (c) $\bar{\mathfrak{f}}_{i,k} \in \Sigma(\mathfrak{d}'_{i,k})$.

$$(5) \text{ pos}(\mathbf{c}_N^*) = \{\bigcup_{k < K} f_{c_k} : \langle \mathbf{c}_k : k < K \rangle \in \Sigma_*(\mathbf{c}_N^*)\}.$$

Let (CR'_p, Σ'_p) be the partial \aleph_1 -CP at $\delta+1$ such that $CR'_p = CR_p \cup \{\mathbf{c}_N^* : N \geq 1 \text{ is a power of } 2\}$ with $\text{dom}(\mathbf{c}_N^*)$, $\text{pos}(\mathbf{c}_N^*)$ and $f_{c_N^*}$ as above, $\Sigma'_p \upharpoonright CR_p = \Sigma_p$ and $\Sigma'_p(\mathbf{c}_N^*) = \Sigma_*(\mathbf{c}_N^*)$. We say that (CR'_p, Σ'_p) is the result of adding the countable join $\mathbf{c}_1 = \bigoplus_{i \geq 1} \bar{\mathbf{d}}_i$ of $\langle \bar{\mathbf{d}}_i : i \geq 1 \rangle$ to (Σ_p, CR_p) .

Note that (CR'_p, Σ'_p) is indeed a partial \aleph_1 -CP at $\delta + 1$ because Σ'_p satisfies transitivity, cuts and finite joins.

Lemma 3.3. *Let (CR'_p, Σ'_p) be as in Definition ???. Let (CR, Σ) be an \aleph_1 -CP such that $CR'_p = \{\mathbf{c} \in CR : \text{dom}(\mathbf{c}) \subseteq \delta\}$ and $\Sigma'_p = \Sigma \upharpoonright CR'_p$. Let $\mathbb{Q} = \mathbb{Q}_{CR, \Sigma}$, $p = \{\mathbf{c}_1^* = \bigoplus_{i \geq 1} \bar{\mathbf{d}}_i\}$ and $p_i = \{\bar{\mathbf{d}}_{i,k} : k < n\}$. Then*

$$p \Vdash_{\mathbb{Q}} \lim_j \frac{|\{i < j : p_i \in G_{\mathbb{Q}}\}|}{j} = 1$$

Proof of Lemma ???: It suffices to show that for every $p_1 \geq p$ and $j_* \geq 2^{10}$ there exists $p_2 \geq p_1$ such that

$$p_2 \Vdash_{\mathbb{Q}} \frac{|\{i < j_* : p_i \in G_{\mathbb{Q}}\}|}{j_*} > 1 - \frac{8}{\log_2 j_*}$$

Since $p_1 \geq p = \{\mathbf{c}_1^*\}$, we can find $p_2 \geq p_1$ and $j_0 > j_*$, such that j_0 is a power of 2 and

$$\bigcup \{\bar{\mathbf{g}}_k : k < m\} \cup \bigcup \{\bar{\mathbf{f}}_{i,k} : i < j_0, m \leq k < n\} \cup \{\mathbf{c}_{j_0}^*\} \subseteq p_2$$

where $\langle \bar{\mathbf{d}}'_{i,k} : i < j_0, m \leq k < n \rangle$, $\langle \bar{\mathbf{f}}_{i,k} : i < j_0, m \leq k < n \rangle$ and $\langle \bar{\mathbf{g}}_k : k < m \rangle$ are as in Definition ???(3)(ii).

Choose $N \geq 10$ such that $2^N \leq j_* < 2^{N+1}$. Then p_2 forces that

$$\frac{|\{i < j_* : p_i \in G_{\mathbb{Q}}\}|}{j_*} \geq 1 - \left(\sum_{1 \leq j < N} \frac{2^{j+1} - 2^j}{j j_*} \right) - \frac{2^{N+1} - 2^N}{N j_*} \geq 1 - \left(\sum_{1 \leq j < N} \frac{1}{j 2^{N-j}} \right) - \frac{1}{N}$$

Since $\sum_{1 \leq j < N/2} 1/(j 2^{N-j}) \leq N/2^{N/2} \leq 4/N$ (as $N \geq 10$) and $\sum_{N/2 \leq j < N} 1/(j 2^{N-j}) \leq 2/N$, it follows that

$$p_2 \Vdash_{\mathbb{Q}} \frac{|\{i < j_* : p_i \in G_{\mathbb{Q}}\}|}{j_*} \geq 1 - \left(\frac{4}{N} + \frac{2}{N} + \frac{1}{N} \right) > 1 - \frac{8}{N}$$

□

Definition 3.4. (CR, Σ) is a thin \aleph_1 -CP if (CR, Σ) is an \aleph_1 -CP and there exist S and $\langle \mathbf{c}_\delta : \delta \in S \rangle$ such that the following hold.

(a) $S \subseteq \bigcup_{k \geq 1} S_k$.

(b) $\mathbf{c}_\delta \in CR$.

(c) For every $k_* \geq 1$ and $\delta \in S \cap S_{k_*}$, letting (CR_p, Σ_p) be the partial \aleph_1 -CP at δ satisfying $CR_p = CR \upharpoonright \delta = \{\mathbf{c} \in CR : \sup(\text{dom}(\mathbf{c})) < \delta\}$ and $\Sigma_p = \Sigma \upharpoonright CR_p$, there exists a joinable candidate $\langle \bar{\mathfrak{d}}_i : i \geq 1 \rangle$ for (CR_p, Σ_p) at δ such that

(i) $\mathbf{c}_\delta = \bigoplus_{i \geq 1} \bar{\mathfrak{d}}_i$ and

(ii) $CR'_p = \{\mathbf{c} \in CR : \text{dom}(\mathbf{c}) \subseteq \delta\}$ and $\Sigma'_p = \Sigma \upharpoonright CR'_p$ where (CR'_p, Σ'_p) is the result of adding $\bigoplus_{i \geq 1} \bar{\mathfrak{d}}_i$ to (CR_p, Σ_p) .

(d) $\mathbf{c} \in CR$ iff \mathbf{c} is a finite join of $\{\mathfrak{d} \in CR : \mathfrak{d} \text{ is finite}\} \cup \bigcup \{\Sigma(\mathbf{c}_\delta) : \delta \in S\}$.

Claim 3.5. Suppose (CR, Σ) is an \aleph_1 -CP as witnessed by $S, \langle \mathbf{c}_\delta : \delta \in S \rangle$. Suppose $\mathbf{c} \in CR$, $k_* \geq 1$, $\delta \in S_{k_*}$, $\text{dom}(\mathbf{c})$ is an unbounded subset of δ . Then there exist $\bar{\mathbf{c}} = \langle \mathbf{c}_k : k \leq k_1 \rangle \in \Sigma(\mathbf{c})$ and $\bar{\mathfrak{d}} = \langle \mathfrak{d}_k : k \leq k_2 \rangle \in \Sigma(\mathbf{c}_\delta)$ such that $\mathbf{c}_{k_1} = \mathfrak{d}_{k_2}$.

Proof of Claim ???: Easily follows from Definition ???. □

4 Guided products

Definition 4.1. Suppose $\langle \mathbb{Q}_\alpha : \alpha < \omega_2 \rangle$ and $\langle p_\delta^* : \delta \in S_{\aleph_0}^{\omega_2} \rangle$ satisfy the following.

(i) $\mathbb{Q}_\alpha = \mathbb{Q}_{CR_\alpha, \Sigma_\alpha}$ where $(CR_\alpha, \Sigma_\alpha)$ is a thin \aleph_1 -CP.

(ii) p_δ^* is a function whose domain is a countable unbounded subset of δ and for every $\alpha \in \text{dom}(p_\delta^*)$, $p_\delta^*(\alpha) \in \mathbb{Q}_\alpha$.

For $\gamma \leq \omega_2$, define a forcing \mathbb{P}_γ as follows.

(1) $p \in \mathbb{P}_\gamma$ iff

(a) p is a function, $\text{dom}(p)$ is a countable subset of γ ,

(b) for every $\alpha \in \text{dom}(p)$, $p(\alpha) \in \mathbb{Q}_\alpha$ and

(c) for every $\delta \leq \gamma$ with $\text{cf}(\delta) = \aleph_0$, if $\text{dom}(p) \cap \delta$ is unbounded in δ , then for some $\eta < \delta$, $p \upharpoonright (\eta, \delta) = p_\delta^* \upharpoonright (\eta, \delta)$.

(2) For $p, q \in \mathbb{P}_\gamma$, define $p \leq q$ iff $\text{dom}(p) \subseteq \text{dom}(q)$ and for every $\alpha \in \text{dom}(p)$, $p(\alpha) \leq_{\mathbb{Q}_\alpha} q(\alpha)$.

We say that \mathbb{P}_{ω_2} is the countable support product of $\langle \mathbb{Q}_\alpha : \alpha < \omega_2 \rangle$ guided by $\langle p_\delta^* : \delta \in S_{\aleph_0}^{\omega_2} \rangle$. Note that for $\text{cf}(\gamma) = \aleph_1$, \mathbb{P}_γ is completely determined by $\langle \mathbb{Q}_\alpha : \alpha < \gamma \rangle$ and $\langle p_\delta^* : \delta < \gamma, \text{cf}(\delta) = \aleph_0 \rangle$.

Claim 4.2. Let $\langle \mathbb{Q}_\alpha : \alpha < \omega_2 \rangle$, $\langle p_\delta^* : \delta \in S_{\aleph_0}^{\omega_2} \rangle$ and \mathbb{P}_γ for $\gamma \leq \omega_2$ be as in Definition ???. Then the following hold.

(a) $\mathbb{P}_{\gamma+1} = \mathbb{P}_\gamma \times \mathbb{Q}_\gamma$.

(b) \mathbb{P}_γ satisfies ccc.

Proof of Claim ??: (a) is obvious from the definition of \mathbb{P}_γ . (b) follows from Claim ?? below and the fact that each \mathbb{Q}_α has \aleph_1 as a precaliber (Claim ??).

Claim 4.3. *Suppose $\gamma \leq \omega_2$ and $\langle p_i : i < \omega_1 \rangle$ is a sequence of conditions in \mathbb{P}_γ . Then there exists $X \in [\omega_1]^{\aleph_1}$ and a finite $F \subseteq \omega_2$ such that for every $\alpha \in \omega_2 \setminus F$, if there are $i < j$ in X such that $\alpha \in \text{dom}(p_i) \cap \text{dom}(p_j)$, then $(\forall i \in X)(\alpha \in \text{dom}(p_i) \text{ and } p_i(\alpha) \text{ does not depend on } i \in X)$.*

Proof of Claim ??: By induction on $\gamma \leq \omega_2$. If γ is a successor or $\gamma = \omega_2$, this is trivial.

Suppose $\text{cf}(\gamma) = \aleph_0$ and let $\langle p_i : i < \omega_2 \rangle$ be a sequence of conditions in \mathbb{P}_γ . Let $\langle \gamma_n : n < \omega \rangle$ be increasing cofinal in γ . For each $i < \omega_1$, choose $n = n_i < \omega$ such that either $p_i \in \mathbb{P}_{\gamma_n}$ or $p_i \upharpoonright (\gamma_n, \gamma) = p_\gamma^* \upharpoonright (\gamma_n, \gamma)$. Choose $X \in [\omega_1]^{\aleph_1}$ and $n_* < \omega$, such that $(\forall i \in X)(n_i = n_*)$ and apply the inductive hypothesis to $\langle p_i \upharpoonright \gamma_{n_*} : i \in X \rangle$.

Next suppose $\text{cf}(\gamma) = \omega_1$ and let $\langle p_i : i < \omega_2 \rangle$ be a sequence of conditions in \mathbb{P}_γ . Choose $\langle \gamma_i : i < \omega_1 \rangle$ continuously increasing and cofinal in γ such that $\text{cf}(\gamma_i) = \aleph_0$ for every $i < \omega_1$. For each $i < \omega_1$, choose $j = j_i < i$ such that either $p_i \upharpoonright \gamma_j \in \mathbb{P}_{\gamma_j}$ or $p_i \upharpoonright (\gamma_j, \gamma) = p_\gamma^* \upharpoonright (\gamma_j, \gamma)$. By Fodor's lemma we can get $S \in [\omega_1]^{\aleph_1}$ and $j_* < \omega_1$ such that $(\forall i \in S)(j_i = j_*)$. Choose $X \in [S]^{\aleph_1}$ such that for every $i < j$ in X , $\text{dom}(p_i) \cap \text{dom}(p_j) \subseteq \gamma_{j_*}$. Now apply the inductive hypothesis to $\langle p_i \upharpoonright \gamma_{j_*} : i \in S \rangle$. \square

Lemma 4.4. *Let $\langle \mathbb{Q}_\alpha : \alpha < \omega_2 \rangle$, $\langle p_\delta^* : \delta \in S_{\aleph_0}^{\omega_2} \rangle$ and \mathbb{P}_{ω_2} be as in Definition ??. Then $V^{\mathbb{P}_{\omega_2}} \models \neg \clubsuit^1$.*

Proof of Lemma ??: Towards a contradiction, suppose $p_0 \in \mathbb{P}_{\omega_2}$, $\langle \dot{A}_\delta = \{\dot{\alpha}_{\delta,n} : n < \omega\} : \delta \in \bullet \text{Lim}(\omega_1) \rangle \in V^{\mathbb{P}_{\omega_2}}$ are such that $p_0 \Vdash$ “ $(\forall \delta \in \bullet \text{Lim}(\omega_1))(\{\dot{\alpha}_{\delta,n} : n < \omega\}$ is increasing cofinal in δ) and $\langle \dot{A}_\delta : \delta < \omega_1 \rangle$ is a \clubsuit^1 witnessing sequence”. Since \mathbb{P}_{ω_2} satisfies ccc, we can find $\gamma < \omega_2$ such that $p_0 \in \mathbb{P}_\gamma$ and each $\dot{\alpha}_{\delta,n}$ is a \mathbb{P}_γ -name.

Let $\dot{X} = \{\alpha < \omega_1 : f_{\mathbb{Q}_\gamma}^\alpha = 1\}$. Then $\dot{X} \in V^{\mathbb{P}_{\gamma+1}}$ and $V^{\mathbb{P}_{\gamma+1}} \models \dot{X} \in [\omega_1]^{\aleph_1}$. So there exist $p_1 \in \mathbb{P}_\gamma$, $q \in \mathbb{Q}_\gamma$, $\delta \in \bullet \text{Lim}(\omega_1)$ and $n_* < \omega$ such that $p_1 \geq p_0$ and $(p_1, q) \Vdash_{\mathbb{P}_{\gamma+1}} (\forall n \geq n_*)(\dot{\alpha}_{\delta,n} \in \dot{X})$. Note that we must have that $\text{dom}(q) \cap \delta$ is unbounded in δ otherwise we can easily extend (p_1, q) to get a contradiction. By possibly extending q , by Definition ??(D)(ii), we can assume that $q = \{\mathbf{c}_k : k < K_*\}$ where $\text{dom}(\mathbf{c}_k) < \text{dom}(\mathbf{c}_{k+1})$ for every $k < K_* - 1$ and for some $K < K_*$, $\text{dom}(\mathbf{c}_K)$ is an unbounded subset of δ . Let S_γ and $\langle \mathbf{c}_{\gamma,\delta} : \delta \in S_\gamma \rangle$ witness that $(\text{CR}_\gamma, \Sigma_\gamma)$ is a thin \aleph_1 -CP. By Claim ??, we can further assume that $\mathbf{c}_K = \mathbf{c}'_{K'}$ for some $\langle \mathbf{c}'_n : n \leq K' \rangle \in \Sigma(\mathbf{c}_{\gamma,\delta})$.

Let $m < n < \omega$ and $\bar{\mathbf{d}}_i = \langle \mathbf{d}_{i,k} : k < n \rangle$ for $i \geq 1$ be as in Definition ?? and $\mathbf{c}_{\gamma,\delta} = \bigoplus_{i \geq 1} \bar{\mathbf{d}}_i$. Then as $\langle \mathbf{c}'_n : n \leq K' \rangle \in \Sigma(\mathbf{c}_{\gamma,\delta})$, we can find $N \geq 1$ a power of 2 such that $\mathbf{c}_K = \mathbf{c}'_{K'} = \mathbf{c}_N^*$ in the notation of Definition ??.

Choose $p_2 \in \mathbb{P}_\gamma$, $p_2 \geq p_1$, $n(1) > n_*$ and $\alpha > \min(\text{dom}(\mathbf{c}_N^*))$ such that $p_2 \Vdash_{\mathbb{P}_\gamma} \dot{\alpha}_{\delta,n(1)} = \alpha$. We can assume that $\alpha \in \text{dom}(\mathbf{c}_N^*)$ - Otherwise letting \mathbf{c}_* be a creature with domain $\{\alpha\}$ and $f_{\mathbf{c}_*}(\alpha) = 1$, we have $q' = q \cup \{\mathbf{c}_*\} \Vdash_{\mathbb{Q}_\gamma} \alpha \notin \dot{X}$ so that (p_2, q') forces a contradiction.

Choose $i(\star) \geq N$ and $m \leq k(\star) < n$ such that $\alpha \in \text{dom}(\mathfrak{d}_{i(\star),k(\star)})$. Let $N_1 \geq N$ be the largest power of 2 such that $i(\star) \geq N_1$ and let $j > i(\star)$ be a power of 2. Choose a creature $\mathfrak{d}'_{i(\star),k(\star)}$ such that $\text{dom}(\mathfrak{d}'_{i(\star),k(\star)}) = \text{dom}(\mathfrak{d}_{i(\star),k(\star)})$ and $\bar{f} \in \Sigma(\mathfrak{d}'_{i(\star),k(\star)})$ such that for some finite $\mathbf{c}_\star \in \bar{f}$, $\text{dom}(\mathbf{c}_\star) = \{\alpha\}$ and $f_{\mathbf{c}_\star}(\alpha) = 0$. It follows that, under appropriate order

$$\{\mathfrak{d}_{i,k} : N \leq i < j, m \leq k < n, (i, k) \neq (i(\star), k(\star))\} \cup \bar{f} \cup \{\mathbf{c}_j^\star\} \in \Sigma(\mathbf{c}_N^\star)$$

Let $q' = (q \setminus \{\mathbf{c}_K\}) \cup \{\mathfrak{d}_{i,k} : N \leq i < j, m \leq k < n, (i, k) \neq (i(\star), k(\star))\} \cup \bar{f} \cup \{\mathbf{c}_j^\star\}$. Then $(p_2, q') \geq (p, q)$ and $q' \Vdash_{\mathbb{Q}_\gamma} \alpha \notin \dot{X}$ - Contradiction. \square

5 \clubsuit^{lim} and $\neg\clubsuit^1$

We define a preparatory forcing \mathbb{R} which generically adds $\langle \mathbb{Q}_\alpha : \alpha < \omega_2 \rangle$ and $\langle p_\delta^\star : \delta \in S_{\aleph_0}^{\omega_2} \rangle$ satisfying Definition ??(i)-(ii) using countable approximations. This ensures that the resulting guided product \mathbb{P}_{ω_2} preserves a \clubsuit^{lim} witnessing sequence \bar{A} which is also added by \mathbb{R} via countable approximations.

Definition 5.1. Let \mathbb{R} be a forcing whose conditions are $r = (u_r, \delta_r, \langle \mathbb{Q}_{r,\alpha} : \alpha \in u_r \rangle, v_r, \langle p_{r,\alpha}^\star : \alpha \in v_r \rangle, \bar{A}_r)$ where

- (a) $u_r \in [\omega_2]^{\leq \aleph_0}$, $\delta_r < \omega_1$,
- (b) $\mathbb{Q}_{r,\alpha} = \bigcup_{\xi < \delta_r} (\mathbb{Q}_{\mathcal{C}R_{r,\alpha}, \Sigma_{r,\alpha}} \upharpoonright \xi)$ for some thin \aleph_1 -CP $(\mathcal{C}R_{r,\alpha}, \Sigma_{r,\alpha})$ as witnessed by $(S_{r,\alpha}, \langle \mathbf{c}_{r,\alpha,\delta} : \delta \in S_{r,\alpha} \rangle)$ - So only $S_{r,\alpha} \cap \delta_r$ and $\langle \mathbf{c}_{r,\alpha,\delta} : \delta \in S_{r,\alpha} \cap \delta_r \rangle$ are relevant,
- (c) $v_r \subseteq u \cap \bullet\text{Lim}(\omega_1)$ and for every $\alpha \in v_r$, $u_r \cap \alpha$ is unbounded in α ,
- (d) $p_{r,\alpha}^\star$ is a function with domain an unbounded subset of $u_r \cap \alpha$ and for each $\xi \in \text{dom}(p_{r,\alpha}^\star)$, $p_{r,\alpha}^\star(\xi) \in \mathbb{Q}_{r,\alpha}$ and
- (e) $\bar{A}_r = \langle A_{r,\gamma} : \gamma \in \bullet\text{Lim}(\omega_1) \cap \delta_r \rangle$ where each $A_{r,\gamma}$ is an unbounded subset of γ of order type ω .

For $r, s \in \mathbb{R}$, define $r \leq s$ iff the following hold.

- (i) $u_r \subseteq u_s$, $\delta_r \leq \delta_s$.
- (ii) For every $\alpha \in u_r$, $S_{r,\alpha} \cap \delta_r = S_{s,\alpha} \cap \delta_r$ and $\mathbf{c}_{r,\alpha,\delta} = \mathbf{c}_{s,\alpha,\delta}$ for every $\delta \in S_r \cap \delta_r$. It follows that $\mathbb{Q}_{r,\alpha} \subseteq \mathbb{Q}_{s,\alpha}$ and for every $p \in \mathbb{Q}_{s,\alpha}$, if $\text{dom}(p)$ is bounded below δ_r , then $p \in \mathbb{Q}_{r,\alpha}$.
- (iii) $v_r \subseteq v_s$ and for every $\alpha \in v_r$, $p_{r,\alpha}^\star = p_{s,\alpha}^\star$.
- (iv) $\bar{A}_r = \bar{A}_s \upharpoonright (\bullet\text{Lim}(\omega_1) \cap \delta_r)$.

Claim 5.2. \mathbb{R} is countably closed and hence it preserves stationary subsets of ω_1 . Under CH, it satisfies \aleph_2 -c.c. and therefore preserves all cofinalities.

Proof of Claim ???: It is clear that \mathbb{R} is countably closed. Next let $\{r_i : i < \omega_2\} \subseteq \mathbb{R}$. Using CH, we can find $X_0 \in [\omega_2]^{\aleph_2}$ such that $\langle u_{r_i} : i \in X_0 \rangle$ forms a Δ -system with root u_\star . By possibly extending each r_i , we can assume that $u_{r_i} \setminus u_\star \neq \emptyset$ for every $i \in X_0$. Choose $X \in [X_0]^{\aleph_2}$ such that the following hold.

- (i) For every $i, j \in X$ with $i < j$, $\sup(u_\star) < \min(u_{r_i} \setminus u_\star) \leq \sup(u_{r_j} \setminus u_\star) < \inf(u_{r_j} \setminus u_\star)$.
- (ii) $\langle v_{r_i} : i \in X \rangle$ forms a Δ -system with root $v_\star \subseteq u_\star$.
- (iii) $\delta_{r_i} = \delta_\star$ does not depend on $i \in X$.
- (iv) For every $\alpha \in u_\star$, $\mathbb{Q}_{r_i, \alpha} = \mathbb{Q}_\alpha$ does not depend on $i \in X$.
- (v) For every $\alpha \in v_\star$, $p_{r_i, \alpha}^\star = p_\alpha^\star$ does not depend on $i \in X$.
- (vi) $\bar{A}_{r_i} = \bar{A}_\star$ does not depend on $i \in X$.

For clauses (iv), (v) and (vi), we use CH. It is clear that any two conditions in $\{r_i : i \in X\}$ have a common extension. \square

From now on we assume CH. The next claim is easily verified.

Claim 5.3. *Each of the following sets is dense in \mathbb{R} .*

- (a) $\{r \in \mathbb{R} : \alpha \in u_r\}$ for $\alpha < \omega_2$.
- (b) $\{r \in \mathbb{R} : \delta_r > \delta\}$ for $\delta < \omega_1$.
- (c) $\{r \in \mathbb{R} : \delta \in v_r\}$ for $\delta \in S_{\aleph_0}^{\omega_2}$.

Let $G_{\mathbb{R}}$ be \mathbb{R} -generic over V . Work in $V_1 = V[G_{\mathbb{R}}]$. For $\alpha < \omega_2$, define $\mathbb{Q}_\alpha = \bigcup\{\mathbb{Q}_{r, \alpha} : r \in G_{\mathbb{R}}, \alpha \in u_r\}$. Note that for every $\alpha < \omega_2$, $S_\alpha = \bigcup\{S_{r, \alpha} \cap \delta_r : r \in G_{\mathbb{R}}, \alpha \in u_r\}$ is a stationary subset of $\bigcup_{k \geq 1} S_k$ and $V_1 \models “(\forall \alpha < \omega_2)(\mathbb{Q}_\alpha = \mathbb{Q}_{\text{CR}_\alpha, \Sigma_\alpha} \text{ for some thin } \aleph_1\text{-CP } (\text{CR}_\alpha, \Sigma_\alpha))”$. For $\delta \in S_{\aleph_0}^{\omega_2}$, let $p_\delta^\star = p_{r, \delta}^\star$ for some $r \in G_{\mathbb{R}}$ with $\delta \in v_r$. Let $\bar{A} = \langle A_\delta : \delta \in \bullet \text{Lim}(\omega_1) \rangle = \bigcup\{A_r : r \in G_{\mathbb{R}}\}$. Let $\{\alpha_{\delta, n} : n < \omega\}$ list A_δ in increasing order.

Let $\mathbb{P}_{\omega_2} \in V_1$ be the countable support product of $\langle \mathbb{Q}_\alpha : \alpha < \omega_2 \rangle$ guided by $\langle p_\delta^\star : \delta \in S_{\aleph_0}^{\omega_2} \rangle$. Note that, since \mathbb{R} is countably closed, the set of conditions $(r, p) \in \mathbb{R} \star \mathbb{P}_{\omega_2}$ satisfying the following is dense in $\mathbb{R} \star \mathbb{P}_{\omega_2}$.

- (a) p is an actual object.
- (b) $\text{dom}(p) \subseteq u_r$.
- (c) $(\forall \alpha \in \text{dom}(p))(p(\alpha) \in \mathbb{Q}_{r, \alpha})$.
- (d) For every $\alpha < \omega_2$ of cofinality \aleph_0 , if $\text{dom}(p) \cap \alpha$ is unbounded in α , then $\alpha \in v_r$.

So we can assume that our conditions in $\mathbb{R} \star \mathbb{P}_{\omega_2}$ have this form.

Theorem 5.4. $V_1^{\mathbb{P}_{\omega_2}} \models \clubsuit^{\text{lim}} \wedge \neg \clubsuit^1$

Proof of Theorem ???: That $V_1^{\mathbb{P}_{\omega_2}} \models \neg \clubsuit^1$ follows from Lemma ??. We'll show that \bar{A} witnesses \clubsuit^{lim} in $V_1^{\mathbb{P}_{\omega_2}}$. Suppose $(r_*, p_*) \Vdash_{\mathbb{R} \star \mathbb{P}_{\omega_2}} \dot{A} \in [\omega_1]^{\aleph_1}$. We'll construct $(r, p) \geq (r_*, p_*)$ and $\delta < \omega_1$ such that

$$(r, p) \Vdash_{\mathbb{R} \star \mathbb{P}_{\omega_2}} \lim_n \frac{|\{k < n : \dot{\alpha}_{\delta, k} \in \dot{A}\}|}{n} = 1$$

Choose $\langle (r_i, p_i, \gamma_i) : i < \omega_1 \rangle$ such that the following hold.

- (i) $(r_i, p_i) \geq (r_*, p_*)$.
- (ii) For all $i < j < \omega_1$, $r_i \leq_{\mathbb{R}} r_j$, $\text{sup}(u_{r_i}) < \text{sup}(u_{r_j})$ and $i \leq \delta_{r_i} < \delta_{r_j}$.
- (iii) For every $i < \omega_1$, $\text{sup}(\bigcup_{j < i} \text{dom}(p_j)) < \text{sup}(\text{dom}(p_i))$.
- (iv) For every $i < \omega_1$ $i \in u_i$ and for every $\alpha < \text{sup}(u_{r_i})$, there exists $j \in (i, \omega_1)$ such that $\alpha \in u_{r_j}$. So $\bigcup_{i < \omega_1} u_{r_i} = \alpha_* \in [\omega_1, \omega_2)$ and $\text{cf}(\alpha_*) = \aleph_1$.
- (v) For every $\delta < \alpha_*$ with $\text{cf}(\delta) = \aleph_0$, there exists $i < \omega_1$ such that $\delta \in v_{r_i}$. Hence $\bigcup_{i < \omega_1} v_{r_i} = \{\delta < \alpha_* : \text{cf}(\delta) = \aleph_0\}$.
- (vi) $\langle \gamma_i : i < \omega_1 \rangle$ is a strictly increasing sequence in ω_1 .
- (vii) $(r_i, p_i) \Vdash \gamma_i \in \dot{A}$.

Claim 5.5. *There exist $F \subseteq \omega_2$ finite and $X \in [\omega_1]^{\aleph_1}$ such that for every $\alpha \in \omega_2 \setminus F$, if $\alpha \in \text{dom}(p_i) \cap \text{dom}(p_j)$ for some $i < j$ in X , then $(\forall i \in X)(\alpha \in \text{dom}(p_i) \text{ and } p_i(\alpha) \text{ does not depend on } i \in X)$.*

Proof of Claim ??: For $\alpha < \alpha_*$, let $\mathbb{Q}'_{\alpha} = \bigcup \{\mathbb{Q}_{r_i, \alpha} : i < \omega_1, \alpha \in u_{r_i}\}$. Then \mathbb{Q}'_{α} is a thin \aleph_1 -CP. For $\delta < \alpha_*$ with $\text{cf}(\delta) = \aleph_0$, let $p_{\delta}^* = p_{r_i, \delta}^*$ where $i < \omega_1$ and $\delta \in v_{r_i}$. Let \mathbb{P}_{α_*} be the countable support product of $\langle \mathbb{Q}'_{\alpha} : \alpha < \alpha_* \rangle$ guided by $\langle p_{\delta}^* : \delta < \alpha_*, \text{cf}(\delta) = \aleph_0 \rangle$ so that each $p_i \in \mathbb{P}_{\alpha_*}$. Now apply Claim ??. \square

By shrinking X and F , we can assume that for every $i \in X$, $F \subseteq \text{dom}(p_i)$. Let $W = \bigcap_{i \in X} (\text{dom}(p_i) \setminus F)$ and $Y_i = \text{dom}(p_i) \setminus (F \cup W)$. Then $\langle Y_i : i \in X \rangle$ is a sequence of pairwise disjoint non empty countable sets. By shrinking X , we can also assume that for every $i < j$ in X , $\text{sup}(Y_i) < \min(Y_j)$ and $\text{otp}(\text{dom}(p_i))$ does not depend on $i \in X$.

By Claim ??, we can find $X_1 \in [X]^{\aleph_1}$ such that for every $\alpha \in F$ exactly one of the following holds.

- (A) For every $i \in X_1$, $p_i(\alpha) = q_{\alpha}$ does not depend on i .

- (B) There are $m = m_\alpha$, $n = n_\alpha$, $m < n < \omega$ and $\langle q_{i,\alpha} : i \in X_1 \rangle$ such that for every $i \in X_1$,
- (i) $q_{i,\alpha} \in \mathbb{Q}_{r_i,\alpha}$, $\text{dom}(q_{i,\alpha}) = \text{dom}(p_i(\alpha))$ and $r_i \Vdash_{\mathbb{R}} p_i(\alpha) \leq_{\mathbb{Q}_\alpha} q_{i,\alpha}$,
 - (ii) $q_{i,\alpha} = \{\mathfrak{d}_{i,\alpha,k} : k < n\}$ and for every $k < n - 1$, $\text{dom}(\mathfrak{d}_{i,\alpha,k}) < \text{dom}(\mathfrak{d}_{i,\alpha,k+1})$,
 - (iii) for every $k < m$, $\mathfrak{d}_{i,\alpha,k} = \mathfrak{d}_{\alpha,k}$ does not depend on $i \in X_1$,
 - (iv) for every $j < j'$ in X , $\text{dom}(\mathfrak{d}_{j,\alpha,n-1}) < \text{dom}(\mathfrak{d}_{j',\alpha,m})$ and
 - (v) $\text{otp}(\mathfrak{d}_{i,\alpha,k}) = \theta_{\alpha,k}$ does not depend on $i \in X_1$ and $1 \leq k_\alpha < \omega$ is such that $\theta_{\alpha,k} < \omega^{k_\alpha}$.

Let F_0 be the set of $\alpha \in F$ for which case (A) holds and $F_1 = F \setminus F_0$.

By reindexing, we can assume that $X_1 = \omega_1$. Let $k_\star = \max(\{k_\alpha + 2 : \alpha \in F\})$. Put $Y = \bigcup_{i < \omega_1} Y_i$. Choose a club $E \subseteq \omega_1$ such that for every $\delta \in E$, the following hold.

- (a) For every $i < \delta$, there exists $j < \delta$ such that $\text{sup}(u_{r_i} \cap Y) < \text{sup}(Y_j)$.
- (b) $\text{sup}(\{\delta_{r_i} : i < \delta\}) = \delta$.
- (c) For every $\alpha \in F_1$, $\text{sup}(\{\text{dom}(q_i(\alpha)) : i < \delta\}) = \delta$.
- (d) $\text{sup}(\{\gamma_i : i < \delta\}) = \delta$.

Fix $\delta \in S_{k_\star} \cap E$ and let $\langle i(n) : n < \omega \rangle$ be increasing cofinal in δ . Let $\alpha_\star = \text{sup}(\{Y_{i(n)} : n < \omega\})$. We can assume that $\alpha_\star \notin F \cup W$ - Just pick a sufficiently large $\delta \in S_{k_\star} \cap E$. Define $r \in \mathbb{R}$ as follows.

- (a) $u_r = \bigcup_{n < \omega} u_{r_{i(n)}} \cup \{\alpha_\star\}$, $\delta_r = \delta + 1$.
- (b) For $\alpha \in u_r$, choose $\mathbb{Q}_{r,\alpha}$, $(\text{CR}_{r,\alpha}, \Sigma_{r,\alpha})$ and $(S_{r,\alpha}, \langle \mathfrak{c}_{r,\alpha,\delta} : \delta \in S_{r,\alpha} \rangle)$ as follows.
 - (i) If $\alpha \in u_r \setminus (F_1 \cup \{\alpha_\star\})$, choose a thin \aleph_1 -CP $(\text{CR}_{r,\alpha}, \Sigma_{r,\alpha})$ with witnessing pair $(S_{r,\alpha}, \langle \mathfrak{c}_{r,\alpha,\delta} : \delta \in S_{r,\alpha} \rangle)$ such that for every $n < \omega$, $S_{r,\alpha} \cap \delta_{i(n)} = S_{r_{i(n)},\alpha} \cap \delta_{i(n)}$ and $\mathfrak{c}_{r,\alpha,\delta} = \mathfrak{c}_{r_{i(n)},\alpha,\delta}$ for every $\delta \in S_{r,\alpha} \cap \delta_{i(n)}$. So $\bigcup_{n < \omega} \mathbb{Q}_{r_{i(n)},\alpha} \subseteq \mathbb{Q}_{r,\alpha} = \mathbb{Q}_{\text{CR}_{r,\alpha}, \Sigma_{r,\alpha}} \upharpoonright \delta$.
 - (ii) If $\alpha = \alpha_\star$, choose $\mathbb{Q}_{r,\alpha}$, $(\text{CR}_{r,\alpha}, \Sigma_{r,\alpha})$ and $(S_{r,\alpha}, \langle \mathfrak{c}_{r,\alpha,\delta} : \delta \in S_{r,\alpha} \rangle)$ arbitrarily.
 - (iii) If $\alpha \in F_1$, choose a thin \aleph_1 -CP $(\text{CR}_{r,\alpha}, \Sigma_{r,\alpha})$ with witnessing pair $(S_{r,\alpha}, \langle \mathfrak{c}_{r,\alpha,\delta} : \delta \in S_{r,\alpha} \rangle)$ such that for every $n < \omega$, $S_{r,\alpha} \cap \delta_{i(n)} = S_{r_{i(n)},\alpha} \cap \delta_{i(n)}$, $\mathfrak{c}_{r,\alpha,\delta} = \mathfrak{c}_{r_{i(n)},\alpha,\delta}$ for every $\delta \in S_{r,\alpha} \cap \delta_{i(n)}$, $\delta \in S_{r,\alpha}$ and $\mathfrak{c}_{r,\alpha,\delta} = \bigoplus_{n \geq 1} \langle \mathfrak{d}_{i(n),\alpha,k} : k < n_\alpha \rangle$ where $\langle \mathfrak{d}_{i(n),\alpha,k} : k < n_\alpha \rangle$ is from clause (B)(ii) above. Put $\mathbb{Q}_{r,\alpha} = \mathbb{Q}_{\text{CR}_{r,\alpha}, \Sigma_{r,\alpha}} \upharpoonright \delta$.
- (c) $v_r = \bigcup_{n < \omega} v_{r_{i(n)}} \cup \{\alpha_\star\}$.
- (d) For $\alpha \in v_{r_{i(n)}}$, $p_{r,\alpha}^\star = p_{r_{i(n)},\alpha}^\star$ and $p_{r,\alpha_\star}^\star = \bigcup_{n < \omega} p_{i(n)} \upharpoonright Y_{i(n)}$. So $\text{dom}(p_{r,\alpha_\star}^\star)$ is an unbounded subset of $u_r \cap \alpha_\star$.
- (e) $\bar{A}_r = \bigcup_{n < \omega} \bar{A}_{r_{i(n)}} \cup \{(\delta, \{\gamma_{i(n)} : n < \omega\})\}$.

Next define p as follows.

- (i) $\text{dom}(p) = F \cup W \cup \bigcup_{n < \omega} Y_n$.
- (ii) If $\alpha \in F_0$, then $p(\alpha) = q_\alpha$ where q_α is from clause (B) above.
- (iii) If $\alpha \in F_1$, then $p(\alpha) = \{\mathfrak{c}_{r,\alpha,\delta}\}$.
- (iv) If $\alpha \in W$, then $p(\alpha) = p_{i(n)}(\alpha)$ which does not depend on $n < \omega$.
- (v) For every $n < \omega$, $p \upharpoonright Y_{i(n)} = p_{i(n)} \upharpoonright Y_i$.

It is clear that $(r_\star, p_\star) \leq_{\mathbb{R} \star \mathbb{P}_{\omega_2}} (r, p)$. By Lemma ??,

$$(r, p) \Vdash_{\mathbb{R} \star \mathbb{P}_{\omega_2}} \lim_n \frac{|\{k < n : (r_{i(k)}, p_{i(k)}) \in G_{\mathbb{R} \star \mathbb{P}_{\omega_2}}\}|}{n} = 1$$

Hence

$$(r, p) \Vdash_{\mathbb{R} \star \mathbb{P}_{\omega_2}} \lim_n \frac{|\{k < n : \gamma_{i(k)} \in \mathring{A}\}|}{n} = 1$$

Since $A_{r,\delta} = \{\gamma_{i(n)} : n < \omega\}$, the result follows. \square

6 On $\clubsuit^{\text{inf} \geq a}$

Definition 6.1. For $a \in (0, 1]$, the principle $\clubsuit^{\text{inf} > a-}$ says the following. There exists $\bar{A} = \langle A_\delta : \delta \in \bullet \text{Lim}(\omega_1) \rangle$ such that each $A_\delta = \{\alpha_{\delta,n} : n < \omega\}$ where $\alpha_{\delta,n}$'s are increasing cofinal in δ and for every $A \in [\omega_1]^{\aleph_1}$ and $b < a$, there exists some δ such that

$$\liminf_n \frac{|\{k < n : \alpha_{\delta,k} \in A\}|}{n} \geq b$$

Theorem 6.2. Let $0 < a \leq 1$ and suppose for every $b < a$, $\clubsuit^{\text{inf} \geq b}$ holds. Then $\clubsuit^{\text{inf} > a-}$ holds.

We need two lemmas.

Lemma 6.3. Suppose $\clubsuit_S^{\text{inf} \geq a}$ holds. Then there exists a partition $\langle S_i : i < \omega_1 \rangle$ of S into stationary sets such that for every $i < \omega_1$, $\clubsuit_{S_i}^{\text{inf} \geq a}$ holds.

Proof of Lemma ??: Fix a witness $\bar{A} = \langle A_\delta : \delta \in S \rangle$ for $\clubsuit_S^{\text{inf} \geq a}$ where each $A_\delta = \{\alpha_{\delta,n} : n < \omega\}$ and $\alpha_{\delta,n}$'s are increasing cofinal in δ . Note that if $a \in (0.5, 1]$, this is easy - Choose $\langle X_i : i < \omega_1 \rangle$ where X_i 's are pairwise disjoint unbounded subsets of ω_1 and let

$$S_i = \{\delta \in S : \liminf_n \frac{|\{k < n : \alpha_{\delta,k} \in X_i\}|}{n} \geq a\}$$

Since $a > 0.5$, S_i 's are pairwise disjoint and for every $Y \in [X_i]^{\aleph_1}$, there are stationary many $\delta \in S_i$ such that

$$\liminf_n \frac{|\{k < n : \alpha_{\delta,k} \in Y\}|}{n} \geq a$$

Fix $i < \omega_1$ and let $\{\alpha_\xi : \xi < \omega_1\}$ list X_i in increasing order. Choose a club $E \subseteq \omega_1$ such that for every $\delta \in E$, $\sup_{\xi < \delta} \alpha_\xi = \delta$. Define $\bar{C} = \langle C_\delta : \delta \in S_i \rangle$ as follows. If $\delta \in E \cap S_i$, put $C_\delta = \{\xi : \alpha_\xi \in A_\delta\}$, otherwise choose C_δ arbitrarily. It is clear that \bar{C} witnesses $\clubsuit_{S_i}^{\text{inf} \geq a}$.

In the general case, S_i 's may not be pairwise disjoint but for any $F \in [\omega_1]^K$, where $Ka > 1$, we have $\bigcap_{i \in F} S_i = \emptyset$. For $Y \subseteq \omega_1$, let $S(Y)$ be the set of $\delta \in S$ such that

$$\liminf_n \frac{|\{k < n : \alpha_{\delta,k} \in Y\}|}{n} \geq a$$

Claim 6.4. *There exists $\langle Y_i : i \in W \rangle$ such that $W \in [\omega_1]^{\aleph_1}$, each $Y_i \in [X_i]^{\aleph_1}$ and for every $i \in W$ and $Z \in [Y_i]^{\aleph_1}$, $S(Z) \setminus \bigcup_{j \in W \cap i} S(Y_j)$ is stationary.*

Proof of Claim ???: Let \mathcal{F} be the set of $\bar{Y} = \langle Y_i : i \in W \rangle$ where $W \in [\omega_1]^{\aleph_1}$ and each $Y_i \in [X_i]^{\aleph_1}$. For $\bar{Y} = \langle Y_i : i \in W \rangle \in \mathcal{F}$, let $n(\bar{Y})$ be the least n such that for every $F \in [W]^n$, $\bigcap_{i \in F} S(Y_i)$ is non-stationary - So $2 \leq n(\bar{Y}) \leq K$. Let $N = \min\{n(\bar{Y}) : \bar{Y} \in \mathcal{F}\}$ and fix $\bar{Y} = \langle Y_i : i \in W \rangle$ with $n(\bar{Y}) = N$. It suffices to show that for every $i_* \in W$, there exists $j \in W$ such that $j > i_*$ and for every $Z \in [Y_j]^{\aleph_1}$, $S(Z) \setminus \bigcup\{S(Y_i) : i \leq i_*, i \in W\}$ is stationary. Towards a contradiction, suppose this fails for some $i_* \in W$. Let $W' = W \setminus (i_* + 1)$. For each $j \in W'$, choose $Z_j \in [Y_j]^{\aleph_1}$ such that $S(Z_j) \setminus \bigcup\{S(Y_i) : i \leq i_*, i \in W\}$ is non-stationary. Let $\bar{Z} = \langle Z_j : j \in W' \rangle$. Then $n(\bar{Z}) \geq N$, so we can find $F \in [W']^{N-1}$ and such that $\bigcap_{j \in F} S(Z_j)$ is stationary. It follows that there exists $i \in W$ such that $i \leq i_*$ and $\bigcap_{j \in F} S(Z_j) \cap S(Y_i)$ is stationary. Hence $\bigcap_{j \in F \cup \{i_*\}} S(Y_j)$ is also stationary: Contradiction. \square

Let $\langle Y_i : i \in W \rangle$ be as in Claim ??. For $i \in W$, let $T_i = S(Y_i) \setminus \bigcup_{j \in W \cap i} S(Y_j)$. Then each T_i is stationary and for every $Z \in [Y_i]^{\aleph_1}$, there are stationary many $\delta \in T_i$ such that

$$\liminf_n \frac{|\{k < n : \alpha_{\delta,k} \in Z\}|}{n} \geq a$$

We can now proceed as before to get a $\clubsuit_{T_i}^{\text{inf} \geq a}$ witnessing sequence from $\langle A_\delta : \delta \in T_i \rangle$. This completes the proof of Lemma ??. \square

Lemma 6.5. *Suppose $\clubsuit_S^{\text{inf} \geq a}$ holds and $S = S_1 \cup S_2$. Then one of $\clubsuit_{S_1}^{\text{inf} \geq a}$, $\clubsuit_{S_2}^{\text{inf} \geq a}$ holds.*

Proof of Lemma ??: Fix a witness $\bar{A} = \langle A_\delta : \delta \in S \rangle$ for $\clubsuit_S^{\text{inf} \geq a}$ where each $A_\delta = \{\alpha_{\delta,n} : n < \omega\}$ and $\alpha_{\delta,n}$'s are increasing cofinal in δ . Suppose $\clubsuit_{S_1}^{\text{inf} \geq a}$ fails and choose $A \in [\omega_1]^{\aleph_1}$ such that for every $\delta \in S_1$

$$\liminf_n \frac{|\{k < n : \alpha_{\delta,k} \in A\}|}{n} < a$$

Since \bar{A} is $\clubsuit_S^{\text{inf} \geq a}$ witnessing sequence, it follows that for every $B \in [A]^{\aleph_1}$, there are stationary many $\delta \in S_2$ such that

$$\liminf_n \frac{|\{k < n : \alpha_{\delta,k} \in B\}|}{n} \geq a$$

Now we can construct a $\clubsuit_{S_2}^{\text{inf} \geq a}$ witnessing sequence as above. \square

Proof of Theorem ?? : Let $\langle a_n : n < \omega \rangle$ be an increasing sequence with $\lim_n a_n = a$. For each n , using Lemma ??, choose a sequence $\langle S_{n,i} : i < \omega_1 \rangle$ of pairwise disjoint stationary sets such that $\clubsuit_{S_{n,i}}^{\text{inf} \geq a_n}$ holds. For $m < n < \omega$, define $W_{m,n} = \{i < \omega_1 : \clubsuit_{S_{m,i}}^{\text{inf} \geq a_n} \text{ holds}\}$.

First suppose that for some $m < \omega$, there are infinitely many $n > m$ such that $W_{m,n}$ is infinite. Let $\langle n(k) : k < \omega \rangle$ list such n 's in increasing order. Inductively choose $i(k) \in W_{m,n(k)}$ such that $i(k)$'s are pairwise distinct and $\clubsuit_{S_{m,i(k)}}^{\text{inf} \geq a_{n(k)}}$ holds. Since $\langle S_{m,i(k)} : k < \omega \rangle$ consists of pairwise disjoint sets, the result follows.

So we can assume that there is no such m . Inductively choose a strictly increasing sequence $\langle m(k) : k < \omega \rangle$ such that for every $n \geq m(k+1)$, $W_{m(k),n}$ is finite. Let $W = \bigcup \{W_{m(j),m(k)} : j < k < \omega\}$ and choose $i > \sup(W)$. Put $T_k = S_{m(k),i} \setminus \bigcup_{l < k} S_{m(l),i}$ and $T'_k = S_{m(k),i} \setminus T_k$. Then T_k 's are pairwise disjoint, $S_{m(k),i} = T_k \cup T'_k$ and by our choice of i , $\clubsuit_{T'_k}^{\text{inf} \geq a_{m(k)}}$ does not hold. Hence, by Lemma ??, $\clubsuit_{T_k}^{\text{inf} \geq a_{m(k)}}$ must hold and we are done. \square

Proof of Theorem ??(2): Fix $0 < a < 1$. We indicate the essential changes in the proof of Theorem ??(1) to get a model of $\clubsuit^{\text{inf} \geq a} \wedge (\forall b \in (a, 1]) \neg \clubsuit^{\text{inf} \geq b}$. Define a modified countable join as follows. In Definition ??, replace Clause (3)(ii)(b) by (b_\star) and Clause (4)(ii)(b) by $(b_{\star\star})$ below.

- (b_\star) $|\{i \in [2, j_1) : (\exists k \in [m, n))(\mathfrak{d}'_{i,k} \neq \mathfrak{d}_{i,k})\}| \leq j_1(1 - a)$ for every $2 < j_1 \leq j$.
- $(b_{\star\star})$ $|\{i \in [N, j_1) : (\exists k \in [m, n))(\mathfrak{d}'_{i,k} \neq \mathfrak{d}_{i,k})\}| \leq (j_1 - N)(1 - a)$ for every $N < j_1 \leq j$.

Note that this gives rise to a transitive Σ'_p there. Lemma ?? gets modified to the following.

Lemma 6.6. *Let (CR'_p, Σ'_p) be as in Definition ?? with (b_\star) in place of Clause (3)(ii)(b) and $(b_{\star\star})$ in place of Clause (4)(ii)(b). Let (CR, Σ) be an \aleph_1 -CP such that $CR'_p = \{\mathfrak{c} \in CR : \text{dom}(\mathfrak{c}) \subseteq \delta\}$ and $\Sigma'_p = \Sigma \upharpoonright CR'_p$. Let $\mathbb{Q} = \mathbb{Q}_{CR, \Sigma}$, $p = \{\mathfrak{c}_1^* = \bigoplus_{i \geq 1} \bar{\mathfrak{d}}_i\}$ and $p_i = \{\mathfrak{d}_{i,k} : k < n\}$. Then*

$$p \Vdash_{\mathbb{Q}} \liminf_j \frac{|\{i < j : p_i \in G_{\mathbb{Q}}\}|}{j} \geq a$$

Next, Lemma ?? gets replaced by the following.

Lemma 6.7. *For every $b \in (a, 1]$, $V^{\mathbb{P}_{\omega_2}} \models \neg \clubsuit^{\text{inf} \geq b}$.*

Proof of Lemma ?? : Fix $b' \in (a, 1]$. Towards a contradiction, suppose $p_0 \in \mathbb{P}_{\omega_2}$, $\langle \dot{A}_\delta = \{\dot{\alpha}_{\delta,n} : n < \omega\} : \delta \in \bullet \text{Lim}(\omega_1) \rangle \in V^{\mathbb{P}_{\omega_2}}$ are such that $p_0 \Vdash$ “ $(\forall \delta \in \bullet \text{Lim}(\omega_1))(\{\dot{\alpha}_{\delta,n} : n < \omega\}$ is increasing cofinal in δ) and $\langle \dot{A}_\delta : \delta < \omega_1 \rangle$ is a $\clubsuit^{\text{inf} \geq b'}$ witnessing sequence”. Since \mathbb{P}_{ω_2} satisfies ccc, we can find $\gamma < \omega_2$ such that $p_0 \in \mathbb{P}_\gamma$ and each $\dot{\alpha}_{\delta,n}$ is a \mathbb{P}_γ -name. Fix $b \in (a, b')$.

Let $\dot{X} = \{\alpha < \omega_1 : \dot{f}_{\mathbb{Q}_\gamma} = 1\}$. Then $\dot{X} \in V^{\mathbb{P}_{\gamma+1}}$ and $V^{\mathbb{P}_{\gamma+1}} \models \dot{X} \in [\omega_1]^{\aleph_1}$. So there exist $p_1 \in \mathbb{P}_\gamma$, $q \in \mathbb{Q}_\gamma$, $\delta \in \bullet \text{Lim}(\omega_1)$ and $n_0 < \omega$ such that $p_1 \geq p_0$ and $(p_1, q) \Vdash_{\mathbb{P}_{\gamma+1}} (\forall j \geq$

$n_0)(|\{i < j : \dot{\alpha}_{\delta,i} \in \dot{X}\}| \geq jb)$. We must have that $\text{dom}(q) \cap \delta$ is unbounded in δ otherwise we can easily extend (p_1, q) to get a contradiction. By possibly extending q , by Definition ??(D)(ii), we can assume that $q = \{\mathbf{c}_k : k < K_\star\}$ where $\sup(\text{dom}(\mathbf{c}_k)) < \inf(\text{dom}(\mathbf{c}_{k+1}))$ for every $k < K_\star - 1$ and for some $K < K_\star$, $\text{dom}(\mathbf{c}_K)$ is an unbounded subset of δ . Let S_γ and $\langle \mathbf{c}_{\gamma,\delta} : \delta \in S_\gamma \rangle$ witness that $(\text{CR}_\gamma, \Sigma_\gamma)$ is a thin \aleph_1 -CP. By Claim ??, we can further assume that $\mathbf{c}_K = \mathbf{c}'_{K'}$ for some $\langle \mathbf{c}'_n : n \leq K' \rangle \in \Sigma(\mathbf{c}_{\gamma,\delta})$.

Let $m < n < \omega$ and $\bar{\mathbf{d}}_i = \langle \mathbf{d}_{i,k} : k < n \rangle$ for $i \geq 1$ be as in Definition ?? and $\mathbf{c}_{\gamma,\delta} = \bigoplus_{i \geq 1} \bar{\mathbf{d}}_i$. Then as $\langle \mathbf{c}'_n : n \leq K' \rangle \in \Sigma(\mathbf{c}_{\gamma,\delta})$, we can find $N \geq 1$ a power of 2 such that $\mathbf{c}_K = \mathbf{c}'_{K'} = \mathbf{c}^*_N$ in the notation of Definition ??.

Choose $p_2 \in \mathbb{P}_\gamma$, $p_2 \geq p_1$, $n_\star > n_0$ a power of 2 and $\alpha_{n_\star} > \min(\text{dom}(\mathbf{c}^*_N))$ such that $p_2 \Vdash_{\mathbb{P}_\gamma} \dot{\alpha}_{\delta,n_\star} = \alpha_{n_\star}$. Put $c = (a + b)/2$. Let $n_{\star\star} > n_\star$ be a power of 2 such that $n_\star/n_{\star\star} < (b - c)/(1 - c)$. Choose $p_3 \geq p_2$ and $\langle \alpha_n : n \in [n_\star, n_{\star\star}] \rangle$ such that for every $n \in [n_\star, n_{\star\star}]$, $p_3 \Vdash_{\mathbb{P}_\gamma} \dot{\alpha}_{\delta,n} = \alpha_n$. Let $F = \{\alpha_n \notin \text{dom}(q) : n \in [n_\star, n_{\star\star}]\}$. Let $q' = q \cup \bigcup_{\alpha \in F} \{\mathfrak{d}_\alpha\}$ where $\text{dom}(\mathfrak{d}_\alpha) = \{\alpha\}$ and $f_{\mathfrak{d}_\alpha}(\alpha) = 0$. If F is empty, put $q' = q$.

Now it is possible to choose $\bar{\mathbf{g}} \in \Sigma(\mathbf{c}^*_N)$ such that letting $q'' = (q' \setminus \{\mathbf{c}^*_N\}) \cup \bar{\mathbf{g}}$ forces $\{n \in [n_\star, n_{\star\star}] : \alpha_n \notin \dot{X}\} \geq (1 - c)(n_{\star\star} - n_\star)$ - We leave the details of this to the reader. This means that (p_3, q'') forces that $|\{i < n_{\star\star} : \dot{\alpha}_{\delta,i} \in \dot{X}\}| \leq n_\star + c(n_{\star\star} - n_\star) < bn_{\star\star}$ which is a contradiction. \square

Now the remainder of the proof is exactly the same except for the fact that at the end of the proof of $\clubsuit^{\text{inf} \geq a}$, we use Lemma ?? in place of Lemma ??. \square

Proof of Theorem ??(3): Let $\langle a_k : k \geq 1 \rangle$ be an increasing sequence with limit a . Proceed as in the proof of Theorem ??(2) with the following modification for countable joins. In Definition ??, replace Clause (3)(ii)(b) by (b \star) and Clause (4)(ii)(b) by (b $\star\star$) below.

- (b \star) $|\{i \in [2, j_1] : (\exists k \in [m, n])(\mathfrak{d}'_{i,k} \neq \mathfrak{d}_{i,k})\}| \leq j_1(1 - a_{k_\star})$ for every $2 < j_1 \leq j$.
 (b $\star\star$) $|\{i \in [N, j_1] : (\exists k \in [m, n])(\mathfrak{d}'_{i,k} \neq \mathfrak{d}_{i,k})\}| \leq (j_1 - N)(1 - a_{k_\star})$ for every $N < j_1 \leq j$.

The rest of the proof is similar to that of Theorem ??(2). We leave the details to the reader. \square

7 On \clubsuit^{sup}

Definition 7.1. For $a \in (0, 1]$ and $S \subseteq \bullet\text{Lim}(\omega_1)$ stationary, the principle $\clubsuit^{\text{sup} \geq a}_S$ says the following: There exists $\bar{A} = \langle A_\delta : \delta \in S \rangle$ such that

- (a) each $A_\delta = \{\alpha_{\delta,n} : n < \omega\}$ and $\alpha_{\delta,n}$'s are increasing cofinal in δ and
 (b) for every $A \in [\omega_1]^{\aleph_1}$, there exists $\delta \in S$ such that

$$\limsup_n \frac{|\{k < n : \alpha_{\delta,k} \in A\}|}{n} \geq a$$

As usual, if $S = \bullet \text{Lim}(\omega_1)$, we just write $\clubsuit^{\text{sup} \geq a}$.

The following remark describes the situation in the Cohen and the random reals models.

Remark 7.2. (1) Suppose $V \models \clubsuit$ and let \mathbb{P} be the forcing for adding \aleph_2 Cohen reals. Then $V^{\mathbb{P}} \models \clubsuit^{\text{sup} \geq 1} \wedge (\forall a > 0) \neg \clubsuit^{\text{inf} \geq a}$. Moreover, the following fails in $V^{\mathbb{P}}$: There exists $\bar{A} = \langle A_\delta : \delta \in \bullet \text{Lim}(\omega_1) \rangle$ where each $A_\delta = \{\alpha_{\delta,n} : n < \omega\}$ and $\alpha_{\delta,n}$'s are increasing cofinal in δ such that for every $A \in [\omega_1]^{\aleph_1}$ and $\varepsilon > 0$, there exists some δ such that

$$\liminf_n \frac{|\{k < n : \alpha_{\delta,k} \in A\}|}{n} \geq \varepsilon$$

(2) Suppose $V \models \clubsuit$ and let \mathbb{P} be the forcing for adding \aleph_2 random reals. Then $V^{\mathbb{P}} \models (\forall a > 0) \neg \clubsuit^{\text{sup} \geq a}$. Moreover, the following holds in $V^{\mathbb{P}}$: There exists $\bar{A} = \langle A_\delta : \delta \in \bullet \text{Lim}(\omega_1) \rangle$ where each $A_\delta = \{\alpha_{\delta,n} : n < \omega\}$ and $\alpha_{\delta,n}$'s are increasing cofinal in δ such that for every $A \in [\omega_1]^{\aleph_1}$, there exists $\varepsilon > 0$ and δ such that for every sufficiently large n ,

$$\frac{|\{k < n : \alpha_{\delta,k} \in A\}|}{n} \geq \varepsilon$$

We now prove Theorem ?? - For all $a, b \in (0, 1)$, $\clubsuit_S^{\text{sup} \geq a}$ is equivalent to $\clubsuit_S^{\text{sup} \geq b}$. For this, it is clearly enough to show the following.

Lemma 7.3. Let $a \in (0, 1)$ and $a \leq b < \sqrt{a}$. Then $\clubsuit_S^{\text{sup} \geq a}$ implies $\clubsuit_S^{\text{sup} \geq b}$.

Proof of Lemma ??: Let $\bar{A} = \langle A_\delta : \delta \in S \rangle$ witness $\clubsuit_S^{\text{sup} \geq a}$. We can assume that \bar{A} is not a $\clubsuit_S^{\text{sup} \geq b}$ witnessing sequence. Choose $A \in [\omega_1]^{\aleph_1}$ such that for every $\delta \in S$, for every large enough $\alpha < \delta$

$$\frac{|A \cap A_\delta \cap \alpha|}{|A_\delta \cap \alpha|} < b$$

Let S' be the set of $\delta \in S$ such that

$$\limsup_{\alpha \rightarrow \delta} \frac{|A \cap A_\delta \cap \alpha|}{|A_\delta \cap \alpha|} \geq a$$

Then S' is stationary. For $\delta \in S'$, define $B_\delta = A_\delta \cap A$.

Claim 7.4. For every $B \in [A]^{\aleph_1}$ there are stationary many $\delta \in S'$ such that

$$\limsup_{\alpha \rightarrow \delta} \frac{|B \cap B_\delta \cap \alpha|}{|B_\delta \cap \alpha|} \geq b$$

Proof of Claim ??: Suppose not. Choose $B \in [A]^{\aleph_1}$ and $W \subseteq S'$ non stationary such that for every $\delta \in S' \setminus W$, for every large enough $\alpha < \delta$, we have

$$\frac{|B \cap B_\delta \cap \alpha|}{|B_\delta \cap \alpha|} < b$$

Since $B \subseteq A$, we can choose $\delta \in S' \setminus W$ such that

$$\limsup_{\alpha \rightarrow \delta} \frac{|B \cap A_\delta \cap \alpha|}{|A_\delta \cap \alpha|} \geq a$$

Now for every large enough $\alpha < \delta$, we have

$$\left(\frac{|B \cap B_\delta \cap \alpha|}{|B_\delta \cap \alpha|} \right) \left(\frac{|A \cap A_\delta \cap \alpha|}{|A_\delta \cap \alpha|} \right) < b^2$$

Since $B \cap B_\delta = B \cap A_\delta$ and $B_\delta \cap \alpha = A \cap A_\delta \cap \alpha$, we get

$$\frac{B \cap A_\delta \cap \alpha}{A_\delta \cap \alpha} < b^2 < a$$

which is impossible.

Let $\{\alpha_i : i < \omega_1\}$ list A in increasing order. Let $E \subseteq \omega_1$ be a club such that for every $i \in E$, $\sup_{j < i} \alpha_j = i$. Define $\bar{C} = \langle C_\delta : \delta \in S \rangle$ as follows. If $\delta \in E \cap S'$, then $C_\delta = \{j < \delta : \alpha_j \in B_\delta\}$. Otherwise, choose C_δ to be an arbitrary unbounded subset of δ of order type ω . It is easy to check that \bar{C} witnesses $\clubsuit_S^{\sup \geq b}$. \square

8 $\neg \clubsuit^{\sup \geq 1}$ and $\clubsuit^{\sup > 1-}$

Definition 8.1. The principle $\clubsuit^{\sup > 1-}$ says the following: There exists $\bar{A} = \langle A_\delta : \delta \in \bullet \text{Lim}(\omega_1) \rangle$ such that

- (a) each $A_\delta = \{\alpha_{\delta,n} : n < \omega\}$ and $\alpha_{\delta,n}$'s are increasing cofinal in δ and
- (b) for every $A \in [\omega_1]^{\aleph_1}$ and $\varepsilon > 0$, there exists some δ such that

$$\limsup_n \frac{|\{k < n : \alpha_{\delta,k} \in A\}|}{n} \geq 1 - \varepsilon$$

To prove Theorem ??, it is enough to show that

Theorem 8.2. $\neg \clubsuit^{\sup \geq 1} \wedge \clubsuit^{\sup > 1-}$ is consistent.

Definition 8.3. Suppose $\bar{A} = \langle A_\delta : \delta \in \bullet \text{Lim}(\omega_1) \rangle$ satisfies: For every δ , $A_\delta = \{\alpha_{\delta,n} : n < \omega\}$ where $\alpha_{\delta,n}$'s are increasing and cofinal in δ . Define $\mathbb{Q} = \mathbb{Q}_{\bar{A}}$ as follows: $p \in \mathbb{Q}$ iff $p = (f_p, u_p, \bar{\varepsilon}_p)$ where

- (i) f_p is a finite partial function from ω_1 to $\{0, 1\}$,
- (ii) u_p is a finite subset of $\bullet \text{Lim}(\omega_1)$ and
- (iii) $\bar{\varepsilon}_p = \langle \varepsilon_{p,\delta} : \delta \in u_p \rangle$ where each $\varepsilon_{p,\delta}$ is a positive rational < 1 .

For $p, q \in \mathbb{Q}$ define $p \leq q$ iff

- (a) $f_p \subseteq f_q$,

- (b) $u_p \subseteq u_q$,
- (c) $\bar{\varepsilon}_p = \bar{\varepsilon}_q \upharpoonright u_p$ and
- (d) for every $\delta \in u_p$, letting $W = \{n < \omega : \alpha_{\delta,n} \in \text{dom}(f_q) \setminus \text{dom}(f_p)\}$, for every $N < \omega$ either $W \cap [0, N) = \emptyset$ or

$$\frac{|\{n \in W \cap [0, N) : f_q(\alpha_{\delta,n}) = 1\}|}{|W \cap [0, N)|} \leq 1 - \varepsilon_{p,\delta}$$

Claim 8.4. Let \bar{A} and $\mathbb{Q} = \mathbb{Q}_{\bar{A},a}$ be as in Definition ???. Then \mathbb{Q} has \aleph_1 as a precaliber.

Proof of Claim ???: Suppose $\{p_i = (f_i, u_i, \bar{\varepsilon}_i) : i < \omega_1\} \subseteq \mathbb{Q}$. By thinning down we can assume the following.

- (a) $\langle \text{dom}(f_i) : i < \omega_1 \rangle$ is a Δ -system with root R and $f_i \upharpoonright R$ does not depend on i .
- (b) $\langle u_i : i < \omega_1 \rangle$ is a Δ -system with root u_* and $\bar{\varepsilon}_i \upharpoonright u_*$ does not depend on i .
- (c) For every $i < j < \omega_1$ and $\delta \in u_i$, $\text{dom}(f_j) \cap A_\delta \subseteq R$.

Let $E \subseteq \omega_1$ be a club such that for every $i \in E$, for every $j < i$, $\text{dom}(f_j) \cup u_j \subseteq i$. Choose $S \subseteq E$ stationary such that for every $i \in S$, $\text{dom}(f_i) \cap i = R$, $u_i \cap i = u_*$ and $\bigcup \{A_\delta \cap i : \delta \in u_i, \delta > i\} = F$ where F does not depend on $i \in S$. Note that for every infinite $X \subseteq S$ and $i \in S$, if $i > \sup(X)$, then for all but finitely many $j \in X$, $\text{dom}(f_j) \cap A_i \subseteq R$. Let $X \in [S]^{\aleph_1}$ be such that for every increasing sequence $\langle \alpha_n : n < \omega \rangle$ in X , $\sup_n \alpha_n \notin X$. Define $c : [X]^2 \rightarrow \{0, 1\}$ by $c(\{i, j\}) = 1$ iff $i < j$ and $A_j \cap \text{dom}(f_i) \subseteq R$. By Erdos-Dushnik-Miller, either there exists $Y \in [X]^{\aleph_1}$ such that $c[[Y]^2] = \{1\}$ or there exists $Y' \subseteq X$ such that $\text{otp}(Y') = \omega + 1$ and $c[[Y']^2] = \{0\}$. Since the latter is impossible, we can find $Y \in [X]^{\aleph_1}$ such that $c[[Y]^2] = \{1\}$. Hence

- (d) For every $i \neq j$ in Y and $\delta \in u_j$, $\text{dom}(f_i) \cap A_\delta \subseteq R$.

It follows that $\{p_i : i \in Y\}$ is centered. □

Let $\mathring{f}_\mathbb{Q} = \bigcup \{f_p : p \in G_\mathbb{Q}\}$. Then $\Vdash_\mathbb{Q} \mathring{f}_\mathbb{Q} : \omega_1 \rightarrow \{0, 1\}$. Let $\mathring{X}_\mathbb{Q} = \{\alpha < \omega_1 : \mathring{f}_\mathbb{Q}(\alpha) = 1\}$. Then $\Vdash_\mathbb{Q} \mathring{X}_\mathbb{Q} \in [\omega_1]^{\aleph_1}$.

Claim 8.5. $\mathring{X}_\mathbb{Q}$ witnesses that \bar{A} is not a $\clubsuit^{\text{sup} \geq 1}$ witnessing sequence in $V^\mathbb{Q}$.

Proof: Easy. □

Claim 8.6. Suppose $V \models \clubsuit^{\text{sup} > 1-}$ holds and let $\bar{C} = \langle C_\delta : \delta \in \bullet \text{Lim}(\omega_1) \rangle$ be a witness where $C_\delta = \{\beta_{\delta,n} : n < \omega\}$ and $\beta_{\delta,n}$'s are increasing cofinal in δ . Then $V^\mathbb{Q} \models \clubsuit^{\text{sup} > 1-}$ holds with \bar{C} as witness.

Proof of Claim ???: Suppose $p \Vdash_\mathbb{Q} \mathring{A} \in [\omega_1]^{\aleph_1}$ and $\varepsilon > 0$. Choose $\langle (p_i, \gamma_i) : i < \omega_1 \rangle$ such that γ_i 's are increasing and for each $i < \omega_1$, $p \leq p_i \Vdash_\mathbb{Q} \gamma_i \in \mathring{A}$. Arguing as in the proof of Claim ??, we can assume the following.

- (a) $\langle \text{dom}(f_i) : i < \omega_1 \rangle$ is a Δ -system with root R , $f_i \upharpoonright R = f_\star$ and $|\text{dom}(f_i) \setminus R| = n_\star$ do not depend on i .
- (b) If $i < j$, then $R < \text{dom}(f_i) \setminus R < \text{dom}(f_j) \setminus R$.
- (c) $\langle u_i : i < \omega_1 \rangle$ is a Δ -system with root u_\star , $\bar{\varepsilon}_i \upharpoonright u_\star = \bar{e}_\star$ does not depend on i and $i < j$ implies $u_i \setminus u_\star < u_j \setminus u_\star$.
- (c) For every $i \neq j$ and $\delta \in u_i$, $\text{dom}(f_j) \cap A_\delta \subseteq R$.

Put $X = \{\gamma_i : i < \omega_1\}$. Let $E \subseteq \omega_1$ be a club such that for every $i \in E$ and $j < i$, $\gamma_j < i$ and $u_\star \cup \text{dom}(f_j) \subseteq i$. Choose $\delta \in E$ such that

$$\limsup_n \frac{|\{k < n : \beta_{\delta,k} \in X\}|}{n} \geq 1 - \varepsilon/10$$

Let $q = (f_\star, u_\star \cup \{\delta\}, \bar{\varepsilon}_\star \cup \{(\delta, \varepsilon/5)\})$. It suffices to show that for any $q_1 \geq q$ and $N_0 < \omega$, there exist $r \geq q_1$ and $N_2 > N_0$ such that

$$r \Vdash_{\mathbb{Q}} \frac{|\{n < N_2 : \beta_{\delta,n} \in \dot{A}\}|}{N_2} \geq 1 - \varepsilon$$

So fix $q_1 \geq q$ and $N_0 < \omega$. For each $n < \omega$, define

$$r_n = \begin{cases} p_i & \text{if } \beta_{\delta,n} = \gamma_i \\ q & \text{if } \beta_{\delta,n} \notin X \end{cases}$$

Let $W'_n = \text{dom}(f_{r_n}) \setminus R$ and $W_n = W'_n \cap A_\delta$. Choose $N_1 > N_0$ such that for every $n \geq N_1$, if $\delta' \in u_{q_1} \setminus \{\delta\}$, then $W'_n \cap A_{\delta'} = \emptyset$. We need a lemma.

Lemma 8.7. *Suppose $0 < a_1 < a_2 < 1$ and $1 \leq K < \omega$. Then for all sufficiently large $N < \omega$, the following holds. For every $\langle W_k : k < N \rangle$ where each W_k is an interval in ω such that $|W_k| \leq K$, $W_k < W_{k+1}$ and $\bigcup_{k < n} W_k = [0, M)$, there exists $F \subseteq N$ such that*

(i) $|F| \geq Na_1$ and

(ii) For every $m \leq M$, $|[0, m) \cap \bigcup_{k \in F} W_k| \leq ma_2$

Proof of Lemma ???: First assume that $|W_k| = K$ for every $k < N$ - So $M = NK$. Let $m_1 < N$ be least such that $Km_1 \geq M(1 - a_2)$. Then $F = [m_1, N)$ is as required. For the general case, for each $K' \leq K$, put $S_{K'} = \{k < N : |W_k| = K'\}$ and find a suitable $F_{K'} \subseteq S_{K'}$ for $\langle W_k : k \in S_{K'} \rangle$. Then $F = \bigcup \{F_{K'} : 1 \leq K' \leq K\}$ is as required. \square

Choose $N_2 > N_1$ such that $(1 - N_1/N_2)(1 - \varepsilon/2) \geq 1 - \varepsilon$ and $|\{k \in [N_1, N_2) : \beta_{\delta,k} \in X\}| \geq (1 - \varepsilon/4)(N_2 - N_1)$. Using Lemma ??, choose $F \subseteq [N_1, N_2)$ such that the following hold.

(a) $|F| \geq (N_2 - N_1)(1 - \varepsilon/4)$.

- (b) $r = (f_r, u_r, \bar{\varepsilon}_r)$ extends each condition in $\{q_1, r_n : n \in F\}$ where
- (i) $u_r = u_{q_1} \cup \bigcup_{n \in F} u_{r_n}$,
 - (ii) $\text{dom}(f_r) = \text{dom}(f_{q_1}) \cup \bigcup_{n \in F} W'_n \cup \bigcup \{W_n : n \in [N_1, N_2) \setminus F\}$,
 - (iii) $f_{q_1} \subseteq f_r$,
 - (iv) $f_r \upharpoonright \bigcup \{W_n : n \in [N_1, N_2) \setminus F\} \equiv 0$,
 - (v) for every $n \in F$, $f_r \upharpoonright W'_n = f_{r_n}$ and
 - (vi) $\bar{\varepsilon}_r = \bar{\varepsilon}_{q_1} \cup \bigcup_{n \in F} \bar{\varepsilon}_{r_n}$.

Note that $r \Vdash |\{k < N_2 : \beta_{\delta,k} \in \mathring{A}\}| \geq (N_2 - N_1)(1 - \varepsilon/2)$. By our choice of N_2 , it follows that

$$r \Vdash_{\mathbb{Q}} \frac{|\{n < N_2 : \beta_{\delta,n} \in \mathring{A}\}|}{N_2} \geq 1 - \varepsilon$$

□

Let $\eta \geq 1$ and suppose $\langle (\mathbb{P}_\xi, \mathbb{Q}_\xi, \bar{A}_\xi) : \xi < \eta \rangle$ satisfies the following.

- (1) $\langle (\mathbb{P}_\xi, \mathbb{Q}_\xi) : \xi < \eta \rangle$ is a finite support iteration with limit \mathbb{P}_η .
- (2) $\bar{A}_\xi \in V^{\mathbb{P}_\xi}$ and $\Vdash_{\mathbb{P}_\xi} \text{“}\bar{A}_\xi = \langle A_{\xi,\delta} : \delta \in \bullet \text{Lim}(\omega_1) \rangle$, $A_{\xi,\delta} = \{\alpha_{\xi,\delta,n} : n < \omega\}$ where $\alpha_{\xi,\delta,n}$'s are increasing cofinal in δ ”.
- (3) $V^{\mathbb{P}_\xi} \models \mathbb{Q}_\xi = \mathbb{Q}_{\bar{A}_\xi}$.

Note that \mathbb{P}_η is ccc.

Claim 8.8. *Suppose $V \models \clubsuit^{\text{sup} > 1-}$ holds and let $\bar{C} = \langle C_\delta : \delta \in \bullet \text{Lim}(\omega_1) \rangle$ be a witness where $C_\delta = \{\beta_{\delta,n} : n < \omega\}$ and $\beta_{\delta,n}$'s are increasing cofinal in δ . Then $V^{\mathbb{P}_\eta} \models \clubsuit^{\text{sup} \geq 1-}$ via the same witness.*

Proof of Claim ??: By induction on η . If η is a successor or $\text{cf}(\eta) > \aleph_1$, this follows from Claim ??.

Suppose $\text{cf}(\eta) = \aleph_0$. Let $\langle \eta(n) : n < \omega \rangle$ be increasing cofinal in η . Suppose $p \Vdash_{\mathbb{P}_\eta} \mathring{X} \in [\omega_1]^{\aleph_1}$. Choose $n_\star < \omega$ such that $p \in \mathbb{P}_{\eta(n_\star)}$. For each $n < \omega$, let $\mathring{X}_n = \{\alpha < \omega_1 : (\exists p \in G_{\mathbb{P}_{\eta(n)}})(p \Vdash_{\mathbb{P}_\eta} \alpha \in \mathring{X})\}$ - So $\mathring{X}_n \in V^{\mathbb{P}_{\eta(n)}}$ and $\Vdash_{\mathbb{P}_\eta} \mathring{X}_n \subseteq \mathring{X}$. Then for some $n \in [n_\star, \omega)$, $p \Vdash_{\mathbb{P}_{\eta(n)}} \mathring{X}_n \in [\omega_1]^{\aleph_1}$. Now apply the inductive hypothesis.

Next suppose $\text{cf}(\eta) = \aleph_1$, $\varepsilon > 0$ and $p \Vdash_{\mathbb{P}_\eta} \mathring{X} \in [\omega_1]^{\aleph_1}$. Choose $\langle (p_i, \gamma_i) : i < \omega_1 \rangle$ such that the following hold.

- (a) γ_i 's are increasing.
- (b) $p_i \in \mathbb{P}_\eta$, $p_i \geq p$ and $p_i \Vdash_{\mathbb{P}_\eta} \gamma_i \in \mathring{X}$.
- (c) $\langle \text{dom}(p_i) : i < \omega_1 \rangle$ is a Δ -system with root W .

Choose $\theta < \eta$ such that $W \subseteq \theta$. Since \mathbb{P}_θ is ccc, we can find $q \in \mathbb{P}_\theta$ such that $q \geq p$ and $q \Vdash_{\mathbb{P}_\theta}$ “ $\{i < \omega_1 : p_i \upharpoonright \theta \in G_{\mathbb{P}_\theta}\}$ is uncountable”. Let $\dot{Y} = \{\gamma_i : i < \omega_1 \wedge p_i \upharpoonright \theta \in G_{\mathbb{P}_\theta}\}$. Then $\dot{Y} \in V^{\mathbb{P}_\theta}$ and $q \Vdash_{\mathbb{P}_\theta} \dot{Y} \in [\omega_1]^{\aleph_1}$. By the inductive hypothesis, we can find $r \in \mathbb{P}_\theta$ and $\delta \in \bullet\text{Lim}(\omega_1)$ such that $r \geq q$ and

$$r \Vdash_{\mathbb{P}_\theta} \limsup_n \frac{|\{k < n : \beta_{\delta,k} \in \dot{Y}\}|}{n} \geq 1 - \varepsilon/2$$

Since $\langle \text{dom}(p_i) \setminus \theta : i < \omega_1 \rangle$ is a sequence of pairwise disjoint sets, it also follows that

$$r \Vdash_{\mathbb{P}_\eta} \limsup_n \frac{|\{k < n : \beta_{\delta,k} \in \dot{X}\}|}{n} \geq 1 - \varepsilon$$

□

Proof of Theorem ???: Starting with a model of $2^{\aleph_1} = \aleph_2$ and $\clubsuit^{\text{sup}>1-}$ construct $\langle (\mathbb{P}_\xi, \mathbb{Q}_\xi, \bar{A}_\xi) : \xi < \omega_2 \rangle$ such that the following hold.

- (1) $\langle (\mathbb{P}_\xi, \mathbb{Q}_\xi) : \xi < \omega_2 \rangle$ is a finite support iteration with limit \mathbb{P}_{ω_2} .
- (2) $\bar{A}_\xi \in V^{\mathbb{P}_\xi}$ and $\Vdash_{\mathbb{P}_\xi}$ “ $\bar{A}_\xi = \langle A_{\xi,\delta} : \delta \in \bullet\text{Lim}(\omega_1) \rangle$, $A_{\xi,\delta} = \{\alpha_{\xi,\delta,n} : n < \omega\}$ where $\alpha_{\xi,\delta,n}$ ’s are increasing cofinal in δ ”.
- (3) $V^{\mathbb{P}_\xi} \models \mathbb{Q}_\xi = \mathbb{Q}_{\bar{A}_\xi}$.
- (4) For every $\eta < \omega_2$ and $\bar{A} \in V^{\mathbb{P}_\eta}$ satisfying $\Vdash_{\mathbb{P}_\eta}$ “ $\bar{A} = \langle A_\delta : \delta \in \bullet\text{Lim}(\omega_1) \rangle$ where each A_δ is an unbounded subset of δ of order type ω ”, there exists $\xi \in [\eta, \omega_2)$ such that $\Vdash_{\mathbb{P}_\xi} \bar{A} = \bar{A}_\xi$.

To see why clause (4) can be satisfied, use $2^{\aleph_1} = \aleph_2$ and the fact that for each $\eta < \omega_2$, \mathbb{P}_η is a ccc forcing with a dense subset of size \aleph_1 . □

We conclude with some questions.

Question 8.9. (1) Is $\clubsuit^{\text{sup} \geq 0.5} \wedge \neg \clubsuit^{\text{sup} > 1-}$ consistent? What if CH holds?

(2) Assume CH. Does $\clubsuit^{\text{sup} \geq 0.5}$ imply $\clubsuit^{\text{sup} \geq 1}$? Does $\clubsuit^{\text{sup} > 1-}$ imply $\clubsuit^{\text{sup} \geq 1}$?

(3) For $a \in (0, 1)$, is $\clubsuit^{\text{inf} \geq a} \wedge \neg \clubsuit^{\text{sup} \geq 1}$ consistent?

References

- [1] M. Dzamonja, S. Shelah, Similar but not the same: various versions of \clubsuit do not coincide, *Journal of Symbolic Logic* 64 (1999), 180-198
- [2] A. Kumar, S. Shelah, RVM, RVC revisited: Clubs and Lusin sets, Submitted
- [3] A. Kumar, S. Shelah, Clubs on quasi measurable cardinals, *Mathematical Logic Quarterly*, To appear
- [4] K. Kunen, *Set theory: An introduction to independence proofs*, Studies in Logic and the Foundations of Mathematics, Vol. 102, North-Holland Publishing Company, 1980