

On Hanf numbers of the infinitary order property*

Rami Grossberg
Department of Mathematics
Carnegie Mellon University
Pittsburgh, PA 15213

Saharon Shelah[†]
Institute of Mathematics
The Hebrew University of Jerusalem
Jerusalem, 91094 ISRAEL

&
Department of Mathematics
Rutgers University
New Brunswick, NJ 08902

October 5, 2020

*The authors thank the United States - Israel Binational Science foundation for supporting this research as well as the Mathematics department of Rutgers University where part of this research was carried out.

[†]This paper replaces item # 259 from Shelah's list of publications.

Abstract

We study several cardinal, and ordinal-valued functions that are relatives of Hanf numbers. Let κ be an infinite cardinal, and let $T \subseteq L_{\kappa^+, \omega}$ be a theory of cardinality $\leq \kappa$, and let γ be an ordinal $\geq \kappa^+$. For example we look at

1. $\mu_T^*(\gamma, \kappa) := \min\{\mu^* : \forall \phi \in L_{\infty, \omega}, \text{ with } rk(\phi) < \gamma, \text{ if } T \text{ has the } (\phi, \mu^*)\text{-order property then there exists a formula } \phi'(x; y) \in L_{\kappa^+, \omega}, \text{ such that for every } \chi \geq \kappa, T \text{ has the } (\phi', \chi)\text{-order property}\}$.
2. $\mu^*(\gamma, \kappa) := \sup\{\mu_T^*(\gamma, \kappa) \mid T \in L_{\kappa^+, \omega}\}$.

We discuss several other related functions, sample results are:

- It turns out that if T has the $(\phi, \mu^*(\gamma, \kappa))$ -order property for some $\phi \in L_{\infty, \omega}$, with $rk(\phi) < \gamma$ then for every $\chi > \kappa$ we have that $I(\chi, T) = 2^\chi$ holds.
- For every κ and γ as above there exists an ordinal $\delta^*(\gamma, \kappa)$ such that $\mu^*(\gamma, \kappa) = \beth_{\delta^*(\gamma, \kappa)}$,
- $\delta^*(\gamma, \kappa) \leq (|\gamma|^\kappa)^+$,
- for κ with uncountable cofinality, we have that $\delta^*(\gamma, \kappa) > |\gamma|^\kappa$ and
- the ordinal $\delta^*(\gamma, \kappa)$ is bounded below by the Galvin–Hajnal rank of a reduced product.

For many cardinalities we have better bounds, some of the bounds obtained using Shelah's *PCF theory*. The function $\mu^*(\gamma, \kappa)$ is used to compute bounds to the values of the function $\bar{\mu}(\lambda, \kappa)$ we studied in a previous paper.

1 Introduction

Let χ be an infinite cardinality, and suppose that $T \subseteq L_{\chi,\omega}$ (notice that when $\chi = \omega$ we are dealing with first-order theories).

The fundamental meta-problem in the area of classification theory can be stated as:

Problem 1.1 *What is the structure of $Mod(T)$?*

A more precise (and concrete) test-question is:

Problem 1.2 *What are the possible functions $I(\cdot, T) : Card \rightarrow Card$? (where $I(\lambda, T)$ stands for the number of isomorphism types of models for T of cardinality λ)*

A much more precise (and a very difficult) particular case of 1.2 is the following

Conjecture 1.3 (Shelah about 1976) *Let $\psi \in L_{\omega_1,\omega}$ be given. If there exists a cardinality $\mu > \beth_{\omega_1}$ such that $I(\mu, \psi) = 1$ then for every $\mu > \beth_{\omega_1}$, $I(\mu, \psi) = 1$ holds.*

A possible approach to Problem 1.1 and its relatives, is to try to imitate Classification theory for elementary classes (see [Sh c]). Namely it would be desirable to find properties parallel to *stability*, *superstability* etc. Much work has been done in the last 25 years (see for example – [Sh 48],[Sh 87a],[Sh 87b],[Sh 88],[Sh 300],[MaSh 285],[GrSh 238], or [Sh 299] for a general survey). In this article we concentrate on dealing with the parallel (for infinitary languages) to instability. The following can be viewed as a definition of stability for first-order theories:

Fact 1.4 ([Sh 16]) *Let T be a complete first-order theory. The following are equivalent:*

1. *T is unstable*
2. *There exist a formula $\phi(x; y) \in L(T)$, a model M for T , and a set $\{a_n : n < \omega\} \subseteq M$ such that $l(x) = l(y) = l(a_n)$ for every $n < \omega$, and for all $n, k < \omega$ we have $n < k \iff M \models \phi[a_n; a_k]$.*

Condition 2 in Fact 1.4 is sometimes called the order-property. One of the most important properties of unstable theories is the following:

Fact 1.5 [Sh 12] *Let T be a complete first-order theory. If T is unstable then for every $\mu > |T|$ we have that $I(\mu, T) = 2^\mu$.*

An inspection of the proof of 1.5 shows that the hypothesis that T is a complete first-order unstable theory could be replaced by the following property:

- There is an expansion L' of $L(T)$ with built-in Skolem functions and an L' -structure M , a “formula” $\phi(x; y)$, and there exists $I := \{a_i : i < \omega\} \subseteq M$ a sequence of L' -indiscernibles such that $l(x) = l(y) = l(a_n)$ for every $n < \omega$, M is the Skolem Hull of I , $M \upharpoonright L(T) \models T$, and for every $n, m < \omega$ we have
- $$n < m \iff M \models \phi[a_n; a_m].$$

By “formula” we mean that ϕ is in any logic (over the vocabulary L') such that ϕ is preserved by isomorphisms of $L(T)$ -structures.

The condition in Fact 1.4 seems to be a natural candidate for a definition of instability for infinitary logics. Since the compactness theorem fails even for $L_{\omega_1, \omega}$ the next definition is the replacement of the (the first-order) order-property.

Definition 1.6 *Let $T \subseteq L_{\chi, \omega}$, $\phi(x; y) \in L_{\infty, \omega}$, and let μ be a cardinality.*

1. *We say that M has the (ϕ, μ) -order property iff there exists $\{a_i : i < \mu\} \subseteq M$ such that $l(x) = l(y) = l(a_i) < \omega$, and for every $i, j < \mu$ we have $i < j \iff M \models \phi[a_i; a_j]$.*
2. *T has the (ϕ, μ) -order property iff there exists $M \models T$ such that M has the (ϕ, μ) -order property.*
3. *T has the (ϕ, ∞) -order property iff for every μ , T has the (ϕ, μ) -order property.*
4. *Let λ and μ be cardinalities, we say that T has the $(L_{\lambda, \omega}, \mu)$ -order property iff there exists $\phi \in L_{\lambda, \omega}$ such that T has the (ϕ, μ) -order property.*

Remark 1.7 1. In light of the last definition, Fact 1.4 can be restated as (for first-order complete T): T is unstable iff T has the $(L_{\omega, \omega}, \aleph_0)$ -order property.

2. It is not difficult to see (using [Mo], see 1.10 below) that the following implication is true: If T has the $(L_{\lambda^+, \omega}, \infty)$ -order property then $(*)_T$ holds.

The natural question to ask in this context is: Given a theory T and a cardinality μ , does T have the $(L_{\lambda^+, \omega}, \mu)$ -order property? The main object of study in [GrSh] was the function $\bar{\mu}(\lambda, \kappa)$. The following $\mu^*(\lambda, \kappa)$ is a relative of $\bar{\mu}(\lambda, \kappa)$ from [GrSh].

Definition 1.8 *Let $\kappa \leq \lambda$.*

1. *Let $\psi \in L_{\kappa^+, \omega}$, $\mu_\psi^*(\lambda, \kappa) := \min\{\mu^* : \forall \phi \in L_{\lambda^+, \omega}$ if ψ has the (ϕ, μ^*) -order property, then $\exists \phi'(x; y) \in L_{\kappa^+, \omega}$, such that ψ has the (ϕ', ∞) -order property $\}$.*
2. $\mu^*(\lambda, \kappa) := \sup\{\mu_\psi^*(\lambda, \kappa) \mid \psi(x; y) \in L_{\kappa^+, \omega}\}$.

Remark 1.9 The idea behind Definition 1.8 is that when ψ has the $(L_{\lambda^+, \omega}, \mu^*(\lambda, \kappa))$ -order property then (by Remark 1.7 and $(*)_\psi$) for every $\chi > \kappa$ $I(\chi, \psi) = 2^\chi$.

Already in [Sh 16] Shelah realized the importance of the above concept (it did not appear there explicitly. Only in [Gr] (see [GrSh]) we realized the importance of functions of this kind). The previous definition is a generalization of one of our definitions from [GrSh], see Definition 1.8. Shelah's fundamental result from [Sh 16] can be restated as:

Fact 1.10 ([Sh 16]) *For every $\kappa \leq \lambda$, we have $\mu^*(\lambda, \kappa) \leq \mu_0(\lambda, 1)^1$.*

Let us mention here the following dramatic improvement (for $\kappa = \aleph_0$) of 1.10:

Theorem 1.11 ([GrSh]) *For every $\lambda \geq \aleph_0$, we have $\mu^*(\lambda, \aleph_0) \leq \beth_{\lambda^+}$.*

It turns out that even for first-order theories the above question is interesting (for $\kappa = |L| = \aleph_0$, T is a complete first-order theory in L , we could ask what is an upper bound of $\mu_T^*(\lambda, \aleph_0)$?). Since there are cases when T is stable (i.e. there is no first-order formula defining an ω -sequence in a model of T) but still T has a hidden instability (like in the case of stable theories with the omitting-types order property).

Namely there is a natural class of examples of theories that do not have a first-order formula exemplifying the order-property but do have an infinitary order property. Any stable first order theory that has the omitting types

¹ $\mu_0(\lambda, \lambda)$ is the usual Morley number to be introduced in Definition 2.2 below. It is easy to see that $\mu_0(\lambda, \lambda) = \mu_0(\lambda, 1)$

order-property has the $(L_{\omega_1, \omega}, \infty)$ -order property but not the $(L_{\omega, \omega}, \aleph_0)$ -order property (see [Sh 200]).

Already from Morley's omitting-types theorem it follows that given T and ϕ as above there exists $\mu := \mu(T, \phi)$ such that if T has the (ϕ, μ) -order property then $\forall \lambda \geq \chi$, T has the (ϕ, λ) -order property. The bound obtained from repeating the argument in the proof of Morley's omitting types theorem (see [Sh 16]) is: $\mu(T, \phi) \leq \max\{Hanf(T), Hanf(\phi)\}$. Where $Hanf(T)$ and $Hanf(\phi)$ are the Hanf numbers of T and the logic containing ϕ (respectively).

Let $\chi > \aleph_0$ (T still may be first-order). Our object is to find upper bounds on μ . It turns out that for $\phi \in L_{\infty, \omega} - L_{\chi, \omega}$ there is a cardinality $\mu^* := \mu^*(T, \phi)^2$, such that the following implication holds: If T has the (ϕ, μ^*) -order property then there exists a formula $\phi' \in L_{\chi, \omega}$ (it is a collapse of ϕ) such that T has the (ϕ', λ) -order property for every $\lambda \geq \chi$.

In this paper we present a systematic study of several cardinal and ordinal valued functions related to the infinitary order property. This is a continuation of [GrSh], we deal with similar problems and improve many of our results. This is achieved via a generalization of the original problem (dealing with new cases) while obtaining often better estimates to our earlier bounds. The reader is not expected to be familiar with [GrSh].

Notation: Everything is standard. Often x, y , and z will denote free variables or finite sequences of variables, when x is a sequence $l(x)$ denote its length. It should be clear from the context whether we deal with variables or sequences of variables. L will denote a similarity type (also known as-language or signature), Δ will stand for a set of L formulas. M and N will stand for L -structures, $|M|$ the universe of the structure M , $\|M\|$ the cardinality of the universe of M . Given a fixed structure M , subsets of its universe will be denoted by A, B, C , and D . So when we write $A \subseteq M$ we really mean that $A \subseteq |M|$, while $N \subseteq M$ stands for “ N is a submodel of M ”. Let M be a structure. By $a \in M$ we mean $a \in |M|$, when a is a finite sequence of elements then $a \in M$ stands for “all the elements of the sequence a are elements of $|M|$ ”. For cardinalities $\kappa \leq \lambda$, let $S_{<\kappa}(\lambda) := \{X \subseteq \lambda : |X| < \kappa\}$. When T is a first-order theory, Γ denotes a set of T -types over the empty set (not necessarily complete types). $EC(T, \Gamma) := \{M : M \models T, \forall p \in \Gamma M \text{ omits the type } p\}$. When T is first-order, $L \subseteq L(T)$, and Γ is a set of T -types by $PC(T, \Gamma, L)$ we denote the following $\{M \upharpoonright L : M \models T, \forall p \in \Gamma M \text{ omits the type } p\}$; namely

² The surprise is that often $\mu^*(T, \phi)$ is much smaller than $\mu(T, \phi)$.

$EC(T, \Gamma) = PC(T, \Gamma, L(T))$. λ, μ, κ , and χ will stand for infinite cardinalities; $\alpha, \beta, \gamma, \delta, \zeta$, and ξ are ordinals. References of the form “Theorem IV 3.12” are to [Sh c]. For $\phi \in L_{\infty, \omega}$, let $Sub(\phi)$ be the set of subformulas of ϕ , now let

$$rk(\phi) := \begin{cases} 0 & \text{if } \phi \text{ is atomic} \\ \text{Sup}\{rk(\chi) + 1 : \chi \in Sub(\phi)\} & \text{otherwise.} \end{cases}$$

2 Review

C.C. Chang in [Ch] made the following fundamental observation:

Fact 2.1 *Let κ be an infinite cardinality, and let L be a similarity type of cardinality no more than κ . Given $\psi \in L_{\kappa^+, \omega}$, there exist a similarity type $L' \supseteq L$, a first-order theory T in L' , and a set of T -types Γ (all three of cardinality less than or equal to κ) such that $\text{Mod}(\psi) = \text{PC}(T, \Gamma, L)$.*

Namely instead of studying $\text{Mod}(\psi)$ directly for an infinitary theory ψ it is enough to look at a class of reducts of models of a first-order theory that omits a set of types.

W. Hanf and M. Morley [Mo], recognized the importance of the following concept:

Definition 2.2 *Let T be a first-order theory, and let Γ be a set of T -types. The Morley number³ of T and Γ , is the following:*

1. $\mu_0(T, \Gamma) := \min\{\mu : \exists M \in \text{EC}(T, \Gamma) \ \|M\| \geq \mu \Rightarrow \forall \chi \geq |T| \ \exists N \in \text{EC}(T, \Gamma) \text{ of cardinality } \geq \chi\}$.
2. Let λ, κ be cardinalities.
 $\mu_0(\lambda, \kappa) := \sup\{\mu_0(T, \Gamma) : |T| \leq \lambda, \Gamma \text{ a set of } T\text{-types of cardinality } \leq \kappa\}$ ⁴.

Morley (among other things) have shown that $\mu_0(\aleph_0, \aleph_0) = \beth_{\omega_1}$. His most general result is stated as Theorem 2.4 below. Shelah in [Sh 78] have dealt with what can be viewed to be an interpolant of $\mu_0(T, \Gamma)$ and $\mu_0(\lambda, \kappa)$:

$$\mu_0(T, \kappa) := \sup\{\mu_0(T, \Gamma) : \Gamma \text{ a set of } T\text{-types, } |\Gamma| \leq \kappa\}.$$

It is not difficult to conclude from the proof of Morley's categoricity theorem that when T is a countable and \aleph_0 -stable theory then $\mu_0(T, \cdot) \leq \aleph_1$. Shelah in [Sh 78] studied the effect that stability of T has on the upper bounds on $\mu_0(T, \kappa)$. This work was continued about ten years later by Hrushovski and Shelah in [HrSh 334].

In this paper, since our main goal is the study of unstable theories (or theories that are not stable in a weak sense) we will ignore the effect that the stability of T may have on the function $\mu_0(T, \kappa)$.

³Some authors call this the Hanf number of T and the Γ

⁴We hope that the reader is not bothered by this abuse of notation. We are using the same letter μ_0 to denote entirely different (but related) functions. They can be distinguished by the type of the arguments they take.

The modern era in the study of Hanf numbers began with the paper of Barwise and Kunen [BaKu]. They studied systematically the relationship between the function μ_0 and the first ordinal that exemplify the undefinability of well ordering in classes of models that omit a set of types. Below we introduce an ordinal-valued function $\delta_0(\lambda, \kappa)$ that turns out to be related to $\mu_0(\lambda, \kappa)$ in a nice way.

Definition 2.3 *Let λ and κ be infinite cardinalities, T varies over consistent first-order theories such that $L(T) \supseteq \{P, <\}$ when P is a unary predicate and $T \vdash \text{“} < \text{ linearly orders } P \text{”}$.*

$\delta_0(\lambda, \kappa) := \min\{\delta : |T| \leq \lambda, \Gamma$ a set of T -types, $|\Gamma| \leq \kappa$ if for every $\delta' < \delta$ there exists $M \in EC(T, \Gamma)$ such that $otp(P^M, <^M) \geq \delta'$ then there exists $N \in EC(T, \Gamma)$ s.t. $(P^N, <^N)$ is not well ordered $\}$.

The following is a restatement of Morley’s “other” important theorem:

Fact 2.4 (Theorem VII 5.5) $\mu_0(\lambda, \kappa) = \beth_{\delta_0(\lambda, \kappa)}$.

The following ordinal and cardinal-valued functions are from §4 of [GrSh]:

Definition 2.5 *Suppose T is a first-order theory such that $L(T)$ contains $\{<, P\}$ and*

$T \vdash [< \text{ is a linear order}] \wedge [< \upharpoonright P \text{ is a linear order on the unary predicate } P]$.

1. $\delta_1(\theta, \lambda, \kappa) := \min\{\delta : \Gamma$ a set of T -types, $|\Gamma| \leq \lambda, |T| \leq \kappa$ if $\forall \delta' < \delta \exists M \in EC(T, \Gamma)$ with $otp(P^M, <^M) \in On \cap \theta^+$ and $otp(M - P, <^M) \geq \delta'$, then $\exists N \in EC(T, \Gamma)$ s.t. $otp(P^N, <^N) \in On \cap \kappa^+$ and $(N - P^N, <^N)$ is not well ordered $\}$.
2. $\mu_1(\theta, \lambda, \kappa) := \min\{\mu : |\Gamma| \leq \lambda, |T| \leq \kappa$ Γ a set of T -types, if $\exists M \in EC(T, \Gamma)$ $\|M\| \geq \mu$ with $otp(P^M, <^M) \in On \cap \theta^+$ then for every $\chi \geq \kappa \exists N \in EC(T, \Gamma)$ of cardinality at least χ such that $otp(P^N, <^N) \in On \cap \kappa^+$ $\}$.
3. When $\theta = \lambda$ we will omit the first parameter- θ

We will prove the following equality:

Theorem 2.6 *For every $\kappa \leq \lambda \leq \theta$ we have $\mu_1(\theta, \lambda, \kappa) = \beth_{\delta_1(\theta, \lambda, \kappa)}$.*

From now on we concentrate on the case that $\theta = \lambda$ and we will work with the functions $\delta_1(\lambda, \kappa)$ and $\mu_1(\lambda, \kappa)$. The arguments for the functions with three parameters are essentially similar (they require an additional technical effort, but require no new ideas). Note that by [Ch] working with two parameter functions is sufficient for $L_{\lambda^+, \omega}$. The new point is that we are able to show that $\mu_1(\lambda, \kappa) \geq \beth_{\delta_1(\lambda, \kappa)}$. The proof of Theorem 2.6 is similar to that of Theorem VII 5.5, we skip its proof, since later we will prove a related theorem (Th. 2.25) whose proof is similar (and little harder).

The next proposition provides us with a lower bound for $\delta_1(\lambda, \kappa)$, it follows immediately from the definitions (in Theorem 2.27 we show a better lower bound).

Proposition 2.7 *For $\lambda \geq \kappa$ we have that $\delta_1(\lambda, \kappa) \geq \delta_1(\kappa, \kappa) \geq \delta_0(\kappa, \kappa) = \delta_0(\kappa, 1)$.*

In the following proposition the connection between the last definition and the order property is clarified.

Theorem 2.8 *Let $\kappa \leq \lambda$, be cardinalities. $\beth_{\lambda^+} \leq \mu^*(\lambda, \kappa) \leq \mu_1(\lambda, \kappa)$.*

Proof. First we show that $\mu^*(\lambda, \kappa) \leq \mu_1(\lambda, \kappa)$. Let $\psi \in L_{\kappa^+, \omega}$, and $\phi(x; y) \in L_{\lambda^+, \omega}$ be given. Suppose ψ has the $(\phi, \mu_1(\lambda, \kappa))$ -order property we need to find a formula $\phi' \in L_{\kappa^+, \omega}$ such that ψ has the (ϕ', ∞) -order property.

By Fact 2.1 there exists a first-order theory T in a similarity type $L(T)$ that extends L , and there is a set Γ of T -types of cardinality $\leq \kappa$ such that $PC(T, \Gamma, L) = Mod(\psi)$. By following the inductive definition of the formula ϕ we may identify ϕ with a function f from the set P into the set $L \cup \{\wedge, \neg, (,), =\} \cup \{x_i : i < \kappa\}$.

Let χ be a regular large enough such that

$$\{L_{\lambda^+, \omega}, L_{\kappa^+, \omega}, T, \Gamma, L, \phi, f, P, \psi, \lambda^+, \mu_1(\lambda, \kappa), \delta_1(\lambda, \kappa)\} \cup \mu_1(\lambda, \kappa) \subseteq H(\chi).$$

In addition we require that the structure $\langle H(\chi), \in \rangle$ reflects all the relevant properties of the above sets. Let P be the rank of the formula ϕ , note that it is an ordinal less than λ^+ . Let $\mathfrak{A}' \prec \langle H(\chi), \in, \dots \rangle$ of cardinality $\mu_1(\lambda, \kappa)$ such that $\mu_1(\lambda, \kappa)^{\mathfrak{A}'} = \mu_1(\lambda, \kappa)$ (so $\mu_1(\lambda, \kappa) + 1 \subseteq \mathfrak{A}'$), fix a bijection G from $\mu_1(\lambda, \kappa)$ onto the universe of \mathfrak{A}' , and let $\mathfrak{A} := \langle \mathfrak{A}', G \rangle$. By the definition of $\mu_1(\lambda, \kappa)$, for every $\chi \geq \kappa$ there exists $\mathfrak{B}_\chi \equiv \mathfrak{A}$ of cardinality χ such that \mathfrak{B}_χ omits the types from Γ , $\kappa^{\mathfrak{B}_\chi} = \kappa$, and P is an ordinal less than κ^+ (just apply the Mostowski collapse on \mathfrak{B}_χ). Using $P^{\mathfrak{B}_\chi}$, and $f^{\mathfrak{B}_\chi}$ we know (in

\mathfrak{B}_χ) that $\phi^{\mathfrak{B}_\chi} \in L_{\kappa^+, \omega}^{\mathfrak{B}_\chi}$, but since $\kappa^{\mathfrak{B}_\chi} = \kappa$ we have that $\phi^{\mathfrak{B}_\chi} \in L_{\kappa^+, \omega}$ is a formula as required in the definition of $\mu^*(\lambda, \kappa)$.

To see that $\mu^*(\lambda, \kappa) \geq \beth_{\lambda^+}$: It is enough to show that for every $\alpha < \lambda^+$ there exist a sentence $\psi_\alpha \in L_{\kappa^+, \omega}$, and a formula $\phi_\alpha \in L_{\lambda^+, \omega}$ such that ψ_α has the $(\phi_\alpha, \beth_\alpha)$ -order property and ψ_α does not have the $(L_{\kappa^+, \omega}, \infty)$ -order property.

Before proving this we will introduce several definitions.

Notation: The sentence ψ_α will be defined as a the theory of a well founded tree. We deal with well-founded trees whose vertices are decreasing sequences of ordinals, the root of the tree \mathcal{T} is denoted by $rt(\mathcal{T})$, for an element $x \in \mathcal{T}$ let $Suc_{\mathcal{T}}(x)$ stand for the set of immediate successors of x , and $\mathcal{T}[x]$ stands for the subtree of \mathcal{T} consisting of the elements that are greater than or equal to x .

Definition 2.9 *Let \mathcal{T} be a well founded tree.*

1. For $x \in \mathcal{T}$ let $Dp_{\mathcal{T}}(x) = \beta$ the depth of x in \mathcal{T} defined by induction on β :
 - (a) if $Suc_{\mathcal{T}}(x) = \emptyset$ then $Dp_{\mathcal{T}}(x) = 0$.
 - (b) if for every $y \in Suc_{\mathcal{T}}(x)$ we have $Dp_{\mathcal{T}}(y) < \beta$, and for every $\gamma < \beta$ there exists $z \in Suc_{\mathcal{T}}(x)$ of such that $Dp_{\mathcal{T}}(z) \geq \gamma$ then $Dp_{\mathcal{T}}(x) = \beta$.
2. The depth of \mathcal{T} is $Dp(\mathcal{T}) := \sup\{Dp_{\mathcal{T}}(x) : x \in \mathcal{T}\}$.

Proposition 2.10 *Let \mathcal{T} be a well-founded tree, $Dp(\mathcal{T}) = Dp_{\mathcal{T}}(rt(\mathcal{T}))$.*

Proof. Trivial. □_{2.10}

Claim 2.11 *For every α there exists a well-founded tree \mathcal{T}_α of depth α such that $\|\mathcal{T}_\alpha\| \leq |\alpha| + \aleph_0$.*

Proof. By induction on α :

For $\alpha = 0$; Simply let $\mathcal{T}_0 := \langle \rangle$.

For $\alpha = \beta + 1$; Suppose \mathcal{T}_β is a tree of depth β .

Let $\mathcal{T}_\alpha := \{\langle \rangle\} \cup \{\langle \rangle \hat{\ } \eta : \eta \in \mathcal{T}_\beta\}$. The order on \mathcal{T}_α is the obvious.

For α a limit ordinal; By the induction hypothesis let $\{\mathcal{T}_\beta : \beta < \alpha\}$ be pairwise disjoint trees, each of depth β .

Define \mathcal{T}_α to be the tree $\{\langle \rangle\} \cup \{\langle \rangle \hat{\ } \eta : \eta \in \mathcal{T}_\beta, \beta < \alpha\}$. □_{2.11}

Definition 2.12 1. Let $\mathcal{T}_1, \mathcal{T}_2$ be well-founded trees, and let α be an ordinal. By induction on α define when $\mathcal{T}_1 \approx_\alpha \mathcal{T}_2$:

- (a) For $\alpha = 0$, always $\mathcal{T}_1 \approx_\alpha \mathcal{T}_2$.
- (b) For $\alpha \neq 0$, if for every $\beta < \alpha$ and for every $x_1 \in \text{Suc}_{\mathcal{T}_1}(\text{rt}(\mathcal{T}_1))$ there exists $x_2 \in \text{Suc}_{\mathcal{T}_2}(\text{rt}(\mathcal{T}_2))$ such that $\mathcal{T}_1[x_1] \approx_\beta \mathcal{T}_2[x_2]$, and for every $x_2 \in \text{Suc}_{\mathcal{T}_2}(\text{rt}(\mathcal{T}_2))$ there exists $x_1 \in \text{Suc}_{\mathcal{T}_1}(\text{rt}(\mathcal{T}_1))$ such that $\mathcal{T}_2[x_2] \approx_\beta \mathcal{T}_1[x_1]$.

- 2. A tree \mathcal{T} is called simple iff there are no distinct $x_1, x_2 \in \text{Suc}_{\mathcal{T}}(\text{rt}(\mathcal{T}))$ such that $Dp_{\mathcal{T}}(x_1) = Dp_{\mathcal{T}}(x_2)$ and $\mathcal{T}[x_1] \approx_{Dp_{\mathcal{T}}(x_1)} \mathcal{T}[x_2]$.

Proposition 2.13 Let $\mathcal{T}_1, \mathcal{T}_2$ be trees, and let α be an ordinal. If $\mathcal{T}_1 \approx_\alpha \mathcal{T}_2$ then one of the following conditions holds:

- 1. $Dp(\mathcal{T}_1) = Dp(\mathcal{T}_2)$, or
- 2. $Dp(\mathcal{T}_1) \geq \alpha$ and $Dp(\mathcal{T}_2) \geq \alpha$.

Proof. Easy, by induction on α . □_{2.13}

Claim 2.14 For every ordinal α there exists a family of simple trees $\{\mathcal{T}_i : i < \beth_\alpha\}$, such that for every $i < \beth_\alpha$

- 1. $\|\mathcal{T}_i\| \leq \beth_\alpha$,
- 2. $i \neq j \Rightarrow \mathcal{T}_i \not\approx_{\omega+\alpha} \mathcal{T}_j$,
- 3. $Dp(\mathcal{T}_i) = \omega + \alpha + 1$.

Proof. By induction on α :

For $\alpha = 1$; First construct \aleph_0 simple trees $\{\mathcal{T}_n \subseteq^{>\omega} \omega : n < \omega\}$ such that $\mathcal{T}_0 := \{\langle \rangle\}$, $\mathcal{T}_{n+1} := \{\langle n+1 \rangle\} \cup \mathcal{T}_n$ when the order is an extension of the order on \mathcal{T}_n ; $\langle \rangle$ is the root, and $\langle n+1 \rangle$ is a new immediate successor of the root incomparable with the elements of $\text{Suc}_{\mathcal{T}_n}(\langle \rangle)$. Now for every $A \subset \omega$ let

$$\mathcal{T}_A := \{\langle \omega \rangle \hat{\ } \eta : \eta \in \mathcal{T}_n, n \in A\} \cup \{\nu \in \mathcal{T}_k : k \notin A\}.$$

The order of \mathcal{T}_A is defined as follows: $\langle \omega \rangle$ is a new immediate successor of the root, the elements $\langle \omega \rangle \hat{\ } \eta$ and ν are pairwise incomparable when $\eta \in \mathcal{T}_n$, ($n \in A$) and $\nu \in \mathcal{T}_k$, ($k \notin A$), and we require that

$$\langle \omega \rangle \hat{\ } \eta_1 < \langle \omega \rangle \hat{\ } \eta_2 \Leftrightarrow \eta_1 <_{\mathcal{T}_n} \eta_2.$$

In order to see that $A \neq B \subseteq \omega \Rightarrow \mathcal{T}_A \not\approx_{\omega+\alpha} \mathcal{T}_B$: W.l.o.g. we may assume

that $\exists n \in A - B$. Since $\mathcal{T}_A[\langle \omega, n \rangle] \not\approx_\omega \mathcal{T}_B[\nu]$ for any $\nu \in \text{Suc}_{\mathcal{T}_B}(\langle \omega \rangle)$ (this is because \mathcal{T}_k and \mathcal{T}_n are inequivalent for $k \neq n$).

For $\alpha \neq 1$; By the inductive hypothesis let $\{\{\mathcal{T}_i^\beta : i < \beth_\beta\} : \beta < \alpha\}$ be disjoint trees satisfying the statement of the Theorem. Denote by S the set $\{\langle \beta, i \rangle : i < \beth_\beta, \beta < \alpha\}$. Fix an injective mapping from S into $On - \text{Sup}(\bigcup_{\beta, i} \mathcal{T}_i^\beta)$, denote by $\gamma_{\beta, i}$ the image of the pair $\langle \beta, i \rangle$. For every $\gamma < \alpha$, and for every $A \subseteq S$ cardinality \beth_γ define

$$\mathcal{T}_A := \{\langle \rangle\} \cup \{\langle \gamma_{\beta, i} \rangle \hat{\ } \eta : \eta \in \mathcal{T}_i^\beta, \langle \beta, i \rangle \in A\}.$$

The order on \mathcal{T}_A is defined in the natural way: $\langle \rangle$ is the root, and $\langle \beta, i \rangle \hat{\ } \eta_1 <_{\mathcal{T}_A} \langle \beta, i \rangle \hat{\ } \eta_2$ iff $\eta_1 <_{\mathcal{T}_i^\beta} \eta_2$. The verification that $Dp(\mathcal{T}_A) = \omega + \alpha + 1$ is left to the reader. Suppose $A \neq B \subseteq \beth_\alpha$ both of cardinality \beth_β for some $\beta < \alpha$. We need to show that $\mathcal{T}_A \not\approx_{\omega + \alpha} \mathcal{T}_B$. W.l.o.g. there exists $\gamma_{\beta, i} \in A - B$. Since $\{\langle \gamma_{\beta, i} \rangle \hat{\ } \eta : \eta \in \mathcal{T}_i^\beta\}$ is a subtree of \mathcal{T}_A and for all $j < \beth_\beta$ we have that $j \neq i \Rightarrow \mathcal{T}_i^\beta \not\approx_{\omega + \beta} \mathcal{T}_j^\beta$, from the definition of the relation \approx , and the fact that it follows that there is no ordinal ϵ such that the tree $\{\epsilon \hat{\ } \eta : \eta \in \mathcal{T}_i^\beta\}$ does not appear as a subtree of \mathcal{T}_B it is clear that $\mathcal{T}_A \not\approx_{\omega + \alpha} \mathcal{T}_B$. □_{2.14}

Back to the proof of Theorem 2.8: Let $\alpha < \lambda^+$ be given. By Claim 2.14 there exists a family of nonequivalent simple trees $\{\mathcal{T}_i : i < \beth_\alpha\}$. By renaming, we may assume that the above trees do not contain sequences of ordinals which are less than \beth_α . We define a new tree: M_α its set of elements consists of

$$\{\langle \rangle\} \cup \{\langle i \rangle : i < \beth_\alpha\} \cup \{\langle i, i' \rangle \hat{\ } \eta : \eta \in \mathcal{T}_{i'}, i' < i, i < \beth_\alpha\}.$$

We can view M_α as a partially ordered set (by “being initial segment”). We view M_α as a model in a language consisting of single function symbol: A unary function f whose interpretation is the predecessor of its argument (if the argument is the root than the value is defined to be the root). Notice that the following formula (of $L_{\omega_1, \omega}$), $\bigvee_{k < \omega} [x = f^k(y)]$ ⁵ defines the relation of “being an initial segment” on well founded trees.

Let $\psi_\alpha := \bigwedge Th_{\omega, \omega}(M_\alpha) \wedge (\forall x) \bigvee_{n < \omega} [f^n(x) = \langle \rangle]$. Namely ψ_α is the first-order theory of M_α together with the statement that say that every element is of finite distance from the root.

Let $\phi_\alpha(x, y)$ be the following statement: $x, y \in \text{Suc}(\langle \rangle)$, and for every $x' \in \text{Suc}(x)$ there exists $y' \in \text{Suc}(y)$ such that $\mathcal{T}[x'] \approx_{\omega + \alpha + 1} \mathcal{T}[y']$, and

⁵ When $f^k(y)$ stands for $f(\dots f(y)\dots)$ k -many times.

there exists $y' \in \text{Suc}(y)$ such that for every $x' \in \text{Suc}(x)$ we have that $\mathcal{T}[x'] \not\approx_{\omega+\alpha+1} \mathcal{T}[y']$ holds.

In order to complete the proof of Theorem 2.8, it suffices to prove the following:

- SubClaim 2.15**
1. $\phi_\alpha(x, y) \in L_{\lambda^+, \omega}$,
 2. M_α has the $(\phi_\alpha, \beth_\alpha)$ -order property,
 3. There do not exist a formula $\phi'(x, y) \in L_{\kappa^+, \omega}$ such that ψ_α has the $(\phi'(x, y), \infty)$ -order property.

Proof.

1. Let \mathcal{T} be a well founded tree, and let $\alpha < \lambda^+$ be given, it is enough to show by induction on α that there exists a formula $\chi(x, y) \in L_{\lambda^+, \omega}$ such that for every $a, b \in \mathcal{T}$ we have that $\mathcal{T} \models \chi[a, b]$ iff $\mathcal{T}[a] \approx_\alpha \mathcal{T}[b]$. It is easy to check that the relation \approx_α is definable in $L_{\lambda^+, \omega}$.
2. Check that for every $i_1, i_2 < \beth_\alpha$ we have that $i_1 < i_2$ iff $M_\alpha \models \phi_\alpha[\langle i_1 \rangle, \langle i_2 \rangle]$.
3. For the sake of contradiction suppose that there exists a formula $\phi'(x, y) \in L_{\kappa^+, \omega}$ such that ψ_α has the (ϕ', ∞) -order property. Suppose that γ is a limit ordinal $< \kappa^+$ such that the formula ϕ' has quantifier depth $< \gamma$. Denote by μ the cardinality $(\beth_{\gamma+1}(|L|))^+$. Let $N \models \psi_\alpha$ be a model of cardinality μ such that there exists $\{a_i : i < \mu\}$ such that $l(x) = l(y) = l(a_i) = n < \omega$ and for every $i_1, i_2 < \mu$ we have $i_1 < i_2 \iff N \models \phi'[a_{i_1}, a_{i_2}]$ holds. For every $i < \mu$ fix $\langle b_l^i : l < n \rangle = a_i$. By the $L_{\omega_1, \omega}$ -part of the definition of ψ_α we have that $N \models (\forall x) \bigvee_{m < \omega} f^m(x) = f^{m+1}(x)$. For every $c \in N$ let $m(c) := \min\{m : N \models f^m(c) = f^{m+1}(c)\}$.

Since μ is regular, after renaming we may assume that for every $l < n$ there are $k_l < \omega$ such that for every $i < \mu$ we have $m(b_l^i) = k_l$. By increasing n we may assume that for every $i < \mu$ we have that $f(b_l^i) \in \{b_k^i : k < n\}$, and for every $i_1, i_2 < \mu$ and every $l_1, l_2 < n$ we have

$$N \models f(b_{l_1}^{i_1}) = b_{l_2}^{i_1} \iff N \models f(b_{l_1}^{i_2}) = b_{l_2}^{i_2} \bigwedge N \models b_{l_1}^{i_1} = b_{l_2}^{i_1} \iff N \models b_{l_1}^{i_2} = b_{l_2}^{i_2}.$$

We may also assume that $\langle b_l : l < n \rangle$ has no repetition. By the Δ -system lemma there exists $s \subseteq n$ such that for every $i_1, i_2 < \mu$ and every $l_1, l_2 < n$ we have that $b_{l_1}^{i_1} = b_{l_2}^{i_2} \iff l_1 = l_2 \in s$.

Let Φ_γ be the set of $L_{\infty,\omega}$ formulas of quantifier depth $< \gamma$ with finitely many free variables. Clearly $|\Phi_\gamma| \leq \beth_\gamma(|L|)$ and $|\mathcal{P}(\Phi_\gamma)| \leq \beth_{\gamma+1}(|L|) < \mu = cf(\mu)$.

Let $tp_\gamma(b_0, \dots, b_{m-1}; M) := \{\phi(\bar{x}) \in \Phi_\gamma : M \models \phi[b_0, \dots, b_{m-1}]\}$. Without loss of generality we may assume that for every $i, j < \mu$ we have $tp_\gamma(b_0^i, \dots, b_{m-1}^i; N) = tp_\gamma(b_0^j, \dots, b_{m-1}^j; N)$.

We obtain a contradiction to the assumption that ψ_α has the (ϕ', ∞) -order property by proving the following:

Claim 2.16 *For every $i, j < \mu$ we have*

$$N \models \phi'[b_0^i, \dots, b_{n-1}^i, b_0^j, \dots, b_{n-1}^j] \iff N \models \phi'[b_0^j, \dots, b_{n-1}^j, b_0^i, \dots, b_{n-1}^i]$$

Proof. Left to the reader. □_{2.15}

Remark 2.17 In Definition 2.5 we have introduced a third parameter, but since it does not add anything of substance (just complicates the notation that may be already little heavy) we decided to limit our treatment to the above particular case. At the end of this section we discuss several generalizations.

Theorem 2.8 provides a better upper bound than the one in Fact 1.10:

Corollary 2.18 *For every $\kappa \leq \lambda$, we have $\mu^*(\lambda, \kappa) \leq \beth_{\delta_1(\lambda, \kappa)}$.*

Remark 2.19 Using Facts 1.10 and 2.18, one can show that for $\kappa \leq \lambda$ we have $\delta_1(\lambda, \kappa) \leq \delta_0(\lambda, \kappa)$. In [GrSh] we have shown that in many instances the ordinal $\delta_1(\lambda, \kappa)$ is much smaller than $\delta_0(\lambda, \lambda)$ [e.g. when $\kappa = \aleph_0$, we have that $\delta_1(\lambda, \aleph_0) = \lambda^+$, while for $\lambda = \beth_{\omega_1}$, we have $\delta_0(\lambda, \lambda) > 2^\lambda$.]

In Theorem 2.6 we reduced the problem of finding estimates for $\mu_1(\cdot, \cdot)$ to finding bounds for $\delta_1(\cdot, \cdot)$. In Fact 2.21, below we state an important result from [GrSh], first we need the following:

Definition 2.20 *For uncountable κ , and $\lambda \geq \kappa$, denote by*

$$\kappa^* := \begin{cases} \kappa & \text{if } cf\kappa = \aleph_0 \\ \kappa^+ & \text{if } cf\kappa > \aleph_0. \end{cases}$$

$cov(\lambda, \kappa) := \min\{|F| : F \subseteq S_{<\kappa^*}(\lambda), \forall X \in S_{<\kappa^*}(\lambda) \exists \{w_l : l < \omega\} \subseteq F, \text{ such that } X \subseteq \bigcup_{l < \omega} w_l\}$.

Clearly $cov(\lambda, \kappa) \leq \lambda^\kappa$. But often $cov(\lambda, \kappa) < \lambda^\kappa$. In [Sh g] Shelah has a more general function. Our $cov(\lambda, \kappa)$ is the same as $cov(\lambda, \kappa^*, \kappa^*, \aleph_1)$ from Definition II 5.2 of [Sh g].

Fact 2.21 (Theorem 4.4 of [GrSh]) Let $\kappa \leq \lambda$ be infinite cardinalities.

1. if $\kappa = \aleph_0$ then $\delta_1(\lambda, \kappa) \leq \lambda^+$.
2. if $cf\kappa > \aleph_0$ then $\delta_1(\lambda, \kappa) \leq (cov(\lambda, \kappa) + 2^\kappa)^+$.
3. if $cf\kappa = \aleph_0$ then $\delta_1(\lambda, \kappa) \leq (cov(\lambda, \kappa) + 2^{<\kappa} + \aleph_0)^+$.

Note that the above innocent looking results are quite powerful! E.g. By a result of [Sh g] (from Chapter XI), if $(\forall \mu < \chi)[\mu^\kappa < \lambda] \wedge cf(\chi) = \aleph_0 \wedge \chi \leq \lambda < \chi^{\delta+\omega_1}$ then we have that $cov(\lambda, \kappa) = \lambda$, thus $\mu_1(\lambda, \kappa) \leq \beth_{\lambda^+}$, while using Morley's methods we get only $\mu_1(\lambda, \kappa) \leq \beth_{(2^\lambda)^+}$.

The following is a generalization of the cardinal-valued function we have introduced in Definition 1.8. Here instead of assuming that $\phi(x; y)$ is an $L_{\lambda^+, \omega}$ formula we look at all $\phi \in L_{\infty, \omega}$ with quantifier depth $< \gamma$, we take into consideration only the rank of the formula ϕ .

Definition 2.22 Let κ be an infinite cardinality, and let γ be an ordinal greater or equal to κ^+ , $T \in L_{\kappa^+, \omega}$

1. $\mu_T^*(\gamma, \kappa) := \min\{\mu^* : \forall \phi \in L_{\infty, \omega}, \text{ with } rk(\phi) < \gamma, \text{ if } T \text{ has the } (\phi, \mu^*)\text{-order property, then } \exists \phi'(x; y) \in L_{\kappa^+, \omega}, \text{ such that } T \text{ has the } (\phi', \infty)\text{-order property}\}$.
2. $\mu_2^*(\gamma, \kappa) := \sup\{\mu_T^*(\gamma, \kappa) \mid T \in L_{\kappa^+, \omega}\}$ ⁶.

The improvement in comparison to what we have seen before is that instead of limiting attention to formulas with the order-property to be from $L_{\lambda^+, \omega}$ we consider what may look as a weaker order-property, by considering formulas with the order property to be from the logic $L_{\infty, \omega}$ (with rank bounded by γ).

Definition 2.23 Let $T, <, <^P, P$ be as in Definition 2.5. For an ordinal $\gamma > \kappa$ let

⁶Note that similarly to what we did in the previous section with the function $\mu_0(\cdot, \cdot)$ above, the functions $\mu^*(\gamma, \kappa)$ and $\mu^*(\lambda, \kappa)$ are different objects, we distinguish between them by using different arguments.

1. $\delta_2(\lambda, \gamma, \kappa) := \min\{\delta : \Gamma \text{ is a set of } T\text{-types}, |\Gamma| \leq \kappa, |T| \leq \lambda$
if $\forall \delta' < \delta \exists M \in EC(T, \Gamma)$ with $otp(P^M, <^{P^M}) < \gamma$ and $otp(M - P, <) \geq \delta'$, *then* $\exists N \in EC(T, \Gamma)$ s.t. $otp(P^N, <^{P^N}) \in On \cap \kappa^+$ and $(N - P^N, <^N)$ is not well ordered}.
2. $\mu_2(\lambda, \gamma, \kappa) := \min\{\mu : \Gamma \text{ is a set of } T\text{-types}, |\Gamma| \leq \kappa, |T| \leq \lambda$ *if* $\exists M \in EC(T, \Gamma)$ $\|M\| \geq \mu$ with $otp(P^M, <^M) < \gamma$ *then for every* $\chi \geq \kappa \exists N \in EC(T, \Gamma)$ of cardinality at least χ such that $otp(P^N, <^N) \in On \cap \kappa^+$ }.
3. When $\lambda = \kappa$ we may omit λ . By the discussion after Theorem 2.6 this case is interesting enough.

The following is an analog of Proposition 2.8:

Proposition 2.24 *Let κ and μ be cardinalities, and let γ be an ordinal $\geq \kappa$. Then (1) \Rightarrow (2) \Rightarrow (3) where*

1. $\mu \geq \mu_2(\gamma, \kappa)$
2. for every $\psi \in L_{\kappa^+, \omega}$, and for every $\phi(x; y) \in L_{\infty, \omega}$ of quantifier depth $< \gamma$ *if* ψ has the (ϕ, μ) -order property *then there exists* $\phi' \in L_{\kappa^+, \omega}$ such that ψ has the (ϕ', ∞) -order property.
3. $\mu \geq \beth_\gamma$.

Theorem 2.25 *For every κ and every ordinal $\gamma \geq \kappa$ we have $\mu_2(\gamma, \kappa) = \beth_{\delta_2(\gamma, \kappa)}$.*

Theorem 2.25 will be proved in the next section.

The following theorem connects δ_2 to the Galvin–Hajnal rank and provides a lower bound for $\delta_2(\gamma, \kappa)$:

Theorem 2.26 (A lower bound): *Suppose κ is an uncountable regular cardinality. Let J be the ideal of nonstationary subsets of κ . For every ordinal $\gamma > \kappa$ we have $\|\gamma\|_J < \delta_2(\gamma, \kappa)$, when $\|\gamma\|_J$ is the Galvin–Hajnal rank of the constant function $f : \kappa \rightarrow \gamma + 1$ whose value is γ .*

Instead of proving the above theorem, we prove a more general result. It turns out that the ideal J of nonstationary subsets can be replaced by almost any other ideal satisfying rather weak conditions:

Theorem 2.27 (A better lower bound): *Suppose J is an \aleph_1 -complete ideal on κ such that*

() J as an ideal is generated by $\leq \kappa$ sets or at least we have*

*(**) there exists a model \mathcal{B} (of an expansion of set theory) with universe κ , $|L(\mathcal{B})| \leq \kappa$ and $\psi(P) \in L_{\kappa^+, \omega}$, when $L = L(\mathcal{B}) \cup \{P\}$, P is a unary predicate; having the following property:*

$$\otimes_J \text{ for every } A \subseteq \kappa, \text{ we have that } A \in J \iff \langle \mathcal{B}, A \rangle \models \psi(P)$$

or at least

$$\otimes_J^- \text{ for every } A \subseteq \kappa, \text{ we have that } A \in J \text{ iff for some } A' \text{ we have that } A \subseteq A' \in J, \langle \mathcal{B}, A' \rangle \models \psi(P)$$

then for every ordinal $\gamma > \kappa$ we have that $\|\gamma\|_J < \delta_2(\gamma, \kappa)$.

Remark 2.28 1. One way to see that Theorem 2.26 is a special case of Theorem 2.27 is by using the same argument. Another formal argument (using the statement of 2.27) we can take $\mathcal{B} := \langle \kappa, < \rangle$ and $\psi(P)$ will say that $\{x : P(x)\}$ is a closed unbounded set. This satisfy \otimes_J^- but not \otimes_J .

2. Note that \otimes_J^- is equivalent to: for some $\psi(P, \bar{R}) \in L_{\kappa^+, \omega}$, we have that $A \in J \iff (\exists \bar{R}) \langle \mathcal{B}, A \rangle \models \psi(P, \bar{R})$.

Proof. Let $\gamma^* := \|\gamma\|_J$, and let $ds(\gamma^*)$ stand for the set $\{\nu \mid \nu \text{ is strictly decreasing sequence of ordinals } < \gamma^*\}$. There exists a family of functions $\{f_\eta : \kappa \rightarrow On \mid \eta \in ds(\gamma^*)\}$ with the following properties:

1. f_\emptyset is constantly γ .
2. if $\eta \hat{i} \in ds(\gamma^*)$ then $f_{\eta \hat{i}} <_J f_\eta$, and for every $\zeta < \kappa$ we have $\neg[f_{\eta \hat{i}}(\zeta) < f_\eta(\zeta)] \Rightarrow f_{\eta \hat{i}}(\zeta) = f_\eta(\zeta) = 0$.
3. if $\eta \neq \emptyset$ then $\forall \zeta < \kappa [f_\eta(\zeta) < \gamma]$.
4. $\eta \hat{i} \in ds(\gamma^*) \Rightarrow \|f_{\eta \hat{i}}\|_J \geq i$.
5. $\|f_\emptyset\|_J = \gamma^*$.

This is possible: Define the function f_η by induction on $l(\eta)$:

For $l(\eta) = 0$; Let f_\emptyset be the constant function as in requirement (1).

For $l(\eta) > 0$; If $\eta \hat{i} \in ds(\gamma^*)$ then f_η is defined, and by the inductive hypothesis we have that $\|f_{\eta \hat{i}}\|_J > i$ (as $\eta \hat{i} \in ds(\gamma^*)$ and $\|f_\emptyset\|_J = \gamma^*$), by the definition of the Galvin–Hajnal rank there exists $f' <_J f_\eta$ such that $\|f'\|_J \geq i$. Now for $\zeta < \kappa$ let

$$f_{\eta \hat{i}}(\zeta) := \begin{cases} f'(\zeta) & \text{if } f'(\zeta) < f_\eta(\zeta) \\ 0 & \text{otherwise.} \end{cases}$$

This is enough: Denote by \bar{f} the sequence $\langle f_\eta : \eta \in ds(\gamma^*) \rangle$.

Let χ^* be a sufficiently large regular cardinal such that $H(\chi^*)$ contains all relevant sets and the structure $\langle H(\chi^*), \in \rangle$ reflects all relevant properties. Let $\mathfrak{C} := \langle H(\chi^*), \in, <_{\chi^*}^*, \bar{f}, \kappa, D, \mathcal{B}, \psi(\cdot), J, Q, P, i \rangle_{i \leq \kappa}$, when $<_{\chi^*}^*$ is a well ordering of the set $H(\chi^*)$, P is the unary predicate $\{i : i < \gamma\}$, Q is the unary predicate interpreted by the set $\{j : j < \gamma^*\}$, D interpreted by $ds(\gamma^*)$. Let $T := Th(\mathfrak{C})$, and Γ is a set of types consisting only of the following type – $\{x \in \kappa \wedge x \neq i : i < \kappa\}$. Suppose $N \in EC(T, \Gamma)$ is such that $(P^N, <^{P^N})$ is well ordered, and we will show that $(Q^N, <)$ is well ordered.

W.l.o.g. we may assume that $A^N := \{x \in N : N \models rk(x) \in P\}$ is a transitive set and $\in^N \upharpoonright A^N = \in \upharpoonright A^N$ (by taking the Mostowski's collapse). So $P^N = \gamma'$ for some γ' , and since N omits the type in Γ we have $\kappa^N = \kappa$, since the universe of \mathcal{B} is κ we have $\mathcal{B}^N = \mathcal{B}$. So necessarily $\psi(\cdot)^N = \psi(\cdot)$.

By the axioms of T it follows that

(*) if $N \models \eta \hat{x} \in D$ then $N \models (\exists X \subseteq \{i < \kappa : f_\eta(i) \leq f_{\eta \hat{x}}(i)\}) \mathcal{B} \models \psi(X)$. Now since $\eta \hat{x} \in D^N$, we have $f_\eta^N, f_{\eta \hat{x}}^N \in A^N$. So by the functions from κ into γ' . Also $\psi^N = \psi$, by absoluteness we have $N \models (\exists X \subseteq \{i < \kappa : f_\eta(i) \leq f_{\eta \hat{x}}(i)\}) \mathcal{B} \models \psi(X)$. So by (**) we have

(*) $\eta \hat{x} \in D^N \Rightarrow f_{\eta \hat{x}} <_J f_\eta$.

Now if $(Q^N, <)$ is not well ordered then we can find $\{x_n \in Q^N : N \models [x_{n+1} < x_n]\}$. From T 's axioms it follows that there are $\{y_n : n < \omega\}$ such that $y_0 = \langle \rangle$, $y_{n+1} = y_n \hat{x}_n \in D$ for all $n < \omega$. So we have that $\{f_{y_n} : \kappa \rightarrow \gamma' \mid n < \omega\}$ and for every $n < \omega$ $f_{y_{n+1}} <_J f_{y_n}$ (in V). Since J is an \aleph_1 -complete ideal we have a contradiction. We have shown that there exists a pair T, Γ of the appropriate cardinalities such that

1. $N \in EC(T, \Gamma)$, $(P^N, <)$ is well ordered $\Rightarrow (Q^N, <)$ is well ordered.
2. there is $N \in EC(T, \Gamma)$ with $otp(P^N, <) \leq \gamma$ and $(Q^N, <)$ of order type γ^* (take $N = \mathfrak{C}$).

This establishes that $\gamma^* < \delta_2(\gamma, \kappa)$. □_{2.27}

3 Concluding Remarks

It is natural to ask whether the lower bound from Theorem 2.27 is equal to the one in Fact 2.21. The following seems to be a reasonable

Conjecture 3.1 *For cardinalities κ of uncountable cofinality and λ such that $2^\kappa < \lambda$ we have $\delta_1(\lambda, \kappa) = (\text{cov}(\lambda, \kappa) + 2^\kappa)^+$.*

Remark 3.2 1. Notice that when κ is strong limit singular of cofinality \aleph_0 then the conjecture holds.

2. Why $2^\kappa < \lambda$ – See Barwise-Kunen for independence results.

3. The conjecture can to large extent be translated to a one on *pcf*; it is evident that e.g. (***) below is a sufficient condition:

(***) for any set \mathbf{a} of $\leq \kappa$ regular cardinals which are $> 2^\kappa$ the set *pcf*(\mathbf{a}) has cardinality at most κ , or at least the set *pcf* $_{\aleph_1}$ -complete(\mathbf{a}) has cardinality at most κ .

This is because by [Sh g] II 5.4 if $2^\kappa < \lambda$ then $\mu = \text{cov}(\lambda, \kappa)$ is the first μ such that if $\{\lambda_i : i < \kappa\}$ is a set of regular cardinalities in the interval $(2^\kappa, \lambda)$ and J is an \aleph_1 -complete ideal on κ and *cf*($\prod_{i < \kappa} \lambda_i, < J$) is well defined then it is $\leq \mu$.

The problem is that the ideal may not satisfy even \otimes_J^- . However by [Sh g] VII, 2.6 the ideal J is generated by a family of $\leq |\text{pcf}\{\lambda_i : i < \kappa\}|$ sets and even by a family of just $\leq |\text{pcf}_{\aleph_1}\text{-complete}\{\lambda_i : i < \kappa\}|$ sets, so we have $|\text{pcf}(\{\lambda_i : i < \kappa\})| \leq \kappa \Rightarrow \otimes_J^-$

References

- [BaKu] K. J. Barwise and K. Kunen, Hanf numbers for fragments of $L_{\infty, \omega}$, *Israel J. of Mathematics*, **10**, 306–320, 1971
- [Ch] C. C. Chang, Some remarks on the model theory of infinitary languages, **The Syntax and Semantics of Infinitary Languages**, Springer-Verlag Lecture Notes in Mathematics **72**, edited by J. Barwise (1968) pages 36–63.
- [Gr] Rami Grossberg, **Classification theory for non elementary classes**, Ph.D. Thesis (in Hebrew), The Hebrew University of Jerusalem 1986.
- [GrSh] Rami Grossberg and Saharon Shelah, On the number of nonisomorphic models of an infinitary theory which has the infinitary order property. I, *The Journal of Symbolic Logic*, **51**:302–322, 1986. — **MR**: 87j:03037, (03C45)
- [GrSh 238] Rami Grossberg and Saharon Shelah, A nonstructure theorem for an infinitary theory which has the unsuperstability property, *Illinois Journal of Mathematics*, **30**:364–390, 1986. Volume dedicated to the memory of W.W. Boone; ed. Appel, K., Higman, G., Robinson, D. and Jockush, C. — **MR**: 87j:03036, (03C45)
- [HrSh 334] Ehud Hrushovski and Saharon Shelah, Stability and omitting types, *Israel Journal of Mathematics*, **74**:289–321, 1991. — **MR**: 92m:03049, (03C45)
- [Ma] Menachem Magidor, Chang’s conjecture and powers of singular cardinals, *The Journal of Symbolic Logic*, **42**: 272–276, 1977.
- [MaSh 285] Michael Makkai and Saharon Shelah, Categoricity of theories in $L_{\kappa\omega}$, with κ a compact cardinal, *Annals of Pure and Applied Logic*, **47**:41–97, 1990. — **MR**: 92a:03054, (03C75)
- [Mo] Michael Morley, Omitting classes of elements, **The Theory of Models**, edited by Addison, Henkin and Tarski, North-Holland Publ Co (1965) pages 265–273.
- [Sh c] Saharon Shelah. *Classification theory and the number of nonisomorphic models*, volume 92 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, xxxiv+705 pp, 1990.

- [Sh g] Saharon Shelah. *Cardinal Arithmetic*, volume 29 of *Oxford Logic Guides*. Oxford University Press, 1994.
- [Sh 12] Saharon Shelah. The number of non-isomorphic models of an unstable first-order theory. *Israel Journal of Mathematics*, **9**:473–487, 1971. — **MR**: 43-4652.
- [Sh 16] Saharon Shelah, A combinatorial problem; stability and order for models and theories in infinitary languages, *Pacific Journal of Mathematics*, **41**:247–261, 1972. — **MR**: 46:7018, (02H10)
- [Sh 87a] Saharon Shelah. Classification theory for nonelementary classes, I. The number of uncountable models of $\psi \in L_{\omega_1, \omega}$. Part A. *Israel Journal of Mathematics*, **46**:212–240, 1983.
- [Sh 87b] Saharon Shelah. Classification theory for nonelementary classes, I. The number of uncountable models of $\psi \in L_{\omega_1, \omega}$. Part B. *Israel Journal of Mathematics*, **46**:241–273, 1983.
- [Sh 88] Saharon Shelah. Classification of nonelementary classes. II. Abstract elementary classes. In *Classification theory (Chicago, IL, 1985)*, volume 1292 of *Lecture Notes in Mathematics*, pages 419–497. Springer, Berlin, 1987. Proceedings of the USA–Israel Conference on Classification Theory, Chicago, December 1985; ed. Baldwin, J.T.
- [Sh 300] Saharon Shelah. Universal classes. In *Classification theory (Chicago, IL, 1985)*, volume 1292 of *Lecture Notes in Mathematics*, pages 264–418. Springer, Berlin, 1987. Proceedings of the USA–Israel Conference on Classification Theory, Chicago, December 1985; ed. Baldwin, J.T.
- [Sh 299] Saharon Shelah. Taxonomy of universal and other classes. In *Proceedings of the International Congress of Mathematicians (Berkeley, Calif., 1986)*, volume 1, pages 154–162. Amer. Math. Soc., Providence, RI, 1987. ed. Gleason, A.M.
- [Sh 78] Saharon Shelah. Hanf number of omitting type for simple first-order theories. *The Journal of Symbolic Logic*, **44**:319–324, 1979.
- [Sh 48] Saharon Shelah. Categoricity in \aleph_1 of sentences in $L_{\omega_1, \omega}(Q)$. *Israel Journal of Mathematics*, **20**:127–148, 1975.

- [Sh 200] Saharon Shelah. Classification of first order theories which have a structure theorem. *American Mathematical Society. Bulletin. New Series*, **12**:227–232, 1985.