

PERFECTLY MEAGER SETS AND UNIVERSALLY NULL SETS

TOMEK BARTOSZYNSKI AND SAHARON SHELAH

ABSTRACT. We will show that there is no ZFC example of a set distinguishing between universally null and perfectly meager sets.

1. INTRODUCTION

Consider the following three families of sets of reals:

Definition 1. Let $X \subseteq \mathbb{R}$.

- (1) X is perfectly meager if for every perfect set $P \subseteq \mathbb{R}$, $P \cap X$ is meager in P .
- (2) X is universally meager if every Borel isomorphic image of X is meager
- (3) X is universal null if every Borel isomorphic image of X has Lebesgue measure zero.

Let **PM**, **UM** and **UN** denote these families respectively.

The family **UM** was studied recently by Zakrzewski [13], and identified as an analog of **UN**.

One gets an equivalent definition of **UN** by replacing “Borel isomorphic” by “homeomorphic”, but this is not the case with **UM**.

Let \mathcal{M} and \mathcal{N} denote the σ -ideals of meager and of measure zero subsets of the reals, respectively.

For a σ -ideal $\mathcal{J} \subseteq P(\mathbb{R})$ let

$$\text{non}(\mathcal{J}) = \min\{|X| : X \subseteq \mathbb{R} \text{ \& } X \notin \mathcal{J}\}.$$

There are many ZFC examples of uncountable sets that are in $\mathbf{UM} \cap \mathbf{UN}$. These include $\omega_1 \omega_1^*$ -gaps, a selector from the constituents of a non-Borel $\mathbf{\Pi}_1^1$ set, etc. (see [9]) All these sets have size \aleph_1 , since Miller [8] showed that, consistently, no set of size 2^{\aleph_0} is in $\mathbf{UM} \cup \mathbf{UN}$.

Grzegorek found other constructions in ZFC that produce sets of (consistently) different sizes.

Theorem 2 (Grzegorek, [6]).

- (1) There exists a set $X \in \mathbf{UN}$ such that $|X| = \text{non}(\mathcal{N})$,
- (2) There exists a set $X \in \mathbf{UM}$ such that $|X| = \text{non}(\mathcal{M})$.

The problem whether the equality $\mathbf{UM} = \mathbf{UN}$ is consistent is open. However, both inclusions are consistent with ZFC; $\mathbf{UM} \subsetneq \mathbf{UN}$ holds in a model obtained by

1991 *Mathematics Subject Classification.* 03E17.

Key words and phrases. perfectly meager, universally null, consistency.

The first author was partially supported by NSF grant DMS 9971282 and Alexander von Humboldt Foundation.

The second author was partially supported by Israel Science Foundation. Publication 732.

adding \aleph_2 Cohen reals, and $\mathbf{UN} \subsetneq \mathbf{UM}$ holds in a model obtained by adding \aleph_2 random reals (side-by-side) (see [4], [9], [8]).

In this paper we investigate the connection between families \mathbf{UN} and \mathbf{PM} , and show that both inclusions $\mathbf{PM} \subseteq \mathbf{UN}$ and $\mathbf{UN} \subseteq \mathbf{PM}$ are consistent with ZFC as well. Observe that trivially $\mathbf{UM} \subseteq \mathbf{PM}$, thus we only need to check that $\mathbf{PM} \subseteq \mathbf{UN}$ is consistent. Recall that $\mathbf{PM} \neq \mathbf{UM}$ is consistent ([12]) as well as $\mathbf{PM} = \mathbf{UM}$ ([2]). We will show that:

Theorem 3. *It is consistent with ZFC that $\mathbf{PM} \subseteq [\mathbb{R}]^{\leq \aleph_1} \subseteq \mathbf{UN}$.*

2. FORCING

Suppose that $X \subseteq 2^\omega$ is a perfectly meager set in \mathbf{V} . Let \tilde{P} be a fixed closed subset of $2^\omega \times 2^\omega$ which is universal for perfect sets in 2^ω . In other words, for every perfect set $P \subseteq 2^\omega$ there exists an x such that $P = (\tilde{P})_x = \{y : (x, y) \in \tilde{P}\}$. Note that this property is absolute. Since X is perfectly meager, we can find sets $\tilde{Q}^n \subseteq 2^\omega \times 2^\omega$ such that for every $x \in 2^\omega$ and $n \in \omega$,

- (1) $(\tilde{Q}^n)_x$ is a closed nowhere dense subset of $(\tilde{P})_x$,
- (2) $X \cap (\tilde{P})_x \subseteq \left(\bigcup_{n \in \omega} \tilde{Q}^n\right)_x$.

Clearly, the set $\bigcup_{n \in \omega} \tilde{Q}^n$ witnesses that $X \in \mathbf{PM}$ since

$$X \subseteq 2^\omega \setminus \bigcup_{x \in 2^\omega} (\tilde{P} \setminus \bigcup_{n \in \omega} \tilde{Q}^n)_x.$$

Note that the last inclusion makes sense even if X is not a subset of \mathbf{V} . Suppose that $\mathbf{V}' \subseteq \mathbf{V}$ and $X \subseteq \mathbf{V}$ is a set of reals. We will say $\mathbf{V}' \models X \in \mathbf{PM}$ if there exists a family $\{\tilde{Q}^n : n \in \omega\} \in \mathbf{V}'$ such that $X \cap (\tilde{P})_x \subseteq \left(\bigcup_{n \in \omega} \tilde{Q}^n\right)_x$ for every real $x \in \mathbf{V}'$.

The property of being perfectly meager is not absolute so whether X is perfectly meager in \mathbf{V}' has no bearing onto whether X is perfectly meager in \mathbf{V} . For example, if $x \in \mathbf{V}$ is a Cohen real over \mathbf{V}' then the set $\{x\}$ is perfectly meager in \mathbf{V} but not in \mathbf{V}' .

Lemma 4. *Let $\langle \mathcal{P}_\alpha, \dot{Q}_\alpha : \alpha < \omega_2 \rangle$ be a countable support iteration of proper forcing notions over $\mathbf{V} \models \mathbf{CH}$. Suppose that $X \subseteq \mathbf{V}^{\mathcal{P}_{\omega_2}} \cap \mathbb{R}$ is a perfectly meager set. Then there exists an ω_1 -club $C \subseteq \omega_2$ such that for every $\alpha \in C$,*

$$\mathbf{V}^{\mathcal{P}_\alpha} \models X \in \mathbf{PM}.$$

Proof. Let $\{\tilde{Q}^n : n \in \omega\} \in \mathbf{V}^{\mathcal{P}_{\omega_2}}$ be a family witnessing that X is perfectly meager. Let C consist of those ordinals of cofinality ω_1 that for every n , $\tilde{Q}^n \cap ((2^\omega \cap \mathbf{V}^{\mathcal{P}_\alpha}) \times 2^\omega) \in \mathbf{V}^{\mathcal{P}_\alpha}$. The usual argument involving Skolem-Löwenheim theorem shows that C has the required property. \square

Our objective is to find a set of general conditions on a forcing notion \mathbb{P} such that the countable support iteration of \mathbb{P} of length ω_2 produces a model where $\mathbf{PM} \subseteq [\mathbb{R}]^{\leq \aleph_1} \subseteq \mathbf{UN}$. These conditions are sufficient for the class of forcing notions defined using norms [10].

These conditions are the following:

- (1) $\mathbf{V}^{\mathbb{P}} \models \mathbf{V} \cap 2^\omega \in \mathcal{N}$,
- (2) $\mathbf{V}^{\mathbb{P}} \models \mathbf{V} \cap 2^\omega \notin \mathcal{M}$,

- (3) \mathbb{P} is ω^ω -bounding, that is $\omega^\omega \cap \mathbf{V}$ is a dominating family in $\omega^\omega \cap \mathbf{V}^{\mathbb{P}}$,
- (4) \mathbb{P} adds a real $x_{\mathbb{P}} \in 2^\omega$ such that $\mathbf{V} \models \{x_{\mathbb{P}}\} \notin \mathbf{PM}$.
- (5) \mathbb{P} generic real is minimal, that is, if g is \mathbb{P} -generic over \mathbf{V} and $x \in \mathbf{V}[g] \cap 2^\omega$ then $x \in \mathbf{V}$ or $g \in \mathbf{V}[x]$.

Condition (1) is necessary to make all sets of size \aleph_1 universally null, and condition (2) is necessary to avoid making all \aleph_1 sets perfectly meager. Recall that (2) and (3) together are essentially equivalent to $\mathbf{V}^{\mathbb{P}} \models \mathbf{V} \cap \mathcal{M}$ is cofinal in \mathcal{M} .

For the forcing notions \mathbb{P} that we have in mind the following property holds: for every real $x \in \mathbf{V}^{\mathbb{P}}$ there exists a continuous function $f \in \mathbf{V}$ such that $x = f(x_G)$, where x_G is a generic real.

Condition (5), guarantees that in the above context f can be chosen to be a homeomorphism. In particular, if X is a set of reals of size \aleph_2 then X will contain a homeomorphic image of a sequence of generic reals.

The following forcing notion appeared in [5], it is similar (but not identical) to the infinitely equal real forcing from [7].

For a tree p and $t \in p$, let $\text{succ}_p(t)$ be the set of all immediate successors of t in p , $p_t = \{v \in p : t \subseteq v \text{ or } v \subseteq t\}$ the subtree of p determined by t , $p|n$ the n -th level of p , and let $[p]$ be the set of branches of p . By identifying $s \in \omega^{<\omega}$ with the full-branching tree having root s , we can also denote $[s] = \{f \in \omega^\omega : s \subseteq f\}$.

Fix a strictly increasing function $f \in \omega^\omega$ and let $\mathbf{X} = \prod_{n \in \omega} f(n)$. Note that \mathbf{X} is a Polish space homeomorphic to 2^ω . For technical reasons we require that $f(n) = 2^{\dot{f}(n)}$ for $n \in \omega$.

Let $\mathbb{E}\mathbb{E}$ be the following forcing notion: $p \in \mathbb{E}\mathbb{E}$ if

- (1) p is a nonempty subtree of $\omega^{<\omega}$,
- (2) $s(n) < f(n)$ for all $s \in p$ and $n \in \text{dom}(s)$,
- (3) for all $s \in p$ there exists an extension t of s such that $t \hat{\ } n \in p$ for all $n < f(|t|)$.

For $p, q \in \mathbb{E}\mathbb{E}$, $p \geq q$ if $p \subseteq q$. Without loss of generality we can assume that $|\text{succ}_p(s)| = 1$ or $\text{succ}_p(s) = f(|p|)$ for all $p \in \mathbb{E}\mathbb{E}$ and $s \in p$. Conditions of this type form a dense subset of $\mathbb{E}\mathbb{E}$.

Let

$$\text{split}(p) = \{s \in p : |\text{succ}_p(s)| > 1\} = \bigcup_{n \in \omega} \text{split}_n(p),$$

where $\text{split}_n(p) = \left\{ s \in \text{split}(p) : \left| \{t \subsetneq s : t \in \text{split}(p)\} \right| = n \right\}$.

For $p, q \in \mathbb{E}\mathbb{E}$, $n \in \omega$, we let

$$p \geq_n q \iff p \geq q \ \& \ \text{split}_n(q) = \text{split}_n(p).$$

Lemma 5 ([5]). (1) $\mathbb{E}\mathbb{E}$ satisfies Axiom A, so it is proper,

- (2) $\mathbf{V}^{\mathbb{E}\mathbb{E}} \models \mathbf{V} \cap 2^\omega \in \mathcal{N}$,
- (3) $\mathbf{V}^{\mathbb{E}\mathbb{E}} \models \mathbf{V} \cap 2^\omega \notin \mathcal{M}$,
- (4) for every maximal antichain $\mathcal{A} \subseteq \mathbb{E}\mathbb{E}$, $p \in \mathbb{E}\mathbb{E}$, and $n \in \omega$ there exists $q \geq_n p$ such that $\{r \in \mathcal{A} : r \text{ is compatible with } q\}$ is finite.
- (5) for every family of maximal antichains $\{\mathcal{A}_n : n \in \omega\}$ and $p \in \mathbb{E}\mathbb{E}$ there exists $q \geq p$ such that for every n , $\{r \in \mathcal{A}_n : r \text{ is compatible with } q\}$ is finite.
- (6) $\mathbb{E}\mathbb{E}$ is ω^ω bounding,
- (7) $\mathbf{V}^{\mathbb{E}\mathbb{E}} \models \mathbf{V} \cap \mathcal{M}$ is cofinal in \mathcal{M} . \square

Note that for $p \in \mathbb{E}\mathbb{E}$ the set $[p]$ is a compact subset of $\mathbf{X} = \prod_n f(n)$. Moreover, there is a canonical isomorphism between $[p]$ and 2^ω defined as follows:

For every n let $\{s_0^n, \dots, s_{f(n)}^n\}$ be a fixed enumeration of 0-1 sequences of length $\tilde{f}(n)$ (recall that $f(n) = 2^{\tilde{f}(n)}$). Define $F : [p] \rightarrow 2^\omega$ as

$$F(x) = s_{x(n_0+1)}^{n_0} \widehat{\ } s_{x(n_1+1)}^{n_1} \widehat{\ } \dots,$$

where n_0, n_1, \dots is the increasing enumeration of the set $\{n : x \upharpoonright n \in \text{split}(p)\}$.

Lemma 6. *Let $p \in \mathbb{E}\mathbb{E}$ and suppose that $H \subseteq [p]$ is a meager set in $[p]$. For every $n \in \omega$ there exists $q \geq_n p$ such that $[q] \cap H = \emptyset$. In particular, $\Vdash_{\mathbb{E}\mathbb{E}} \mathbf{V} \models \{\dot{g}\} \notin \mathbf{PM}$.*

Proof. Let $H \subseteq [p]$ be a meager set, and let $n \in \omega$. Fix a descending sequence of open sets $\langle U_k : k \in \omega \rangle$ such that each U_k is dense in $[p]$ and $H \cap \bigcap_k U_k = \emptyset$. By induction build a sequence $\langle p_k : k \in \omega \rangle$ such that $p_0 = p$, and for every k ,

- (1) $p_{k+1} \geq_{n+k+1} p_k \in \mathbb{E}\mathbb{E}$,
- (2) $[p_{k+1}] \subseteq U_k$.

Suppose that p_k is given. For every $v \in \text{split}_{n+k+1}(p_k)$ find $q_v \geq (p_k)_v$ such that $[q_v] \subseteq U_k$. Let $p_{k+1} = \bigcup \{q_v : v \in \text{split}_{n+k+1}(p_k)\}$. Condition $q = \lim_k p_k$ has the required property.

Suppose that $\{\widetilde{Q}^n : n \in \omega\} \in \mathbf{V}$ is a possible witness that $\{\dot{g}\}$ is perfectly meager, and let $p \in \mathbb{E}\mathbb{E}$. Find $x \in \mathbf{V}$ such that $[p] = (P)_x$ and let $q \geq p$ be such that $[q] \cap \left(\bigcup_n \widetilde{Q}^n\right)_x = \emptyset$. Clearly,

$$q \Vdash_{\mathbb{E}\mathbb{E}} \{\dot{g}\} \in \bigcup_{x \in \mathbf{V}} \left(P \setminus \bigcup_n \widetilde{Q}^n \right)_x.$$

In particular, $q \Vdash_{\mathbb{E}\mathbb{E}} \mathbf{V} \models \{\dot{g}\} \notin \mathbf{PM}$. □

Lemma 7. *Suppose that $p \in \mathbb{E}\mathbb{E}$ and $p \Vdash_{\mathbb{E}\mathbb{E}} \dot{x} \in 2^\omega$. For every $n \in \omega$ there exists $q \geq_n p$ and a continuous function $F : [q] \rightarrow 2^\omega$ such that $q \Vdash_{\mathbb{E}\mathbb{E}} \dot{x} = F(\dot{g})$, where \dot{g} is the canonical name for the generic real.*

Moreover, we can require that for every $v \in \text{split}_n(q)$ and any $x_1, x_2 \in [q_v]$, $F(x_1) \upharpoonright n = F(x_2) \upharpoonright n$.

Proof. The first part is a special case of a more general fact. For $n \in \omega$ let $\mathcal{A}_n \subseteq \mathbb{E}\mathbb{E}$ be a maximal antichain below p such that $\forall r \in \mathcal{A}_n \exists s \in 2^n r \Vdash_{\mathbb{E}\mathbb{E}} \dot{x} \upharpoonright n = s$. Use 5(5) to find $q \geq p$ such that for every $n \in \omega$, $\{r \in \mathcal{A}_n : r \text{ is compatible with } q\}$ is finite. Let $\mathcal{A}'_n = \{r \in \mathcal{A}_n : r \text{ is compatible with } q\}$. Without loss of generality we can assume that $[q] \subseteq \bigcup_{r \in \mathcal{A}'_n} [r]$. It follows that $[r] \cap [q]$ is clopen in $[q]$ for every $r \in \mathcal{A}'_n$. Define $F : [q] \rightarrow 2^\omega$ as $F(x) = y$ if for every $n \in \omega$ there exists $r \in \mathcal{A}'_n$ such that $x \in [r]$ and $r \Vdash_{\mathbb{E}\mathbb{E}} \dot{x} \upharpoonright n = y \upharpoonright n$. It is easy to see that F is a continuous function that has the required properties.

To show the second part we need to build q in such a way that for every $v \in \text{split}_n(q)$, there is $r \in \mathcal{A}'_n$ such that $q_v \geq r$. □

Lemma 8. *Suppose that $p \in \mathbb{E}\mathbb{E}$, $n \in \omega$ and $p \Vdash_{\mathbb{E}\mathbb{E}} \dot{x} \in 2^\omega$. Let $F : [q] \rightarrow 2^\omega$ be a continuous function such that $p \Vdash_{\mathbb{E}\mathbb{E}} \dot{x} = F(\dot{g})$.*

There exists $q \geq p$ such that $F \upharpoonright [q]$ is constant, or there exists $q \geq_n p$ such that $F \upharpoonright [q]$ is one-to-one. In particular, the generic real is minimal.

Proof. Consider the following two cases.

CASE 1 $p \Vdash_{\mathbb{E}\mathbb{E}} \dot{x} \notin \mathbf{V}$. Let $x \in \mathbf{V}$ and $q \geq p$ be such that $q \Vdash_{\mathbb{E}\mathbb{E}} \dot{x} = x$. Clearly $F \upharpoonright [q]$ is constant with value x .

CASE 2 $p \Vdash_{\mathbb{E}\mathbb{E}} \dot{x} \notin \mathbf{V}$.

Build by induction a sequence of conditions $\langle p_k : k \in \omega \rangle$ such that $p_0 = p$ and for every k ,

- (1) $p_{k+1} \geq_{n+k+1} p_k$,
- (2) sets $\left\{ F^{\omega} \left([(p_{k+1})_s] \right) : s \in \text{split}_{n+k+1}(p_{k+1}) \right\}$ are pairwise disjoint and have diameter $< 2^{-k}$

Suppose that p_k is given. Note that $F^{\omega} \left([(p_k)_s] \right)$ is uncountable for every $s \in p_k$. For $v \in \text{split}_{n+k+1}(p_k)$ choose pairwise different reals $x_v \in F^{\omega} \left([(p_k)_v] \right)$. It is not important now but will be relevant in the sequel, that we can choose these reals “effectively” from a fixed countable subset of $[p_k]$. Let $\ell > k$ be such that sequences $x_v \upharpoonright \ell$ are also pairwise different. For every $v \in \text{split}_{n+k+1}(p_k)$ let $s_v \in \text{split}(p_k)$ be such that for every $z \in [(p_k)_{s_v}]$, $F(z) \upharpoonright \ell = x_v \upharpoonright \ell$. If F is as in the second part of lemma 7 then we can find s_v in $\text{split}_{\ell}(p_k)$. Define $p_{k+1} = \bigcup \{ (p_k)_{s_v} : v \in \text{split}_{n+k+1}(p_k) \}$. Observe that $q = \lim_k p_k$ has the required property. \square

Note that the above lemma shows that the reals added by $\mathbb{E}\mathbb{E}$ are minimal. Infinitely equal forcing from [7] or [4] does not have this property.

3. ITERATION OF $\mathbb{E}\mathbb{E}$.

Let $\alpha \leq \omega_2$ be an ordinal and suppose that $\mathbb{E}\mathbb{E}_{\alpha}$ is a countable support iteration of $\mathbb{E}\mathbb{E}$ of length α . In other words, $p \in \mathbb{E}\mathbb{E}_{\alpha}$ is

- (1) p is a function and $\text{dom}(p) = \alpha$,
- (2) $\text{supp}(p) = \{ \beta : p(\beta) \neq \emptyset \}$ is countable,
- (3) $\forall \beta < \alpha$ $p \upharpoonright \beta \Vdash_{\mathbb{E}\mathbb{E}_{\beta}} p(\beta) \in \mathbb{E}\mathbb{E}$.

For $F \in [\alpha]^{<\omega}$, $n \in \omega$, and $p, q \in \mathbb{E}\mathbb{E}_{\alpha}$ define

$$q \geq_{F,n} p \iff q \geq p \ \& \ \forall \beta \in F \ q \upharpoonright \beta \Vdash_{\mathbb{E}\mathbb{E}_{\beta}} q(\beta) \geq_n p(\beta).$$

The following fact is well-known.

Theorem 9 ([5], [7], [3]). *Suppose that $p \in \mathbb{E}\mathbb{E}_{\alpha}$, $F \in [\alpha]^{<\omega}$, and $n \in \omega$.*

- (1) *for every maximal antichain $\mathcal{A} \subseteq \mathbb{E}\mathbb{E}_{\alpha}$, there exists $q \geq_{F,n} p$ such that $\{ r \in \mathcal{A} : r \text{ is compatible with } q \}$ is finite.*
- (2) *for every family of maximal antichains $\{ \mathcal{A}_n : n \in \omega \}$ there exists $q \geq p$ such that for every n , $\{ r \in \mathcal{A}_n : r \text{ is compatible with } q \}$ is finite.*
- (3) $\mathbf{V}^{\mathbb{E}\mathbb{E}_{\omega_2}} \models [\mathbb{R}]^{<2^{\aleph_0}} \subseteq \mathcal{N}$.
- (4) $\mathbf{V}^{\mathbb{E}\mathbb{E}_{\omega_2}} \models \mathcal{M} \cap \mathbf{V}$ is cofinal in \mathcal{M} . \square

For $p \in \mathbb{E}\mathbb{E}_{\alpha}$ let $\text{cl}(p)$ be the smallest set $w \subseteq \alpha$ such that p can be evaluated using generic reals $\langle \dot{g}_{\beta} : \beta \in w \rangle$. In other words, $\text{cl}(p)$ consists of those $\beta < \alpha$ such that the transitive closure of p contains $\mathbb{E}\mathbb{E}_{\beta}$ name for an element of $\mathbb{E}\mathbb{E}$. It is well-known [11] that $\{ p \in \mathbb{E}\mathbb{E}_{\alpha} : \text{cl}(p) \in [\alpha]^{<\omega} \}$ is dense in $\mathbb{E}\mathbb{E}_{\alpha}$.

Suppose that $p \in \mathbb{E}\mathbb{E}_{\alpha}$, $w = \text{cl}(p)$ is countable and $\alpha_p = \text{ot}(\text{cl}(p))$. Let $\mathbb{E}\mathbb{E}_w$ be the countable support iteration of $\mathbb{E}\mathbb{E}$ with the domain w . In other words, consider

The countable support iteration $\langle \mathcal{P}_\beta, \dot{Q}_\beta : \beta < \text{sup}(w) \rangle$ such that

$$\forall \beta < \text{sup}(w) \Vdash_{\mathcal{P}_\beta} \dot{Q}_\beta \simeq \begin{cases} \mathbb{E}\mathbb{E} & \text{if } \beta \in w \\ \emptyset & \text{if } \beta \notin w \end{cases}.$$

It is clear that $\mathbb{E}\mathbb{E}_w \simeq \mathbb{E}\mathbb{E}_{\alpha_p}$. Moreover, we can view condition p as a member of $\mathbb{E}\mathbb{E}_w$.

For the rest of the section we will consider only the iteration of $\mathbb{E}\mathbb{E}$ of countable length α and show that $\mathbb{E}\mathbb{E}_\alpha$ has the same properties that $\mathbb{E}\mathbb{E}$ has.

Let α be a countable ordinal and $p \in \mathbb{E}\mathbb{E}_\alpha$. Define $\bar{p} \subseteq \mathbf{X}^\alpha$ as follows:

$\langle x_\beta : \beta < \alpha \rangle \in \bar{p}$ if for every $\beta < \alpha$,

$$x_\beta \in \left[p(\beta) \left[\langle x_\gamma : \gamma < \beta \rangle \right] \right].$$

Note that $p(\beta) \left[\langle x_\gamma : \gamma < \beta \rangle \right]$ is the interpretation of $p(\beta)$ using reals $\langle x_\gamma : \gamma < \beta \rangle$ so may be undefined if these reals are not sufficiently generic.

For a set $G \subseteq \mathbf{X}^\alpha$, $u \subseteq \alpha$, and $x \in \mathbf{X}^u$ let

$$(G)_x = \{y \in \mathbf{X}^{\alpha \setminus u} : \exists z \in G \ z \upharpoonright u = x \ \& \ z \upharpoonright (\alpha \setminus u) = y\},$$

and for $\beta \in \alpha$ let $(G)_\beta = \{x(\beta) : x \in G\}$.

We say that $p \in \mathbb{E}\mathbb{E}_\alpha$ is good if

- (1) \bar{p} is compact,
- (2) for every $\beta < \alpha$ and $x \in \overline{p \upharpoonright \beta}$, $\overline{p[x]} = (p)_x$ and $\overline{p(\beta)[x]} = ((p)_x)_\beta$.
- (3) \bar{p} is homeomorphic to \mathbf{X}^α via a homeomorphism h such that for every $\beta < \alpha$ and $x \in \overline{p \upharpoonright \beta}$, $h \upharpoonright ((p)_x)_\beta$ is a homeomorphism between $((p)_x)_\beta$ and \mathbf{X} .

Lemma 10. $\{p \in \mathbb{E}\mathbb{E}_\alpha : \bar{p} \text{ is good}\}$ is dense in $\mathbb{E}\mathbb{E}_\alpha$.

Proof. CASE 1. $\alpha = \beta + 1$.

Fix $p \in \mathbb{E}\mathbb{E}_\alpha$ and for $n \in \omega$ let \mathcal{A}_n be a maximal antichain below $p \upharpoonright \beta$ such that

- (1) $\forall r \in \mathcal{A}_n \ \bar{r}$ is compact.
- (2) $\forall r \in \mathcal{A}_n \ \exists t \subseteq \prod_{j < n} f(j) \ r \Vdash_{\mathbb{E}\mathbb{E}_\beta} p(\beta) \upharpoonright n = t$.

Fix a sequence $\langle F_n : n \in \omega \rangle$ such that for $n \in \omega$,

- (1) $F_n \in [\beta]^{<\omega}$,
- (2) $F_n \subseteq F_{n+1}$,
- (3) $\bigcup_n F_n = \beta$.

By induction build a sequence $\langle q_n : n \in \omega \rangle$ such that for $n \in \omega$,

- (1) $\overline{q_n}$ is compact,
- (2) $q_{n+1} \geq_{F_n, n} q_n$,
- (3) $\exists \mathcal{A}'_n \in [\mathcal{A}_n]^{<\omega} \ \overline{q_n} \subseteq \bigcup_{r \in \mathcal{A}'_n} \bar{r}$.

Let $q_\omega = \lim_n q_n$. As in the proof of 7 we show that there exists a continuous function $F : \overline{q_\omega} \rightarrow \mathbb{E}\mathbb{E}$ (encode elements of $\mathbb{E}\mathbb{E}$ as reals) such that

$$q_\omega \Vdash_{\mathbb{E}\mathbb{E}_\beta} p(\beta) = F(\langle \dot{g}_\gamma : \gamma < \beta \rangle).$$

Consider $q = q_\omega \widehat{\ } p(\beta) \geq p$. Clearly, $\bar{q} = \{\langle x, y \rangle : x \in \overline{q_\omega}, y \in [F(x)]\}$ is compact in \mathbf{X}^α . Remaining requirements are met as well.

CASE 2. α is limit.

Given $p \in \mathbb{E}\mathbb{E}_\alpha$ fix sequences $\langle F_n : n \in \omega \rangle$ and $\langle \alpha_n : n \in \omega \rangle$ such that

- (1) $F_n \in [\alpha_n]^{<\omega}$,
- (2) $F_n \subseteq F_{n+1}$,

- (3) $\bigcup_n F_n = \alpha$,
- (4) $\sup_n \alpha_n = \alpha$.

By induction build a sequence $\langle q_n : n \in \omega \rangle$ such that for $n \in \omega$,

- (1) $q_n \in \mathbb{E}\mathbb{E}_\alpha$,
- (2) $\text{supp}(q_n) \subseteq \alpha_n$,
- (3) $q_{n+1} \geq_{F_n, n} q_n$,
- (4) $q_n \upharpoonright \alpha_n \geq p \upharpoonright \alpha_n$,
- (5) $q_n \upharpoonright \alpha_n$ is compact in \mathbf{X}^{α_n} .

Let $q = \lim_n q_n$. Note that $\bar{q} = \bigcap_n \overline{q_n \upharpoonright \alpha_n} \times \mathbf{X}^{\alpha \setminus \alpha_n}$ is as required. \square

From now on we will always work with conditions p such that \bar{p} is good. We noticed earlier that for every condition $p \in \mathbb{E}\mathbb{E}$, $[p]$ is canonically isomorphic to 2^ω , in exactly the same way we can verify that if $p \in \mathbb{E}\mathbb{E}_\alpha$ and \bar{p} is good then \bar{p} is isomorphic to $(2^\omega)^\alpha$.

As in the lemma 7 we show that:

Lemma 11. *Suppose that $p \in \mathbb{E}\mathbb{E}_\alpha$ and $p \Vdash_{\mathbb{E}\mathbb{E}_\alpha} \dot{x} \in 2^\omega$. Then there exists $q \geq p$ and a continuous function $F : \bar{p} \rightarrow 2^\omega$ such that $q \Vdash_{\mathbb{E}\mathbb{E}_\alpha} \dot{x} = F(\dot{\mathbf{g}})$, where $\dot{\mathbf{g}} = \langle \dot{g}_\beta : \beta < \alpha \rangle$ is the sequence of generic reals.*

Lemma 12. *Let $p \in \mathbb{E}\mathbb{E}_\alpha$ and suppose that $H \subseteq \bar{p}$ is a meager set in \bar{p} . For every $F \in [\alpha]^{<\omega}$ and $n \in \omega$ there exists $q \geq_{F, n} p$ such that $\bar{q} \cap H = \emptyset$.*

Proof. As before, without loss of generality we can assume that α is countable.

Induction on α .

CASE 1. $\alpha = \beta + 1$.

Suppose that $p \in \mathbb{E}\mathbb{E}_\alpha$ and $H \subseteq \bar{p} \subseteq \mathbf{X}^\beta \times \mathbf{X}$ is meager, and let $F \in [\alpha]^{<\omega}$ and $n \in \omega$ be given.

Let

$$H' = \{x \in \bar{p} \upharpoonright \beta : (H)_x \text{ is not meager in } [p(\beta)[x]] = ((\bar{p})_x)_\beta\}.$$

Using the fact that \bar{p} is homeomorphic to $(2^\omega)^\alpha$ via homeomorphism respecting vertical sections, and by Kuratowski-Ulam theorem, we conclude that H' is a meager set in $p \upharpoonright \beta$.

Recall the following classical lemma:

Lemma 13 ([1]). *Suppose that $H \subseteq 2^\omega \times 2^\omega$ is a Borel set.*

- (1) *Assume $(H)_x$ is meager for all x . Then there exists a sequence of Borel sets $\{G_n : n \in \omega\} \subseteq 2^\omega \times 2^\omega$ such that*
 - (a) $(G_n)_x$ is a closed nowhere dense set for all $x \in 2^\omega$,
 - (b) $H \subseteq \bigcup_{n \in \omega} G_n$.

By the inductive hypothesis we can find $q^* \geq_{F \cap \beta, n} p \upharpoonright \beta$ such that $\bar{q}^* \cap H' = \emptyset$. By lemma 6 for every $x \in \bar{q}^*$ there exists $q_x \geq_n p(\beta)[x]$ such that $[q_x] \cap (H)_x = \emptyset$. Moreover, by 13, the mapping $x \mapsto q_x$ is can be chosen to be Borel, and subsequently, by shrinking q^* , continuous. Let $q \in \mathbb{E}\mathbb{E}_\alpha$ be defined such that $q \upharpoonright \beta = q^*$ and $q^* \Vdash_{\mathbb{E}\mathbb{E}_\beta} q(\beta) = q_{\dot{g}_\beta}$. It is clear that q has the required properties.

CASE 2. α is limit.

Fix sequences $\langle F_n : n \in \omega \rangle$ and $\langle \alpha_n : n \in \omega \rangle$ such that

- (1) $F_n \in [\alpha_n]^{<\omega}$,
- (2) $F_n \subseteq F_{n+1}$,

- (3) $\bigcup_n F_n = \alpha$,
- (4) $\sup_n \alpha_n = \alpha$.

By induction build a sequence $\langle q_n : n \in \omega \rangle$ such that for $n \in \omega$,

- (1) $q_n \in \mathbb{E}\mathbb{E}_\alpha$,
- (2) $\text{supp}(q_n) \subseteq \alpha_n$,
- (3) $q_{n+1} \geq_{F_n, n} q_n$,
- (4) $q_n \upharpoonright \alpha_n \geq p \upharpoonright \alpha_n$,
- (5) $\overline{q_n \upharpoonright \alpha_n} \cap H_n = \emptyset$, where $H_n = \left\{ x \in \overline{q_n \upharpoonright \alpha_n} : (H)_x \text{ is not meager in } \overline{p \upharpoonright x} \right\}$.

As before (5) is possible by Kuratowski-Ulam theorem. Let $q = \lim_n q_n$. It is clear that $\overline{q} \cap H = \emptyset$. \square

The following lemma is an analog of lemma 8.

Lemma 14. *Suppose that $p \in \mathbb{E}\mathbb{E}_\alpha$, $n \in \omega$ and $p \Vdash_{\mathbb{E}\mathbb{E}_\alpha} \dot{x} \in 2^\omega$. Let $F : \overline{p} \rightarrow 2^\omega$ be a continuous function such that $p \Vdash_{\mathbb{E}\mathbb{E}_\alpha} \dot{x} = F(\dot{\mathbf{g}})$, where $\dot{\mathbf{g}} = \langle \dot{g}_\beta : \beta < \alpha \rangle$ is the sequence of generic reals. There exists $q \geq p$ such that exactly one of the following conditions hold:*

- (1) $F \upharpoonright \overline{q}$ is constant,
- (2) there exists $\beta < \alpha$ such that $F \upharpoonright \overline{q \upharpoonright \beta}$ is one-to-one and for every $x \in \overline{q \upharpoonright \beta}$, $F \upharpoonright (q \upharpoonright \beta)_x$ is constant,
- (3) $F \upharpoonright \overline{q}$ is one-to-one.

Proof. We have three cases:

CASE 1. There exists $q \geq p$ such that $q \Vdash_{\mathbb{E}\mathbb{E}_\alpha} \dot{x} \in \mathbf{V}$. Without loss of generality we can assume that for some $x \in \mathbf{V} \cap 2^\omega$ $q \Vdash_{\mathbb{E}\mathbb{E}_\alpha} \dot{x} = x$. It follows that $F \upharpoonright \overline{q}$ is constant.

CASE 2. There exists $q \geq p$ such that $q \Vdash_{\mathbb{E}\mathbb{E}_\alpha} \exists \beta < \alpha \dot{x} \in \mathbf{V}^{\mathbb{E}\mathbb{E}_\beta}$. By shrinking q we can assume that there exists a continuous function $G : \overline{q \upharpoonright \beta} \rightarrow 2^\omega$ such that $q \Vdash_{\mathbb{E}\mathbb{E}_\alpha} \dot{x} = G(\dot{\mathbf{g}} \upharpoonright \beta)$. In particular, for $x \in [q]$, $F(x) = G(x \upharpoonright \beta)$. If β was minimal then, using the argument below, we can also assume that G is one-to-one.

Suppose that $q \in \mathbb{E}\mathbb{E}_\alpha$, $F \in [\alpha]^{<\omega}$, and $n \in \omega$. Without loss of generality we can assume that for every $\beta \in F$, $q \upharpoonright \beta$ determines the value of $\text{split}_n(q(\beta))$ (up to finitely many values). Suppose that $\sigma : F \rightarrow \omega^{<\omega}$ is a function such that $\sigma(\beta) \in \text{split}_n(q(\beta))$ for $\beta \in F$. Let $(q)_\sigma$ be the condition defined as

$$\forall \beta < \alpha (q)_\sigma \upharpoonright \beta \Vdash_{\mathbb{E}\mathbb{E}_\beta} (q)_\sigma(\beta) = \begin{cases} q(\beta) & \text{if } \beta \notin F \\ (q(\beta))_{\sigma(\beta)} & \text{if } \beta \in F \end{cases} .$$

Let $\Sigma_{F,n}$ be the finite set of all mappings σ satisfying the requirements.

Lemma 15. *Suppose that $F \in [\alpha]^{<\omega}$, $n \in \omega$ and*

$$p \Vdash_{\mathbb{E}\mathbb{E}_\alpha} \dot{x} = F(\dot{\mathbf{g}}) \ \& \ \forall \beta < \alpha \dot{x} \notin \mathbf{V}^{\mathbb{E}\mathbb{E}_\beta} .$$

There exists $q \geq_{F,n} p$ such that the sets $\left\{ F''(\overline{(q)_\sigma}) : \sigma \in \Sigma_{F,n} \right\}$ are pairwise disjoint.

Proof. Induction on $|F|$ and α . If $F = \{\beta\}$ this is essentially lemma 8.

Let $\{v_j : j < k^*\}$ be an enumeration of $\text{split}_n(p(\beta))$. For $v \in \text{split}_n(p)$ choose pairwise different reals $x_v \in F''(\overline{(p)_v})$. Note that this choice can be made canonically from, for example, the countable dense set of leftmost branches of subtrees

of p . Let $\ell > k$ be such that sequences $x_v \upharpoonright \ell$ are also pairwise different. Define conditions $\langle r_j : j \leq k^* \rangle, \langle q_j : j \leq k^* \rangle$ such that for every $j \leq k^*$,

- (1) $r_j \in \mathbb{E}\mathbb{E}_\beta$,
- (2) $r_{j+1} \geq r_j$,
- (3) $r_j \Vdash_{\mathbb{E}\mathbb{E}_\beta} q_j \geq (p)_{v_j} \upharpoonright [\beta, \alpha)$,
- (4) $\forall z \in \overline{r_j} \widehat{q_j}, F(z) \upharpoonright \ell = F(x_{v_j}) \upharpoonright \ell$.

Let $q \upharpoonright \beta = q_{k^*}$ and $q \upharpoonright [\beta, \alpha) = \bigcup_{j < k^*} q_j$.

Suppose that $|F| = k + 1$ and let $\beta = \max(F)$.

By the part already proved, for each $\mathbf{x} = \langle x_\gamma : \gamma < \beta \rangle \in \overline{p \upharpoonright \beta}$ find a condition $q_{\mathbf{x}} \geq_n p \upharpoonright [\beta, \alpha)[\mathbf{x}]$ such that the sets $\left\{ F''(\overline{(q_{\mathbf{x}})_s}) : s \in \text{split}_n(q_{\mathbf{x}}) \right\}$ are pairwise disjoint. Note that we can do it in such a way that the mapping $\mathbf{x} \mapsto q_{\mathbf{x}}$ is continuous (As before we first choose $q_{\mathbf{x}}$ in a Borel way, and then shrink $p \upharpoonright \beta$ to make this mapping continuous). That defines a $\mathbb{E}\mathbb{E}_\beta$ -name for an element of $\mathbb{E}\mathbb{E}_{\beta, \alpha}$, which we call q^* .

Next, let $F' = F \setminus \{\beta\}$ and apply the inductive hypothesis to find $q' \geq_{F', n} p \upharpoonright \beta$ such that $\left\{ F''(\overline{(q')_\sigma}) : \sigma \in \Sigma_{F', n} \right\}$ are pairwise disjoint. Let $q \in \mathbb{E}\mathbb{E}_\alpha$ be defined as $q \upharpoonright \beta = q'$ and $q \upharpoonright \beta \Vdash_{\mathbb{E}\mathbb{E}_\beta} q \upharpoonright [\beta, \alpha) = q^*$.

It is clear that q is as required. \square

CASE 3. $p \Vdash_{\mathbb{E}\mathbb{E}_\alpha} \forall \beta < \alpha \dot{x} \notin \mathbf{V}^{\mathbb{E}\mathbb{E}_\beta}$.

Let $\langle F_n : k \in \omega \rangle$ be an increasing sequence of finite sets such that $\bigcup_n F_n = \alpha$.

By induction build a sequence of conditions $\langle p_n : n \in \omega \rangle$ such that $p_0 = p$ and for every n ,

- (1) $p_{n+1} \geq_{F_n, n} p_n$,
- (2) sets $\left\{ F''(\overline{(p_n)_\sigma}) : \sigma \in \Sigma_{F_n, n} \right\}$ are pairwise disjoint.

Let $q = \lim_n p_n$.

Suppose that $\mathbf{x} = \langle x_\beta : \beta < \alpha \rangle$ and $\mathbf{x}' = \langle x'_\beta : \beta < \alpha \rangle$ are two distinct points in \overline{q} . Let β be the first ordinal such that $x_\beta \neq x'_\beta$. Let n be so large that $\beta \in F_n$ and there are two distinct $\sigma, \sigma' \in \Sigma_{F_n, n}$ such that $\mathbf{x} \in \overline{(p_n)_\sigma}$ and $\mathbf{x}' \in \overline{(p_n)_{\sigma'}}$. Since $F''(\overline{(p_n)_\sigma}) \cap F''(\overline{(p_n)_{\sigma'}}) = \emptyset$, it follows that $F(\mathbf{x}) \neq F(\mathbf{x}')$. \square

4. A MODEL WHERE $\mathbf{PM} \subseteq \mathbf{UN}$.

Let $\mathbb{E}\mathbb{E}_{\omega_2}$ be the countable support iteration of $\mathbb{E}\mathbb{E}$ of length \aleph_2 . We will show that in $\mathbf{V}^{\mathbb{E}\mathbb{E}_{\omega_2}}$, $\mathbf{PM} \subseteq \mathbf{UN}$.

By theorem 9(2), $\mathbf{V}^{\mathbb{E}\mathbb{E}_{\omega_2}} \models [\mathbb{R}]^{< 2^{\aleph_0}} \subseteq \mathbf{UN}$, thus we have to show that

$$\mathbf{V}^{\mathbb{E}\mathbb{E}_{\omega_2}} \models \mathbf{PM} \subseteq [\mathbb{R}]^{< 2^{\aleph_0}}.$$

Suppose that $X \in \mathbf{V}^{\mathbb{E}\mathbb{E}_{\omega_2}}$ is a set of reals of size \aleph_2 . Let $\{\dot{x}_\alpha : \alpha < \omega_2\}$ be the set of names for elements of X such that $\Vdash_{\mathbb{E}\mathbb{E}_{\omega_2}} \forall \alpha \neq \beta \dot{x}_\alpha \neq \dot{x}_\beta$. Apply lemma 11 and find for each $\alpha < \omega_2$ a set $w_\alpha \in [\omega_2]^{< \omega}$, a condition $p_\alpha \in \mathbb{E}\mathbb{E}_{w_\alpha}$, and a continuous function $F_\alpha : \overline{p_\alpha} \rightarrow 2^\omega$ such that $p_\alpha \Vdash_{\mathbb{E}\mathbb{E}_{\omega_2}} \dot{x}_\alpha = F_\alpha(\langle \dot{g}_\beta : \beta \in w_\alpha \rangle)$. We can assume that w_α is minimal. In other words, $p_\alpha \Vdash_{\mathbb{E}\mathbb{E}_{\omega_2}} \forall \beta < \sup(w_\alpha) \dot{x}_\alpha \notin \mathbf{V}^{\mathbb{E}\mathbb{E}_\beta}$. In particular, without loss of generality we can assume F_α is one-to-one, so it is a homeomorphism.

By thinning out we can assume that $\text{ot}(w_\alpha) = \gamma$, $F_\alpha = F$ and $\overline{p_\alpha} = \overline{p}$. Moreover, since $\mathbf{V} \models \text{CH}$, we can assume that $w_\alpha \cap w_\beta = w^*$ for $\alpha \neq \beta$. Finally, without loss of generality we can assume that $w^* = \emptyset$.

Let $P = F''(\overline{p})$. Since F is a homeomorphism, P is perfect. We will show that $X \cap P$ is not meager in $\mathbf{V}^{\mathbb{E}\mathbb{E}_{\omega_2}}$ (relative to P).

Assume otherwise and let $H \subseteq P$ be a meager set such that for some $p^* \in \mathbb{E}\mathbb{E}_{\omega_2}$, $p^* \Vdash_{\mathbb{E}\mathbb{E}_{\omega_2}} X \cap P \subseteq H$. By 9(4) we can assume that $H \in \mathbf{V}$. Set $G = (F)^{-1}(H)$ and notice that G is a meager subset of \overline{p} .

Find $\alpha < \omega_2$ such that $w_\alpha \cap \text{cl}(p^*) = \emptyset$. By lemma 12 there exists $q \geq p$, $q \in \mathbb{E}\mathbb{E}_{w_\alpha} \simeq \mathbb{E}\mathbb{E}_\gamma$ such that $\overline{q} \cap G = \emptyset$.

Since p^* and q are compatible let $r \geq p^*, q$. It follows that

$$r \Vdash_{\mathbb{E}\mathbb{E}_{\omega_2}} \dot{x}_\alpha = F_\alpha(\langle \dot{q}_\beta : \beta \in w_\alpha \rangle) \notin H,$$

which finishes the proof.

Acknowledgments: The work was done while the first author was spending his sabbatical year at the Rutgers University and the College of Staten Island, CUNY, and their support is gratefully acknowledged.

REFERENCES

- [1] Tomek Bartoszyński. Measure and Category Invariants. In M. Magidor M.Foreman, A. Kanamori, editor, *Handbook of Set Theory*. submitted.
- [2] Tomek Bartoszyński. On perfectly meager sets. available from front.math.ucdavis.edu.
- [3] Tomek Bartoszyński and Haim Judah. *Set Theory: on the structure of the real line*. A.K. Peters, 1995.
- [4] Paul Corazza. The generalized Borel conjecture and strongly proper orders. *Transactions of the American Mathematical Society*, 316(1):115–140, 1989.
- [5] Martin Goldstern, Haim Judah, and Saharon Shelah. Strong measure zero sets without Cohen reals. *The Journal of Symbolic Logic*, 58(4):1323–1341, 1993.
- [6] Edward Grzegorek. Solution of a problem of Banach on σ -fields without continuous measures. *L'Academie Polonaise des Sciences. Bulletin. Serie des Sciences Mathematiques*, 28(1-2):7–10, 1980.
- [7] Arnold W. Miller. Some properties of measure and category. *Transactions of the American Mathematical Society*, 266(1):93–114, 1981.
- [8] Arnold W. Miller. Mapping a set of reals onto the reals. *The Journal of Symbolic Logic*, 48(3):575–584, 1983.
- [9] Arnold W. Miller. Special subsets of the real line. In K. Kunen and J. E. Vaughan, editors, *Handbook of Set Theoretic Topology*, pages 201–235. North-Holland, Amsterdam, 1984.
- [10] Andrzej Roslanowski and Saharon Shelah. *Norms on possibilities I: forcing with trees and creatures*. Memoirs of the American Mathematical Society. American Mathematical Society, 1999.
- [11] Saharon Shelah. *Proper and Improper Forcing*. Perspectives in Logic. Springer-Verlag, 1998.
- [12] Wacław Sierpiński. Sur une probleme de M.Kuratowski concernant la propriete de Baire des ensembles. *Fundamenta Mathematicae*, 22:262–266, 1934.
- [13] Piotr Zakrzewski. Universally meager sets. to appear in Proceedings of the American Mathematical Society.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, BOISE STATE UNIVERSITY, BOISE, IDAHO 83725 U.S.A.

Email address: tomek@math.boisestate.edu, <http://math.boisestate.edu/~tomek>

DEPARTMENT OF MATHEMATICS, HEBREW UNIVERSITY, JERUSALEM, ISRAEL AND DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW BRUNSWICK, NJ

Email address: shelah@math.huji.ac.il, <http://math.rutgers.edu/~shelah/>