

THE COMBINATORICS OF REASONABLE ULTRAFILTERS

SAHARON SHELAH

ABSTRACT. We are interested in generalizing part of the theory of ultrafilters on ω to larger cardinals. Here we set the scene for further investigations introducing properties of ultrafilters in strong sense dual to being normal.

0. INTRODUCTION

Questions concerning ultrafilters on ω have occurred to be very stimulating for research in several subareas of Set Theory and Topology. We hope that this success story could be repeated for ultrafilters on uncountable regular cardinals λ , particularly if λ is strongly inaccessible. Our aim in the present paper is to introduce new properties of ultrafilters and argue that these properties could play the stimulating role that was once played by *P*-points on ω .

In a long run, we plan to find generalizations of the following results:

- (a) Consistently, some ultrafilters on ω are generated by $< 2^{\aleph_0}$ many sets.
- (b) *P*-points are preserved by some forcing notions (see, e.g., [She98, V], [RS99]).
- (c) Consistently, there is no *P*-point.
- (d) For a function $f : \omega \rightarrow \omega$ and ultrafilter D on ω , let

$$D/f \stackrel{\text{def}}{=} \{A \subseteq \omega : f^{-1}(A) \in D\};$$

it is an ultrafilter on ω (of course, we are interested in the cases when D and D/f are uniform, which in this case is the same as non-principal). By Blass and Shelah [BS87], consistently for any two non-principal ultrafilters D_1, D_2 on ω there are finite-to-one non-decreasing functions $f_1, f_2 : \omega \rightarrow \omega$ such that $D_1/f_1 = D_2/f_2$.

Date: September 2005.

1991 *Mathematics Subject Classification.* Primary 03E05; Secondary: 03E20.

The author acknowledges support from the United States-Israel Binational Science Foundation (Grant no. 2002323). Publication 830.

- (f) For a significant family of forcing notions built according to the scheme of creatures of [RS99] we may consider an appropriate filter, i.e., if $\langle p_\alpha : \alpha < \omega_1 \rangle$ is \leq^* -increasing it may define an ultrafilter (see [RS99, §5,6]) which is not necessarily generated by \aleph_1 -sets, so we may ask on this.

There are many works on normal ultrafilters; their parallel on ω are Ramsey ultrafilters. Now, every Ramsey ultrafilter on ω is a P -point but there are P -points of very different characters, e.g., P -point with no Ramsey ultrafilter below. Gitik [Git81] has investigated generalizations of P -points for normal ultrafilters. But this paper goes in a different direction (which up to recently I have not considered to be fruitful) and we restrict our attention to ultrafilters which are very non-normal — *the weakly reasonable ultrafilters*. What is a weakly reasonable ultrafilter on λ ? It is a uniform ultrafilter on a regular cardinal λ which does not contain some club of λ and such that this property is preserved if we divide it by a non-decreasing $f : \lambda \rightarrow \lambda$ with unbounded range (see Definition 1.4 below).

We also want that our ultrafilters generalize P -points on ω and in the second section we introduce *reasonable* and *very reasonable* ultrafilters. The property defining P -points is that countable families of sets from the ultrafilter have pseudo-intersections in the ultrafilter. We modify this property so that we involve some description of how the considered ultrafilter is generated, and we postulate that *the generating systems* are suitably directed. This is a replacement for the existence of pseudo-intersections and it is the essence of Definition 2.5(4,5). The third section shows that the number of generating systems (of our type) for somewhat reasonable ultrafilters cannot be too small. We conclude the paper with a section listing open problems and describing further research.

Notation: Our notation is rather standard and compatible with that of classical textbooks (like Jech [Jec03]). In forcing we keep the older convention that *a stronger condition is the larger one*. (However, in the present paper we use forcing notions only for combinatorial constructions and almost every mention of forcing just means that we are dealing with a transitive reflexive relation $\mathbb{P} = (\mathbb{P}, \leq_{\mathbb{P}})$.)

- (1) Ordinal numbers will be denoted by the lower case initial letters of the Greek alphabet ($\alpha, \beta, \gamma, \delta \dots$) and also by i, j (with possible sub- and superscripts).
- (2) Cardinal numbers will be called κ, λ, μ ; λ will be always assumed to be a regular uncountable cardinal (we may forget to mention it).
- (3) D, \mathcal{U} will denote filters on λ , G, G^*, G_ℓ^* will be subsets of specific partial orders used to generate filters on λ .

- (4) A bar above a letter denotes that the object considered is a sequence; usually \bar{X} will be $\langle X_i : i < \zeta \rangle$, where ζ is the length $\text{lh}(\bar{X})$ of \bar{X} . Sometimes our sequences will be indexed by a set of ordinals, say $S \subseteq \lambda$, and then \bar{X} will typically be $\langle X_\delta : \delta \in S \rangle$.

Definition 0.1. A dominating family in ${}^\lambda\lambda$ is a family $\mathcal{F} \subseteq {}^\lambda\lambda$ such that

$$(\forall g \in {}^\lambda\lambda)(\exists f \in \mathcal{F})(\exists \alpha < \lambda)(\forall \beta > \alpha)(g(\beta) < f(\beta)).$$

The λ -dominating number \mathfrak{d}_λ is defined as

$$\mathfrak{d}_\lambda = \min \{ \|\mathcal{F}\| : \mathcal{F} \subseteq {}^\lambda\lambda \text{ is a dominating family in } {}^\lambda\lambda \}.$$

A club-dominating family in ${}^\lambda\lambda$ is a family $\mathcal{F} \subseteq {}^\lambda\lambda$ such that

$$(\forall g \in {}^\lambda\lambda)(\exists f \in \mathcal{F})(\{\beta < \lambda : g(\beta) \geq f(\beta)\} \text{ is non-stationary in } \lambda).$$

The $\text{cl}(\lambda)$ -dominating number $\mathfrak{d}_{\text{cl}}(\lambda)$ is defined as

$$\mathfrak{d}_{\text{cl}}(\lambda) = \min \{ \|\mathcal{F}\| : \mathcal{F} \subseteq {}^\lambda\lambda \text{ is a } \text{cl}(\lambda)\text{-dominating family in } {}^\lambda\lambda \}.$$

On $\mathfrak{d}_\lambda, \mathfrak{d}_{\text{cl}}(\lambda)$ see, e.g., in Cummings and Shelah [CS95].

Acknowledgment: I thank Tomek Bartoszyński and Andrzej Rosłanowski for stimulating discussions.

1. WEAKLY REASONABLE ULTRAFILTERS

In Definition 1.4(1) we formulate the main property of ultrafilters on λ which is of interest to us: being a weakly reasonable ultrafilter. In the spectrum of all ultrafilters, weakly reasonable ultrafilters are at the opposite end to the one occupied by normal ultrafilters. We show that there exist (in ZFC) weakly reasonable ultrafilters (see 1.10) and we also give some properties of such ultrafilters.

Definition 1.1. For a cardinal λ ,

- (a) $\text{ulf}(\lambda)$ is the set of all ultrafilters on λ ,
- (b) $\text{uuf}(\lambda)$ is the family of all uniform ultrafilters on λ ,
- (c) if D is an ultrafilter on λ and $f \in {}^\lambda\lambda$, then

$$D/f \stackrel{\text{def}}{=} \{A \subseteq \lambda : f^{-1}(A) \in D\}.$$

Let us note that in the literature D/f is also denoted by $f(D)$ or $f_*(D)$ and it is called *the image* or *the projection* of the ultrafilter D by f . We use the quotient notation and terminology because we will deal mostly with D/C , where:

Definition 1.2. Assume D is an ultrafilter on λ .

- (1) If E is an equivalence relation on λ , then $f_E \in {}^\lambda\lambda$ is defined by

$$f_E(\alpha) = \text{otp}\left(\{\beta < \alpha : \beta = \min(\beta/E) < \min(\alpha/E)\}\right),$$

and D/E is D/f_E . (Here, α/E stands for the E -equivalence class of α .)

- (2) For a club C of λ let E_C be the following equivalence relation on λ :

$$\alpha E_C \beta \quad \text{iff} \quad (\forall \gamma \in C)(\alpha < \gamma \Leftrightarrow \beta < \gamma)$$

(so E_C is the equivalence relation determined by the partition of λ into intervals $[\xi, \zeta)$ for consecutive members $\xi < \zeta$ of $C \cup \{0\}$). Let D/C be D/E_C .

- (3) \mathcal{F}_λ is the family of all non-decreasing unbounded functions from λ to λ .

Observation 1.3. *Assume that λ is a regular cardinal, $D \in \text{ufl}(\lambda)$.*

- (1) *If $f : \lambda \rightarrow \lambda$, then $D/f \in \text{ufl}(\lambda)$.*
- (2) *If $f \in \mathcal{F}_\lambda$ and D is uniform, then also D/f is a uniform ultrafilter on λ .*
- (3) *If C is a club of λ and $\langle \delta_\xi : \xi < \lambda \rangle$ is the increasing enumeration of $C \cup \{0\}$, then for a set $A \subseteq \lambda$,*

$$A \in D/C \quad \text{if and only if} \quad \bigcup \{[\delta_\xi, \delta_{\xi+1}) : \xi \in A\} \in D.$$

Definition 1.4. Let D be a uniform ultrafilter on λ .

- (1) We say that D is *weakly reasonable* if for every $f \in \mathcal{F}_\lambda$ there is a club C of λ such that

$$\bigcup \{[\delta, \delta + f(\delta)) : \delta \in C\} \notin D.$$

- (2) We define a game \mathfrak{D}_D between two players, Odd and Even, as follows. A play of \mathfrak{D}_D lasts λ steps and during a play an increasing continuous sequence $\bar{\alpha} = \langle \alpha_i : i < \lambda \rangle \subseteq \lambda$ is constructed. The terms of $\bar{\alpha}$ are chosen successively by the two players so that Even chooses the α_i for even i (including limit stages i where she has no free choice) and Odd chooses α_i for odd i .

Even wins the play if and only if

$$\bigcup \{[\alpha_{2i+1}, \alpha_{2i+2}) : i < \lambda\} \in D.$$

Observation 1.5. *Let $D \in \text{ufl}(\lambda)$. Then the following conditions are equivalent:*

- (A) *D is weakly reasonable,*

(B) for every increasing continuous sequence $\langle \delta_\xi : \xi < \lambda \rangle \subseteq \lambda$ there is a club C^* of λ such that

$$\bigcup \{[\delta_\xi, \delta_{\xi+1}) : \xi \in C^*\} \notin D,$$

(C) for every club C of λ the quotient D/C does not extend the filter generated by clubs of λ .

Proposition 1.6. *Assume $D \in \text{uuf}(\lambda)$.*

- (1) *If λ is strongly inaccessible and Odd has a winning strategy in \mathfrak{D}_D , then D is not weakly reasonable.*
- (2) *If D is not weakly reasonable, then Odd has a winning strategy in the game \mathfrak{D}_D .*
- (3) *In part (1) instead “ λ is strongly inaccessible”, it suffices to assume \diamond_λ^* .*

Proof. (1) Suppose towards contradiction that λ is strongly inaccessible, Odd has a winning strategy \mathbf{st} in the game \mathfrak{D}_D but D is weakly reasonable. By induction on $\varepsilon < \lambda$ choose an increasing continuous sequence $\langle N_\varepsilon : \varepsilon < \lambda \rangle$ of elementary submodels of $\mathcal{H}(\lambda^{++})$ so that for each ε :

- (a) $N_\varepsilon \prec (\mathcal{H}(\lambda^{++}), \in, <^*, \|N_\varepsilon\| < \lambda, N_\varepsilon \cap \lambda \in \lambda,$
- (b) ${}^\varepsilon N_{\varepsilon+1} \subseteq N_{\varepsilon+1},$
- (c) $\langle N_\zeta : \zeta \leq \varepsilon \rangle \in N_{\varepsilon+1},$
- (d) \mathbf{st}, λ, D belong to N_0 .

Let $\delta_\varepsilon = N_\varepsilon \cap \lambda$ (for $\varepsilon < \lambda$). Then $\langle \delta_\varepsilon : \varepsilon < \lambda \rangle$ is an increasing continuous sequence of limit ordinals. Let $f(\alpha) = \delta_{\alpha+1}$ for $\alpha < \lambda$, so $f \in \mathcal{F}_\lambda$.

Since D is a weakly reasonable ultrafilter, there is a club C of λ such that

$$\bigcup \{[\delta, \delta + f(\delta)) : \delta \in C\} \notin D.$$

Let

$$C^* = \{\varepsilon \in C : \varepsilon = \delta_\varepsilon \text{ is a limit ordinal}\}$$

(it is a club of λ). Then for $\varepsilon \in C^*$ we have $[\delta_\varepsilon, \delta_{\varepsilon+1}) \subseteq [\varepsilon, \varepsilon + f(\varepsilon))$ and hence

$$\bigcup \{[\delta_\varepsilon, \delta_{\varepsilon+1}) : \varepsilon \in C^*\} \notin D.$$

Let us define a strategy \mathbf{st}' for Even in the game \mathfrak{D}_D as follows. For an even ordinal $i < \lambda$, in the i -th move of a play, if $\langle \alpha_j : j < i \rangle$ has been played so far then Even plays

$$\alpha_i = \begin{cases} \sup\{\alpha_j : j < i\} & \text{if } i \text{ is limit,} \\ \min\{\varepsilon \in C^* : (\forall j < i)(\alpha_j < \varepsilon)\} & \text{otherwise.} \end{cases}$$

Now consider a play $\langle \alpha_i : i < \lambda \rangle$ in which Even uses the strategy \mathbf{st}' and Odd plays according to \mathbf{st} . Then for each $i < \lambda$ we have $\alpha_{2i} \in C^*$ and thus $\alpha_{2i} = \delta_{\alpha_{2i}} \in N_{\alpha_{2i}+1}$, and also $\{\alpha_j : j < 2i\} \subseteq \alpha_{2i} \subseteq N_{\alpha_{2i}+1}$. Since the

model $N_{\alpha_{2i+1}}$ is closed under forming sequences of length $\alpha_{2i} + 1$ (by (b)), we conclude that $\langle \alpha_j : j \leq 2i \rangle \in N_{\alpha_{2i+1}}$. Since $\mathbf{st} \in N_0 \prec N_{\alpha_{2i+1}}$, clearly $\alpha_{2i+1} \in N_{\alpha_{2i+1}} \cap \lambda$ and therefore $\alpha_{2i+1} < \delta_{\alpha_{2i+1}}$. Hence

$$\bigcup \{[\alpha_{2i}, \alpha_{2i+1}) : i < \lambda\} \subseteq \bigcup \{[\delta_{\alpha_{2i}}, \delta_{\alpha_{2i+1}}) : i < \lambda\} \subseteq \bigcup \{[\delta_\varepsilon, \delta_{\varepsilon+1}) : \varepsilon \in C^*\} \notin D.$$

But \mathbf{st} is a winning strategy for Odd, so he wins the play and

$$\bigcup \{[\alpha_{2i+1}, \alpha_{2i+2}) : i < \lambda\} \notin D,$$

a contradiction.

(2) Suppose that $D \in \text{uuf}(\lambda)$ is not weakly reasonable. Then we may find $f \in \mathcal{F}_\lambda$ such that for every club C of λ we have

$$\bigcup \{[\delta, \delta + f(\delta)) : \delta \in C\} \in D.$$

Let \mathbf{st} be a strategy of Odd in \mathcal{D}_D which instructs him to play as follows. For an odd ordinal $i = i_0 + 1 < \lambda$, in the i -th move of a play, if $\langle \alpha_j : j \leq i_0 \rangle$ has been played so far, then Odd plays $\alpha_i = \alpha_{i_0} + f(\alpha_{i_0}) + 1$.

We claim that \mathbf{st} is a winning strategy for Odd (in \mathcal{D}_D). To this end suppose that $\langle \alpha_j : j < \lambda \rangle \subseteq \lambda$ is a result of a play of \mathcal{D}_D in which Odd uses the strategy \mathbf{st} . Let $C' = \{\alpha_i : i < \lambda \text{ is limit}\}$ – it is a club of λ , so by the choice of f we have

$$\bigcup \{[\delta, \delta + f(\delta)) : \delta \in C'\} \in D.$$

Since $\bigcup \{[\delta, \delta + f(\delta)) : \delta \in C'\} \subseteq \bigcup \{[\alpha_{2i}, \alpha_{\alpha_{2i+1}}) : i < \lambda\}$ we may now conclude that Odd indeed wins the play. \square

Remark 1.7. Let us note that some assumptions on λ in 1.6(1) are needed. This will be shown in the subsequent paper Rosłanowski and Shelah [RS08].

Lemma 1.8. *Suppose that λ is a regular uncountable cardinal, $D \in \text{uuf}(\lambda)$ is a weakly reasonable ultrafilter and $\langle \beta_i : i < \lambda \rangle$ is an increasing continuous sequence of ordinals below λ . Then there is an increasing continuous sequence $\langle \delta_\xi : \xi < \lambda \rangle \subseteq \lambda$ consisting of limit ordinals and such that*

$$\bigcup \{[\beta_{\delta_{2\xi+1}}, \beta_{\delta_{2\xi+2}}) : \xi < \lambda\} \in D.$$

Proof. It follows from 1.5 that we may find a club C^* of λ such that all members of C^* are limit ordinals and $\bigcup \{[\beta_\xi, \beta_{\xi+1}) : \xi \in C^*\} \notin D$. Let $C^+ = C^* \cup \{\xi + 1 : \xi \in C^*\}$ (clearly it is a club of λ) and let $\langle \delta_\xi : \xi < \lambda \rangle$ be the increasing enumeration of C^+ . Note that $C^* = \{\delta_\xi : \xi < \lambda \text{ is even}\}$ and, for an even ordinal $\xi < \lambda$, $\delta_{\xi+1} = \delta_\xi + 1$. Hence

$$\bigcup \{[\beta_{\delta_\xi}, \beta_{\delta_{\xi+1}}) : \xi < \lambda \text{ is even}\} = \bigcup \{[\beta_{\delta_\xi}, \beta_{\delta_{\xi+1}}) : \xi < \lambda \text{ is even}\} = \bigcup \{[\beta_\zeta, \beta_{\zeta+1}) : \zeta \in C^*\} \notin D$$

Consequently, $\bigcup \{[\beta_{\delta_\xi}, \beta_{\delta_{\xi+1}}] : \xi < \lambda \text{ is odd}\} \in D$. \square

Theorem 1.9. *If λ is a regular uncountable cardinal and $D \in \text{uuf}(\lambda)$ is weakly reasonable, then D is a regular ultrafilter.*

Proof. Using Lemma 1.8 we may choose by induction on $\varepsilon < \lambda$ a sequence $\langle \bar{\delta}^\varepsilon : \varepsilon < \lambda \rangle$ so that

- (a) $\bar{\delta}^\varepsilon = \langle \delta_i^\varepsilon : i < \lambda \rangle$ is an increasing continuous sequence of non-successor ordinals below λ , $\delta_0^\varepsilon = 0$,
- (b) the set $A_\varepsilon \stackrel{\text{def}}{=} \bigcup \{[\delta_{2i+1}^\varepsilon, \delta_{2i+2}^\varepsilon] : i < \lambda\}$ belongs to D ,
- (c) if $\zeta < \varepsilon$, $i < \lambda$, then $\delta_i^\varepsilon \in \{\delta_j^\zeta : j < \lambda \text{ is a limit ordinal or zero}\}$.

For $\varepsilon < \lambda$ let $f_\varepsilon : A_\varepsilon \rightarrow \lambda$ be such that

$$\alpha \in [\delta_{2i+1}^\varepsilon, \delta_{2i+2}^\varepsilon) \Rightarrow f_\varepsilon(\alpha) = \delta_{2i+1}^\varepsilon.$$

Note that

- (\otimes) if $\zeta < \varepsilon < \lambda$, $\alpha \in A_\zeta \cap A_\varepsilon$, then $f_\varepsilon(\alpha) < f_\zeta(\alpha)$.

[Why? Let $f_\zeta(\alpha) = \delta_{2i+1}^\zeta$ (so $\alpha \in [\delta_{2i+1}^\zeta, \delta_{2i+2}^\zeta)$). It follows from (c) that $f_\varepsilon(\alpha) \in \{\delta_j^\zeta : j < \lambda \text{ is a limit ordinal or zero}\}$ and hence (since also $f_\varepsilon(\alpha) \leq \alpha$) we may conclude that $f_\varepsilon(\alpha) < f_\zeta(\alpha)$.]

For $\alpha < \lambda$, let $w_\alpha = \{\varepsilon < \lambda : \alpha \in A_\varepsilon\}$. It follows from (\otimes) that (for every $\alpha < \lambda$) the sequence $\langle f_\varepsilon(\alpha) : \varepsilon \in w_\alpha \rangle$ is strictly decreasing, so necessarily each w_α is finite. Since $A_\varepsilon \in D$ for each $\varepsilon < \lambda$ (by (b)), we have shown the regularity of D . \square

Theorem 1.10. *Let $\lambda > \aleph_0$ be a regular cardinal. Then there is a uniform weakly reasonable ultrafilter D on λ .*

Proof. Let $\{f_\varepsilon : \varepsilon < \mathfrak{d}_\lambda\} \subseteq {}^\lambda \lambda$ be a dominating family and for $\varepsilon < \mathfrak{d}_\lambda$ let C_ε be a club of λ such that members of C_ε are limit ordinals and

$$(\forall \delta \in C_\varepsilon)(\forall \alpha < \delta)(f_\varepsilon(\alpha) < \delta).$$

Let $\langle \alpha_{\varepsilon,i} : i < \lambda \rangle$ be the increasing enumeration of C_ε .

By induction on ε we will choose sets $E_\varepsilon, A_\varepsilon$ so that for each $\varepsilon < \mathfrak{d}_\lambda$:

- (a) A_ε is an unbounded subset of λ and $E_\varepsilon \subseteq C_\varepsilon$ is a club of λ ,
- (b) $A_\varepsilon \cap \bigcup \{[\alpha_{\varepsilon,\gamma}, \alpha_{\varepsilon,\gamma+1}) : \gamma \in E_\varepsilon\} = \emptyset$,
- (c) if $n < \omega$, $\zeta_0 < \dots < \zeta_{n-1} < \varepsilon$, then $\|A_\varepsilon \cap \bigcap_{i < n} A_{\zeta_i}\| = \lambda$.

So suppose that we have chosen A_ζ, E_ζ for $\zeta < \varepsilon < \mathfrak{d}_\lambda$ so that the respective reformulations of (a)–(c) hold true. For a finite sequence $\bar{\zeta} = \langle \zeta_i : i < n \rangle$ of ordinals below ε let $A^{\bar{\zeta}} = \bigcap_{i < n} A_{\zeta_i}$ (note that $\|A^{\bar{\zeta}}\| = \lambda$ by the demand in

(c)). Let $g_{\bar{\zeta}}^\varepsilon \in {}^\lambda \lambda$ be such that

(\oplus) if $\alpha_{\varepsilon,i} \leq \alpha < \alpha_{\varepsilon,i+1}$, then $g_{\bar{\zeta}}^{\varepsilon}(\alpha) = \min\{\delta > \alpha_{\varepsilon,i+1} : [\alpha_{\varepsilon,i+1}, \delta) \cap A^{\bar{\zeta}} \neq \emptyset\}$.

The family $\{g_{\bar{\zeta}}^{\varepsilon} : \bar{\zeta} \in {}^{\omega}>\varepsilon\}$ is a subset of ${}^{\lambda}\lambda$ of cardinality $\leq |\varepsilon| + \aleph_0 < \mathfrak{d}_{\lambda}$, so it cannot be a dominating family. Therefore we may pick a function $h_{\varepsilon} \in {}^{\lambda}\lambda$ such that

$$(\forall \bar{\zeta} \in {}^{\omega}>\varepsilon)(\exists^{\lambda} \alpha < \lambda)(g_{\bar{\zeta}}^{\varepsilon}(\alpha) < h_{\varepsilon}(\alpha)).$$

Put

$$E_{\varepsilon} = \{\delta < \lambda : \delta = \alpha_{\varepsilon,\delta} \text{ is a limit ordinal and } (\forall \alpha < \delta)(h_{\varepsilon}(\alpha) < \delta)\} \quad \text{and} \\ A_{\varepsilon} = \bigcup \{[\alpha_{\varepsilon,\gamma+1}, \alpha_{\varepsilon,\delta}) : \gamma < \delta \text{ are successive members of } E_{\varepsilon}\}.$$

It should be clear that $E_{\varepsilon}, A_{\varepsilon}$ satisfy demands (a), (b).

Let us argue that also condition (c) holds true. Let $\bar{\zeta} \in {}^{\omega}>\varepsilon$ and we shall prove that $A_{\varepsilon} \cap A^{\bar{\zeta}}$ is unbounded in λ . By the choice of h_{ε} , the set $B = \{\alpha < \lambda : g_{\bar{\zeta}}^{\varepsilon}(\alpha) < h_{\varepsilon}(\alpha)\}$ is of cardinality λ . Let us fix for a moment $\alpha \in B$ and let $i < \lambda$ be such that $\alpha_{\varepsilon,i} \leq \alpha < \alpha_{\varepsilon,i+1}$. Let $\sup(E_{\varepsilon} \cap \alpha_{\varepsilon,i+1}) = \gamma = \alpha_{\varepsilon,\gamma}$ and $\min(E_{\varepsilon} \setminus \alpha_{\varepsilon,\gamma+1}) = \delta = \alpha_{\varepsilon,\delta}$. Then γ, δ are successive members of E_{ε} and

$$\gamma \leq \alpha_{\varepsilon,i} \leq \alpha < \alpha_{\varepsilon,i+1} < \delta.$$

Hence (by the definition of E_{ε} and by $\alpha \in B$) we get

$$[\alpha_{\varepsilon,i+1}, g_{\bar{\zeta}}^{\varepsilon}(\alpha)) \subseteq [\alpha_{\varepsilon,i+1}, h_{\varepsilon}(\alpha)) \subseteq [\alpha_{\varepsilon,\gamma+1}, \alpha_{\varepsilon,\delta}) \subseteq A_{\varepsilon}.$$

It follows from (\oplus) that $[\alpha_{\varepsilon,i+1}, g_{\bar{\zeta}}^{\varepsilon}(\alpha)) \cap A^{\bar{\zeta}} \neq \emptyset$, and consequently $A_{\varepsilon} \cap A^{\bar{\zeta}} \setminus \alpha \neq \emptyset$. Since $\|B\| = \lambda$ we may now easily conclude that $\|A_{\varepsilon} \cap A^{\bar{\zeta}}\| = \lambda$, showing that $A_{\varepsilon}, E_{\varepsilon}$ are as required.

After the construction is carried out (and we have the sequence $\langle E_{\varepsilon}, A_{\varepsilon} : \varepsilon < \mathfrak{d}_{\lambda} \rangle$) we may find a uniform ultrafilter D on λ such that $\{A_{\varepsilon} : \varepsilon < \mathfrak{d}_{\lambda}\} \subseteq D$ (remember the demand in (c)). We claim that D is weakly reasonable. To this end suppose that C is a club of λ and $\langle \delta_{\xi} : \xi < \lambda \rangle \subseteq \lambda$ is the increasing enumeration of C . By the choice of $f_{\varepsilon}, C_{\varepsilon}$ (for $\varepsilon < \mathfrak{d}_{\lambda}$) we may find $\varepsilon < \mathfrak{d}_{\lambda}$ and $j_0 < \lambda$ such that

$$(\forall i \geq j_0)(\| [\alpha_{\varepsilon,i}, \alpha_{\varepsilon,i+1}) \cap C \| > 2).$$

Let

$$C^* = \{\gamma \in E_{\varepsilon} \cap C \setminus j_0 : \gamma = \alpha_{\varepsilon,\gamma} = \delta_{\gamma} \text{ is a limit ordinal}\}$$

(it is a club of λ). Since for $\gamma \in C^*$ we have that $\alpha_{\varepsilon,\gamma} = \delta_{\gamma} < \delta_{\gamma+1} < \alpha_{\varepsilon,\gamma+1}$ we may easily conclude from (b) that

$$\bigcup \{[\delta_{\gamma}, \delta_{\gamma+1}) : \gamma \in C^*\} \notin D,$$

completing the proof (remember 1.5). □

2. MORE REASONABLE ULTRAFILTERS

In this section we propose a property of ultrafilters stronger than being weakly reasonable (see Definition 2.5(5)). We believe that the notion of *very reasonable ultrafilters* is the right re-interpretation of being a P -point in the setting of “very non-normal ultrafilters” on an uncountable regular cardinal λ . We start with describing a forcing notion \mathbb{Q}_λ^1 which motivated our choice of *generating systems* of 2.5.

Like before, λ is always assumed to be an uncountable regular cardinal.

Definition 2.1. We define a forcing notion \mathbb{Q}_λ^1 as follows.

A condition in \mathbb{Q}_λ^1 is a tuple $p = (\gamma^p, C^p, \langle Z_\delta^p : \delta \in C^p \rangle, \langle d_\delta^p : \delta \in C^p \rangle)$ such that

- (i) $\gamma^p < \lambda$, C^p a club of λ consisting of limit ordinals only, and for $\delta \in C^p$:
- (ii) $Z_\delta^p = [\delta, \min(C^p \setminus (\delta + 1)))$ and
- (iii) $d_\delta^p \subseteq \mathcal{P}(Z_\delta^p)$ is a proper ultrafilter on Z_δ^p .

The order $\leq_{\mathbb{Q}_\lambda^1} = \leq$ of \mathbb{Q}_λ^1 is given by

$p \leq_{\mathbb{Q}_\lambda^1} q$ if and only if

- (a) $\gamma^p \leq \gamma^q$, $C^p \cap \gamma^p \subseteq C^q \subseteq C^p$ and
- (b) if $\delta < \varepsilon$ are successive members of C^q (so $Z_\delta^q = [\delta, \varepsilon)$), then

$$(\forall A \in d_\delta^q)(\exists \zeta \in C^p \cap [\delta, \varepsilon))(A \cap Z_\zeta^p \in d_\zeta^p).$$

Remark 2.2. The forcing notion \mathbb{Q}_λ^1 can be represented according to the framework of [RS07, §B.5].

Proposition 2.3. (1) \mathbb{Q}_λ^1 is a partial order, $\|\mathbb{Q}_\lambda^1\| = 2^{2^{<\lambda}}$.

(2) If $p, q \in \mathbb{Q}_\lambda^1$, $p \leq q$, $\delta < \varepsilon$ are two successive members of C^p , $\delta, \varepsilon \in C^q$, then $Z_\delta^q = Z_\delta^p$ and $d_\delta^q = d_\delta^p$.

(3) \mathbb{Q}_λ^1 is $(<\lambda)$ -complete (so it does not add bounded subsets of λ).

(4) If $p \in \mathbb{Q}_\lambda^1$, $A \subseteq \lambda$, then there is a condition $q \in \mathbb{Q}_\lambda^1$ stronger than p and such that

$$\text{either } (\forall \delta \in C^q \setminus \gamma^p)(A \cap Z_\delta^q \in d_\delta^q) \quad \text{or} \quad (\forall \delta \in C^q \setminus \gamma^p)(A \cap Z_\delta^q \notin d_\delta^q).$$

Proof. (1), (2) Straightforward.

(3) Assume that $\delta < \lambda$ is a limit ordinal and a sequence $\langle p_i : i < \delta \rangle \subseteq \mathbb{Q}_\lambda^1$ is $\leq_{\mathbb{Q}_\lambda^1}$ -increasing. Let E be a uniform ultrafilter on δ . Let us put:

- $\gamma = \sup\{\gamma^{p_i} : i < \delta\}$, $C = \bigcap_{i < \delta} C^{p_i}$, and for $\alpha \in C$ let
- $Z_\alpha = [\alpha, \min(C \setminus (\alpha + 1)))$ and
- $d_\alpha = \{A \subseteq Z_\alpha : \{i < \delta : A \cap Z_\alpha^{p_i} \in d_\alpha^{p_i}\} \in E\}$.

It is easy to check that $p = (\gamma, C, \langle Z_\alpha : \alpha \in C \rangle, \langle d_\alpha : \alpha \in C \rangle)$ belongs to \mathbb{Q}_λ^1 and that it is a condition stronger than all p_i (for $i < \delta$).

(4) Let $p \in \mathbb{Q}_\lambda^1$ and $A \subseteq \lambda$. Just for simplicity we may assume that $\gamma^p \in C^p$ (as we may always increase γ^p). Put

$$Y \stackrel{\text{def}}{=} \{\alpha \in C^p : A \cap Z_\alpha^p \in d_\alpha^p\}$$

and let us consider two cases.

CASE 1: Y is unbounded in λ .

Then we may choose an increasing continuous sequence $\langle \delta_i : i < \lambda \rangle \subseteq C^p$ such that $\delta_0 = \gamma^p$ and $(\forall i < \lambda)([\delta_i, \delta_{i+1}) \cap Y \neq \emptyset)$. Put

- $\gamma = \gamma^p$, $C = \{\delta_i : i < \lambda\} \cup (C^p \cap \gamma^p)$,
- if $\alpha \in C^p \cap \gamma^p$, then $Z_\alpha = Z_\alpha^p$ and $d_\alpha = d_\alpha^p$,
- if $\alpha = \delta_i$, $i < \lambda$, then $Z_\alpha = [\delta_i, \delta_{i+1})$ and

$$d_\alpha = \{B \subseteq Z_\alpha : B \cap Z_{\min(Y \setminus \alpha)}^p \in d_{\min(Y \setminus \alpha)}^p\}.$$

It is straightforward to verify that $q = (\gamma, C, \langle Z_\alpha : \alpha \in C \rangle, \langle d_\alpha : \alpha \in C \rangle) \in \mathbb{Q}_\lambda^1$ is a condition stronger than p and it is also clear that $(\forall \alpha \in C \setminus \gamma^p)(A \cap Z_\alpha \in d_\alpha)$.

CASE 2: Y is bounded in λ .

Then the set $\lambda \setminus Y$ is unbounded, so we may apply the construction of q from Case 1 replacing Y by its complement $\lambda \setminus Y$. It should be clear that the condition q which we will get then satisfies $(\forall \alpha \in C^q \setminus \gamma^p)(A \cap Z_\alpha^q \notin d_\alpha^q)$. \square

Remark 2.4. The following discussion presents our motivations for the definitions and concepts presented later in this section.

Suppose that $G \subseteq \mathbb{Q}_\lambda^1$ is a generic filter over \mathbf{V} . In $\mathbf{V}[G]$ we define $C = \bigcup \{C^p \cap \gamma^p : p \in G\}$ and for $\alpha \in C$ we let $d_\alpha = d_\alpha^p$ for some (equivalently: all) $p \in G$ such that $\alpha < \gamma^p$ and $C^p \cap (\alpha, \gamma^p) \neq \emptyset$. Then C is a club of λ and (for $\alpha \in C$) d_α is an ultrafilter on $[\alpha, \min(C \setminus (\alpha + 1)))$. Let

$$D = \{A \in \mathcal{P}(\lambda)^{\mathbf{V}} : (\exists \varepsilon < \lambda)(\forall \alpha > \varepsilon)(A \cap [\alpha, \min(C \setminus (\alpha + 1))) \in d_\alpha)\}.$$

It follows from 2.3(4) that D is an ultrafilter on the Boolean algebra $\mathcal{P}(\lambda)^{\mathbf{V}}$.

Let \underline{D} be a \mathbb{Q}_λ^1 -name for the D defined as above. Note that if $p \in \mathbb{Q}_\lambda^1$, $A \subseteq \lambda$ and $(\exists \varepsilon < \lambda)(\forall \delta \in C^p \setminus \varepsilon)(A \cap Z_\delta^p \in d_\delta^p)$, then $p \Vdash_{\mathbb{Q}_\lambda^1} \text{“} A \in \underline{D} \text{”}$. Plainly, the family $\{A \subseteq \lambda : p \Vdash_{\mathbb{Q}_\lambda^1} \text{“} A \in \underline{D} \text{”}\}$ is a uniform filter on λ , and, of course, for a generic filter $G \subseteq \mathbb{Q}_\lambda^1$ over \mathbf{V} ,

$$D^G = \bigcup \left\{ \{A \subseteq \lambda : p \Vdash_{\mathbb{Q}_\lambda^1} \text{“} A \in \underline{D} \text{”}\} : p \in G \right\}.$$

Definition 2.5. (1) We define a forcing notion \mathbb{Q}_λ^0 as follows.

A condition in \mathbb{Q}_λ^0 is a tuple $p = (C^p, \langle Z_\delta^p : \delta \in C^p \rangle, \langle d_\delta^p : \delta \in C^p \rangle)$

such that $(0, C^p, \langle Z_\delta^p : \delta \in C^p \rangle, \langle d_\delta^p : \delta \in C^p \rangle) \in \mathbb{Q}_\lambda^1$;

The order $\leq_{\mathbb{Q}_\lambda^0} = \leq$ **of** \mathbb{Q}_λ^0 is inherited from \mathbb{Q}_λ^1 in a natural way.

- (2) We define a relation $\leq_{\mathbb{Q}_\lambda^0}^* = \leq^*$ on \mathbb{Q}_λ^0 as follows:

$p \leq^* q$ if and only if for some $\alpha < \lambda$ we have

$$(C^p \setminus \alpha, \langle Z_\delta^p : \delta \in C^p \setminus \alpha \rangle, \langle d_\delta^p : \delta \in C^p \setminus \alpha \rangle) \leq_{\mathbb{Q}_\lambda^0} (C^q \setminus \alpha, \langle Z_\delta^q : \delta \in C^q \setminus \alpha \rangle, \langle d_\delta^q : \delta \in C^q \setminus \alpha \rangle).$$

- (3) For a condition $q \in \mathbb{Q}_\lambda^0$ we let

$$\text{fil}(q) \stackrel{\text{def}}{=} \{A \subseteq \lambda : (\exists \varepsilon < \lambda)(\forall \delta \in C^q \setminus \varepsilon)(A \cap Z_\delta^q \in d_\delta^q)\},$$

and for a set $G^* \subseteq \mathbb{Q}_\lambda^0$ we let $\text{fil}(G^*) \stackrel{\text{def}}{=} \bigcup \{\text{fil}(p) : p \in G^*\}$. We also define a binary relation \leq^0 on \mathbb{Q}_λ^0 by

$$p \leq^0 q \text{ if and only if } \text{fil}(p) \subseteq \text{fil}(q).$$

- (4) We say that an ultrafilter D on λ is *reasonable* if it is weakly reasonable (see 1.4(1)) and there is a directed (with respect to \leq^0) set $G^* \subseteq \mathbb{Q}_\lambda^0$ such that $D = \text{fil}(G^*)$. The family G^* may be called *the generating system for* D .
- (5) An ultrafilter D on λ is said to be *very reasonable* if it is weakly reasonable and there is a $(< \lambda^+)$ -directed (with respect to \leq^0) set $G^* \subseteq \mathbb{Q}_\lambda^0$ such that $D = \text{fil}(G^*)$.

Remark 2.6. Note that $\|\text{fil}(p)\| = 2^\lambda$ whenever $p \in \mathbb{Q}_\lambda^0$ is such that all ultrafilters d_α^p (for $\alpha \in C^p$) are not principal. Thus even if $D = \text{fil}(G^*)$ for some small generating system $G^* \subseteq \mathbb{Q}_\lambda^0$, the minimal number of generators for D as a filter may be 2^λ .

Observation 2.7. (1) If $p \leq_{\mathbb{Q}_\lambda^0}^* q$, then $\text{fil}(p) \subseteq \text{fil}(q)$ (so $p \leq^0 q$).

- (2) If a set $G^* \subseteq \mathbb{Q}_\lambda^0$ is directed with respect to \leq^0 , then $\text{fil}(G^*)$ is a filter of subsets of λ containing all co-bounded subsets of λ .

Definition 2.8. Suppose that

- (a) X is a non-empty set and e is an ultrafilter on X ,
 (b) d_x is an ultrafilter on a set Z_x (for $x \in X$).

We let

$$\bigoplus_{x \in X}^e d_x = \{A \subseteq \bigcup_{x \in X} Z_x : \{x \in X : Z_x \cap A \in d_x\} \in e\}.$$

(Clearly, $\bigoplus_{x \in X}^e d_x$ is an ultrafilter on $\bigcup_{x \in X} Z_x$.)

Proposition 2.9. Let $p, q \in \mathbb{Q}_\lambda^0$. Then the following are equivalent:

- (a) $p \leq^0 q$,

(b) *there is $\varepsilon < \lambda$ such that*

$$(\forall \alpha \in C^q \setminus \varepsilon) (\forall A \in d_\alpha^q) (\exists \beta \in C^p) (A \cap Z_\beta^p \in d_\beta^p),$$

(c) *there is $\varepsilon < \lambda$ such that*

if $\alpha \in C^q \setminus \varepsilon$, $\beta_0 = \sup(C^p \cap (\alpha + 1))$, $\beta_1 = \min(C^p \setminus \min(C^q \setminus (\alpha + 1)))$,

then there is an ultrafilter e on $[\beta_0, \beta_1) \cap C^p$ such that

$$d_\alpha^q = \{A \cap Z_\alpha^q : A \in \bigoplus^e \{d_\beta^p : \beta \in [\beta_0, \beta_1) \cap C^p\}\}.$$

Proof. (a) \Rightarrow (b) Assume towards contradiction that $p \leq^0 q$, but (b) fails. Then we may pick a sequence $\langle \alpha_\xi, A_\xi : \xi < \lambda \rangle$ such that for each $\xi < \lambda$,

- (i) $\alpha_\xi \in C^q$, $A_\xi \in d_{\alpha_\xi}^q$,
- (ii) if $\xi < \zeta < \lambda$, $\beta \in C^p \cap \min(C^q \setminus (\alpha_\xi + 1))$, then $\min(C^p \setminus (\beta + 1)) < \alpha_\zeta$,
- (iii) $(\forall \beta \in C^p) (A_\xi \cap Z_\beta^p \notin d_\beta^p)$.

It follows from (ii) that for every $\beta \in C^p$ there is at most one $\xi < \lambda$ such that $Z_\beta^p \cap Z_{\alpha_\xi}^q \neq \emptyset$. Also if $\beta \in C^p$ and $Z_\beta^p \cap Z_{\alpha_\xi}^q \in d_\beta^p$, then $(Z_{\alpha_\xi}^q \setminus A_\xi) \cap Z_\beta^p \in d_\beta^p$.

Put $A = \bigcup_{\xi < \lambda} A_\xi$. By what we have said above, for all $\beta \in C^p$ we have $(\lambda \setminus A) \cap Z_\beta^p \in d_\beta^p$, and hence $\lambda \setminus A \in \text{fil}(p) \subseteq \text{fil}(q)$. This contradicts (i).

(b) \Rightarrow (c) Assume that (b) holds true as witnessed by $\varepsilon < \lambda$. Let $\alpha \in C^q \setminus \varepsilon$, $\alpha' = \min(C^q \setminus (\alpha + 1))$, $\beta_0 = \sup(C^p \cap (\alpha + 1))$ and $\beta_1 = \min(C^p \setminus \alpha')$. For $A \in d_\alpha^q$ put

$$w(A) = \{\beta \in [\beta_0, \beta_1) \cap C^p : A \cap Z_\beta^p \in d_\beta^p\}.$$

It follows from (b) that $w(A) \neq \emptyset$. Plainly $w(A \cap A') = w(A) \cap w(A')$ for $A, A' \in d_\alpha^q$, so we may pick an ultrafilter e on $[\beta_0, \beta_1) \cap C^p$ such that $\{w(A) : A \in d_\alpha^q\} \subseteq e$. Now it should be clear that

$$d_\alpha^q \subseteq \{B \cap Z_\alpha^q : B \in \bigoplus^e \{d_\beta^p : \beta \in [\beta_0, \beta_1) \cap C^p\}\}$$

and (since clearly $Z_\alpha^q \in \bigoplus^e \{d_\beta^p : \beta \in [\beta_0, \beta_1) \cap C^p\}$) the set on the right-hand side is a proper filter on Z_α^q . Consequently the two sets are equal.

(c) \Rightarrow (a) Assume that (c) holds true as witnessed by $\varepsilon < \lambda$, and suppose that $A \in \text{fil}(p)$. Pick $\varepsilon' < \lambda$ such that $\varepsilon < \varepsilon'$ and

$$(\forall \beta \in C^p \setminus \varepsilon') (A \cap Z_\beta^p \in d_\beta^p).$$

Suppose $\alpha \in C^q \setminus (\min(C^p \setminus \varepsilon') + 1)$ and let $\beta_0 = \sup(C^p \cap (\alpha + 1))$, $\beta_1 = \min(C^p \setminus \min(C^q \setminus (\alpha + 1)))$. Let e be an ultrafilter on $[\beta_0, \beta_1) \cap C^p$

such that

$$d_\alpha^q = \{B \cap Z_\alpha^q : B \in \bigoplus^e \{d_\beta^p : \beta \in [\beta_0, \beta_1) \cap C^p\}\}.$$

Note that $\beta_0 \geq \varepsilon'$ and hence $A \cap Z_\beta^p \in d_\beta^p$ for all $\beta \in [\beta_0, \beta_1) \cap C^p$. Consequently

$$A \cap [\beta_0, \beta_1) \in \bigoplus^e \{d_\beta^p : \beta \in [\beta_0, \beta_1) \cap C^p\}$$

and therefore also

$$A \cap Z_\alpha^q = (A \cap [\beta_0, \beta_1)) \cap Z_\alpha^q \in d_\alpha^q.$$

Now we easily conclude that $A \in \text{fil}(q)$. □

Definition 2.10. Let $p \in \mathbb{Q}_\lambda^0$. Suppose that $X \in [C^p]^\lambda$ and $C \subseteq C^p$ is a club of λ such that

if $\alpha < \beta$ are successive elements of C ,
then $|\llbracket \alpha, \beta \rrbracket \cap X| = 1$.

(In this situation we say that p is *restrictable to* $\langle X, C \rangle$.) We define *the restriction of p to $\langle X, C \rangle$* as an element $q = p \upharpoonright \langle X, C \rangle \in \mathbb{Q}_\lambda^0$ such that $C^q = C$, and if $\alpha < \beta$ are successive elements of C , $x \in [\alpha, \beta) \cap X$, then $Z_\alpha^q = [\alpha, \beta)$ and $d_\alpha^q = \{A \subseteq Z_\alpha^q : A \cap Z_x^p \in d_x^p\}$.

Proposition 2.11. (1) Assume that $G^* \subseteq \mathbb{Q}_\lambda^0$ is \leq^0 -directed and \leq^0 -downward closed, $p \in G^*$, $X \in [C^p]^\lambda$ and $C \subseteq C^p$ is a club of λ such that p is restrictable to $\langle X, C \rangle$. If $\bigcup_{x \in X} Z_x^p \in \text{fil}(G^*)$, then $p \upharpoonright \langle X, C \rangle \in G^*$.
(2) If $G^* \subseteq \mathbb{Q}_\lambda^0$ is \leq^0 -directed and $\|G^*\| \leq \lambda$, then G^* has a \leq^0 -upper bound. (Hence, in particular, $\text{fil}(G^*)$ is not an ultrafilter.)

Proof. (1) Suppose that G^*, p, X, C are as in the assumptions and $\bigcup_{x \in X} Z_x^p \in \text{fil}(G^*)$. Since G^* is \leq^0 -directed (and $p \in G^*$) we may pick $r \in G^*$ such that $p \leq^0 r$ and $\bigcup_{x \in X} Z_x^p \in \text{fil}(r)$. We are going to show that $q \stackrel{\text{def}}{=} p \upharpoonright \langle X, C \rangle \leq^0 r$ (which will imply that $q \in G^*$ as G^* is downward closed).

Since $\bigcup_{x \in X} Z_x^p \in \text{fil}(r)$, there is $\varepsilon < \lambda$ such that

$$(\forall \alpha \in C^r \setminus \varepsilon) \left(\bigcup_{x \in X} Z_x^p \cap Z_\alpha^r \in d_\alpha^r \right)$$

and

$$(\alpha \in C^r \setminus \varepsilon) (\forall A \in d_\alpha^r) (\exists \beta \in C^p) (A \cap Z_\beta^p \in d_\beta^p)$$

(remember 2.9(b)). Now suppose that $\alpha \in C^r \setminus \varepsilon$ and $A \in d_\alpha^r$. Then $\bigcup_{x \in X} Z_x^p \cap A \in d_\alpha^r$ so there is $\beta \in C^p$ such that $\bigcup_{x \in X} Z_x^p \cap A \cap Z_\beta^p \in d_\beta^p$. In particular, $\bigcup_{x \in X} Z_x^p \cap Z_\beta^p \cap A \neq \emptyset$, so necessarily $\beta \in X$. Let $\beta_0 < \beta_1$ be the successive elements of C such that $\beta_0 \leq \beta < \beta_1$. Since

$$Z_\beta^p \cap A = \bigcup_{x \in X} Z_x^p \cap Z_\beta^p \cap A \in d_\beta^p,$$

we also have $A \cap Z_{\beta_0}^q \in d_{\beta_0}^q$. Thus we have shown that

$$\begin{aligned} & \text{if } \alpha \in C^r \setminus \varepsilon \text{ and } A \in d_\alpha^r, \\ & \text{then there is } \beta_0 \in C^q \text{ such that } A \cap Z_{\beta_0}^q \in d_{\beta_0}^q. \end{aligned}$$

Consequently, $q \leq^0 r$ (remember 2.9).

(2) Let $\langle p_\xi : \xi < \lambda \rangle$ list (with possible repetitions) all members of G^* . For $\xi < \lambda$ let $C_\xi = \{\delta < \lambda : \delta = \sup(\delta \cap C^{p_\xi})\}$ (it is a club of λ), and for $\xi, \zeta < \lambda$ let $\varepsilon(\{\xi, \zeta\}) < \lambda$ be such that

if $p_\xi \leq^0 p_\zeta$, then

$$(\forall \alpha \in C^{p_\zeta} \setminus \varepsilon(\{\xi, \zeta\})) (\forall A \in d_\alpha^{p_\zeta}) (\exists \beta \in C^{p_\xi}) (A \cap Z_\beta^{p_\xi} \in d_\beta^{p_\xi})$$

(remember 2.9). Let

$$C^* = \{\delta < \lambda : \delta \text{ is limit and } \{p_\xi : \xi < \delta\} \text{ is } \leq^0\text{-directed}\}$$

(again, it is a club of λ). Finally, let

$$C = \{\delta \in C^* \cap \bigtriangleup_{\xi < \lambda} C_\xi : (\forall \xi, \zeta < \delta) (\varepsilon(\{\xi, \zeta\}) < \delta)\}.$$

Plainly, C is a club of λ . Now, suppose that $\delta < \gamma$ are two successive members of C . Put $Z_\delta = [\delta, \gamma)$ and let

$$I_\delta = \{A \subseteq Z_\delta : (\exists \xi < \delta) (\forall \alpha \in C^{p_\xi} \setminus \delta) (A \cap Z_\alpha^{p_\xi} \notin d_\alpha^{p_\xi})\}.$$

It easily follows from the definition of C that I_δ is a proper ideal on Z_δ , so we may pick an ultrafilter d_δ on Z_δ disjoint from I_δ . Let $q = (C, \langle Z_\delta : \delta \in C \rangle, \langle d_\delta : \delta \in C \rangle)$. Clearly $q \in \mathbb{Q}_\lambda^0$ and we will argue that q is a \leq^0 -upper bound to G^* . So let $\xi < \lambda$. Suppose that $\delta \in C \setminus (\xi + 1)$ and $A \in d_\delta$. Then $A \notin I_\delta$, so there is $\alpha \in C^{p_\xi} \setminus \delta$ such that $A \cap Z_\alpha^{p_\xi} \in d_\alpha^{p_\xi}$. Now we may use 2.9 to conclude that $p_\xi \leq^0 q$. \square

Proposition 2.12. *If $2^\lambda = \lambda^+$, then there is a $\leq_{\mathbb{Q}_\lambda^*}^*$ -increasing sequence $\bar{p} = \langle p_\varepsilon : \varepsilon < \lambda^+ \rangle \subseteq \mathbb{Q}_\lambda^0$ such that*

$$\text{fil}(\bar{p}) \stackrel{\text{def}}{=} \bigcup \{\text{fil}(p_\varepsilon) : \varepsilon < \lambda^+\}$$

is a uniform ultrafilter on λ .

Proof. Straightforward induction using 2.3(4) and the proof of 2.11(2). \square

For basic information on the ideal of meager subsets of ${}^\lambda\lambda$ and its covering number we refer the reader e.g. to Matet, Rosłanowski and Shelah [MRS05, §4]. Here we state only the definitions we will need.

- Definition 2.13.** (1) The space ${}^\lambda\lambda$ is endowed with the topology obtained by taking as basic open sets \emptyset and O_s for $s \in {}^{\lambda>}\lambda$, where $O_s = \{f \in {}^\lambda\lambda : s \subseteq f\}$.
- (2) The $(<\lambda)$ -complete ideal of subsets of ${}^\lambda\lambda$ generated by nowhere dense subsets of ${}^\lambda\lambda$ is denoted by $\mathbf{M}_{\lambda,\lambda}^\lambda$.
- (3) $\text{cov}(\mathbf{M}_{\lambda,\lambda}^\lambda)$ is the minimal size of a family $\mathcal{A} \subseteq \mathbf{M}_{\lambda,\lambda}^\lambda$ such that $\bigcup \mathcal{A} = {}^\lambda\lambda$.

Theorem 2.14. *Assume that $\lambda^{<\lambda} = \lambda$ and $\text{cov}(\mathbf{M}_{\lambda,\lambda}^\lambda) = 2^\lambda$. Then there exists a very reasonable ultrafilter on λ .*

Proof. Fix a model $N \prec \mathcal{H}(\chi)$ (for some large regular cardinal χ) such that $\|N\| = \lambda$ and ${}^{\lambda>}N \subseteq N$.

For $p \in \mathbb{Q}_\lambda^0$ let $\langle \delta_\alpha^p : \alpha < \lambda \rangle$ be the increasing enumeration of C^p and let η^p be the sequence of length λ such that

$$(\forall \alpha < \lambda) (\eta^p(\alpha) = \langle Z_{\delta_\alpha^p}^p, d_{\delta_\alpha^p}^p \rangle).$$

Next let

$$\mathcal{T}_\alpha = \{\eta^p \upharpoonright \alpha : p \in \mathbb{Q}_\lambda^0\} \cap N \quad (\text{for } \alpha < \lambda) \quad \text{and} \quad \mathcal{T} = \bigcup_{\alpha < \lambda} \mathcal{T}_\alpha.$$

Clearly \mathcal{T} is a tree isomorphic to ${}^{\lambda>}\lambda$ by an isomorphism preserving the levels (i.e., mapping \mathcal{T}_α onto ${}^\alpha\lambda$). Also, every λ -branch $\eta \in \lim(\mathcal{T})$ determines a condition $p \in \mathbb{Q}_\lambda^0$ such that $\eta = \eta^p$. Let $Q^* = \{p \in \mathbb{Q}_\lambda^0 : \eta^p \in \lim(\mathcal{T})\}$.

A family $G^* \subseteq Q^*$ is *linked* if it is $(<\omega)$ -linked with respect to the partial order \leq^0 -restricted to Q^* , that is if every finite subset of G^* has a \leq^0 -upper bound in Q^* (but the bound does not have to be in G^*). Note that if $p_0, \dots, p_n \in Q^*$ have a \leq^0 -upper bound in \mathbb{Q}_λ^0 , then they have a \leq^0 -upper bound in Q^* as well. For $p_0, \dots, p_n \in Q^*$, $\delta < \delta' < \lambda$ and an ultrafilter d on $[\delta, \delta')$ let $(\oplus)^{p_0, \dots, p_n}(\delta, \delta', d)$ mean

- $(\oplus)^{p_0, \dots, p_n}$ (a) $\delta, \delta' \in C^{p_0} \cap \dots \cap C^{p_n}$, and
 (b) if $B \in d$, $i \leq n$, then there is $\xi \in [\delta, \delta') \cap C^{p_i}$ such that $B \cap Z_\xi^{p_i} \in d_\xi^{p_i}$.

Claim 2.14.1. *If $G^* \subseteq Q^*$ is a linked family, $\|G^*\| < \text{cov}(\mathbf{M}_{\lambda,\lambda}^\lambda)$, and $A \subseteq \lambda$, then there is $p \in Q^*$ such that*

- (a) $G^* \cup \{p\}$ is linked, and
 (b) either $A \in \text{fil}(p)$ or $\lambda \setminus A \in \text{fil}(p)$.

Proof of the Claim. We will consider two cases.

CASE 1: For every $p_0, \dots, p_n \in G^*$, $n < \omega$, there is $p \in Q^*$ such that $A \in \text{fil}(p)$ and $p_0 \leq^0 p, \dots, p_n \leq^0 p$.

Note that the assumption of the present case is equivalent to

- (\otimes) for every $p_0, \dots, p_n \in G^*$, $n < \omega$, and $\alpha < \lambda$ there are $\delta < \delta' < \lambda$ and an ultrafilter $d \in N$ on $[\delta, \delta')$ such that $(\oplus)^{p_0, \dots, p_n}(\delta, \delta', d)$ holds true and $\alpha < \delta$ and $A \cap [\delta, \delta') \in d$.

We let

$$\mathcal{T}_A = \{ \eta \in \mathcal{T} : (\forall \alpha < \text{lh}(\eta)) (\forall Z, d) (\eta(\alpha) = \langle Z, d \rangle \Rightarrow A \cap Z \in d) \}.$$

Clearly, \mathcal{T}_A is a λ -branching subtree of \mathcal{T} and \mathcal{T}_A is isomorphic to ${}^{\lambda}>\lambda$. Now, for $p_0, \dots, p_n \in G^*$, $n < \omega$, and $\alpha < \lambda$ let

$$I_\alpha^A(p_0, \dots, p_n) \stackrel{\text{def}}{=} \{ \eta \in \text{lim}(\mathcal{T}_A) : (\exists \beta > \alpha) (\exists \delta, \delta', d) ((\oplus)^{p_0, \dots, p_n}(\delta, \delta', d) \ \& \ \eta(\beta) = \langle [\delta, \delta'), d \rangle) \}.$$

It should be clear that $I_\alpha^A(p_0, \dots, p_n)$ is an open dense subset of $\text{lim}(\mathcal{T}_A)$ (remember (\otimes)). Therefore (as $\|G^*\| < \text{cov}(\mathbf{M}_{\lambda, \lambda}^\lambda)$) we know that

$$\bigcap \{ I_\alpha^A(p_0, \dots, p_n) : n < \omega \ \& \ p_0, \dots, p_n \in G^* \ \& \ \alpha < \lambda \} \neq \emptyset$$

and we may choose η from the set on the left-hand side above. Let $p \in Q^*$ be such that $\eta = \eta^p$. Since $\eta \in \text{lim}(\mathcal{T}_A)$ we know that $A \in \text{fil}(p)$. Also, for every $p_0, \dots, p_n \in G^*$ we have $\eta \in \bigcap_{\alpha < \lambda} I_\alpha^A(p_0, \dots, p_n)$ and hence

$$\| \{ \delta \in C^p : \text{if } \delta' = \min(C^p \setminus (\delta + 1)) \text{ then } (\oplus)^{p_0, \dots, p_n}(\delta, \delta', d_\delta^p) \} \| = \lambda.$$

So one may easily construct $p^* \in Q^*$ which is \leq^0 -stronger than p, p_0, \dots, p_n (remember 2.9). Thus we have justified that $G^* \cup \{p\}$ is linked.

CASE 2: There are $p_0, \dots, p_n \in G^*$, $n < \omega$, such that

$$\text{if } p \in Q^* \text{ is } \leq^0\text{-stronger than } p_0, \dots, p_n, \text{ then } A \notin \text{fil}(p).$$

It follows from the proof of 2.3(4) that then

$$\text{for every } q_0, \dots, q_m \in G^*, \ m < \omega, \text{ there is } q \in Q^* \text{ such that} \\ \lambda \setminus A \in \text{fil}(q) \text{ and } q_0 \leq^0 q, \dots, q_m \leq^0 q$$

(remember G^* is linked and that bounded subsets of λ are in N). Thus we may repeat the arguments of Case 1 for $\lambda \setminus A$ and we find $p \in Q^*$ such that $G^* \cup \{p\}$ is linked and $\lambda \setminus A \in \text{fil}(p)$. \square

Claim 2.14.2. *If $G^* \subseteq Q^*$ is linked, $\|G^*\| < \text{cov}(\mathbf{M}_{\lambda, \lambda}^\lambda)$ and $p_0, p_1 \in G^*$, then there is $p \in Q^*$ such that*

- (a) $G^* \cup \{p\}$ is linked, and
- (b) $p_0 \leq^0 p$ and $p_1 \leq^0 p$.

Proof of the Claim. Let $p_0, p_1 \in G^*$. Note that

- (\odot) for every $p_2, \dots, p_n \in G^*$, $2 \leq n < \omega$, and $\alpha < \lambda$ there are $\delta < \delta' < \lambda$ and an ultrafilter $d \in N$ on $[\delta, \delta']$ such that $(\oplus)^{p_0, p_1, p_2, \dots, p_n}(\delta, \delta', d)$ holds true and $\alpha < \delta$.

We let \mathcal{T}^{p_0, p_1} be the set

$$\{\eta \in \mathcal{T} : (\forall \alpha < \text{lh}(\eta)) (\forall \delta, \delta', d) (\eta(\alpha) = \langle [\delta, \delta'], d \rangle \Rightarrow (\oplus)^{p_0, p_1}(\delta, \delta', d))\}$$

and we note that \mathcal{T}^{p_0, p_1} is a λ -branching subtree of \mathcal{T} isomorphic to ${}^{\lambda}>\lambda$.

For $p_2, \dots, p_n \in G^*$, $2 \leq n < \omega$, and $\alpha < \lambda$ we let

$$I_{\alpha}^{p_0, p_1}(p_2, \dots, p_n) \stackrel{\text{def}}{=} \{\eta \in \text{lim}(\mathcal{T}^{p_0, p_1}) : (\exists \beta > \alpha) (\exists \delta, \delta', d) ((\oplus)^{p_2, \dots, p_n}(\delta, \delta', d) \ \& \ \eta(\beta) = \langle [\delta, \delta'], d \rangle)\}.$$

Then $I_{\alpha}^{p_0, p_1}(p_2, \dots, p_n)$ is an open dense subset of $\text{lim}(\mathcal{T}^{p_0, p_1})$ (remember (\odot)). Since $\|G^*\| < \text{cov}(\mathbf{M}_{\lambda, \lambda}^{\lambda})$, we may choose $p \in Q^*$ such that

$$\eta^p \in \bigcap \{I_{\alpha}^{p_0, p_1}(p_2, \dots, p_n) : 2 \leq n < \omega \ \& \ p_2, \dots, p_n \in G^* \ \& \ \alpha < \lambda\} \neq \emptyset.$$

Like in the proof of 2.14.1 we argue that $G^* \cup \{p\}$ is linked. Since $\eta^p \in \text{lim}(\mathcal{T}^{p_0, p_1})$ we easily see that p is \leq^0 -stronger than both p_0 and p_1 . \square

Claim 2.14.3. *If $G^* \subseteq Q^*$ is a linked family, $\|G^*\| < \text{cov}(\mathbf{M}_{\lambda, \lambda}^{\lambda})$, $\xi \leq \lambda$ is a limit ordinal and a sequence $\langle p_{\zeta} : \zeta < \xi \rangle \subseteq G^*$ is \leq^0 -increasing, then there is $p \in Q^*$ such that*

- (a) $G^* \cup \{p\}$ is linked, and
- (b) $(\forall \zeta < \xi)(p_{\zeta} \leq^0 p)$.

Proof of the Claim. First let us consider the case when $\xi < \lambda$. Suppose that a sequence $\bar{p} = \langle p_{\zeta} : \zeta < \xi \rangle \subseteq G^*$ is \leq^0 -increasing and let

$$\mathcal{T}_{\bar{p}} = \{\eta \in \mathcal{T} : (\forall \alpha < \text{lh}(\eta)) (\forall \delta, \delta', d) (\eta(\alpha) = \langle [\delta, \delta'], d \rangle \Rightarrow (\forall \zeta < \xi) (\oplus)^{p_{\zeta}}(\delta, \delta', d))\}.$$

By arguments similar to that of 2.3(3) we verify that $\mathcal{T}_{\bar{p}}$ is a λ -branching subtree of \mathcal{T} and it is isomorphic to ${}^{\lambda}>\lambda$. Like in the previous claims, for $p'_0, \dots, p'_n \in G^*$, $n < \omega$, and $\alpha < \lambda$ we put

$$I_{\alpha}^{\bar{p}}(p'_0, \dots, p'_n) \stackrel{\text{def}}{=} \{\eta \in \text{lim}(\mathcal{T}_{\bar{p}}) : (\exists \beta > \alpha) (\exists \delta, \delta', d) ((\oplus)^{p'_0, \dots, p'_n}(\delta, \delta', d) \ \& \ \eta(\beta) = \langle [\delta, \delta'], d \rangle)\}.$$

Then each $I_{\alpha}^{\bar{p}}(p'_0, \dots, p'_n)$ is an open dense subset of $\text{lim}(\mathcal{T}_{\bar{p}})$. [Why? Let $\eta \in \mathcal{T}_{\bar{p}}$. We may assume that for each $\varepsilon < \zeta < \xi$ and $\beta \in C^{p_{\zeta}} \setminus \text{lh}(\eta)$ and $A \in d_{\beta}^{p_{\zeta}}$ there is $\gamma \in C^{p_{\varepsilon}}$ such that $A \cap Z_{\gamma}^{p_{\varepsilon}} \in d_{\gamma}^{p_{\varepsilon}}$. We also may demand that

$$\delta_0 \stackrel{\text{def}}{=} \sup(\delta' < \lambda : (\exists \alpha < \text{lh}(\eta)) (\exists \delta, d) (\eta(\alpha) = \langle [\delta, \delta'], d \rangle) \in \bigcap_{\varepsilon < \xi} C^{p_{\varepsilon}} \cap \bigcap_{i \leq n} C^{p'_i}).$$

Choose inductively a sequence $\langle \delta_{\zeta}, d_{\zeta} : \zeta < \xi \rangle$ so that

- (a) $\langle \delta_\zeta : \zeta < \xi \rangle$ is an increasing continuous sequence of ordinals below λ ,
- (b) $d_\zeta \in N$ is an ultrafilter on $[\delta_\zeta, \delta_{\zeta+1})$, $\delta_{\zeta+1} \in \bigcap_{\varepsilon < \xi} C^{p_\varepsilon}$,
- (c) $(\bigoplus)^{p'_0, \dots, p'_n, p_\zeta}(\delta_\zeta, \delta_{\zeta+1}, d_\zeta)$ holds true (for each $\zeta < \xi$).

Let $\delta_\xi = \sup(\delta_\zeta : \zeta < \xi)$ and let $e \in N$ be an ultrafilter on ξ containing all co-bounded subsets of ξ . Put $d = \bigoplus_{\zeta < \xi}^e d_\zeta$ — it is an ultrafilter on $[\delta_0, \delta_\xi)$,

$d \in N$ and $(\bigoplus)^{p'_0, \dots, p'_n, p_\zeta}(\delta_0, \delta_\xi, d)$ holds true for each $\zeta < \xi$. Consequently $\eta \cup \{(\text{lh}(\eta), \langle [\delta_0, \delta_\xi), d \rangle)\} \in \mathcal{T}_{\bar{p}}$ and every member of $\lim(\mathcal{T}_{\bar{p}})$ extending it belongs to $I_{\alpha}^{\bar{p}}(p'_0, \dots, p'_n)$.

Thus we may pick $p \in Q^*$ such that

$$\eta^p \in \bigcap \{I_{\alpha}^{\bar{p}}(p'_0, \dots, p'_n) : n < \omega \ \& \ p'_0, \dots, p'_n \in G^* \ \& \ \alpha < \lambda\}.$$

Since $\eta^p \in \lim(\mathcal{T}_{\bar{p}})$ we easily see that $p_\zeta \leq^0 p$ for all $\zeta < \xi$, and like in the proof of 2.14.1 we argue that $G^* \cup \{p\}$ is linked.

If $\xi = \lambda$ and $\bar{p} = \langle p_\zeta : \zeta < \lambda \rangle$ is \leq^0 -increasing, then we proceed in a similar manner except that we work in the tree

$$\mathcal{T}_{\bar{p}}^* = \{\eta \in \mathcal{T} : (\forall \alpha < \text{lh}(\eta)) (\forall \delta, \delta', d) (\eta(\alpha) = \langle [\delta, \delta'), d \rangle \Rightarrow (\forall \zeta < \alpha) (\bigoplus)^{p_\zeta}(\delta, \delta', d))\}.$$

□

Claim 2.14.4. *Assume that $G^* \subseteq Q^*$ is a linked family, $\|G^*\| < \text{cov}(\mathbf{M}_{\lambda, \lambda}^\lambda)$, $C \subseteq \lambda$ is a club and $\langle \delta_\xi : \xi < \lambda \rangle$ is the increasing enumeration of C . Then there is $p \in Q^*$ and a club C^* of λ such that*

- (a) $G^* \cup \{p\}$ is linked, and
- (b) $\bigcup \{[\delta_{\xi+1}, \delta_\xi) : \xi < \zeta \text{ are successive members of } C^*\} \in \text{fil}(p)$.

Proof of the Claim. Let

$$\mathcal{T}_C = \{\eta \in \mathcal{T} : \text{for each } \alpha < \text{lh}(\eta) \text{ such that } \alpha = \delta_\alpha \text{ and for every } \alpha', d \\ \eta(\alpha) = \langle [\alpha, \alpha'), d \rangle \Rightarrow \delta_{\alpha+1} < \alpha' \ \& \ [\delta_\alpha, \delta_{\alpha+1}) \notin d\}.$$

One easily verifies that \mathcal{T}_C is a λ -branching subtree of \mathcal{T} which is isomorphic to ${}^\lambda \lambda$. Like before, for $p_0, \dots, p_n \in G^*$, $n < \omega$, and $\alpha < \lambda$ we put

$$I_{\alpha}^C(p_0, \dots, p_n) \stackrel{\text{def}}{=} \{\eta \in \lim(\mathcal{T}_C) : (\exists \beta > \alpha) (\exists \delta, \delta', d) ((\bigoplus)^{p_0, \dots, p_n}(\delta, \delta', d) \ \& \ \eta(\beta) = \langle [\delta, \delta'), d \rangle)\}.$$

Each $I_{\alpha}^C(p_0, \dots, p_n)$ is an open dense subset of $\lim(\mathcal{T}_C)$ and hence there is $p \in Q^*$ such that

$$\eta^p \in \bigcap \{I_{\alpha}^C(p_0, \dots, p_n) : n < \omega \ \& \ p_0, \dots, p_n \in G^* \ \& \ \alpha < \lambda\}.$$

Like in the proof of 2.14.1 we argue that $G^* \cup \{p\}$ is linked. Put

$$C^* = \{\alpha < \lambda : \alpha = \delta_\alpha \text{ is limit \& } (\exists \alpha', d)(\eta^p(\alpha) = \langle [\alpha, \alpha'], d \rangle)\}$$

and note that C^* is a club of λ . Note that if $\alpha \in C^*$ and $\eta^p(\alpha) = \langle [\alpha, \alpha'], d \rangle$, then $\delta_{\alpha+1} < \alpha'$ and $[\delta_\alpha, \delta_{\alpha+1}] \notin d$. Consequently,

$$\bigcup \{[\delta_{\alpha+1}, \delta_\beta] : \alpha < \beta \text{ are successive members of } C^*\} \in \text{fil}(p).$$

□

To prove the theorem we construct inductively a sequence $\langle q_\zeta : \zeta < 2^\lambda \rangle$ of elements of Q^* such that

- for each $\xi < 2^\lambda$ the family $\{q_\zeta : \zeta < \xi\}$ is linked,
- for each $A \subseteq \lambda$ there is $\zeta < 2^\lambda$ such that either $A \in \text{fil}(q_\zeta)$ or $\lambda \setminus A \in \text{fil}(q_\zeta)$,
- for each $\zeta < \xi < 2^\lambda$ there is $\alpha < 2^\lambda$ such that $q_\zeta \leq^0 q_\alpha$ and $q_\xi \leq^0 q_\alpha$,
- if $\xi \leq \lambda$ and $\langle p_\zeta : \zeta < \xi \rangle$ is a \leq^0 -increasing sequence of elements of $\{q_\zeta : \zeta < 2^\lambda\}$, then there is $\alpha < 2^\lambda$ such that q_α is a \leq^0 -upper bound to all p_ζ 's,
- if a sequence $\langle \delta_\xi : \xi < \lambda \rangle \subseteq \lambda$ is increasing continuous, then for some $\zeta < 2^\lambda$ and a club C^* of λ we have

$$\bigcup \{[\delta_{\xi+1}, \delta_{\xi'}] : \xi < \xi' \text{ are successive members of } C^*\} \in \text{fil}(q_\zeta).$$

The construction is a straightforward application of a suitable bookkeeping device and Claims 2.14.1–2.14.4. After it is carried out put $G^* = \{q_\zeta : \zeta < 2^\lambda\}$ and note that $\text{fil}(G^*)$ is a very reasonable ultrafilter on λ . □

Let us finish this section with an observation showing that the assumption $\lambda^{<\lambda} = \lambda$ in Theorem 2.14 is very natural in the given context.

Proposition 2.15. *Assume $\theta < \lambda = \text{cf}(\lambda) < 2^\theta$. Then $\text{cov}(\mathbf{M}_{\lambda, \lambda}^\lambda) = \lambda^+$.*

Proof. Let $\langle \nu_\xi : \xi < \lambda^+ \rangle$ be a sequence of distinct functions from θ to 2. Let $\langle \delta_\alpha : \alpha < \lambda \rangle \subseteq \lambda$ be an increasing continuous sequence such that $\delta_0 = 0$, $\delta_{\alpha+1} = \delta_\alpha + \theta$ (for $\alpha < \lambda$). Now, for $\xi < \lambda^+$ we define

$$F_\xi = \{\eta \in {}^\lambda \lambda : (\forall \alpha < \lambda)(\exists i < \theta)(\eta(\delta_\alpha + i) \neq \nu_\xi(i))\}.$$

Plainly, each F_ξ is a closed nowhere dense subset of ${}^\lambda \lambda$. We claim that

$\bigcup_{\xi < \lambda^+} F_\xi = {}^\lambda \lambda$. To this end suppose that $\eta \in {}^\lambda \lambda$ and consider the restrictions $\eta \upharpoonright [\delta_\alpha, \delta_{\alpha+1})$ for $\alpha < \lambda$. These restrictions determine λ functions from θ to 2, so we may find $\xi < \lambda^+$ such that ν_ξ is distinct from all these functions, i.e., $(\forall \alpha < \lambda)(\exists i < \theta)(\eta(\delta_\alpha + i) \neq \nu_\xi(i))$. Then $\eta \in F_\xi$. □

3. $\text{fil}(G^*)$ AND DOMINATING FAMILIES

In this section we show that families $G^* \subseteq \mathbb{Q}_\lambda^0$ generating reasonable ultrafilters cannot be too small.

Theorem 3.1. *For $p \in \mathbb{Q}_\lambda^0$ let $f_p \in {}^\lambda \lambda$ be such that*

$$(\forall \alpha < \lambda) (f_p(\alpha) \in C^p \ \& \ \text{otp}(C^p \cap f_p(\alpha)) = \omega \cdot \alpha + \omega).$$

- (1) *Suppose that $G_0^* \subseteq \mathbb{Q}_\lambda^0$ is $(<\aleph_1)$ -directed (with respect to \leq^0) and $\text{fil}(G_0^*)$ is a weakly reasonable ultrafilter. Then $\mathcal{F}_0 = \{f_p : p \in G_0^*\}$ is a dominating family in ${}^\lambda \lambda$.*
- (2) *Suppose that $G_1^* \subseteq \mathbb{Q}_\lambda^0$ is directed (with respect to \leq^0) and $\text{fil}(G_1^*)$ is a weakly reasonable ultrafilter on λ . Then $\mathcal{F}_1 = \{f_p : p \in G_1^*\}$ is a club-dominating family in ${}^\lambda \lambda$.*

Proof. (1) First note that if $p, q \in G_0^*$, $p \leq^0 q$, then for some $\varepsilon < \lambda$, if $\alpha < \beta < \gamma$ are successive members of $C^q \setminus \varepsilon$, then $(\alpha, \gamma) \cap C^p \neq \emptyset$. Thus $p \leq^0 q$ implies that for all sufficiently large $\alpha < \lambda$ we have $f_p(\alpha) \leq f_q(\alpha)$. Consequently the family \mathcal{F}_0 is $(<\aleph_1)$ -directed (with respect to \leq^*).

Suppose towards contradiction that \mathcal{F}_0 is not a dominating family. Then we may choose an increasing continuous sequence $\bar{\alpha}^0 = \langle \alpha_\xi^0 : \xi < \lambda \rangle$ such that

$$(\forall p \in G_0^*) (\exists^\lambda \varepsilon < \lambda) (f_p(\alpha_\varepsilon^0) < \alpha_{\varepsilon+1}^0).$$

Now, by induction on $n < \omega$, choose increasing continuous sequences $\bar{\alpha}^n = \langle \alpha_\xi^n : \xi < \lambda \rangle$ so that letting $C_n = \{\alpha_\xi^n : \xi < \lambda\}$ we have

- (i) $\bar{\alpha}^0$ is the one chosen earlier,
- (ii) $C_{n+1} \subseteq \{\alpha_\varepsilon^n : \varepsilon = \alpha_\varepsilon^n \text{ is a limit ordinal}\}$,
- (iii) $\bigcup \{[\alpha_\varepsilon^n, \alpha_{\varepsilon+1}^n] : \varepsilon \in C_{n+1}\} \notin \text{fil}(G_0^*)$.

It should be clear that the construction of $\bar{\alpha}^n$'s is possible (remember that $\text{fil}(G_0^*)$ is a weakly reasonable ultrafilter; use 1.5). Let $C_\omega = \bigcap_{n < \omega} C_n$ and

let $\langle \alpha_\xi^\omega : \xi < \lambda \rangle$ be the increasing enumeration of C_ω . It follows from (ii) that for every $\varepsilon < \lambda$ the sequence $\langle \alpha_{\varepsilon+1}^n : n < \omega \rangle$ is strictly increasing and $\sup(\alpha_{\varepsilon+1}^n : n < \omega) \in C_\omega$, and if $\xi = \alpha_\varepsilon^\omega$, then $\sup(\alpha_{\xi+1}^n : n < \omega) = \alpha_{\varepsilon+1}^\omega$. It follows from (iii) that for every $n < \omega$

$$A_n \stackrel{\text{def}}{=} \bigcup \{[\alpha_\varepsilon^\omega, \alpha_{\xi+1}^n] : \varepsilon < \lambda \ \& \ \xi = \alpha_\varepsilon^\omega\} \notin \text{fil}(G_0^*).$$

Fix $p \in G_0^*$ for a moment. By the choice of $\bar{\alpha}^0$ we know that the set $\{\xi < \lambda : f_p(\alpha_\xi^0) < \alpha_{\xi+1}^0\}$ is unbounded in λ , and hence also the set $\{\varepsilon < \lambda : f_p(\alpha_\varepsilon^\omega) < \alpha_{\varepsilon+1}^\omega\}$ is unbounded in λ . Therefore for some $n < \omega$ we have

$$\|\{\varepsilon < \lambda : \xi = \alpha_\varepsilon^\omega \Rightarrow f_p(\alpha_\varepsilon^\omega) < \alpha_{\xi+1}^n\}\| = \lambda;$$

let $n(p)$ be the first such $n < \omega$.

Note that if $p \leq^0 q$ are from G_0^* , then $n(p) \leq n(q)$ (as $f_p \leq^* f_q$). Consequently, since G_0^* is $(<\aleph_1)$ -directed, there is $n^* < \omega$ such that $(\forall p \in G_0^*)(n(p) \leq n^*)$. Look at the set A_{n^*} : for every $p \in G_0^*$ there are λ many $\varepsilon < \lambda$ such that $\alpha_\varepsilon^\omega < f_p(\alpha_\varepsilon^\omega) < \alpha_{\xi+1}^{n^*}$, where $\xi = \alpha_\varepsilon^\omega$, and so (by the definition of f_p) we get $A_{n^*} \in (\text{fil}(p))^+$. Since $\text{fil}(G_0^*)$ is an ultrafilter we get an immediate contradiction with $A_{n^*} \notin \text{fil}(G_0^*)$.

(2) Suppose towards contradiction that \mathcal{F}_1 is not club-dominating in ${}^\lambda\lambda$. Then we may find an increasing function $h \in {}^\lambda\lambda$ such that

$$(\forall p \in G_1^*)(\{\varepsilon < \lambda : f_p(\varepsilon) < h(\varepsilon)\} \text{ is stationary in } \lambda).$$

Pick an increasing continuous sequence $\langle \delta_\xi : \xi < \lambda \rangle \subseteq \lambda$ such that $(\forall \xi < \lambda)(h(\delta_\xi) < \delta_{\xi+1})$. Since $\text{fil}(G_1^*)$ is weakly reasonable, we may use 1.5 to pick a club C of λ such that $C \subseteq \{\xi < \lambda : \delta_\xi = \xi \text{ is a limit ordinal}\}$ and

$$\bigcup \{[\delta_\xi, \delta_{\xi+1}) : \xi \in C\} \notin \text{fil}(G_1^*).$$

Since $\text{fil}(G_1^*)$ is an ultrafilter, for some $p \in G_1^*$ we have

$$\lambda \setminus \bigcup \{[\delta_\xi, \delta_{\xi+1}) : \xi \in C\} \in \text{fil}(p).$$

However, by the choice of h , the set $\{\xi < \lambda : \delta_\xi = \xi \in C \ \& \ f_p(\xi) < h(\xi) < \delta_{\xi+1}\}$ is stationary (so of size λ), and we get an immediate contradiction with the definition of f_p . \square

Corollary 3.2. (1) *If $G_0^* \subseteq \mathbb{Q}_\lambda^0$ is $(<\aleph_1)$ -directed (with respect to \leq^0) and $\text{fil}(G^*)$ is a weakly reasonable ultrafilter, then $\|G_0^*\| \geq \mathfrak{d}_\lambda$.*

(2) *If $G_1^* \subseteq \mathbb{Q}_\lambda^0$ is directed (with respect to \leq^0) and $\text{fil}(G_1^*)$ is a weakly reasonable ultrafilter on λ , then $\|G_1^*\| \geq \mathfrak{d}_{\text{cl}}(\lambda)$.*

Proposition 3.3. *Suppose that $G_0^* \subseteq \mathbb{Q}_\lambda^0$ is $(<\lambda)$ -directed (with respect to $\leq_{\mathbb{Q}_\lambda^0}$) and $\text{fil}(G^*)$ is an ultrafilter. For $p \in G_0^*$ let $f_p \in {}^\lambda\lambda$ be defined as in 3.1. If $\mathcal{F}_0 = \{f_p : p \in G_0^*\}$ is not a dominating family in ${}^\lambda\lambda$, then λ is measurable.*

Proof. Similarly as in the proof of 3.1(1), we note that \mathcal{F}_0 is $(<\lambda)$ -directed (with respect to \leq^*). Assume \mathcal{F}_0 is not dominating family. Then may choose an increasing continuous sequence $\langle \alpha_\xi : \xi < \lambda \rangle$ such that

$$(\forall p \in G_0^*)(\exists^\lambda \varepsilon < \lambda)(f_p(\alpha_\varepsilon) < \alpha_{\varepsilon+1}).$$

Let

$$\mathcal{U} = \left\{ A \subseteq \lambda : (\exists p \in G_0^*)(\exists \delta < \lambda)(\forall \varepsilon > \delta)(f_p(\alpha_\varepsilon) < \alpha_{\varepsilon+1} \Rightarrow \varepsilon \in A) \right\}.$$

We are going to show that \mathcal{U} is a λ -complete uniform ultrafilter on λ . It should be clear that \mathcal{U} includes all co-bounded subsets of λ and that it is a

λ -complete filter (remember that \mathcal{F}_0 is $<\lambda$ -directed). To show that it is an ultrafilter suppose that $A \subseteq \lambda$ and let

$$B = \bigcup \{[\alpha_\varepsilon, \alpha_{\varepsilon+1}) : \varepsilon \in A\} \subseteq \lambda.$$

Since $\text{fil}(G_0^*)$ is an ultrafilter, then either $B \in \text{fil}(G_0^*)$ or $\lambda \setminus B \in \text{fil}(G_0^*)$. Suppose that the former happens, so we may choose $p \in G_0^*$ such that $B \in \text{fil}(p)$. Then for some $\delta < \lambda$ we have

$$(\forall \beta \in C^p \setminus \delta)(B \cap Z_\beta^p \in d_\beta^p).$$

Now, if $\varepsilon > \delta$ and $f_p(\alpha_\varepsilon) < \alpha_{\varepsilon+1}$, then $[\alpha_\varepsilon, \alpha_{\varepsilon+1}) \cap B \neq \emptyset$ and thus $[\alpha_\varepsilon, \alpha_{\varepsilon+1}) \subseteq B$, so $\varepsilon \in A$. Consequently $A \in \mathcal{U}$ (as witnessed by p, δ). In the same manner one shows that if $\lambda \setminus B \in \text{fil}(G_0^*)$, then $\lambda \setminus A \in \mathcal{U}$. \square

4. OPEN PROBLEMS AND FURTHER INVESTIGATIONS

It may well be that our forcing techniques for uncountable λ are still not strong enough to carry out the arguments parallel to the consistency results for ultrafilters on ω . However, we feel that the recent progress in the theory of forcing iterated with uncountable supports (as exemplified by [She03], Roslanowski and Shelah [RS01], [RS07], [RS06] and Eisworth [Eis03]) may prove to be useful in developing iterated forcing for “killing” and/or “preserving” some subfamilies of the class of reasonable ultrafilters. In particular, in Roslanowski and Shelah [RS11] we continue the research of the present paper and we introduce *super reasonable ultrafilters* which are stronger than very reasonable ultrafilters. We show that for inaccessible λ it is consistent that there are such ultrafilters determined by generating systems of size less than 2^λ , and we also prove a result on preserving them in λ -support iterations. We also show that consistently there are no super reasonable ultrafilters. These results may be interpreted as some progress towards generalizing (a), (b) and (c) from the Introduction. However, several other natural problems remain untouched. One of the main questions we are interested in is

Problem 4.1. Let λ be a regular uncountable cardinal. Is it provable in ZFC that there exist reasonable ultrafilters on λ ? Very reasonable? (See 2.5(4,5).)

Problem 4.2. Is it consistent that there exists a very reasonable ultrafilter D on λ such that for every very reasonable ultrafilter D' on λ for some function $f \in \mathcal{F}_\lambda$ we have $D/f = D'/f$?

Since in the present paper we deal with dividing by $f \in \mathcal{F}_\lambda$, and the normal ultrafilters are fixed points for this operation, the natural question is:

Problem 4.3. Is it consistent that for every $D \in \text{uuf}$ there is $f \in \mathcal{F}_\lambda$ such that either D/f is normal or D/f is reasonable (or even very reasonable)?

We may also re-interpret our aim as follows.

Definition 4.4. (1) Let $\text{UE}_{\lambda,\mu}^*$ be the family of all $(<\mu)$ -directed (with respect to \leq^0) subsets G^* of \mathbb{Q}_λ^0 such that $\text{fil}(G^*)$ is a proper ultrafilter on λ ;

(2) $\text{UF}_{\lambda,\mu}^* = \{\text{fil}(G^*) : G^* \in \text{UE}_{\lambda,\mu}^*\}$; $\text{UF}_\lambda^* = \text{UF}_{\lambda,\lambda^+}^*$ and $\text{UF}_\lambda = \text{UF}_{\lambda,\aleph_0}^*$.

Aim 4.5. Investigate UF_λ^* , UF_λ ; in particular can we have: any two of them have common quotients.

We expect that the forcing theorems needed for further research will be similar to [She03] and more so to [RS01, RS07, RS06], in some respects, and for others to [She].

Let us also note that combinatorial aspects of generating ultrafilters by means similar to generating systems of Definition 2.5 will be studied in Rosłanowski and Shelah [RS08].

REFERENCES

- [BS87] Andreas R. Blass and Saharon Shelah, *There may be simple P_{\aleph_1} - and P_{\aleph_2} -points and the Rudin-Keisler ordering may be downward directed*, Ann. Pure Appl. Logic **33** (1987), no. 3, 213–243. MR 879489
- [CS95] James Cummings and Saharon Shelah, *Cardinal invariants above the continuum*, Ann. Pure Appl. Logic **75** (1995), no. 3, 251–268, arXiv: math/9509228. MR 1355135
- [Eis03] Todd Eisworth, *On iterated forcing for successors of regular cardinals*, Fundamenta Mathematicae **179** (2003), 249–266.
- [Git81] Moti Gitik, *On nonminimal p -points over a measurable cardinal*, Annals of Mathematical Logic **20** (1981), 269–288.
- [Jec03] Thomas Jech, *Set theory*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2003, The third millennium edition, revised and expanded.
- [MRS05] Pierre Matet, Andrzej Rosłanowski, and Saharon Shelah, *Cofinality of the non-stationary ideal*, Trans. Amer. Math. Soc. **357** (2005), no. 12, 4813–4837, arXiv: math/0210087. MR 2165389
- [RS99] Andrzej Rosłanowski and Saharon Shelah, *Norms on possibilities. I. Forcing with trees and creatures*, Mem. Amer. Math. Soc. **141** (1999), no. 671, xii+167, arXiv: math/9807172. MR 1613600
- [RS01] ———, *Iteration of λ -complete forcing notions not collapsing λ^+* , Int. J. Math. Math. Sci. **28** (2001), no. 2, 63–82, arXiv: math/9906024. MR 1885053
- [RS06] ———, *Reasonably complete forcing notions*, Set theory: recent trends and applications, Quad. Mat., vol. 17, Dept. Math., Seconda Univ. Napoli, Caserta, 2006, arXiv: math/0508272, pp. 195–239. MR 2374767
- [RS07] ———, *Sheva-Sheva-Sheva: large creatures*, Israel J. Math. **159** (2007), 109–174, arXiv: math/0210205. MR 2342475
- [RS08] ———, *Generating ultrafilters in a reasonable way*, MLQ Math. Log. Q. **54** (2008), no. 2, 202–220, arXiv: math/0607218. MR 2402629

- [RS11] ———, *Reasonable ultrafilters, again*, Notre Dame J. Form. Log. **52** (2011), no. 2, 113–147, arXiv: math/0605067. MR 2794647
- [She] Saharon Shelah, *Many forcing axioms for all regular uncountable cardinals*, Preprint, arXiv: 1306.5399.
- [She98] ———, *Proper and improper forcing*, second ed., Perspectives in Mathematical Logic, Springer-Verlag, Berlin, 1998. MR 1623206
- [She03] ———, *Not collapsing cardinals $\leq \kappa$ in $(< \kappa)$ -support iterations*, Israel J. Math. **136** (2003), 29–115, arXiv: math/9707225. MR 1998104

EINSTEIN INSTITUTE OF MATHEMATICS, EDMOND J. SAFRA CAMPUS, GIVAT RAM,
THE HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM, 91904, ISRAEL, AND DE-
PARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW BRUNSWICK, NJ 08854,
USA

Email address: shelah@math.huji.ac.il

URL: <http://shelah.logic.at>