

# Examples of Non-locality

John T. Baldwin\*

Department of Mathematics, Statistics and Computer Science  
University of Illinois at Chicago

Saharon Shelah†

Department of Mathematics  
Hebrew University of Jerusalem  
Rutgers University

October 5, 2020

## Abstract

We use  $\kappa$ -free but not Whitehead Abelian groups to construct Abstract Elementary Classes (AEC) which satisfy the amalgamation property but fail various conditions on the locality of Galois-types. We introduce the notion that an AEC admits intersections. We conclude that for AEC which admit intersections, the amalgamation property can have no positive effect on locality: there is a transformation of AEC's which preserves non-locality but takes any AEC which admits intersections to one with amalgamation. More specifically we have: Theorem 5.3. There is an AEC with amalgamation which is not  $(\aleph_0, \aleph_1)$ -tame but is  $(2^{\aleph_0}, \infty)$ -tame; Theorem 3.3. It is consistent with ZFC that there is an AEC with amalgamation which is not  $(\leq \aleph_2, \leq \aleph_2)$ -compact.

A primary object of study in first order model theory is a *syntactic* type: the type of  $a$  over  $B$  in a model  $N$  is the collection of formulas  $\phi(x, \mathbf{b})$  which are true of  $a$  in  $N$ . Usually the  $N$  is suppressed because a preliminary construction has established a universal domain for the investigation. In such a *homogeneous* universal domain one can identify the type of  $a$  over  $B$  as the orbit of  $a$  under automorphisms which fix  $B$  pointwise.

An abstract elementary class is a pair  $(\mathbf{K}, \prec_{\mathbf{K}})$ , a collection of structures of a fixed similarity type and a partial order on  $\mathbf{K}$  which refines substructure and satisfies natural axioms which are enumerated in many places such as ([She87, Bal00, She99, Gro02, GV06b]). In this case, ‘Galois’ types (introduced in [She87] and named in [Gro02]) are defined as equivalence classes of triples  $(M, a, N)$  where  $a \in N - M$  under the equivalence relation generated by  $(M_1, a_1, N_1) \sim$

---

\*Partially supported by NSF-0500841.

†This is paper 862 in Shelah’s bibliography. The authors thank the Binational Science Foundation for partial support of this research.

$(M_2, a_2, N_2)$  if  $M_1 = M_2$  and there is an amalgam of  $N_1$  and  $N_2$  over  $M_1$  where  $a_1$  and  $a_2$  have the same image. If  $\mathbf{K}$  has the amalgamation property the equivalence classes, i.e. the Galois types, of this equivalence relation can again be identified as orbits of automorphisms of a universal domain which fix the domain of the type. The notions and definitions which appear in this paper stem from a long series of papers by Shelah ([She87, She99, She01] etc.) They occur in the form used here in [Bal00]. Grossberg and Vandieren [GV06b] isolated the notion of tame as a fruitful object of study. Recent work by such authors Grossberg, Kolesnikov, VanDieren, Villaveces [BKV06, GV06b, GV06a, GV06c, GK] either assume or derive tameness. In particular, a number of results on the stability spectrum and transfer of categoricity have been proved for tame AEC.

Unless we specifically add hypotheses,  $\mathbf{K}$  denotes an arbitrary AEC. The Löwenheim-Skolem number of an AEC  $\mathbf{K}$  is denoted  $\text{LS}(\mathbf{K})$ . We introduce in this paper two new notions: admitting intersections and model completeness. As explained in Section 1, in an AEC which admits intersections the notion of Galois type is better behaved. Model completeness is the natural analog of the first order notion in this context.

Syntactic types have certain natural locality properties. Any increasing chain of types has at most one upper bound; two distinct types differ on a finite set; an increasing chain of types has a realization. The translations of these conditions to Galois types do not hold in general. But there have been few specific examples of their failure. In the first section, we first give precise names to these three notions (in order): locality, tameness, and compactness and attach certain cardinal parameters to them. Precise statements of the results depend on these definitions and occur with the proofs. But vaguely speaking, in Section 2 we show there is an AEC with the amalgamation property which is not  $\aleph_0$ -tame and does not attain tameness at any small cardinal. In Section 3 we find a  $\mathbf{K}$  which is not compact at one of  $\aleph_1, \aleph_2$ . These results were proved before Baldwin and Kolesnikov [BK] generated a number of examples of tameness failing at different low cardinalities. These two papers seem to represent different kinds of failures of tameness. Here, we code non-continuity of ‘freeness’. The key to [BK] is requiring a number of parameters before structure imposed by affine maps becomes apparent; this appearance manifests itself in failures of  $k$ -goodness for larger  $k$  and failure of tameness at larger  $\aleph_k$ . In Section 4 we introduce a general construction which shows that one can transform a failure of locality in an AEC which admits intersections to a failure in an AEC with amalgamation. And in Section 5, we combine Sections 2 and 4 and answer a question of [GV06b] by providing an example which is not  $\aleph_0$ -tame but is  $2^{\aleph_0}$ -tame.

In the presence of amalgamation, the subject of this paper can be considered as a study of the automorphism group of the monster model. For example, compactness is the assertion: if  $M_i$  is an increasing sequence of strong submodels of  $\mathcal{M}$ ,  $G_i = \text{aut}_{M_i}(\mathcal{M})$ , and  $X_i$  is a decreasing sequence of orbits under  $G_i$ , then the intersection of the  $X_i$  is nonempty. The cardinal parameters of the formal definition fix the cardinality of the  $M_i$  and the length of the chain.

## 1 Some notions of locality

We work throughout in an abstract elementary class  $(\mathbf{K}, \prec_{\mathbf{K}})$ . A strong embedding is an injective homomorphism  $f$  from  $M$  into  $N$  such that  $fM \prec_{\mathbf{K}} N$ . A chain  $\langle A_i : i < \delta \rangle$  is a sequence of members of  $\mathbf{K}$  such that if  $i < j$ ,  $A_i \prec_{\mathbf{K}} A_j$ ; the chain is continuous if for each limit ordinal  $\alpha$ ,  $A_\alpha = \bigcup_{\beta < \alpha} A_\beta$ .

Although we compare the properties of Galois types and syntactic types, the types that actually occur in this paper are all Galois-types. So we fix the following notation.

**Definition 1.1** 1. For  $M \prec_{\mathbf{K}} N_1 \in \mathbf{K}$ ,  $M \prec_{\mathbf{K}} N_2 \in \mathbf{K}$  and  $a \in N_1 - M$ ,  $b \in N_2 - M$ , write  $(M, a, N_1) \sim_{AT} (M, b, N_2)$  if there exist strong embeddings  $f_1, f_2$  of  $N_1, N_2$  into some  $N^*$  which agree on  $M$  and with  $f_1(a) = f_2(b)$ .

2. Let  $\sim$  be the transitive closure of  $\sim_{AT}$  (as a binary relation on triples).

3. We say the Galois type of  $a$  over  $M$  in  $N_1$  is the same as the Galois type  $a$  over  $M$  in  $N_2$  if  $(M, a, N_1) \sim (M, b, N_2)$  and write

$$\text{tp}(a/M, N_1) = \text{tp}(b/M, N_2)$$

We define the notions below with two cardinal parameters: the first is the size of a certain submodel or the length of a sequence of types; the second is the size of the models under consideration. In a rough sense, the first parameter is the important one; ideally the second can be replaced by  $\infty$ . But the main theorem which derives locality from categoricity (without identifying Galois types with syntactic types of some sort) [She99], chapter 12 of [Bal00]) does so only for models of fixed size. So we use the fastidious notation. Replacing  $(\lambda, \kappa)$  by e.g.  $(< \lambda, \kappa)$  has the obvious meaning. The following property holds of all AEC considered in this paper.

**Definition 1.2** We say the AEC  $(\mathbf{K}, \prec_{\mathbf{K}})$  admits intersections if for every  $X \subseteq M \in \mathbf{K}$ , there is a minimal closure of  $X$  in  $M$ . That is,  $\text{cl}_M(X) \prec_{\mathbf{K}} M$  is the substructure of  $M$  with universe  $\bigcap \{N : X \subseteq N \prec_{\mathbf{K}} M\}$ .

Any relation defined by such an intersection will have the monotonicity, finite character, and transitivity properties of a closure relation. Note that this property is nontrivial even if one restricts to first order theories with elementary submodel. In that case it applies to strongly minimal or  $o$ -minimal theories; the first order case was characterized by Rabin [Rab62]. And of course the condition is satisfied when one has Skolem functions. But we work in a more general situation. If an AEC admits intersections we have a natural way to check equality of Galois types.

**Lemma 1.3** Let  $(\mathbf{K}, \prec_{\mathbf{K}})$  admit intersections.

1. Suppose  $M_0 \prec_{\mathbf{K}} M_1, M_2$  with  $a_i \in M_i$  for  $i = 1, 2$ . Then  $\text{tp}(a_1/M_0, M_1) = \text{tp}(a_2/M_0, M_2)$  if and only if there is an isomorphism over  $M_0$  from  $M_1 \upharpoonright_{\text{cl}_{M_1}(M_0 a_1)}$  onto  $M_2 \upharpoonright_{\text{cl}_{M_2}(M_0 a_2)}$  which maps  $a_1$  to  $a_2$ .
2.  $(M_1, a_1, N_1)$  and  $(M_2, a_2, N_2)$  represent the same Galois type over  $M_1$  iff  $M_1 = M_2$  and there is an amalgam of  $N_1$  and  $N_2$  over  $M_1$  where  $a_1$  and  $a_2$  have the same image.

Proof. Immediate. □<sub>1.3</sub>

That is, while in general Galois equivalence may result from a finite composition of maps, in this context only one step is required. If  $M \prec_{\mathbf{K}} N$ ,  $p \in \text{ga} - \text{S}(M)$ ,  $q \in \text{ga} - \text{S}(M)$ , the notion that  $q$  extends  $p$  is similarly complicated in an arbitrary AEC. Lemma 1.3 yields a simpler characterization.

**Fact 1.4** *Suppose  $\mathbf{K}$  admits intersections or has the amalgamation property. If  $M \prec_{\mathbf{K}} N$ ,  $p \in \text{ga} - \text{S}(M)$ ,  $q \in \text{ga} - \text{S}(M)$ , then  $q$  extends  $p$  if and only if for each  $(N, b, N')$  realizing  $p$  there is a  $\mathbf{K}$ -map fixing  $M$  and taking  $(N, b, N')$  to an  $(M, a, M')$  realizing  $q$ .*

We illustrate in this article that unions of increasing chains of Galois types do not behave as increasing chains of syntactic types. The problem is that to guarantee an increasing chain of types has an upper bound we need that it is coherent in the following sense.

**Definition 1.5** *Let  $\langle M_i : i < \gamma \rangle$  be an increasing  $\prec_{\mathbf{K}}$ -chain of submodels of  $\mathcal{M}$ . A coherent chain of Galois types of length  $\gamma$  is an increasing chain of types  $p_i \in \text{ga} - \text{S}(M_i)$  equipped with realizations  $a_i$  of  $p_i$  and for  $i < j < \gamma$  functions  $f_{ij} \in \text{aut}(\mathcal{M})$  such that  $f_{ij}$  fixes  $M_i$ ,  $f_{ij}(a_j) = a_i$  and for  $i < j < k < \gamma$ ,  $f_{ij} \circ f_{jk} = f_{ik}$ .*

Here is a characterization of realizing the union of a chain of types in terms of maps. The straightforward justification is in [Bal00] in the chapter on locality and tameness.

**Fact 1.6** 1. *If  $p_i \in \text{ga} - \text{S}(M_i)$  for  $i < \delta$  is a coherent chain of Galois types, there is a  $p_\delta \in \text{ga} - \text{S}(M_\delta)$  that extends each  $p_i$  so that  $\langle p_i : i \leq \delta \rangle$  is a coherent sequence.*

2. *Conversely,  $p_\delta \in \text{ga} - \text{S}(M_\delta)$  extends each  $p_i$ , there is a choice of  $f_{i,j}$  for  $i \leq j \leq \delta$  that witness  $\langle p_i : i \leq \delta \rangle$  is a coherent sequence.*

**Definition 1.7** *Galois types are  $(\kappa, \lambda)$ -compact in  $\mathbf{K}$  if for every continuous increasing chain  $M = \bigcup_{i < \kappa} M_i$  of members of  $\mathbf{K}$  with  $|M| = \lambda$  and every increasing chain  $\{p_i : i < \kappa\}$  of members  $\text{ga} - \text{S}(M_i)$  there is a  $p \in \text{ga} - \text{S}(M)$  with  $p \upharpoonright M_i = p_i$  for every  $i$ .*

**Definition 1.8**  *$\mathbf{K}$  has  $(\kappa, \lambda)$ -local galois types if for every continuous increasing chain  $M = \bigcup_{i < \kappa} M_i$  of members of  $\mathbf{K}$  with  $|M| = \lambda$  and for any  $p, q \in \text{ga} - \text{S}(M)$ : if  $p \upharpoonright M_i = q \upharpoonright M_i$  for every  $i$  then  $p = q$ .*

The following results were stated by Shelah in e.g. [She99]; a full proof appears in [Bal00].

**Lemma 1.9** *For any  $\lambda$ , if  $\mathbf{K}$  has  $(< \kappa, \leq \lambda)$ -local Galois types, then Galois types are  $(\leq \kappa, \leq \lambda)$ -compact in  $\mathbf{K}$ .*

Now we turn to the notion of tameness. The property was first isolated in [She99] in the midst of a proof. Grossberg and VanDieren [GV06b] focused attention on the notion as a general property of AEC's. We introduce a parameterized version in hopes of deriving tameness from categoricity by an induction. And weakly tame is the version that can actually be proved. That is, the best result now known [She99, Bal00] is that if  $\mathbf{K}$  is categorical in some regular  $\lambda$  greater than  $H_1 = H(\mathbf{K})$  (the Hanf number for  $\mathbf{K}$ ), then for each  $\mu < \lambda$ ,  $\mathbf{K}$  is  $(\chi, \mu)$ -weakly tame for some  $\chi < H_1$ .

**Definition 1.10** 1. *We say  $\mathbf{K}$  is  $(\chi, \mu)$ -weakly tame if for any saturated  $N \in \mathbf{K}$  with  $|N| \leq \mu$  if  $p, q \in \text{ga} - \text{S}(N)$  and for every  $N_0 \leq N$  with  $|N_0| \leq \chi$ ,  $p \upharpoonright N_0 = q \upharpoonright N_0$  then  $p = q$ .*

2. *We say  $\mathbf{K}$  is  $(\chi, \mu)$ -tame if the previous condition holds for all  $N$  with cardinality  $\mu$ .*

3.  *$(\chi, \mu)$ -weakly compact and  $(\chi, \mu)$ -weakly local are defined analogously.*

Thus the vague notion of  $\kappa$ -tame in the introduction is formally  $(\kappa, \infty)$ -tame. Finally, we say  $\mathbf{K}$  is  $\kappa$  (weakly)-tame or  $(\kappa, \infty)$  (weakly)-tame if it  $(\kappa, \lambda)$ - (weakly)-tame for every  $\lambda \geq \kappa$ . There are a few relations between tameness and locality. The second was observed in conversation by Olivier Lessmann.

**Lemma 1.11** *If  $\lambda \geq \kappa$  and  $\text{cf}(\kappa) > \chi$ , then  $(\chi, \lambda)$ -tame implies  $(\kappa, \lambda)$ -local. In particular,  $(\aleph_0, \aleph_1)$ -tame implies  $(\aleph_1, \aleph_1)$ -local.*

Proof. Suppose  $\langle M_i, p_i : i < \kappa \rangle$  is an increasing chain with  $\bigcup_i M_i = M$  and  $|M| \leq \lambda$ . If both  $p, q \in \text{ga} - \text{S}(M)$  extend each  $p_i$ , by  $(\chi, \lambda)$ -tameness, there is a model  $N$  of cardinality  $\chi$  on which they differ. Since  $\text{cf}(\kappa) > \chi$ ,  $N$  is contained in some  $M_i$ . □<sub>1.11</sub>

**Lemma 1.12** *If  $\mathbf{K}$  is  $(< \mu, < \mu)$ -local and  $\mu \geq \text{LS}(\mathbf{K})$  then  $\mathbf{K}$  is  $(\text{LS}(\mathbf{K}), \mu)$ -tame.*

Proof. We prove the result by induction on  $\mu$  and it is clear for  $\mu = \text{LS}(\mathbf{K})$ . Suppose it holds for all  $\kappa < \mu$ . Let  $p, q$  be distinct types in  $\text{ga} - \text{S}(M)$  where  $|M| = \mu$  and write  $M$  as an increasing chain  $\langle M_i : i < \mu \rangle$  with  $|M_i| \leq |i| + \text{LS}(\mathbf{K})$ . Let  $p_i$ , respectively  $q_i$  denote the restriction to  $M_i$ . Since  $p \neq q$ , locality gives an  $M_j$  with  $p_j \neq q_j$  and  $|M_j| < \mu$ . By induction there exists an  $N \prec_{\mathbf{K}} M_j$  with  $|N| = \text{LS}(\mathbf{K})$  and  $p_j \upharpoonright N \neq q_j \upharpoonright N$ . But then,  $p \upharpoonright N \neq q \upharpoonright N$  and we finish. □<sub>1.12</sub>

## 2 A Concrete Example of Non-tameness

In this section we find a concrete example of a class which is not  $(\aleph_1, \aleph_1)$ -local and so not  $(\aleph_0, \aleph_1)$ -tame. We encode some well-known ‘incompactness’ phenomena for Abelian groups. Recall again that all types are Galois types.

**Definition 2.1** *We say  $A$  is a Whitehead group if  $\text{Ext}(A, \mathbb{Z}) = 0$ . That is, every short exact sequence*

$$0 \rightarrow \mathbb{Z} \rightarrow H \rightarrow A \rightarrow 0,$$

*splits or in still another formulation,  $H$  is the direct sum of  $A$  and  $\mathbb{Z}$ .*

Every free group is Whitehead and a Whitehead group of power  $\aleph_1$  is  $\aleph_1$ -free, i.e., every countable subgroup is free. Recall that J.H.C. Whitehead conjectured that every Whitehead group of cardinality  $\aleph_1$  is free. We do not rely in this section on Shelah’s argument that the Whitehead conjecture is independent of ZFC. But we use some of the techniques of the argument and more appear in the next section. Now we contradict locality. We rely on Shelah’s construction, reported on page 228 of [EM90] of a group with the following properties. The moreover clause is not used in this section but is crucial for Section 5.

**Fact 2.2** *There is an  $\aleph_1$ -free group  $G$  of cardinality  $\aleph_1$  which is not Whitehead. Moreover, there is a countable subgroup  $R$  of  $G$  such that  $G/R$  is  $p$ -divisible for each prime  $p$ .*

**Example 2.3** Let  $\mathbf{K}$  be the class of structures  $M = \langle G, Z, I, H \rangle$ , where each of the listed sets is the solution set of one of the unary predicates  $(\mathbf{G}, \mathbf{Z}, \mathbf{I}, \mathbf{H})$ .  $G$  is a torsion-free Abelian Group.  $Z$  is a copy of  $(\mathbb{Z}, +)$ .  $I$  is an index set and  $H$  is a family of infinite Abelian groups. The vocabulary also includes function symbols  $\mathbf{F}, \mathbf{k}$  and  $\mathbf{\pi}$ , naming functions  $F, k$ , and  $\pi$ .  $F$  maps  $H$  onto  $I$  and for  $s \in I$ ,  $+(\cdot, \cdot, s)$  is a group operation on  $H_s = F^{-1}(s)$ . Finally,  $\pi$  maps  $H$  onto  $G$  so that  $\pi_s = \pi|_{H_s}$  is a projection from  $H_s$  onto  $G$ . The kernel of each  $\pi_s$  is isomorphic to  $Z$  via a map  $\mathbf{k}(\cdot, s)$  where  $k : Z \times I \mapsto H$ .

Further, we write  $M_0 \prec_{\mathbf{K}} M_1$  if  $M_0$  is a substructure of  $M$ , but  $\mathbf{Z}^{M_0} = \mathbf{Z}^{M_1}$  and  $\mathbf{G}^{M_0}$  is a pure subgroup of  $\mathbf{G}^{M_1}$ .

Since each  $\mathbf{G}^M$  is torsion-free, it follows that if  $M_0 \prec_{\mathbf{K}} M_1$ , for each  $t \in I^{M_0}$ ,  $\mathbf{H}_t^{M_0}$  is pure in  $\mathbf{H}_t^{M_1}$ . (Recall that if the abelian group  $G$  is elementarily equivalent to  $G_1$  and  $G$  is pure in  $G_1$  then  $G$  is an elementary submodel of  $G_1$ .) The class  $\mathbf{K}$  is almost first order definable; we require some infinitary logic to keep  $\mathbb{Z}$  standard. But the notion of  $\prec_{\mathbf{K}}$  is much weaker than elementary submodel. The models are essentially many exact sequences. They all have the same kernel  $\mathbb{Z}$ ; and there may be many with the same image  $G$ , but the middle terms  $H$  are all disjoint. It is fruitful (see Section 5) to restrict the class of image groups. But it is delicate to do so while keeping  $\mathbf{K}$  closed under unions and amalgamation.

It is easy to check that under these definitions

**Lemma 2.4** *The class  $(\mathbf{K}, \prec_{\mathbf{K}})$  defined in Example 2.3 is an abstract elementary class.*

We defined the notion of an AEC admitting intersections in Definition 1.2

**Lemma 2.5** *The class  $(\mathbf{K}, \prec_{\mathbf{K}})$  defined in Definition 2.3 admits intersections.*

Proof. To find the the required closure of a subset  $A$  of  $M$ , first close  $A$  under the functions of the language:  $\mathbf{F}^M, \pi^M$  and the group operation in  $G$  and in each fiber of  $\mathbf{H}^M$  to form a set  $X'$ . Then take the pure closure of  $X' \cap \mathbf{G}^M$  in  $\mathbf{G}^M$  as  $X''$ . Finally add those  $y \in \mathbf{H}^M$  which satisfy both  $\pi^M(y) \in X''$  and  $\mathbf{F}^M(y) \in X''$ . If  $N$  is the substructure of  $M$  with this universe it is easy to check that  $N \prec_{\mathbf{K}} M$  and  $N$  is contained in any  $N'$  with  $A \subset N' \prec_{\mathbf{K}} M$ .  $\square_{2.5}$

The next easy lemma provides a nice characterization for this example of equality for certain Galois types.

**Lemma 2.6** *Suppose  $M_0 \prec_{\mathbf{K}} M_1, M_2$  and the group  $G = \mathbf{G}^{M_0}$  is the same in each of the three structures. Let  $t_1, t_2$  be in  $\mathbf{I}^{M_1} - \mathbf{I}^{M_0}, \mathbf{I}^{M_2} - \mathbf{I}^{M_0}$  respectively. The following are equivalent.*

1.  $\text{tp}(t_1/M_0, M_1) = \text{tp}(t_2/M_0, M_2)$ .
2. *There is an isomorphism  $h$  from  $\mathbf{H}_{t_1}^{M_1}$  onto  $\mathbf{H}_{t_2}^{M_2}$  such that*
  - (a) *For  $z \in \mathbf{Z}^{M_0}$ ,  $h(\mathbf{k}^{M_1}(z, t_1)) = \mathbf{k}^{M_2}(z, t_2)$  and*
  - (b) *for  $y \in \mathbf{H}_{t_1}^{M_1}$ ,  $h(\pi^{M_1}(y, t_1)) = \pi^{M_2}(h(y), t_2)$ .*

Recall that the class of torsion-free abelian groups has the amalgamation property for pure embeddings. Specifically, to amalgamate  $G_1$  and  $G_2$  over  $G_0$  just form  $G_1 \times G_2$  and factor out the subgroup of elements  $\{(x, -x) : x \in G_0\}$ . The purity of  $G_0$  in  $G_1$  and  $G_2$  guarantees the amalgam is torsion free. Note that if  $H_0 \subset H_1, H_2$  and we have maps from  $H_1$  onto  $G_1$  and from  $H_2$  onto  $G_2$  with common kernel contained in  $H_0$ , these maps extend coordinate-wise to maps from the amalgam of the  $H$ 's to the amalgam of the  $G$ 's.

**Lemma 2.7**  *$(\mathbf{K}, \prec_{\mathbf{K}})$  has the amalgamation property.*

Proof. We want to amalgamate  $M_1$  and  $M_2$  over  $M_0$  to construct  $M_3$ . Without loss of generality we can assume  $M_1$  and  $M_2$  intersect in  $M_0$ . So just take disjoint union on  $\mathbf{I}$ , the group amalgams on  $\mathbf{G}$  and also the group amalgam of each  $\mathbf{H}_t^1$  and  $\mathbf{H}_t^2$  if  $t \in \mathbf{I}^{M_0}$  and extend the functions naturally. If  $t \in \mathbf{I}^1 - \mathbf{I}^2$ ,  $\mathbf{H}_t^1 = \mathbf{H}_t^3$ . (The case  $\mathbf{I}^1 - \mathbf{I}^2 \neq \emptyset$  is similar.) In particular,  $\pi^{M_3}(y_1, y_2)$  is only defined if  $\mathbf{F}^{M_1}(y_1) = \mathbf{F}^{M_2}(y_2)$  and in that case, the value is  $((\pi^{M_1}(y_1), (\pi^{M_2}(y_2))) / \mathbf{G}^{M_0})$ . There is no interaction between the problems for  $H_s$  and  $H_t$  if  $s \neq t \in \mathbf{I}^{M_1} \cup \mathbf{I}^{M_2}$ .

$\square_{2.7}$

**Lemma 2.8** *Let  $(\mathbf{K}, \prec_{\mathbf{K}})$  be as in Example 2.3.  $(\mathbf{K}, \prec_{\mathbf{K}})$  is not  $(\aleph_1, \aleph_1)$ -local. That is, there is an  $M^0 \in \mathbf{K}$  of cardinality  $\aleph_1$ , a continuous increasing chain of models  $M_i^0$  for  $i < \aleph_1$ , and two distinct types  $p, q \in \text{ga} - \text{S}(M^0)$  with  $p \upharpoonright M_i^0 = q \upharpoonright M_i^0$  for each  $i$ .*

*Proof.* We define  $p$  and  $q$ , show they are distinct, and then show their restrictions are the same.

Let  $G$  be the Abelian group from Fact 2.2 of cardinality  $\aleph_1$  which is  $\aleph_1$ -free but not a Whitehead group. Then, there is an  $H$  such that,

$$0 \rightarrow \mathbb{Z} \rightarrow H \rightarrow G \rightarrow 0$$

is exact but does not split. Say  $g_1 : H \rightarrow G$ . But, we can write  $G$  as a continuous increasing chain  $G = \bigcup_{i < \aleph_1} G_i$  of countable free groups such that each exact sequence:

$$0 \rightarrow \mathbb{Z} \rightarrow H_i \rightarrow G_i \rightarrow 0,$$

splits, where  $H_i = g_1^{-1}(G_i)$ , because  $G_i$  is free.

Let  $M^0$  have  $\mathbf{G}^{M^0} = G$ ,  $\mathbf{Z}^{M^0} = \mathbb{Z}$ ,  $\mathbf{I}^{M^0} = \{a\}$ , and  $\mathbf{H}^{M^0}$  a copy of  $G \oplus \mathbb{Z}$ . Now define  $M^1, M^2$  which have one additional point  $t_i \in \mathbf{I}^{M^i}$ . The key point is that  $\mathbf{H}^{M^1}$  is  $H$  and  $\mathbf{H}^{M^2}$  is  $G \oplus \mathbb{Z}$ . For ease of reading let  $\pi^{M^1}(-, t_1)$  be  $g_1$  and  $\pi^{M^2}(-, t_2)$  be the projection map  $g_2$  from  $G \oplus \mathbb{Z}$  onto  $G$ . Let  $p = \text{tp}(t_1/M^0, M^1)$  and  $q = \text{tp}(t_2/M^0, M^2)$ . Since the exact sequence for  $\mathbf{H}^{M^2}$  splits and that for  $\mathbf{H}^{M^1}$  does not, it is immediate from Lemma 2.6 that  $p \neq q$ .

Now we define the models  $M_i^\ell$  for  $i < \aleph_1$  and  $\ell = 0, 1, 2$ . Then  $M_i^0$  is naturally obtained by letting  $\mathbf{G}^{M_i^0} = G_i$  and leaving the other components as in  $M^0$ . For  $\ell = 1, 2$ , let  $M_i^\ell$  be the restriction of  $M^\ell$  to

$$\langle G_i, \mathbb{Z}, \{t_\ell\}, \{y \in \mathbf{H}^{M^\ell} : g_\ell(y) \in G_i\} \rangle.$$

By Lemma 1.3,  $\text{tp}(t_\ell/M_i^\ell, M^\ell) = \text{tp}(t_\ell/M_i^\ell, M^\ell)$  for each  $i$  and  $\ell = 1, 2$ . By the choice of  $H$ , for  $\ell = 1$  and by the restriction of  $g_2$  for  $\ell = 2$  each of the exact sequences:

$$0 \rightarrow \mathbb{Z} \rightarrow H_i^\ell \rightarrow G_i^\ell \rightarrow 0$$

splits. This implies there is an isomorphism  $h$  from  $M_i^1$  onto  $M_i^2$  over  $M_i^0$  mapping  $t_1$  to  $t_2$ . That is,  $\text{tp}(t_1/M_i^0, M_i^1) = \text{tp}(t_2/M_i^0, M_i^2)$ . Thus,  $p \upharpoonright M_i^0 = q \upharpoonright M_i^0$ . We have the required counterexample.  $\square_{2.8}$

**Remark 2.9** While the existence of an  $\aleph_1$ -free group which is not Whitehead can be done in ZFC,  $\kappa$ -free but not Whitehead groups for larger regular  $\kappa$  become sensitive to set theory. But if, for example  $V = L$  (much weaker conditions suffice), the class  $\mathbf{K}$  is not  $(\kappa, \kappa)$ -local for arbitrary regular  $\kappa$ .



### 3 Incompactness

In this section we construct an increasing sequence of Galois types which has no upper bound. The model theoretic example is the same as Section 2 but the choice of groups for the counterexample is different. In contrast to nonlocality we obtain only a consistency result. The set theoretic hypotheses,  $\square$  and the various diamonds, used below follow from  $V = L$ .

**Definition 3.1**  $\square$  holds if there exists  $\langle C_\alpha : \alpha < \omega_2, \alpha \text{ limit} \rangle$  such that:

1.  $C_\alpha$  is a club of  $\alpha$ ;
2.  $|C_\alpha| \leq \aleph_0$  when  $\text{cf}(\alpha) = \aleph_0$ ;
3. and if  $\beta \in C_\alpha$  is a limit ordinal,  $C_\beta = C_\alpha \cap \beta$ .

**Notation 3.2** Assuming  $\square$ , we fix this notation for subsets of  $\aleph_2$ .

1. For each  $\alpha$ , let

$$C'_\alpha = \{\beta \in C_\alpha : \beta = \sup(C_\alpha \cap \beta)\}$$

and

$$S = \{\alpha \in \omega_2 : \alpha = \sup(C'_\alpha)\}.$$

2.  $S_0$  is the elements of  $S$  with cofinality  $\omega$  and  $S_1$  is the elements of  $S$  with cofinality  $\omega_1$ . (Note that, in fact,  $S_1$  contains all ordinals less than  $\aleph_2$  of cofinality  $\omega_1$ ; i.e.  $S_1 = S_1^2$ .)
3. Choose a ladder system  $\{\eta_\delta : \delta \in S_0\}$  from the  $C'_\delta$ . That is, each  $\eta_\delta$  is an increasing  $\omega$  sequence with limit  $\delta$  of elements of  $C'_\delta$ .

We will use the  $\square$ -principle in the proof of Lemma 3.9.

**Theorem 3.3** Assume  $2^{\aleph_0} = \aleph_1$ , and  $\diamond_{\aleph_1}$ ,  $\square_{\aleph_2}$ , and  $\diamond_{S_1^2}$ , where

$$S_1^2 = \{\delta < \aleph_2 : \text{cf}(\delta) = \aleph_1\}.$$

If the  $\mathbf{K}$  from Section 2 is  $(\aleph_1, \aleph_1)$ -compact then it is not  $(\aleph_2, \aleph_2)$ -compact.

Note that this is a more precise version of the statement in the abstract that  $\mathbf{K}$  is not  $(\leq \aleph_2, \leq \aleph_2)$ -compact.

We will use several times the following fact, which is one of the equivalent conditions in Lemma IV.2.3 of [EM90].

**Fact 3.4 (Pontryagin's criterion)**  $G$  is an  $\aleph_1$ -free Abelian group if and only if  $G$  is torsion free and every finite subset of  $G$  is contained in a finitely generated pure subgroup of  $G$ .

The set theoretic hypotheses of Theorem 3.3 allow us to define subsets of  $\aleph_2$  satisfying the following conditions. From them we will define a family of groups to establish the result. Note that  $\diamond$  implies every Whitehead group of power  $\aleph_1$  is free [EM90, She74].

**Definition 3.5** For  $\alpha \leq \aleph_2$ , let  $G_\alpha$  be the Abelian group generated by

$$\{x_\beta : \beta < \alpha\} \cup \{y_{\delta,n} : \delta \in S_0 \cap \alpha, n < \omega\}$$

subject only to the relations  $n!y_{\delta,n+1} = y_{\delta,n} - x_{\eta_\delta(n)}$ , where  $n!$  is the factorial of  $n$ .

We use interval notation in the ordinals, writing  $\{\gamma : \alpha + 1 \leq \gamma < \beta\}$  as  $[\alpha + 1, \beta)$ .

**Lemma 3.6** With the notation above,

1. The  $G_\alpha$  form an increasing continuous sequence (under pure embedding) of  $\aleph_1$ -free abelian groups.
2. For  $\alpha < \beta \leq \aleph_2$ ,  $G_{\alpha+1}$  is a direct summand of  $G_\beta$ .

Proof. Check 1) using Pontryagin's criteria. We now prove 2. For  $\delta \in S_0 \cap [\alpha + 1, \beta)$ , choose  $b(\delta) = b < \omega$  so that  $\eta_\delta(b) > \alpha + 1$ . Let  $G_{\alpha+1,\beta}$  be the group generated by

$$\{x_\gamma : \alpha + 1 \leq \gamma < \beta\} \cup \{y_{\delta,m} : \delta \in S_0 \cap [\alpha + 1, \beta), b(\delta) \leq m\}.$$

Then  $G_\beta = G_{\alpha+1} \oplus G_{\alpha+1,\beta}$  as required, since there are no relations between the generators of  $G_{\alpha+1}$  and  $G_{\alpha+1,\beta}$  and each  $y_{\delta,n}$  with  $\eta_\delta(n) < b(\delta)$  can be written as a sum of elements from  $G_{\alpha+1}$  and  $G_{\alpha+1,\beta}$ .  $\square_{3.6}$

**Notation 3.7** ( $\diamond_{S_1^2}$ ) Let  $\langle F_\delta : \delta \in S_1 \rangle$  be a diamond sequence, i.e.,

1.  $F_\delta$  is a two-place function with domain  $\delta$ ,  $F_\delta(\gamma_1, \gamma_2)$  is a permutation of some  $\beta_{\delta,\gamma_1,\gamma_2} < \delta$  for  $\gamma_1 < \gamma_2 < \delta$ .
2. If  $\bar{f} = \langle f_{\gamma_1,\gamma_2} : \gamma_1 < \gamma_2 < \aleph_2 \rangle$ ,  $f_{\gamma_1,\gamma_2}$  is a permutation of some  $\beta_{\gamma_1,\gamma_2} < \aleph_2$  then  $\{\delta \in S_1 : (\forall \gamma_1 \gamma_2)(\gamma_1 < \gamma_2 < \delta) f_{\gamma_1,\gamma_2} = F_\delta(\gamma_1, \gamma_2)\}$  is stationary.

We can assume the universe of  $G_\alpha$  is an ordinal  $\delta_\alpha < \aleph_2$ , since the  $G_\alpha$  are a continuous increasing sequence.

Now we construct by induction on  $\alpha$  an array of Abelian groups indexed by the pairs below the diagonal in  $\aleph_2 \times \aleph_2$ :  $\langle \langle H_{\beta,\alpha}, g_{\beta,\alpha} \rangle : \beta \leq \alpha < \aleph_2 \rangle$  and  $\langle \pi_{\beta,\alpha} : \beta \leq \alpha < \aleph_2 \rangle$  which satisfy the following pair of conditions:

- $A_\alpha$ :
  1.  $H_{\alpha,\alpha}$  is an abelian group with universe  $\delta_\alpha$ .
  2.  $g_{\alpha,\alpha} = g_\alpha$  is a homomorphism from  $H_{\alpha,\alpha}$  onto  $G_\alpha$  with kernel  $\mathbf{Z}$ .
  3. For  $\beta < \alpha$ ,  $H_{\beta,\alpha} = H_{\alpha,\alpha} \upharpoonright \{x \in H_{\alpha,\alpha} : g_{\alpha,\alpha}(x) \in G_\beta\}$ ;  $H_\beta$  denotes  $H_{\beta,\beta}$ .
  4.  $g_{\beta,\alpha} = g_{\alpha,\alpha} \upharpoonright H_{\beta,\alpha}$ .

5. If  $\beta \leq \alpha < \aleph_2$ ,  $\pi_{\beta,\alpha}$  is an isomorphism from  $H_\beta$  onto  $H_{\beta,\alpha}$  such that  $g_{\beta,\beta} = g_{\beta,\alpha} \circ \pi_{\beta,\alpha}$ . (Each  $\pi_{\beta,\beta}$  is the identity map).

- $B_\alpha$ : If  $\alpha \in S_1$ , the sequence of maps  $F_\alpha$  (given by the diamond on  $S_1$ ) does not satisfy  $B(\alpha, F_\alpha)$  (Definition 3.8).

**Definition 3.8** [ $B(\alpha, F_\alpha)$ ] Suppose  $\alpha \in S_1$ , and for each  $\beta_1 \leq \beta_2 < \alpha$ ,  $F_\alpha(\beta_1, \beta_2)$  is a permutation of  $H_{\beta_1}$ .  $B(\alpha, F_\alpha)$  holds if

$\langle H_\beta; \pi_{\beta_2, \beta_1} \circ F_\alpha(\beta_1, \beta_2) : \beta_1 \leq \beta_2 \leq \alpha \rangle$  form a sequence of commuting maps. That is, for  $\beta_1 \leq \beta_2 \leq \beta_3 \leq \alpha$  and any  $x \in H_{\beta_1}$ :

$$(\pi_{\beta_3, \beta_1} \circ F_\alpha(\beta_1, \beta_3))(x) = (\pi_{\beta_3, \beta_2} \circ F_\alpha(\beta_2, \beta_3))((\pi_{\beta_2, \beta_1} \circ F_\alpha(\beta_1, \beta_2))(x)).$$

These automorphisms and projections are a slightly different formalism for describing the realization of unions of types than that described in Fact 1.6. We work directly with the groups; in Section 2 we represented the extension groups by a single element using the map  $\pi$ .

We need one more lemma concerning the structure of the  $G_\alpha$ ; it is here that we invoke  $\square$ .

**Lemma 3.9** If  $\alpha \in S_1$  then  $G_\alpha$  can be decomposed as  $G_\alpha = G'_\alpha \oplus G''_\alpha$  where  $G'_\alpha$  is countable and free.

Proof. Let

$$G'_\alpha =: \langle \{x_\beta : \beta \in C'_\alpha\} \cup \{y_{\beta,n} : \beta \in S_0 \cap C'_\alpha, n < \omega\} \rangle_{G_\alpha}$$

and

$$G''_\alpha = \langle \{x_\beta : \beta \in \alpha \setminus C'_\alpha\} \cup \{y_{\delta,n} : \delta \in \alpha \cap S_0 \setminus C'_\alpha \text{ and } \eta_\delta(n) > \sup(C'_\alpha \cap C'_\delta)\} \rangle_{G_\alpha}.$$

Now if  $\delta \in \alpha \cap S_0 \setminus C'_\alpha$  then  $\{n : \eta_\delta(n) \in C'_\alpha\}$  is finite (otherwise  $\delta = \sup(C'_\delta \cap C'_\alpha)$ , whence  $\delta \in C'_\alpha$ ). And the  $y_{\delta,n}$  with  $\eta_\delta(n) \leq \sup(C'_\alpha \cap C'_\delta)$  are represented as sums of elements in  $G'_\alpha$  and  $G''_\alpha$ . Since  $C'_\alpha$  is countable,  $G'_\alpha$  is countable and free since  $G_\alpha$  is  $\aleph_1$ -free.  $\square_{3.9}$

**Construction 3.10** We construct groups  $H_{\beta,\alpha}$  and functions  $g_{\alpha,\beta}$ ,  $\pi_{\alpha,\beta}$  for  $\beta \leq \alpha \leq \aleph_2$  by induction to satisfy conditions  $A_\alpha$  and  $B_\alpha$ .

Let  $H_0$  be  $\mathbb{Z} \oplus G_0$ ;  $g_0$  is the projection of  $H_0$  onto  $G_0$ ;  $\pi_{0,0}$  is the identity.

To satisfy  $A_\alpha$  in limit stages of cofinality  $\omega$ : (One is tempted to just take unions; the  $\pi_{\beta,\alpha}$  have not been constructed to commute so this may fail.) For  $\delta \notin S_0$ , choose a sequence  $\eta_\delta(n)$  with limit  $\delta$ ; for  $\delta \in S_0$ , we already have one. For the moment, we consider only the structures  $H_{\eta_\delta(m), \eta_\delta(k)}$  with  $m, k < \omega$ . We form new maps  $\pi_{\eta_\delta(m), \eta_\delta(k)}^*$  by the composition of  $\pi_{\eta_\delta(r), \eta_\delta(r+1)}$  for  $m \leq r \leq k$ . Now for each  $k < \omega$  the  $\langle H_{\eta_\delta(s)}, \pi_{\eta_\delta(s), \eta_\delta(t)}^* : s \leq t < \omega \rangle$  form a direct system and we can choose  $H_\delta$  as the direct limit of this system with limit maps  $\pi_{\eta_\delta(s), \delta}^*$  from  $H_{\eta_\delta(s)}$  into  $H_\delta$ . Denote the range of  $\pi_{\eta_\delta(s), \delta}^*$  as  $H_{\eta_\delta(s), \delta}$ . Now define  $g_{\eta_\delta(r), \delta}$

from  $H_{\eta_\delta(r),\delta}$  onto  $G_{\eta_\delta(r)}$  as  $g_{\eta_\delta(r),\eta_\delta(r)} \circ (\pi^*)_{\eta_\delta(r),\delta}^{-1}$ . This gives 1) through 3) of  $A_\delta$ . Now we satisfy 4) and 5) by defining  $H_{\beta,\delta}$  and  $g_{\beta,\delta}$  in the natural manner. It remains to define  $\pi_{\delta,\gamma}^*$  when  $\gamma$  is not in the range of  $\eta_\delta$ . Choose  $m$  such that  $\eta_\delta(m) > \gamma$  and let

$$\pi_{\delta,\gamma}^* = \pi_{\delta,\eta_\delta(m)}^* \circ \pi_{\eta_\delta(m),\gamma}.$$

To satisfy  $A_{\alpha+1}$  in a successor stage: given  $0 \rightarrow \mathbb{Z} \rightarrow H_\alpha \rightarrow G_\alpha \rightarrow 0$ , we proceed as follows. Let  $\alpha = \beta + 1$ . If  $\beta$  is not in  $S_0$ ,  $G_\alpha = G_\beta \oplus \langle x_\beta \rangle$  and we just extend  $H_\beta$  freely by a single generator. If  $\beta \in S_0$ , choose elements  $x'_{\beta,n} \in H_\beta$  with  $g_\beta(x'_{\beta,n}) = x_{\eta_\beta(n)}$ . Now form  $H_\alpha$  by adding to  $H_\beta$  elements  $x'_\beta$  and  $y'_{\beta,n}$  subject only to the relations:

$$n!y'_{\beta,n} = y'_{\beta,n-1} - x'_{\beta,n}.$$

Now if we map  $H_\alpha$  to  $G_\alpha$  by  $g_\beta$  on  $H_\beta$  and just dropping the primes on the generators of  $H_\beta$  over  $G_\beta$ , we have the required homomorphism.

We now consider  $\alpha$  of cofinality  $\aleph_1$ . Let  $C'_\alpha = \{\gamma_\varepsilon : \varepsilon < \aleph_1\}$ ,  $\gamma_\varepsilon$  increasing continuous with  $\varepsilon$ . We choose by induction objects  $\langle H_{\beta,\alpha}^0, g_{\beta,\alpha}^0, \pi_{\beta,\alpha}^0 : \beta \leq \gamma_\varepsilon \rangle$  to satisfy the relevant parts of  $A_\alpha$ . Let  $H_{\alpha,\alpha}^0 = \cup\{H_{\beta,\alpha}^0 : \beta < \alpha\}$ ,  $\pi_{\alpha,\alpha}^0 = \text{id}_{H_{\alpha,\alpha}^0}$ ,  $g_{\beta,\alpha}^0 = \cup\{g_{\beta,\alpha}^0 : \beta < \alpha\}$ . Consider an  $M \in \mathbf{K}$  with  $\mathbf{G}^M = G_\alpha$  and for some  $t \in I$ ,  $\mathbf{H}_t^M = H_{\alpha,\alpha}^0$ . We are assuming that  $\mathbf{K}$  is  $(\aleph_1, \aleph_1)$ -compact so Condition  $A_\alpha$  holds. If  $B_\alpha$  holds for this choice, i.e.  $F_\alpha(\alpha) \neq \text{id}_{H_{\alpha,\alpha}^0}$ , we are done. If not, recall the decomposition of  $G_\alpha$  in Lemma 3.9.

Let

$$H'_\alpha = \{x \in H_{\alpha,\alpha}^0 : g_{\alpha,\alpha}^0(x) \in G'_\alpha\}$$

$$H''_\alpha = \{x \in H_{\alpha,\alpha}^0 : g_{\alpha,\alpha}^0(x) \in G''_\alpha\}$$

(so their intersection is the copy of  $\mathbb{Z}$ ).

Since we have assumed  $\diamond_{\aleph_1}$ , every Whitehead group of power  $\aleph_1$  is free.  $\text{Ext}(G'_\alpha, \mathbb{Z}) \neq 0$  as  $G'_\alpha$  is not free. Hence, we can find  $(H_\alpha^*, g_\alpha^*)$  such that

- (a)  $H_\alpha^*$  is an abelian group;
- (b)  $g_\alpha^*$  is a homomorphism from  $H_\alpha^*$  onto  $G'_\alpha$ , which does not split;
- (c)  $\text{Ker}(g_\alpha^*) = \text{Ker}(g_{\alpha,\alpha}^0) \subseteq H_{\alpha,\alpha}$ ;
- (d)  $H_\alpha^* \cap H''_\alpha = \text{Ker}(g_\alpha^*)$ ;

Now we have a new candidate for  $H_{\alpha,\alpha}$ :

$$H_{\alpha,\alpha}^1 = H_\alpha^* \bigoplus_{\text{Ker}(g_{\alpha,\alpha}^0)} H''_\alpha$$

where we define  $g_{\alpha,\alpha}^1 \in \text{Hom}(H_{\alpha,\alpha}^1, G_\alpha)$  by extending  $g_{\alpha,\alpha}^0$  on  $H''_\alpha$  and  $g_\alpha^*$  on  $H_\alpha^*$ .

It remains to construct  $\pi_{\beta,\alpha}^1$  for  $\beta < \alpha$ . Note that

$$G_\beta = (G'_\alpha \cap G_\beta) \oplus (G''_\alpha \cap G_\beta).$$

On those  $x \in H_\beta$  with  $g_\beta(x) \in G'_\alpha \cap G_\beta$ , let  $\pi_{\beta,\alpha}^1(x) = \pi_{\beta,\alpha}^0(x)$ . We need to define  $\pi_{\beta,\alpha}^1(x)$  on  $\hat{H}_\beta = \{x \in H_\beta : g_\beta(x) \in \hat{G} = G''_\alpha \cap G_\beta\}$ . Let  $H_\beta^* = \{x \in H_\alpha^* : g_\alpha^1(x) \in \hat{G}\}$ . Since  $\hat{G}$  is free,  $0 \rightarrow \mathbb{Z} \rightarrow H_\beta^* \rightarrow \hat{G} \rightarrow 0$  splits;  $g_\alpha^*$  has an inverse  $f_\beta$ . Let  $\pi_{\beta,\alpha}^1 = f_\beta \circ g_\beta$ .

This shows  $B_\alpha$  is satisfied since  $F_\alpha(\alpha)$  cannot commute with both the  $\pi_{\beta,\alpha}^0$  and  $\pi_{\beta,\alpha}^1$ . Thus we choose  $(H_{\alpha,\alpha}, g_{\beta,\alpha}, \pi_{\beta,\alpha})$  from  $(H_{\alpha,\alpha}^0, g_{\beta,\alpha}^0, \pi_{\beta,\alpha}^0)$  and  $(H_{\alpha,\alpha}^1, g_{\beta,\alpha}^1, \pi_{\beta,\alpha}^1)$  as the sequence that satisfies  $B_\alpha$ .

Since we assumed that  $\mathbf{K}$  is  $(\aleph_1, \aleph_1)$  compact, able to carry out the construction for  $\aleph_2$  steps to obtain a sequence  $\langle H_{\beta,\aleph_2}, g_{\aleph_2,\beta} \circ \pi_{\beta,\aleph_2} : \beta \leq \aleph_2 \rangle$ . We finish as follows:

**Claim 3.11** *It is not possible to define  $\pi_{\beta,\aleph_2} : \beta \leq \aleph_2$  so that:*

- *The sequence  $\langle H_{\beta,\aleph_2}, g_{\aleph_2,\beta} \circ \pi_{\beta,\aleph_2} : \beta \leq \aleph_2 \rangle$  satisfies:*
  - (a)  $H_{\aleph_2,\aleph_2}$  an abelian group;
  - (b)  $g_{\aleph_2,\aleph_2}$  maps  $H_{\aleph_2,\aleph_2}$  onto  $G_{\aleph_2}$ ;
  - (c)  $H_{\beta,\aleph_2} = \{x \in H_{\aleph_2,\aleph_2} : g_{\aleph_2,\aleph_2}(x) \in G_\beta\}$ ;
  - (d)  $g_{\beta,\aleph_2} = g_{\aleph_2,\aleph_2} \upharpoonright H_{\beta,\aleph_2}$ ;
  - (e)  $\pi_{\beta,\aleph_2}$  is an isomorphism from  $H_{\beta,\aleph_2}$  onto  $H_{\beta,\beta}$ ;
  - (f) if  $x \in H_{\beta,\aleph_2}$  then  $g_{\beta,\aleph_2}(x) = g_{\beta,\beta}(\pi_{\aleph_2,\beta}(x)) \in G_\beta$ .

Proof. Suppose for contradiction that we have constructed such a sequence. Then letting,

$$\rho_{\beta_2,\beta_1} = \pi_{\beta_2,\beta_1}^{-1} \circ \pi_{\aleph_2,\beta_2}^{-1} \circ \pi_{\aleph_2,\beta_1}$$

we have a system  $\langle H_{\beta,\beta}, (\pi_{\beta_2,\beta_1} \circ \rho_{\beta_2,\beta_1}) : \beta < \aleph_2, \beta_1 \leq \beta_2 < \aleph_2 \rangle$  of commuting maps. But by the choice of  $\langle F_\delta : \delta \in S_1 \rangle$  and since, by  $\diamond_{S_1^2}$  and  $S_1 = S_1^2$ ,  $S_1$  is stationary in  $\aleph_2$  for some  $\delta^*$  we have:

$$(\forall \gamma_1 \leq \gamma_2 < \delta^*) F_{\delta^*}(\gamma_1, \gamma_2) = \rho_{\gamma_2,\gamma_2}.$$

This contradicts  $B_{\delta^*}$  in the construction and we finish. □<sub>3.11</sub>

Now we have a counterexample to  $(\aleph_2, \aleph_2)$ -compactness. The  $H_{\beta,\aleph_2}$  from Claim 3.11 give rise to a sequence of Galois types (of singletons via the coding spelled out in Section 2) over the  $G_\beta$  which have no common extension over  $G_{\aleph_2}$ . □<sub>3.3</sub>

Fact 1.9 implies that if  $\mathbf{K}$  were  $(\aleph_0, \aleph_1)$ -local then it would be  $(\aleph_1, \aleph_1)$ -compact and we would have an example of non- $(\aleph_1, \aleph_1)$ -compactness. But this route is not open to us. We now show that  $\mathbf{K}$  even fails  $(\aleph_0, \aleph_0)$ -locality (and failure of  $(\aleph_0, \aleph_1)$ -locality is an easy consequence in this case).

**Lemma 3.12**  $K$  is not  $(\aleph_0, \aleph_0)$ -local.

Proof. We construct a sequence of pairs of Abelian group  $(H_\alpha, G_\alpha)$  such that for  $\alpha < \omega$ ,  $H_\alpha = Z \oplus G_\alpha$ , but  $H_\omega$  is not a split extension of  $G_\omega$ . Since  $Z \oplus G_\omega$  is another limit of this chain, we contradict locality.

Let  $H^+$  be the  $\mathbb{Q}$ -vector space generated by elements  $x, z, y_n$  for  $n < \omega$ . Fix distinct odd primes  $p_n$  and  $q_{n,k}$  for  $n, k < \omega$ . We denote  $(p_n - 1)/2$  by  $r_n$ .

For each  $n < \omega$ ,  $H_n$  is the subgroup of  $H^+$  generated by  $x, z, y_n$  for  $n < \omega$  and the elements  $\frac{p_n y_n + x - r_n z}{q_{n,k}}$  for  $k < \omega$ .

Clearly  $H_n$  is a pure subgroup of  $H_{n+1}$ ; let  $H_\omega = \cup_n H_n$

**Claim 3.13**  $\mathbb{Z}z = \{nz : n \in \mathbb{Z}\}$  is a direct summand of  $H_n$ .

Proof. Since  $\mathbb{Z}$  is free every projection onto  $\mathbb{Z}z$  splits. So we need only construct a homomorphism  $h_n$  from  $H_n$  onto  $\mathbb{Z}z$ . Choose, by the Chinese remainder theorem  $r'_n$  such that  $r'_n \equiv r_\ell \pmod{p_\ell}$  for all  $\ell < n$ . Now let  $h_n(z) = z$ ,  $h_n(x) = r'_n z$ ,  $h_n(y_\ell) = -\frac{(r'_n - r_n)z}{p_\ell}$ , and  $h_n(\frac{p_n y_n + x - r_n z}{q_{n,k}}) = 0$ .

The choice of  $r'_n$  guarantees that each  $h_n(y_\ell) \in \mathbb{Z}z$  (the coefficient is an integer). Clearly  $h_n$  maps onto  $\mathbb{Z}z$ ; the danger is that it is not well defined. It suffices to show that from the values of  $h_n$  on  $z, x$  and the  $y_l$ ,  $h_n(p_n y_n + x - r_n z) = 0$  since that makes our definition consistent. For this, we compute:

$$h_n(p_n y_n + x - r_n z) = -p_n \frac{(r'_n - r_n)z}{p_n} + r'_n z - r_n z = 0.$$

□<sub>3.13</sub>

**Claim 3.14**  $\mathbb{Z}z = \{nz : n \in \mathbb{Z}\}$  is not a direct summand of  $H_\omega$ .

Proof: Suppose for contradiction that  $h$  retracts  $H_\omega$  onto  $\mathbb{Z}z$  (i.e.  $h(z) = z$ ). Now, for any  $n$ ,  $h(p_n y_n + x - r_n z) \in \mathbb{Z}z$  is divisible by  $q_{k,n}$  for all  $k$ . This implies that

$$h(p_n y_n + x - r_n z) = 0.$$

That is,  $h(x - r_n z) \in p_n \mathbb{Z}z$ . Since  $h(x) = r z$  some  $z \in \mathbb{Z}$ , this implies  $r \equiv r_n \equiv (p_n - 1)/2 \pmod{p_n}$ . But it impossible for this to happen for infinitely many  $n$  so  $\mathbb{Z}z$  is not a direct summand of  $H_\omega$ . □<sub>3.14</sub>

With these two claims we complete the proof of Lemma 3.12. □<sub>3.12</sub>

## 4 A General Construction for Amalgamation

Let  $(\mathbf{K}, \prec_{\mathbf{K}})$  be an aec in a relational language  $\tau$  which admits intersections and is model complete. In this section we construct from  $(\mathbf{K}, \prec_{\mathbf{K}})$  an AEC  $(\mathbf{K}', \prec_{\mathbf{K}'})$  which satisfies the amalgamation property and has the same non-locality properties as  $\mathbf{K}$ . The construction will apply to all AEC which admit intersections. We proceed in three steps; we first make a cosmetic change in

$\mathbf{K}$  to guarantee that it has quantifier free closures (Definition 4.1). Then we establish some important properties of AEC with quantifier free closures and finally make the main construction.

Throughout this section we assume that  $\mathbf{K}$  admits intersections (Definition 1.2); this simplifies the notions of Galois type and extension of Galois type (Lemma 1.3).

We require some preliminary definitions and a lemma for our main construction. Note that throughout we write boldface  $\mathbf{a}$  for a finite sequence of elements of a model and  $a$  for a single element.

**Definition 4.1** *Suppose  $(\mathbf{K}, \prec_{\mathbf{K}})$  admits intersections.*

1.  $\mathbf{K}$  is said to have quantifier-free closure if the satisfaction of ‘ $b \in \text{cl}_M(\mathbf{b})$ ’ depends only on the quantifier free (syntactic) type of  $\mathbf{bb}$ .
2.  $\mathbf{K}$  is model complete if  $N \subset M$  and  $N \in \mathbf{K}$ , implies  $N \prec_{\mathbf{K}} M$ .

**Lemma 4.2** *For any AEC  $(\mathbf{K}, \prec_{\mathbf{K}})$  which admits intersections there is an associated AEC  $(\mathbf{K}', \prec_{\mathbf{K}'})$  with exactly the same spectrum of models which has quantifier-free closure.*

Proof. Add to the language  $\tau$  of  $\mathbf{K}$ ,  $n + 1$ -ary relation symbols for each  $n$  and expand  $M \in \mathbf{K}$  to  $M' \in \mathbf{K}'$  by making  $R_n(\mathbf{a}, a)$  hold just if  $a \in \text{cl}_M(\mathbf{a})$ ;  $\mathbf{a}$  has length  $n$ . Let  $\mathbf{K}'$  be exactly the models of this form and define  $M' \prec_{\mathbf{K}'} N'$  if and only if  $M' \upharpoonright \tau \prec_{\mathbf{K}} N' \upharpoonright \tau$ . The isomorphism of  $\mathbf{K}$  and  $\mathbf{K}'$  is immediate and we have introduced quantifier-free closure by fiat.  $\square_{4.2}$

We now introduce a property that will be key in establishing amalgamation and show that it follows from either model completeness or having quantifier-free closure.

**Definition 4.3** *A class  $(\mathbf{K}, \prec_{\mathbf{K}})$ , which admits intersections, is said to be nice for unions if whenever  $\langle M_i : i \leq \delta \rangle$  is a continuous increasing chain of  $\mathbf{K}$ -extensions and  $A$  is a finite subset of  $N \subseteq M_\delta$  with  $N \in \mathbf{K}$ , there is an  $N'$  and an  $i < \delta$  such that  $A \subseteq N' \prec_{\mathbf{K}} N$  and  $N' \prec_{\mathbf{K}} M_i$ .*

Note that for  $A \subset M_1 \prec_{\mathbf{K}} M_2$ ,  $\text{cl}_{M_1}(A) = \text{cl}_{M_2}(A)$ . Moreover, if  $\mathbf{K}$  admits intersections for any  $A \subseteq N \in \mathbf{K}$ ,  $\text{cl}_N(A)$  is defined: take direct limits of the closures of finite sets.

**Lemma 4.4** *Let  $(\mathbf{K}, \prec_{\mathbf{K}})$  be an AEC which admits intersections. If  $\mathbf{K}$*

1. *is model complete or*
2. *has quantifier free closure*

*then it is nice for unions.*

Proof. As  $A$  is finite  $\text{cl}_{M_\delta}(A) = \text{cl}_{M_i}(A)$  when  $i = \min\{j < \delta : A \subseteq M_j\}$ . Thus,  $\text{cl}_{M_\delta}(A) \prec_{\mathbf{K}} M_i$ .

Case 1: model complete

So as  $N, M_\delta \in \mathbf{K}$ ,  $N \subseteq M_\delta$  we have  $N \prec_{\mathbf{K}} M_\delta$  so

$$\text{cl}_N(A) = \text{cl}_{M_\delta}(A) \prec_{\mathbf{K}} M_i.$$

Case 2: has quantifier free closure

Clearly  $\text{cl}_N(A) = \text{cl}_{M_\delta}(A)$  and as we observed to start the proof  $\text{cl}_{M_\delta}(A) \prec_{\mathbf{K}} M_i$ .

□<sub>4.4</sub>

Now we pass to the main construction.

**Definition 4.5** *Let  $\mathbf{K}$  be an AEC with a relational vocabulary  $\tau$ . The vocabulary  $\tau'$  of  $\mathbf{K}'$  is obtained by adding one additional binary relation  $R$ . We say the domain of a  $\tau'$ -structure  $A$  is an  $R$ -set if  $R$  induces a complete graph on  $A$ .*

1. *The class  $\mathbf{K}'$  is those  $\tau'$ -structures  $M$  such that:*
  - (a) *If the finite subset  $A$  of  $M$  is an  $R$ -set there is a  $\tau'$ -structure  $M_A$  such that  $A \subseteq M_A \subseteq M$  with  $|M_A| \leq \text{LS}(\mathbf{K})$ ,  $M_A$  is an  $R$ -set, and  $M_A \upharpoonright \tau \in \mathbf{K}$ .*
  - (b) *If  $N \subset M$  satisfies the conditions on  $M_A$  in requirement 1), then  $M_A \upharpoonright \tau \prec_{\mathbf{K}} N \upharpoonright \tau$ .*
  - (c) *For each  $M$  and  $A$ , we denote  $M_A$  by  $\text{cl}'_M(A)$ .*
2. *If  $M_1 \subseteq M_2$  are each in  $\mathbf{K}'$ , then  $M_1 \prec_{\mathbf{K}'} M_2$  if for each finite  $R$ -set  $A$  in  $M_1$ ,  $\text{cl}'_{M_1}(A) = \text{cl}'_{M_2}(A)$ .*

If  $M \subset N$  are  $\tau'$ -structures in  $\mathbf{K}'$  and  $M \prec_{\mathbf{K}'} N$  then for any finite  $A \subset M$ ,  $\text{cl}'_M(A) = \text{cl}'_N(A)$ .  $M \prec_{\mathbf{K}'} N$  does not imply  $M \upharpoonright \tau \prec_{\mathbf{K}} N \upharpoonright \tau$ ; indeed  $N \upharpoonright \tau$  is not necessarily in  $\mathbf{K}$ .

**Lemma 4.6** *Let  $(\mathbf{K}, \prec_{\mathbf{K}})$  be an aec in a relational language which admits intersections and is nice for unions. Then,  $(\mathbf{K}', \prec_{\mathbf{K}'})$  is an AEC with amalgamation.*

*Proof.* The axioms for an AEC which do not involve unions are easy. For example, we show ‘coherence’ in  $\mathbf{K}'$ . Suppose  $M \subset N \prec_{\mathbf{K}'} N'$  are  $\tau'$ -structure and also  $M \prec_{\mathbf{K}'} N'$ . Fix any finite  $A \subset M$ . Since  $N \prec_{\mathbf{K}'} N'$ ,  $\text{cl}'_N(A) = \text{cl}'_{N'}(A)$ . Since  $M \prec_{\mathbf{K}'} N'$ ,  $\text{cl}'_M(A) = \text{cl}'_{N'}(A)$ . Thus,  $\text{cl}'_M(A) = \text{cl}'_N(A)$  and  $M \prec_{\mathbf{K}'} N$ .

Suppose  $\langle M_i : i \leq \delta \rangle$  is a continuous increasing chain of  $\mathbf{K}'$ -extensions. Let  $A$  be a finite  $R$ -set contained in  $M_\delta$ . Fix the least  $j < \delta$  with  $A \subset M_j$ . Then, each  $i \geq j$ ,  $\text{cl}_{M_i}(A) = \text{cl}_{M_j}(A)$ ; call this set  $M_A$ ; it satisfies the conditions of Definition 4.5 1) a). Consider any other  $N \subset M_\delta$  which satisfies the conditions of Definition 4.5 1) a); that is,  $N$  is an  $R$ -set containing  $A$  and  $N \upharpoonright \tau \in \mathbf{K}$ . To



show  $M_\delta \in \mathbf{K}'$ , we must show  $M_A \upharpoonright \tau \prec_{\mathbf{K}} N \upharpoonright \tau$ . Since  $\mathbf{K}$  is nice for unions there is a  $k < \delta$  and a  $\tau$ -structure  $N'$  with  $N' \prec_{\mathbf{K}} M_k \upharpoonright \tau$ ,  $A \subset N'$  and  $N' \prec_{\mathbf{K}} N \upharpoonright \tau$ . We don't know whether the  $\tau'$  structure with universe  $N'$  is in  $\mathbf{K}'$ . But  $M_A$  is an  $R$ -set;  $M_A \upharpoonright \tau \prec_{\mathbf{K}} M_k$ ,  $M_A \subseteq N'$  and  $N' \prec_{\mathbf{K}} M_k \upharpoonright \tau$ . So, by coherence in  $\mathbf{K}$ ,  $M_A \upharpoonright \tau \prec_{\mathbf{K}} N'$  and we know  $N' \prec_{\mathbf{K}} N \upharpoonright \tau$ . By transitivity of  $\prec_{\mathbf{K}}$ , this is exactly what is needed.

Now we show the second union axiom. Suppose  $\langle M_i : i < \delta \rangle$  is a continuous increasing  $\mathbf{K}'$ -chain with each  $M_i \prec_{\mathbf{K}'} M$ . Then  $M_\delta \prec_{\mathbf{K}'} M$ , since we have shown for any finite  $A \subset M_\delta$  there is an  $i$  with  $\text{cl}'_{M_\delta}(A) = \text{cl}'_{M_i}(A)$ .

To show amalgamation suppose  $M_0 \prec_{\mathbf{K}'} M_1, M_2$ . Without loss of generality  $M_1 \cap M_2 = M_0$ . Now, form the no-edges amalgamation of the underlying graphs of  $M_1$  and  $M_2$  over  $M_0$ . The structure with this domain is in  $\mathbf{K}'$ , as each finite  $R$ -subset  $A$  of it is in either  $M_1$  or  $M_2$ ; the closure of each such  $A$  to satisfy Definition 4.5 is easily found. □<sub>4.6</sub>

Since that it is easy to obtain the hypothesis in Lemma 4.6 that  $\mathbf{K}$  is nice for unions by the transformation in Lemma 4.2, we have shown the first part of the following theorem.

**Theorem 4.7** *To any AEC  $(\mathbf{K}, \prec_{\mathbf{K}})$  which admits intersections, we can assign a  $(\mathbf{K}', \prec_{\mathbf{K}'})$  which has the amalgamation property but so that if  $\mathbf{K}$  is not  $(\delta, \lambda)$ -local then  $\mathbf{K}'$  is not  $(\delta, \lambda)$ -local.*

Proof. We now show the ‘but’. Let  $|M| = \lambda$  and suppose  $p, q \in \text{ga} - \text{S}(M)$ ,  $M = \bigcup_{i < \delta} M_i$ ,  $p \neq q$  and  $M, M_i \in \mathbf{K}$ . Let  $M^1, M^2$  be  $\mathbf{K}$ -extensions of  $M$  and let  $a \in M_1$  realize  $p$  and  $b \in M^2$  realize  $q$ . Now consider the models  $\overline{M}, \overline{M}_1, \overline{M}_2$  in  $\mathbf{K}'$  obtained by adding a complete  $R$ -graph to  $M, M_1, M_2$ . Then for each  $i < \delta$ ,  $\text{tp}(a/\overline{M}_i, \overline{M}^1) = \text{tp}(b/\overline{M}_i, \overline{M}^2)$ . (Witness with the same maps just adding a complete  $R$ -graph.) But  $\text{tp}(a/\overline{M}, \overline{M}^1) \neq \text{tp}(b/\overline{M}, \overline{M}^2)$  since any  $\tau'$ -isomorphism taking  $a$  and  $b$  to the same point would restrict to a similar  $\tau$ -isomorphism. □<sub>4.7</sub>

As opposed to Lemma 4.2, this second transformation, although preserving non-tameness, plays havoc with the spectrum function. We spelled out the result for locality but non-tameness is preserved in the same way.

Recall that we use the adjective ‘weakly’ for locality phenomena where the larger model is required to be saturated. This is an intriguing issue as in [She99] only weak tameness rather than tameness is deduced from categoricity. The construction at hand does not appear to distinguish ‘weak’. More precisely, we now show that at least for regular  $\lambda$ , if  $\mathbf{K}'$  is not  $(\lambda, \lambda)$ -local then it is not weakly  $(\lambda, \lambda)$ -local.

**Definition 4.8** *For any property  $P$  which can hold of models, we say that  $(\kappa, \lambda)$ -almost all models of  $\mathbf{K}$  satisfy  $P$  if Player II has a winning strategy for the following game. The game lasts  $\kappa$  moves. At each stage each player must*

choose a model of cardinality  $\lambda$  extending all the preceding models in the chain. Player II wins if the union satisfies  $P$ .

Note that if two properties are satisfied by  $(\kappa, \lambda)$ -almost all  $M$ , then  $(\kappa, \lambda)$ -almost all  $M$  satisfy both of them. Also when  $\lambda$  is regular, if there is a saturated model in power  $\lambda$ ,  $(\lambda, \lambda)$ -almost all  $M$  are saturated.

We say a model is, e.g.  $(\kappa, \lambda)$ -compact if every union of types over a decomposition of the model has a limit.

**Lemma 4.9** *Let  $\mathbf{K}$  and  $\mathbf{K}'$  be as in Definition 4.5.*

1. *If  $\mathbf{K}$  is not  $(\kappa, \lambda)$ -compact then  $(\kappa, \lambda)$ -almost all models of  $\mathbf{K}'$  are not  $(\kappa, \lambda)$ -compact.*
2. *If  $\mathbf{K}$  is not  $(\kappa, \lambda)$ -local then  $(\kappa, \lambda)$ -almost all models of  $\mathbf{K}'$  are not  $(\kappa, \lambda)$ -local.*

Proof. First we consider compactness. Let  $\langle M_i, p_i \rangle$  for  $i < \kappa$  be a continuous increasing sequence of  $\tau$ -structures that witnesses the incompactness. Expand  $M_\kappa$  to a  $\tau'$  structure by making it an  $R$ -set. Let  $\langle N_i \rangle$  for  $i < \kappa$ , be player I's moves. At stage  $\alpha$ , let player II choose for his move  $M'_\alpha$  so that  $N_\alpha \prec_{\mathbf{K}'} M'_\alpha$  and  $M_\alpha \prec_{\mathbf{K}} M'_\alpha \upharpoonright \tau$ . Let  $p_\alpha = \text{tp}(a_\alpha/M_\alpha, L_\alpha)$ . Let  $L'_\alpha$  be an amalgam of the expansion of  $L_\alpha$  to an  $R$ -set with  $M'_\alpha$  (over  $M'_{\alpha-1}$  at successors; take unions at limits.); let  $p'_\alpha = \text{tp}(a_\alpha/M'_\alpha, L'_\alpha)$ . Now there is no realization of  $\bigcup p'_\alpha$  since it would reduce to a realization of  $\bigcup p_\alpha$ . So Player II wins the game that asks each model to be extended to one which witnesses incompactness. For locality, do the same argument but choose  $L$  so that there are  $a, b \in L$  such that  $\text{tp}(a/M_i, L) = \text{tp}(b/M_i, L)$  for each  $i < \kappa$  but  $\text{tp}(a/M, L) \neq \text{tp}(b/M, L)$  and finish as in Theorem 4.7.  $\square_{4.9}$

It is immediate from Lemma 4.9 and the remark before it that

**Corollary 4.10** *If  $\mathbf{K}$  is not  $(\kappa, \lambda)$ -compact then  $\mathbf{K}'$  is not weakly- $(\kappa, \lambda)$ -compact.*

## 5 Gaining tameness

We gave in Section 2 an example of an AEC with the amalgamation property and Lowenheim-Skolem number  $\aleph_0$  which is not  $\aleph_0$ -tame. But at least consistently there are arbitrarily large  $\kappa$  for which it is not  $(\kappa, \infty)$ -tame. Here, we respond to a question of Grossberg and VanDieren [GV06b] and provide an example of an AEC with the amalgamation property and Lowenheim-Skolem number  $\aleph_0$  which is not  $\aleph_0$ -tame but is  $(2^{\aleph_0}, \infty)$ -tame. The example is very close to that in Section 2 but we bound the size of the image group  $G$ .

Now we use the moreover clause from Fact 2.2: There is a countable subgroup  $R$  of  $G$  such that every element of  $G/R$  is divisible by every prime. See [EM90].

**Example 5.1** Let  $\mathbf{K}^s$  be the class of structures  $M = \langle G, Z, I, H, R \rangle$ , where each of the listed sets is the solution set of one of the unary predicates  $(\mathbf{G}, \mathbf{Z}, \mathbf{I}, \mathbf{H}, \mathbf{R})$ . The first four predicates are interpreted exactly as in Example 2.3 but  $\mathbf{R}$  is interpreted as the subgroup  $R$  described above. Crucially, we require that the group  $G$  be not merely torsion-free but  $\aleph_1$ -free. The notion of strong submodel is as before except in addition  $M \prec_{\mathbf{K}^s} N$  implies  $\mathbf{R}^M = \mathbf{R}^N$ .

**Lemma 5.2** *The class  $(\mathbf{K}^s, \prec_{\mathbf{K}^s})$  defined in Definition 5.1 is an AEC which admits intersections.*

Proof. To show that  $(\mathbf{K}^s, \prec_{\mathbf{K}^s})$  is closed under unions of chains (the interesting case is countable unions) apply Pontryagin’s criterion. We can construct closures exactly as in Lemma 2.5.  $\square_{5.2}$

Since we have required that  $G$  is  $\aleph_1$ -free, the amalgamation property is no longer true. (The amalgamation of torsion-free groups is torsion-free. But, Shelah had observed long ago that even under pure embedding the class of  $\aleph_1$ -free groups does not have the amalgamation property. The argument will appear in [BCG<sup>+</sup>00].) But we are saved by Section 4.

**Theorem 5.3** *There is an AEC with the amalgamation property in a countable language with Lowenheim-Skolem number  $\aleph_0$  which is not  $(\aleph_0, \aleph_1)$ -tame but is  $(2^{\aleph_0}, \infty)$ -tame.*

Proof. Since  $\mathbf{K}^s$  (Example 5.1) admits intersections we can get the desired example with the amalgamation property from Corollary 4.7, provided we show Example 5.1 is  $(2^{\aleph_0}, \infty)$ -tame. Since the source of non-tameness is types over the target group  $G$ , it suffices to show the cardinality of  $\mathbf{G}^M$  is at most the continuum for any  $M \in \mathbf{K}^s$ . But if  $G$  is torsion free and for some countable subgroup  $R$ ,  $G/R$  is  $p$ -divisible for all  $p$ , then  $|G| \leq 2^{\aleph_0}$ . For, if  $G$  is larger there exist  $x, y \in G$  which realize the same first order type over  $R$ . Thus for each  $p$  there is an  $x_1$  such that  $px_1 - x = r \in R$  and a  $y_1$  such that  $py_1 - y = r \in R$ . But then  $x - y$  is a non-zero element of  $G$  which is divisible by every prime. By Pontryagin’s criterion, there is no such element in an  $\aleph_1$ -free group.  $\square_{5.3}$

## 6 Conclusion

This paper has several messages. The notion of an AEC admitting intersections is rather natural; it has come up without being named in investigations of the Hrushovski construction. It simplifies the treatment of Galois-types while being much weaker than the amalgamation property; we think it deserves further investigation.

We have shown that ‘locality’ has several facets. There has been considerable work on categoricity transfer for tame AEC [GV06a, GV06c, Les05]; under further locality assumptions [HK06] begins a ‘geometric stability’ theory. This paper shows that in general these are real assumptions. But could  $(\text{LS}(\mathbf{K}), \infty)$ -tameness be a consequence of categoricity? Let  $\kappa$  be the Löwenheim-Skolem

number of  $\mathbf{K}$  and let  $H_1$  denote  $\beth_{(2^{\aleph_1})}^+$ . Analysis of the Hart-Shelah examples [BK] (done after this paper) gave examples of an AEC  $\mathbf{K}$  with the amalgamation property which is categorical in small cardinals and fails tameness in a cardinality. In contrast, the next question should be quite hard. Shelah proved (see [Bal00] for a short account):

**Theorem 6.1** [She99] *Suppose  $\mathbf{K}$  has the amalgamation property and arbitrarily large models. Suppose  $\mathbf{K}$  is  $\lambda^+$ -categorical with  $\lambda > H_1$ . For each  $\mu \leq \lambda$ ,  $\mathbf{K}$  is  $(\chi, \mu)$ -weakly tame for some  $\chi < H_1$ .*

**Question 6.2** *Suppose  $\mathbf{K}$  has the amalgamation property and arbitrarily large models. Suppose  $\mathbf{K}$  is  $\lambda^+$  categorical with  $\lambda > H_1$ .*

*Is there any way to reduce the upper bound on  $\chi$  in Theorem 6.1?*

*Can one prove  $\mathbf{K}$  is  $(\chi, \mu)$ -weakly tame for some  $\chi < H_1$ ?*

Here are questions which naturally arise in extending this work.

**Problem 6.3** *Find an example of non-compactness in ZFC.*

**Question 6.4** *Are the examples in Section 3 (Section 5),  $(\aleph_1, \aleph_1)$ -compact?*

**Question 6.5** *Find examples of  $\mathbf{K}$  with the amalgamation property which gain tameness as in Section 5 but for inherent algebraic reasons rather than through the transformation of Section 4.*

## References

- [Bal00] J.T. Baldwin. Categoricity. Available at [www.math.uic.edu/~jbaldwin/](http://www.math.uic.edu/~jbaldwin/), 200?
- [BCG<sup>+</sup>00] J. Baldwin, W. Calvert, J. Goodrick, A. Villaveces, and A. Walczak-Typke. Abelian groups as aec's. preprint, 200?
- [BK] John T. Baldwin and Alexei Kolesnikov. Categoricity, amalgamation, and tameness. preprint: [www.math.uic.edu/~jbaldwin/](http://www.math.uic.edu/~jbaldwin/), to appear in Israel Journal of Mathematics.
- [BKV06] J.T. Baldwin, D.W. Kueker, and M. VanDieren. Upward stability transfer theorem for tame abstract elementary classes. *Notre Dame Journal of Formal Logic*, 47:291–298, 2006.
- [EM90] P. Eklof and Alan Mekler. *Almost Free Modules: Set theoretic Methods*. North Holland, 1990.
- [GK] Rami Grossberg and Alexei Kolesnikov. Excellent abstract elementary classes are tame. preprint.
- [Gro02] Rami Grossberg. Classification theory for non-elementary classes. In Yi Zhang, editor, *Logic and Algebra*, pages 165–204. AMS, 2002. Contemporary Mathematics 302.

- [GV06a] R. Grossberg and M. VanDieren. Categoricity from one successor cardinal in tame abstract elementary classes. *The Journal of Mathematical Logic*, 6:181–201, 2006.
- [GV06b] R. Grossberg and M. VanDieren. Galois stability for tame abstract elementary classes. *Journal of Mathematical Logic*, 6:1–24, 2006.
- [GV06c] R. Grossberg and M. VanDieren. Shelah’s categoricity conjecture from a successor for tame abstract elementary classes. *The Journal of Symbolic Logic*, 71:553–568, 2006.
- [HK06] T. Hyttinen and M. Kesälä. Independence in finitary abstract elementary classes. *Ann. Pure Appl. Logic*, 143(1-3):103–138, 2006.
- [Les05] Olivier Lessmann. Upward categoricity from a successor cardinal for an abstract elementary class with amalgamation. *Journal of Symbolic Logic*, 70:639–661, 2005.
- [Rab62] Michael Rabin. Classes of structures and sets of sentences with the intersection property. In *Actes du Colloque de mathématiques a l’Occasion de Tricentenaire de la mort de B. Pascal*, pages 39–53. Université de Clermont, 1962.
- [She74] S. Shelah. Infinite abelian groups, whitehead problem and some constructions. *Israel Journal of Mathematics*, 18:243–256, 1974.
- [She87] Saharon Shelah. Classification of nonelementary classes II, abstract elementary classes. In J.T. Baldwin, editor, *Classification theory (Chicago, IL, 1985)*, pages 419–497. Springer, Berlin, 1987. paper 88: Proceedings of the USA–Israel Conference on Classification Theory, Chicago, December 1985; volume 1292 of *Lecture Notes in Mathematics*.
- [She99] S. Shelah. Categoricity for abstract classes with amalgamation. *Annals of Pure and Applied Logic*, 98:261–294, 1999. paper 394. Consult Shelah for post-publication revisions.
- [She01] S. Shelah. Categoricity of abstract elementary class in two successive cardinals. *Israel Journal of Mathematics*, 126:29–128, 2001. paper 576.