

Topological Partition Relations of the Form $\omega^* \rightarrow (Y)_2^1$

W. W. Comfort ^{a,b,c}, Wesleyan University (USA)

Akio Kato, National Defense Academy (Japan)

Saharon Shelah ^d, Hebrew University (Israel)

ABSTRACT

Theorem. The topological partition relation $\omega^* \rightarrow (Y)_2^1$

- (a) fails for every space Y with $|Y| \geq 2^{\mathfrak{c}}$;
- (b) holds for Y discrete if and only if $|Y| \leq \mathfrak{c}$;
- (c) holds for certain non-discrete P -spaces Y ;
- (d) fails for $Y = \omega \cup \{p\}$ with $p \in \omega^*$;
- (e) fails for Y infinite and countably compact.

AMS Classification Numbers: Primary 54A25, 54D40, 05A17; Secondary 54A10, 54D30, 04A20

Key words and phrases: Topological Ramsey theory; topological partition relation; Stone-Čech remainder; P -space; countably compact space.

[Footnotes]

^a Presented at the Madison Conference by this co-author.

^b Member, New York Academy of Sciences.

^c This author gratefully acknowledges support received from the Technische Hochschule Darmstadt and from the Deutscher Akademischer Austauschdienst (=DAAD) of the Federal Republic of Germany.

^d This author acknowledges partial support received from the Basic Research Fund of the Israeli Academy of Sciences, Publ. 440.

§1. Introduction.

For topological space X and Y we write $X \approx Y$ if X and Y are homeomorphic, and we write $f : X \approx Y$ if f is a homeomorphism of X onto Y . The “topological inclusion relation” is denoted by \subseteq_h ; that is, we write $Y \subseteq_h X$ if there is $Y' \subseteq Y$ such that $Y \approx Y'$.

The symbol ω denotes both the least infinite cardinal and the countably infinite discrete space; the Stone-Čech remainder $\beta(\omega) \setminus \omega$ is denoted ω^* .

For a space X we denote by wX and dX the weight and density character of X , respectively. Following [7], for $A \subseteq \omega$ we write $A^* = (\text{cl}_{\beta(\omega)} A) \setminus \omega$.

For proofs of the following statements, and for other basic information on topological and combinatorial properties of the space ω^* , see [7], [3], [12].

1.1. Theorem. (a) $\{\text{cl}_{\beta(\omega)} A : A \subseteq \omega\}$ is a basis for the open sets of $\beta(\omega)$; thus $w(\beta(\omega)) = \mathfrak{c}$.

(b) There is an (almost disjoint) family \mathcal{A} of subsets of ω such that $|\mathcal{A}| = \mathfrak{c}$ and $\{A^* : A \in \mathcal{A}\}$ is pairwise disjoint.

(c) ω^* contains a family of $2^{\mathfrak{c}}$ -many pairwise disjoint copies of $\beta(\omega)$.

(d) Every infinite, closed subspace Y of ω^* contains a copy of $\beta(\omega)$, so $|Y| = |\beta(\omega)| = 2^{\mathfrak{c}}$. \square

For cardinals κ and λ and topological spaces X and Y , the symbol $X \rightarrow (Y)_{\lambda}^{\kappa}$ means that if the set $[X]^{\kappa}$ of all κ -membered subsets of X is written in the form $[X]^{\kappa} = \cup_{i < \lambda} P_i$, then there are $i < \lambda$ and $Y' \subseteq X$ such that $Y \approx Y'$ and $[Y']^{\kappa} \subseteq P_i$. Our present primary interest is in topological arrow relations of the form $X \rightarrow (Y)_2^1$ (with $X = \omega^*$). For spaces X and Y , the relation $X \rightarrow (Y)_2^1$ reduces to this: if $X = P_0 \cup P_1$, then either $Y \subseteq_h P_0$ or $Y \subseteq_h P_1$.

The relation $X \rightarrow (Y_0, Y_1)^1$ indicates that if $X = P_0 \cup P_1$, then either $Y_0 \subseteq_h P_0$ or $Y_1 \subseteq_h P_1$.

It is obvious that if X and Y are spaces such that $Y \subseteq_h X$ fails, then $X \rightarrow (Y)_2^1$

fails.

By way of introduction it is enough here to observe that the classical theorem of F. Bernstein, according to which there is a subset S of the real line \mathbf{R} such that neither S nor its complement $\mathbf{R} \setminus S$ contains an uncountable closed set, is captured by the assertion that the relation $\mathbf{R} \rightarrow (\{0, 1\}^\omega)_2^1$ fails; in the positive direction, it is easy to see that the relation $\mathbf{Q} \rightarrow (\mathbf{Q})_2^1$ holds for \mathbf{Q} the space of rationals.

For a report on the present-day “state of the art” concerning topological partition relations, and for references to the literature and open questions, the reader may consult [14], [15], [16].

This paper is organized as follows. §2 shows that $\omega^* \rightarrow (Y)_2^1$ fails for every infinite compact space Y . §3 characterizes those discrete spaces Y for which $\omega^* \rightarrow (Y)_2^1$, and §4 shows that $\omega^* \rightarrow (Y)_2^1$ holds for certain non-discrete spaces Y . §5 shows that $\omega^* \rightarrow (Y)_2^1$ fails for spaces of the form $Y = \omega \cup \{p\}$ with $p \in \omega^*$, hence fails for every infinite countably compact space Y . The results of §§2–5 prompt several questions, and these are given in §6.

We are grateful to Jan van Mill, K. P. S. Bhaskara Rao, and W. A. R. Weiss for helpful conversations.

We announced some of our results in the abstract [2]. See also [1] for related results.

§2. $\omega^* \not\rightarrow (Y)_2^1$ for $|Y| \geq 2^{\mathfrak{c}}$.

2.1. Lemma. If $Y \subseteq_h \omega^*$ then $|\{A \subseteq \omega^* : A \approx Y\}| = 2^{\mathfrak{c}}$.

Proof. The inequality \geq is immediate from Theorem 1.1(c). For \leq , it is enough to fix (a copy of) $Y \subset \omega^*$ and to notice that since $dY \leq wY \leq w(\omega^*) = \mathfrak{c}$ (by Theorem 1.1(a)), the number of continuous functions from Y into ω^* does not exceed $|(\omega^*)^{dY}| \leq (2^{\mathfrak{c}})^{\mathfrak{c}} = 2^{\mathfrak{c}}$. \square

2.2. Theorem. If Y is a space such that $|Y| \geq 2^{\mathfrak{c}}$, then $\omega^* \not\rightarrow (Y)_2^1$.

Proof. We assume $Y \subseteq_h \omega^*$ (in particular we assume $|Y| = |\omega^*| = 2^{\mathfrak{c}}$) since

otherwise $\omega^* \not\rightarrow (Y)_2^1$ is obvious. Following Lemma 2.1 let $\{A_\xi : \xi < 2^{\mathfrak{c}}\}$ enumerate $\{A \subseteq \omega^* : A \approx Y\}$, choose distinct $p_0, q_0 \in A_0$ and recursively, if $\xi < 2^{\mathfrak{c}}$ and p_η, q_η have been chosen for all $\eta < \xi$ choose distinct

$$p_\xi, q_\xi \in A_\xi \setminus (\{p_\eta : \eta < \xi\} \cup \{q_\eta : \eta < \xi\}).$$

It is then clear, writing

$$P_0 = \{p_\xi : \xi < 2^{\mathfrak{c}}\} \text{ and } P_1 = \omega^* \setminus P_0,$$

that the relations $Y \subseteq_h P_0$ and $Y \subseteq_h P_1$ both fail. \square

The following statement is an immediate consequence of Theorems 2.2 and 1.1(d).

2.3. Corollary. The relation $\omega^* \rightarrow (Y)_2^1$ fails for every infinite compact space Y . \square

By less elementary methods we strengthen Corollary 2.3 in Theorem 5.14 below.

§3. Concerning the Relation $\omega^* \rightarrow (Y)_2^1$ for Y Discrete.

The very simple result of this section, included in the interest of completeness, shows for discrete spaces Y that $\omega^* \rightarrow (Y)_2^1$ if and only if $Y \subseteq_h \omega^*$.

3.1. Theorem. For a discrete space Y , the following conditions are equivalent.

- (a) $|Y| \leq \mathfrak{c}$;
- (b) $\omega^* \rightarrow (Y)_{\mathfrak{c}}^1$;
- (c) $\omega^* \rightarrow (Y)_2^1$;
- (d) $Y \subseteq_h \omega^*$.

Proof. (a) \Rightarrow (b). [Here we profit from a suggestion offered by the referee.] Given $\omega^* = \cup_{i < \mathfrak{c}} P_i$, recall from [10](2.2) or [12](3.3.2) this theorem of Kunen: there is a matrix $\{A_i^\xi : \xi < \mathfrak{c}, i < \mathfrak{c}\}$ of clopen subsets of ω^* such that

- (i) for each $i < \mathfrak{c}$ the family $\{A_i^\xi : \xi < \mathfrak{c}\}$ is pairwise disjoint, and
- (ii) each $f \in \mathfrak{c}^{\mathfrak{c}}$ satisfies $\cap_{i < \mathfrak{c}} A_i^{f(i)} \neq \emptyset$.

Now if one of the sets P_i meets A_i^ξ for each $\xi < \mathfrak{c}$ (say $p_\xi \in A_i^\xi$) then the discrete set $D = \{p_\xi : \xi < \mathfrak{c}\}$ satisfies $Y \subseteq_h D \subseteq P_i$; otherwise for each $i < \mathfrak{c}$ there is $f(i)$ such that $P_i \cap A_i^{f(i)} = \emptyset$, so $\emptyset \neq \cap_{i < \mathfrak{c}} A_i^{f(i)} \subseteq \omega^* \setminus \cup_{i < \mathfrak{c}} P_i$.

That (b) \Rightarrow (c) and (c) \Rightarrow (d) and clear.

(d) \Rightarrow (a). Theorem 1.1(a) gives $|Y| = wY \leq w(\beta(\omega)) = \mathbf{c}$.

§4. $\omega^* \rightarrow (Y)_2^1$ for Certain Non-Discrete Y .

For an infinite cardinal κ we denote by P_κ the ordinal space $\kappa + 1 = \kappa \cup \{\kappa\}$ topologized to be “discrete below κ ” and with a neighborhood base at κ the same as in the usual interval topology. That is, a subset U of $\kappa + 1$ is open in P_κ if and only if either $U \subseteq \kappa$ or some $\xi < \kappa$ satisfies $(\xi, \kappa] \subseteq U$.

4.1. Theorem. For cardinals $\kappa \geq \omega$ and $m_0, m_1 < \omega$, the space P_κ satisfies $P_\kappa^{m_0+m_1} \rightarrow (P_\kappa^{m_0}, P_\kappa^{m_1})^1$.

Proof. Let $P^I = X_0 \cup X_1$ and $|I| = m_0 + m_1$ and suppose without loss of generality that the point $c = \langle c_i \rangle_{i \in I}$ with $c_i = \kappa$ (all $i \in I$) satisfies $c \in X_0$. Let $I = I_0 \cup I_1$ with $|I_0| = m_0$, $|I_1| = m_1$, and set $D = P_\kappa \setminus \{\kappa\}$, and for $x \in D^{I_0}$ define

$$S(x) = \{x\} \times \{y \in P_\kappa^{I_1} : \max\{x_i : i \in I_0\} < \min\{y_i : i \in I_1\}\}.$$

If some $x \in D^{I_0}$ satisfies $S(x) \subseteq X_1$ we have $P_\kappa^{m_1} \approx S(x) \subseteq X_1$ and the proof is complete. Otherwise for each $x \in D^{I_0}$ there is $p(x) \in S(x) \cap X_0$ and then

$$P_\kappa^{m_0} \approx \{p(x) : x \in D^{I_0}\} \cup \{c\} \subseteq X_0,$$

as required. \square

4.2. Corollary. Every infinite cardinal κ satisfies $P_\kappa \times P_\kappa \rightarrow (P_\kappa)_2^1$. \square

We say as usual that a topological space $X = \langle X, \mathcal{T} \rangle$ is a P -space if each $\mathcal{U} \subseteq \mathcal{T}$ with $|\mathcal{U}| \leq \omega$ satisfies $\bigcap \mathcal{U} \in \mathcal{T}$. Since (clearly) P_κ is a non-discrete P -space if and only if $\text{cf}(\kappa) > \omega$, the following theorem shows the existence of a nondiscrete Y such that $X \rightarrow (Y)_2^1$.

4.3. Theorem. Let $\omega_1 \leq \kappa \leq \mathbf{c}$ satisfy $\text{cf}(\kappa) > \omega$. Then $\omega^* \rightarrow (P_\kappa)_2^1$.

Proof. It is a theorem of E. K. van Douwen that every P -space X such that $wX \leq \mathbf{c}$ satisfies $X \subseteq_h \omega^*$. (For a proof of this result see [4] or [12].) Thus for κ as hypothesized we have $P_\kappa \times P_\kappa \subseteq_h \omega^*$, so the relation $\omega^* \rightarrow (P_\kappa)_2^1$ is immediate from Corollary 4.2. \square

4.4. Remarks. (a) The following simple result, suggested by the proof of Theorem 4.2, is peripheral to the principal thrust of our paper. Here as usual for a space $X = \langle X, \mathcal{T} \rangle$ we denote by $PX = \langle PX, P\mathcal{T} \rangle$ the set X with the smallest topology $P\mathcal{T}$ such that $P\mathcal{T} \supseteq \mathcal{T}$ and PX is a P -space; thus $\{\cap \mathcal{U} : \mathcal{U} \subseteq \mathcal{T}, |\mathcal{U}| \leq \omega\}$ is a base for $P\mathcal{T}$.

Theorem. For a P -space Y , the following conditions are equivalent.

- (i) $\omega^* \rightarrow (Y)_2^1$;
- (ii) $\{0, 1\}^c \rightarrow (Y)_2^1$;
- (iii) $P(\omega^*) \rightarrow (Y)_2^1$;
- (iv) $P(\{0, 1\}^c) \rightarrow (Y)_2^1$.

Proof. The implications (iii) \Rightarrow (iv) \Rightarrow (i) \Rightarrow (ii) follow respectively from the inclusions $P(\omega^*) \subseteq_h P(\{0, 1\}^c) \subseteq_h \omega^* \subseteq_h \{0, 1\}^c$. (Of these three inclusions the third follows from Theorem 1.1, the first from the third, and the second from van Douwen's theorem cited above.) That (ii) \Rightarrow (iii) follows from $P(\{0, 1\}^c) \subseteq_h \omega^*$ (whence $P(\{0, 1\}^c) \subseteq_h P(\omega^*)$) and the case $A = \{0, 1\}^c$, $B = Y = PY$ of this general observation: if $A \rightarrow (B)_2^1$ then $PA \rightarrow (PB)_2^1$. \square

(b) We note in passing the following result, from which (with 4.1) it follows that for $\kappa \geq \omega$ the space P_κ satisfies $P_\kappa^{2^n} \rightarrow (P_\kappa)_{n+1}^1$.

Theorem. Let S be a space such that $S^{m_0+m_1} \rightarrow (S^{m_0}, S^{m_1})^1$ for $m_0, m_1 < \omega$. Then $S^{2^n} \rightarrow (S)_{n+1}^1$ for $n < \omega$. (*)

Proof. Statement (*) is trivial when $n = 0$, and is given by the case $m_0 = m_1 = 1$ of the hypothesis when $n = 1$.

Now suppose (*) holds for $n = k$, and let $S^{2^{k+1}} = \cap_{i=0}^{k+1} X_i$. With $Y_0 = X_0$ and $Y_1 = \cup_{i=1}^{k+1} X_i$, it follows from $S^{2^k+2^k} \rightarrow (S^{2^k}, S^{2^k})$ that there is $T \subseteq S^{2^{k+1}}$ such that $T \approx S^{2^k}$ and either $T \subseteq Y_0$ or $T \subseteq Y_1$. In the first case we have $S \subseteq_h T \subseteq X_0$, and in the second case from $T \subseteq \cup_{i=1}^{k+1} X_i$ and (*) at k there exists i such that $1 \leq i \leq k+1$ and $S \subseteq_h X_i$, as required. \square

(c) The method of proof of 4.1 and 4.2 applies to many spaces other than those

of the form P_κ . The reader may easily verify, for example, denoting by C_κ the one-point compactification of the discrete space κ , that $C_\kappa \times C_\kappa \rightarrow (C_\kappa)_2^1$, and hence $\{0, 1\}^\kappa \rightarrow (C_\kappa)_2^1$, for all $\kappa \geq \omega$. For a proof due to S. Todorćević of a much stronger topological partition relation, namely $\{0, 1\}^\kappa \rightarrow (C_\kappa)_{cf(\kappa)}^1$, see Weiss [15].

§5. $\omega^* \not\rightarrow (Y)_2^1$ for Y Infinite and Countably Compact.

To prove this result, we show first that the relation $\omega^* \rightarrow (\omega \cup \{p\})_2^1$ fails for every $p \in \omega^*$. While this can be proved directly by combinatorial arguments, we find it convenient (given $p \in \omega^*$) to introduce and use as a tool a new topology $\mathcal{T}(p)$ on ω^* .

Given $f : \omega \rightarrow \omega^*$, we denote by $\bar{f} : \beta(\omega) \rightarrow \omega^*$ the Stone extension of f . For $X \subseteq \omega^*$ we set

$$X^p = X \cup \{\bar{f}(p) : f : \omega \approx f[\omega] \subseteq X\};$$

that is, X^p is X together with its “ p -limits through discrete countable sets.”

5.1. Lemma. There is a topology $\mathcal{T}(p)$ for ω^* such that each $X \subseteq \omega^*$ satisfies: X is $\mathcal{T}(p)$ -closed if and only if $X = X^p$.

Proof. It is enough to show

- (a) $\emptyset = \emptyset^p$;
- (b) $\omega^* = (\omega^*)^p$;
- (c) $X_0 \cup X_1 = (X_0 \cup X_1)^p$ if $X_i = X_i^p$ ($i = 0, 1$) and
- (d) $\bigcap_{i \in I} X_i = (\bigcap_{i \in I} X_i)^p$ if each X_i satisfies $X_i = X_i^p$.

Now (a) and (b) are obvious, as are the inclusions \subseteq of (c) and (d).

(c) (\supseteq) If $f : \omega \approx f[\omega] \subseteq X_0 \cup X_1$ satisfies $\bar{f}(p) = x \in (X_0 \cup X_1)^p$ then with $A_i = \{n < \omega : f(n) \in X_i\}$ we have $A_0 \cup A_1 \in p$ and hence $A_{\bar{i}} \in p$ for suitable $\bar{i} \in \{0, 1\}$; changing the values of f on $\omega \setminus A_{\bar{i}}$ if necessary (to ensure $f[\omega] \subseteq A_{\bar{i}}$), we conclude that $x = \bar{f}(p) \in X_{\bar{i}}^p = X_{\bar{i}} \subseteq X_0 \cup X_1$.

(d) (\supseteq). If $x = \bar{f}(p)$ with $f : \omega \approx f[\omega] \subseteq \bigcap_i X_i$ then $x \in \bigcap_i (X_i^p) = \bigcap_i X_i$. \square

5.2. Remarks. (a) In the terminology of Lemma 5.1, the topology $\mathcal{T}(p)$ is defined by the relation

$$\mathcal{T}(p) = \{\omega^* \setminus X : X \subseteq \omega^*, X \text{ is } \mathcal{T}(p)\text{-closed}\}.$$

(b) For notational convenience we denote by $I(p)$ the set of $\mathcal{T}(p)$ -isolated points of ω^* , and we write $A(p) = \omega^* \setminus I(p)$. Clearly $x \in I(p)$ if and only if x is not a “discrete limit” of points in $\omega^* \setminus \{x\}$, that is, if and only if every $f : \omega \approx f[\omega] \subseteq \omega^* \setminus \{x\}$ satisfies $\bar{f}(p) \neq x$. The fact that $I(p) \neq \emptyset$ has been known for many years. Indeed, Kunen [10] has shown that there exist 2^c -many points $x \in \omega^*$ such that $x \notin \text{cl}_{\beta(\omega)} A$ whenever $A \subseteq \omega^* \setminus \{x\}$ and $|A| \leq \omega$. (These are the so-called weak- P -points of ω^* .)

As a mnemonic device one may think of $A(p)$ and $I(p)$ as the sets of p -accessible and p -inaccessible points, respectively.

(c) For $X \subseteq \omega^*$ the set X^p may fail to be closed. Indeed, the $\mathcal{T}(p)$ -closure of $X \subseteq \omega^*$ is determined by the following iterative procedure (cf. also [1]).

5.3. Lemma. Let $X \subseteq \omega^*$. For $\xi \leq \omega^+$ define X_ξ by :

$$X_0 = X;$$

$$X_\xi = \cup_{\eta < \xi} X_\eta \text{ if } \xi \text{ is a limit ordinal};$$

$$X_{\xi+1} = X_\xi^p.$$

Then $X_{\omega^+} = \mathcal{T}(p) - \text{cl } X$. \square

The following fact, noted in [8], [5], [6], is crucial to many studies of ω^* (see also [3](16.13) for a proof). One may capture the thrust of this lemma by paraphrasing the picturesque terminology of Frolík [6]: “No type produces itself.”

5.4. Lemma. No homeomorphism from $\beta(\omega)$ into ω^* has a fixed point. \square

5.5. Lemma. Let A and B be countable, discrete subsets of ω^* , with $A \subseteq B^*$.

Then $A^p \cap B^p = \emptyset$.

Proof. If $x \in A^p \cap B^p$ we may suppose without loss of generality that there are $f : \omega \approx A$ and $g : \omega \approx B$ such that $x = \bar{f}(p) = \bar{g}(p)$. The function

$h = f \circ g^{-1} : B \approx A \subseteq B^*$ satisfies

$$\bar{f} \circ \bar{g}^{-1} = \bar{h} : \beta(B) \approx \beta(A) \subseteq B^*$$

and $\bar{h}(x) = x \in B^*$, contrary to Lemma 5.4. \square

5.6. Corollary. Let A and B be countably infinite, discrete subsets of ω^* such that $A \cap B = \emptyset$. Then $A^p \cap B^p = \emptyset$.

Proof. Let $x \in A^p \cap B^p$ and let $f : \omega \rightarrow f[\omega] \subseteq A$ and $g : \omega \rightarrow g[\omega] \subseteq B$ satisfy $x = \bar{f}(p) = \bar{g}(p)$. Leaving f and g unchanged on suitably chosen elements of p , but making modifications elsewhere if necessary, we assume without loss of generality that either $f[\omega] \subseteq (g[\omega])^*$ or $g[\omega] \subseteq (f[\omega])^*$ or $f[\omega] \cap (g[\omega])^* = (f[\omega])^* \cap g[\omega] = \emptyset$. By Lemma 5.5 the first of these possibilities, and by symmetry the second, cannot occur. We conclude that $f[\omega] \cup g[\omega]$ is a countable, discrete subset of ω^* such that $f[\omega] \cap g[\omega] = \emptyset$; it follows that $(f[\omega])^* \cap (g[\omega])^* = \emptyset$, since every countable (discrete) subset of ω^* is C^* -embedded (cf. [7](14.27, 14N.5), [3](16.15)). This contradicts the relation $x \in (f[\omega])^* \cap (g[\omega])^*$. \square

5.7. Corollary. If $\omega^* \supseteq X \in \mathcal{T}(p)$, then $X^p \in \mathcal{T}(p)$.

Proof. If $\omega^* \setminus X^p$ is not $\mathcal{T}(p)$ -closed then there is $f : \omega \approx f[\omega] = A \subseteq \omega^* \setminus X^p$ such that $x = \bar{f}(p) \in X^p$. Since $X \in \mathcal{T}(p)$ we have $x \in X^p \setminus X$ so there is $g : \omega \approx g[\omega] = B \subseteq X$ such that $x = \bar{g}(p)$. From $A \cap B = \emptyset$ and 5.6 now follows $x \in A^p \cap B^p = \emptyset$, a contradiction. \square

5.8. Corollary. If $\omega^* \supseteq X \in \mathcal{T}(p)$ then $\mathcal{T}(p) - \text{cl } X \in \mathcal{T}(p)$.

Proof. This is immediate from 5.3 and 5.7. \square

Our goal is to 2-color the points of ω^* in such a way that every copy of $\omega \cup \{p\}$ receives two colors. First we consider how to extend a given coloring function.

5.9. Lemma. Let $\omega^* \supseteq X \in \mathcal{T}(p)$ and let $c : X \rightarrow 2 = \{0, 1\}$ be a function with no monochromatic copy of $\omega \cup \{p\}$ (that is, if $X \supseteq Y \approx \omega \cup \{p\}$ then $c^{-1}(\{i\}) \cap Y \neq \emptyset$ for $i \in \{0, 1\}$). Then c extends to $\tilde{c} : X^p \rightarrow 2$ with no monochromatic copy of $\omega \cup \{p\}$.

Proof. Set $X_i = c^{-1}(\{i\})$ for $i \in 2 = \{0, 1\}$, so that $X^p = X_0^p \cup X_1^p$ by 5.1(c) and $(X_0^p \setminus X) \cap (X_1^p \setminus X) = \emptyset$

by 5.6. Since $\{X, X_0^p \setminus X, X_1^p \setminus X\}$ is a partition of X , the function $\tilde{c} : X^p \rightarrow 2$, given by the rule

$$\begin{aligned}\tilde{c}(x) &= c(x) \text{ if } x \in X \\ &= 1 \text{ if } x \in X_0^p \setminus X \\ &= 0 \text{ if } x \in X_1^p \setminus X,\end{aligned}$$

is well-defined. To see that \tilde{c} is as required let $h : \omega \cup \{p\} \approx A \cup \{x\} \subseteq X^p$ with $h : \omega \approx A$, $h(p) = x$. Modifying h (as before) if necessary, we assume without loss of generality that either (i) $A \subseteq X_0$ or (ii) $A \subseteq X_0 \setminus X$ (the cases $A \subseteq X_1$, $A \subseteq X_1^p \setminus X$ are treated symmetrically). In case (i) we have $\tilde{c} \equiv 0$ on A and $\tilde{c}(x) = 1$ (since either $x \in X$ or $x \in X_0^p \setminus X$); case (ii) cannot arise, since $x \in X$ violates $X \in \mathcal{T}(p)$ while $x \in X^p \setminus X$ violates Corollary 5.6. \square

Combining Lemmas 5.9 and 5.3 yields this.

5.10. Lemma. Let $\omega^* \supseteq X \in \mathcal{T}(p)$ and let $c : X \rightarrow \{0, 1\}$ be a function with no monochromatic copy of $\omega \cup \{p\}$. Then c extends to $\tilde{c} : \mathcal{T}(p) - \text{cl } X \rightarrow \{0, 1\}$ with no monochromatic copy of $\omega \cup \{p\}$. \square

The preceding lemma indicates how to extend a coloring function from $X \in \mathcal{T}(p)$ over $\mathcal{T}(p) - \text{cl } X$, but it remains to initiate the coloring procedure. For this purpose it is convenient to consider a particular base $\mathcal{S}(p)$ for the topology $\mathcal{T}(p)$. We call the elements of $\mathcal{S}(p)$ the *p-satellite sets*.

5.11. Definition. Let $x \in \omega^*$. A set $S = S(x)$ is a *p-satellite set* based at x if there are a tree $T \subseteq \omega^{<\omega} = \bigcup_{n < \omega} \omega^n$ (ordered by containment) and for $s \in T$ a point $x_s \in S$ and $U_s \subseteq \omega^*$ such that

- (i) U_s is open-and-closed in the usual topology of ω^* ;
- (ii) $x = x_\langle \rangle$ with $\langle \rangle$ the empty sequence;
- (iii) $U_\langle \rangle = \omega^*$;
- (iv) if $x_s \in S(x)$ and $x_s \in A(p)$ then: $\{x_{s \hat{\ } n} : n < \omega\}$ enumerates the range of a function f such that $f : \omega \approx f[\omega] \subseteq \omega^*$ with $\bar{f}(p) = x_s$, and $\{U_{s \hat{\ } n} : n < \omega\}$ is a pairwise disjoint family such that $x_{s \hat{\ } n} \in U_{s \hat{\ } n} \subseteq U_s$;
- (v) if $x_s \in S(x)$ and $x_s \in I(p)$ then s is a maximal node in T (and $x_{s \hat{\ } n}$, $U_{s \hat{\ } n}$ are

defined for no $n < \omega$).

5.12. Remark. It is not difficult to see that for every $x \in X \in \mathcal{T}(p)$ there is $S = S(x) \in \mathcal{S}(p)$ such that $x \in S \subseteq X$. (If $x \in I(p)$ one takes $S = \{x\}$; if $x_s \in S \cap X$ has been defined one uses (iv) and $X \in \mathcal{T}(p)$ to choose $x_{s \hat{\ } n} \in S \cap X$ if $x_s \in A(p)$.) That each of the sets $S(x)$ is $\mathcal{T}(p)$ -open is immediate from Corollary 5.6 above. It follows that $\mathcal{S}(p)$ is indeed a base for $\mathcal{T}(p)$.

5.13. Theorem. Every $p \in \omega^*$ satisfies $\omega^* \not\rightarrow (\omega \cup \{p\})_2^1$.

Proof. Let $\{S(x(i)) : i \in I\}$ be a maximal pairwise disjoint subfamily of $\mathcal{S}(p)$. For each $i \in I$ define $c_i : S(x(i)) \rightarrow 2$ by

$$\begin{aligned} c_i(x(i)_s) &= 0 \text{ if length of } s \text{ is even} \\ &= 1 \text{ if length of } s \text{ is odd.} \end{aligned}$$

It is clear from Corollary 5.6 that not only each function c_i on $S(x(i))$, but also the function

$$c = \cup_{i \in I} c_i : \cup_{i \in I} S(x(i)) \rightarrow 2,$$

is monochromatic on no copy of $\omega \cup \{p\}$. Since $\cup_{i \in I} S(x(i))$ is $\mathcal{T}(p)$ -open and $\mathcal{T}(p)$ -dense in ω^* , the desired result follows from Lemma 5.10. \square

5.14. Theorem. The relation $\omega^* \rightarrow (Y)_2^1$ fails for every infinite, countably compact space Y .

Proof. Given infinite $Y \subseteq \omega^*$ there is $f : \omega \approx f[\omega] \subseteq Y$, and if Y is countably compact there is $p \in \omega^*$ such that $\bar{f}(p) \in Y$. Since $f[\omega]$ is C^* -embedded in ω^* we have

$$\omega \cup \{p\} \approx f[\omega] \cup \{\bar{f}(p)\} \subseteq Y,$$

so $\omega^* \not\rightarrow (Y)_2^1$ follows from $\omega^* \not\rightarrow (\omega \cup \{p\})_2^1$. \square

5.15. Remarks. (a) We cite three facts which (taken together) show that the index set I used in the proof of Theorem 5.13 satisfies $|I| = 2^c$: (i) The set W of weak- P -points of ω^* introduced by Kunen [10] satisfies $|W| = 2^c$; (ii) each $S(x) \in \mathcal{S}(p)$ satisfies $|S(x)| \leq \omega$; (iii) $W \subseteq I(p)$, so $W \subseteq \cup_{i \in I} S(x(i))$.

(b) With no attempt at a complete topological classification, we note five

elementary properties enjoyed by each of our topologies $\mathcal{T}(p)$ on ω^* .

(i) $\mathcal{T}(p)$ refines the usual topology of ω^* , so $\mathcal{T}(p)$ is a Hausdorff topology.

(ii) $\mathcal{T}(p)$ has $2^{\mathfrak{c}}$ -many isolated points. (Indeed, we have noted already that the set W of weak- P -points satisfies $|W| = 2^{\mathfrak{c}}$ and $W \subseteq I(p)$.)

(iii) Since $\mathcal{S}(p)$ is a base for $\mathcal{T}(p)$ and each $S(x) \in \mathcal{S}(p)$ satisfies $|S(x)| \leq \omega$, the topology $\mathcal{T}(p)$ is locally countable.

(iv) From Theorem 1.1(b) it is easy to see that if $S(x) \in \mathcal{S}(p)$ and $|S(x)| = \omega$, then $|\mathcal{T}(p) - \text{cl } S(x)| = \mathfrak{c}$. Thus $\mathcal{T}(p)$ is not a regular topology for ω^* .

(v) According to Corollary 5.8, the $\mathcal{T}(p)$ -closure of each $\mathcal{T}(p)$ -open subset of ω^* is itself $\mathcal{T}(p)$ -open. Such a topology is said to be extremally disconnected.

(c) In our development of $\mathcal{T}(p)$ and its properties we did not introduce explicitly the Rudin-Frolík pre-order \sqsubseteq on ω^* (see [5], [6], or [13], or [3] for an expository treatment) since doing so does not appear to simplify the arguments. We note however (as in [1]) that the relation \sqsubseteq lies close to our work: For $x, p \in \omega^*$ one has $p \sqsubset x$ if and only if some $f : \omega \approx f[\omega] \subseteq \omega^*$ satisfies $\bar{f}(p) = x$. \square

§6. Questions.

Perhaps this paper is best viewed as establishing some boundary conditions which may help lead to a solution of the following ambitious general problem.

6.1. Problem. Characterize those spaces Y such that $\omega^* \rightarrow (Y)_2^1$. \square

There are P -spaces Y such that $|Y| = 2^{\mathfrak{c}}$ and $Y \subseteq_h \omega^*$. (For example, according to van Douwen's theorem cited above, one may take $Y = P(\omega^*)$.) According to Theorem 2.2, the relation $\omega^* \rightarrow (Y)_2^1$ fails for each such Y . This situation suggests the following question.

6.2. Question. Does $\omega^* \rightarrow (Y)_2^1$ for every P -space Y such that $Y \subseteq_h \omega^*$ and $|Y| < 2^{\mathfrak{c}}$? What if $|Y| = \mathfrak{c}$? \square

We have no example of a non- P -space Y such that $\omega^* \rightarrow (Y)_2^1$, so we are compelled to ask:

6.3. Question. If Y is a space such that $\omega^* \rightarrow (Y)_2^1$, must Y be a P -space? \square

For $|Y| = \omega$, Question 6.3 takes this simple form:

6.4. Question. If Y is a countable space such that $\omega^* \rightarrow (Y)_2^1$, must Y be discrete? \square

6.5. Remark. In connection with Question 6.4 it should be noted that there exists a countable, dense-in-itself subset C of ω^* such that every $x \in C$ satisfies

$$(*) \ x \notin \text{cl}_{\beta(\omega)} D \text{ whenever } D \text{ is discrete and } D \subseteq C \setminus \{x\}$$

(equivalently: $\omega \cup \{p\} \subseteq_h C$ fails for every $p \in \omega^*$). To find such C we follow the construction of van Mill [11](3.3, pp. 53-54). Let E be the absolute (i.e., the Gleason cover) of the Cantor set $\{0, 1\}^\omega$, let $\pi : E \rightarrow \{0, 1\}^\omega$ be perfect and irreducible, and embed E into ω^* as a \mathbf{c} -OK set; then every countable $F \subseteq \omega^* \setminus E$ satisfies $E \cap \text{cl}_{\beta(\omega)} F = \emptyset$. Now by the method of [11](3.3) for $t \in \{0, 1\}^\omega$ choose $x_t \in \pi^{-1}(\{t\})$ such that every discrete $D \subseteq E \setminus \{x_t\}$ satisfies $x_t \notin \mathbf{cl}_{\beta(\omega)} D$, and take $C = \{x_t : t \in C_0\}$ with C_0 a countable, dense subset of $\{0, 1\}^\omega$. Since π is irreducible the set C is dense in E and is dense-in-itself, and it is easy to see that condition (*) is satisfied.

Of course no element of C is a P -point of ω^* . The existence in ZFC of non- P -points $x \in \omega^*$ such that $x \notin \text{cl}_{\beta(\omega)} D$ whenever D is a countable, discrete, subspace of $\omega^* \setminus \{x\}$ is given explicitly by van Mill [11]; see also Kunen [9] for a construction in ZFC + CH (or, in ZFC + MA) of a set C as above.

For the set C constructed above the relation $\omega \cup \{p\} \subseteq_h C$ fails for every $p \in \omega^*$, so the following question, closely related to Question 6.4, is apparently not answered by the methods of this paper.

6.6. Question. Let C be a countable, dense-in-itself subset of ω^* such that $\omega \cup \{p\} \subseteq_h C$ fails for every $p \in \omega^*$. Is the relation $\omega^* \rightarrow (C)_2^1$ valid? \square

List of References

- [1] W. W. Comfort and Akio Kato. Non-homeomorphic disjoint spaces whose union is ω^* . Rocky Mountain J. Math. To appear.
- [2] W. W. Comfort and Akio Kato. Topological partition relations of the form $\omega^* \rightarrow (Y)_2^1$. Abstracts Amer. Math. Soc. 74 (1991), 288-289 (= Abstract #91T-54-25).
- [3] W. W. Comfort and S. Negrepointis. The Theory of Ultrafilters. Grundlehren der math. Wissenschaften Band 211. Springer-Verlag. Berlin, Heidelberg and New York. 1974.
- [4] Alan Dow and Jan van Mill. An extremally disconnected Dowker space. Proc. Amer. Math. Soc. 86 (1982), 669-672.
- [5] Zdeňek Frolík. Sums of ultrafilters. Bull. Amer. Math. Soc. 73 (1967), 87-91.
- [6] Zdeňek Frolík. Fixed points of maps of $\beta\mathbf{N}$. Bull. Amer. Math. Soc. 74 (1968), 187-191.
- [7] Leonard Gillman and Meyer Jerison. Rings of Continuous Functions. D. Van Nostrand Co., Inc. Princeton, Toronto, New York and London. 1960.
- [8] M. Katětov. A theorem on mappings. Commentationes Math. Universitatis Carolinae 8 (1967), 431-433.
- [9] Kenneth Kunen. Some points in $\beta\mathbf{N}$. Math. Proc. Cambridge Philosophical Soc. 80 (1976), 385-398.
- [10] Kenneth Kunen. Weak P -points in \mathbf{N}^* . Colloquia Mathematica Societatis János Bolyai 23 (1978), 741-749.
- [11] Jan van Mill. Sixteen topological types in $\beta\omega \setminus \omega$. Topology and Its Applications 13 (1982), 43-57.
- [12] Jan van Mill. An introduction to $\beta\omega$. In: Handbook of Set-theoretic Topology, pp. 503-567. Edited by K. Kunen and J. E. Vaughan. North-Holland Publ. Co. Amsterdam, New York, Oxford. 1984.

[13] Mary Ellen Rudin. Partial orders on the types in $\beta\mathbf{N}$. Trans. Amer. Math. Soc. 155 (1971), 353-362.

[14] Stevo Todorćevic. Partition Problems in Topology. Contemporary Mathematics, vol. 84. American Mathematical Society. Providence, Rhode Island. 1989.

[15] William Weiss. Partitioning topological spaces. In: Mathematics of Ramsey Theory, pp. 154-171. Edited by J. Nešetřil and V. Rödl. Springer-Verlag. Berlin. 1990.

[16] W. A. R. Weiss. Weiss's questions. In: Open Problems in Topology, pp. 77-84. Edited by Jan van Mill and G. M. Reed. North-Holland Publ. Co. Amsterdam. 1990.