

On Gross Spaces

Saharon Shelah ^{1 2} and Otmar Spinas ¹

ABSTRACT: A *Gross space* is a vector space E of infinite dimension over some field F , which is endowed with a symmetric bilinear form $\Phi : E^2 \rightarrow F$ and has the property that every infinite dimensional subspace $U \subseteq E$ satisfies $\dim U^\perp < \dim E$. Gross spaces over uncountable fields exist (in certain dimensions) (see [G/O]). The existence of a Gross space over countable or finite fields (in a fixed dimension not above the continuum) is independent of the axioms of ZFC. This was shown in [B/G], [B/Sp] and [Sp2]. Here we continue the investigation of Gross spaces. Among other things we show that if the cardinal invariant \mathfrak{b} equals ω_1 a Gross space in dimension ω_1 exists over every infinite field, and that it is consistent that Gross spaces exist over every infinite field but not over any finite field. We also generalize the notion of a Gross space and construct generalized Gross spaces in ZFC.

0 Introduction

Let E be a vector space of infinite dimension over some field F , and let $\Phi : E \times E \rightarrow F$ be a symmetric bilinear form. If U is a subspace of E , the *orthogonal* complement is

1991 Mathematics Subject Classification: Primary 11E04, 03E35; secondary 12L99, 15A36

¹ The authors are supported by the Basic Research Foundation of the Israel Academy of Science

² Publication number 468

denoted by U^\perp . If U has finite dimension, then clearly we have $\text{codim}_E U^\perp \leq \dim U$, where $\text{codim}_E U^\perp$ is the dimension of some *linear* complement of U^\perp in E . As is well-known from Hilbert space, this is completely false if U is of infinite dimension.

In [G/O], the investigation of quadratic spaces sharing the following strong property has been started:

(*) for all subspaces $U \subseteq E$ of infinite dimension: $\dim U^\perp < \dim E$.

Such a space we call a *Gross space*. In [G/O], the motivation for this was that Gross spaces are natural candidates to have a small orthogonal group in the sense that every isometry is the product of finitely many hyperplane reflections. If a Gross space is the orthogonal sum of two subspaces, then one of them must be finite dimensional. Hence, a Gross space is far from having an orthogonal basis, and so its dimension must be uncountable.

In [G/O], over every uncountable field F a *strong Gross space* has been constructed, i.e. a space sharing the following stronger property:

(**) for all subspaces $U \subseteq E$ of infinite dimension: $\dim U^\perp \leq \aleph_0$.

Such a construction has been achieved in every uncountable dimension less or equal the cardinality of the field.

To construct a Gross space gets the more difficult the smaller the cardinality of the field is compared with the dimension of the space.

In [B/G], a Gross space of dimension \aleph_1 (so Gross=strongly Gross) has been constructed over every countable or finite field. But for this the Continuum Hypothesis (CH) has been assumed, and hence the question was raised whether CH is necessary or a construction in ZFC is possible.

In [Sp1], [B/Sp] and [Sp2], it turned out that this question leads into set theory, and that in fact it is independent of ZFC.

Here we answer several open questions from these papers, and we also continue the investigation of Gross spaces over uncountable fields in [G/O].

A difficult result in [B/Sp] says that if $\mathfrak{b} = \omega_1$, then a Gross space of dimension \aleph_1 exists over every field which is the extension of some finite or countable field by countably many transcendentals. Here \mathfrak{b} is defined as the minimal cardinality of a family of functions

from the natural numbers to themselves which is unbounded under eventual dominance. Clearly $\omega_1 \leq \mathbf{b} \leq \mathbf{c}$. But ZFC does not decide where \mathbf{b} lies exactly. In §2 we will show that if $\mathbf{b} = \omega_1$, then a Gross space of dimension \aleph_1 exists over *every* infinite field.

In §3 we show that $\mathbf{b} = \omega_1$ may hold but no symmetric bilinear space of dimension \aleph_1 over any finite field is a Gross space. We prove three variants of this. The first uses a forcing from [Sp2] which kills Gross spaces, the second involves the splitting number \mathfrak{s} , and the third uses a model from [B/Sh] where simultaneously P_{\aleph_1} - and P_{\aleph_2} -points exist. In this model there exist no Gross spaces over finite fields at all.

For a couple of years the main open problem about Gross spaces has been whether there exists a ZFC model where there exists no Gross space over any countable or finite field in any dimension. The first author thinks that he has constructed a model for this, using a new iteration technique. The paper has not yet been written.

In §4 we investigate a natural generalization of the Gross property:

(***) for all subspaces $U \subseteq E$ of dimension $\geq \lambda$: $\dim U^\perp < \dim E$

Here λ is an infinite cardinal less or equal the dimension of E . A space sharing the property (***) is called a λ -Gross space. Hence a Gross space is an ω -Gross space.

We concentrate on results in ZFC. We show that ω_1 -Gross spaces of dimension \aleph_1 and \aleph_2 always exist over every countably infinite field. We also show that $|F|^\lambda$ is an upper bound for the dimension of a λ -Gross space over F . Hence, in ZFC these results are maximal. We do not know whether these results hold also for finite fields. But we show that if $\lambda > \omega_1$ is regular uncountable, then a λ^+ -Gross space of dimension λ^+ exists over any field.

We also investigate the situation for uncountable fields. The construction in [G/O] yields strong Gross spaces of dimension at most as large as the cardinality of the base field. Is it possible to enlarge the dimension of the space while keeping the size of the field small? Over every field of uncountable cardinality λ we construct a λ -Gross space of dimension λ^+ , as well as a λ^+ -Gross space of dimension λ^{++} . Again, by the upper bound for the dimension of a λ -Gross space mentioned above, these results are maximal in ZFC.

We remark that all the investigations mentioned above may take place in a rather more general context. First, everything remains true if we skip to orthosymmetric sesquilinear forms over a division ring endowed with an involutory antiautomorphism (see [G] for

the definitions). Second, using a representation theorem for AC-lattices equipped with a polarity, the results can be transferred to the level of abstract ortho-lattices. This has been announced in [Sp3].

For a survey on the whole subject refer to [Sp4].

1 Notation and definitions

1.0 Forms

Let F be a commutative field of arbitrary characteristic. Let E be a vector space over F , endowed with a symmetric bilinear form $\Phi : E \times E \rightarrow F$, i.e. Φ is linear in both arguments and $\Phi(x, y) = \Phi(y, x)$ always. Most of the spaces we construct will have *isotropic* vectors, i.e. nonzero vectors x such that $\Phi(x, x) = 0$. For a subspace $U \subseteq E$ the *orthogonal* complement U^\perp is the subspace $\{x \in E : \forall y \in U \Phi(x, y) = 0\}$. We call $\langle E, \Phi \rangle$ *nondegenerate* if $E^\perp = \{0\}$. By E^* we denote the F -vectorspace of linear functionals $E \rightarrow F$. If in E , we have fixed a basis $\langle e_\alpha : \alpha \in I \rangle$, then for a vector $x \in E$ the *support* of x , denoted by $\text{supp}(x)$, is the unique finite set of α 's such that, in the representation of x by $\langle e_\alpha : \alpha \in I \rangle$, e_α has a nonzero coefficient.

1.1 Set theory

For the theory of forcing refer to [B], [J], [K] or [Sh1].

For sets A, B , the set of functions from A to B is denoted by ${}^A B$. If κ, λ are cardinals, then κ^λ denotes the cardinality of the set ${}^\lambda \kappa$, and $[A]^\lambda$ denotes the set of all subsets of A which have cardinality λ . By \mathfrak{c} we denote the cardinality of the continuum. If A is well-ordered, then $\text{o.t.}(A)$ is its order type.

If $f, g \in {}^\omega \omega$, then we say f *eventually dominates* g , and we write $g <^* f$, if $\exists k \forall n \geq k g(n) < f(n)$. A family F of members of ${}^\omega \omega$ is *<*-unbounded* if there is no $f \in {}^\omega \omega$ such that $\forall g \in \text{ran} F g <^* f$. F is called *dominating*, if every function from ${}^\omega \omega$ is eventually dominated by some member of F .

Then \mathfrak{b} is defined as the minimal cardinality of a *<*-unbounded* family in ${}^\omega \omega$.

The cardinal invariant \mathfrak{d} is defined as the minimal cardinality of a dominating family.

For $A, B \in [\omega]^\omega$, we write $A \subseteq^* B$ if $A \setminus B$ is finite. A family \mathcal{S} of members of $[\omega]^\omega$ is called a *splitting family*, if for every $A \in [\omega]^\omega$ there exists $S \in \text{ran}\mathcal{S}$ such that $A \cap S$ and $A \setminus S$ are both infinite.

The splitting number \mathfrak{s} is defined as the minimal cardinality of a splitting family.

Here \mathfrak{b} , \mathfrak{d} and \mathfrak{s} are instances of so called cardinal invariants of the continuum which have found lots of applications throughout mathematics. See [vD] for an introduction.

The following theorems form part of the folklore in set theory. Proofs of the first two of them may be found in [vD].

Theorem 1. \mathfrak{b} , \mathfrak{d} and \mathfrak{s} are uncountable cardinals $\leq \mathfrak{c}$; $\mathfrak{b}, \mathfrak{s} \leq \mathfrak{d}$, and \mathfrak{b} is regular.

The axioms of ZFC do not decide where exactly \mathfrak{b} , \mathfrak{d} and \mathfrak{s} lie.

Theorem 2. Let κ and λ be regular cardinals with $\omega_1 \leq \kappa \leq \lambda$. There exists a forcing preserving all cardinals such that in the extension obtained by forcing with it, $\mathfrak{c} = \lambda$ and $\mathfrak{b} = \mathfrak{d} = \mathfrak{s} = \kappa$ hold.

Theorem 3 [Sh2]/[BS]. $\omega_1 = \mathfrak{b} < \mathfrak{s}$ is consistent with ZFC as well as is $\omega_1 = \mathfrak{s} < \mathfrak{b}$.

2 When $\mathfrak{b} = \omega_1$, a Gross space exists over every infinite field

In [B/Sp, §4] it has been shown that the assumption $\mathfrak{b} = \omega_1$ implies that a Gross space of dimension ω_1 exists over any field which is the extension of an arbitrary finite or countable field by countably many transcendentals. Here we show that $\mathfrak{b} = \omega_1$ is enough to construct a Gross space of dimension ω_1 over any countably infinite field. In §3, we will show that $\mathfrak{b} = \omega_1$ may hold but no Gross spaces exist over finite fields.

The construction will use the filtrations given by the following Lemma.

Lemma. There exists a family $\langle A_k^\alpha : \alpha < \omega_1, k < \omega \rangle$ of finite sets such that for any $\alpha, \beta < \omega_1$ and $k, l < \omega$ the following requirements are satisfied:

- (1) $\alpha = \bigcup_{k < \omega} A_k^\alpha$
- (2) if $k < l$ then $A_k^\alpha \subseteq A_l^\alpha$
- (3) if $\beta < \alpha$ and $\beta \in A_k^\alpha$ then $A_k^\beta = A_k^\alpha \cap \beta$

Proof. We define A_k^α by induction on α .

For $\alpha = 0$ let $A_k^\alpha = \emptyset$.

For $\alpha = \beta + 1$ let $A_k^\alpha = A_k^\beta \cup \{\beta\}$

For α a limit let $\langle \alpha_n : n < \omega \rangle$ be an increasing sequence such that $\alpha = \sup_{n < \omega} \alpha_n$.

Now choose $\langle k_n : n < \omega \rangle$ increasing such that for all n we have $\alpha_n \in A_{k_n}^{\alpha_{n+1}}$. Define

$$A_m^\alpha = A_m^{\alpha_0}$$

for $m \leq k_0$ and

$$A_m^\alpha = A_m^{\alpha_{l+1}}$$

for $k_l < m \leq k_{l+1}$. It is easy to check that this works. □

Theorem. *Assume $\mathfrak{b} = \omega_1$. Then a Gross space of dimension \aleph_1 exists over every infinite field.*

Proof. Let F be a countable, infinite field, and let E be a vector space over F , spanned by a basis $\langle e_\alpha : \alpha < \omega_1 \rangle$.

Let $\langle p_n : n \in \omega \rangle$ enumerate all polynomials in finitely many variables with coefficients in F .

In F choose $\langle a_n : n < \omega \rangle$ such that for any polynomial $p[X_1, \dots, X_k]$, for any n larger than some integer depending on p we have $a_n \notin \{p(a_{l_1}, \dots, a_{l_k}) : l_1, \dots, l_k < n\}$.

Finally, let $\langle f_\alpha : \alpha < \omega_1 \rangle$ be an unbounded, well-ordered family of increasing functions in ω^ω .

For any $\alpha < \omega_1$ we define a function $h_\alpha : \alpha \rightarrow \omega$ by induction, using the filtrations from the Lemma. Assume $\gamma \in A_{k+1}^\alpha \setminus A_k^\alpha$ and assume h_β for $\beta < \alpha$, $h_\alpha \upharpoonright A_k^\alpha$ and $h_\alpha \upharpoonright (A_{k+1}^\alpha \cap \gamma)$ have been defined.

Let $h_\alpha(\gamma) = n$ such that a_n is distinct from all the finitely many scalars of the form

$$p_i(a_{l_1}, a_{l_2}, \dots)$$

where $i < f_\alpha(k+1)$ and any l_j belongs to the range of one of the following functions:

$h_\alpha \upharpoonright A_k^\alpha$, $h_\alpha \upharpoonright (A_{k+1}^\alpha \cap \gamma)$, $h_\beta \upharpoonright A_{k+1}^\beta$ for $\beta \in A_{k+1}^\alpha$.

Now define a symmetric bilinear form $\Phi : E \times E \rightarrow F$ as follows: For $\alpha < \beta < \omega_1$ define:

$$\Phi(e_\alpha, e_\beta) = a_n \text{ if and only if } h_\beta(\alpha) = n$$

$\Phi(e_\beta, e_\beta)$ may be defined arbitrarily.

We have to show that $\langle E, \Phi \rangle$ is Gross. Assume that this is not true. So there is a subspace U of infinite dimension, spanned by a basis $\langle y_k : k < \omega \rangle$, such that $\dim U^\perp$ is uncountable. Each y_k has a representation

$$y_k = \sum_{l=1}^{m_k} b_{kl} e_{\alpha(k,l)}$$

with nonzero coefficients b_{kl} . Choose $\alpha^* < \omega_1$ such that $U \subseteq \text{span}\langle e_\alpha, \alpha < \alpha^* \rangle$. Using a Δ -system argument and the bilinearity of the form, in U^\perp we may certainly find vectors $\langle z_\iota, \iota < \omega_1 \rangle$, all of them having the same nonzero coefficients in their representation, say

$$z_\iota = \sum_{l=1}^n c_l e_{\beta(\iota,l)}$$

such that for all $\iota_1 < \iota_2 < \omega_1$ we have $\alpha^* < \beta(\iota_1, 1) < \dots < \beta(\iota_1, n) < \beta(\iota_2, 1) < \dots < \beta(\iota_2, n)$. Furthermore, we may assume that for some $k^* < \omega$, for all $\iota < \omega_1$, $1 \leq l_1 < l_2 \leq n$ and $1 \leq l \leq n$ we have

$$\beta(\iota, l_1) \in A_{k^*}^{\beta(\iota, l_2)} \text{ and } \alpha^* \in A_{k^*}^{\beta(\iota, l)} \quad (1)$$

Since $\langle f_\alpha, \alpha < \omega_1 \rangle$ is unbounded and well-ordered there exists $j < \omega$ such that $\langle f_{\beta(\iota, n)}(j), \iota \in \omega_1 \rangle$ is unbounded in ω . Since the f_α are increasing this remains true if j is replaced by any $j' \geq j$.

Now choose $k < \omega$ such that the minimal j' with the property

$$\{\alpha(k, 1), \dots, \alpha(k, m_k)\} \subseteq A_{j'+1}^{\alpha^*} \quad (2)$$

is larger than $\max\{j, k^*\}$. Note that by (1) and property (3) of the filtrations from the Lemma, j' is minimal such that (2) holds for any $\beta(\iota, l)$ instead of α^* .

Choose $l_0 \in \{1, \dots, m_k\}$ such that $\alpha(k, l_0)$ is maximal in $A_{j'+1}^{\alpha^*} \setminus A_{j'}^{\alpha^*}$. Then the polynomial

$$- \sum_{\langle l, l' \rangle \in \{1, \dots, n\} \times \{1, \dots, m_k\} \setminus \langle n, l_0 \rangle} b_{kl_0}^{-1} c_n^{-1} b_{kl'} c_l X_{ll'}$$

has its number, say i , in the enumeration of all polynomials fixed in the beginning. There exists $\iota < \omega_1$ such that $f_{\beta(\iota, n)}(j' + 1) > i$. From the definition of Φ , using (1), we conclude

$$\Phi(e_{\beta(\iota, n)}, e_{\alpha(k, l_0)}) \neq - \sum_{\langle l, l' \rangle \in \{1, \dots, n\} \times \{1, \dots, m_k\} \setminus \langle n, l_0 \rangle} b_{kl_0}^{-1} c_n^{-1} b_{kl'} c_l \Phi(e_{\beta(\iota, l)}, e_{\alpha(k, l')})$$

But then $\Phi(z_\iota, y_k) \neq 0$, a contradiction. □

3 Gross spaces over finite fields

The first Lemma shows that \mathbf{c} is an upper bound for the dimension of a Gross space over finite or countable fields.

Lemma 1. *Let $\langle E, \Phi \rangle$ be a symmetric bilinear space over F , and let λ be an infinite cardinal \leq the dimension of E . If $\langle E, \Phi \rangle$ has the property*

$$(***) \text{ for all subspaces } U \subseteq E \text{ of dimension } \geq \lambda: \dim U^\perp < \dim E$$

then $\dim E \leq |F|^\lambda$.

Proof. Let $\dim E = \kappa$ and let $\langle e_\alpha : \alpha < \kappa \rangle$ be a basis. For $\alpha < \kappa$ define

$$f_\alpha : \lambda \rightarrow F, \beta \mapsto \Phi(e_\alpha, e_\beta)$$

We get $\kappa = \bigcup_{f \in {}^\lambda F} A_f$, where $A_f = \{\alpha < \kappa : f_\alpha = f\}$ and hence $\kappa = \sum_{f \in {}^\lambda F} |A_f|$.

So if $\kappa > |F|^\lambda$ there exists $B \subseteq {}^\lambda F$ such that $\sum_{f \in B} |A_f| = \kappa$ and $|A_f| \geq 2$, for all $f \in B$. Fix $\alpha_f \in A_f$ for $f \in B$. We conclude that

$$\bigcup_{f \in B} \{e_\alpha - e_{\alpha_f} : \alpha \in A_f \text{ and } \alpha \neq \alpha_f\}$$

is linearly independent, of cardinality κ and a subset of $\langle e_\beta : \beta < \lambda \rangle^\perp$. □

In [Sp2], the following forcing to kill Gross spaces was introduced.

Let F be a field. The direct sum $\bigoplus_{n < \omega} F$ of ω copies of F naturally bears a vector space structure over F

The forcing P^F consists of pairs $\langle s, A \rangle$ where

- (1) $s = \langle s_0, \dots, s_k \rangle$ is a finite sequence of vectors $s_i \in \bigoplus_{n < \omega} F$
- (2) $A \subseteq (\bigoplus_{n < \omega} F)^*$ is a finite set of linear functionals.

The ordering on P^F is defined as follows:

$\langle s', A' \rangle$ extends $\langle s, A \rangle$ if and only if $s' \supseteq s$ and $A' \supseteq A$ and

$$\forall s'_i \in \text{ran}(s') \setminus \text{ran}(s) \forall f \in A \ f(s'_i) = 0$$

It is easily seen that two conditions with the same first coordinate are compatible. Hence, if F is countable or finite P^F has the countable chain condition.

Let $\langle E, \Phi \rangle$ be an inner product space over F and U an arbitrary subspace of infinite dimension. Then forcing with P^F introduces in U an infinite set of linearly independent vectors such that every vector in E is orthogonal on all but finitely many of them. Moreover, iterating forcing with P^F such that F again and again runs through all finite and countable fields produces a model where Gross spaces of any dimension less than the length of the iteration (which is supposed to be a regular cardinal) do not exist (see [Sp2] for the proofs).

By the Theorem in §2 above, if $\mathfrak{b} = \omega_1$ holds a Gross space of dimension ω_1 exists over every infinite field. Hence if F is infinite, the forcing P^F which kills Gross spaces over F adds a dominating function (see [Sp2]). Here we show that in case F is finite P^F does not destroy unbounded families, more exactly it is almost ${}^\omega\omega$ -bounding. Hence, by a preservation theorem in [Sh1] or [Sh2], iterating forcings P^F where F runs again and again through all finite fields yields a model where a Gross space in dimension ω_1 exists over every infinite field but not over any finite field, provided that we start with a ground model satisfying $\mathfrak{b} = \omega_1$.

Lemma 2. *Assume that F is a finite field and τ is a P^F -name for a function in ${}^\omega\omega$. Then there exists $f \in {}^\omega\omega$ in the ground model such that for every $p \in P^F$ there are only finitely many n such that $p \vdash \tau(n) > f(n)$.*

Proof. At first, let us introduce the following notation. If $s = \langle s_0, \dots, s_k \rangle$ is a sequence of vectors in $\bigoplus_{n < \omega} F$ and ϕ is a formula in the language of forcing with P^F we write $s \vdash^* \phi$ if and only if there exists a finite $A \subset (\bigoplus_{n < \omega} F)^*$ such that $\langle s, A \rangle \vdash \phi$.

By induction on $i < \omega$ we will define an increasing sequence $\langle n_i : i < \omega \rangle$ such that:

(1) $n_0 = 0$ and

(2) for every i , if $s = \langle s_0, \dots, s_k \rangle$ is a sequence of vectors in $\bigoplus_{j < n_i} F$ and f_0, \dots, f_{n_i} are linear functionals on $\bigoplus_{j < n_{i+1}} F$, then there is a sequence s' of vectors in $\bigoplus_{j < n_{i+1}} F$ which extends s , such that $\text{ran}(s') \setminus \text{ran}(s) \subseteq \bigcap_{k \leq n_i} \ker f_k$ and for some $\sigma \in {}^{n_i}\omega$ we have $s' \vdash^* \tau \upharpoonright n_i = \sigma$.

We have to show that for given n_i there exists $n_{i+1} > n_i$ such that (2) holds. Assume that this is not true. So we may find s with $\text{ran}(s) \subseteq \bigoplus_{j < n_i} F$ such that for infinitely many $j > n_i$ there are $f_0^j, \dots, f_{n_i}^j \in (\bigoplus_{u < j} F)^*$ such that there is no s' as in the conclusion of (2).

Now every $(\bigoplus_{u < j} F)^*$ is finite. Hence by König's Lemma, there exist $f_0, \dots, f_{n_i} \in (\bigoplus_{j < \omega} F)^*$ such that for some $B \in [\omega]^\omega$ we have $\forall j \in B \exists j' > j \forall k \leq n_i f_k \upharpoonright \bigoplus_{u < j} F = f_k^{j'} \upharpoonright \bigoplus_{u < j} F$.

Since $\langle s, \{f_0, \dots, f_{n_i}\} \rangle$ is a condition in P^F there exists a stronger condition $\langle s', A' \rangle$ which decides the first n_i values of τ . Choose $j \in B$ so large that $\text{ran}(s') \subseteq \bigoplus_{u < j} F$. Now for some $j' > \max\{j, n_i\}$ we know $\forall k \leq n_i f_k \upharpoonright \bigoplus_{u < j} F = f_k^{j'} \upharpoonright \bigoplus_{u < j} F$. But then s' satisfies the conclusion of (2) for s and $f_0^{j'}, \dots, f_{n_i}^{j'}$, a contradiction.

Hence we have shown that $\langle n_i : i < \omega \rangle$ can be chosen increasing and satisfying (1) and (2).

Next we define $f : \omega \rightarrow \omega$ as follows: If $n_i \leq n < n_{i+1}$, then

$$f(n) = \max\{\sigma(n) : \text{there exists } s \text{ such that } \text{ran}(s) \subseteq \bigoplus_{j < n_{i+2}} F \text{ and } s \vdash^* \tau \upharpoonright n_{i+1} = \sigma\}$$

We claim that f is as desired. So let $\langle s, \{f_0, \dots, f_m\} \rangle \in P^F$ and let $i < \omega$ such that $n_i > m$ and $\text{ran}(s) \subseteq \bigoplus_{j < n_i} F$. Assume now $n \geq n_i$. We will find an extension of $\langle s, \{f_0, \dots, f_m\} \rangle$ forcing " $f(n) \geq \tau(n)$ ".

Choose j such that $n_j \leq n < n_{j+1}$. By construction, there is s' extending s such that $\text{ran}(s') \subseteq \bigoplus_{u < n_{j+2}} F$, $\text{ran}(s') \setminus \text{ran}(s) \subseteq \bigcap_{k=1}^m \ker f_k$ and $s' \vdash^* \tau \upharpoonright n_{j+1} = \sigma$, for some σ . Choose a finite $A \subseteq (\bigoplus_{u < \omega} F)^*$ containing $\{f_0, \dots, f_m\}$ such that $\langle s', A \rangle \vdash \tau \upharpoonright n_{j+1} = \sigma$. Then $\langle s', A \rangle$ extends $\langle s, \{f_0, \dots, f_m\} \rangle$ and $\langle s', A \rangle \vdash f(n) \geq \tau(n)$. \square

Remark: It is easy to see that Lemma 2 says much more than that P^F is almost ${}^\omega\omega$ -bounding (this notion is defined in [Sh2]). In [Sh2] it is proved that this property is preserved under countable support iterations. In the new edition of [Sh1] it is proved that this is true also for finite support iterations. A more accessible proof of this preservation theorem (for the finite support case) can be found in [Go, Example 8.5]. It is not difficult to see that P^F adds a Cohen real, hence it is not ${}^\omega\omega$ -bounding.

Corollary. *Assume P is a finite or countable support iteration of forcings P^F where F is a finite field. If in V , $\langle f_\alpha : \alpha < \kappa \rangle$ is an unbounded family of functions in ω^ω , then $\langle f_\alpha : \alpha < \kappa \rangle$ is unbounded in V^P .*

Proof. Preservation of unboundedness at successor steps of the iteration follows from Lemma 1: Assume τ is a P -name for a function in ω^ω such that some $p \in P$ forces “ τ bounds $\langle f_\alpha : \alpha < \kappa \rangle$ ”. Choose $f \in \omega^\omega \cap V$ for τ as in the Lemma. Since $\langle f_\alpha : \alpha < \kappa \rangle$ is unbounded in V there exists α such that for infinitely many n we have $f(n) < f_\alpha(n)$. Find $p' \in P$ extending p and n_0 such that $p' \vdash \forall n \geq n_0 \tau(n) > f_\alpha(n)$. But then for infinitely many n we would have $p' \vdash \tau(n) > f(n)$, a contradiction.

The limit steps are handled by a general preservation theorem due to the first author (see [Sh2] or [Sh1-new ed., VI§3]). \square

From the Corollary, the Theorem in §2 and the results about forcing with P^F 's in [Sp2] we obtain the following theorem:

Theorem 1. *Assume that the ground model V satisfies $\mathfrak{b} = \omega_1$. Let $P = \lim \langle P_\alpha, Q_\alpha : \alpha < \kappa \rangle$ be a finite or countable support iteration of length κ where $\kappa > \omega_1$ is regular such that*

- (1) *each Q_α is P^F defined in V^{P_α} where F is a finite field, and*
- (2) *for each finite field F , cofinally many times we have $Q_\alpha = P^F$.*

Then in V^P , there exists a Gross space of dimension ω_1 over every infinite field but there is no Gross space of dimension κ over any finite field.

Another way to get a model for the conclusion of the Theorem above is shown by the following Lemma combined with Theorem 3, §1, and the Theorem in §2.

Lemma 3. *Assume that $\langle E, \Phi \rangle$ is a symmetric bilinear space over a finite field and the dimension of E has uncountable cofinality. If $\mathfrak{s} > \dim E$ then $\langle E, \Phi \rangle$ is not Gross.*

Proof. Let $\dim E = \kappa$. In E , choose a basis $\langle e_\alpha, \alpha < \kappa \rangle$ and let $\langle y_n, n < \omega \rangle$ span a subspace. For every $\alpha < \kappa$ let

$$f_\alpha : \omega \rightarrow F, n \mapsto \Phi(e_\alpha, y_n)$$

where F is the base field. Let $\mathcal{A} = \langle f_\alpha^{-1}\{a\} : \alpha < \kappa, a \in F \rangle$. Since $\mathfrak{s} > \kappa$, \mathcal{A} is not a splitting family. Hence there is an infinite $A \subseteq \omega$ such that for all $B \in \text{ran } \mathcal{A}$, either $A \subseteq^* B$ or $A \cap B$ is finite. For any $\alpha < \kappa$, $\langle f_\alpha^{-1}\{a\}, a \in F \rangle$ is a finite partition of ω . So there exist a_α and n_α such that $A \setminus n_\alpha \subseteq f_\alpha^{-1}\{a_\alpha\}$. By $\text{cf}(\kappa) > \omega$ there exist a, n and $X \in [\kappa]^\kappa$ such that for every $\alpha \in X$ we have $a_\alpha = a$ and $n_\alpha = n$. Now it is easy to see that every vector $e_\alpha - e_\beta$ with $\alpha, \beta \in X$ is in the orthogonal complement of $\text{span}\langle y_m, m \in A \setminus n \rangle$. \square

The model from [B/Sh] where simultaneously simple P_{\aleph_1} - and P_{\aleph_2} -points exist shows that a stronger statement than the conclusion of Theorem 1 is true. A P_κ -point, where κ is a regular uncountable cardinal, is a filter on ω which is generated by a tower (i.e. a \subseteq^* -decreasing family in $[\omega]^\omega$) of length κ .

Theorem 2. *It is consistent that a Gross space of dimension ω_1 exists over every infinite field but no Gross space exists over any finite field in any dimension.*

Proof. In the model of [B/Sh] where simple P_{\aleph_1} - and P_{\aleph_2} -points exist, $\mathfrak{b} = \omega_1$ holds. Hence by the Theorem in §2, a Gross space of dimension \aleph_1 exists over any infinite field. Using a similar argument as in the proof of Lemma 3, one shows that a P_{\aleph_1} -, P_{\aleph_2} -point rules out the existence of a Gross space over any finite field in dimension \aleph_1, \aleph_2 respectively. But in that model $\mathfrak{c} = \omega_2$. Hence by Lemma 1 we are done. \square

Question: Does there exist a ZFC-model where there exists no Gross space over any finite or countable field in any dimension?

In view of the Theorem in §2, a natural question to ask is whether the assumption $\mathfrak{s} = \omega_1$ is strong enough to construct a Gross space of dimension \aleph_1 over a finite field. The answer is “no”.

Theorem 3. *It is consistent that $\mathfrak{s} = \omega_1$ holds and there exists no Gross space in dimension ω_1 over any finite field.*

Proof. Since every finite or countable field can be coded by a real, it is not difficult to see that the forcing notion P^F , where F is finite or countable, is Souslin in the sense of [J/Sh]; i.e., the set of conditions can be viewed as an analytic set of reals, whereas the ordering and the incompatibility relation are analytic subsets of the plane. In [J/Sh] it is proved that if we start with a model satisfying CH, then in any finite support extension of Souslin forcing notions having the countable chain condition, the reals of the ground model remain a splitting family. Hence in V^P , where P is the finite support iteration of Theorem 1 and V satisfies CH, $\mathfrak{s} = \omega_1$ holds and there are no Gross spaces in any dimension $< \mathfrak{c}$ of uncountable cofinality.

Is there another cardinal invariant such that its being \aleph_1 implies that Gross spaces over finite fields exist? The largest of the “classical” cardinal invariants is \mathfrak{d} . The model in [Sh3, §2] shows that even $\mathfrak{d} = \omega_1$ does not suffice for our purpose. In that model, if $\langle E, \Phi \rangle$ is a quadratic space spanned by a basis $\langle e_\alpha : \alpha < \omega_1 \rangle$ over some finite field F , then the following holds:

Fact. *If $\langle u_n : n < \omega \rangle$ is a sequence of pairwise disjoint subsets of ω such that every u_n has size $|F|^n + 1$, then we can find $\langle H_n : n < \omega \rangle$, where each H_n is a family of at most n functions $u_n \rightarrow F$, such that*

$$\forall \alpha < \omega_1 \exists n_\alpha < \omega \forall n \geq n_\alpha \exists h \in H_n \forall m \in u_n [\Phi(e_\alpha, e_m) = h(m)]$$

It is not difficult to see that this implies that $\langle E, \Phi \rangle$ is not Gross (solve many systems of homogeneous equations).

Next we will show that if there is a family of \aleph_1 many meagre sets of reals which is cofinal with respect to inclusion in the set of all meagre sets, then a Gross space of

dimension \aleph_1 exists over every finite field. We work in 2^ω considered as topological product of the discrete space $2 = \{0, 1\}$. Denote by \mathcal{M} the set of all meagre sets in 2^ω . Now $\text{cof}(\mathcal{M})$ is the cardinal invariant defined as follows:

$$\text{cof}(\mathcal{M}) = \min\{|\mathcal{F}| : \mathcal{F} \text{ is a family of meagre sets such that } \forall A \in \mathcal{M} \exists B \in \text{ran } \mathcal{F} \ A \subseteq B\}$$

Theorem 4. *Assume $\text{cof}(\mathcal{M}) = \omega_1$. A Gross space of dimension \aleph_1 exists over every finite field.*

Proof. Using the assumption it is standard work to construct a family $\langle r_\alpha : \alpha < \omega_1 \rangle$ of reals such that, if $A \subseteq (2^\omega)^n$ is a meagre set for some n , then $\exists \alpha < \omega_1 \forall \alpha < \alpha_1 < \dots < \alpha_n \langle r_{\alpha_1}, \dots, r_{\alpha_n} \rangle \notin A$.

By induction on $\alpha < \omega_1$ choose one-to-one functions $h_\alpha : \alpha \rightarrow \omega$ such that $\omega \setminus \text{ran}(h_\alpha)$ is infinite and for every $\beta < \alpha$ the set $\{\gamma < \beta : h_\beta(\gamma) \neq h_\alpha(\gamma)\}$ is finite. (This is a well-known construction of an Aronszajn tree due to Todorcevic.)

Now let F be a field and E a vector space over F of dimension \aleph_1 , spanned by a basis $\langle e_\alpha : \alpha < \omega_1 \rangle$. Define a symmetric bilinear form Φ on E as follows: For $\alpha < \beta < \omega_1$ set:

$$\Phi(e_\alpha, e_\beta) = r_\beta(h_\beta(\alpha))$$

We claim that $\langle E, \Phi \rangle$ is Gross. If this is not true there are families of vectors $\langle y_k : k < \omega \rangle$ and $\langle z_\iota : \iota < \omega_1 \rangle$ such that $\Phi(y_k, z_\iota) = 0$ always, $\langle \text{dom}(y_k) : k < \omega \rangle$ and $\langle \text{dom}(z_\iota) : \iota < \omega_1 \rangle$ are families of pairwise disjoint sets and the sets of the latter have all the same cardinality and are disjoint from some $\alpha^* < \omega_1$ such that $\text{span}\langle y_k : k < \omega_1 \rangle \subseteq \text{span}\langle e_\alpha : \alpha < \alpha^* \rangle$. Let

$$y_k = \sum_{l=1}^{m_k} b_{kl} e_{\alpha(k,l)}$$

$$z_\iota = \sum_{l=1}^n c_l e_{\beta(\iota,l)}$$

such that each $m_k > 0$ and $n > 0$, each $b_{kl}, c_l \neq 0$ and $\beta(\iota, 1) < \dots < \beta(\iota, n)$ always.

Define $A \subseteq (2^\omega)^n$ as follows:

$$A = \{\langle r_1, \dots, r_n \rangle : \{k : \sum_{1 \leq l \leq m_k} \sum_{1 \leq l' \leq n} b_{kl} c_{l'} r_{l'}(h_{\alpha^*}(\alpha(k, l))) \neq 0\} \text{ is finite}\}$$

It is not difficult to see that A is meagre. Hence by construction we may choose ι so large that $\langle r_{\beta(\iota, 1)}, \dots, r_{\beta(\iota, n)} \rangle \notin A$. Consequently, by the choice of the h_α we may find k such that

$$\begin{aligned} \Phi(y_k, z_\iota) &= \sum_{1 \leq l \leq m_k} \sum_{1 \leq l' \leq n} b_{kl} c_{l'} r_{\beta(\iota, l')}(h_{\beta(\iota, l')}(y_k)) = \\ &= \sum_{1 \leq l \leq m_k} \sum_{1 \leq l' \leq n} b_{kl} c_{l'} r_{\beta(\iota, l')}(h_{\alpha^*}(\alpha(k, l))) \neq 0 \end{aligned}$$

This is a contradiction. □

Remark. Theorem 4 is true if the assumption $\text{cof}(\mathcal{M}) = \omega_1$ is replaced by $\text{cof}(\mathcal{N}) = \omega_1$ where \mathcal{N} denotes the set of all Lebesgue measure zero sets of reals. The same proof works if we replace “meagre” by “measure zero”. The proof also works for arbitrary (not necessarily finite) fields. All this is not astonishing since Cichoń’s diagram (see [F]) tells us that $\mathfrak{b} \leq \mathfrak{d} \leq \text{cof}(\mathcal{M}) \leq \text{cof}(\mathcal{N})$.

4 Generalized Gross spaces exist in ZFC

Here we consider a natural generalization of the Gross property and will obtain results in ZFC.

Let F be a field of arbitrary cardinality and $\langle E, \Phi \rangle$ a symmetric bilinear space over F , and let λ be an infinite cardinal. We say that $\langle E, \Phi \rangle$ is a λ -Gross space if it has the following property:

$$(***) \text{ for all subspaces } U \subseteq E \text{ of dimension } \geq \lambda: \dim U^\perp < \dim E$$

So a Gross space is a ω -Gross space. By Lemma 1 in §3, $|F|^\lambda$ is an upper bound for the dimension of a λ -Gross space over F .

By results from [B/G] and [B/Sp] the existence of Gross spaces of dimension \aleph_1 over countable or finite fields is independent of ZFC. We will prove here that in ZFC an ω_1 -Gross space of dimension \aleph_1 can be constructed over any countably infinite field.

We need the following fancy Lemma:

Lemma 1. *Let F be an infinite field. There exists a sequence $\langle a_n : n < \omega \rangle$ of elements of F such that, whenever $\langle k_{ij} : 1 \leq i, j \leq m \rangle$ is a finite sequence of pairwise distinct integers, then*

$$\begin{vmatrix} a_{k_{11}} & \cdots & a_{k_{1m}} \\ \vdots & & \vdots \\ a_{k_{m1}} & \cdots & a_{k_{mm}} \end{vmatrix} \neq 0$$

Proof. By a fusion argument. Start with a sequence $\langle a_n^1 : n < \omega \rangle$ of pairwise distinct scalars in $F \setminus \{0\}$. Then clearly every subsequence satisfies the conclusion of the Lemma for $m = 1$.

Now assume $\langle a_n^m : n < \omega \rangle$ has been constructed such that the conclusion of the Lemma holds for any $1 \leq m' \leq m$. Let $n_i = i$ for any $i \leq (m+1)^2 - 1$. Assume n_0, \dots, n_l have been chosen.

Let $\langle i_0, j_0 \rangle \in \{1, \dots, m+1\}^2$ and let $\langle k_{ij} : \langle i, j \rangle \in \{1, \dots, m+1\}^2 \setminus \langle i_0, j_0 \rangle \rangle$ be a sequence of pairwise distinct integers in $\{n_0, \dots, n_l\}$. By induction hypothesis, the equation

$$i_0 \rightarrow \begin{vmatrix} & \begin{matrix} j_0 \\ \downarrow \end{matrix} & \\ a_{k_{11}}^m & \cdots & a_{k_{1m+1}}^m \\ \vdots & X & \vdots \\ a_{k_{m+11}}^m & \cdots & a_{k_{m+1m+1}}^m \end{vmatrix} = 0$$

in one variable X which has coordinates $\langle i_0, j_0 \rangle$ has exactly one solution. (Expand the determinant by the i_0 th row. Then the cofactors are all nonzero.)

Now choose $n_{l+1} > n_l$ minimal such that for any $n \geq n_{l+1}$, a_n^m is distinct from all the finitely many solutions of the equation above obtained by running through all possible $\langle i_0, j_0 \rangle$'s and $\langle k_{ij} : \langle i, j \rangle \in \{1, \dots, m+1\}^2 \setminus \langle i_0, j_0 \rangle \rangle$'s.

Now define $a_l^{m+1} = a_{n_l}^m$, $l < \omega$.

Finally define

$$a_n = a_n^m \text{ iff } n \in \{m^2 - 1, \dots, (m + 1)^2 - 2\}$$

It is easily checked that $\langle a_n : n < \omega \rangle$ is as desired. □

For uncountable fields an analog of Lemma 1 is true which is more obvious. This is essentially [B/Sp, Lemma 2, §4].

Lemma 2. *Let F be a field of uncountable cardinality λ . There exists a sequence $\langle a_\nu : \nu < \lambda \rangle$ of elements of F such that such that, whenever $\langle \nu_{ij} : 1 \leq i, j \leq m \rangle$ is a finite sequence of pairwise distinct ordinals $< \lambda$, then*

$$\begin{vmatrix} a_{\nu_{11}} & \dots & a_{\nu_{1m}} \\ \vdots & & \vdots \\ a_{\nu_{m1}} & \dots & a_{\nu_{mm}} \end{vmatrix} \neq 0$$

Proof. Let $\langle a_\nu : \nu < \lambda \rangle$ be a one-to-one enumeration of a transcendence base of F over its prime field. Prove by induction on the matrix size m that the conclusion is satisfied. Expand the determinant by, say, the first row. Then by induction hypothesis the cofactors are all nonzero, and the transcendentals of the first row do not occur in them. □

Theorem 1. *A ω_1 -Gross space of dimension \aleph_1 exists over every countably infinite field.*

Proof. Let F be a countably infinite field. Choose scalars $\langle a_n : n < \omega \rangle$ in F as in Lemma 1. Let $\langle A_\alpha : \alpha < \omega_1 \rangle$ be an almost disjoint family of subsets of $\text{ran}\langle a_n : n < \omega \rangle$. For each $\alpha < \omega_1$ choose a one-to-one function $f_\alpha : \alpha \rightarrow A_\alpha$. Let E be a vector space over F of dimension \aleph_1 and let $\langle e_\alpha : \alpha < \omega_1 \rangle$ be a basis of E .

On E , define a symmetric bilinear form as follows: For $\alpha < \beta < \omega_1$ define

$$\Phi(e_\alpha, e_\beta) = f_\beta(\alpha)$$

$\Phi(e_\alpha, e_\alpha)$ may be defined arbitrarily.

We claim that $\langle E, \Phi \rangle$ is as desired. Using a Δ -system argument and the bilinearity of Φ , it is not difficult to see that it is enough to prove the following:

(\star) Assume that U is a subspace spanned by a basis $\langle y_k : k < \omega \rangle$ such that the sets in $\langle \text{supp}(y_k) : k < \omega \rangle$ are pairwise disjoint and of the same cardinality, then $\dim U^\perp < \omega_1$.

By way of contradiction assume that U and $\langle y_k : k < \omega \rangle$ are as in (\star) but U^\perp contains ω_1 many linearly independent vectors $\langle z_\iota : \iota < \omega_1 \rangle$.

Let

$$y_k = \sum_{l=1}^m b_{kl} e_{\alpha(k,l)}$$

We may assume that each z_ι has the same nonzero coefficients in its representation, say

$$z_\iota = \sum_{l=1}^n c_l e_{\beta(\iota,l)}$$

and for any $\iota < \nu < \omega_1$ the sets $\text{supp}(z_\iota)$ and $\text{supp}(z_\nu)$ are disjoint and $\text{supp}(z_\iota)$ is disjoint from $\text{sup} \bigcup_{k < \omega} \text{supp}(y_k)$.

We claim that not even the first $m \cdot n$ many z_ι 's are in U^\perp . In order to see this, let us compute

$$\begin{aligned} \Phi(z_\iota, y_k) = & c_1 b_{k1} f_{\beta(\iota,1)}(\alpha(k,1)) + \dots + c_1 b_{km} f_{\beta(\iota,1)}(\alpha(k,m)) \\ & \vdots \\ & \vdots \\ & + c_n b_{k1} f_{\beta(\iota,n)}(\alpha(k,1)) + \dots + c_n b_{km} f_{\beta(\iota,n)}(\alpha(k,m)) \end{aligned}$$

Since the sets A_α are almost disjoint, the sets $\text{supp}(y_k) = \{\alpha(k,1), \dots, \alpha(k,m)\}$ are pairwise disjoint, the functions f_α are one-to-one and the scalars $\langle a_n : n < \omega \rangle$ are chosen satisfying the conclusion of Lemma 1, there is $k < \omega$ such that for $\iota < m \cdot n$ the sets

$$\{f_{\beta(\iota,i)}(\alpha(k,j)) : 1 \leq i \leq n, 1 \leq j \leq m\}$$

are pairwise disjoint, contain $m \cdot n$ elements and

$$\begin{vmatrix} f_{\beta(0,1)}(\alpha(k,1)) & \dots & f_{\beta(0,n)}(\alpha(k,m)) \\ \vdots & & \vdots \\ f_{\beta(mn-1,1)}(\alpha(k,1)) & \dots & f_{\beta(mn-1,n)}(\alpha(k,m)) \end{vmatrix} \neq 0 \quad (1)$$

But then y_k is not orthogonal on every z_ι , $\iota < m \cdot n$, since otherwise the vector

$$\langle c_1 b_{k1}, \dots, c_n b_{km} \rangle \in F^{mn}$$

would be a nontrivial solution of the system of homogeneous equations $Ax = 0$ where A is the $mn \times mn$ -matrix in (1). \square

The upper bound for the dimension of a ω_1 -Gross space over a countable field is $\omega^{\omega_1} = 2^{\omega_1}$, by Lemma 1, §3. Hence, the largest dimension in which such a space conceivably can be constructed in ZFC is ω_2 . Theorem 6 below will show that in fact such a construction is possible.

For the following question we have no answer:

Question. Does there exist a ω_1 -Gross space of uncountable dimension over any finite field?

Next we will show that this question has a positive answer for regular cardinals larger than ω_1 . For this, recent work of the first author on colouring pairs of ordinals will be applied. Let $\lambda, \mu, \kappa, \theta$ be cardinals such that λ is infinite and $\lambda \geq \mu \geq \kappa + \theta$. Let $\text{Pr}_0(\lambda, \mu, \kappa, \theta)$ be the following statement:

$\text{Pr}_0(\lambda, \mu, \kappa, \theta)$: There exists a function $c : [\lambda]^2 \rightarrow \kappa$ such that, if $\xi < \theta$ and for $\nu < \mu$, $\langle \alpha_{\nu, \zeta} : \zeta < \xi \rangle$ is a strictly increasing sequence of ordinals $< \lambda$, the $\alpha_{\nu, \zeta}$ distinct, and $h : \xi \times \xi \rightarrow \kappa$, then there are $\nu_1 < \nu_2 < \mu$ such that for all $\zeta_1, \zeta_2 < \xi$ we have $c\{\alpha_{\nu_1, \zeta_1}, \alpha_{\nu_2, \zeta_2}\} = h\langle \zeta_1, \zeta_2 \rangle$.

The following theorem, which is due to the first author, states that $\text{Pr}_0(\lambda^+, \lambda^+, \lambda^+, \omega)$ holds for all regular uncountable cardinals. For regular $\lambda > \omega_1$ this is proved in [Sh4, Corollary (a), Section 4, p.100]; see Definition 1, p.95. For $\lambda = \omega_1$ this was proved in [Sh7, Theorem 1.1].

Theorem 2. $\text{Pr}_0(\lambda^+, \lambda^+, \lambda^+, \omega)$ and hence $\text{Pr}_0(\lambda^+, \lambda^+, 2, \omega)$ holds for every regular uncountable cardinal λ .

For more on Theorem 2 (for $\lambda > \omega_1$) see also [Sh5]. It appears (when we specify $\theta = \omega$) in [Sh5, III, 4.8(1), p.177]. Moreover, 4.8(2) gives similar results for inaccessibles with stationary subsets not reflecting in inaccessibles.

Theorem 3. *Let λ be a regular uncountable cardinal. A λ^+ -Gross space of dimension λ^+ exists over any field.*

Proof. Let F be an arbitrary field of size at most λ (for larger fields see Theorem 4 below) and E a vector space over F of dimension λ^+ . Choose a basis $\langle e_\alpha : \alpha < \lambda^+ \rangle$. We define a symmetric bilinear form on E using the colouring given by $\text{Pr}_0(\lambda^+, \lambda^+, 2, \omega)$. For $\{\alpha, \beta\} \in [\lambda^+]^2$ we set

$$\Phi(e_\alpha, e_\beta) = c\{\alpha, \beta\}$$

The angles $\Phi(e_\alpha, e_\alpha)$ may be defined arbitrarily.

Assume that $\langle E, \Phi \rangle$ is not λ^+ -Gross. So we may find families $\langle y_\nu : \nu < \lambda^+ \rangle$ and $\langle z_\nu : \nu < \lambda^+ \rangle$ of linearly independent vectors such that $\Phi(y_\nu, z_{\nu'}) = 0$ always. Without loss of generality we may assume that $\langle \text{dom}(y_\nu) : \nu < \lambda^+ \rangle$ and $\langle \text{dom}(z_\nu) : \nu < \lambda^+ \rangle$ are families of pairwise disjoint sets, the sets in the first one all of size n and those in the latter all of size m , and if $\nu < \nu'$ and $\alpha \in \text{dom}(y_\nu)$, $\beta \in \text{dom}(z_\nu)$, $\gamma \in \text{dom}(y_{\nu'})$ then $\alpha < \beta < \gamma$. Furthermore we may assume that each y_ν and each z_ν has the same nonzero coefficients in its representation, say

$$y_\nu = \sum_{l < n} b_l e_{\beta(\nu, l)}$$

$$z_\nu = \sum_{l < m} c_l e_{\gamma(\nu, l)}$$

We assume that $\beta(\nu, l)$ and $\gamma(\nu, l)$ increase with l . Let $\langle \alpha_{\nu, i} : i < n + m \rangle$ be the increasing enumeration of $\text{dom}(y_\nu) \cup \text{dom}(z_\nu)$.

It is easy to define $h : (n + m) \times (n + m) \rightarrow 2$ such that

$$\sum_{l < n} \sum_{l' < m} b_l c_{l'} h\langle l, n + l' \rangle \neq 0$$

But now by $\text{Pr}_0(\lambda^+, \lambda^+, 2, \omega)$ we may find $\nu_1 < \nu_2 < \lambda^+$ such that for all $i, j < n + m$ we have $c\{\alpha_{\nu_1, i}, \alpha_{\nu_2, j}\} = h\langle i, j \rangle$. We conclude $\Phi(y_{\nu_1}, z_{\nu_2}) \neq 0$, a contradiction. \square

For uncountable fields of cardinality equal to the dimension of the space a much stronger version of Theorems 1 and 3 was proved in [G/O]:

Theorem 4 [G/O, Thm. 1, §1]. *Let F be a field of uncountable cardinality λ . A strong Gross space of dimension λ over F exists.*

Here a strong Gross space is a space $\langle E, \Phi \rangle$ satisfying the following property:

(**) for all subspaces $U \subseteq E$ of infinite dimension: $\dim U^\perp \leq \aleph_0$

For more on strong Gross spaces see [Sp4].

Constructing a Gross space gets the more difficult, the larger its dimension is compared with the size of its base field. For $\lambda = \omega$, the existence of a λ -Gross space of dimension λ^+ over some field of size λ is independent of ZFC. The next theorem shows that if λ is uncountable, then such a space can be constructed in ZFC. If $\text{cf}(\lambda) > \omega$, a similar construction as in Theorem 1 is possible. If $\text{cf}(\lambda) = \omega$, the almost disjoint sets used for the construction must be chosen more carefully. For this we will need the following result from the first author's work on cardinal arithmetic. It is obtained from [Sh6, 6.2] with the ideal of bounded subsets of κ and the f_α 's coming from [Sh5, II, 1.5, p.50].

Lemma 3 [Sh6, 6.2]. *Assume that λ is a singular cardinal and $\text{cf}(\lambda) = \kappa$. There exists an increasing sequence $\langle \lambda_\iota : \iota < \kappa \rangle$ of regular cardinals $\lambda_\iota > \kappa$ with limit λ and a family $\langle f_\alpha : \alpha < \lambda^+ \rangle$ of functions in $\prod_{\iota < \kappa} \lambda_\iota$ such that for all $\alpha < \beta < \lambda^+$:*

(1) $f_\alpha < f_\beta$ modulo bounded subsets of κ , and

(2) if $\langle u_\zeta : \zeta < \lambda^+ \rangle$ is a sequence of pairwise disjoint nonempty subsets of λ^+ , each u_ζ of size $< \kappa$, and $\mu < \lambda$ is a cardinal, then there exists $B \subseteq \lambda^+$ of size μ and $\iota^* < \kappa$ such that for all $\zeta, \xi \in B$ and $\iota < \kappa$:

(i) if $\zeta < \xi$, then $\sup u_\zeta < \min u_\xi$, and

(ii) if $\alpha, \beta \in \bigcup_{\zeta \in B} u_\zeta$, $\alpha < \beta$ and $\iota^* \leq \iota < \kappa$, then $f_\alpha(\iota) < f_\beta(\iota)$.

Theorem 5. *Let λ be an uncountable cardinal and F a field of size λ . Then a λ -Gross space over F of dimension λ^+ exists.*

Proof. Let F be a field of cardinality λ . Let $\langle a_\nu : \nu < \lambda \rangle$ be a transcendence base of F over its prime field F_0 . Hence, F is an algebraic extension of $F_0(\langle a_\nu : \nu < \lambda \rangle)$. Let E be a F -vector space of dimension λ^+ , spanned by a basis $\langle e_\alpha : \alpha < \lambda^+ \rangle$.

Case 1: Assume $\text{cf}(\lambda) > \omega$. Choose $\langle A_\alpha : \alpha < \lambda^+ \rangle$, a family of almost disjoint subsets of λ . I.e. each A_α has size λ and if α, β are distinct then $|A_\alpha \cap A_\beta| < \lambda$ (see [J, p.252]). For each $\alpha < \lambda^+$ choose a one-to-one function $g_\alpha : \alpha \rightarrow A_\alpha$.

On E , define a symmetric bilinear form as follows: For $\alpha < \beta < \lambda^+$ let

$$\Phi(e_\alpha, e_\beta) = a_\nu \text{ if and only if } g_\beta(\alpha) = \nu$$

The angles $\Phi(e_\alpha, e_\alpha)$ may be defined arbitrarily.

We claim that $\langle E, \Phi \rangle$ is a λ -Gross space. Since $\text{cf}(\lambda) > \omega$, in every subspace $U \subseteq E$ of dimension λ we may find λ many vectors such that the supports of each of them has the same size. We also may assume that these supports are pairwise disjoint (see Case 2). Hence it is enough to prove the following:

($\star\star$) Assume that U is a subspace spanned by a basis $\langle y_\nu : \nu < \lambda \rangle$ such that the sets in $\langle \text{supp}(y_\nu) : \nu < \lambda \rangle$ are pairwise disjoint and of the same cardinality. Then $\dim U^\perp < \dim E$.

The proof of this is completely analogous to that of (\star) in the proof of Theorem 1, if we use Lemma 2 instead of Lemma 1.

Case 2: Assume $\text{cf}(\lambda) = \omega$. In this case, we have to choose the almost disjoint sets $\langle A_\alpha : \alpha < \lambda^+ \rangle$ more carefully. Let $\langle \lambda_n : n < \omega \rangle$ and $\langle f_\alpha : \alpha < \lambda^+ \rangle$ be as in Lemma 3 where $\kappa = \omega$.

Let $\langle \mu_n : n < \omega \rangle$ be increasing, continuous and with limit λ , such that $\mu_n < \lambda_n$ for every n . For each $\alpha < \lambda^+$ define:

$$A_\alpha = \bigcup_{n < \omega} [\mu_n + \mu_n \cdot f_\alpha(n), \mu_n + \mu_n \cdot (f_\alpha(n) + 1))$$

and let $g_\alpha : \alpha \rightarrow A_\alpha$ be one-to-one and onto.

Now let E be a vector space over F of dimension λ^+ , spanned by a basis $\langle e_\alpha : \alpha < \lambda^+ \rangle$. On E , define a symmetric bilinear form as follows: If $\alpha < \beta < \lambda^+$, let:

$$\Phi(e_\alpha, e_\beta) = a_{g_\beta(\alpha)}$$

The angles $\Phi(e_\alpha, e_\alpha)$ may be defined arbitrarily.

We claim that $\langle E, \Phi \rangle$ is a λ -Gross space. Assume that this is not true. So there is a subspace U , spanned by a basis $\langle y_\nu : \nu < \lambda \rangle$, such that U^\perp contains a family $\langle z_\zeta : \zeta < \lambda^+ \rangle$ of linearly independent vectors. Without loss of generality we may and do assume that $\langle \text{supp}(y_\nu) : \nu < \lambda \rangle$ is a family of pairwise disjoint sets. This may be seen as follows: For each $n < \omega$ there exists $Y_n \subseteq [\lambda_n, \lambda_n^+)$ of size λ_n^+ such that $\langle \text{supp}(y_\nu) : \nu \in Y_n \rangle$ is a Δ -system. By forming linear combinations, we find $X_n \subseteq \text{span}\langle y_\nu : \nu \in Y_n \rangle$ such that $\langle \text{supp}(x) : x \in X_n \rangle$ is a pairwise disjoint family. By cutting off from each X_n at most $\sum_{n' < n} \lambda_{n'}$ many elements, we obtain a family as desired.

Let

$$y_\nu = \sum_{l=1}^{m_\nu} b_{\nu l} e_{\alpha(\nu, l)}$$

where each $b_{\nu l}$ is nonzero. Choose $\alpha^* < \lambda^+$ such that $\forall \nu < \lambda \forall 1 \leq l \leq m_\nu \alpha(\nu, l) < \alpha^*$.

Without loss of generality we may assume that there are nonzero c_1, \dots, c_n such that each z_ζ has a representation

$$z_\zeta = \sum_{l=1}^n c_l e_{\beta(\zeta, l)}$$

and for all $\zeta < \eta$ we have $\alpha^* < \beta(\zeta, 1) < \dots < \beta(\zeta, n) < \beta(\eta, 0) < \dots < \beta(\eta, n)$.

By applying Lemma 3 to the family $\langle \text{supp}(z_\zeta) : \zeta < \lambda^+ \rangle$ and $\mu = \omega$, we find $B \in [\lambda^+]^\omega$ and $n^* < \omega$ such that (ii) from the conclusion of Lemma 3 holds.

Note that now $\langle A_\beta \setminus \lambda_{n^*} : \beta \in \bigcup_{\zeta \in B} \text{dom}(z_\zeta) \rangle$ is a family of disjoint sets. Since also $\langle \text{dom}(y_\nu) : \nu < \lambda \rangle$ is a disjoint family and the g_α 's are one-to-one we may find y_ν such that for all $\beta \in \bigcup_{\zeta \in B} \text{dom}(z_\zeta)$ we have $g_\beta(\text{dom}(y_\nu)) \subseteq A_\beta \setminus \lambda_{n^*}$.

But now not even $m_\nu \cdot n$ many vectors $z_{\zeta_1}, \dots, z_{\zeta_{m_\nu n}}$, where each $\zeta_i \in B$, are orthogonal on y_ν . For if this would be true, then $\langle b_1 c_1, \dots, b_{m_\nu} c_n \rangle \in F^{m_\nu n}$ would be a nontrivial solution of the equation $Ax = 0$ where A is the $(m_\nu n \times m_\nu n)$ -matrix

$$\begin{pmatrix} \Phi(e_{\alpha(\nu,1)}, e_{\beta(\zeta_1,1)}) & \cdots & \Phi(e_{\alpha(\nu,m_\nu)}, e_{\beta(\zeta_1,n)}) \\ \vdots & & \vdots \\ \Phi(e_{\alpha(\nu,1)}, e_{\beta(\zeta_{m_\nu n},1)}) & \cdots & \Phi(e_{\alpha(\nu,m_\nu)}, e_{\beta(\zeta_{m_\nu n},n)}) \end{pmatrix}$$

But by construction and Lemma 2, this matrix has nonzero determinant, a contradiction. □

By Lemma 1 in §3, the upper bound for a λ -Gross space over F is $|F|^\lambda$. Hence in ZFC, the dimension of a space as in Theorem 5 cannot be enlarged. But the following theorem shows that it can be enlarged at the loss that we only obtain a λ^+ -Gross space. An upper bound for a λ^+ -Gross space over a field of size λ is $\lambda^{\lambda^+} = 2^{\lambda^+}$. So again, in ZFC we cannot expect a better result. Theorem 6 is true also for countable fields, so we get a complement to Theorem 1.

The construction will use filtrations given by the following Lemma:

Lemma 4. *Let λ be a cardinal.*

(A) *There exists a family $\langle A_\gamma^\alpha : \gamma < \lambda^+, \alpha < \lambda^{++} \rangle$ of sets of size $\leq \lambda$ such that for any $\alpha, \beta < \lambda^{++}$ and $\gamma, \delta < \lambda^+$ the following requirements are satisfied:*

- (1) $\langle A_\gamma^\alpha : \gamma < \lambda^+ \rangle$ is increasing, continuous such that $\alpha = \bigcup_{\gamma < \lambda^+} A_\gamma^\alpha$ and $A_0^\alpha = \emptyset$.
- (2) if $\gamma < \delta$ then $A_\gamma^\alpha \subseteq A_\delta^\alpha$
- (3) if $\alpha < \beta$ and $\alpha \in A_\gamma^\beta$ then $A_\gamma^\alpha = A_\gamma^\beta \cap \alpha$.

(B) *Let $\langle C_\alpha : \alpha < \lambda^{++} \rangle$ be a family of clubs of λ^+ . There exists a family $\langle C'_\alpha : \alpha < \lambda^{++} \rangle$ of clubs and a sequence $\langle A_\gamma^\alpha : \gamma \in C'_\alpha, \alpha < \lambda^{++} \rangle$ of sets of size $\leq \lambda$ such that for any $\alpha, \beta < \lambda^{++}$ and $\gamma, \delta < \lambda^+$ we have $C'_\alpha \subseteq C_\alpha$, and if $\alpha < \beta$ the set $C'_\beta \setminus C'_\alpha$ is bounded and the following requirements are satisfied:*

- (1) $\langle A_\gamma^\alpha : \gamma \in C'_\alpha \rangle$ is increasing, continuous such that $\alpha = \bigcup_{\gamma \in C'_\alpha} A_\gamma^\alpha$ and $A_0^\alpha = \emptyset$.

(2) if $\gamma, \delta \in C'_\alpha$ and $\gamma < \delta$ then $A_\gamma^\alpha \subseteq A_\delta^\alpha$

(3) if $\alpha < \beta$ and $\alpha \in A_\gamma^\beta$ then $C'_\beta \setminus C'_\alpha \subseteq \gamma$ and $A_\gamma^\alpha = A_\gamma^\beta \cap \alpha$.

Proof. We prove only (A). The proof of (B) is similar.

Assume that $\langle A_\gamma^\nu : \gamma < \lambda^+ \rangle$ for $\nu < \alpha$ have been constructed satisfying the requirements. Let $\langle \alpha_\zeta : \zeta < \lambda^+ \rangle$ be an enumeration of α . Then the set

$$\{\gamma < \lambda^+ : (\forall \zeta_1, \zeta_2 < \lambda^+) \text{ if } \zeta_1 < \zeta_2 \text{ and } \alpha_{\zeta_1} < \alpha_{\zeta_2} \text{ then } \alpha_{\zeta_1} \in A_\gamma^{\alpha_{\zeta_2}}\}$$

clearly is a club of λ^+ . Let $\langle \gamma_\varepsilon : \varepsilon < \lambda^+ \rangle$ be its increasing enumeration and define $A_0^\alpha = \emptyset$, and

$$A_\gamma^\alpha = \bigcup_{\zeta < \gamma_\varepsilon} A_\gamma^{\alpha_\zeta}$$

if and only if $\gamma_\varepsilon < \gamma \leq \gamma_{\varepsilon+1}$.

It is not difficult to see that this works. □

Theorem 6. *Let λ be an infinite cardinal. A λ^+ -Gross space of dimension λ^{++} exists over every field of size λ .*

Proof. The proof is divided into two cases:

Case 1: There exists a family $\langle C_\alpha : \alpha < \lambda^{++} \rangle$ of clubs of λ^+ such that for every club C of λ^+ , for some α and λ^+ many $\gamma \in C$ we have $\min(C \setminus (\gamma + 1)) \leq \min(C_\alpha \setminus (\gamma + 1))$. Notice that this is equivalent to the assumption that there exist λ^{++} clubs of λ^+ such that no club gets eventually inside all of them.

By Lemma 4(B), without loss of generality we may assume that for $\alpha < \beta$ the set $C_\beta \setminus C_\alpha$ is bounded in λ^+ and we have families $\langle A_\gamma^\alpha : \gamma \in C_\alpha \rangle$ satisfying (1), (2) and (3).

Now let F be a field of size λ and E a vector space over F , spanned by a basis $\langle e_\alpha : \alpha < \lambda^{++} \rangle$.

On E , we will define a symmetric bilinear form Φ by induction on $\alpha < \lambda^{++}$, using the filtrations above. Assume that $\Phi(e_\beta, e_{\beta'})$ has been defined for $\beta \leq \beta' < \alpha$ and $\Phi(e_\beta, e_\alpha)$ has been defined for $\beta \in A_\gamma^\alpha$.

Let δ be the successor of γ in C_α . If $A_\delta^\alpha \setminus A_\gamma^\alpha$ has size less than λ , then $\Phi(e_\beta, e_\alpha)$, for $\beta \in A_\delta^\alpha \setminus A_\gamma^\alpha$, and $\Phi(e_\alpha, e_\alpha)$ may be defined arbitrarily. Otherwise, choose one-to-one

enumerations such that $F = \text{ran}\langle a_\nu : \nu < \lambda \rangle$, $A_\gamma^\alpha = \text{ran}\langle \zeta_\nu : \nu < \lambda \rangle$ and $A_\delta^\alpha \setminus A_\gamma^\alpha = \text{ran}\langle \xi_\nu : \nu < \lambda \rangle$.

Assume that $\Phi(e_{\xi_{\nu'}}, e_\alpha)$ has been defined for $\nu' < \nu$.

Define $\Phi(e_{\xi_\nu}, e_\alpha)$ such that $\Phi(z, y) \neq 0$ whenever z and y are vectors such that $\alpha \in \text{dom}(z) \subseteq \{\alpha\} \cup \{\zeta_{\nu'} : \nu' < \nu\}$ and $\xi_\nu \in \text{dom}(y) \subseteq \{\xi_{\nu'} : \nu' \leq \nu\}$ and all coefficients of z and y are contained in $\{a_{\nu'} : \nu' < \nu\}$. There are less than λ many such z and y . Hence, this definition can be fulfilled.

We will show that $\langle E, \Phi \rangle$ is a λ^+ -Gross space. Assume that this is not true. So there is a subspace $U \subseteq E$ of dimension λ^+ , say $U = \text{span}\langle y_\nu : \nu < \lambda^+ \rangle$, such that in U^\perp , there is a family of λ^{++} linearly independent vectors, say $\langle z_\iota : \iota < \lambda^{++} \rangle$.

Without loss of generality, we may assume that $\langle \text{supp}(y_\nu) : \nu < \lambda^+ \rangle$ is a family of pairwise disjoint sets of the same cardinality and each y_ν has the same nonzero coefficients in its representation. Mutatis mutandis, we may assume the same for the z_ι 's.

Choose $\alpha^* < \lambda^{++}$ such that $U \subseteq \text{span}\langle e_\alpha : \alpha < \alpha^* \rangle$.

We may certainly find a club $C \subseteq C_{\alpha^*}$ such that, if $\gamma < \delta$ are successive members of C , then

$$|\{\nu < \lambda^+ : \text{dom}(y_\nu) \subseteq A_\delta^{\alpha^*} \setminus A_\gamma^{\alpha^*}\}| = \lambda \quad (1)$$

Since we are in Case 1 there exists $\alpha_C \in (\alpha^*, \lambda^{++})$ such that for λ^+ many $\gamma \in C$ we have $\min(C \setminus (\gamma + 1)) \leq \min(C_{\alpha_C} \setminus (\gamma + 1))$.

Now let $\iota < \lambda^{++}$ such that $\text{dom}(z_\iota) \not\subseteq \alpha_C$ and let α the largest member of $\text{dom}(z_\iota)$. Hence $\alpha \geq \alpha_C$, $C_\alpha \setminus C_{\alpha_C}$ is bounded, and hence $S = \{\gamma \in C : \min(C \setminus (\gamma + 1)) \leq \min(C_\alpha \setminus (\gamma + 1))\}$ has size λ^+ .

Now choose $\gamma' \in S$ so large that, if γ is the maximal element of C_α such that $\gamma \leq \gamma'$, then

$$\{\alpha^*\} \cup \text{dom}(z_\iota) \setminus \{\alpha\} \subseteq A_\gamma^\alpha \quad (2)$$

Let δ be the successor of γ in C_α . So clearly $\min(C \setminus (\gamma' + 1)) \leq \delta$.

By construction of the filtrations $\langle A_\nu^\alpha : \nu \in C_\alpha \rangle$ and by (1) and (2) we know that

$$|\{\nu < \lambda^+ : \text{dom}(y_\nu) \subseteq A_\delta^\alpha \setminus A_\gamma^\alpha\}| = \lambda \quad (3)$$

Now look at the definition of $\Phi(., e_\alpha)$ on $A_\delta^\alpha \setminus A_\gamma^\alpha$. Choose $\nu < \lambda$ and $\mu < \lambda^+$ such that

- (i) the coefficients of z_ν and those of y_μ (and hence of any $y_{\mu'}$) are contained in $\{a_{\nu'} : \nu' < \nu\}$,
- (ii) $\text{dom}(z_\nu) \setminus \{\alpha\} \subseteq \{\zeta_{\nu'} : \nu' < \nu\}$, and
- (iii) $\xi_\nu \in \text{dom}(y_\mu)$ and $\text{dom}(y_\mu) \setminus \{\xi_\nu\} \subseteq \{\xi_{\nu'} : \nu' < \nu\}$.

By (3), such a choice is possible. But now $\Phi(e_{\xi_\nu}, e_\alpha)$ was defined such that $\Phi(z_\nu, y_\mu) \neq 0$, a contradiction.

Case 2: There exists no family of clubs as in Case 1.

Let $\langle A_\gamma^\alpha : \gamma < \lambda^+, \alpha < \lambda^{++} \rangle$ be as in Lemma 4(A).

Let $f_\alpha : \lambda^+ \rightarrow \lambda^+$ be defined by $f_\alpha(\gamma) = \text{o.t.}(A_{\gamma+1}^\alpha)$. Note that if $\alpha < \beta$, then $f_\alpha(\gamma) < f_\beta(\gamma)$ for every γ so large that $\alpha \in A_\gamma^\beta$.

Let $C_\alpha = \{\gamma < \lambda^+ : \gamma \text{ is a limit ordinal and } (\forall \delta < \gamma) f_\alpha(\delta) < \gamma\}$. Clearly C_α is a club.

Since $\langle C_\alpha : \alpha < \lambda^{++} \rangle$ cannot serve for Case 1, there exists a club C of λ^+ such that

$$\forall \alpha < \lambda^{++} \exists \gamma < \lambda^+ \forall \delta \in (\gamma, \lambda^+) \cap C [\min(C \setminus (\delta + 1)) > \min(C_\alpha \setminus (\delta + 1))] \quad (4)$$

Let $\langle \gamma_\varepsilon : \varepsilon < \lambda^+ \rangle$ be the increasing enumeration of $C \cup \{0\}$. Now define $f^* : \lambda^+ \rightarrow \lambda^+$ by:

$$f^*(\gamma) = \gamma_{\varepsilon+2} \text{ if and only if } \gamma_\varepsilon \leq \gamma < \gamma_{\varepsilon+1}$$

Using the definition of C_α and (4), it is easy to see that:

$$\forall \alpha < \lambda^{++} \exists \gamma < \lambda^+ \forall \delta \in (\gamma, \lambda^+) [f_\alpha(\gamma) < f^*(\gamma)] \quad (5)$$

Now let F be a field of size λ and E a vector space over F of dimension λ^{++} , spanned by a basis $\langle e_\alpha : \alpha < \lambda^{++} \rangle$.

Let $\langle F_\gamma : \gamma < \lambda^+ \rangle$ be an almost disjoint family of subsets of F (so each F_γ has size λ and $F_\gamma \cap F_\delta$ has size $< \lambda$ for any $\gamma \neq \delta$) such that:

in case λ is uncountable, $\bigcup F_\gamma$ is a set of elements which are algebraically independent over the prime field of F

in case $\lambda = \omega$, $\bigcup F_\gamma$ is a subset of $\text{ran}\langle a_n : n < \omega \rangle$ where $\langle a_n : n < \omega \rangle$ is a sequence as in Lemma 1.

For each $\gamma < \lambda^+$, let $\langle a_{\varepsilon, \zeta}^\gamma : \varepsilon < \zeta < f^*(\gamma) \rangle$ be a one-to-one enumeration of F_γ .

In difference to the previous constructions in this paper, here the angles $\Phi(e_\alpha, e_\beta)$ will be defined upwards, as follows:

For $\alpha < \lambda^{++}$ and $\gamma < \lambda^+$ define

$$W_\gamma^\alpha = \{\beta < \lambda^{++} : \alpha \in A_{\gamma+1}^\beta \setminus A_\gamma^\beta\}$$

So $\lambda^{++} \setminus \alpha = \bigcup_{\gamma < \lambda^+} W_\gamma^\alpha$, and this is a disjoint union.

Now let $\alpha < \beta < \lambda^{++}$. We define

$$\Phi(e_\alpha, e_\beta) = a_{\text{o.t.}(A_{\gamma+1}^\alpha), \text{o.t.}(A_{\gamma+1}^\beta)}^\gamma$$

in case $\text{o.t.}(A_{\gamma+1}^\beta) < f^*(\gamma)$ where γ is the uniquely determined ordinal such that $\beta \in W_\gamma^\alpha$. Note that $\text{o.t.}(A_{\gamma+1}^\alpha) < \text{o.t.}(A_{\gamma+1}^\beta)$ in this case. Otherwise, $\Phi(e_\alpha, e_\beta)$ may be defined arbitrarily. Also $\Phi(e_\alpha, e_\alpha)$ may be defined arbitrarily.

We claim that $\langle E, \Phi \rangle$ is λ^+ -Gross. If this is not true, then there is a subspace $U \subseteq E$ of dimension λ^+ , say $U = \text{span}\langle y_\nu : \nu < \lambda^+ \rangle$, such that in U^\perp , there is a family of λ^{++} linearly independent vectors, say $\langle z_\iota : \iota < \lambda^{++} \rangle$.

Again, we may assume that $\langle \text{supp}(y_\nu) : \nu < \lambda^+ \rangle$ is a family of pairwise disjoint sets of the same cardinality and each y_ν has the same nonzero coefficients in its representation. Mutatis mutandis, we may assume the same for the z_ι 's. We may also assume that each $\text{supp}(z_\iota)$ is disjoint from α^* where α^* is chosen such $U \subseteq \text{span}\langle e_\alpha : \alpha < \alpha^* \rangle$. Let

$$y_\nu = \sum_{l=1}^m b_l e_{\alpha(\nu, l)}$$

$$z_\iota = \sum_{l=1}^n c_l e_{\beta(\iota, l)}$$

By (5) and property A(3), Lemma 4, of the filtrations $\langle A_\gamma^\alpha : \gamma < \lambda^+, \alpha < \lambda^{++} \rangle$ we can choose $\gamma^* < \lambda^+$ such that

(j) for each $\beta \in \bigcup_{\iota < \lambda} \text{dom}(z_\iota)$ and $\gamma \in (\gamma^*, \lambda^+)$ $f_\beta(\gamma) < f^*(\gamma)$, and

(jj) if $\beta_1, \beta_2 \in \bigcup_{\iota < \lambda} \text{dom}(z_\iota)$ and $\beta_1 < \beta_2$, then $\beta_1 \in A_{\gamma^*}^{\beta_2}$ and $\alpha^* \in A_{\gamma^*}^{\beta_1}$.

Now find y_ν such that $\text{dom}(y_\nu) \cap A_{\gamma^*}^{\alpha^*} = \emptyset$. There are finitely many $\gamma_1, \dots, \gamma_p > \gamma^*$ such that

$$\text{dom}(y_\nu) \subseteq \bigcup_{i=1}^p (A_{\gamma_{i+1}}^{\alpha^*} \setminus A_{\gamma_i}^{\alpha^*})$$

Then for all $\beta \in \bigcup_{\iota < \lambda} \text{dom}(z_\iota)$ and $1 \leq i \leq p$ we have

$$\text{dom}(y_\nu) \cap (A_{\gamma_{i+1}}^{\alpha^*} \setminus A_{\gamma_i}^{\alpha^*}) = \text{dom}(y_\nu) \cap (A_{\gamma_{i+1}}^\beta \setminus A_{\gamma_i}^\beta)$$

Call this set M_i . So for $\alpha \in M_i$ and $\beta \in \bigcup_{\iota < \lambda} \text{dom}(z_\iota)$ we have $\beta \in W_{\gamma_i}^\alpha$ and hence, by (j) above, $\Phi(e_\alpha, e_\beta) = a_{\text{o.t.}(A_{\gamma_{i+1}}^\alpha), \text{o.t.}(A_{\gamma_{i+1}}^\beta)}^{\gamma_i} \in F_{\gamma_i}$. Notice that for distinct $\alpha, \alpha' \in M_i$ we have $\text{o.t.}(A_{\gamma_{i+1}}^\alpha) \neq \text{o.t.}(A_{\gamma_{i+1}}^{\alpha'})$, and for distinct $\beta, \beta' \in \bigcup_{\iota < \lambda} \text{dom}(z_\iota)$ we have $\text{o.t.}(A_{\gamma_{i+1}}^\beta) \neq \text{o.t.}(A_{\gamma_{i+1}}^{\beta'})$. Hence, by almost-disjointness of the F_{γ_i} , we may find $\iota_1, \dots, \iota_{m \cdot n} < \lambda$ such that for $1 \leq j \leq mn$ the sets

$$\{a_{\text{o.t.}(A_{\gamma_{i+1}}^\alpha), \text{o.t.}(A_{\gamma_{i+1}}^{\beta(\iota_j, l)})}^{\gamma_i} : \alpha \in M_i, 1 \leq i \leq p, 1 \leq l \leq n\} \quad (6)$$

are pairwise disjoint and each of them contains $m \cdot n$ elements. If now

$$\Phi(y_\nu, z_{\iota_j}) = \sum_{1 \leq l \leq m, 1 \leq k \leq n} b_l c_k \Phi(e_{\alpha(\nu, l)}, e_{\beta(\iota_j, k)}) = 0$$

would be true for every $1 \leq j \leq mn$, then $\langle b_1 c_1, \dots, b_m c_n \rangle \in F^{mn}$ would be a nontrivial solution of the equation $Ax = 0$ where A is the $(mn \times mn)$ -matrix

$$\begin{pmatrix} \Phi(e_{\alpha(\nu, 1)}, e_{\beta(\iota_1, 1)}) & \dots & \Phi(e_{\alpha(\nu, m)}, e_{\beta(\iota_1, n)}) \\ \vdots & & \vdots \\ \Phi(e_{\alpha(\nu, 1)}, e_{\beta(\iota_{mn}, 1)}) & \dots & \Phi(e_{\alpha(\nu, m)}, e_{\beta(\iota_{mn}, n)}) \end{pmatrix}$$

But the rows of this matrix are the sets in (6), hence its determinant is nonzero, a contradiction. \square

References

- [B/S] B. Balcar and P. Simon, Cardinal invariants in Boolean spaces, *General Topology and its Relations to Modern Analysis and Algebra V*, Proc. Fifth Prague Topol. Symp. Symp. 1981, ed. J. Novak, Heldermann Verlag, Berlin, 1982, 39-47.
- [B] J.E. Baumgartner, Iterated forcing, *Surveys of set theory* (A.R.D. Mathias, editor), London Mathematical Society Lecture Note Series, no.87, Cambridge University Press, Cambridge, 1983, 1-59.
- [B/Sp] J.E. Baumgartner and O. Spinas, Independence and consistency proofs in quadratic form theory, *Journal of Symbolic Logic*, vol.57, no.4, 1991, 1195-1211.
- [B/G] W. Baur and H. Gross, Strange inner product spaces, *Comment. Math. Helv.* 52, 1977, 491-495.
- [B/Sh] A. Blass and S. Shelah, There may be simple P_{\aleph_1} - and P_{\aleph_2} -points and the Rudin-Keisler ordering may be downward directed, *Annals of Pure and Applied Logic*, vol.53, 1987, 213-243.
- [vD] E.K. van Douwen, The integers and topology, *Handbook of set-theoretic topology*, K. Kunen and J.E. Vaughan (editors), North-Holland, Amsterdam, 1984, 111-167.
- [F] D.H. Fremlin, Cichoń's diagram, *Initiation à l'Analyse*, Université Pire et Marie Curie, Paris, 1985.
- [G] H. Gross, Quadratic forms in infinite dimensional vector spaces, *Progress in Mathematics*, vol.1, Birkhäuser, Boston, 1979.
- [G/O] H. Gross and E. Ogg, Quadratic spaces with few isometries, *Comment. Math. Helv.* 48, 1973, 511-519.
- [Go] M. Goldstern, Tools for your forcing construction, in: *Set theory of the reals*, Proceedings of the Bar Ilan Conference in honour of Abraham Fraenkel 1991, H. Judah (editor), 305-360.
- [J] T. Jech, *Set theory*, Academic Press, New York, 1978.
- [J/Sh] H. Judah and S. Shelah, Souslin forcing, *Journal of Symbolic Logic*, vol.53, no.4, 1988, 1188-1207.
- [K] K. Kunen, *Set theory. An introduction to independence proofs*, North Holland, Amsterdam, 1980.
- [Sh1] S. Shelah, *Proper forcing*, *Lecture Notes in Mathematics*, vol.942, Springer.

- [Sh2] S. Shelah, On cardinal invariants of the continuum, Proceedings of the 6/83 Boulder conference in set theory, ed. J. Baumgartner, D. Martin and S. Shelah, Contemporary mathematics, vol.31, AMS, 1984, 183-207.
- [Sh3] S. Shelah, Vive la difference I, Nonisomorphism of ultrapowers of countable models, in: Set theory of the continuum, ed. H. Judah, W. Just, H. Woodin, Springer, New York, 1992, 357-405.
- [Sh4] S. Shelah, Strong negative partition relations below the continuum, Acta Math. Hungar. 58, no. 1-2, 1991, 95-100.
- [Sh5] S. Shelah, There are Jońsson algebras in many inaccessible cardinals, in: Cardinal Arithmetic, Oxford University Press, 1994.
- [Sh6] S. Shelah, Further cardinal arithmetic, Israel Journal of mathematics, in press ([Sh430] in Shelah's list of publications).
- [Sh7] S. Shelah, Colouring and \aleph_2 -c.c. not productive, in preparation ([Sh572] in Shelah's list of publications).
- [Sp1] O. Spinas, Konsistenz- und Unabhangigkeitsresultate in der Theorie der quadratischen Formen, Dissertation, University of Zurich, 1989.
- [Sp2] O. Spinas, Iterated forcing in quadratic form theory, Israel Journal of Mathematics 79, 1991, 297-315.
- [Sp3] O. Spinas, An undecidability result in lattice theory, Abstracts of papers presented to the AMS, vol.11, no.2. p.161, March 1990.
- [Sp4] O. Spinas, Cardinal invariants and quadratic forms, in: Set theory of the reals, Proceedings of the Bar Ilan Conference in honour of Abraham Fraenkel 1991, H. Judah (editor), 563-581.

Addresses: *First author:* Department of Mathematics, The Hebrew University of Jerusalem, Givat Ram, Jerusalem, ISRAEL.

Second author: Department of Mathematics, University of California, Irvine, CA 92717, USA