

AN  $\aleph_2$ -SOUSLIN TREE FROM A STRANGE HYPOTHESIS  
ABSTRACTS OF AMS  
(1985)P.198(84-T-03-160)  
SHE:4

SAHARON SHELAH

Institute of Mathematics  
The Hebrew University  
Jerusalem, Israel

Rutgers University  
Mathematics Department  
New Brunswick, NJ USA

---

I would like to thank Alice Leonhardt for the beautiful typing.  
First Typed - 98/Jan/14  
Latest Revision - 98/Mar/19

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{T}\mathcal{E}\mathcal{X}$

## §1

**Theorem.** *Suppose CH holds and the filter  $\mathcal{D}_{\omega_1}$  (see below) is  $\aleph_2$ -saturated. Then there is an  $\aleph_2$ -Souslin tree.*

Notation:  $\mathcal{D}_{\omega_1}$  is the filter generated by the closed unbounded subset of  $\omega_1$ . Let  $S_\beta^\alpha = \{\delta < \aleph_\alpha : \text{cf}(\delta) = \aleph_\beta\}$ .

*Proof.* It is known that the assumption implies  $2^{\aleph_1} = \aleph_2$ . By Gregory [Gre] we know that if there is a stationary  $S \subseteq S_0^2$  with no initial segment stationary, then there is an  $\aleph_2$ -Souslin tree. So assume there is no such  $S$ . By Gregory [Gre],  $\diamond(S_0^2)$  holds, and let  $\langle A_\delta : \delta \in S_0^2 \rangle$  exemplify this. For each  $\alpha \in S_1^2$  define  $\mathcal{P}_\alpha = \{B \subseteq \alpha : \{\delta < \alpha : B \cap \delta = A_\delta\}$  is a stationary subset of  $\alpha\}$ . If  $|\mathcal{P}_\alpha| > \aleph_1$  let  $B_i \in \mathcal{P}_\alpha (i < \aleph_2)$  be pairwise distinct and  $\langle \gamma(\zeta) : \zeta < \omega_1 \rangle$  be an increasing continuous sequence of ordinals such that  $\gamma(\zeta) < \alpha$  and  $\bigcup_{\zeta} \gamma(\zeta) = \alpha$ ; let  $S_i =: \{\zeta < \omega_1 : B_i \cap \gamma(\zeta) = A_{\gamma(\zeta)}\}$ . Now  $S_i$  is a stationary subset of  $\omega_1$  (as  $B_i \in \mathcal{P}_\alpha$ ) and  $S_i \cap S_j$  is bounded for  $i \neq j$  (as for some  $\zeta_0 < \omega_1, B_i \cap \gamma(\zeta_0) \neq B_j \cap \gamma(\zeta_0)$ ) hence  $\langle S_i : i < \omega_2 \rangle$  exemplifies that  $\mathcal{D}_{\omega_1}$  is  $\aleph_2$ -saturated, contradiction, hence  $|\mathcal{P}_\alpha| \leq \aleph_1$ . Also for every  $A \subseteq \omega_2, \{[\delta \in S_0^2 : A \cap \delta = A_\delta]\}$  is stationary hence for some  $\alpha \in S_1^2, \{\delta \in S_0^2 \cap \alpha : A \cap \delta = A_\delta\}$  is stationary below  $\alpha$ . So  $\langle \mathcal{P}_\alpha : \alpha \in S_1^2 \rangle$  exemplify a variant of  $\diamond(S_1^2)$  which by Kunen implies  $\diamond(S_1^2)$ ; together with  $2^{\aleph_0} = \aleph_1, 2^{\aleph_1} = \aleph_2$  we finish.

*Remark.* We can replace  $(\aleph_1, \mathcal{D}_{\omega_1})$  by  $(\aleph_\alpha, D(\aleph_{\alpha+1}) + S_\alpha^{\alpha+1})$  if  $\aleph_\alpha$  is regular. (Received December 7, 1983).

AN  $\aleph_2$ -SOUSLIN TREE FROM A STRANGE HYPOTHESIS ABSTRACTS OF AMS (1985)P.198(84-T-03-160) S

REFERENCES.

- [Gre] John Gregory. Higher Souslin trees and the generalized continuum hypothesis.  
*Journal of Symbolic Logic*, **41**(3):663–671, 1976.