

On Monk's questions

by

Saharon Shelah (Jerusalem and New Brunswick, N.J.)

Abstract. We deal with Boolean algebras and their cardinal functions: π -weight π and π -character $\pi\chi$. We investigate the spectrum of π -weights of subalgebras of a Boolean algebra B . Next we show that the π -character of an ultraproduct of Boolean algebras may be different from the ultraproduct of the π -characters of the factors.

Annotated content

1. Introduction

2. Existence of subalgebras with a preassigned algebraic density. We first note (in 2.1) that if $\pi(B) \geq \theta = \text{cf}(\theta)$ then for some $B' \subseteq B$ we have $\pi(B') = \theta$. Call this statement (*). Then we give a criterion for $\pi(B) = \mu > \text{cf}(\mu)$ (in 2.2) and conclude for singular μ that for a club of $\theta < \mu$ the (*) above holds (2.2A), and investigate the criterion (in 2.3). Our main aim is, starting with $\mu = \mu^{<\mu}$, $\text{cf}(\lambda) < \lambda$, to force the existence of a Boolean algebra B such that $\pi(B) > \theta$ but for no $B' \subseteq B$ do we have $\pi(B') = \lambda$ (in fact $(\exists B' \subseteq B)[\pi(B') = \theta \Leftrightarrow \theta = \text{cf}(\theta) \vee \text{cf}(\theta) \leq \mu]$ for every $\theta \leq |B|$). Toward this, we define the forcing (Definition 2.5: a condition p tells us how $\langle x_\alpha : \alpha \in W^p \rangle$ generate a Boolean algebra, $\text{BA}[p]$, $W^p \in [\lambda]^{<\mu}$ with $x_\alpha > \theta$ having no non-zero member of $\langle x_\beta : \beta \in W^p \cap \alpha \rangle_{\text{BA}[p]}$ below it). We prove the expected properties of the generic (2.6), also the forcing has the expected properties (μ -complete, μ^+ -c.c.) (in 2.7). The main theorem (2.9) stated, the main point being that if $\mu < \text{cf}(\theta) < \theta$ for $B \subseteq \text{BA}[G]$, then $\pi(B) \neq \theta$; we use the above criterion, and a lemma related to Δ -systems (see [Sh 430], 6.6D, [Sh 513], 6.1) quoted in 2.4, to reduce the problem to some special amalgamation of finitely many copies (the exact number is in relation to the arity of the term defining the relevant elements from the x_α 's). The existence of such amalgamation was done separately earlier (2.8).

Lastly, in 2.10 we show that the $\text{cf}(\theta) > \mu$ above was necessary by proving the existence of a subalgebra with prescribed singular algebraic density λ satisfying $\pi(B) > \lambda$ and $(\forall \mu < \lambda)[\mu^{<\text{cf}(\lambda)} < \lambda]$.

1991 *Mathematics Subject Classification*: Primary 03G05, 03E35; Secondary 03E05, 04A20.

Partially supported by the Deutsche Forschungsgemeinschaft, Grant No. 490/7-1.

I would like to thank Alice Leonhardt for the beautiful typing.

Publication number 479.

3. On π and $\pi\chi$ of products of Boolean algebras. If e.g. $\aleph_0 < \kappa = \text{cf}(\chi) < \chi < \lambda = \text{cf}(\lambda) < \chi^\kappa$ and $(\forall \theta < \chi)[\theta^\kappa < \chi]$ we show that for some Boolean algebras B_i (for $i < \kappa$), $\chi = \sum_{i < \kappa} \pi\chi(B_i) < \lambda$ and (for D a regular ultrafilter on κ) $\lambda = \pi\chi(\prod_{i < \kappa} B_i/D)$ but $\prod_{i < \kappa} (\pi\chi(B_i)/D) = \chi^\kappa$. For this we use interval Boolean algebras on orders of the form $\lambda_i \times \mathbb{Q}$.

We also prove for infinite Boolean algebras B_i (for $i < \kappa$) and D an ultrafilter on κ that if $n_i < \aleph_0$ and $\mu = \prod_{i < \kappa} n_i/D$ is a regular (infinite) cardinal then $\pi\chi(\prod_{i < \kappa} B_i/D) \geq \mu$.

1. Introduction. Monk [M] asks (problems 13, 15 in his list; π is the algebraic density, see 1.1 below): For a (Boolean algebra) B with $\aleph_0 \leq \theta \leq \pi(B)$, does B have a subalgebra B' with $\pi(B') = \theta$?

If θ is regular the answer is easily seen to be positive (see 2.1). We show that in general it may be negative (see 2.9(3)), but for quite many singular cardinals, it is positive (2.10); the theorems are quite complementary. This is dealt with in §2.

In §3 we mainly deal with $\pi\chi$ (see Definition 3.2) and show that the $\pi\chi$ of an ultraproduct of Boolean algebras is not necessarily the ultraproduct of the $\pi\chi$'s. Note that in Koppelberg–Shelah [KpSh 415], Theorem 1.1, we prove that if SCH holds and $\pi(B_i) > 2^\kappa$ for $i < \kappa$ then

$$\pi\left(\prod_{i < \kappa} B_i/D\right) = \prod_{i < \kappa} (\pi(B_i)/D).$$

We also prove that for infinite Boolean algebras A_i ($i < \kappa$) and a non-principal ultrafilter D on κ , if $n_i < \aleph_0$ for $i < \kappa$ and $\mu := \prod_{i < \kappa} n_i/D$ is regular, then $\pi\chi(A) \geq \mu$. Here $A := \prod_{i < \kappa} A_i/D$. By a theorem of Peterson [P] the regularity of μ is necessary.

1.1. NOTATION. Boolean algebras are denoted by B and sometimes A . For a Boolean algebra B , set

$$B^+ := \{x \in B : x \neq 0\},$$

$$\pi(B) := \min\{|X| : X \subseteq B^+ \text{ is such that } (\forall y \in B^+)(\exists x \in X)[x \leq y]\}.$$

X like that is called *dense in B* . More generally, if $X, Y \subseteq B$ we say X is *dense in Y* if $y \in Y \ \& \ y \neq 0 \Rightarrow (\exists x \in X)[0 < x \leq y]$. For a $Y \subseteq B$, $\langle Y \rangle_B$ is the subalgebra of B which Y generates.

0_A is the constant function with domain A and value zero. 1_A is defined similarly.

2. Existence of subalgebras with a preassigned algebraic density

2.1. OBSERVATION. If $\pi(B) > \theta = \text{cf}(\theta) \geq |Y| + \aleph_0$ and $Y \subseteq B$ then for some subalgebra A of B , $Y \subseteq A$, $\pi(A) = \theta$ and $|A| = \theta$.

PROOF. Without loss of generality $|Y| = \theta$. Let $Y = \{y_\alpha : \alpha < \theta\}$. Choose by induction on $\alpha \leq \theta$ subalgebras A_α of B , increasing continuously in α ,

with $|A_\alpha| < \theta$ and $y_\alpha \in A_{\alpha+1}$, such that for each $\alpha < \theta$, some $x_\alpha \in A_{\alpha+1}^+$ is not above any $y \in A_\alpha^+$. This is possible because for no $\alpha < \theta$ can A_α^+ be dense in B . Now $A = A_\theta$ is as required. ■_{2.1}

2.2. CLAIM. Assume B is a Boolean algebra with $\pi(B) = \mu > \text{cf}(\mu) \geq \aleph_0$ (see Notation 1.1). Then for arbitrarily large regular $\theta < \mu$,

$(*)_\theta^B$ for some set Y we have:

$(*)_\theta^B[Y]$ $Y \subseteq B^+, |Y| = \theta$, and there is no $Z \subseteq B^+$ of cardinality $< \theta$, dense in Y (i.e. $(\forall y \in Y)(\exists z \in Z)[z \leq y]$).

2.2A. CONCLUSION. If B is a Boolean algebra, $\pi(B) = \mu > \text{cf}(\mu) > \aleph_0$ and $\langle \mu_\zeta : \zeta < \text{cf}(\mu) \rangle$ is increasing continuously with limit μ (so $\mu_\zeta < \mu$), then for some club C of $\text{cf}(\mu)$, for every $\zeta \in C$, for some $B'_\zeta \subseteq B$ we have $\pi(B'_\zeta) = \mu_\zeta$.

Proof of 2.2. Let $Z^* \subseteq B^+$ be dense with $|Z^*| = \mu$. If the conclusion fails, then for some $\theta^* < \mu$, for no regular $\theta \in (\theta^*, \mu)$ does $(*)_\theta^B$ hold. We now assume we chose such a θ^* , and show by induction on $\lambda \leq \mu$ that

(\otimes_λ) if $Y \subseteq B^+$ and $|Y| \leq \lambda$, then for some $Z \subseteq B^+$, $|Z| \leq \theta^*$ and Z is dense in Y .

Case 1: $\lambda \leq \theta^*$. Let $Z = Y$.

Case 2: $\theta^* < \lambda \leq \mu$ and $\text{cf}(\lambda) < \lambda$. Let $Y = \bigcup \{Y_\zeta : \zeta < \text{cf}(\lambda)\}$, $|Y_\zeta| < \lambda$. By the induction hypothesis for each $\zeta < \text{cf}(\lambda)$ there is $Z_\zeta \subseteq B^+$ of cardinality $\leq \theta^*$ which is dense in Y_ζ .

Now $Z' := \bigcup_{\zeta < \text{cf}(\lambda)} Z_\zeta$ has cardinality $\leq \theta^* + \text{cf}(\lambda) < \lambda$, hence by the induction hypothesis there is $Z \subseteq B^+$ dense in Z' with $|Z| \leq \theta^*$. Clearly Z is dense in Y , $|Z| \leq \theta^*$ and $Z \subseteq B^+$ so we finish the case.

Case 3: $\theta^* < \lambda \leq \mu$, λ regular. If for this Y , $(*)_\lambda^B[Y]$ holds, we get the conclusion of the claim. We are assuming not so; so there is $Z' \subseteq B^+$ dense in Y with $|Z'| < \lambda$. Apply the induction hypothesis to Z' and get Z as required.

So we have proved (\otimes_λ) .

We apply (\otimes_λ) to $\lambda = \mu$, $Y = Z^*$ and get a contradiction. ■_{2.2}

2.3. CLAIM. (1) If B , μ , θ , Y are as in 2.2 (so $(*)_\theta^B[Y]$ holds and θ is regular) then we can find $\bar{y} = \langle y_\alpha : \alpha < \theta \rangle$ whose range is contained in B^+ , and a proper θ -complete filter D on θ containing all cobounded subsets of θ such that

$(\otimes_{\bar{y}, D}^B)$ for every $z \in B^+$, $\{\alpha < \theta : z \leq y_\alpha\} = \emptyset \text{ mod } D$.

(2) If in addition θ is a successor cardinal ⁽¹⁾ then we can demand that D is normal.

Remark. Part (2) is for curiosity only.

Proof. (1) Let $Y = \{y_\alpha : \alpha < \theta\}$. Define D as follows: for $\mathcal{U} \subseteq \theta$,

$\mathcal{U} \in D$ iff for some $0 < \zeta < \theta$ and $z_\varepsilon \in B^+$ for $\varepsilon < \zeta$,
we have $\mathcal{U} \supseteq \{\alpha < \theta : (\forall \varepsilon < \zeta)[z_\varepsilon \not\leq y_\alpha]\}$.

Trivially D is closed under supersets and intersections of $< \theta$ members and every cobounded subset of θ belongs to it. Now $\emptyset \notin D$ because $(*)_\theta^B[Y]$.

(2) Let $\theta = \sigma^+$. Assume there are no such \bar{y}, D . We try to choose by induction on $n < \omega$, Y_α^n ($\alpha < \theta$) and club E_n of θ such that:

- (a) Y_α^n is a subset of B^+ of cardinality $< \theta$, increasing continuous in α ,
- (b) $Y_\alpha^n \subseteq Y_\alpha^{n+1}$,
- (c) $Y_\alpha^0 = \{y_\beta : \beta < \alpha\}$ (where $\{y_\alpha : \alpha < \theta\}$ are taken from part (1)),
- (d) E_n is a club of θ , $E_{n+1} \subseteq E_n$, $E_0 = \{\delta < \theta : \delta \text{ divisible by } \sigma\}$,
- (e) if $\delta \in E_{n+1}$ and $\delta \leq \alpha < \min(E_n \setminus (\delta + 1))$ then for every $y \in Y_\alpha^n$ there is $z \in Y_\delta^{n+1}$ with $z \leq y$.

If we succeed, let $\beta^* = \bigcup_{n < \omega} \min(E_n)$ ($< \theta$), and we shall prove that $\bigcup_{n < \omega} Y_{\beta^*}^n$ is dense in Y , getting a contradiction. For every $y \in \bigcup_{n < \omega, \alpha < \theta} Y_\alpha^n$ let $\beta(y)$ be the minimal $\beta < \theta$ such that $(\exists z \in \bigcup_{n < \omega} Y_\beta^n)[z \leq y]$. Now β is well defined as $y \in \bigcup_{\beta < \theta} \bigcup_{n < \omega} Y_\beta^n$. If $\beta(y) \leq \beta^*$ for every $y \in Y$ ($\subseteq \bigcup_{\alpha < \theta} Y_\alpha^0$) we are done, as $\langle \bigcup_{n < \omega} Y_\beta^n : \beta < \theta \rangle$ is increasing continuous; assume not, so some $y^* \in Y = \bigcup_{\alpha < \theta} Y_\alpha^0$ exemplifies this. Now let $\beta = \beta(y^*)$ and let $z \in \bigcup_{n < \omega} Y_\beta^n$ exemplify this. Clearly $\langle \sup(\beta \cap E_n) : n < \omega \rangle$ is well defined; clearly it is a non-increasing sequence of ordinals, hence eventually constant, say $n \geq n^* \Rightarrow \sup(\beta \cap E_n) = \gamma$. Now, without loss of generality $z \in Y_\beta^{n^*}$ (by clause (b)); note $\gamma \in E_n$ for $n \geq n^*$ (hence for every n). But by clause (e) there is $z' \in Y_\gamma^{n^*+1}$ with $z' \leq z$, contradicting the choice of β .

So we cannot carry out the construction, that is, we are stuck at some n . Fix such an n . Let $E_n \cup \{0\} = \{\delta_\varepsilon : \varepsilon < \theta\}$ (increasing with ε). Let $Y_{\delta_{\varepsilon+1}}^n \setminus Y_{\delta_\varepsilon}^n \subseteq \{y_\zeta^\varepsilon : \zeta < \sigma\}$. For each $\zeta < \sigma$, let D_ζ be the normal filter generated by the family of subsets of θ of the form $\{\varepsilon < \theta : z \not\leq y_\zeta^\varepsilon\}$ for $z \in B^+$. If $\emptyset \in D_\zeta$ for every $\zeta < \sigma$, we can define Y_α^{n+1} and E_{n+1} , a contradiction. So for some ζ , $\bar{y}^\zeta := \langle y_\zeta^\varepsilon : \zeta < \sigma \rangle$ and D_ζ are as required in $(\otimes_{\bar{y}^\zeta, D_\zeta}^B)$. ■_{2.3}

2.4. CLAIM. Suppose D is a σ -complete filter on θ , $\theta = \text{cf}(\theta) \geq \sigma > 2^\kappa$, and for each $\alpha < \theta$, $\bar{\beta}^\alpha = \langle \beta_\varepsilon^\alpha : \varepsilon < \kappa \rangle$ is a sequence of ordinals. Then

⁽¹⁾ But see the end of the paper.

for every $\mathcal{U} \subseteq \theta$ with $\mathcal{U} \neq \emptyset \bmod D$ there are $\langle \beta_\varepsilon^* : \varepsilon < \kappa \rangle$ (a sequence of ordinals) and $w \subseteq \kappa$ such that:

- (a) $\varepsilon \in \kappa \setminus w \Rightarrow \text{cf}(\beta_\varepsilon^*) \leq \theta$,
- (b) if $(\forall \alpha < \sigma)[|\alpha|^\kappa < \sigma]$ then $\varepsilon \in \kappa \setminus w \Rightarrow \sigma \leq \text{cf}(\beta_\varepsilon^*)$,
- (c) if $\beta'_\varepsilon \leq \beta_\varepsilon^*$ for all ε and $[\varepsilon \in w \equiv \beta'_\varepsilon = \beta_\varepsilon^*]$ then $\{\alpha \in \mathcal{U} : \beta'_\varepsilon \leq \beta_\varepsilon^\alpha \leq \beta_\varepsilon^* \text{ for all } \varepsilon \text{ and } [\varepsilon \in w \equiv \beta_\varepsilon^\alpha = \beta_\varepsilon^*]\} \neq \emptyset \bmod D$.

Proof. [Sh 430], 6.1D, and better presented in [Sh 513], 6.1.

2.5. DEFINITION. (1) If $F \subseteq {}^w 2$ let

$cl(F) = \{g \in {}^w 2 : \text{for every finite } u \subseteq w \text{ and some } f \in F \text{ we have } g \upharpoonright u = f \upharpoonright u\}$.

If $f \in {}^w 2$, $w \subseteq \text{Ord}$ and $\alpha \in \text{Ord}$ let $f^{[\alpha]}$ be $(f \upharpoonright (w \cap \alpha)) \cup 0_{w \setminus \alpha}$; let $f^{[\infty]} = f$.

(2) Let $\mu = \mu^{<\mu} < \lambda$. We define a forcing notion $Q = Q_{\lambda, \mu}$:

- (a) the members are pairs $p = (w, F) = (w^p, F^p)$, $w \subseteq \lambda$, $|w| < \mu$, and F is a family of $< \mu$ functions from w to $\{0, 1\}$ satisfying
 - (α) for every $\alpha \in w$ and some $f \in F$, $f(\alpha) = 1$,
 - (β) if $f \in F$ and $\alpha \in w$ then $f^{[\alpha]} \in F$,
- (b) the order: $p \leq q$ iff $w^p \subseteq w^q$ and
 - (α) $f \in F^q \Rightarrow f \upharpoonright w^p \in cl(F^p)$,
 - (β) $(\forall f \in F^p)(\exists g \in F^q)[f \subseteq g]$.

(3) For $w \subseteq \lambda$ and $F \subseteq {}^w 2$ let $\text{BA}[w, F] = \text{BA}[(w, F)]$ be the Boolean algebra freely generated by $\{x_\alpha : \alpha \in w\}$ except that if u and v are finite subsets of w and $1_u \cup 0_v \subseteq f$ for no $f \in F$, then $\bigcap_{\alpha \in u} x_\alpha - \bigcup_{\beta \in v} x_\beta = 0$.

(4) If $G \subseteq Q_{\lambda, \mu}$ is generic over V then $\text{BA}[G]$ is $\bigcup_{p \in G} \text{BA}[p]$ (see 2.6(2), (3) below). Here $\text{BA}[p] := \text{BA}[w^p, F^p]$.

2.6. CLAIM. (0) For $p \in Q_{\lambda, \mu}$, $\text{BA}[p]$ is a Boolean algebra; also for $f \in F^p$ and ordinal α (or ∞) we have $f^{[\alpha]} \in F^p$.

(1) If $f \in F^p$ and $p \in Q_{\lambda, \mu}$ then f induces a homomorphism f^{hom} from $\text{BA}[p]$ to the two-member Boolean algebra $\{0, 1\}$. In fact, for a term τ in $\{x_\alpha : \alpha \in w^p\}$, $\text{BA}[p] \models \tau \neq 0$ iff for some $f \in F^p$, $f^{\text{hom}}(\tau) = 1$.

(2) If $p \leq q$ then $\text{BA}[p]$ is a Boolean subalgebra of $\text{BA}[q]$.

(3) Hence $\text{BA}[G]$ is well defined, $p \Vdash \text{“BA}[p] \text{ is a Boolean subalgebra of BA}[G]\text{”}$.

(4) For $p \in Q_{\lambda, \mu}$ and $\alpha \in w^p$, x_α is a non-zero element which is not in the subalgebra generated by $\{x_\beta : \beta < \alpha\}$ nor is there below it a non-zero member of $\langle x_\beta : \beta < \alpha \rangle_{\text{BA}[p]}$.

Proof. Part (0) should be clear, and also part (1). Now part (2) follows by 2.5(2)(b) and the definition of $\text{BA}[p]$; so (3) should become clear. Lastly, concerning part (4), x_α is a non-zero member of $\text{BA}[p]$ by clause (α) of

2.5(2)(a). For $\alpha \in w^p$, by 2.5(2)(a)(α) there is $f^1 \in F^p$ with $f^1(\alpha) = 1$, and by 2.5(2)(a)(β) there is $f^0 \in F^p$ with $f^0(\alpha) = 0$, $f^0 \upharpoonright (w \cap \alpha) = f^1 \upharpoonright (w \cap \alpha)$; together with part (1) this proves the second phrase of part (4). As for the third phrase, let τ be a non-zero element of the subalgebra generated by $\{x_\beta : \beta < \alpha\}$, so for some $f \in F^p$, $f^{\text{hom}}(\tau) = 1$. By 2.5(2)(a)(β), letting $f_1 = f^{[\alpha]}$, we have $f_1(\alpha) = 0$ and $f_1 \in F^p$ and $f_1 \upharpoonright (w \cap \alpha) \subseteq f$. Hence $f_1^{\text{hom}}(\tau) = f^{\text{hom}}(\tau) = 1$ and $f_1(\alpha) = 0$, hence $f_1^{\text{hom}}(x_\alpha) = 0$. This proves $\text{BA}[p] \models \text{“}\tau \not\leq x_\alpha\text{”}$. ■_{2.6}

2.7. CLAIM. Assume $\mu = \mu^{<\mu} < \lambda$.

- (1) $Q_{\lambda,\mu}$ is a μ -complete forcing notion of cardinality $\leq \lambda^{<\mu}$.
- (2) $Q_{\lambda,\mu}$ satisfies the μ^+ -c.c.

Proof. (1) The number of elements of $Q_{\lambda,\mu}$ is at most

$$\begin{aligned} & |\{(w, F) : w \subseteq \lambda, |w| < \mu \text{ and } F \text{ is a family of } < \mu \text{ functions} \\ & \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \text{from } w \text{ to } \{0, 1\}\}| \\ & \leq \sum_{w \subseteq \lambda, |w| < \mu} |\{F : F \subseteq {}^w 2 \text{ and } |F| < \mu\}| \\ & \leq \sum_{w \subseteq \lambda, |w| < \mu} (2^{|w|})^{<\mu} \leq |\{w : w \subseteq \lambda \text{ and } |w| < \mu\}| \times \mu \\ & = \lambda^{<\mu} + \mu = \lambda^{<\mu}. \end{aligned}$$

As for the μ -completeness, let $\langle p_\zeta : \zeta < \delta \rangle$ be an increasing sequence of members of $Q_{\lambda,\mu}$ with $\delta < \mu$. Let $p_\zeta = (w_\zeta, F_\zeta)$, let $F'_\zeta = \text{cl}(F_\zeta)$, let $w = \bigcup_{\zeta < \delta} w_\zeta$ and let $F' = \{f \in {}^w 2 : \text{for every } \zeta < \delta \text{ we have } f \upharpoonright w_\zeta \in F'_\zeta\}$. Clearly for every $\zeta < \delta$ and $f \in F'_\zeta$ there is $g = g_f \in F'$ extending f . Lastly, let $F = \{g_f : f \in \bigcup_{\zeta < \delta} F'_\zeta\}$. Then $p = (w, F) \in Q_{\lambda,\mu}$ is an upper bound of $\langle p_\zeta : \zeta < \delta \rangle$, as required.

(2) By the Δ -system argument it suffices to prove that p^0, p^1 are compatible when:

- (a) $\text{otp}(w^{p^0}) = \text{otp}(w^{p^1})$ and (letting $H = H_{w^{p^1}, w^{p^0}}^{\text{OP}}$ be the unique order preserving function from w^{p^0} onto w^{p^1}),
- (b) H maps p^0 onto p^1 , i.e. $f \in F^{p^0} \Leftrightarrow (f \circ H^{-1}) \in F^{p^1}$,
- (c) $\alpha \in w^{p^0} \Rightarrow \alpha \leq H(\alpha)$,
- (d) for $\alpha \in w^{p^0}$ we have $\alpha \in w^{p^1}$ iff $\alpha = H(\alpha)$.

We now define $q \in Q$ by setting $w^q = w^{p^0} \cup w^{p^1}$ and

$$F^q = \{(f \cup (f \circ H))^{[\beta]} : f \in F^{p^1}, \beta \in w^q \cup \{\infty\}\}. \quad \blacksquare_{2.7}$$

2.8. CLAIM. Assume $\mu = \mu^{<\mu} < \lambda$. Suppose $Q = Q_{\lambda, \mu}$ and:

- (a) $p^l \in Q$ for $l < m < \aleph_0$,
- (b) $\text{otp}(w^{p^l}) = \text{otp}(w^{p^0})$, and $H_{l,k} = H_{w^{p^l}, w^{p^k}}^{\text{OP}}$ (see the proof of 2.7(2)),
- (c) $H_{l,k}$ maps p^k onto p^l ,
- (d) for $\alpha \in w^{p^0}$ the sequence $\langle H_{l,0}(\alpha) : l = 1, \dots, m-1 \rangle$ is either strictly increasing or constant; and $\{\alpha, \beta\} \subseteq w^{p^0}$ & $l, k < m$ & $H_{l,0}(\alpha) = H_{k,0}(\beta)$ implies $\alpha = \beta$; lastly, letting $w^* = w^{p^0} \cap w^{p^1}$ we have $[l \neq k \Rightarrow w^{p^l} \cap w^{p^k} = w^*]$ and $H_{l,k} \upharpoonright w^*$ is the identity,
- (e) $\tau(x_1, \dots, x_n)$ is a Boolean term, $\alpha_i^0 \in w^{p^0}$ for $i \in \{1, \dots, n\}$, $\alpha_1^0 < \dots < \alpha_n^0$ and $\alpha_i^l = H_{l,0}(\alpha_i^0)$,
- (f) in $\text{BA}[p^0]$, $\tau(x_{\alpha_1^0}, \dots, x_{\alpha_n^0})$ is not zero and even not in the subalgebra generated by $\{x_\alpha : \alpha \in w^*\}$,
- (g) $m-1 > n+1$.

Then there is $q \in Q$ such that:

- (α) $p^l \leq q$ for $l < m$, and $w^q = \bigcup_{l < m} w^{p^l}$,
- (β) $q \Vdash$ "in $\text{BA}[\underline{G}]$, there is a non-zero Boolean combination τ^* of $\{\tau(x_{\alpha_1^l}, \dots, x_{\alpha_n^l}) : 1 \leq l < m\}$ which is $\leq \tau(x_{\alpha_1^0}, \dots, x_{\alpha_n^0})$ ".

Proof. By assumption (f) (and 2.6(0), (4)) there are $f_0^*, f_1^* \in \text{cl}(F^{p^0})$ such that:

- (A) $f_0^* \upharpoonright w^* = f_1^* \upharpoonright w^*$,
- (B) in the two-member Boolean algebra $\{0, 1\}$ we have

$$\tau(f_0^*(\alpha_1^0), \dots, f_0^*(\alpha_n^0)) = 0, \quad \tau(f_1^*(\alpha_1^0), \dots, f_1^*(\alpha_n^0)) = 1.$$

Now there is $\gamma \in w^{p^0} \cup \{\infty\}$ such that $(f_0^*)^{[\gamma]} = f_0^*$ & $(f_1^*)^{[\gamma]} = f_1^*$ (e.g. $\gamma = \infty$). Choose such a (γ, f_0^*, f_1^*) with γ minimal. Let $w^q = \bigcup_{l=0}^{m-1} w^{p^l}$. We define a function $g \in {}^{(w^q)}2$ as follows:

- $g \upharpoonright w^{p^0} = f_1^*$,
- for odd $l \in [1, m)$, $g \upharpoonright w^{p^l} = f_1^* \circ H_{0,l}$, and
- for even $l \in [1, m)$ (but not $l = 0$), $g \upharpoonright w^{p^l} = f_0^* \circ H_{0,l}$.

Now g is well defined by clause (A) above. Let us define q :

$$F^q = \left\{ \left(\bigcup_{l=0}^{m-1} f \circ H_{0,l} \right)^{[\alpha]} : \alpha \in w^q \cup \{\infty\} \text{ and } f \in F^{p_0} \right\} \\ \cup \{g^{[\alpha]} : \alpha \in w^q \cup \{\infty\}\}, \\ q = (w^q, F^q).$$

We first check that $q \in Q$. Clearly $w^q \in [\lambda]^{<\mu}$. Also $F^q \subseteq (w^q)2$ and $|F^q| < \mu$ so we have to check the conditions (α) and (β) of Definition 2.5(2)(a):

(α) If $\alpha \in w^q$ then $\alpha \in w^{p^l}$ for some $l < m$, so as $p^l \in Q$ there is $f_l \in F^{p^l}$ such that $f_l(\alpha) = 1$. Now $f_0 = f_l \circ H_{0,l}$ for some $f_0 \in F^{p^0}$ so

$$f := \bigcup_{k < m} (f_0 \circ H_{0,k}) = \left(\bigcup_{k < m} (f_0 \circ H_{0,k}) \right)^{[\infty]}$$

belongs to F^q and $f(\alpha) = (f_0 \circ H_{0,l})(\alpha) = f_l(\alpha) = 1$.

(β) As for $\alpha, \beta \in w \cup \{\infty\}$ and $f \in (w^q)2$ we have $(f^{[\alpha]})^{[\beta]} = f^{[\min\{\alpha, \beta\}]}$ and as $(\bigcup_{l < m} f_l)^{[\alpha]} = \bigcup_{l < m} (f_l)^{[\alpha]}$, this condition holds by the way we have defined F^q .

We now check that $p^l \leq q$ for $l < m$. By the choice of q clearly $w^{p^l} \subseteq w^q$. Also if $f \in F^{p^l}$ then $f \circ H_{l,0} \in F^{p^0}$ and

$$\bigcup_{k < m} ((f \circ H_{l,0}) \circ H_{0,k})^{[\infty]}$$

belongs to F^q and extends f . Lastly, if $f \in F^q$ we prove that $f \upharpoonright w^{p^l} \in \text{cl}(F^{p^l})$ (in fact, $\in F^{p^l}$). Let $w^l := w^{p^l}$. We have two cases: in the first case $f = (\bigcup_{l < m} (f_0 \circ H_{0,l}))^{[\alpha]}$ for some $f_0 \in F^{p^0}$; let $\beta = \min[w^l \cup \{\infty\} \setminus \alpha]$, so $f^{[\alpha]} \upharpoonright w^l = (f_0 \circ H_{0,l})^{[\beta]}$; clearly $f_0 \circ H_{0,l} \in F^{p^l}$, hence $(f_0 \circ H_{0,l})^{[\beta]} \in F^{p^l}$ is as required. The second case is $f = g^{[\alpha]}$. Let $\beta = \min[w^{p^l} \cup \{\infty\} \setminus \alpha]$; now $f \upharpoonright w^{p^l}$ is $f_0^* \circ H_{0,l}$ or $f_1^* \circ H_{0,l}$ so $f \upharpoonright w^{p^l}$ is $(f_0^* \circ H_{0,l})^{[\beta]}$ or $(f_1^* \circ H_{0,l})^{[\beta]}$ and hence belongs to F^{p^l} .

Finally, we check that there is a non-zero Boolean combination of $\{\tau(x_{\alpha_1^l}, \dots, x_{\alpha_n^l}) : l = 1, \dots, m-1\}$ which is $\leq \tau(x_{\alpha_1^q}, \dots, x_{\alpha_n^q})$ in $\text{BA}[q]$. The required Boolean combination will be

$$\tau^* = \bigcap_{l=0}^{[(m-2)/2]} \tau(x_{\alpha_1^{2l+1}}, \dots, x_{\alpha_n^{2l+1}}) - \bigcup_{l=1}^{[(m-1)/2]} \tau(x_{\alpha_1^{2l}}, \dots, x_{\alpha_n^{2l}}).$$

So we have to prove the following two assertions.

ASSERTION 1. $\text{BA}[q] \models \text{“}\tau^* \neq 0\text{”}$.

Now $g = g^{[\infty]} \in F^q$ satisfies, for each $l \in [0, [(m-2)/2]]$,

$$g^{\text{hom}}(\tau(x_{\alpha_1^{2l+1}}, \dots, x_{\alpha_n^{2l+1}})) = (f_1^* \circ H_{0,2l+1})^{\text{hom}}(\tau(x_{\alpha_1^{2l+1}}, \dots, x_{\alpha_n^{2l+1}})) = 1;$$

also for each $l \in [1, [(m-1)/2]]$,

$$g^{\text{hom}}(\tau(x_{\alpha_1^{2l}}, \dots, x_{\alpha_n^{2l}})) = (f_0^* \circ H_{0,2l})^{\text{hom}}(\tau(x_{\alpha_1^{2l}}, \dots, x_{\alpha_n^{2l}})) = 0.$$

Putting the two together, we get the assertion.

ASSERTION 2. $\text{BA}[q] \models \text{``}\tau(x_{\alpha_1^0}, \dots, x_{\alpha_n^0}) \geq \tau^*\text{''}$.

So we have to prove just that

$$f \in F^q \Rightarrow f^{\text{hom}}(\tau^* - \tau(x_{\alpha_1^0}, \dots, x_{\alpha_n^0})) = 0.$$

Case 1: For some $\alpha \in w^q \cup \{\infty\}$ and $f_0 \in F^{p^0}$ we have

$$f = \left(\bigcup_{l < m} f_0 \circ H_{0,l} \right)^{[\alpha]}.$$

Let $\beta_l = \min(w^{p^l} \cup \{\infty\} \setminus \alpha)$, and let $\gamma_l \in w^{p^0}$ be such that $\gamma_l = H_{0,l}(\beta_l)$ or $\gamma_l = \beta_l = \infty$. Now by the assumption on $\langle w^{p^l} : l < m \rangle$, $\langle \gamma_l : l < m \rangle$ is non-increasing. For $l < m$, let $j_l = \min\{j : [j = n + 1] \text{ or } [j \in \{1, \dots, n\} \text{ and } \alpha_j^0 \geq \gamma_l]\}$. So $\langle j_l : l < m \rangle$ is non-increasing and there are $\leq n + 1$ possible values for each j_l . But by assumption (g), $m - 1 > n + 1$, so for some $k, 0 < k < k + 1 < m$ and $j_k = j_{k+1}$. So (as $\alpha_1^i < \dots < \alpha_n^i$)

$$(\forall j = 1, \dots, n)[f(x_{\alpha_j^k}) = f(x_{\alpha_j^{k+1}})],$$

hence

$$(\forall j = 1, \dots, n)[f^{\text{hom}}(x_{\alpha_j^k}) = f^{\text{hom}}(x_{\alpha_j^{k+1}})],$$

hence

$$f^{\text{hom}}(\tau(x_{\alpha_1^k}, \dots, x_{\alpha_n^k})) = f^{\text{hom}}(\tau(x_{\alpha_1^{k+1}}, \dots, x_{\alpha_n^{k+1}})),$$

hence (see definition of τ^*) $f^{\text{hom}}(\tau^*) = 0$, hence

$$f^{\text{hom}}(\tau^* - \tau(x_{\alpha_1^0}, \dots, x_{\alpha_n^0})) = 0,$$

as required.

Case 2: For some $\alpha \in w^q \cup \{\infty\}$, $f = g^{[\alpha]}$.

Let again $\beta_l = \min(w^{p^l} \cup \{\infty\} \setminus \alpha)$, $\gamma_l = H_{0,l}(\beta_l)$ (or $\gamma_l = \beta_l = \infty$), and

$$\gamma'_l = \begin{cases} \gamma_l & \text{if } \gamma_l < \gamma, \\ \infty & \text{if } \gamma_l \geq \gamma, \end{cases}$$

and let $j_l = \min\{j : [j = n + 1] \text{ or } [j \in \{1, \dots, n\} \text{ and } \alpha_j^0 \geq \gamma'_l]\}$. So $\langle \gamma_l : l < m \rangle$ and $\langle \gamma'_l : l < m \rangle$ are non-increasing and so is $\langle j_l : l < m \rangle$. Here γ is the ordinal we chose before defining q_1 , just after (B) in the proof.

If for some $k, 0 < k < k + 1 < m$ and $j_k = j_{k+1} \leq n$ (hence $\gamma'_{k+1} \leq \gamma'_k < \gamma$), then

$$f^{\text{hom}}(\tau^* - \tau(x_{\alpha_1^0}, \dots, x_{\alpha_n^0})) = 0$$

as $f^{\text{hom}}(\tau^*) = 0$, which holds because

$$f^{\text{hom}}(\tau(x_{\alpha_1^k}, \dots, x_{\alpha_n^k})) = f^{\text{hom}}(\tau(x_{\alpha_1^{k+1}}, \dots, x_{\alpha_n^{k+1}}))$$

(the last equality holds by the choice of γ ; i.e. if inequality holds then the triple $(\gamma_k, (f_0^*)^{[\gamma_k]}, (f_1^*)^{[\gamma_k]})$ contradicts the choice of γ as minimal). But

j_l ($l = 1, \dots, m-1$) is non-increasing, hence we can show inductively on $l = 1, \dots, n+1$ that $j_{m-l} \geq l$. So necessarily $j_1 = n+1$ but as j_l is non-increasing, clearly $j_0 = n+1$ and hence

$$\begin{aligned} f^{\text{hom}}(\tau(x_{\alpha_1^0}, \dots, x_{\alpha_n^0})) &= g^{[\alpha]}(\tau(x_{\alpha_1^0}, \dots, x_{\alpha_n^0})) = g(\tau(x_{\alpha_1^0}, \dots, x_{\alpha_n^0})) \\ &= f_1^*(\tau(x_{\alpha_1^0}, \dots, x_{\alpha_n^0})) = 1, \end{aligned}$$

hence

$$f^{\text{hom}}(\tau^* - \tau(x_{\alpha_1^0}, \dots, x_{\alpha_n^0})) = 0,$$

as required. $\blacksquare_{2.8}$

2.9. THEOREM. *Suppose $\mu = \mu^{<\mu} < \lambda$, $Q = Q_{\lambda, \mu}$ and $V \models G.C.H.$ (for simplicity). Then:*

- (1) Q is μ -complete, μ^+ -c.c. (hence forcing with Q preserves cardinals and cofinalities).
- (2) $\Vdash_Q "2^\mu = (\lambda^\mu)^V"$, $|Q| = \lambda^{<\mu}$, so cardinal arithmetic in V^Q is easily determined.
- (3) Let $G \subseteq Q$ be generic over V . Then $\text{BA}[G]$ (see Definition 2.5(4)) is a Boolean algebra of cardinality λ such that:
 - (a) if $\theta \leq \lambda$ is regular then for some subalgebra B of $\text{BA}[G]$, $\pi(B) = \theta$,
 - (b) if $\theta \leq \lambda$ and $\theta > \text{cf}(\theta) > \mu$ then for no $B \subseteq \text{BA}[G]$ is $\pi(B) = \theta$,
 - (c) $\text{BA}[G]$ has μ non-zero pairwise disjoint elements but no μ^+ such elements (so its cellularity is μ),
 - (d) if $a \in B^+$ then $\text{BA}[G] \upharpoonright a$ satisfies (a), (b), (c) above (and also (e)),
 - (e) if $\theta \leq \lambda$ and $\text{cf}(\theta) \leq \mu$ then for some $B' \subseteq \text{BA}[G]$ we have $\pi(B') = \theta$,
 - (f) in $\text{BA}[G]$ for every $\alpha < \lambda$, $\{x_\beta : \alpha \leq \beta < \alpha + \mu\} \subseteq B^+$ is dense in $\langle \{x_\beta : \beta < \alpha\} \rangle_{\text{BA}[G]}$.

2.9A. Remark. (1) This shows the consistency of a negative answer to problems 13 and 15 of Monk [M].

(2) We could of course make 2^μ bigger by adding the right number of Cohen subsets of μ .

Proof of Theorem 2.9. By Claim 2.7 clearly parts (1), (2) hold. We are left with part (3). By 2.6(3), $\text{BA}[G]$ is a Boolean algebra, by 2.6(4) it has cardinality λ . As for clause (a), it is exemplified by $\langle x_\alpha : \alpha < \theta \rangle_{\text{BA}[G]}$ (by 2.6(4)). The first statement of (c) is easy by the genericity of G (i.e. for $p \in Q$ and $\alpha \in \lambda \setminus w^p$ we can find q such that $p \leq q \in Q$, $w^q = w^p \cup \{\alpha\}$ and in $\text{BA}[q]$, x_α is disjoint from all $y \in J$, for any ideal J of $\text{BA}[p]$). The second statement of (c) follows from the Δ -system argument and the proof of 2.7(2).

Concerning the generalization of clause (a) in (d), let $a \in (\text{BA}[G])^+$, so we can find finite disjoint $u, v \subseteq \lambda$ such that $0 < \bigcap_{\alpha \in u} x_\alpha - \bigcap_{\alpha \in v} x_\alpha \leq a$, choose $\beta = \sup(u \cup v)$, and let

$$U = \left\{ x_\gamma : \beta < \gamma < \beta + \theta, \text{ and } \text{BA}[G] \models "x_\gamma \leq \left(\bigcap_{\alpha \in u} x_\alpha - \bigcap_{\alpha \in v} x_\alpha \right)" \right\}.$$

This set is forced to be of cardinality θ and the subalgebra of $(\text{BA}[G]) \upharpoonright a$ generated by $\{x_\gamma \cap a : \gamma \in [\beta, \beta + \theta)\}$ is as required.

The generalization of (b) in (d) follows from clause (b). For the generalization of clause (c) in (d), the cellularity being $\leq \mu$ follows from (c), and the existence of $\min\{|a|, \mu\}$ pairwise disjoint elements follows from the fact that for every $p \in Q_{\lambda, \mu}$, $\alpha < \lambda$ and $a^* \in \text{BA}[p]$ such that $a^* \in \langle x_\beta : \beta \in w^p \cap \alpha \rangle_{\text{BA}[p]}$ and $\beta \in [\alpha, \lambda) \setminus w^p$ there is q such that $p \leq q \in Q_{\lambda, \mu}$ and $\text{BA}[q] \models "(\exists \gamma \in w^p \setminus \alpha)[x_\beta \cap x_\gamma = 0] \ \& \ x_\beta \leq a^*" .$

As for clause (e) (and the generalization in clause (d)), let $a \in (\text{BA}[G])^+$, and let $u \subseteq \lambda$ be finite such that $a \in \langle x_\alpha : \alpha \in u \rangle_{\text{BA}[G]}$. Then we can find $\langle a_i : i < \mu \rangle$, pairwise disjoint non-zero members of $\langle x_\alpha : \alpha \in (\sup(u), \sup(u) + \mu) \rangle_{\text{BA}[G]}$ which are below a . Let $\theta = \sum_{\zeta < \text{cf}(\theta)} \theta_\zeta$ with each θ_ζ regular, let $B_\zeta \subseteq \text{BA}[G] \upharpoonright a_\zeta$ be a subalgebra with $\pi(B_\zeta) = \theta_\zeta$, and lastly let B be the subalgebra of $\text{BA}[G] \upharpoonright a$ generated by $\{a_i : i < \text{cf}(\theta)\} \cup \bigcup_{\zeta < \text{cf}(\theta)} B_\zeta$; check that $\pi(B) = \theta$.

Clause (f) follows by a density argument. The real point (and the only one left) is to prove clause (b) of part (3). So suppose toward a contradiction that $\mu < \text{cf}(\chi) < \chi \leq \lambda$ and $p \in Q$ but $p \Vdash_Q "B \subseteq \text{BA}[\underline{G}]$ is a subalgebra, $\pi(B) = \chi"$. Then by Claim 2.2+2.3(1), $p \Vdash$ "for arbitrarily large regular $\theta < \chi$, there is $\bar{y} = \langle \bar{y}_\alpha : \alpha < \theta \rangle$ (a sequence of non-zero elements of B) and a θ -complete proper filter D on θ (containing the cobounded subsets of θ) such that $(\otimes_{\bar{y}, D}^B)$ holds (see 2.3(1))".

Let $\kappa = \text{cf}(\chi)$. Then we can find regular $\theta_\zeta \in (\text{cf}(\chi), \chi)$ (so $\theta_\zeta > \mu$) increasing with ζ such that $\chi = \sum_{\zeta < \kappa} \theta_\zeta$, and for $i < \kappa$, $(\sum_{j < i} \theta_j)^\kappa < \theta_i$ (remember $V \models \text{G.C.H.}$) and for each $\zeta < \kappa$, a condition p_ζ with $p \leq p_\zeta \in Q$, and $\bar{y}^\zeta = \langle \bar{y}_\alpha^\zeta : \alpha < \theta_\zeta \rangle$, and (a Q -name of a) proper θ_ζ -complete filter D_ζ on θ_ζ containing the cobounded subsets of θ_ζ such that $p_\zeta \Vdash "(\otimes_{\bar{y}^\zeta, D_\zeta}^B)"$ (and without loss of generality $\Vdash "y_\alpha^\zeta \in (\text{BA}[\underline{G}])^+"$). For each $\zeta < \kappa$ and $\alpha < \theta_\zeta$ there is a maximal antichain $\bar{p}^{\zeta, \alpha} = \langle p_{\zeta, \alpha, \varepsilon} : \varepsilon < \mu \rangle$ of members of Q above p_ζ and terms $\tau_{\zeta, \alpha, \varepsilon} = \tau'_{\zeta, \alpha, \varepsilon}(x_{\beta(\zeta, \alpha, \varepsilon, 0)}, \dots, x_{\beta(\zeta, \alpha, \varepsilon, n_\alpha(\zeta, \varepsilon))})$ (i.e. Boolean terms in $\{x_\alpha : \alpha < \lambda\}$) such that $p_\zeta \leq p_{\zeta, \alpha, \varepsilon}$ and $p_{\zeta, \alpha, \varepsilon} \Vdash_Q "y_\alpha^\zeta = \tau_{\zeta, \alpha, \varepsilon}"$. Without loss of generality $\{\beta(\zeta, \alpha, \varepsilon, l) : l \leq n_\alpha(\zeta, \varepsilon)\} \subseteq w[p_{\zeta, \alpha, \varepsilon}]$.

Clearly for each $\zeta < \kappa$, $p_\zeta \Vdash$ " θ_ζ is the disjoint union of $\{\alpha < \theta_\zeta : p_{\zeta, \alpha, \varepsilon} \in \underline{G}\}$ for $\varepsilon < \mu$ " so for some Q -name $\underline{\varepsilon}_\zeta < \mu$, we have $p_\zeta \Vdash_Q "\{\alpha < \theta_\zeta : p_{\zeta, \alpha, \underline{\varepsilon}_\zeta} \in \underline{G}\} \neq \emptyset \text{ mod } D_\zeta"$. So there are $\varepsilon_\zeta < \mu$ and q_ζ satisfying $p_\zeta \leq q_\zeta \in Q$ such

that $q_\zeta \Vdash \text{“}\varepsilon_\zeta \text{ is as above”}$ and let $p_{\zeta,\alpha} := p_{\zeta,\alpha,\varepsilon_\zeta}$. So we have a Q -name \underline{A}_ζ such that $q_\zeta \Vdash_Q \text{“}\underline{A}_\zeta \subseteq \theta_\zeta, \underline{A}_\zeta \neq \emptyset \text{ mod } \underline{D}_\zeta \text{ and } \alpha \in \underline{A}_\zeta \Leftrightarrow p_{\zeta,\alpha} \in \underline{G}_Q\text{”}$.

By possibly replacing $\theta_\zeta, \underline{A}_\zeta$ by $A_\zeta^* \in [\theta_\zeta]^{\theta_\zeta}$, $\underline{A}'_\zeta = \underline{A}_\zeta \cap A_\zeta^*$ respectively, and increasing q_ζ , we can assume that $\text{otp}(w^{p_{\zeta,\alpha}}) = i_\zeta^* (< \mu)$, and letting $w^{p_{\zeta,\alpha}} = \{\beta_{\alpha,\zeta,i} : i < i_\zeta^*\}$ (increasing with i) and (by Claim 2.4) for some $w_\zeta^* \subseteq i_\zeta^*$, $\langle \beta_{\zeta,i}^* : i < i_\zeta^* \rangle$ and τ_ζ^* we have $\tau'_{\zeta,\alpha,\varepsilon_\zeta} = \tau_\zeta^*$, and for some strictly increasing $\langle j(\zeta, l) : l \leq n_\zeta \rangle$ we have

- $q_\zeta \Vdash_Q$ “(a) $\alpha \in \underline{A}_\zeta$ & $i \in w_\zeta^* \Rightarrow \beta_{\alpha,\zeta,i} = \beta_{\zeta,i}^*$,
- (b) $\beta(\zeta, \alpha, \varepsilon_\zeta, l) = \beta_{\zeta,\alpha,j(\zeta,l)}$ and $n_\alpha(\zeta, \varepsilon_\zeta) = n_\zeta$,
- (c) for every $\beta'_{\zeta,i} < \beta_{\zeta,i}^*$ (for $i \in i_\zeta^* \setminus w_\zeta^*$) we have
- $\{\alpha < \theta_\zeta : \alpha \in \underline{A}_\zeta, [i \in w_\zeta^* \Rightarrow \beta_{\alpha,\zeta,i} = \beta_{\zeta,i}^*]$ and for $i \in i_\zeta^* \setminus w_\zeta^*$
we have $\beta'_{\zeta,i} < \beta_{\alpha,\zeta,i} < \beta_{\zeta,i}^*\} \neq \emptyset \text{ mod } \underline{D}_\zeta\text{”}$.

Also

$$i \in i_\zeta^* \setminus w_\zeta^* \Rightarrow \left(2 + \sum_{j < i} \theta_j\right)^\kappa < \text{cf}(\beta_{\zeta,i}^*) \leq \theta_\zeta$$

(remember \underline{D}_ζ is a θ_ζ -complete filter on θ_ζ).

As we can replace $\langle \theta_\zeta : \zeta < \kappa \rangle$ by any subsequence of length κ , and $\kappa = \text{cf}(\kappa) > \mu$, without loss of generality $i_\zeta^* = i^*$, $w_\zeta^* = w^*$ and $\text{otp}(w^{q_\zeta}) = j^*$. Let $w^{q_\zeta} = \{\beta_{\zeta,i}^* : i^* \leq i < j^*\}$. Now we apply 2.4 to κ , D_κ (filter of closed unbounded sets) and $\langle \langle \beta_{\zeta,i}^* : i < j^* \rangle : \zeta < \kappa \rangle$ and get $\langle \beta_i^\otimes : i < j^* \rangle$ and $w^\otimes \subseteq j^*$. Without loss of generality the q_ζ are pairwise isomorphic. Note

$$i \in i^* \setminus w^* \text{ \& } \zeta < \xi < \kappa \Rightarrow \beta_{\zeta,i}^* \neq \beta_{\xi,i}^*$$

(as $\text{cf}(\beta_{\zeta,i}^*) \leq \theta_\zeta$ and $\text{cf}(\beta_{\xi,i}^*) > \theta_\zeta$). Hence $w^\otimes \cap i^* \subseteq w^*$. For every $\zeta < \kappa$ and $i \in i^* \setminus w^*$, let $\beta_{\zeta,i}^- < \beta_{\zeta,i}^*$ be such that the interval $[\beta_{\zeta,i}^-, \beta_{\zeta,i}^*)$ is disjoint from $\{\beta_{\xi,j}^* : \xi < \kappa, j < i^*\} \cup \{\beta_i^\otimes : i < i^*\}$, and as we can omit an initial segment of $\langle \theta_i : i < \kappa \rangle$, without loss of generality $[\beta_{\zeta,i}^-, \beta_i^\otimes)$ is disjoint from $\{\beta_j^\otimes : j < i^*\}$. For each $\zeta < \kappa$, choose $\alpha_\zeta \in A_\zeta^*$ such that

$$i \in i^* \setminus w^* \Rightarrow \beta_{\zeta,\alpha_\zeta,i} \in [\beta_{\zeta,i}^-, \beta_{\zeta,i}^*).$$

Let $\underline{Y} =$ the Boolean subalgebra generated by $\{\tau_{\zeta,\alpha_\zeta} : q_\zeta \in \underline{G} \text{ and } p_{\zeta,\alpha_\zeta} \in \underline{G}\}$. This set has cardinality $\leq \kappa$, and we shall prove

- (*) $q_0 \Vdash_Q \text{“}\underline{Y} \setminus \{0\} \text{ is dense in } \{\tau_{0,\beta} : \beta \in \underline{A}_0\}\text{”}$.

This contradicts the choice $q_0 \Vdash \text{“}(\otimes_{\underline{y}_0}^B) \text{ \& } \underline{A} \neq \emptyset \text{ mod } \underline{D}_0\text{”}$.

To prove (*) assume $q_0 \leq r_0 \in Q$. We can choose $\zeta^* < \kappa$ and r_ζ^+ for $\zeta \in [\zeta^*, \kappa)$ such that $r_0 \leq r_0^+$, $q_\zeta \leq r_\zeta^+$, $p_{\zeta,\alpha_\zeta} \leq r_\zeta^+$ and $\langle r_\zeta : \zeta = 0 \text{ or } \zeta \in [\zeta^*, \kappa) \rangle$ is as in 2.8; apply 2.8 and get a contradiction. $\blacksquare_{2.9}$

A theorem complementary to 2.9 is:

2.10. THEOREM. *Suppose $\pi(B) > \lambda$ and either*

(A) $\text{cf}(\lambda) = \aleph_0$ *or*

(B) $(\forall \mu < \lambda)[\mu^{<\text{cf}(\lambda)} < \lambda]$ *or*

(C) $(\forall \mu < \lambda)[2^\mu < \pi(B)]$ *(or just)* $(\forall \mu < \lambda)[2^\mu < |B|]$ & $\lambda \leq \pi(B)$.

Then B has a subalgebra B' such that $\lambda = \pi(B') = |B'|$.

Remark. The conclusion of 2.10 implies that $\lambda \in \pi_{S_S}(B) := \{\pi(A) : A \subseteq B\}$.

Proof of Theorem 2.10. Case (C) is easier so we ignore it. By 2.1 without loss of generality $\pi(B) = \lambda^+ = |B|$. By induction on $\alpha < \lambda$, we try to choose a_α such that:

(a) $a_\alpha \in B^+$,

(b) for $\beta < \alpha$ we have $B \models "a_\alpha \cap a_\beta = 0"$,

(c) $\pi(B \upharpoonright a_\alpha) < \lambda^+$.

Let a_α be defined iff $\alpha < \alpha^*$.

Case 1: $\alpha^* \geq \lambda$. Let B' be the subalgebra generated by $\{a_\alpha : \alpha < \lambda\}$. Clearly $|B'| = \pi(B') = \lambda$.

Case 2: *Not Case 1 but* $\sum_{\alpha < \alpha^*} \pi(B \upharpoonright a_\alpha) \geq \lambda$. So we can find distinct $\alpha_\zeta < \alpha^*$ for $\zeta < \text{cf}(\lambda)$ such that $\sum_\zeta \pi(B \upharpoonright a_{\alpha_\zeta}) \geq \lambda$. We can find regular $\theta_\zeta \leq \pi(B \upharpoonright a_{\alpha_\zeta})$ such that $\sup_{\zeta < \text{cf}(\lambda)} \theta_\zeta = \lambda$ and then find $B_\zeta \subseteq B \upharpoonright a_{\alpha_\zeta}$ such that $|B_\zeta| = \theta_\zeta$ and $\pi(B_\zeta) = \theta_\zeta$ (by 2.1). Let B' be the subalgebra of B generated by $\bigcup_{\zeta < \text{cf}(\lambda)} B_\zeta \cup \{a_{\alpha_\zeta} : \zeta < \text{cf}(\lambda)\}$. Clearly $|B'| = \pi(B') = \lambda$.

Case 3: *Cases 1 and 2 fail.* Let $I = \{a \in B : (\forall \alpha < \alpha^*)[a \cap a_\alpha = 0]\}$, so I is an ideal of B and $a \in I \Rightarrow \pi(B \upharpoonright a) \geq \lambda$. Also $I \neq \{0\}$ (since if $I = \{0\}$ then $\pi(B) \leq \sum_{\alpha < \alpha^*} \pi(B \upharpoonright a_\alpha) < \lambda$). So easily without loss of generality:

(*) if $a \in I \cap B^+$ then $\pi(B \upharpoonright a) > \lambda$,

(**) if $a \in I \cap B^+$ then $B \upharpoonright a$ is an atomless Boolean algebra.

Let $B^* = I \cup \{-b : b \in I\}$, a subalgebra of B . Now without loss of generality B^* satisfies $(\text{cf}(\lambda))$ -c.c. (otherwise act as in Case 2), so we have finished if Case (A) of the hypothesis holds.

So Case (B) of the hypothesis holds, hence we can use Lemma 4.9, p. 88 of [Sh:92] and find a free subalgebra B' of B^* of cardinality $(\lambda^{<\text{cf}(\lambda)})^+$, hence of cardinality λ ; this B' is as required. ■_{2.10}

3. On π and $\pi\chi$ of products of Boolean algebras

3.1. THEOREM. *Suppose*

(\otimes) $\aleph_0 < \kappa = \text{cf}(\chi) < \chi < \lambda = \text{cf}(\lambda) < \chi^\kappa$ and $(\forall \theta < \chi)[\theta^\kappa < \chi]$.

Then there are Boolean algebras B_i (for $i < \kappa$) such that (for $\pi(F, B)$, $\pi\chi(B)$ see below)

- (*) (a) $\pi\chi(B_i) < \chi = \sum_{j < \kappa} \pi\chi(B_j)$,
 (b) for any uniform ultrafilter D on κ , $\lambda = \pi\chi(\prod_{i < \kappa} B_i/D)$,
 (c) if D is a regular ultrafilter on κ then $\prod_{i < \kappa} (\pi\chi(B_i))/D = \chi^\kappa$.

3.2. DEFINITION. (1) For a Boolean algebra B and an ultrafilter F of B , let

$$\pi(F, B) = \min\{|X| : X \subseteq B^+ \text{ and } (\forall y \in F)(\exists x \in X)[x \leq y]\}.$$

We say X as above is *dense* in F (though possibly $x \notin F$).

(2) For a Boolean algebra B ,

$$\begin{aligned} \pi\chi(B) &= \sup\{\pi(F, B) : F \text{ an ultrafilter of } B\}, \\ \pi\chi^+(B) &= \bigcup\{\pi(F, B)^+ : F \text{ an ultrafilter of } B\}. \end{aligned}$$

3.3. Remark. (1) If $\kappa = \aleph_0$ the theorem holds almost always and probably always, but we omit this case to simplify the statement. (The theorem holds for $\kappa = \aleph_0$ e.g. if $\chi < \lambda = \text{cf}(\lambda) < (\text{first fix point } > \chi)$, more generally if

$$(\otimes') \quad \kappa = \text{cf}(\chi) < \chi < \lambda = \text{cf}(\lambda) < \text{pp}_{\mathcal{J}_{\text{bd}}^+}(\chi) \quad \text{and} \quad 2^\kappa < \chi$$

(see [Sh:g], VIII, §1). The point is that [Sh 355], 5.4, deals with uncountable cofinalities.)

3.4. Proof of Theorem 3.1. For a linear order \mathcal{I} , let $\text{BA}[\mathcal{I}]$ be the Boolean algebra of subsets of \mathcal{I} generated by the closed-open intervals $[a, b) = \{x \in \mathcal{I} : a \leq x < b\}$ where we allow $a \in \{-\infty\} \cup \mathcal{I}$, $b \in \mathcal{I} \cup \{\infty\}$ (and $a \leq b$). Now clearly

- (*)₁ if F is an ultrafilter on $\text{BA}[\mathcal{I}]$, then there is a Dedekind cut $(\mathcal{I}^d, \mathcal{I}^u)$ of \mathcal{I} (i.e. $\mathcal{I}^d \cap \mathcal{I}^u = \emptyset$, $\mathcal{I}^d \cup \mathcal{I}^u = \mathcal{I}$ and $(\forall x_0 \in \mathcal{I}^d)(\forall x_1 \in \mathcal{I}^u)[x_0 < x_1]$) such that for $x \in \text{BA}[\mathcal{I}]$, $x \in F$ iff for some $a_0 \in \mathcal{I}^d$, $a_1 \in \mathcal{I}^u$ we have $[a_0, a_1) \leq x$,
 (*)₂ if \mathcal{I}, F and $(\mathcal{I}^d, \mathcal{I}^u)$ are as above then

$$\pi(F, \text{BA}(\mathcal{I})) = \begin{cases} \max\{\text{cf}(\mathcal{I}^d), \text{cf}((\mathcal{I}^u)^*)\} & \text{if } \text{cf}(\mathcal{I}^d), \text{cf}((\mathcal{I}^u)^*) \geq \aleph_0, \\ \text{cf}(\mathcal{I}^d) & \text{if } \text{cf}((\mathcal{I}^u)^*) \leq 1, \\ \text{cf}((\mathcal{I}^u)^*) & \text{if } \text{cf}(\mathcal{I}^d) \leq 1, \\ 1 & \text{if } \text{cf}(\mathcal{I}^d) = \text{cf}((\mathcal{I}^u)^*) = 1. \end{cases}$$

Note also

$$\pi\chi(\text{BA}(\mathcal{I})) = \sup\{\text{cf}(\mathcal{I}^d), \text{cf}((\mathcal{I}^u)^*) : (\mathcal{I}^d, \mathcal{I}^u) \text{ a Dedekind cut of } \mathcal{I}\}.$$

Now by the assumption (\otimes) (and [Sh:g], II,5.4 + VIII, §1), we can find a (strictly) increasing sequence $\langle \lambda_i : i < \kappa \rangle$ of regular cardinals with $\kappa < \lambda_i < \chi$ and $\chi = \sum_{i < \kappa} \lambda_i$ such that $\prod_{i < \kappa} \lambda_i / J_\kappa^{\text{bd}}$ has true cofinality λ (where J_κ^{bd} is the ideal of bounded subsets of κ).

Let \mathbb{Q} be the rational order and \mathcal{I}_i be $\lambda_i \times \mathbb{Q}$ (i.e. the set of elements is $\{(\alpha, q) : \alpha < \lambda_i, q \in \mathbb{Q}\}$, and the order is lexicographical). Let $B_i = \text{BA}[\mathcal{I}_i]$. By $(*)_1$ and $(*)_2$ we know that $\pi\chi(B_i) = \lambda_i$. Moreover, if F is an ultrafilter of B_i , then $\pi(F, B) = \aleph_0$ except when F is the ultrafilter F_i generated by $\{x_\alpha^i = [(\alpha, 0), \infty) : \alpha < \lambda_i\}$. Let $x_{\alpha, q}^i := [(\alpha, q), \infty)$. Let D be a uniform ultrafilter on κ , so $\prod_{i < \kappa} \lambda_i / D$ has cofinality λ . Also if D is regular, then (see [CK]) we know that $\chi^\kappa = \prod_{i < \kappa} \lambda_i / D = \prod_{i < \kappa} (\pi\chi(B_i)) / D$. So parts (a) and (c) of $(*)$ of Theorem 3.1 are satisfied.

To prove part (b) of $(*)$, let D be a uniform ultrafilter on κ and let $B := \prod_{i < \kappa} B_i / D$. Let F be such that $(B, F) = \prod_{i < \kappa} (B_i, F_i) / D$. Clearly F is an ultrafilter of B ; it is generated by $X = \prod_{i < \kappa} X_i / D$, where $X_i = \{x_{\alpha, q}^i : \alpha < \lambda, q \in \mathbb{Q}\} \subseteq B_i$, which is linearly ordered in B , and this linear order has the same cofinality as $\prod_{i < \kappa} \lambda_i / D$, which has cofinality λ . So $\pi\chi(F, B) = \lambda$, hence $\pi\chi(B) \geq \lambda$.

Let F' be an ultrafilter of B with $F' \neq F$. Let $X_d := \{x \in X : x \in F'\}$ and $X_u := \{x \in X : x \notin F' \text{ (i.e. } 1_B - x \in F')\}$. Clearly (X_d, X_u) is a Dedekind cut of X (which is linearly ordered: as a subset of B , or as $\prod_{i < \kappa} X_i / D$, where $X_i \subseteq B_i$ inherit the order from B_i , so $x_{\alpha, a}^i < x_{\beta, b}^i \Leftrightarrow (\alpha, a) < (\beta, b)$). If $X_d = X$ then clearly $F' = F$, a contradiction, so $X_u \neq \emptyset$.

We now prove that $\pi\chi(F, B) \leq 2^\kappa$. If not, we shall choose by induction on $\zeta < (2^\kappa)^+$ a set Y_ζ , subsets Z_ζ^i , of λ_i for $i < \kappa$, increasing continuous in ζ , and y_ζ such that:

- $|Z_\zeta^i| \leq 2^\kappa$,
- $\xi < \zeta \Rightarrow Z_\xi^i \subseteq Z_\zeta^i$,
- $Y_\zeta = \prod_{i < \kappa} (Z_\zeta^i \times \mathbb{Q}) / D \setminus \{0\}$, so $|Y_\zeta| \leq 2^\kappa$,
- $y_\zeta \in F'$,
- $y_\zeta \in Y_{\zeta+1}$,
- $(\forall y \in Y_\zeta)[y > 0 \Rightarrow B \models \neg y \leq y_\zeta]$.

There is no problem in doing this for $i = 0$: let $Z_\zeta^0 = \{0\}$, and for i limit let $Z_\zeta^i = \bigcup_{\varepsilon < \zeta} Z_\varepsilon^i$. Now having defined $\langle Z_\zeta^i : i < \kappa \rangle$ (hence Y_ζ), choose appropriate y_ζ and let

$$y_\zeta = \langle y_\zeta^i : i < \kappa \rangle / D, \quad y_\zeta^i = \bigcup_{l < n_{i, \zeta}} [x_{\alpha_{i, \zeta, 2l}, q_{i, \zeta, 2l}}^i, x_{\alpha_{i, \zeta, 2l+1}, q_{i, \zeta, 2l+1}}^i),$$

where $\langle (\alpha_{i, \zeta, l}, q_{i, \zeta, l}) : l < 2n_{i, \zeta} \rangle$ is a strictly increasing sequence of members

of $Z_\zeta^i \cup \{-\infty, +\infty\}$ (we write $-\infty = (-\infty, 0)$, $+\infty = (\infty, 0)$). Let

$$Z_{\zeta+1}^i = Z_\zeta^i \cup \{\alpha_{i,\zeta,l} : l < 2n_{i,\zeta}\}.$$

For some unbounded $\mathcal{U} \subseteq (2^\kappa)^+$ we have

$$(a) \quad q_{i,\zeta,l} = q_{i,l} \quad \text{and} \quad n_{i,\zeta} = n_i;$$

applying 2.4, we get an easy contradiction. $\blacksquare_{3.4}$

3.4A. Remark. We can similarly analyze (when $B_i = \text{BA}[\mathcal{I}_i]$)

$$\left\{ \pi \left(F, \prod_{i < \kappa} B_i / D \right) : F \text{ an ultrafilter of } \prod_{i < \kappa} B_i / D \right\} \setminus (2^\kappa)^+$$

$$= \left\{ \lambda : \lambda'_i = \text{cf}(\lambda'_i) > 2^\kappa \text{ and in } \mathcal{I}_i \text{ there is a Dedekind cut} \right.$$

$$(X_d, X_u) \text{ such that } (\text{cf}(X_d), \text{cf}(X_u^*)) = (\lambda_i^d, \lambda_i^u) \text{ such that}$$

$$\left. \lambda = \min \left[\left\{ \text{cf} \left(\prod \lambda_i^u / D \right), \text{cf} \left(\prod \lambda_i^d / D \right) \right\} \setminus \{1\} \right] \right\}.$$

Note in comparison that by Koppelberg–Shelah [KpSh 415], Th. 1.1, we have

3.5. THEOREM. Assume D is an ultrafilter on κ , and for $i < \kappa$, A_i is a Boolean algebra with $\lambda_i = \pi(A_i)$. Assume the Strong Hypothesis [Sh 420], 6.2, i.e. $\text{pp}(\mu) = \mu^+$ for all singulars or just SCH. If $2^\kappa < \lambda_i$ (or just $\{i : 2^\kappa < \lambda_i\} \in D$) then

$$\pi \left(\prod_{i < \kappa} A_i / D \right) = \prod_{i < \kappa} \lambda_i / D.$$

3.6. CLAIM. Assume that for $i < \kappa$, A_i is an infinite Boolean algebra, D is a non-principal ultrafilter on κ and $A := \prod_{i < \kappa} A_i / D$. If $n_i < \omega$ for $i < \kappa$ and $\mu := |\prod_{i < \kappa} n_i / D|$ and μ is a regular cardinal then $\pi\chi(A) \geq \mu$, even $\pi\chi^+(A) > \mu$.

Proof. Let χ be a large enough regular cardinal (i.e. such that κ, D, A_i, A belong to $\underline{H}(\chi)$). Let $\mathfrak{C}_i = (\underline{H}(\chi), \in, <^*)$ and $\mathfrak{C} = \prod_i \mathfrak{C}_i / D$, so A is a member of \mathfrak{C} .

Clearly $\omega^* := \langle \omega : i < \kappa \rangle / D$ is considered by \mathfrak{C} a limit ordinal and, from the outside, has a cofinality, which we call λ . Without loss of generality, $i < \kappa \Rightarrow n_i > 2$.

The proof is divided into two cases.

Case 1: There are no $\mu_0 < \mu$ and $n_i^0 < n_i$ such that $\aleph_0 \leq \mu_0 = |\prod_{i < \kappa} n_i^0 / D|$. We can find n_i^* such that (we shall not use any further properties of the n_i^*):

$$(1) \quad \mu = |\prod_{i < \kappa} n_i^* / D|,$$

$$(2) \mu = |\prod_{i < \kappa} 2^{(n_i^*)^{(n_i^*)}} / D|.$$

For $i < \kappa$, let $\langle a_k^i : k < 2^{(n_i^*)^{(n_i^*)}} \rangle$ be pairwise disjoint non-zero members of A_i with union 1_{A_i} . Let P^i be the Boolean subalgebra of A_i generated by $\{a_k^i : k < 2^{(n_i^*)^{(n_i^*)}}\}$. Let $R^i := \{a_k^i : k < 2^{(n_i^*)^{(n_i^*)}}\}$. For $k < n_i^*$, let $Q_k^i \subseteq P^i$ be a set of n_i^* pairwise disjoint non-zero elements of P^i such that if $\langle b_k : k < n_i^* \rangle \in \prod_{k < n_i^*} Q_k^i$ then $\bigcap \{b_k : k < n_i^*\}$ is not zero.

Let $F^i(x) := \bigcup \{a_k^i : x \cap a_k^i \neq 0_{A_i} \text{ and } l < k \Rightarrow x \cap a_l^i = 0_{A_i}\}$ so the union is on at most one element and $F^i(x) = 0_{A_i} \Leftrightarrow x = 0_{A_i}$.

Let $(A, P, Q, R, F, n^*) := \prod_{i < \kappa} (A_i, P^i, Q^i, R^i, F^i, n_i^*)$. (We consider Q as a two-place relation.) Note that

(*)₁ P is a Boolean subalgebra of A ,

(*)₂ if D is a subset of P^+ then its density in A is equal to its density in P .

[Why? If $Y \subseteq A^+$ is dense in D , then $\{F(c) : c \in Y\}$ is a subset of P^+ dense in D of cardinality $\leq |Y|$; for the other direction use the same set.]

Now let us enumerate the members of n^* as $\{k_\alpha : \alpha < \mu\}$ (no repetitions); we also list the members of P^+ as $\{c_\alpha : \alpha < \mu\}$. Now by induction on $\alpha < \mu$ we choose a member b_α of Q_{k_α} which contains (in A) no one of $\{c_\beta : \beta < \alpha\}$. As each c_β can "object" to at most one $b \in Q_{k_\alpha}$ (as the candidates are pairwise disjoint) and Q_{k_α} has cardinality $\mu > |\alpha|$, we can do this. Also by the choice of the Q^i 's there is a filter of P to which b_α belongs for every $\alpha < \mu$, so we are done as μ is regular.

Case 2 ⁽²⁾: There are $\mu_0 < \mu$ and $n_i^0 < n_i$ such that $\aleph_0 \leq \mu_0 = \prod_{i < \kappa} n_i^0 / D$. We can define X_i, Y_i such that X_i is the family of those subsets of Y_i with exactly n_i^0 elements and $|Y_i| = n_i \times n_i^0 + 1$ and e.g. Y_i is a set of natural numbers; note that $|X_i| > n_i$. Let $X := \langle X_i : i < \kappa \rangle / D$ and $Y := \langle Y_i : i < \kappa \rangle / D$. For $y \in Y$ (in \mathfrak{C} 's sense) let $S_y := \{x \in X : y \in x\}$. Let $n_i^1 := |X_i|$, note $|\prod_{i < \kappa} n_i^1 / D| \geq \mu$; let $\langle a_k^i : k < n_i^1 \rangle$ be a partition of 1_{A_i} to non-zero members of A_i and h_i be a one-to-one function from X_i onto $R_i := \{a_k^i : k < n_i^1\}$, and for $y \in Y_i$ let $b_y^i := \bigcup \{h_i(x) : x \in S_y\} \in A_i$; we define $h, n^1, \langle b_y : y \in Y \rangle \in \mathfrak{C}$ naturally. Let P_i^* be the subalgebra of A_i generated by $\{a_k^i : k < n_i^1\}$ and $P^* = \prod_{i < \kappa} P_i^* / D$ as in the other case. By a cardinality argument if $k < \omega$, then

$$n_i^0 > k \ \& \ y_0, \dots, y_{k-1} \in Y \Rightarrow A_i \models "b_{y_0}^i \cap \dots \cap b_{y_{k-1}}^i \neq 0_{A_i}",$$

hence $\{b_y : y \in Y\} \subseteq P^*$ generates a filter of P^* . Let D be an ultrafilter of P^* containing b_y for $y \in Y$. If $Z \subseteq A \setminus \{0\}$ exemplifies the density of D in A^* and is of cardinality $\mu_2 < \mu$, as in Case 1 without loss of

⁽²⁾ In this case the regularity of μ is not used.

generality $Z = \{a_f : f \in F\} \subseteq P^*$, where $F \subseteq \prod_{i < \kappa} n_i^1$ with $|F| < \mu$ and $a_f := \langle a_{f(i)}^i : i < \kappa \rangle / D$. Let $n = \langle n_i : i < \kappa \rangle / D$ and $n^0 = \langle n_i^0 : i < \kappa \rangle / D$.

For each $f \in F$, $h^{-1}(a_f)$ is from X , so is a subset of Y of cardinality n_0 from the inside (“considered” by \mathfrak{C} to be so) and μ_0 from the outside; there is a set W of n members of X pairwise disjoint; now from the inside W has cardinality n and from the outside it has cardinality μ so there is a member of X disjoint from all the $h^{-1}(a_f)$, a contradiction to density. So D has density μ in A , hence for every ultrafilter F of A extending D , $\pi(F, A) \geq \mu$.

Hence $\pi\chi(A) \geq \mu$ as required. ■_{3.6}

3.7. Remark. (1) If we ignore regularity, Case 1 suffices as for every $\bar{n}/D \in \omega^\kappa/D$ and $\mu = \prod \bar{n}/D \geq \aleph_0$ we can find $\bar{n}^l/D \in \omega^\kappa/D$ such that $\omega^\kappa/D \models “\bar{n} \leq 2^{\bar{n}^{l+1}/D} < \bar{n}^l/D”$, so $\langle |\bar{n}^l/D| : l < \omega \rangle$ is eventually constant.

(2) If each A_i is of cardinality \aleph_0 and $\mu = \aleph_0^\kappa/D$ is regular the proof above gives $\pi\chi(\prod_{i < \kappa} A_i/D) = \mu$ (if 3.6 does not apply, ω^κ/D is μ -like, so we can apply Case 2 with $|X_i| = |Y_i| = \aleph_0$).

(3) By Peterson [P] the regularity of μ is necessary. For singular μ our proof still gives $\pi\chi^+(A) > \text{cf}(\mu)$.

References

- [CK] C. C. Chang and H. J. Keisler, *Model Theory*, Stud. Logic Found. Math. 73, North-Holland, Amsterdam, 1973.
- [KpSh 415] S. Koppelberg and S. Shelah, *Densities of ultraproducts of Boolean algebras*, Canad. J. Math. 47 (1995), 132–145.
- [M] D. Monk, *Cardinal functions of Boolean algebras*, circulated notes.
- [P] D. Peterson, *Cardinal functions on ultraproducts of Boolean algebras*, J. Symbolic Logic, to appear.
- [Sh:92] S. Shelah, *Remarks on Boolean algebras*, Algebra Universalis 11 (1980), 77–89.
- [Sh 355] —, $\aleph_{\omega+1}$ has a Jonsson algebra, in: Cardinal Arithmetic, Oxford Logic Guides 29, Chapter II, Oxford Univ. Press, 1994, 34–116.
- [Sh 420] —, *Advances in cardinal arithmetic*, in: Finite and Infinite Combinatorics in Sets and Logic, N. W. Sauer *et al.* (eds.), Kluwer Acad. Publ., 1993, 355–383.
- [Sh 430] —, *Further cardinal arithmetic*, Israel J. Math., to appear.
- [Sh 513] —, *PCF and infinite free subsets*, Arch. Math. Logic, to appear.
- [Sh:g] —, *Cardinal Arithmetic*, Oxford Logic Guides 29, Oxford Univ. Press, 1994.

Institute of Mathematics
The Hebrew University
Jerusalem, Israel
E-mail: shelah@math.huji.ac.il

Department of Mathematics
Rutgers University
New Brunswick, New Jersey 08903, U.S.A.
E-mail: shelah@math.rutgers.edu

Received 2 February 1994;
in revised form 17 January 1996

Added in proof (June 1996). We can add to Claim 2.3:

CLAIM 2.3(3). Assume that B is a Boolean algebra and $\sigma < \theta = \text{cf}(\theta)$ and $(*)_{\sigma}^B[Y]$ and for no $\tau \in (\sigma, \theta)$ and Y' do we have $(*)_{\tau}^B[Y']$. Then in the conclusion of 2.3(1) we can add: D is normal (hence in 2.2 we get that for arbitrarily large $\theta < \mu$, there are a normal filter D on θ and $\langle y_i : i < \theta \rangle$ as in 2.3(1)).

REMARK. If $\theta = \tau^+$ then 2.3(2) gives the conclusion.

PROOF. Let $Y = \{y_i : i < \theta\}$. We choose by induction on $n < \omega$ a club E_n of θ and a sequence $\bar{y}^n = \langle y_i^n : i < \theta \rangle$ of non-zero members of B such that:

- (a) letting $Y_n = \{y_i^n : i < \theta\}$, we have $Y = Y_0$, $Y_n \subseteq Y_{n+1}$ and $E_{n+1} \subseteq E_n$,
- (b) if $\delta \in E_{n+1}$ and if $\delta < \alpha < \min(E_n \setminus (\delta + 1))$ then for some $\beta < \delta$, $y_{\beta}^{n+1} \leq_B y_{\alpha}^n$.

For $n = 0$, let \bar{y}^0 list Y and $E_0 = \{\delta < \theta : \delta \text{ a limit ordinal divisible by } \sigma\}$.

For $n = m + 1$, for each $\delta \in E_n$, let $\gamma_{\delta} = \min(E_n \setminus (\delta + 1))$ and let Z_{δ}^n be a subset of B^+ of cardinality $\leq \sigma$ dense in $\{y_i^n : \delta \leq i < \gamma_{\delta}\}$, which exists by the proof of 2.2. Let $Z_{\delta}^n = \{z_{\delta+i}^n : i < \sigma\}$ (no double use of the same index).

Also for each $\zeta < \sigma$ let D_{ζ}^n be the normal filter on θ generated by the subsets of θ of the form $\{i < \theta : z \not\leq y_{i+\zeta}^n\}$ for $z \in B^+$; by our assumption toward contradiction there are $z_{\zeta, \varepsilon}^n \in B^+$ for $\varepsilon < \theta$ and club E_{ζ}^n of θ such that if $\delta \in E_{\zeta}^n$ and $\zeta < \sigma$ then for some $\varepsilon < \delta$ we have $z_{\zeta, \varepsilon}^n \leq y_{\delta+\zeta}^n$.

Let E_{n+1} be a club of θ included in E_n and in each E_{ζ}^n and choose \bar{y}^{n+1} such that: its range includes the range of \bar{y}^n and for $\delta_1 < \delta_2 \in E_{n+1}$, $\{y_{\delta+\xi}^{n+1} : \xi < \sigma\}$ includes $\{y_{\delta+\xi}^n : \xi < \sigma\} \cup Z_{\delta}^n$ and $\{y_i^{n+1} : i < \delta\}$ includes each $z_{\zeta, \varepsilon}^n$ for $\zeta < \sigma$ and $\varepsilon < \delta$. The rest is as in the proof of 2.3(2).