Trends in Mathematics, © 1999 Birkhäuser Verlag Basel/Switzerland

## Almost disjoint pure subgroups of the Baer-Specker group

Oren Kolman \* and Saharon Shelah <sup>†</sup>

## Abstract

We prove in ZFC that the Baer-Specker group  $\mathbf{Z}^{\omega}$  has  $2^{\aleph_1}$  non-free pure subgroups of cardinality  $\aleph_1$  which are almost disjoint: there is no non-free subgroup embeddable in any pair.

In this short paper we prove the following result.

**Theorem 1** There exists a family  $\mathbf{G} = \{G_{\alpha} : \alpha < 2^{\aleph_1}\}$  of non-isomorphic non-free pure subgroups of the Baer-Specker group  $\mathbf{Z}^{\omega}$  such that: (1.1) each  $G_{\alpha}$  has cardinality  $\aleph_1$ ; (1.2) if  $\alpha < \beta$ , then  $G_{\alpha}$  and  $G_{\beta}$  are almost disjoint: if H is isomorphic to subgroups of  $G_{\alpha}$  and  $G_{\beta}$ , then H is free. In particular,  $G_{\alpha} \cap G_{\beta}$  is free.

Recall that the Baer-Specker group  $\mathbf{Z}^{\omega}$  is the abelian group of functions from the natural numbers into the integers (see [1] and [18]). It contains the canonical pure free subgroup  $\mathbf{Z}_{\omega} = \bigoplus_{n < \omega} \mathbf{Z}$ . The group  $\mathbf{Z}^{\omega}$  is not  $\kappa$ -free for any cardinal  $\kappa > \aleph_1$ , but it is  $\aleph_1$ -free, so the groups  $G_{\alpha}$  in Theorem 1 are almost free.

Theorem 1 answers a question of the second author, and has its place in the line of recent research dealing with the lattice structure of the pure subgroups of  $\mathbf{Z}^{\omega}$  (see [2], [3], and [5]–[8]). For example, Irwin asked whether there is a subgroup of  $\mathbf{Z}^{\omega}$  with uncountable dual but no free summands of infinite rank. This problem was resolved recently by Corner and Göbel [5] who proved the following stronger fact.

**Theorem 2** [5] The Baer-Specker group  $\mathbf{Z}^{\omega}$  contains a pure subgroup G whose endomorphism ring splits as  $End(G) = \mathbf{Z} \oplus Fin(G)$ , with  $|G^*| = 2^{\aleph_0}$ , where  $\mathbf{Z}$  is the scalar multiplication by integers and Fin(G) is the ideal of all endomorphisms of G of finite rank.

<sup>\*</sup>We thank the referee for constructive comments.

<sup>&</sup>lt;sup>†</sup>This research was partially supported by the German-Israel Foundation project No. G-545-173.06/97; publication number 683.

Quotient-equivalent and almost disjoint abelian groups have been studied by Eklof, Mekler and Shelah in [9]–[11], who showed that under various settheoretic hypotheses, there exist families of maximal possible size of almost free abelian groups which are pairwise almost disjoint. Following [11], we say that two groups A and B are almost disjoint if whenever H is embeddable as a subgroup in both A and B, then H is free. Clearly if A and B are non-free and almost disjoint, then they are non-isomorphic in a very strong way. On the other hand, the intersection of two almost disjoint groups of size  $\aleph_1$  need not necessarily be countable, so group-theoretic almost disjointness differs from its set-theoretic homonym. Theorem 1 establishes in ZFC that the Baer-Specker group contains large families of almost disjoint almost free non-free pure uncountable subgroups.

Our group and set-theoretic notation is standard and can be found in [10] and [14]. For example,  $\omega_1 > 2$  is the set of partial functions from  $\omega_1$  into  $\{0, 1\}$  whose domains are at most countable;  $\omega_1 2$  is the set of all functions from  $\omega_1$  into  $\{0, 1\}$ ; for a regular cardinal  $\chi$ ,  $H(\chi)$  is the family of all sets of hereditary cardinality less than  $\chi$ .

For a set  $A \subseteq H(\chi)$  for  $\chi$  large enough, we write  $\operatorname{dcl}_{(H(\chi), \in, <)}[A]$  for the Skolem closure (Skolem hull) of A in the structure  $(H(\chi), \in, <)$ , where

< is a well-ordering of  $H(\chi)$  (for details, see [16], 400-402, or [15], 165-170). In proving Theorem 1, we shall appeal to the well-known Engelking-Karłowicz theorem from set-theoretic topology:

**Theorem 3** [13] If  $|Y| = \mu = \mu^{<\sigma} < \lambda = |X| \leq 2^{\mu}$ , then there are functions  $h_{\alpha}: X \to Y$  for  $\alpha < \mu$  such that for every partial function f from X to Y of cardinality less than  $\sigma$ , for some  $\alpha < \mu$ ,  $f \subseteq h_{\alpha}$ .

A self-contained short proof can be found in [17], 422-423. We shall need just the case when  $\mu = \sigma = \aleph_0$ , and  $\lambda = 2^{\mu}$ . Since it may be less familiar to algebraists, for convenience we deduce the fact to which we appeal later on (although it also appears as Corollary 3.17 in [4]).

**Lemma 4** There exists a family  $\{f_{\eta} : \eta \in {}^{\omega_1 > 2}\}$  such that  $f_{\eta} : \omega \to \mathbb{Z}$ , and whenever  $\eta_1, \ldots, \eta_k$  are distinct and  $a_1, \ldots, a_k \in \mathbb{Z}$ , then  $\{i < \omega : (\forall l \leq k)(f_{\eta_l}(i) = a_l)\}$  is infinite.

**Proof.** Take  $\mu = \sigma = \aleph_0$ ,  $\lambda = 2^{\mu}$ ,  $X = {}^{\omega_1>2}$  and  $Y = \mathbb{Z}$  in the Engelking-Karłowicz theorem. Since  $|{}^{\omega_1>2}| = 2^{\aleph_0}$  and  $|\mathbb{Z}| = \aleph_0$ , we know that there exist functions  $h_n : {}^{\omega_1>2} \to \mathbb{Z}$  for  $n < \omega$  such that for every partial function f from  ${}^{\omega_1>2}$  to  $\mathbb{Z}$  whose domain is finite, there is some  $m < \omega$  such that  $f \subseteq h_m$ . Let  $\{g_i : i < \omega\}$  be an enumeration with infinitely many repetitions of each  $h_n$  for  $n < \omega$ .

For each  $\eta \in {}^{\omega_1>2}$ , define  $f_{\eta} : \omega \to \mathbb{Z}$  by  $f_{\eta}(i) = g_i(\eta)$ . The family  $\{f_{\eta} : \eta \in {}^{\omega_1>2}\}$  is as required: for if  $\eta_1, \ldots, \eta_k$  are distinct and  $a_1, \ldots, a_k \in$ 

Sh:683

Almost disjoint subgroups

**Z** are given, then the set  $f = \langle (\eta_1, a_1), \ldots, (\eta_k, a_k) \rangle$  is a finite function, so there is some m such that  $f \subseteq h_m$  and it is now easy to see that  $\{i < \omega : (\forall l \leq k) (f_{\eta_l}(i) = a_l)\}$  is infinite.  $\Box$ 

A well-known algebraic fact will also be useful:

**Lemma 5** Let C be a closed unbounded subset of the regular uncountable cardinal  $\kappa$ . Suppose that H is an abelian group of cardinality  $\kappa$ , and  $\langle H_{\alpha} : \alpha < \kappa \rangle$  is a  $\kappa$ -filtration of H (a continuous increasing chain of subgroups  $H_{\alpha}$ ,  $|H_{\alpha}| < \kappa$ , whose union is H). Let  $S = \{\alpha \in C : H/H_{\alpha} \text{ is not } \kappa\text{-free}\}$ . Then H is free if and only if S is non-stationary in  $\kappa$ .

**Proof.** Well known: see Proposition IV.1.7 in [10].  $\Box$ 

We refer the reader to [14] for the definitions of the characteristic  $\chi(g)$  and the type  $\tau(g)$  of an element g in a group.

Now we prove Theorem 1.

**Proof.** Let **P** be the set of prime numbers, and let  $\{P_{\eta} : \eta \in {}^{\omega_1 > 2}\}$  be a family of almost disjoint (infinite) subsets of **P**:  $\eta \neq \nu \in {}^{\omega_1 > 2} \Rightarrow |P_{\eta} \cap P_{\nu}| < \aleph_0$ . By Lemma 4, there exists  $\{f_{\eta} : \eta \in {}^{\omega_1 > 2}\}$  such that  $f_{\eta} : \omega \to \mathbf{Z}$ , and if  $\eta_1, \ldots, \eta_k$  are distinct and  $a_1, \ldots, a_k \in \mathbf{Z}$ , then  $\{i < \omega : (\forall l \leq k)(f_{\eta_l}(i) = a_l)\}$  is infinite.

Define functions  $x_{\eta}$  and  $x_{\eta,j}$  in  $\mathbb{Z}^{\omega}$  as follows. Let  $x_{\eta} = \langle \pi_{\eta,i} \cdot f_{\eta}(i) : i < \omega \rangle$  where  $\pi_{\eta,i} = \Pi\{p \in P_{\eta} : p < i\}$ , and let  $x_{\eta,j} = \langle \pi_{\eta,i}^{j} \cdot f_{\eta}(i) : i < \omega \rangle$  where  $\pi_{\eta,i}^{j} = \Pi\{p \in P_{\eta} : j \le p < i\}$  (=0 if  $i \le j$ ). Note that  $x_{\eta} = x_{\eta,0}$ .

For  $\eta \in {}^{\omega_1}2$ , let  $G_{\eta}$  be the subgroup of  $\mathbf{Z}^{\omega}$  generated by  $\mathbf{Z}_{\omega} \cup \{x_{\eta \mid \alpha, j} : \alpha < \omega_1, 0 \le j < \omega\}$ .

We show that the family  $\mathbf{G} = \{G_{\eta} : \eta \in {}^{\omega_1}2\}$  satisfies the conclusions of Theorem 1.

**Claim 1:**  $G_{\eta}$  is pure in  $\mathbf{Z}^{\omega}$ .

**Proof of Claim 1:** Suppose that rx = g for some  $x \in \mathbf{Z}^{\omega}$ ,  $r \in \mathbf{N}$ , and  $g \in G_{\eta}$ . Say  $g = y + n_1 x_{\eta \mid \alpha_1, j_1} + \cdots + n_m x_{\eta \mid \alpha_m, j_m}$ ,  $n_l \neq 0$ , with  $y \in \mathbf{Z}_{\omega}$ . Without loss of generality (adding more elements from  $\mathbf{Z}_{\omega}$  to the RHS if necessary),  $(\forall l \leq m)(j_l = j)$  for some  $j < \omega$ , j > r, y(i) = 0  $(\forall i > j)$ , and x(i) = 0  $(\forall i \leq j)$ . Relabelling (if necessary), we may assume that  $\alpha_1 < \cdots < \alpha_m < \omega_1$ , and because  $x_{\eta \mid \alpha_l, j}(i) = 0$  if  $i \leq j$ , we may write

$$rx = ry^* + n_1 x_{\eta|\alpha_1, j} + \dots + n_m x_{\eta|\alpha_m, j}, \quad \text{for some } y^* \in \mathbf{Z}_{\omega}.$$

Fix  $k \in \{1, \ldots, m\}$ . Since  $\eta | \alpha_1, \ldots, \eta | \alpha_m$  are distinct  $(\alpha_1 < \cdots < \alpha_m)$ , letting  $a_l = \delta_{kl}$  (Kronecker delta), we know that the set  $N_k = \{i < \omega :$ 

O. Kolman and S. Shelah

 $(\forall l \neq k)(f_{\eta_l}(i) = 0, f_{\eta_k}(i) = 1)\}$  is infinite. For large enough *i* in this set (e.g.  $i > \max_{1 \leq l \leq m} [\min(P_{\eta|\alpha_l} \setminus \{0, \ldots, j\})]), x_{\eta|\alpha_l,j}(i)$  is zero if and only if  $l \neq k$ . So for infinitely many  $i < \omega$ , for  $l \neq k, x_{\eta|\alpha_l,j}(i) = 0$ , and  $x_{\eta|\alpha_k,j}(i) \neq 0$ .

Unfix k. For each  $k \leq m$ , for infinitely many  $i \in (j, \omega) \cap N_k$ ,  $rx(i) = n_k x_{\eta \mid \alpha_k, j}(i) = n_k \Pi\{p \in P_{\eta \mid \alpha_k} : j \leq p < i\}$ . Since r < j, we must have  $rs_k = n_k$  for some  $s_k$  in  $\mathbb{Z}$ , and therefore  $x = y^* + s_1 x_{\eta \mid \alpha_1, j} + \cdots + s_m x_{\eta \mid \alpha_m, j} \in G_{\eta}$  ( $G_{\eta}$  is torsion-free). Hence  $G_{\eta}$  is pure in  $\mathbb{Z}^{\omega}$ , which establishes Claim 1.

**Claim 2:**  $G_{\eta}$  has cardinality  $\aleph_1$ , so (1.1) holds.

**Proof of Claim 2:** If  $\xi \neq \zeta \in {}^{\omega_1>2}$ , then for some  $j < \omega$ ,  $P_{\xi} \cap P_{\zeta} \subseteq j$ . Pick p, q > j with  $p \in P_{\xi}$  and  $q \in P_{\zeta}$ ; so the set  $B = \{i < \omega : f_{\xi}(i) = p \text{ and } f_{\zeta}(i) = q\}$  is infinite, and if  $i \in B$  is bigger than  $\max\{j, p, q\}$ , then  $x_{\xi,j}(i) \neq x_{\zeta,j}(i)$ , since  $x_{\xi,j}(i)$  is non-zero and divisible by  $p^2$  but by no prime in  $P_{\zeta}$ , and  $x_{\zeta,j}(i)$  is non-zero and divisible by  $q^2$  but by no prime in  $P_{\xi}$ . It follows that  $G_{\eta}$  has cardinality  $\aleph_1$ . After this observation, a second's reflection on the element types of  $G_{\eta_1}$  and  $G_{\eta_2}$  (for  $\eta \neq \nu$ ) should convince the reader that the groups are neither isomorphic nor free.

**Claim 3:** (1.2) holds: if  $\eta_1 \neq \eta_2 \in {}^{\omega_1}2$ , then  $G_{\eta_1}$  and  $G_{\eta_2}$  are almost disjoint.

**Proof of Claim 3:** Suppose (towards a contradiction) that for some  $\eta_1 \neq \eta_2 \in {}^{\omega_1}2$ , for some non-free abelian group H, there exist isomorphisms  $\varphi_l : H \to \operatorname{range}(\varphi_l) \leq G_{\eta_l}, \ l = 1, 2$ . Since  $G_{\eta_l}$  is  $\aleph_1$ -free, H must have cardinality  $\aleph_1$ . Let  $\langle H_i : i < \omega_1 \rangle$  be an  $\omega_1$ -filtration of H. Without loss of generality, we may assume that each  $H_i$  is pure in H, so that  $H/H_i$  is torsion-free.

Let  $G_{\eta,i} = \langle \mathbf{Z}_{\omega} \cup \{ x_{\eta|\beta,j} : j < \omega, \beta < i \} \rangle$  for  $i < \omega_1$  and  $\eta \in \{ \eta_1, \eta_2 \}$ .

Note that  $\langle G_{\eta,i} : i < \omega_1 \rangle$  is a  $\omega_1$ -filtration of  $G_\eta$ , since it is increasing and continuous with union  $G_\eta$ , and each  $G_{\eta,i}$  is countable. For large enough  $\chi$ , the set C defined by  $\{\delta < \omega_1 : \operatorname{dcl}_{(H(\chi), \in, <)} | \delta \cup \{G_{\eta_1}, G_{\eta_2}, \{x_\nu, f_\nu : \nu \in \omega_1 > 2\}, \eta_1, \eta_2, \varphi_1, \varphi_2, \{H_i : i < \omega_1\}\} ] \cap \omega_1 = \delta\}$  is a club of  $\omega_1$  (well known, or see [16], 401). Note that if  $\delta \in C$ , then  $\varphi_l$  maps  $H_\delta$  into  $G_{\eta_l,\delta}$ . Since H is not free, it follows by Lemma 5 that  $S = \{\delta \in C : H/H_\delta \text{ is not } \aleph_1 - free\}$  is stationary. By Pontryagin's Criterion, for each  $\delta \in S$ ,  $H/H_\delta$  has a non-free (torsion-free) subgroup  $K_\delta/H_\delta$  of finite rank  $n_\delta + 1$  such that every subgroup of  $K_\delta/H_\delta$  of rank less than  $n_\delta + 1$  is free. Let  $H_\delta^+/H_\delta$ be a pure subgroup of  $K_\delta/H_\delta$  of rank  $n_\delta$ . Then  $H_\delta^+/H_\delta$  is free with basis  $y_0 + H_\delta, \ldots, y_{n_\delta - 1} + H_\delta$  say. So  $K_\delta/H_\delta^+ \simeq (K_\delta/H_\delta)/(H_\delta^+/H_\delta)$  is a torsionfree rank-1 group which is not free, and hence there is a non-zero element  $y_{n_\delta} + H_\delta^+$  which is divisible in  $K_\delta/H_\delta^+$  by infinitely many natural numbers. Call this set of natural numbers A.

Sh:683

228

For l = 1, 2, for large enough  $j_l(*) < \omega$ , and  $\beta_0^l < \cdots < \beta_{k_l}^l < \omega_1$ ,  $\varphi_l(y_m)$  is an element of the subgroup of  $G_{\eta_l}$  generated by  $G_{\eta_l,\delta} \cup \{x_{\eta_l}|\beta_{0,j_l}(*), \ldots, x_{\eta_l}|\beta_{k_l,j_l}(*)\}$  for all  $m \le n_{\delta}$ .

Taking large enough  $\delta \in S$ , we may assume that  $\min\{\alpha : \eta_1 | \alpha \neq \eta_2 | \alpha\} < \beta^l_0, \ l = 1, 2$ . Since  $\delta \in C$ , we can show the following claims: (\*)<sub>1</sub>: The set A does not contain infinitely many powers of one prime.

(\*)<sub>2</sub>: The set  $Q = (\mathbf{P} \cap A) \subseteq P_{\eta_l \mid \beta^l_0} \cup \cdots \cup P_{\eta_l \mid \beta^l_{k_l}}$ .

Now  $(*)_1$  is true because non-zero sums of elements in  $G_{\eta_l,\delta} \cup \{x_{\eta_l|\beta^l_0,j_l(*)}, \ldots, x_{\eta_l|\beta^l_{k_l},j_l(*)}\}$  are divisible by at most finitely many powers of any given prime (by the definition of the elements  $x_{\eta_l|\beta,j}$ ). Note that  $\chi(y_{n_{\delta}} + H_{\delta}^{+}) = \bigcup_{\{y \in y_{n_{\delta}} + H_{\delta}^{+}\}} \chi(y) \leq \bigcup_{\{y \in y_{n_{\delta}} + H_{\delta}^{+}\}} \chi(\varphi_l(y))$ , where the characteristics are taken relative to  $K_{\delta}/H_{\delta}^{+}$ ,  $K_{\delta}$  and  $G_{\eta_l,\delta} \cup \{x_{\eta_l|\beta^l_0,j_l(*)}, \ldots, x_{\eta_l|\beta^l_{k_l},j_l(*)}\}$  respectively. Hence  $(*)_1$  holds. By  $(*)_1$ , since A is infinite, the set  $Q = \mathbf{P} \cap A$  is infinite.

Also, the same characteristic inequality implies that  $Q \subseteq P_{\eta_l | \beta_0} \cup \cdots \cup P_{\eta_l | \beta_{k_l}}$ . So  $(*)_2$  is true. Hence,  $Q \subseteq \bigcap_{l=1,2} (\bigcup_{k \leq k_l} P_{\eta_l | \beta_k})$  which is finite (since the family  $\{P_{\eta} : \eta \in \omega_1 > 2\}$  is almost disjoint). This is a contradiction, and so Claim 3 follows, completing the proof of Theorem 1.

**Corollary 6** Every non-slender  $\aleph_1$ -free abelian group G has a family  $\{G_{\alpha} : \alpha < 2^{\aleph_1}\}$  of non-free subgroups such that: (6.1) each  $G_{\alpha}$  is almost free of cardinality  $\aleph_1$ ; (6.2) if  $\alpha < \beta$ , then  $G_{\alpha}$  and  $G_{\beta}$  are almost disjoint.

**Proof.** By Nunke's characterisation of slender groups (see Corollary IX.2.5 in [10] for example), G must contain a copy of the Baer-Specker group.

**Remark**: For the same reason, the corollary is true for any non-slender cotorsion-free abelian group.

## References

- R. Baer, Abelian groups without elements of finite order, Duke Math. J. 3 (1937), 68-122.
- [2] A. Blass, Cardinal characteristics and the product of countably many infinite cyclic groups, J. Algebra 169 (1994), 512-540.
- [3] A. Blass and R. Göbel, Subgroups of the Baer-Specker group with few endomorphisms but large dual, Fund. Math. 149 (1996), 19-29.
- [4] W. W. Comfort and S. Negrepontis, The Theory of Ultrafilters, Berlin: Springer-Verlag (1974).

## O. Kolman and S. Shelah

- [5] A. L. S. Corner and R. Göbel, Essentially rigid subgroups of the Baer-Specker group, Manuscripta math. 94 (1997), 319-326.
- [6] A. L. S. Corner and B. Goldsmith, On endomorphisms and automorphisms of some pure subgroups of the Baer-Specker group, in Abelian Group Theory and Related Topics (R. Göbel, P. Hill and W. Liebert, eds.), Contemp. Math. 171 (1994), 69-78.
- [7] M. Dugas and J. Irwin, On pure subgroups of cartesian products of integers, Results in Math. 15 (1989), 35-52.
- [8] K. Eda, A note on subgroups of Z<sup>N</sup>, in Abelian Group Theory (R. Göbel et al, eds.), Springer Lecture Notes in Mathematics 1006 (1983), 371–374.
- [9] P. C. Eklof and A. H. Mekler, Infinitary stationary logic and abelian groups, Fund. Math. 112 (1981), 1-15.
- [10] P. C. Eklof and A. H. Mekler, Almost Free Modules. Set-theoretic Methods, Amsterdam: North-Holland (1990).
- [11] P. C. Eklof, A. H. Mekler, and S. Shelah, Almost disjoint abelian groups, Israel J. Math. 49 (1984), 34-54.
- P. C. Eklof and S. Shelah, A combinatorial principle equivalent to the existence of non-free Whitehead groups, in Abelian Group Theory and Related Topics (R. Göbel, P. Hill and W. Liebert, eds.), Contemp. Math. 171 (1994), 79-98.
- [13] R. Engelking and M. Karłowicz, Some theorems of set theory and their topological consequences, Fund. Math. 57 (1965), 275-285.
- [14] L. Fuchs, Infinite Abelian Groups, Vols. I and II, New York: Academic Press (1970, 1973).
- [15] L. Heindorf and L. B. Shapiro, Nearly Projective Boolean Algebras, Springer Lecture Notes in Mathematics 1596 (1994).
- [16] S. Shelah, Classification Theory and the Number of Non-Isomorphic Models, Revised edition, Studies in Logic and the Foundations of Mathematics, Vol. 92, Amsterdam: North-Holland (1990).
- [17] S. Shelah, Cardinal Arithmetic, Oxford: OUP (1994).
- [18] E. Specker, Additive Gruppen von Folgen ganzer Zahlen, Portugaliae Math. 9 (1950), 131-140.

230

Sh:683