

ON $\text{CON}(\mathfrak{d}_\lambda > \text{COV}_\lambda(\text{MEAGRE}))$
SH:945

SAHARON SHELAH

ABSTRACT. We prove the consistency of: for suitable strongly inaccessible cardinal λ the dominating number, i.e., the cofinality of ${}^\lambda\lambda$, is strictly bigger than $\text{cov}_\lambda(\text{meagre})$, i.e. the minimal number of nowhere dense subsets of ${}^\lambda 2$ needed to cover it. This answers a question of Matet.

Date: 2021-01-11.

2010 *Mathematics Subject Classification.* Primary 03E35, 03E55; Secondary: 03E17.

Key words and phrases. set theory, independence, forcing, cardinal invariants, inaccessible.

The author thanks Alice Leonhardt for the beautiful typing. This research was supported by the United States-Israel Binational Science Foundation. First version (in TeX Sept/16/2005 as F741). First version in LaTeX January 21, 2009.

§ 0. INTRODUCTION

Cardinal characteristics were defined, historically, over the continuum. See celebrated Van Dowen [vD84], for the general topologist perspective and excellent survey Blass [Bla], Bartoszyński [Bar10] for the set theoretic perspective. In recent years there are many results concerning generalized cardinal characteristics. The idea is to imitate the definition of a given characteristic over the continuum, by translating it to uncountable cardinals.

It is reasonable to distinguish regular cardinals and singular cardinals. Among the regular cardinals, it make sense to distinguish limit cardinals from successor cardinals. In this paper we focus on strongly inaccessible cardinals. These cardinals and their characteristics behave, in many cases, much like \aleph_0 , but certainly not always. See Landver [Lan92], Cummings-Shelah [CS95] and Matet-Shelah [MS]. Our main result is the consistency of $\text{cov}_\lambda(\text{meagre}) < \mathfrak{d}_\lambda$ at a supercompact cardinal λ , and we begin with the following definitions:

We shall define three cardinal invariants (but the paper deals, actually, just with two of them):

Definition 0.1. The bounding and dominating numbers.

Let λ be an inaccessible cardinal.

Let $f, g \in {}^\lambda\lambda$

- (a) $f \leq^* g$ if $|\{\alpha < \lambda : f(\alpha) > g(\alpha)\}| < \lambda$
- (b) $A \subseteq {}^\lambda\lambda$ is unbounded if there is no $h \in {}^\lambda\lambda$ so that $f \in A \Rightarrow f \leq^* h$
- (c) $A \subseteq {}^\lambda\lambda$ is dominating when for every $f \in {}^\lambda\lambda$ there exists $g \in A$ so that $f \leq^* g$
- (d) the bounding number for λ , denoted by \mathfrak{b}_λ , is $\min\{|A| : A \text{ is unbounded in } {}^\lambda\lambda\}$
- (e) the dominating number for λ , denoted by \mathfrak{d}_λ , is $\min\{|A| : A \text{ is dominating in } {}^\lambda\lambda\}$.

Notice that the usual definitions of \mathfrak{b} and \mathfrak{d} are \mathfrak{b}_{\aleph_0} and \mathfrak{d}_{\aleph_0} according to Definition 0.1. The definition of $\text{cov}_\lambda(\text{meagre})$ involves some topology.

Definition 0.2. The meagre covering number.

Let λ be a regular cardinal

- (a) ${}^\lambda 2$ is the space of functions from λ into 2
- (b) $({}^\lambda 2)^{[\nu]} = \{\eta \in {}^\lambda 2 : \nu \triangleleft \eta\}$, for $\nu \in {}^{\lambda > 2} 2 := \bigcup_{\alpha < \lambda} {}^\alpha 2$
- (c) $\mathcal{U} \subseteq {}^\lambda 2$ is open in the topology $({}^\lambda 2)_{< \lambda}$, iff for every $\eta \in \mathcal{U}$ there exists $i < \lambda$ so that $({}^\lambda 2)^{[\eta \upharpoonright i]} \subseteq \mathcal{U}$
- (d) $\text{cov}_\lambda(\text{meagre})$ is the minimal cardinality of a family of meagre subsets of $({}^\lambda 2)_{< \lambda}$, which covers this space.

This paper deals with the relationship between \mathfrak{d}_λ and $\text{cov}_\lambda(\text{meagre})$. If λ is a successor cardinal then $\text{cov}_\lambda(\text{meagre}) < \mathfrak{d}_\lambda$ is consistent (see (b) below). Matet asked (a personal communication) whether $\mathfrak{d}_\lambda \leq \text{cov}_\lambda(\text{meagre})$ is provable in ZFC, where λ is strongly inaccessible. We give here a negative answer.

For λ a supercompact cardinal and $\lambda < \kappa = \text{cf}(\kappa) < \mu = \mu^\lambda$, we force large \mathfrak{d}_λ i.e., $\mathfrak{d}_\lambda = \mu$ and small covering number (i.e., $\text{cov}_\lambda(\text{meagre}) = \kappa$). A similar result

should hold also for a wider class of cardinals and we intend to return elsewhere to this subject.

Let us sketch some known results. These results are related to the inequality number and the covering number for category. Recall:

Definition 0.3. The inequality number.

Let κ be an infinite cardinal. The inequality number of $\kappa, \mathfrak{e}_\kappa$, is the minimal cardinal λ satisfying that there is a set $\mathcal{F} \subseteq {}^\kappa\kappa$ of cardinality λ such that there is no $g \in {}^\kappa\kappa$ satisfying $(\forall f \in \mathcal{F})(\exists^\kappa \alpha < \kappa)(f(\alpha) = g(\alpha))$.

For $\kappa = \aleph_0, \mathfrak{e}_\kappa = \text{cov}_{\aleph_0}(\text{meagre})$; see Bartoszyński (in [Bar87]) and Miller (in [Mil82]).

Now

- (a) the statement $\mathfrak{e}_\kappa = \text{cov}_\kappa(\text{meagre})$ is valid for $\kappa > \aleph_0$, in the case that κ is strongly inaccessible, by [Lan92]. But if κ is a successor cardinal, it may fail
- (b) if $\kappa < \kappa^{<\kappa}$, then $\text{cov}_\kappa(\text{meagre}) = \kappa^+$. This is due to Landver (in [Lan92]).

We intend also to address:

Problem 0.4. Can we replace “supercompact” by “strongly inaccessible”?

Problem 0.5. 1) Can we prove the consistency of $\text{cov}_\lambda(\text{meagre}) < \mathfrak{b}_\lambda$?
2) For λ strongly inaccessible (or just Laver indestructible supercompact) is there a non-trivial λ^+ -c.c. ($< \lambda$)-strategically complete forcing notion \mathbb{Q} which is ${}^\lambda\lambda$ -bounding?

We thank the referee, Shimoni Garti and Haim Horowitz for helpful comments and pressuring me to expand some proofs and Johannes Schürz for a helpful comment. We say more in subsequent works [She17], [Shea] and in preparation [Shec].

A point which in a previous version was just a step along the way, the referee asked to justify fully, was analyzed to be serious. This was done but eventually is separated to [Sheb]. A posteriori the point is that in the parallel case for $\lambda = \aleph_0$, for full memory FS iteration such a claim is true. In fact, by Judah-Shelah [JS88], if $\langle \mathbb{P}_\alpha, \mathbb{Q}_\beta : \alpha \leq \alpha(*), \beta < \alpha(*) \rangle$ is FS iteration of Suslin-c.c.c. forcing notion, \mathbb{Q}_β with the generic $\eta_\beta \in {}^\omega\omega$ and for notational transparency, its definition is with no parameter and $\zeta : \beta(*) \rightarrow \alpha(*)$ is increasing and $\mathbb{P} = \langle \mathbb{P}'_\alpha, \mathbb{Q}'_\beta : \alpha \leq \beta(*), \beta < \beta(*) \rangle$ is FS iteration, but \mathbb{Q}'_β is defined exactly as $\mathbb{Q}_{\zeta(\beta)}$ is but now in $\mathbf{V}^{\mathbb{P}'_\beta}$ rather than in $\mathbf{V}^{\mathbb{P}_{\zeta(\beta)}}$ then $\Vdash_{\mathbb{P}_{\alpha(*)}} \langle \eta_{\zeta(\beta)} : \beta < \beta(*) \rangle$ is generic for $\mathbb{P}'_{\beta(*)}$ over \mathbf{V} .

Now this is not clear to us for ($< \lambda$)-support iteration of ($< \lambda$)-strategically complete forcing notions. The solution is essentially to change the iteration: to use a “quite generic” ($< \lambda$)-support iteration which “includes” the one we like and use the complete subforcing it generates; see [Sheb].

We try to use standard notation. We use $\theta, \kappa, \lambda, \mu, \chi$ for cardinals and $\alpha, \beta, \gamma, \delta, \varepsilon, \zeta$ for ordinals. We use also i and j as ordinals. We adopt the Cohen convention that $p \leq q$ means that q gives more information, in forcing notions. The symbol \triangleleft is preserved for “being an initial segment”. Also recall ${}^B A = \{f : f \text{ a function from } B \text{ to } A\}$ and let ${}^\alpha A = \cup\{\beta A : \beta < \alpha\}$, some prefer ${}^{<\alpha} A$, but ${}^\alpha A$ is used systematically in the author’s papers. Lastly, J_λ^{bd} denotes the ideal of the bounded subsets of λ .

For exact references to [Sheb] see the introduction there, just before Def 01.1.

The picture of cardinal invariants related to uncountable λ is related but usually quite different than the one for \aleph_0 , they are more similar if κ is “large” enough, mainly strongly inaccessible.

§ 1. PRELIMINARIES

Definition 1.1. Let λ be supercompact. We say that $h : \lambda \rightarrow \mathcal{H}(\lambda)$ is a Laver diamond (for λ) when for every $x \in \mathbf{V}$ there are a normal fine ultrafilter D over $I = [\mathcal{H}(\chi)]^{<\lambda}$ for some χ such that $x \in \mathcal{H}(\chi)$ and the Mostowski collapse \mathbf{j} on \mathbf{V}^I/D maps $\langle h(\sup(u \cap \kappa)) : u \in I \rangle / D$ to x ; (we can use elementary embeddings instead of an ultrafilter).

Notation 1.2. If \mathbb{P} is a forcing notion in \mathbf{V} then $\mathbf{V}^{\mathbb{P}}$ denotes $\mathbf{V}[\mathbf{G}]$ for $\mathbf{G} \subseteq \mathbb{P}$ generic over \mathbf{V} ; we may write $\mathbf{V}[\mathbb{P}]$ instead.

The most straightforward way to increase \mathfrak{b}_λ in the classical case of \aleph_0 is Hechler forcing = dominating real forcing. A condition is a function $f_p : \omega \rightarrow \omega$ which is separated into a finite stem η_p and the rest of the function. Formally, $p = (\eta_p, f_p)$ where $\eta_p \trianglelefteq f_p$.

If p, q are conditions then $p \leq q$ iff $\eta_p \trianglelefteq \eta_q$ and $f_q(n) \geq f_p(n)$ for every $n \notin \text{dom}(\eta_p)$ hence for every n . A generic objects adds a function $g : \omega \rightarrow \omega$ which dominates the functions from the ground model. By iterating Hechler reals one increases the bounding number \mathfrak{b} .

If $\lambda = \lambda^{<\lambda}$ then one can define the generalized Hechler forcing \mathbb{D}_λ by replacing ω by λ . The basic step is λ -complete and λ^+ -c.c. and actually λ -centered. Hence one can iterate and increase \mathfrak{b}_λ .

In [She92, §1, §2] and then Goldstern-Shelah [GS93] Kellner-Shelah [KS12] consider other invariants. Consider two functions $f, g : \omega \rightarrow (\omega \setminus \{0\})$ going to infinity such that $f \geq g$ and ask about:

- $\mathfrak{c}_{f,g}^+ = \min\{|\mathcal{F}| : \mathcal{F} \subseteq \prod_i [f(i)]^{g(i)} \text{ and } (\forall \eta \in \prod_i f(i)) (\exists g \in \mathcal{F}) [\bigwedge_i \eta(i) \in g(i)]\}$
- $\mathfrak{c}_{f,g}^- = \min\{|\mathcal{F}| : \mathcal{F} \subseteq \prod_i f(i) \text{ and for no } g \in \prod_i [f(i)]^{g(i)} \text{ do we have } (\forall \eta \in \mathcal{F}) (\forall^\infty i) (\eta(i) \in g(i))\}$.

There are relevant forcing notions; we shall use a λ^+ -c.c. one as in c.c.c. creature forcing (see [RS97],[HS]).

For transparency

Convention 1.3. Below $\lambda, \bar{\theta}$ are as in 1.4 below.

Definition 1.4. Let λ be inaccessible, $\bar{\theta} = \langle \theta_\varepsilon : \varepsilon < \lambda \rangle$ be a sequence of regular cardinals $< \lambda$ satisfying $\theta_\varepsilon > \varepsilon$.

1) We define the forcing notion $\mathbb{Q} = \mathbb{Q}_{\bar{\theta}}$ by

- (a) $p \in \mathbb{Q}$ iff
 - (a) $p = (\eta, f) = (\eta^p, f^p)$
 - (b) $\eta \in \prod_{\zeta < \varepsilon} \theta_\zeta$ for some $\varepsilon < \lambda$, (η is called the trunk of p)
 - (c) $f \in \prod_{\zeta < \lambda} \theta_\zeta$
 - (d) $\eta \triangleleft f$

- (β) $p \leq_{\mathbb{Q}} q$ iff
- (a) $\eta^p \trianglelefteq \eta^q$
 - (b) $f^p \leq f^q$, i.e. $(\forall \varepsilon < \lambda) f^p(\varepsilon) \leq f^q(\varepsilon)$
 - (c) if $\ell g(\eta^p) \leq \varepsilon < \ell g(\eta^q)$ then $\eta^q(\varepsilon) \in [f^p(\varepsilon), \lambda)$, actually follows.

2) The generic is $\eta = \cup \{\eta^p : p \in \mathbf{G}_{\mathbb{Q}_{\bar{\delta}}}\}$.

The new forcing defined above is not λ -complete anymore. By fixing a stem η one can define a short increasing sequence of conditions which goes up to some θ_ζ at the ζ -th coordinate and hence has no upper bound in $\prod_{\zeta < \varepsilon} \theta_\zeta$. However, this

forcing is $(< \lambda)$ -strategically complete since the COM (= completeness) player can increase the stem at each move.

Remark 1.5. The forcing is parallel to the creature forcing from [She92, §1,§2], [KS12] but they are ${}^\omega\omega$ -bounding.

Recall

Definition 1.6. 1) We say that a forcing notion \mathbb{P} is α -strategically complete when for each $p \in \mathbb{P}$ in the following game $\mathcal{D}_\alpha(p, \mathbb{P})$ between the players COM and INC, the player COM has a winning strategy.

A play lasts α moves; in the β -th move, first the player COM chooses $p_\beta \in \mathbb{P}$ such that $p \leq_{\mathbb{P}} p_\beta$ and $\gamma < \beta \Rightarrow q_\gamma \leq_{\mathbb{P}} p_\beta$ and second the player INC chooses $q_\beta \in \mathbb{P}$ such that $p_\beta \leq_{\mathbb{P}} q_\beta$.

The player COM wins a play if he has a legal move for every $\beta < \alpha$.

2) We say that a forcing notion \mathbb{P} is $(< \lambda)$ -strategically complete when it is α -strategically complete for every $\alpha < \lambda$.

Basic properties of $\mathbb{Q}_{\bar{\delta}}$ are summarized and proved in [GS12, §2].

The following fact describes some immediate connections between various concepts of completeness:

Fact 1.7.

- (a) if $\bar{\mathbb{Q}} = \langle \mathbb{P}_\alpha, \mathbb{Q}_\beta : \alpha \leq \delta, \beta < \delta \rangle$ is a $(< \lambda)$ -support iteration of $(< \lambda)$ -strategically complete forcing notions, then \mathbb{P}_δ is also $(< \lambda)$ -strategically complete; (see e.g. [She00]).
- (b) If \mathbb{P} is $(< \lambda)$ -strategically complete forcing notion then $({}^{\lambda > \text{Ord}} \mathbf{V})^{\mathbf{V}} = ({}^{\lambda > \text{Ord}} \text{Ord})^{\mathbf{V}^{\mathbb{P}}}$, and consequently λ is strongly inaccessible in $\mathbf{V}^{\mathbb{P}}$
- (c) like (a) replacing “ $(< \lambda)$ -strategically complete” by “ $(< \lambda)$ -complete”
- (d) if \mathbb{P} is $(< \lambda)$ -complete then \mathbb{P} is λ -strategically complete.

Definition 1.8. For an ordinal $\alpha_* = \alpha(*)$ let $\mathbf{Q}_{\lambda, \bar{\theta}, \alpha(*)}$ be the class of quadruples $\mathbf{q} = (\bar{u}, \bar{\mathcal{P}}, \bar{\mathbb{P}}, \bar{\mathbb{Q}}, \bar{\eta})$ consisting of (omitting α_* means for some α_* and $\ell g(\mathbf{q}) = \alpha_{\mathbf{q}} = \alpha_*$):

- (a) $\bar{u} = \langle u_\alpha : \alpha < \alpha_* \rangle$ and $\bar{\mathcal{P}} = \langle \mathcal{P}_\alpha : \alpha < \alpha_* \rangle$ where $\mathcal{P}_\alpha \subseteq [u_\alpha]^{\leq \lambda}$, $u_\alpha \subseteq \alpha$, without loss of generality \mathcal{P}_α is closed under subsets (but is not necessarily an ideal)
- (b) $\langle \mathbb{P}_{0, \alpha}, \mathbb{Q}_{0, \beta} : \alpha \leq \alpha_*, \beta < \alpha_* \rangle$ is a $(< \lambda)$ -support iteration and let $\mathbb{P}_{\mathbf{q}, 0} = \mathbb{P}_{\mathbf{q}, 0, \alpha(\mathbf{q})}$

- (c) each of \mathbb{P}_α is strategically $(< \lambda)$ -complete and λ^+ -c.c.
- (d) $\eta_\beta \in \prod_{\varepsilon < \lambda} \theta_\varepsilon$ is the generic of \mathbb{Q}_β where η_β , the generic of \mathbb{Q}_β (defined in clause (e) below) is $\cup\{\eta_p : p \in \mathbf{G}_{\mathbb{Q}_\beta}\}$
- (e) if $\mathbf{G} \subseteq \mathbb{P}_\beta$ is generic over \mathbf{V} then $\eta_\alpha[\mathbf{G}]$ in $(\prod_{\varepsilon < \lambda} \theta_\varepsilon, <_{J_\lambda^{\text{bd}}})$ dominate every $\nu \in \prod_{\varepsilon < \lambda} \theta_\varepsilon$ from $\mathbf{V}[\langle \eta_\gamma : \gamma \in u \rangle]$ when $u \in \mathcal{P}_\alpha$; moreover, in $\mathbf{V}[\mathbf{G}]$
 - (*) $\mathbb{Q}_\beta[\mathbf{G}]$ is the subforcing of $\mathbb{Q}_{\bar{\theta}}$ consisting of the $p \in \mathbb{Q}_{\bar{\theta}}$ such that: for some $\bar{s}, \underline{f}, \eta_p$ (so $\eta_p = \eta$, etc.) we have:
 - (α) $p = (\eta, f) = (\eta_p, f_p)$ so $\eta \in \prod_{\varepsilon < \zeta} \theta_\varepsilon$ for some $\zeta < \lambda$
 - (β) $\bar{s} = \langle (u_i, f_i) : i < i_* \rangle$
 - (γ) $i_* < \lambda$
 - (δ) for each $i < i_*$ we have $u_i \in \mathcal{P}_{\beta, \eta} \triangleleft f_i \in \prod_{\varepsilon < \lambda} \theta_\varepsilon$ and $f_i \in \mathbf{V}[\langle \eta_\gamma[\mathbf{G}] : \gamma \in u_i \rangle]$
 - (ε) $f = \sup\{f_i : i < i_*\}$, i.e. $\varepsilon < \lambda \Rightarrow f(\varepsilon) = \cup\{f_i(\varepsilon) : i < i_*\}$
- (f) notation: so $u_{\mathbf{q}, \alpha} = u_\alpha, \mathbb{P}_{\mathbf{q}, \alpha} = \mathbb{P}_\alpha$, etc., but when \mathbf{q} is clear from the context we may omit it.

Definition 1.9. For $\mathbf{q} \in \mathbf{Q}_{\lambda, \bar{\theta}, \alpha(*)}$.

1) We let $\alpha \leq \alpha_*$, $\mathbb{P}_{1, \alpha} = \mathbb{P}_{1, \alpha}^{\mathbf{q}}$ be essentially the completion of \mathbb{P}_α ; we express it by:

- (*)₁ the elements of $\mathbb{P}_{1, \alpha} = \mathbb{P}_{\mathbf{q}, 1, \alpha}$ are of the form $\mathbf{B}(\dots, \eta_{\gamma_i}, \dots)_{i < i(*)}$ where:
 - (α) $i_* = i(*) \leq \lambda$
 - (β) $\gamma_i \in \mathcal{U}$ for $i < i_*$
 - (γ) \mathbf{B} is a λ -Borel function from ${}^{i(*)}(\prod_{\varepsilon < \lambda} \theta_\varepsilon)$ into $\{0, 1\} = \{\text{false}, \text{true}\}$; \mathbf{B} is from \mathbf{V} , of course, such that $\Vdash_{\mathbb{P}_{\mathbf{q}}} \text{“}\mathbf{B}(\dots, \eta_{\gamma_i}, \dots)_{i < i(*)} = 0\text{”}$
- (*)₂ the order is natural: $\mathbb{P}_{1, \alpha} \Vdash \text{“}\mathbf{B}_1(\dots, \eta_{\gamma(i, 1)}, \dots)_{i < i(1)} \leq \mathbf{B}_2(\dots, \eta_{\gamma(i, 2)}, \dots)_{i < i(2)}\text{”}$
iff $\Vdash_{\mathbb{P}_\alpha} \text{“if } \mathbf{B}_2(\dots, \eta_{\gamma(i, 2)}[\mathbf{G}], \dots)_{i < i(2)} \text{ is equal to 1 then so is } \mathbf{B}_1(\dots, \eta_{\gamma(i, 1)}, \dots)_{i < i(1)}\text{”}$.

2) For $\mathcal{U} \subseteq \alpha_*$ let $\mathbb{P}_{\mathcal{U}} = \mathbb{P}_{\mathcal{U}}^{\mathbf{q}}$ be the subforcing of $\mathbb{P}_{1, \alpha(\mathbf{q})}$ consist of $\{\mathbf{B}(\dots, \eta_{\gamma(i)}, \dots)_{i < i(*)} \in \mathbb{P}_{\alpha(\mathbf{q})} : i(*) \leq \lambda \text{ and } \gamma_i \in \mathcal{U} \text{ for every } i < i(*)\}$.

Claim 1.10. 1) For any sequence $\langle u_\alpha, \mathcal{P}_\alpha : \alpha < \alpha_* \rangle$ as above, i.e. as in clause (a) of Definition 1.8, there is one and only one $\mathbf{q} \in \mathbf{Q}_{\lambda, \bar{\theta}, \alpha_*}$ with $u_{\mathbf{q}, \alpha} = u_\alpha, \mathcal{P}_{\mathbf{q}, \alpha} = \mathcal{P}_\alpha$ for $\alpha < \alpha_*$.

1A) For $\alpha \leq \alpha_*$, the forcing notions $\mathbb{P}_{\mathbf{q}, \alpha}, \mathbb{P}_{\mathbf{q}, \mathcal{U}}$'s are well defined and are as demanded in Definition 1.9.

2) For every $\alpha \leq \alpha_*$ the set $\mathbb{P}_{\mathbf{q}, \alpha}^\bullet$ of $p \in \mathbb{P}_{\mathbf{q}, \alpha}$ satisfying the following is dense:

- (*) if $\beta \in \text{dom}(p)$, then $q = p(\beta)$ is a \mathbb{P}_β^\bullet -name of a member of \mathbb{Q}_β such that:
 - (a) $\eta_q, i_p, \langle u_{q, i} : i < i_p \rangle$ are objects (not just \mathbb{P}_α -names)
 - (b) $\underline{f}_q = \sup\{f_i : i < i_q\}$, each f_i is a \mathbb{P}_β' -name of a member of $\prod_{\varepsilon < \lambda} \theta_\varepsilon$

(c) each f_i has the form $\mathbf{B}_{q,i}(\dots, \eta_{\gamma(i,j)}, \dots)_{j < j(*) \leq \lambda}$ where $\{\gamma(i,j) : j < j(i)\} \subseteq u_{p,i}$ and \mathbf{B}_q is a Borel function from $\overset{\partial(*)}{\prod}_{\varepsilon < \lambda} \theta_\varepsilon$ into $\prod_{\varepsilon < \lambda} \theta_\varepsilon$

(d) $p(\beta) = (\eta_q, f_q)$.

3) Above for every $v \subseteq \alpha$ and $j_* < \lambda$ the set of $p \in \mathbb{P}_\alpha^\bullet$ such that $v \subseteq \text{dom}(p) \wedge (\forall \beta \in \text{dom}(p))(\ell g(\eta_{p(\beta)}) > j_*)$ is dense.

4) $\mathbb{P}_{\mathbf{q},0,\alpha} \triangleleft \mathbb{P}_{\mathbf{q},1,\alpha}$ moreover $\mathbb{P}_{\mathbf{q},0,\alpha}$ is dense in $\mathbb{P}_{\mathbf{q},1,\alpha}$ and $\mathcal{U}_1 \subseteq \mathcal{U}_2 \subseteq \alpha_{\mathbf{q}} \Rightarrow \mathbb{P}_{\mathcal{U}_1}^{\mathbf{q}} \triangleleft \mathbb{P}_{\mathcal{U}_2}^{\mathbf{q}} \triangleleft \mathbb{P}_{\mathbf{q},\alpha}$ so $\mathbb{P}_{\mathbf{q},\{\beta:\beta < \alpha\}} = \mathbb{P}_{\mathbf{q},2,\alpha}$ and $|\mathbb{P}_{\mathbf{q},\mathcal{U}}| \leq |\mathcal{U}|^\lambda$.

5) If $\alpha < \alpha_*$ and $u \in \mathcal{P}_\alpha$ then $\eta_\alpha \in \prod_{\varepsilon < \lambda} \theta_\varepsilon$ dominate every $\nu \in (\prod_{\varepsilon < \lambda} \theta_\varepsilon)^{\mathbf{V}[\eta \upharpoonright u]}$.

6) Assume $\mathbf{G} \subseteq \mathbb{P}_{\mathbf{q}}$ is generic over \mathbf{V} , $\eta_\alpha = \eta_\alpha[\mathbf{G}]$ and $\eta'_\alpha \in (\prod_{\varepsilon < \lambda} \theta_\varepsilon)^{\mathbf{V}[\mathbf{G}]}$ for $\alpha < \alpha_*$ and $\{(\alpha, \varepsilon) : \alpha < \alpha_*, \varepsilon < \alpha \text{ and } \eta_\alpha(\varepsilon) \neq \eta'_\alpha(\varepsilon)\}$ has cardinality $< \lambda$. Then for some (really unique) \mathbf{G}' we have $\mathbf{G}' \subseteq \mathbb{P}_{\mathbf{q}}$ is generic over \mathbf{V} and α .

7) Like (6) for $\mathbb{P}_{\mathcal{U}}^{\mathbf{q}}$

Proof. See [Sheb, 1.11=Lc8, 1.13=Lc11].

□_{1.10}

Theorem 1.11. For any ordinal α_* there is a quadruple $(\mathbf{q}, \delta_*, \mathcal{U}_*, h)$ such that:

- (A) (a) $\mathbf{q} \in \mathbf{Q}_{\lambda, \bar{\theta}}$ and let $\delta_* = \ell g(\mathbf{q})$
- (b) $\mathcal{U}_* \subseteq \delta_*$ has order type α_*
- (c) h is the order preserving function from α_* onto \mathcal{U}_*
- (d) if $\alpha \in \mathcal{U}_*$ then $\mathcal{U}_* \cap \alpha \in \mathcal{P}_{\mathbf{q},\alpha}$
- (B) if $\mathcal{U}_1 \subseteq \mathcal{U}_*, \mathcal{U}_2 \subseteq \mathcal{U}_*$, $\text{otp}(\mathcal{U}_1) = \text{otp}(\mathcal{U}_2)$ and g is the order preserving function from \mathcal{U}_1 onto \mathcal{U}_2 , then g induces an isomorphism \hat{g} from $\mathbb{P}_{\mathbf{q},\mathcal{U}_1}$ onto $\mathbb{P}_{\mathbf{q},\mathcal{U}_2}$ mapping η_β to $\eta_{g(\beta)}$ for $\beta \in \mathcal{U}_1$.

Proof. By [Sheb, 2.13=Lc51].

□_{1.11}

§ 2. THE FORCING

In this section we prove the main result of the paper, which reads as follows:

Theorem 2.1. *Assume*

- (a) λ is supercompact
- (b) $\lambda < \kappa = \text{cf}(\kappa) < \mu = \text{cf}(\mu) = \mu^\lambda$.

Then for some forcing notion \mathbb{P} not collapsing cardinals $\geq \lambda$, λ is still supercompact in $\mathbf{V}^{\mathbb{P}}$ and $\text{cov}_\lambda(\text{meagre}) = \kappa$, $\mathfrak{d}_\lambda = \mu$.

Proof. By Lemma 2.3(1) we force \square_λ while maintaining the supercompactness of λ . By Lemma 2.7 we force $\mathfrak{d}_\lambda = \mu \wedge \text{cov}_\lambda(\text{meagre}) = \kappa$ using a forcing notion \mathbb{P} which is λ^+ -c.c. and $(< \lambda)$ -strategically complete. Notice that λ is still supercompact in the generic extension, so we are done. $\square_{2.1}$

Definition 2.2. For λ supercompact we define \square_λ by:

- \square_λ for any regular cardinal $\chi > \lambda$ and forcing notion $\mathbb{P} \in \mathcal{H}(\chi)$ which is $(< \lambda)$ -strategically complete (see Definition 1.6(2)) the following set $\mathcal{S} = \mathcal{S}_{\mathbb{P}} = \mathcal{S}_{\chi, \mathbb{P}}$ is a stationary subset of $[\mathcal{H}(\chi)]^{< \lambda}$:
 - $\mathcal{S} = \mathcal{S}_{\mathbb{P}} = \mathcal{S}_{\chi, \mathbb{P}}$ is the set of N 's such that for some $\lambda_N, \chi_N, \mathbf{j} = \mathbf{j}_N, \mathbb{A} = \mathbb{A}_N, M = M_N, \mathbf{G} = \mathbf{G}_N$ we have (and we may say $(\lambda_N, \chi_N, \mathbf{j}_N, \mathbb{A}_N, M_N, \mathbf{G}_N)$ is a witness for $N \in \mathcal{S}_{\chi, \mathbb{P}}$ or for (N, \mathbb{P}, χ)) which means:
 - (a) $N \prec (\mathcal{H}(\chi))^{\mathbf{V}}, \in$ and $\mathbb{P} \in N$
 - (b) the Mostowski collapse of N is \mathbb{A} and let $\mathbf{j}_N : N \rightarrow \mathbb{A}$ be the unique isomorphism
 - (c) $N \cap \lambda = \lambda_N$ and $(\lambda_N)^{> N} \subseteq N$ and λ_N is strongly inaccessible
 - (d) $\mathbb{A} \subseteq M := (\mathcal{H}(\chi_N), \in)$, M is transitive as well as \mathbb{A}
 - (e) $\mathbf{G} \subseteq \mathbf{j}_N(\mathbb{P})$ is generic over \mathbb{A} for the forcing notion $\mathbf{j}_N(\mathbb{P})$
 - (f) $M = \mathbb{A}[\mathbf{G}]$.

Our first lemma is closed to Laver's indestructibility. It consists of two parts. In the first part we prove that one can force \square_λ at a supercompact cardinal λ while preserving its supercompactness. In the second part, we prove that this can be done in an indestructible manner. Namely, any further extension of the universe by a $(< \lambda)$ -directed-closed forcing notion will preserve the principle \square_λ .

Lemma 2.3. 1) *If λ is supercompact then after some preliminary forcing of cardinality λ , λ is still supercompact and \square_λ from 2.2.*

2) *The statement \square_λ holds in $\mathbf{V}^{\mathbb{P}}$ when \mathbf{V} satisfies \square_λ and \mathbb{P} is a $(< \lambda)$ -strategically complete forcing notion and \mathbb{P} is $(< \lambda)$ -directed closed, but see 2.4(2).*

Remark 2.4. 1) Recall “ \mathbb{P} is a $(< \lambda)$ -directed closed” means:

- (*) if J is a directed partial order of cardinality $< \lambda$ and $p_s \in \mathbb{P}$ for $s \in J$ and $s \leq_J t \Rightarrow p_s \leq_{\mathbb{P}} p_t$ then the set $\{p_s : s \in J\}$ has an upper bound in \mathbb{P} .

2) In 2.3(2) we can weaken the assumption “ \square_λ and \mathbb{P} is $(< \lambda)$ -directed closed” to:

⊞ if $\chi > \lambda$ and $\mathbb{P} \in \mathcal{H}(\chi)$ then we have (A) \Rightarrow (B) where:

- (A) $N \prec (\mathcal{H}(\chi), \in)$ and $\lambda_N, \chi_N, \mathbf{j}_N, \mathbf{G}_N, \mathbb{Q}$ satisfies:
- (a) $\mathbb{P}, \mathbb{Q} \in N, N \cap \lambda = \lambda_N < \chi_N < \lambda, [N]^{<\lambda_N} \subseteq N,$
 - (b) \mathbb{Q} is a \mathbb{P} -name of $(< \lambda)$ -strategically closed forcing notion
 - (c) \mathbf{j} is the Mostowski Collapse of N , its range is \mathbb{A}
 - (d) \mathbf{G} is a subset of $\mathbf{j}(\mathbb{P} * \mathbb{Q})$, generic over \mathbb{A}
 - (e) $\mathbb{A}[\mathbf{G}] = \mathcal{H}(\chi_N)$
- (B) $\{p \in \mathbb{P} \cap N: \text{for some } (p', q') \in \mathbf{G} \text{ we have } \mathbf{j}(p) = p'\}$ has a common upper bound in \mathbb{P} .

3) We can e.g. restrict the χ to be strong limit.

Proof. 1) This is similar to the proof in Laver [Lav78] using Laver's diamond, see Definition 1.1, but as requested we elaborate. By Laver [Lav78] without loss of generality there is a Laver diamond $h : \lambda \rightarrow \mathcal{H}(\lambda)$. Let $E = \{\theta : \theta < \lambda \text{ a strong limit cardinal and } \alpha < \theta \Rightarrow h(\alpha) \in \mathcal{H}(\theta)\}$, clearly a club of λ and let $\langle \kappa_\varepsilon : \varepsilon < \lambda \rangle$ list $\{\theta \in E : \theta \text{ is strongly inaccessible}\}$ in increasing order.

As requested, we now define \mathbf{q}_ε and $\bar{\chi}^\varepsilon$ by induction on $\varepsilon \leq \lambda$ such that:

- (*) (a) $\mathbf{q}_\varepsilon = \langle \mathbb{P}_\zeta, \mathbb{Q}_\xi : \zeta \leq \varepsilon, \xi < \varepsilon \rangle$ is an Easton support iteration (so $\mathbb{P}_\zeta, \mathbb{Q}_\xi$ do not depend on ε)
- (b) $\mathbb{P}_\zeta \subseteq \mathcal{H}(\kappa_\zeta)$
- (c) $\bar{\chi}^\varepsilon = \langle \chi_\zeta : \zeta < \varepsilon \rangle$ where each χ_ζ is a regular cardinal $\in [\kappa_\zeta, \kappa_{\zeta+1})$
- (d) $\mathbb{Q}_\xi \in \mathcal{H}(\chi_{\xi+1})$ is a \mathbb{P}_ξ -name of a $(< \kappa_\xi)$ -strategically complete forcing notion
- (e) if $h(\xi) = (\mathbb{Q}, \chi)$ and the pair (\mathbb{Q}, χ) satisfies the requirements on $(\mathbb{Q}_\xi, \chi_\xi)$ in clauses (c),(d) then $(\mathbb{Q}_\xi, \chi_\xi) = h(\xi)$.

Concerning clause (b) which says “ $\mathbb{P}_\zeta \subseteq \mathcal{H}(\kappa_\zeta)$ ”, note that for ζ is a limit ordinal letting $\kappa_{<\zeta} = \bigcup \{\kappa_\xi : \xi < \zeta\}$ we have $\kappa_{<\zeta}$ is strong limit and:

- if $\kappa_{<\zeta}$ is regular, equivalently strongly inaccessible then $\kappa_{<\zeta} = \kappa_\zeta$ and $\mathbb{P}_\zeta = \bigcup \{\mathbb{P}_\xi : \xi < \zeta\}$ and so $\mathbb{P}_\zeta \subseteq \bigcup \{\mathcal{H}(\kappa_\xi) : \xi < \zeta\} = \mathcal{H}(\kappa_{<\zeta}) = \mathcal{H}(\kappa_\zeta)$
- if $\kappa_{<\zeta}$ is singular, then $\mathbb{P}_\zeta \subseteq \mathcal{H}(\kappa_{<\zeta}^+) \subseteq \mathcal{H}(\kappa_\zeta)$ as κ_ζ is inaccessible $> \kappa_{<\zeta}$.

Easily we can carry the induction so \mathbf{q}_λ is well defined, $\mathbb{P}_\lambda = \bigcup \{\mathbb{P}_\varepsilon : \varepsilon < \lambda\} \subseteq \bigcup \{\mathcal{H}(\kappa_\varepsilon) : \varepsilon < \lambda\} = \mathcal{H}(\lambda)$ and “ $\xi < \lambda \Rightarrow \mathbb{P}_\lambda / \mathbb{P}_\xi$ is $(< \kappa_\xi)$ -strategically complete” hence $\mathbb{P}_\lambda / \mathbb{P}_\xi$ adds no new sequence of length $< \kappa_\xi$ of ordinals. Clearly it is enough to prove that in $\mathbf{V}^{\mathbb{P}_\lambda}$ we have \square_λ .

Toward contradiction assume $\chi, \mathbb{P}, \mathcal{S} = \mathcal{S}_{\chi, \mathbb{P}}$ form a counter-example in $\mathbf{V}^{\mathbb{P}_\lambda}$, hence there are $p_* \in \mathbb{P}_\lambda$ and \mathbb{P}_λ -names $\underline{\chi}, \underline{\mathbb{P}}, \underline{\mathcal{S}}, \underline{E}$ such that $p_* \Vdash_{\mathbb{P}_\lambda}$ “ $\chi > \lambda$ is regular, $\mathbb{P} \in \mathcal{H}(\chi)$ and $\mathcal{S}_{\chi, \mathbb{P}}$ is defined as in \square_λ and $\underline{E} \subseteq [\mathcal{H}(\chi)^{\mathbf{V}^{\mathbb{P}_\lambda}}]^{<\lambda}$ is a club disjoint to $\underline{\mathcal{S}}$ ”.

As we can increase p_* , without loss of generality $\underline{\chi} = \chi$ and let $x = (\chi, \underline{\mathbb{P}})$; and as $\mathbf{V} \models$ “ λ is supercompact and h is a Laver diamond” for some $(I, D, \mathbf{M}, \mathbf{j}, \mathbf{j}_0, \mathbf{j}_1)$ we have:

- (*)₁ (a) \mathbf{M} is a transitive class
- (b) \mathbf{M} is a model of ZFC

- (c) ${}^x\mathbf{M} \subseteq \mathbf{M}$
- (d) \mathbf{j} is an elementary embedding from \mathbf{V} into \mathbf{M}
- (e) $\text{crit}(\mathbf{j}) = \lambda$
- (f) $\mathbf{j}(h)(\lambda) = (\chi, \mathbb{P})$
- (g) $I = [\mathcal{H}(\chi_1)]^{<\lambda}$ and $\chi_1 > \chi$
- (h) D is a fine normal ultrafilter on I
- (i) \mathbf{j}_0 is the canonical elementary embedding of \mathbf{V} into \mathbf{V}^I/D
- (j) \mathbf{M} is the Mostowski Collapse of \mathbf{V}^I/D
- (k) \mathbf{j}_1 is the canonical isomorphism from \mathbf{V}^I/D onto \mathbf{M}
- (l) $\mathbf{j} = \mathbf{j}_1 \circ \mathbf{j}_0$.

Moreover, by Definition 1.1

$$(*)_2 \quad x = \mathbf{j}_1(\langle f(\sup(u \cap \lambda) : u \in I) / D \rangle).$$

Let $\mathbf{q} = \mathbf{j}(\mathbf{q}_\lambda)$ so $\mathbf{q} = \langle \mathbb{P}_\zeta, \mathbb{Q}_\xi : \zeta \leq \mathbf{j}(\lambda), \xi < \mathbf{j}(\lambda) \rangle$ and $\zeta < \lambda \Rightarrow \mathbb{P}_\zeta^{\mathbf{q}} = \mathbb{P}_\zeta$, etc.

So

- (*)₃ in \mathbf{M} the pair $x = (\chi, \mathbb{P})$ satisfies:
 - (a) $\chi \in (\lambda, \mathbf{j}(\lambda))$, $\mathbf{j}(\lambda)$ is inaccessible
 - (b) $\mathbb{P} \in \mathcal{H}(\chi)$
 - (c) \mathbb{P} is a \mathbb{P}_λ -name of a $(< \lambda)$ -strategically complete forcing notion.

[Why? Because $[\mathbf{M}]^x \subseteq \mathbf{M}$ hence $\mathcal{H}(\chi^+)^{\mathbf{V}} \subseteq \mathbf{M}$.]

Now

- (*)₄ the following sets belong to D :
 - (a) $\mathcal{S}_1 = \{u \in I : x \in u \text{ and } (\mathcal{H}(\chi_1), \epsilon) \upharpoonright u \prec (\mathcal{H}(\chi_1), \epsilon)\}$
 - (b) $\mathcal{S}_2 = \{u \in \mathcal{S}_1 : u \cap \lambda \text{ is an inaccessible cardinal we call } \lambda_u\}$
 - (c) $\mathcal{S}_3 = \{u \in \mathcal{S}_2 : \text{the Mostowski Collapse } N_u^1 \text{ of } (\mathcal{H}(\chi_1), \epsilon) \upharpoonright u \text{ is isomorphic to some } (\mathcal{H}(\chi_u^1), \epsilon)\}$.

[Why? As D is a fine and normal ultrafilter on I .]

- (*)₅ for every formula $\varphi = \varphi(-) \in \mathbb{L}(\{\in\})$ the following are equivalent:
 - (a) $(\mathcal{H}(\chi_1), \theta) \models \varphi[x]$
 - (b) $(\mathcal{H}(\chi_1), \epsilon)^I / D \models \varphi[\langle h(u \cap \lambda) : u \in I \rangle / D]$
 - (c) $\mathcal{X}_\varphi^1 \in D$ where $\mathcal{X}_\varphi^1 = \{u \in I : x \in u \text{ and } (\mathcal{H}(\chi_1), \epsilon) \upharpoonright u \models \varphi[x]\}$
 - (d) $\mathcal{X}_\varphi^2 \in D$ where $\mathcal{X}_\varphi^2 = \{u \in I : p_*, x \in u \text{ and } (\mathcal{H}(\chi_u), \epsilon) \models \varphi[\mathbf{j}_u(x)]\}$
 - (e) $\mathcal{X}_\varphi^3 \in D$ where $\mathcal{X}_\varphi^3 = \{u \in I : x \in u, \chi_u^1 = \text{otp}(\chi_u \upharpoonright u) \text{ and } (\mathcal{H}(\chi_u^1), \epsilon) \models \varphi[\mathbf{j}_u(x)]\}$.

[Why? We have (a) \Leftrightarrow (c) as D is a fine normal ultrafilter on $I = \mathcal{H}(\chi_1)$; we have (c) \Leftrightarrow (d) as j_u is an isomorphism from $(\mathcal{H}(\chi_1), \epsilon) \upharpoonright u$ onto $\mathcal{H}(\chi_u^1)$; we have (d) \Leftrightarrow (e) by the choice of D ; lastly, (b) \Leftrightarrow (c) by Los theorem.]

Hence

- (*)₆ there is N as required in $\mathbf{V}^{\mathbb{P}^*}$.

[Why? Choose $u \in I$ which belongs to all the sets from D mentioned in $(*)_4 + (*)_5$. Let $\zeta = u \cap \lambda$, so it is inaccessible, even measurable, and $\mathbf{j}_u(x) = \mathbf{j}_u(\chi, \mathbb{P}) = h(\zeta)$ so (by the choice of \mathbf{q}) $h(\zeta) = (\chi, \mathbb{Q}_\zeta)$ and \mathbb{Q}_ζ is a $\mathbb{P}_{\mathbf{q}, \zeta}$ -name.

Let \mathbf{G} be a subset of $\mathbb{P}_{\mathbf{q}} = \mathbb{P}_\lambda$ to which p_* belongs, $\mathbf{G}_\zeta = \mathbf{G} \cap \mathbb{P}_{\mathbf{q}, \zeta}$, hence is a generic subset of $\mathbb{P}_{\mathbf{q}, \zeta}$ over \mathbf{V} hence a generic subset of $\mathbf{j}_u(\mathbb{P}_{\mathbf{q}}) \in \mathcal{H}(\chi_\zeta)$ and let $N = ((\mathcal{H}(\chi_1), \epsilon) \upharpoonright u)[\mathbf{G}], \mathbb{A} = (\mathcal{H}(\chi_\zeta)^{\mathbf{V}[\mathbf{G}_\zeta]}, \epsilon), M = \mathbb{A}^{\mathbb{Q}_\zeta[\mathbf{G}_\zeta]}$. Easily N is as promised, contradiction to the choice of p_* .]

So we are done proving part (1).

2) Let \mathbb{Q} be a forcing notion in $\mathbf{V}^{\mathbb{P}}$ which is $(< \lambda)$ -strategically complete and (even) $(< \lambda)$ -directed closed, $\emptyset \in \mathbb{Q}$ is the weakest condition, χ_1 large enough so that $\lambda, \mathbb{Q}, \underline{E} \in \mathcal{H}(\chi_1)$ and it suffices to prove that in $\mathbf{V}^{\mathbb{P}}$, the set $\mathcal{S}_{\chi_1, \mathbb{Q}}$ is stationary. So let $\mathbb{Q}, \underline{E}$ be \mathbb{P} -names such that for some $p \in \mathbb{P}$ we have $p \Vdash_{\mathbb{P}} \text{“}\mathbb{Q} \in \mathcal{H}(\chi_1)\text{”}$ is $(< \lambda)$ -strategically complete, $(< \lambda)$ -directed closed, forcing notion, \underline{E} a club of $[\mathcal{H}(\chi_1)]^{< \lambda}$ disjoint to $\mathcal{S}_{\chi_1, \mathbb{Q}}$, no need to use a name for χ_1 as we can increase p .

Let $\chi \gg \chi_1$; now $\mathbb{P} * \mathbb{Q} \in \mathcal{H}(\chi)$ is a $(< \lambda)$ -strategically complete forcing notion and without loss of generality code (χ_1, E) . As \square_λ holds in \mathbf{V} we can apply it to the forcing $\mathbb{P}_{\geq p} * \mathbb{Q}$ so we can find a tuple $(N, \lambda_N, \chi_N, \mathbf{j}_N, \mathbb{A}_N, M_N, \mathbf{G}_N)$ witnessing it, in particular, $(p, \emptyset) \in \mathbf{G}_N, \mathbb{P} * \mathbb{Q} \in N$ so $\chi_1, \underline{E} \in N$. Let $\mathbf{G}_{\mathbb{P}}$ be a subset of \mathbb{P} generic over \mathbf{V} which extends $\{p' : (p', q') \in \mathbf{G}_N\}$, possible because \mathbf{G}_N is in \mathbf{V} , a subset of \mathbb{P} which has an upper bound, this is the only place we use “ \mathbb{P} is $(< \lambda)$ -directed closed”.

Next, let $\mathbf{V}_1 = \mathbf{V}[\mathbf{G}_{\mathbb{P}}], N_1 = N[\mathbf{G}_{\mathbb{P}}], E_1 = E[\mathbf{G}_{\mathbb{P}}], \mathbb{A}_1 = \mathbb{A}[\mathbf{j}'_N(\mathbf{G}_{\mathbb{P}} \cap N)] = \mathbb{A}[\{p' : (p', q') \in \mathbf{G}_N\}], \mathbf{G}_1 = \{q[\mathbf{j}''(\mathbf{G}_{\mathbb{P}} \cap N)] : (p, q) \in \mathbf{G}_{\mathbb{P}}\}$.

Let $N_2 = N_1 \upharpoonright \mathcal{H}(\chi_1)^{\mathbf{V}[\mathbf{G}_{\mathbb{P}}]}, \mathcal{S} = \mathcal{S}_{\mathbb{Q}}[\mathbf{G}_{\mathbb{P}}], \mathbf{j}_1$ is the lifting of $(\mathbf{j} \upharpoonright (N \cap \mathcal{H}(\chi)))$, to mapping N_1 onto \mathbb{A}_1 .

Now recalling p forces \underline{E} is disjoint to \mathcal{S} clearly

$$(*) \ N_2 \in E.$$

hence

$$(*) \ N_1 \notin \mathcal{S}_1.$$

But easily in \mathbf{V}_1 we have: $(\lambda_N, \chi_N, \mathbf{j}_1, \mathbb{A}_1, M_1 = M, \mathbf{G}_1)$ witness $N_1 \in \mathcal{S} \cap E_1$, a contradiction to the choice of \underline{E} . $\square_{2.3}$

Discussion 2.5. Suppose that one wishes to force an inequality between two cardinal characteristics. There are two general approaches, which can be labeled as Top-down and Bottom-up In the Bottom-up strategy one begins with a universe in which many characteristics are small, e.g. by assuming $2^\lambda = \lambda^+$, and then increases some of them while trying to keep the smallness of the rest. In the Top-down strategy one begins with a universe in which many characteristics are large. The forcing aims to decrease some of them while keeping the large value of the rest.

We shall use the Top-down approach, so we begin by increasing \mathfrak{b}_λ (and \mathfrak{d}_λ) to some $\mu = \text{cf}(\mu) > \lambda$. Notice that \mathfrak{b}_λ is a relatively small characteristics and, in particular, always $\mathfrak{b}_\lambda \leq \mathfrak{d}_\lambda$. The next step will be to decrease $\text{cov}_\lambda(\text{meagre})$ in such a way that maintains the fact that $\mathfrak{d}_\lambda = \mu$. We shall increase \mathfrak{b}_λ by using the generalization to λ of Hechler forcing. This is a standard way to achieve this goal, but we spell out the proof since it demonstrates the way that we employ Lemma 2.3.

Claim 2.6. *Assume that:*

- (a) λ is supercompact
- (b) $\lambda < \mu = \text{cf}(\mu) = \mu^\lambda$.

Then one can force $\mathfrak{b}_\lambda = \mathfrak{d}_\lambda = \mu$ while keeping the supercompactness of λ and the principle \square_λ .

Proof. Begin with the preparatory forcing of Lemma 2.3 to make λ indestructible and to force \square_λ in such a way that it will be preserved by any further $(< \lambda)$ -directed-closed forcing. By 2.3 as in the applications of Laver-indestructibility we can assume that GCH holds above λ after the preparatory forcing. In particular, if $\mu = \text{cf}(\mu) > \lambda$ then $\mu^\lambda = \mu$ follows.

Let \mathbb{D}_λ be the generalized Hechler forcing. A condition $p \in \mathbb{D}_\lambda$ is a pair (η_p, f_p) such that $\eta_p \in {}^{<\lambda}\lambda$, $f_p \in {}^\lambda\lambda$ and $\eta_p \leq f_p$. If $p, q \in \mathbb{D}_\lambda$ then $p \leq q$ iff $\eta_p \leq \eta_q$ and $f_p(\alpha) \leq f_q(\alpha)$ for every $\alpha \in \lambda$.

Let $\mathbf{q} = \langle \mathbb{P}_\alpha, \mathbb{Q}_\beta : \alpha \leq \mu, \beta < \mu \rangle$ be a $(< \lambda)$ -support iteration of the generalized Hechler forcing notions for λ . Explicitly, \mathbb{Q}_α is the \mathbb{P}_α -name of \mathbb{D}_λ in $\mathbf{V}^{\mathbb{P}_\alpha}$ for every $\alpha < \mu$. Denote the generic λ -Hechler for \mathbb{Q}_α by f_α^* . So \mathbb{P}_μ is the limit and choose a generic $\mathbf{G} \subseteq \mathbb{P}_\mu$. We claim that $\mathbf{V}[\mathbf{G}] \models \text{“}\mathfrak{b}_\lambda = \mathfrak{d}_\lambda = \mu\text{”}$ as witnessed by $\langle f_\alpha^* : \alpha < \mu \rangle$. Notice that $2^\lambda = \mu$ in $\mathbf{V}[\mathbf{G}]$, so it is sufficient to prove that $\mathfrak{b}_\lambda = \mu$ in $\mathbf{V}[\mathbf{G}]$.

Since λ is regular, each \mathbb{Q}_α is $(< \lambda)$ -complete. By Fact 1.7, \mathbb{P}_α is $(< \lambda)$ -complete as well, for every $\alpha \leq \mu$. Likewise, each \mathbb{Q}_α is λ -centered so \mathbb{P}_μ is λ^+ -c.c. (see [She78] or [Shed]). It follows that $\mathbf{V}[\mathbf{G}]$ preserves cardinals and cofinalities. Moreover, no new $(< \lambda)$ -sequences are introduced. Notice also that \mathbb{P}_μ is $(< \lambda)$ -directed-closed and hence $\mathbf{V}[\mathbf{G}] \models \text{“}\lambda \text{ is supercompact and } \square_\lambda \text{ holds”}$.

The main point is that $\{f_\alpha^* : \alpha < \mu\}$ is a cofinal family in $({}^\lambda\lambda)^{\mathbf{V}[\mathbf{G}]}$. For this, assume that $\Vdash_{\mathbb{P}_\mu} \text{“}f \in {}^\lambda\lambda\text{”}$. For every $\alpha < \lambda$ fix a maximal antichain $\langle p_{\alpha,i} : i < i_\alpha \leq \lambda \rangle$ of conditions which force a value to $f(\alpha)$. Let $\delta = \sup(\cup \{\text{dom}(f_{\alpha,i}) : \alpha < \lambda, i < i_\alpha\})$. Since $\lambda < \mu = \text{cf}(\mu)$ we see that $\delta < \mu$, and clearly f is a \mathbb{P}_δ -name. We conclude, therefore, that f is dominated by $f_{\delta+1}^*$ and hence $\{f_\alpha^* : \alpha < \mu\}$ exemplifies $\mathfrak{b}_\lambda = \mu$. This fact completes the proof. $\square_{2.6}$

Our second lemma is the main burden of the proof. The statement of the theorem requires λ to be supercompact, in order to obtain the indestructibility properties given by Lemma 2.3. The combinatorial part given in Lemma 2.7 below requires only strong inaccessibility. However, we assume supercompactness order to keep \square_λ .

Lemma 2.7. *Assume that:*

- (a) λ is supercompact
- (b) \square_λ holds
- (c) $\lambda < \kappa = \text{cf}(\kappa) < \mu = \text{cf}(\mu) = \mu^\lambda$.

Then there exists a λ^+ -c.c. $(< \lambda)$ -strategically complete forcing notion \mathbb{P} such that $\Vdash_{\mathbb{P}} \text{“}\mathfrak{d}_\lambda = \mu \wedge \text{cov}_\lambda(\text{meagre}) = \kappa\text{”}$.

Proof. By claim 2.6 without loss of generality $\mathfrak{b}_\lambda = \mathfrak{d}_\lambda = \mu$. In particular, λ is supercompact and \square_λ holds in the generic extension. Let $\langle f_\alpha^* : \alpha < \mu \rangle$ witness $\mathfrak{b}_\lambda = \mathfrak{d}_\lambda = \mu$ and without loss of generality $\alpha < \beta < \mu \Rightarrow f_\alpha^* <_{J_\lambda^{\text{bd}}} f_\beta^*$.

Recalling Definition 1.8, 1.9, Claim 1.10, Theorem 1.11, in \mathbf{V} there are $\beta(*), \mathbf{q}, \bar{u}, \mathcal{U}_*, \dots$ such that:

- (*)₁(A) $\mathbf{q} \in \mathbf{Q}_{\lambda, \bar{\theta}, \beta(*)}$ so in particular we have (in \mathbf{q}):
- (a) $\langle \mathbb{P}_{0, \alpha}, \mathbb{Q}_{0, \beta} : \alpha \leq \beta(*), \beta < \beta(*) \rangle$ is a $(< \lambda)$ -support iteration
 - (b) $\bar{u} = \langle u_\beta : \beta < \beta(*) \rangle, \mathcal{P} = \langle \mathcal{P}_\beta : \beta < \beta(*) \rangle$
 - (c) $u_\beta \subseteq \beta, \mathcal{P}_\beta \subseteq [u_\beta]^{\leq \lambda}$ is closed under subsets
 - (d) $\mathbb{Q}_{0, \beta}$ has generic $\eta_\beta \in \prod_{\varepsilon < \lambda} \theta_\varepsilon$
 - (e) $\mathbb{Q}_{0, \beta}$ is as in 1.8(e) so is $\subseteq \mathbb{Q}_{\bar{\theta}}^{\mathbf{V}[\langle \eta_\alpha : \alpha \in u_\beta \rangle]}$ and $\Vdash_{\mathbb{P}_{\beta+1}}$ “ $\eta_\beta \in \prod_{\varepsilon < \lambda} \theta_\varepsilon$ ” and $\bar{\eta} = \langle \eta_\beta : \beta < \beta(*) \rangle$
 - (f) $\mathcal{U}_* \subseteq \beta(*)$ has order type $\gamma(*) = \kappa$ and $\langle \beta_i^* : i \leq \kappa \rangle$ lists $\mathcal{U}_* \cup \{\beta(*)\}$ in increasing order
 - (g) if $\beta \in \mathcal{U}_*$ then $[\mathcal{U}_* \cap \beta]^{\leq \lambda} \subseteq \mathcal{P}_\beta$ hence $\subseteq u_\beta$ and $\Vdash_{\mathbb{P}_{0, \beta+1}}$ “if $\nu \in \mathbf{V}[\langle \eta_\alpha : \alpha \in \mathcal{U}_* \cap \beta \rangle] \cap \prod_{\varepsilon < \lambda} \theta_\varepsilon$ then $\nu <_{J_\lambda^{\text{bd}}} \eta_\beta$ ”
 - (h) if $\alpha \leq \beta(*)$ then $\mathbb{P}_{0, \alpha}$ is $(< \lambda)$ -strategically complete and λ^+ -c.c.
 - (i) $\mathbb{P}_{1, \alpha}, \mathbb{P}_{1, \mathcal{U}}$ are as in 1.9
- (B) letting $\mathbb{P}'_i = \mathbb{P}_{\mathbf{q}, 1, \{\beta_j^* : j < i\}}$ for $i \leq \kappa$ we have
- (a) The sequence $\langle \mathbb{P}'_i : i \leq \gamma(*) \rangle$ of forcing notions is \triangleleft -increasing, and is continuous for ordinals $i \leq \gamma(*)$ of cofinality $\geq \lambda$, see [Sheb, 2.5(8)=Lz48(8)], but the continuity will not be used
 - (b) \mathbb{P}'_i is $(< \lambda)$ -strategically complete for $i \leq \gamma(*)$
 - (c) $(\prod_{\varepsilon < \lambda} \theta_\varepsilon)^{\mathbf{V}[\mathbb{P}'_{\gamma(*)}]} = \cup \{ (\prod_{\varepsilon < \lambda} \theta_\varepsilon)^{\mathbf{V}[\mathbb{P}'_i]} : i < \gamma(*) \}$.
 - (d) The sequence $\langle \mathbb{P}_{1, \beta} : \beta \leq \beta(*) \rangle$ is a sequence of forcing notions, \triangleleft -increasing and if $\beta \leq \beta(*)$ then $\mathbb{P}_{0, \beta} \triangleleft \mathbb{P}_{1, \beta}$, in fact is dense in it and if $i \leq \gamma(*)$ then $\mathbb{P}'_i \triangleleft \mathbb{P}_{1, \beta_i^*}$.

We shall mention more properties later.

[Why are there such objects? We apply 1.11 and 1.8 and 1.10, that is [Sheb].]

Also

- (*)₂
- (a) recall $\langle \beta_i^* : i \leq \gamma(*) \rangle$ lists $\mathcal{U}_* \cup \{\beta(*)\}$ in increasing order
 - (b) for $i \leq \gamma(*) = \kappa$, for $i < \gamma(*)$ let \bar{g}'_i be $\eta_{\beta_i^*}$ (to avoid excessive subscripts),
 - (c) let $\bar{g}' = \langle \bar{g}'_i : i < \kappa \rangle$
 - (d) let $\bar{g}_\alpha = \eta_\alpha$ for $\alpha < \beta(*)$ and $\bar{g} = \langle \bar{g}_\beta : \beta < \beta(*) \rangle$
 - (e) $\mathcal{P}_\alpha = \mathcal{P}_{\mathbf{q}, \alpha}$ and without loss of generality $u_\alpha = \cup \{u : u \in \mathcal{P}_\alpha\}$ for $\alpha < \beta(*)$.
- (*)₃ if $u \in \mathcal{P}_\alpha, \alpha < \beta(*)$ then $\Vdash_{\mathbb{P}_{0, \alpha+1}}$ “ $\bar{g}_\alpha \in \prod_{\varepsilon < \lambda} \theta_\varepsilon$ dominates $(\prod_{\varepsilon < \lambda} \theta_\varepsilon)^{\mathbf{V}[\langle \bar{g}_\beta : \beta \in u \rangle]}$ ”, the order being modulo J_λ^{bd} .

[Why? By the choice of the forcing, see 1.4 or $(*)_1(A)(g)$ above.]

$(*)_4 \Vdash_{\mathbb{P}'_\kappa} \text{“}\underline{g}' = \langle g'_i : i < \kappa \rangle \text{ is } <_{J_\lambda^{\text{bd}}}\text{-increasing and cofinal in } (\prod_{\varepsilon < \lambda} \theta_\varepsilon, <_{J_\lambda^{\text{bd}}})\text{”}.$

[Why? By $(*)_3$ noting that $(\prod_{\varepsilon < \lambda} \theta_\varepsilon)^{\mathbf{V}[\mathbb{P}'_\kappa]} = \cup \{(\prod_{\varepsilon < \lambda} \theta_\varepsilon)^{\mathbf{V}[\mathbb{P}'_i]} : i < \kappa\}$ which holds by $(*)_1(B)(c)$.] Now

$(*)_5 \Vdash_{\mathbb{P}'_\kappa} \text{“}\text{cov}_\lambda(\text{meagre}) \leq \kappa\text{”}.$

[Why? First, notice that we can look at $\prod_{\varepsilon < \lambda} \theta_\varepsilon$ instead of ${}^\lambda 2$.

Second, for each $\varepsilon < \lambda, i < \kappa$ the set $B_{\varepsilon, i} = \{\eta \in \prod_{\xi < \lambda} \theta_\xi : \text{for every } \zeta \in [\varepsilon, \lambda) \text{ we have } \eta(\zeta) \leq g'_i(\zeta) < \theta_\zeta\}$ is closed nowhere dense, and by $(*)_4$ we have $\mathbf{V}^{\mathbb{P}'_\kappa} \models \text{“}\prod_{\zeta < \lambda} \theta_\zeta = \cup \{B_{\varepsilon, i} : \varepsilon < \lambda, i < \kappa\}\text{”}.$ In fact, $\langle B_{0, i} : i < \kappa \rangle$ suffice.]

Alternatively we have $\langle g'_i : i < \kappa \rangle$ is $<_{J_\lambda^{\text{bd}}}$ -increasing cofinal in $\prod_{\varepsilon < \lambda} \theta_\varepsilon$ and let $\mathcal{W}_{i, \zeta} := \{\eta : \eta \in {}^\lambda 2 \text{ and for every } \varepsilon \in [\zeta, \lambda) \text{ we have either } \eta \upharpoonright [\Sigma_{\xi < \varepsilon} \theta_\xi, \Sigma_{\xi \leq \varepsilon} \theta_\xi) \text{ is constantly zero or } \min\{\alpha : \Sigma_{\xi < \varepsilon} \theta_\xi + \alpha \in \eta^{-1}(\{1\})\} < g'_i(\varepsilon)\}$. So $\mathcal{W}_{i, \zeta}$ is a closed nowhere dense subset of ${}^\lambda 2$ and $\cup \{\mathcal{W}_{i, \zeta} : i < \kappa, \zeta < \lambda\} = {}^\lambda 2$ and $\kappa \times \lambda$ has cardinality $\lambda + \kappa = \kappa$].

$(*)_6 \Vdash_{\mathbb{P}'_\kappa} \text{“}\text{cov}_\lambda(\text{meagre}) \geq \kappa\text{”}.$

[Why? Let us define the \mathbb{P}'_{i+1} -name η'_i of a member of ${}^\lambda 2$ by $\eta'_i(\varepsilon) = 0$ iff $g'_i(\varepsilon)$ is even. Now clearly $\Vdash_{\mathbb{P}'_{i+1}} \text{“}\eta'_i \text{ is a } \lambda\text{-Cohen sequence over } \mathbf{V}^{\mathbb{P}'_i}\text{”}.$ (But let us elaborate; η'_i is also a $\mathbb{P}^{\beta^*_{i+1}}$ -name and $\Vdash_{\mathbb{P}^{\beta^*_{i+1}}} \text{“}\eta'_i \text{ is } \lambda\text{-Cohen over } \mathbf{V}^{\mathbb{P}^{\beta^*_i}} \text{ hence over } \mathbf{V}^{\mathbb{P}'_i}\text{”};$ the last hence because $\mathbb{P}'_i \triangleleft \mathbb{P}^{\beta^*_i}$. As $\mathbb{P}^{\beta^*_{i+1}} \triangleleft \mathbb{P}^{\beta^*_{i+1}}$ and $\mathbb{P}'_{i+1} \triangleleft \mathbb{P}^{\beta^*_{i+1}}$ we are done.)

Also every closed nowhere dense subset of ${}^\lambda 2$ from $\mathbf{V}^{\mathbb{P}'_{\gamma(*)}}$ is from $\mathbf{V}^{\mathbb{P}'_i}$ for some $i < \gamma(*)$. So if $p \Vdash \text{“}\text{cov}_\lambda(\text{meagre}) < \kappa\text{”}$ then for some $\zeta < \kappa$ and $\underline{A}_\varepsilon(\varepsilon < \zeta)$ we have $p \Vdash \text{“}\underline{A}_\varepsilon \text{ is a closed no-where dense set subset of } {}^\lambda 2 \text{ for } \varepsilon < \zeta\text{”}$ and $p \Vdash \text{“}\bigcup_i \underline{A}_i \text{ is equal to the set of } {}^\lambda 2\text{”}.$ Without loss of generality each $\underline{A}_\varepsilon$ is a $\mathbb{P}_{i(\varepsilon)}$ -name, $i(\varepsilon) < \kappa$ and recall that κ is regular. Hence $i = \sup\{i(\varepsilon) : \varepsilon < \zeta\} < \kappa$ and η'_i gives a contradiction to the choice of $\langle \underline{A}_\varepsilon : \varepsilon < \zeta \rangle$; so $(*)_6$ holds indeed.]

The reader may look at some explanation in 2.9.

Now we come to the main and last point recalling $\langle f_\alpha^* : \alpha < \mu \rangle$ from Claim 2.6

$(*)_7 \Vdash_{\mathbb{P}'_{\gamma(*)}} \text{“no } \underline{f} \in ({}^\lambda \lambda) \text{ dominates } \{f_\alpha^* : \alpha < \mu\}\text{”}.$

We shall show that it suffices to prove $(*)_7$ for proving the Lemma 2.3(2), and that $(*)_7$ holds, thus finishing.

Why it suffices? As $\langle f_\alpha^* : \alpha < \mu \rangle$ is $<_{J_\lambda^{\text{bd}}}$ -increasing and $\text{cf}(\mu) = \mu > \lambda$, this implies $\Vdash_{\mathbb{P}'_\kappa} \text{“}\mathfrak{d}_\lambda \geq \mu\text{”}.$ Also in $\mathbf{V}, \mu^\lambda = \mu > \kappa > \lambda$ and $|\mathbb{P}'_{\gamma(*)}| = \kappa^\lambda$ by (A)(g) of 1.10(4) which is $\leq \mu$ and \mathbb{P}'_κ satisfies the λ^+ -c.c. hence $\Vdash_{\mathbb{P}'_\kappa} \text{“}2^\lambda = \mu\text{”}$, hence together $\Vdash_{\mathbb{P}'_\kappa} \text{“}\mathfrak{d}_\lambda = \mu\text{”}.$ Also by $(*)_1(B)(b)$, $\mathbb{P}'_{\gamma(*)}$ is $(< \lambda)$ -strategically complete $+ \lambda^+$ -c.c.” and by $(*)_5 + (*_6)$ we know that $\text{“}\text{cov}_\lambda(\text{meagre}) = \kappa\text{”}$ so we are done; hence $(*)_7$ is really the last piece missing. The rest of the proof is dedicated to proving that $(*)_7$ holds.

We shall use further nice properties of $\mathbb{P}'_j, \underline{g}'_i (j \leq \gamma(*), i < \gamma(*))$ which hold by $(*)_1 + (*)_2$ (and $(*)_3, (*)_4$) and their proof, i.e. 1.10, 1.11 and see [Sheb, 2.12=LC51].

- \boxplus_1 (a) (α) $\langle \underline{g}'_\gamma : \gamma < \gamma(*) \rangle$ is generic for $\mathbb{P}'_{\gamma(*)}$, i.e., if \mathbf{G} is a subset of $\mathbb{P}'_{\gamma(*)}$ generic over \mathbf{V} and $g'_i = \underline{g}'_i[\mathbf{G}]$ then $\mathbf{V}[\mathbf{G}] = \mathbf{V}[\langle g'_i : i < \gamma(*) \rangle]$
 (β) if $\nu \in (\lambda)^\lambda \mathbf{V}[\mathbf{G}]$ then for some $\rho \in (\lambda)^\lambda \mathbf{V}$ and λ -Borel function $\mathbf{B} \in \mathbf{V}$ we have $\nu = \mathbf{B}(\langle g'_{\rho(\varepsilon)} : \varepsilon < \lambda \rangle)$
- (b) if in $\mathbf{V}[\mathbf{G}]$, $g''_\gamma \in \prod_{\zeta < \lambda} \theta_\zeta$ for $\gamma < \gamma(*)$ and the set $\{(\gamma, \zeta) : \gamma < \gamma(*) \text{ and } \zeta < \lambda \text{ and } g''_\gamma(\zeta) \neq g'_\gamma(\zeta)\}$ has cardinality $< \lambda$ then $\bar{g}'' = \langle g''_\gamma : \gamma < \gamma(*) \rangle$ is generic for $\mathbb{P}'_{\gamma(*)}$ and $\mathbf{V}[\bar{g}''] = \mathbf{V}[\bar{g}']$
- (c) $\Vdash_{\mathbb{P}'_\gamma} \text{“} \underline{g}'_\gamma \text{ dominates } (\prod_{\varepsilon < \lambda} \theta_\varepsilon)^{\mathbf{V}[\mathbb{P}'_\gamma]} \text{”}$
- (d) if $\langle \zeta(\gamma) : \gamma < \gamma(*) \rangle$ is an increasing sequence of ordinals $< \gamma(*)$ (from \mathbf{V} !), then $\langle \underline{g}'_{\zeta(\gamma)} : \gamma < \gamma(*) \rangle$ is generic for $\mathbb{P}'_{\gamma(*)}$ (over \mathbf{V})
- (e) if $\gamma \leq \gamma(*)$ then \mathbb{P}'_γ is $(< \lambda)$ -strategically complete and satisfies the λ^+ -c.c.

We shall use \boxplus_1 freely.

To prove $(*)_7$ assume toward contradiction that this fails, and hence for some condition $p^* \in \mathbb{P}'_{\gamma(*)}$ and $\mathbb{P}'_{\gamma(*)}$ -name \underline{f} and λ -Borel function \mathbf{B} and $\rho \in \lambda^{\gamma(*)}$ we have:

$$\otimes_0 \ p^* \Vdash_{\mathbb{P}'_{\gamma(*)}} \text{“} \underline{f} \in \lambda^\lambda \text{ and dominates } (\lambda)^\lambda \mathbf{V} \text{” and } \underline{f} = \mathbf{B}(\langle \underline{g}'_{\rho(i)} : i < \lambda \rangle).$$

Now let χ be regular large enough and we choose $\bar{N} = \langle N_\varepsilon : \varepsilon < \lambda \rangle$ such that

- \otimes_1 (a) N_ε is as in \square_λ for the forcing notion $\mathbb{P}_{0,\beta(*)}$ (equivalently $\mathbb{P}_{1,\beta(*)}$) that is $N_\varepsilon \in \mathcal{S}_{\chi, \mathbb{P}'_{\gamma(*)}}$, see \square_λ of 2.3
 (b) $\bar{N} \upharpoonright \varepsilon \in N_\varepsilon$ and $\text{otp}(N_\varepsilon \cap \kappa) < \theta_{(\lambda_\varepsilon)}$ hence $\bigcup_{\zeta < \varepsilon} N_\zeta \subseteq N_\varepsilon$ where $\lambda_\varepsilon := \text{otp}(N_\varepsilon \cap \lambda) > \lambda_\varepsilon^- := \Sigma\{\|N_\zeta\| : \zeta < \varepsilon\} \geq \Sigma\{\lambda_\zeta : \zeta < \varepsilon\}$
 (c) $\bar{\theta}, \mathbf{q}, \mathcal{U}_*, p^*, \underline{f}, \mathbf{B}, \rho$ belong to N_ε .

Next choose $f^* \in \lambda^\lambda$, i.e. $\in (\lambda)^\lambda \mathbf{V}$, such that:

$$\otimes_2 \text{ for arbitrarily large } \varepsilon < \lambda \text{ for some } \zeta \in [\lambda_\varepsilon^-, \lambda_\varepsilon) \text{ we have } f^*(\zeta) > \lambda_\varepsilon, \text{ (we can demand more: for every } \varepsilon < \lambda).$$

For $\varepsilon < \lambda$ let $(\lambda_\varepsilon, \chi_\varepsilon, \mathbf{j}_\varepsilon, M_\varepsilon, \mathbb{A}_\varepsilon, \mathbf{G}_\varepsilon)$ be a witness for $(N_\varepsilon, \mathbb{P}_{1,\beta(*)}, \chi)$ recalling \square_λ from Definition 2.2 so $\lambda_\varepsilon \in (\varepsilon, \lambda)$ is strongly inaccessible and $\varepsilon < \zeta < \lambda \Rightarrow \lambda_\varepsilon < \lambda_\zeta^- < \lambda_\zeta$, recalling \otimes_1 and noting $\langle \lambda_\varepsilon^- : \varepsilon < \lambda \rangle$ is an increasing and a continuous sequence of cardinals below λ .

Let

- \otimes_3 (a) $v_\varepsilon = N_\varepsilon \cap \gamma(*)$
 (b) $i_\varepsilon = i(\varepsilon) = \text{otp}(v_\varepsilon)$ and so $i(\varepsilon) = \mathbf{j}_\varepsilon(\gamma(*))$, etc.
 (c) $\bar{\gamma}^\varepsilon = \langle \gamma_i(\varepsilon) : i < i(\varepsilon) \rangle$ list v_ε in increasing order

- (d) for $i < \text{otp}(v_\varepsilon)$, equivalently $i < \mathbf{j}_\varepsilon(\gamma(*))$ let $\eta_i^\varepsilon = (\mathbf{j}_\varepsilon(g'_{\gamma_i(\varepsilon)}))^{\mathbb{A}_\varepsilon[\mathbf{G}_\varepsilon]} \in \prod_{\zeta < \lambda_\varepsilon} \theta_\zeta$ and let $\bar{\eta}^\varepsilon = \langle \eta_i^\varepsilon : i < i_\varepsilon \rangle$.

Note that clearly

- ⊗₄ (a) $\bar{\eta}^\varepsilon$ is generic for $(\mathbb{A}_\varepsilon, \mathbf{j}_\varepsilon(\mathbb{P}'_{\gamma(*)}))$, moreover
 (b) for each $\varepsilon < \lambda$, if we change $\eta_i^\varepsilon(\zeta)$ (legally, i.e. to an ordinal $< \theta_\zeta$) for $< \lambda_\varepsilon$ pairs $(i, \zeta) \in \text{otp}(v_\varepsilon) \times \lambda_\varepsilon$ and get $\bar{\eta}'$, then also $\bar{\eta}'$ is generic for $(\mathbb{A}_\varepsilon, \mathbf{j}_\varepsilon(\mathbb{P}'_{\gamma(*)}))$ and $\mathbb{A}_\varepsilon[\bar{\eta}'] = M_\varepsilon$
 (c) like \boxplus_1 with $\mathbf{V}, \mathbb{P}_{\beta(*)}, \lambda$ there standing for $\mathbb{A}_\varepsilon, \mathbf{j}_\varepsilon(\mathbb{P}_{\beta(*)}), \lambda_\varepsilon$ here.

Hence we have

- ⊗'₄ for $\varepsilon < \lambda$,
 (a) let $\Xi_\varepsilon := \{\bar{\nu} : \bar{\nu} = \langle \nu_i : i < i(\varepsilon) \rangle\}$ and for some $\mathbf{G} \subseteq \mathbb{P}'_{\gamma(*)} \cap N_\varepsilon$ generic over N_ε we have $\nu_i \in \prod_{\xi < \lambda_\varepsilon} \theta_\xi$ satisfies $\xi < \lambda_\varepsilon \Rightarrow$ some $\psi \in \mathbf{G}$ forces $g'_{\gamma_i(\varepsilon)} \upharpoonright \xi = \nu_i \upharpoonright \xi$ and there is a subset of \mathbf{G}^+ of $\mathbb{P}_{\beta(*)}$ generic over N_ε such that the Mostowski collapse of $N_\varepsilon[\mathbf{G}^+]$ is $\mathcal{H}(\chi_\varepsilon)$
 (b) if $\bar{\nu} \in \Xi_\varepsilon$ then there is one and only one $\mathbf{G} = \mathbf{G}_{\bar{\nu}}$ as above
 (c) we can choose $\bar{p}_\varepsilon^* = \langle p_{\varepsilon, \psi, \bar{\nu}, \epsilon}^* : \psi \in \mathbb{P}'_{\gamma(*)} \cap N_\varepsilon, \bar{\nu} \in \Xi_\varepsilon, \epsilon < \lambda_\varepsilon \rangle$ such that
 (α) $p_{\varepsilon, \psi, \bar{\nu}, \epsilon}^* \in \mathbb{P}_{\beta(*)}$ moreover, if $j \leq \sigma$ and $\psi \in \mathbb{P}'_{\gamma_j(\varepsilon)}$ then $p_{\varepsilon, \psi, \bar{\nu}, \epsilon}^* \in \mathbb{P}_{\gamma_j(\varepsilon)}$
 (β) $\mathbb{P}_{\beta(*)} \models \psi \leq p_{\varepsilon, \psi, \bar{\nu}, \epsilon}^*$ moreover, if $\mathbb{P}'_{\beta(*)} \Vdash \psi \leq \phi$ then $p_{\varepsilon, \psi, \bar{\nu}, \epsilon}^*, \phi$ are compatible in $\mathbb{P}_{\gamma(*)}$
 (γ) $\langle \text{dom}(p_{\varepsilon, \psi, \bar{\nu}, \epsilon}^*) \setminus \mathcal{U}_* : \psi \in \mathbb{P}'_{\gamma(*)}, \bar{\nu} \in \Xi_\varepsilon \rangle$ is a sequence of pairwise disjoint sets; in [Sheb] we say more
 (δ) $p_{\varepsilon, \psi, \bar{\nu}, \epsilon}^*(\gamma_i(\varepsilon))$ has trunk of length $\leq \epsilon$ which is an initial segment of ν_ϵ when defined
 (d) assume $\bar{\nu} \in \Xi_\varepsilon$
 (α) for \bar{p}_ε^* as in clause (c) and $\bar{\nu} \in \Xi_\varepsilon$ there is $q \in \mathbb{P}_{\beta(*)}$ which is an upper bound of $\{p_{\varepsilon, \psi, \bar{\nu}, \epsilon}^* : \psi \in \mathbf{G}_{\bar{\nu}}, \epsilon < \lambda_\varepsilon\}$
 (β) if $q \in \mathbb{P}_{\beta(*)}$ is an upper bound of $\mathbf{G}_{\bar{\nu}}$ then q is $(N_\varepsilon, \mathbb{P}'_{\gamma(*)})$ -generic naturally and $q \Vdash_{\mathbb{P}'_{\gamma(*)}} \text{“}\mathbf{j}_\varepsilon \text{ can be extended naturally to an isomorphism from } N_\varepsilon[\mathbf{G}_{\mathbb{P}'_{\gamma(*)}}] = N_\varepsilon[\langle g'_\gamma : \gamma \in v_\varepsilon \rangle] \text{ onto } \mathbb{A}_\varepsilon[\bar{\eta}^\varepsilon]\text{”}$.
 (e) Similarly for $\varepsilon < \lambda, j \leq i(\varepsilon)$ and $\bar{\nu} = \langle \nu_i : i < j \rangle$

[Why? See [Sheb, 3.28-3.32=Le53-Le67]]

By the assumption toward contradiction, ⊗₀, and $\mathbb{P}'_{\gamma(*)}$ being $(< \lambda)$ -strategically complete, recalling \boxplus_1 , there are $\zeta(*), p^{**}$ and p^+ such that (recall $p^* \in \mathbb{P}'_{\gamma(*)} \leq \mathbb{P}_{1, \beta(*)}$ is from ⊗₀):

- ⊗₅ (a) $p^* \leq p^{**} \in \mathbb{P}'_{\gamma(*)}$ and $p^+ \in \mathbb{P}_{0, \beta(*)}$ satisfies $\mathbb{P}_{1, \beta(*)} \models \text{“}p^{**} \leq p^+\text{”}$
 (b) $\zeta(*) < \lambda$
 (c) $p^{**} \Vdash_{\mathbb{P}'_{\gamma(*)}} \text{“}f^*(\zeta) < f(\zeta) \text{ whenever } \zeta(*) \leq \zeta < \lambda\text{”}$ where f^* is from ⊗₂

- (d) if $\gamma \in \text{Dom}(p^+)$ then $\eta^{p^+(\gamma)}$ is an object (not just a $\mathbb{P}_{0,\gamma}$ -name) and has length $\geq \zeta(*)$ (recall that $\eta^{p^+(\gamma)}$ is the trunk of the condition $p^+(\gamma)$, see clause $(\alpha)(b)$ of Definition 1.4(1)).

Note that possibly $\text{Dom}(p^+) \not\subseteq \cup\{v_\varepsilon : \varepsilon < \lambda\}$. Choose $\varepsilon(*) < \lambda$ such that $\lambda_{\varepsilon(*)} > \zeta(*) + |\text{Dom}(p^+)|$ and $\gamma \in \text{Dom}(p^+) \Rightarrow \varepsilon(*) > \ell g(\eta^{p^+(\gamma)})$ recalling clause (d) of \otimes_5 and $|\text{Dom}(p^+)| < \lambda$ as $p^+ \in \mathbb{P}_{0,\beta(*)}$ and $\mathbb{P}_{0,\beta(*)}$ is the limit of a $(< \lambda)$ -support iteration.

By \otimes_2 we can add $(\exists \zeta)[\lambda_{\varepsilon(*)}^- \leq \zeta < \lambda_{\varepsilon(*)} < f^*(\zeta)]$. Our intention is to find $q \in \mathbb{P}_{0,\beta(*)}$ above p^+ which (in $\mathbb{P}_{1,\beta(*)}$) is above some $q' \in \mathbb{P}'_{\gamma(*)}$ which is $(N_{\varepsilon(*)}, \mathbb{P}'_{\gamma(*)})$ -generic and forces $\mathbf{G}_{\mathbb{P}'_{\gamma(*)}}$ to include a generic subset of $(\mathbb{P}'_{\gamma(*)})^{N_{\varepsilon(*)}}$ hence is induced by some $\bar{\nu}$ as in \otimes'_4 , recalling $\otimes_4(b)$. Toward this in \otimes_6 below the intention is that $p_{i(*)}^+$ will serve as q .

Let $i(*) = i(\varepsilon(*))$ and γ_i for $i < i(*)$ be such that¹ $\langle \gamma_i : i < i(*) \rangle$ list $\{\beta_i^* : i \in v_{\varepsilon(*)}\} \subseteq \mathcal{Z}_{**} = N_{\varepsilon(*)} \cap \mathcal{Z}_*$ in increasing order; recall $\mathcal{Z}_* = \{\beta_i^* : i < \gamma(*)\}$ and $i < j < \gamma(*) \Rightarrow \beta_i^* < \beta_j^*$ and $v_{\varepsilon(*)} \subseteq \gamma(*)$ has order type $i(\varepsilon(*))$. Next let $\gamma_{i(*)} = \gamma(*)$ so $\{\mathbf{j}_{\varepsilon(*)}(\gamma) : \gamma \in v_{\varepsilon(*)}\} = i(*) = \mathbf{j}_{\varepsilon(*)}(\gamma(*))$. Recall $\gamma(*) = \kappa = \text{cf}(\kappa) > \lambda$, $\text{otp}(v_{\varepsilon(*)}) = \text{otp}(N_{\varepsilon(*)} \cap \gamma(*)) = \text{otp}(N_{\varepsilon(*)} \cap \kappa)$ hence $N_{\varepsilon(*)} \models$ “ $i(*)$ is a regular cardinal $> \lambda_\varepsilon$ ” hence $i(*)$ is really a regular cardinal so call it σ . Now we define a game \boxplus as follows²:

- \boxplus_2 (A) each play lasts $i(*) + 1 = \sigma + 1$ moves and in the i -th move,
- (a) if $i = j + 1$ the antagonist player chooses $\xi_j = \xi(j) < \sigma$ such that $j_1 < j \Rightarrow \xi(j_1) < \xi(j)$
 - (b) then, if $i = j + 1$ the protagonist chooses $\zeta_j = \zeta(j) \in (\xi(j), \sigma)$, but there are more restrictions implicit in \boxplus_3 below
 - (c) in any case the protagonist also chooses $p_i^+, \bar{\nu}^i$ such that \boxplus_3 below holds;
- (B) in the end of the play the protagonist wins the play iff it always has a legal move and in the end p_σ^+ is $(\mathbb{P}'_{\gamma(*)}, N)$ -generic and $\{\zeta(i) : i \leq i(*)\} \in \mathbb{A}_{\varepsilon(*)}$; note that trivially it belongs to $M_{\varepsilon(*)} = \mathbb{A}_{\varepsilon(*)}[\bar{\eta}^{\varepsilon(*)}]$, see $\otimes_3(d)$

where

- \boxplus_3 (a) $p_i^+ \in \mathbb{P}_{0,\gamma_i}$ and $p_i^+, \langle \nu_j : j < i \rangle, \mathbf{G}_{\varepsilon(*),i} \subseteq \mathbb{P}'_{\gamma_i} \cap N$ are as in $\otimes'_4(e)$
- (b) if $j < i$ then $\mathbb{P}_{0,\gamma_i} \models$ “ $p_j^+ \leq p_i^+$ ”
 - (c) if $\gamma \in \cup\{\text{Dom}(p_j^+) : j < i\}$ then $p_i^+ \upharpoonright \gamma \Vdash_{\mathbb{P}_{0,\gamma_i}}$ “ $\eta^{p_i^+(\gamma)}$ has length $\geq i(*)$ and $\geq \lambda_{\varepsilon(*)}$ ” moreover $\eta^{p_i^+(\gamma)}$ is an object, $\eta^{p_i^+(\gamma)}$
 - (d) $\mathbb{P}_{0,\gamma_i} \models$ “ $p^+ \upharpoonright \gamma_i \leq p_i^+$ ”, (p^+ is from \otimes_5)

¹This is used in \boxplus_3 and the proof of $(*)_6$. Not to be confused with $\bar{\gamma}^\varepsilon$ of $\otimes_3(c)$.

²The idea is to scatter the $\eta_{\gamma_i}^{\varepsilon(*)}$'s. Why not use the original places? as then we shall have a problem in \otimes_6 ; the scattering is helpful because we are relying on 1.10 and 1.11.

- (e) $\bar{\nu}^i = \langle \nu_{\gamma_j} : j < i \rangle$ and $\nu_{\gamma_j} \in \prod_{\iota < \lambda_{\varepsilon(*)}} \theta_\iota$ and p_σ^+ is an upper bound of $\Omega_{\varepsilon(*)}^{\bar{\nu}^i} = \{p_{\varepsilon, \psi, \bar{\nu}, \varepsilon}^* \upharpoonright \gamma_i : \bar{\nu} \in \Xi_{\varepsilon(*)} \text{ satisfies } \bar{\nu}^i \triangleleft \bar{\nu} \text{ and } \psi \in \mathbf{G}_{\bar{\nu}}\}$ and $\varepsilon < \lambda_{\varepsilon(*)}$
- (f) for $j < i$ we have $\nu_{\gamma_j} \leq \eta^{p_i^+(\gamma_j)}$ so $p_i^+ \upharpoonright \gamma_j \Vdash \text{“}\nu_{\gamma_j} \triangleleft \underline{g}'_{\gamma_j}\text{”}$ recalling \boxplus_1
- (g) for $j < i$ we have (recall $\bar{\eta}^\varepsilon$ from $\otimes_3(d)$)
- (α) $\nu_{\gamma_j} = \eta_{\gamma_{\zeta(j)}}^{\varepsilon(*)}$ recalling $\eta_{\gamma_j}^{\varepsilon(*)}$ is from $\otimes_3(d)$ or
- (β) $\gamma_j \in \text{Dom}(p^+)$ and $\{\iota < \lambda_{\varepsilon(*)} : \eta_{\zeta(j)}^{\varepsilon(*)}(\iota) \neq \nu_{\gamma_j}(\iota)\}$ is a bounded subset of $\lambda_{\varepsilon(*)}$.

We shall prove

- \otimes_6 in the game \diamond
- (a) the antagonist has no winning strategy
- (b) at stage i , if $\langle \zeta(j) : j < i \rangle \in N$ then the protagonist has a legal move, moreover for any $\zeta(i) \in (\xi(i), \sigma)$ large enough the protagonist can choose it.

Why \otimes_6 suffice?

By clause (a) of \otimes_6 we can choose a play $\langle (\xi(i), \zeta(i), p_i^+, \bar{\nu}^i) : i \leq \sigma) \rangle$ in which the protagonist wins. Recalling $\mathbb{P}'_{\gamma(*)} \triangleleft \mathbb{P}_{1, \beta(*)}$ and $\mathbb{P}_{0, \beta(*)}$ is a dense subforcing of $\mathbb{P}_{1, \beta(*)}$, clearly

- \otimes_7 there is p such that
- (a) $p \in \mathbb{P}'_{\gamma(*)}$
- (b) if $\mathbb{P}'_{\gamma(*)} \Vdash \text{“}p \leq p'\text{”}$ hence $p' \in \mathbb{P}'_{\beta(*)}$ then p', p_σ^+ are compatible in $\mathbb{P}_{1, \beta(*)}$
- (c) p is above p^{**} and it forces $\underline{g}'_{\gamma_i} \upharpoonright \lambda_{\varepsilon(*)} = \nu_{\gamma_i}$ for $i < i(*)$. $\mathbf{G}_{\mathbb{P}'_{\beta(*)}} \cap N = \mathbf{G}_{\langle \nu_{\gamma_i} : i < \sigma \rangle}$

Then on the one hand

- \otimes'_7 $p \in \mathbb{P}'_{\gamma(*)}$ being above p^{**} forces $f^* \upharpoonright [\zeta(*), \lambda) < \underline{f} \upharpoonright [\zeta(*), \lambda)$ hence $f^* \upharpoonright [\zeta(*), \lambda_{\varepsilon(*)}) < \underline{f} \upharpoonright [\zeta(*), \lambda_{\varepsilon(*)})$ recalling that $\zeta(*) < \lambda_{\varepsilon(*)}$, see \otimes_5 and the choice of $\varepsilon(*)$ immediately after \otimes_5 .

On the other hand,

- \otimes''_7 p is $(N_{\varepsilon(*)}, \mathbb{P}'_{\gamma(*)})$ -generic.

[Why? As it forces $\eta_{\gamma_i} \upharpoonright \lambda_{\varepsilon(*)} = \nu_{\gamma_i}$ for $i < i(*)$ and $\langle \nu_{\gamma_i} : i < i(*) \rangle$ is (see $\boxplus_3(g)$ recalling $\text{Dom}(p^{**})$ has cardinality $< \lambda_{\varepsilon(*)}$) “almost equal” to $\langle \eta_{\zeta(i)}^{\varepsilon(*)} : i < i(*) \rangle$ which is a subsequence of the sequence from \otimes_3 . That is $\{(i, \iota) : \iota < \lambda_{\varepsilon(*)}, i < i(*) = \sigma \text{ and } \nu_{\gamma_i}(\iota) \neq \eta_{\zeta(i)}^{\varepsilon(*)}(\iota)\} \subseteq \cup\{\{(i, \iota) : \iota < \lambda_{\varepsilon(*)} \text{ and } \nu_{\gamma_i}(\iota) \neq \eta_{\zeta(i)}^{\varepsilon(*)}(\iota)\} : \gamma \in v_{\varepsilon(*)} \cap \text{Dom}(p^{**})\}$ so is the union of $\leq |\text{Dom}(p_\sigma^+)| < \lambda_{\varepsilon(*)}$ sets each of cardinality $< \lambda_{\varepsilon(*)}$ hence is of cardinality $< \lambda_{\varepsilon(*)}$. Hence by $\otimes_4(c) + \boxplus_1(d)$ the sequence $\bar{\nu}^{i(*)}$ is generic for $(N_{\varepsilon(*)}, \mathbb{P}'_{\gamma(*)})$. By \boxplus_2 and the choice of p_σ^+ above it is $(N_{\varepsilon(*)}, \mathbb{P}'_{\gamma(*)})$ -generic. By $\otimes_7(b)$ also p is.

As $f \in N_{\varepsilon(*)}$ it follows from \otimes_7'' that

$$\otimes_7''' \ p \Vdash \text{“} f \upharpoonright \lambda_{\varepsilon(*)} \text{ is a function from } \lambda_{\varepsilon(*)} \text{ to } \lambda_{\varepsilon(*)}\text{”}.$$

Together $\otimes_7' + \otimes_7'''$ gives a contradiction by the choice of f^* in \otimes_2 and of $\varepsilon(*)$ above, hence \otimes_6 is enough. In Lemma 2.8 below we show that \otimes_6 is true; so we are done. $\square_{2.7}$

Lemma 2.8. *The statement \otimes_6 is true.*

Proof. Let us prove \otimes_6 ; first, assuming clause (b) which is proved below, for clause (a) choose any strategy \mathbf{st} for the antagonist and fix a partial strategy \mathbf{st}' for the protagonist choosing (p_i^+, \bar{v}^i) depending on the previous choices and $\chi(i) < i_{\varepsilon(*)}$ such that it is a legal move if relevant and possible. So the only freedom left for the protagonist is to choose the $\zeta(i)$. So (recalling $\boxplus_2(A)(a)$) we have in \mathbf{V} a function $F : \sigma > \sigma \rightarrow \sigma$ (so F uses $\mathbf{st} \dots$) such that:

- (*) $_F$ playing the game such that the antagonist uses \mathbf{st} and the protagonist uses \mathbf{st}' , arriving to the i -th move, $\bar{\zeta} = \langle \zeta(j) : j < i \rangle$ is well defined and if $\zeta \in N$ then for the protagonist any choice $\zeta_i \in (F(\bar{\zeta}), \sigma) \cap \mathcal{U}_{**}$ is legal.

Now we have to find an increasing sequence $\bar{\zeta} = \langle \zeta(i) : i < \sigma \rangle$ from $\mathbb{A}_{\varepsilon(*)}$ not just from $M_{\varepsilon(*)} = \mathcal{H}(\chi_{\varepsilon(*)})^{\mathbf{V}}$ such that $F(\bar{\zeta} \upharpoonright i) < \zeta(i) < \sigma$ and $\bar{\zeta} \in \mathbb{A}_{\varepsilon(*)}$. As $F \in \mathcal{H}(\chi_{\varepsilon(*)})$ and $\mathcal{H}(\chi_{\varepsilon(*)}) = M_{\varepsilon(*)} = \mathbb{A}_{\varepsilon(*)}[\mathbf{G}_{\varepsilon(*)}]$ where $\mathbf{G}_{\varepsilon(*)}$ is a subset of $\mathbf{j}_{\varepsilon(*)}(\mathbb{P}'_{\gamma(*)}) \in \mathbb{A}_{\varepsilon(*)}$ generic over $\mathbb{A}_{\varepsilon(*)}$ and $\mathbf{j}_{\varepsilon(*)}(\mathbb{P}_{0,\beta(*)})$ satisfies the $\lambda_{\varepsilon(*)}^+$ -c.c. and $\sigma = \text{cf}(\sigma) > \lambda_{\varepsilon(*)}$ this is possible. That is, there is a $\mathbf{j}_{\varepsilon(*)}(\mathbb{P}_{0,\beta(*)})$ -name $\bar{F}_* \in \mathbb{A}_{\varepsilon(*)}$ such that $F = \bar{F}_*[\mathbf{G}_{\varepsilon(*)}]$ and we define in $\mathbb{A}_{\varepsilon(*)}$ the function $F' : \sigma > \sigma \rightarrow \sigma$ by $F'(\langle \zeta(j) : j < i \rangle) = \sup\{\xi + 1 : \xi \in \{\zeta(j) + 1 : j < i\} \text{ or } \xi < \sigma \text{ and } \mathcal{K}_{\mathbf{j}(\mathbb{P}_{0,\beta(*)})}\}$ “ $\bar{F}(\langle \zeta(j) : j < i \rangle) \neq \xi$ ”; clearly this is O.K.

We are left with proving $\otimes_6(b)$.

Case 1: $i = 0$.

Let $p_0^+ = p^{**} \upharpoonright \gamma_0$.

Case 2: i limit.

By clauses (b) and (c) of \boxplus_3 , there is $p_i^+ \in \mathbb{P}_{0,\gamma_i}$ which is an upper bound (even l.u.b.) of $\{p_j^+ : j < i\} \cup \Omega_{\varepsilon(*)}^{\bar{v}^i}$ and it is easily as required. Also \bar{v}^i is well defined and as required.

Case 3: $i = j + 1$ and $\gamma_j \notin \text{Dom}(p^+)$

Clearly γ_i is in \mathcal{U}_* the successor of γ_j and $(\exists \iota)(\gamma_j = \beta_\iota^* \wedge \iota \in v_{\varepsilon(*)})$. As in case 4 below but easier by the properties of the iteration and [Sheb, §3C].

Case 4: $i = j + 1$ and $\gamma_j \in \text{Dom}(p^+)$

Again γ_i is in \mathcal{U}_* the successor of γ_j and $(\exists \iota)(\gamma_j = \beta_\iota^* \wedge \iota \in v_{\varepsilon(*)})$.

First we find p'_j such that:

- \otimes_8 (a) $p_j^+ \leq p'_j \in \mathbb{P}_{0,\gamma_j}$
 (b) if $\gamma \in \text{Dom}(p_j^+)$ then $p'_j \upharpoonright \gamma \Vdash \text{“} \ell g(\eta^{p'_j(\gamma)}) > i(*) = \sigma \text{”}$ (see $\boxplus_3(c)$)

- (c) p'_j forces ³ a value to the pair $(\eta^{p^+(\gamma_j)}, f^{p^+(\gamma_j)} \upharpoonright \lambda_{\varepsilon(*)})$; we call this pair $q_j = (\eta^{q_j}, f^{q_j})$.

[Why? This should be clear.]

Second

- ⊗₉ p'_j hence p'_j is $(N_{\varepsilon(*)}, \mathbb{P}'_{\gamma_j})$ -generic and $\langle \nu_{\gamma_j(1)} : j(1) < j \rangle$ induces the generic.

[Why? As in the proof of ⊗₇' of Lemma 2.7 when we assume that we have carried the induction, by ⊕₂, clause (g) and ⊗₄.]

Now

- ⊗₁₀ (a) $f^{q_j} \in (\prod_{\zeta < \lambda_{\varepsilon(*)}} \theta_\zeta)^{\mathbb{A}_{\varepsilon(*)}[\bar{\nu}^j]}$
 (b) for every large enough $\zeta \in (\xi(i), \sigma)$ we have
- ₁ $f^{q_j} \leq \eta_\zeta^{\varepsilon(*)} \pmod{J_{\lambda_\varepsilon}^{\text{bd}}}$
 - ₂ $f^{q_j} \in \mathbb{A}_{\varepsilon(*)}[\bar{\eta}^{\varepsilon(*)} \upharpoonright \zeta]$
 - ₃ $\langle \zeta(j_1) : j_1 < j \rangle \in \mathbb{A}_{\varepsilon(*)}[\bar{\eta}^{\varepsilon(*)} \upharpoonright \zeta]$.
- (c) $\eta^{q_j} \triangleleft f^{q_j}$.

[Why? Clause (a) holds because $f^{q_j} \in (\prod_{\zeta < \lambda_{\varepsilon(*)}} \theta_\zeta)^{\mathbf{V}}$, hence belongs to $\mathcal{H}(\chi_{\varepsilon(*)})$

which is the universe of $M_{\varepsilon(*)}$ so $f^{q_j} \in M_{\varepsilon(*)}$. But $M_{\varepsilon(*)} = \mathbb{A}_{\varepsilon(*)}[\bar{\eta}^{\varepsilon(*)}]$; recalling $\bar{\eta}^{\varepsilon(*)}$ is a generic for $\mathbf{j}_\varepsilon(\mathbb{P}'_{\gamma(*)})$. Next as $\mathbb{P}'_{\gamma(*)}$ satisfies the λ^+ -c.c. and $\lambda < \kappa = \text{cf}(\gamma(*))$ so $\mathbf{j}_\varepsilon(\mathbb{P}'_{\gamma(*)})$ satisfies the $\lambda_{\varepsilon(*)}^+$ -c.c. hence for some $\zeta_1 < \sigma$, $f^{q_j} \in \mathbb{A}_{\varepsilon(*)}[\bar{\eta}^{\varepsilon(*)} \upharpoonright \zeta_1]$. Similarly for some $\zeta_2 < \sigma$ we have $\langle \zeta(j_2) : j_2 < j \rangle$ belongs to $N'_{\varepsilon(*)}[\bar{\eta}^{\varepsilon(*)} \upharpoonright \zeta_2]$. Letting $\zeta \geq \max\{\zeta_1, \zeta_2\}$ clearly clauses •₂, •₃ of ⊗₁₀(b) holds, also •₁ there holds by ⊕₁(c) (and \mathbf{j}_ε , etc.).]

- ⊗₁₁ $M_{\varepsilon(*)} = \mathbb{A}_{\varepsilon(*)}[\bar{\eta}^{\varepsilon(*)}]$ satisfies: for every $\rho \in (\lambda_{\varepsilon(*)})^{>}(\lambda_{\varepsilon(*)})$, in particular, $\rho = \eta^{q_j}$, the set $\{\zeta < \sigma : \rho \triangleleft \eta_\zeta^{\varepsilon(*)}\}$ is unbounded in σ .

[Why? Because the iteration $\mathbf{q} = \langle \mathbb{P}_\alpha, \mathbb{Q}_\beta : \alpha \leq \kappa, \beta < \kappa \rangle$ is with support $< \lambda$ and similarly $\mathbf{j}_\varepsilon(\mathbf{q})$.

Lastly, ⊗₁₀(c) holds by ⊗₈(c).]

Now we choose $\zeta(j)$ as in clause (b) of ⊗₁₀ and $\nu_j \in \prod_{\varepsilon < \lambda_{\varepsilon(*)}} \theta_\varepsilon$ such that $\eta^{q_j} \triangleleft$

$\nu_j, \eta^{p^+(j)} \triangleleft \nu_j, f^{q_j} \leq \nu_j$ and $\{\iota < \lambda_{\varepsilon(*)} : \nu_j(\iota) \neq \eta_{\zeta(j)}^{\varepsilon(*)}\}$ is a bounded subset of $\lambda_{\varepsilon(*)}$. Next choose $p_i^+ \in \mathbb{P}'_{\gamma(*)}$ such that $p_i^+ \upharpoonright \gamma_j = p'_j, \eta^{p_i^+(\gamma_i)} = \nu_j$ and $f^{p_i^+(\gamma_i)} \upharpoonright [\lambda_\varepsilon, \lambda) = f^{p^+(\gamma)} \upharpoonright [\lambda_\varepsilon, \lambda)$.

We have carried the induction hence proved ⊗₆(b) so we are done proving 2.8.

□_{2.8}

Discussion 2.9. 1) The reader may justly wonder why we use $\mathbf{V}' = \mathbf{V}[\bar{g}'] = \mathbf{V}[\bar{g} \upharpoonright \mathcal{U}_*]$ rather than simply $\mathbf{V}[\bar{g}]$. Of course, nothing is lost by it, but why the extra complication?

2) The answer is that during the proof we shall use: if $\zeta(i) \in \mathcal{U}_*$ is increasing with $i < \gamma(*)$ then also $\langle g_{\zeta(i)} : i < \kappa \rangle$ is generic over \mathbf{V} for the subforcing of $\mathbb{P}_{1,\beta(*)}$

³recall that $\eta^{p^+(\gamma_j)}$ is an object, not a name and p_j^+ is $(N_{\varepsilon(*)}, \mathbb{P}'_{\gamma_j})$ -generic

generated by $\bar{g} \upharpoonright \mathcal{U}_*$; see $\textcircled{7}''$ inside the proof of $\textcircled{6}$ inside 2.8. But using $\mathcal{U}_* = \beta(*)$, we do not know this.

3) Now in the parallel case for $\lambda = \aleph_0$ with FS iteration with full memory, such claim is true, see §0.

4) But we do not know the parallel of 3) for λ , so we use a substitute using \mathcal{U}_* , i.e. \mathbb{P}'_κ .

REFERENCES

- [Bar87] Tomek Bartoszyński, *Combinatorial aspects of measure and category*, *Fundamenta Mathematicae* **127** (1987), 225–239.
- [Bar10] ———, *Invariants of measure and category*, *Handbook of Set Theory*, Vols. 1, 2, 3 (Matthew Foreman and Akihiro Kanamori, eds.), Springer, Dordrecht, 2010, pp. 491–555.
- [Bla] Andreas Blass, *Combinatorial cardinal characteristics of the continuum*, *Handbook of Set Theory* (Matthew Foreman and Akihiro Kanamori, eds.), vol. 1, Springer, pp. 395–490.
- [CS95] James Cummings and Saharon Shelah, *Cardinal invariants above the continuum*, *Ann. Pure Appl. Logic* **75** (1995), no. 3, 251–268, arXiv: math/9509228. MR 1355135
- [GS93] Martin Goldstern and Saharon Shelah, *Many simple cardinal invariants*, *Arch. Math. Logic* **32** (1993), no. 3, 203–221, arXiv: math/9205208. MR 1201650
- [GS12] Shimon Garti and Saharon Shelah, *A strong polarized relation*, *J. Symbolic Logic* **77** (2012), no. 3, 766–776, arXiv: 1103.0350. MR 2987137
- [HS] Haim Horowitz and Saharon Shelah, *Saccharinity with ccc*, arXiv: 1610.02706.
- [JS88] Haim I. Judah and Saharon Shelah, *Souslin forcing*, *J. Symbolic Logic* **53** (1988), no. 4, 1188–1207. MR 973109
- [KS12] Jakob Kellner and Saharon Shelah, *Creature forcing and large continuum: the joy of halving*, *Arch. Math. Logic* **51** (2012), no. 1-2, 49–70, arXiv: 1003.3425. MR 2864397
- [Lan92] A. Landver, *Baire numbers, uncountable cohen sets and perfect-set forcing*, *Journal of Symbolic Logic* **57** (1992), 1086–1107.
- [Lav78] Richard Laver, *Making the supercompactness of κ indestructible under κ -directed closed forcing*, *Israel J. of Math.* **29** (1978), 385–388.
- [Mil82] Arnold W. Miller, *A characterization of the least cardinal for which the baire category theorem fails*, *Proceedings of the American Mathematical Society* **86** (1982), 498–502.
- [MS] Pierre Matet and Saharon Shelah, *Positive partition relations for $P_\kappa(\lambda)$* , arXiv: math/0407440.
- [RS97] Andrzej Rosłanowski and Saharon Shelah, *Norms on possibilities. II. More ccc ideals on 2^ω* , *J. Appl. Anal.* **3** (1997), no. 1, 103–127, arXiv: math/9703222. MR 1618851
- [Shea] Saharon Shelah, *Bounding forcing with chain conditions for uncountable cardinals*.
- [Sheb] ———, *Corrected Iteration*.
- [Shec] ———, *Creature iteration for inaccessible*.
- [Shed] ———, *Forcing axioms for λ -complete μ^+ -c.c.*, arXiv: 1310.4042.
- [She78] ———, *A weak generalization of MA to higher cardinals*, *Israel J. Math.* **30** (1978), no. 4, 297–306. MR 0505492
- [She92] ———, *Vive la différence. I. Nonisomorphism of ultrapowers of countable models*, *Set theory of the continuum* (Berkeley, CA, 1989), *Math. Sci. Res. Inst. Publ.*, vol. 26, Springer, New York, 1992, arXiv: math/9201245, pp. 357–405. MR 1233826
- [She00] ———, *Was Sierpiński right? IV*, *J. Symbolic Logic* **65** (2000), no. 3, 1031–1054, arXiv: math/9712282. MR 1791363
- [She17] ———, *A parallel to the null ideal for inaccessible λ : Part I*, *Arch. Math. Logic* **56** (2017), no. 3-4, 319–383, arXiv: 1202.5799. MR 3633799
- [vD84] Eric K. van Douwen, *The integers and topology*, *Handbook of Set-Theoretic Topology* (K. Kunen and J. E. Vaughan, eds.), Elsevier Science Publishers, 1984, pp. 111–167.

EINSTEIN INSTITUTE OF MATHEMATICS, EDMOND J. SAFRA CAMPUS, GIVAT RAM, THE HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM, 91904, ISRAEL, AND, DEPARTMENT OF MATHEMATICS, HILL CENTER - BUSCH CAMPUS, RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY, 110 FRELINGHUYSEN ROAD, PISCATAWAY, NJ 08854-8019 USA

Email address: shelah@math.huji.ac.il

URL: <http://shelah.logic.at>