

FAILURE OF SINGULAR COMPACTNESS FOR Hom

MOHSEN ASGHARZADEH, MOHAMMAD GOLSHANI, AND SAHARON SHELAH

ABSTRACT. Assuming Gödel’s axiom of constructibility $\mathbf{V} = \mathbf{L}$, we construct a χ -free abelian group G of singular cardinality for some suitable cardinal χ which is regular and uncountable, equipped with the property that for every nontrivial subgroup $G' \subseteq G$ of smaller cardinality, $\text{Hom}(G', \mathbb{Z}) \neq 0$, while $\text{Hom}(G, \mathbb{Z}) = 0$. This provides a consistent counterexample to the singular compactness of nontrivial duality with respect to the functor $\text{Hom}(-, \mathbb{Z})$.

§ 0. INTRODUCTION

Hill [6] proved that if an abelian group G has cardinality a singular cardinal with cofinality at most ω_1 , and every subgroup of G of smaller cardinality is free, then G itself is free. This result forms one of the foundations of Shelah’s *Singular Compactness Theorem* [9], in which he introduced an abstract notion of freeness and removed the restriction on cofinality. In particular, Shelah showed that if an abelian group G is of size a singular cardinal λ , and if every subgroup of G of cardinality $< \lambda$ is free, then G must be free. General references on singular compactness may be found in [3, 4], while applications are discussed in [5].

Compactness principles, and the corresponding phenomena of incompleteness, occupy a central place in modern set-theoretic algebra. Broadly, a compactness principle asserts that if every small subobject of a given object satisfies a property Pr , then the object itself satisfies Pr . At the ASL Annual Meeting in Irvine (2008), Shelah [10] announced several theorems concerning singular incompleteness, although these results were not published. The aim of this paper is to provide a systematic analysis of incompleteness for abelian groups with respect to the duality functor $\text{Hom}(-, \mathbb{Z})$, focusing on singular cardinals. In particular, we verify a prediction from [10]. The property of interest is:

2010 *Mathematics Subject Classification*. Primary: 03E75, 20A15, Secondary: 20K20; 20K40.

Key words and phrases. Abelian groups; almost-free; singular compactness; trivial duality; set theoretical methods in algebra; relative trees.

The second author’s research has been supported by a grant from IPM (No. 1403030417). The third author research partially supported by the Israel Science Foundation (ISF) grant no: 1838/19, and Israel Science Foundation (ISF) grant no: 2320/23; Research partially supported by the grant “Independent Theories” NSF-BSF, (BSF 3013005232). The third author is grateful to Craig Falls for providing typing services that were used during the work on the paper. This is publication number 1264 of the third author.

Pr_λ : Let G be a group of cardinality λ . If every nontrivial subgroup $G' \subseteq G$ of cardinality $< \lambda$ satisfies $\text{Hom}(G', \mathbb{Z}) \neq 0$, then $\text{Hom}(G, \mathbb{Z}) \neq 0$.

For $\mu \leq \lambda$ we recall that $S_\mu^\lambda = \{\alpha < \lambda : \text{cf}(\alpha) = \mu\}$ is stationary in λ . Given a stationary set $S \subseteq \lambda$, we write \diamond_S for Jensen's diamond on S (see Definition 1.6). When $\lambda > \aleph_0$ is regular and \diamond_S holds for some stationary non-reflecting $S \subseteq S_{\aleph_0}^\lambda$, it is known (see Fact 1.8, recorded from [2]) that one may construct a λ -free abelian group G of cardinality λ such that $\text{Hom}(G, \mathbb{Z}) = 0$. Any subgroup G' of cardinality $< \lambda$ is then free, and hence $\text{Hom}(G', \mathbb{Z}) \neq 0$. Thus Pr_λ fails for such λ . These assumptions entail that λ is not weakly compact, since weak compactness implies reflection of stationary sets, whereas we require a non-reflecting stationary subset. In fact, the non-weak compactness of λ is necessary for the property Pr_λ ; see Fact 1.5. However, the argument above does not extend to singular cardinals.

Our main contribution is to show that the failure of Pr_λ at singular cardinals is consistent with Gödel's axiom of constructibility. Although the main theorem is established in a more general setting, the following special case suffices for the purposes of the introduction.

Theorem 0.1. *Assume $\mathbf{V} = \mathbf{L}$ and that:*

- (a) $\langle \lambda_i : i < \kappa \rangle$ is an increasing sequence of regular cardinals with limit λ ;
- (b) each $\lambda_i = \mu_i^+$ is a successor cardinal, where μ_i is regular and not weakly compact;
- (c) $\aleph_0 < \kappa \leq \chi < \lambda_0$, where κ and χ are regular.

Then there exists a χ -free abelian group G of cardinality λ which is a counterexample to singular compactness at λ for the property $\text{Hom}(-, \mathbb{Z}) \neq 0$.

In fact, the assumptions of Theorem 0.1 can be weakened as follows:

- (a) $\langle \lambda_i : i < \kappa \rangle$ is an increasing sequence of regular cardinals with limit λ , with each $\lambda_i = \mu_i^+$;
- (b) $\aleph_0 < \kappa \leq \chi < \lambda_0$, where κ and χ are regular;
- (c) for each $i < \kappa$, there exist stationary, non-reflecting sets $S_i \subseteq S_{\aleph_0}^{\lambda_i}$ and $T_i \subseteq S_{\aleph_0}^{\mu_i}$ such that \diamond_{S_i} and \diamond_{T_i} hold;
- (d) there is no measurable cardinal $\leq \lambda$.

Our investigation is closely related to the classical Whitehead problem, concerning the vanishing of $\text{Ext}(G, \mathbb{Z}) := \text{Ext}_{\mathbb{Z}}^1(G, \mathbb{Z})$, a property of considerable interest but greater complexity. In our forthcoming work [1], we study singular compactness in the context of Ext . Under the additional assumption $\mathbf{V} = \mathbf{L}$, the situation becomes simpler: by [8], an abelian group G of cardinality $\lambda > \aleph_0$ is free if and only if $\text{Ext}(G, \mathbb{Z}) = 0$. Hence, by Shelah's singular compactness theorem for free groups [9], singular compactness also holds for the property $\text{Ext}(-, \mathbb{Z}) = 0$.

Throughout the paper, all groups are abelian. For standard terminology from set-theoretic algebra, we refer to Eklof–Mekler [3] and Göbel–Trlifaj [5]. Background from group theory may be found in Fuchs [4].

§ 1. PRELIMINARIES

In this section we establish the notation and conventions used throughout the paper, and recall several basic facts from set-theoretic algebra. All groups are abelian and written additively.

For abelian groups G and H , we write $\text{Hom}(G, H) := \text{Hom}_{\mathbb{Z}}(G, H)$.

Notation 1.1. Let u be an index set, and define $\mathbb{Z}_{[u]} := \bigoplus_{\alpha \in u} \mathbb{Z}x_{\alpha}$, so that $\langle x_{\alpha} : \alpha \in u \rangle$ is a basis for $\mathbb{Z}_{[u]}$.

Definition 1.2. Let κ be an infinite cardinal and G be an abelian group.

- (a) The group G is called κ -free if every subgroup of G of cardinality less than κ is free.
- (b) The group G is said to be strongly κ -free if it is κ -free and in addition every subset of G of cardinality $< \kappa$ is contained in a subgroup H of G of cardinality $< \kappa$ such that G/H is κ -free.

Remark 1.3. We note that if κ is a singular cardinal, then G is κ -free if and only if G is λ -free for every regular cardinal $\lambda < \kappa$.

Recall that a ring R is said *left-perfect* if every flat left R -module is projective. Since \mathbb{Q} is flat but not projective, it follows that \mathbb{Z} is not left-perfect. Moreover, the following result shows that for a weakly compact cardinal κ , every κ -free group of cardinality κ is free.

Fact 1.4. (See [3, Theorems VI.3.2 and VII.1.4].) Let λ be a weakly compact cardinal and let R be a non-left-perfect ring. If M is a $\leq \lambda$ -generated module which is λ -free, then M is free.

We now verify that Pr_{λ} holds whenever λ is weakly compact.

Fact 1.5. Assume λ is a weakly compact cardinal and G is a group of size λ such that $\text{Hom}(G', \mathbb{Z}) \neq 0$ for every nontrivial subgroup $G' \subseteq G$ of cardinality $< \lambda$. Then $\text{Hom}(G, \mathbb{Z}) \neq 0$. In particular, Pr_{λ} holds.

Proof. Let $\theta > \lambda$ be sufficiently large and regular, and let M be an elementary submodel of $(\mathcal{H}(\theta), \in)$ of cardinality λ such that

$$M^{<\lambda} \subseteq M \quad \text{and} \quad \lambda, G \in M.$$

Since λ is weakly compact, there exists a model N of cardinality λ with $N^{<\lambda} \subseteq N$ and an elementary embedding $j : M \rightarrow N$ such that

$$j, M \in N, \quad j(\alpha) = \alpha \text{ for all } \alpha < \lambda, \quad j(\lambda) > \lambda.$$

In N , the group $j(G)$ has cardinality $j(\lambda)$, and G is a subgroup of $j(G)$ of size $\lambda < j(\lambda)$. By elementarity,

$$N \models \text{“}\text{Hom}(G, \mathbb{Z}) \neq 0\text{”}.$$

This implies $\text{Hom}(G, \mathbb{Z}) \neq 0$ in the ambient universe, and the claim follows. \square

Definition 1.6. Suppose $\lambda > \mu \geq \aleph_0$ are regular cardinals, and let $S \subseteq \lambda$ be stationary.

- (1) *Jensen's diamond* $\diamond_\lambda(S)$ asserts the existence of a sequence $(S_\alpha \mid \alpha \in S)$ such that for every $X \subseteq \lambda$, the set

$$\{\alpha \in S \mid X \cap \alpha = S_\alpha\}$$

is stationary.

- (2) A useful consequence of $\diamond_\lambda(S)$ is the following. Let $A = \bigcup_{\alpha < \lambda} A_\alpha$ and $B = \bigcup_{\alpha < \lambda} B_\alpha$ be two λ -filtrations with $|A_\alpha|, |B_\alpha| < \lambda$. Then there exists a sequence of functions $(g_\alpha : A_\alpha \rightarrow B_\alpha \mid \alpha < \lambda)$ such that, for any function $g : A \rightarrow B$, the set $\{\alpha \in S \mid g \upharpoonright_{A_\alpha} = g_\alpha\}$ is stationary in λ .
- (3) S is *non-reflecting* if for every limit ordinal $\delta < \lambda$ of uncountable cofinality, the set $S \cap \delta$ is non-stationary in δ .
- (4) We set $S_\mu^\lambda := \{\alpha < \lambda \mid \text{cf}(\alpha) = \mu\}$.

Definition 1.7. Let \mathcal{K} be the class of all uncountable cardinals μ so that if $K_1 \subseteq \mathbb{Z}$ is a subgroup, then there is (H_{K_1}, η_{K_1}) equipped with the following properties:

- (α) H_{K_1} is an abelian group extending $\mathbb{Z}_{[\mu]}$,
- (β) $H_{K_1}/\mathbb{Z}_{[\mu]}$ is μ -free,
- (γ) $\eta_{K_1} \in {}^\mu(K_1)$,
- (δ) there is no homomorphism $f : H_{K_1} \rightarrow K_1$ such that $f(x_\alpha) = \eta_{K_1}(\alpha)$ for $\alpha < \mu$:

$$\begin{array}{ccccc} \mathbb{Z}_{[\{\alpha\}] } & \xrightarrow{\subseteq} & \mathbb{Z}_{[\mu]} & \xrightarrow{\subseteq} & H_{K_1} \\ & & & \searrow \exists f & \\ & & & & K_1 \\ & \tilde{\eta} \downarrow & & & \\ & & & & \end{array}$$

where $\tilde{\eta}(x_\alpha) := \eta_{K_1}(\alpha)$.

We now give sufficient conditions for a cardinal μ being in \mathcal{K} . We first recall the following result of Eklof–Mekler, see [2].

Fact 1.8. Let μ be an uncountable regular cardinal. Assume that $S \subseteq S_{\aleph_0}^\mu$ is stationary and non-reflecting, and that \diamond_S holds. Then there exists an indecomposable, strongly μ -free abelian group H of cardinality μ .

Lemma 1.9. *We have $\mu \in \mathcal{K}$ provided that:*

- (a) $\mu = \text{cf}(\mu) > \aleph_0$;
- (b) $S \subseteq S_{\aleph_0}^\mu$ is stationary and non-reflecting;
- (c) \diamond_S holds.

Proof. By Fact 1.8, there exists an indecomposable, strongly μ -free abelian group H of cardinality μ . We first note that $\text{Hom}(H, \mathbb{Z}) = 0$. Indeed, if $0 \neq f \in \text{Hom}(H, \mathbb{Z})$, then

$\text{Im}(f) \cong \mathbb{Z}$ is free, so the exact sequence

$$0 \longrightarrow \ker(f) \longrightarrow H \longrightarrow \text{Im}(f) \longrightarrow 0$$

splits. Since H is indecomposable, this forces $\ker(f) = 0$, hence H embeds into \mathbb{Z} , contradicting $|H| = \mu > \aleph_0$. As H is strongly μ -free, there exists a subgroup $K \subseteq H$ of cardinality $< \mu$ such that $\mathbb{Z} \subseteq K$ and H/K is μ -free. Without loss of generality, write $K = \bigoplus_{i < \theta} \mathbb{Z}$ for some $\theta < \mu$. Fix a partition $\mu = \bigcup_{i < \mu} I_i$ such that the I_i are pairwise disjoint, $|I_i| = \theta$ for all $i < \mu$, and $I_i < I_j$ whenever $i < j$. Define

$$G := \bigoplus_{i < \mu} G_i,$$

where each $G_i \cong H$ and G_i/\mathbb{Z}_{I_i} is μ -free. Since H is torsion-free, we have $\mathbb{Z} \subseteq H$, and hence

$$\mathbb{Z}_{[\mu]} \subseteq \bigoplus_{i < \mu} G_i = G.$$

Moreover,

$$\text{Hom}(G, \mathbb{Z}) \cong \text{Hom}\left(\bigoplus_{i < \mu} G_i, \mathbb{Z}\right) \cong \prod_{i < \mu} \text{Hom}(H, \mathbb{Z}) = 0.$$

We next verify that G is μ -free. Let $L \leq G$ with $|L| < \mu$. There exists $I \subseteq \mu$ with $|I| < \mu$ and subgroups $L_i \leq G_i$, each of size $< \mu$, such that $L \subseteq \bigoplus_{i \in I} L_i$. Since each L_i is free, so is $\bigoplus_{i \in I} L_i$, and hence L is free.

Finally, note that $G/\mathbb{Z}_{[\mu]} \cong \bigoplus_{i < \mu} G_i/\mathbb{Z}_{I_i}$, and each summand on the right-hand side is μ -free. Thus $G/\mathbb{Z}_{[\mu]}$ is μ -free. Therefore $\mu \in \mathcal{K}$. For any $K_1 \subseteq \mathbb{Z}$, set $(H_{K_1}, \eta_{K_1}) = (G, \eta)$, where $\eta: \mu \rightarrow K_1$ is arbitrary. If $K_1 \neq 0$, then $K_1 \cong \mathbb{Z}$, and hence $\text{Hom}(G, K_1) = 0$, which verifies (δ) from Definition 1.7. \square

We also need the following well-known result of Kurepa.

Fact 1.10. Assume $\text{cf}(\lambda) > \aleph_0$ and let \mathcal{T} be a tree of height λ whose levels are all finite. Then \mathcal{T} has a cofinal branch.

§ 2. CONTROLLING $\text{Hom}(G, \mathbb{Z})$

In this section we prove our main result (see Theorem 2.6).

Discussion 2.1. Recall that a cardinal κ is *measurable* if it is uncountable and there exists a non-principal κ -complete ultrafilter \mathcal{D} on κ , meaning that for every subset $S \subseteq \mathcal{D}$ of cardinality less than κ , the intersection $\bigcap S$ belongs to \mathcal{D} . It is a classical result that the existence of measurable cardinals cannot be established within ZFC.

Definition 2.2. Let G be an abelian group. The *dual* of G is the abelian group $\text{Hom}(G, \mathbb{Z})$, denoted by G^* . For $g \in G$, define $\psi_g: G^* \rightarrow \mathbb{Z}$ by evaluation

$$\psi_g(f) := f(g), \quad f \in G^*.$$

The assignment $g \mapsto \psi_g$ defines a canonical map $\psi : G \rightarrow G^{**}$. We say that G is *reflexive* if ψ is an isomorphism.

Fact 2.3. (Lős–Eda, Shelah; see [3, Corollary III.1.5] and [11]). Let $\mu = \mu_{\text{first}}$ be the first measurable cardinal. The following hold:

- (a) For any $\theta < \mu$, the group $\mathbb{Z}^{(\theta)}$ is reflexive; in fact, its dual is \mathbb{Z}^θ .
- (b) For any $\lambda \geq \mu$, the group $\mathbb{Z}^{(\lambda)}$ is not reflexive.
- (c) There exists a reflexive group $G \subset \mathbb{Z}^\mu$ of cardinality μ .

Let Pr be any property of abelian groups, and let λ be a cardinal. Recall that *compactness for* (λ, Pr) means that if G is a group of cardinality λ and

$$\text{“for all } G' \subseteq G \text{ with } |G'| < \lambda, G' \text{ has Pr”},$$

then G itself has Pr . In this paper, we focus on the following specific property of abelian groups:

Notation 2.4. For a cardinal λ , let Pr_λ denote the property: If G is a group of size λ , and if for any nontrivial subgroup $G' \subseteq G$ of size less than λ , $\text{Hom}(G', \mathbb{Z}) \neq 0$, then $\text{Hom}(G, \mathbb{Z}) \neq 0$.

We now turn to the primary framework for our construction.

Definition 2.5. Let θ be a cardinal.

- (1) Let $\mathbf{M}_{1,\theta}$ be the class of objects

$$\mathbf{m} = (\lambda_{\mathbf{m}}, \langle G_\alpha^{\mathbf{m}} : \alpha \leq \alpha_{\mathbf{m}} \rangle, S_{\mathbf{m}}, \langle f_{\mathbf{m},s} : s \in S_{\mathbf{m}} \rangle)$$

consisting of:

- (a) (α) $\lambda_{\mathbf{m}} = \text{cf}(\lambda_{\mathbf{m}}) > \aleph_0$,
 (β) $\lambda_{\mathbf{m}} \geq \alpha_{\mathbf{m}} := \ell g(\mathbf{m})$, where $\ell g(\mathbf{m})$ denotes the length of \mathbf{m} .
- (b) (α) $\langle G_\alpha^{\mathbf{m}} : \alpha \leq \alpha_{\mathbf{m}} \rangle$ is an increasing and continuous sequence of abelian groups,
 (β) $|G_\alpha^{\mathbf{m}}| < \lambda_{\mathbf{m}}$ for $\alpha < \alpha_{\mathbf{m}}$.
- (c) $G_\alpha^{\mathbf{m}}/G_0^{\mathbf{m}}$ is free.
- (d) The set

$$\{\beta < \alpha_{\mathbf{m}} : G_{\beta+1}^{\mathbf{m}}/G_\beta^{\mathbf{m}} \text{ is not free}\}$$

is non-reflecting and stationary.

- (e) (α) $S_{\mathbf{m}}$ is a set of cardinality $\leq \theta$,
 (β) $f_{\mathbf{m},s} \in \text{Hom}(G_{\alpha_{\mathbf{m}}}^{\mathbf{m}}, \mathbb{Z})$ for each $s \in S_{\mathbf{m}}$.
 - (f) The family $\langle f_{\mathbf{m},s} : s \in S_{\mathbf{m}} \rangle$ is a free basis of a subgroup of $\text{Hom}(G_{\alpha_{\mathbf{m}}}^{\mathbf{m}}, \mathbb{Z})$.
- (2) The class $\mathbf{M}_{2,\theta}$ is defined as above, with the modification that in item (a)(β) we set $\alpha_{\mathbf{m}} = \lambda_{\mathbf{m}}$, and we further require:
- (g) For every $f \in \text{Hom}(G_{\alpha_{\mathbf{m}}}^{\mathbf{m}}, \mathbb{Z})$, there exists $h \in \text{Hom}(\prod_{s \in S_{\mathbf{m}}} \mathbb{Z}, \mathbb{Z})$ such that

$$f(x) = h(\langle f_{\mathbf{m},s}(x) : s \in S_{\mathbf{m}} \rangle) \quad \text{for all } x \in G_{\alpha_{\mathbf{m}}}^{\mathbf{m}}.$$

Namely, h is a factorization of f through

$$\pi \in \text{Hom}(G_{\alpha_{\mathbf{m}}}^{\mathbf{m}}, \mathbb{Z}^{S_{\mathbf{m}}}),$$

defined by $\pi(x) := (f_{\mathbf{m},s}(x) \mid s \in S_{\mathbf{m}})$.

- (h) The homomorphism π is surjective.
- (i) For any nonzero subgroup $G' \subseteq G_{\alpha}^{\mathbf{m}}$ with $\alpha < \alpha_{\mathbf{m}}$, we have $\text{Hom}(G', \mathbb{Z}) \neq 0$.

We are now in a position to state and prove our main result:

Theorem 2.6. *Assume that:*

- (a) $\langle \lambda_i : i < \kappa \rangle$ is an increasing sequence of regular cardinals with limit λ , and $\lambda_i = \mu_i^+$, for some regular cardinal μ_i ,
- (b) $\aleph_0 < \kappa \leq \chi < \lambda_0$ and κ, χ are regular,
- (c) $S_i \subseteq S_{\aleph_0}^{\lambda_i}$ is stationary and non-reflecting, and \diamond_{S_i} holds,
- (d) $T_i \subseteq S_{\aleph_0}^{\mu_i}$ is stationary and non-reflecting, and \diamond_{T_i} holds,
- (e) there is no measurable cardinal $\leq \lambda$.

Then there is a χ -free abelian group G of cardinality λ which is counterexample to singular compactness in λ for Pr_{λ} .

Remark 2.7. The sets T_i and S_i have cardinality μ_i and λ_i , respectively. In particular, they are different, though both of them have concentrated on ordinals of countable cofinality.

Proof. We construct a χ -free abelian group G of cardinality λ endowed with the following property: for any nontrivial subgroup $G' \subseteq G$ of smaller cardinality, we have

$$\text{Hom}(G', \mathbb{Z}) \neq 0,$$

while $\text{Hom}(G, \mathbb{Z}) = 0$.

The idea of the construction is to equip the cardinals λ_i with an underlying tree structure \mathcal{T}_i , which allows precise control of homomorphisms from the building blocks G_i of the group

$$G := \bigcup_{i < \kappa} G_i.$$

The proof is organized in four stages. Stage (A) defines the tree structure \mathcal{T}_i . Stage (B), which is more involved, constructs by induction a system of χ -free groups together with homomorphisms from them to \mathbb{Z} . Stage (C) verifies that subgroups of smaller cardinality have nontrivial duals, and Stage (D) shows that there are no non-zero homomorphisms from G to \mathbb{Z} .

Stage (A): We define a tree \mathcal{T} of height κ , whose i -th level \mathcal{T}_i is defined as follows:

- (*)_Aⁱ : \mathcal{T}_i is the set of all sequences η satisfying:
 - (a) η has length $i + 1$,
 - (b) for each $j \leq i$, $\eta(j) = (\eta(j, 1), \eta(j, 2))$,
 - (c) for each $j \leq i$, $\eta(j, 1) < \lambda_j$ and $\eta(j, 2) < \kappa$,

- (d) if $j_1 < j_2 \leq i$, then $\eta(j_1, 1) \leq \eta(j_2, 1)$ and $\eta(j_1, 2) \leq \eta(j_2, 2)$,
- (e) the range of η is finite,
- (f) if $j_1 < j_2 \leq i$ and the sequence $\langle \eta(j, 1) : j \in [j_1, j_2] \rangle$ is constant, then $j_2 < \eta(j_1, 2)$.

Set $\mathcal{T} := \bigcup_{i < \kappa} \mathcal{T}_i$, ordered by the end-extension relation \triangleleft . Then $(\mathcal{T}, \triangleleft)$ is a tree of height κ , whose i -th level is \mathcal{T}_i . Moreover, if $\eta \in \mathcal{T}_i$ and $i < j < \kappa$, there exists $\nu \in \mathcal{T}_j$ such that $\eta \triangleleft \nu$; that is, $\eta = \nu \upharpoonright (i+1)$.

For truncated trees, we define, for $\alpha \leq \lambda_i$,

$$\mathcal{T}_{i,\alpha} := \{\eta \in \mathcal{T}_i : \eta(i, 1) \leq \alpha\}.$$

In particular, $\mathcal{T}_i = \mathcal{T}_{i,\lambda_i}$.

Claim 2.8. *The tree $(\mathcal{T}, \triangleleft)$ has no branches of length κ .*

Proof. Assume, towards a contradiction, that there exists a branch

$$b := \langle \eta_i : i < \kappa \rangle$$

of \mathcal{T} . Then $\langle \eta_i : i < \kappa \rangle$ is \triangleleft -increasing. It follows that the sequence

$$\langle \eta_i(i, 1) : i < \kappa \rangle$$

is non-decreasing in the ordinals. By clause $(*)_A^i(e)$, every initial segment takes only finitely many values, and since $\kappa = \text{cf}(\kappa) > \aleph_0$, the sequence must eventually stabilize. That is, there exists $i_* < \kappa$ such that

$$\eta_i(i, 1) = \eta_{i_*}(i_*, 1) \quad \text{for all } i \in [i_*, \kappa).$$

On the one hand, by $(*)_A^i(f)$, we have

$$\eta(i_*, 2) > i \quad \text{for all } i < \kappa,$$

while on the other hand, $\eta(i_*, 2) < \kappa$. This is a contradiction. □_{2.8}

Stage (B): We define \mathbf{m}_i by induction on $i < \kappa$ such that:

- $(*)_B^i$: The sequence \mathbf{m}_i satisfies:
 - (a) $\mathbf{m}_i = (\lambda_{\mathbf{m}_i}, \langle G_\alpha^{\mathbf{m}_i} : \alpha \leq \alpha_{\mathbf{m}_i} \rangle, S_{\mathbf{m}_i}, \langle f_{\mathbf{m}_i, s} : s \in S_{\mathbf{m}_i} \rangle) \in \mathbf{M}_{1, \lambda_i}$,
 - (b) $\lambda_{\mathbf{m}_i} = \alpha_{\mathbf{m}_i} = \lambda_i$, and the set of elements of $G_{\lambda_i}^{\mathbf{m}_i}$ has cardinality λ_i ,
 - (c) $G_{< i} := \bigcup \{G_{\lambda_j}^{\mathbf{m}_j} : j < i\} \cup \{0\}$,
 - (d) $G_0^{\mathbf{m}_i} := G_{< i}$,
 - (e) $S_{\mathbf{m}_i} := \mathcal{T}_i = \mathcal{T}_{i, \lambda_i}$,
 - (f) if $j < i$, then $\mathbf{m}_j \leq \mathbf{m}_i$, i.e.,

$$\eta \in \mathcal{T}_j \wedge \nu \in \mathcal{T}_i \wedge \eta \triangleleft \nu \implies f_{\mathbf{m}_j, \eta} \subseteq f_{\mathbf{m}_i, \nu},$$

which can be depicted as:

$$\begin{array}{ccc} 0 & \longrightarrow & G_{\lambda_j}^{\mathbf{m}_j} \xrightarrow{\subseteq} G_{\lambda_i}^{\mathbf{m}_i} \\ & & \downarrow f_{\mathbf{m}_j, \eta} \quad \swarrow f_{\mathbf{m}_i, \nu} \\ & & \mathbb{Z} \end{array}$$

- (g) $\langle f_{\mathbf{m}_i, \eta} : \eta \in \mathcal{T}_i \rangle$ is an independent subset of $\text{Hom}(G_{\lambda_i}^{\mathbf{m}_i}, \mathbb{Z})$,
- (h) $\bigcap \{\text{Ker}(f_{\mathbf{m}_i, \eta}) : \eta \in \mathcal{T}_i\} = \{0\}$,
- (i) for any $f \in \text{Hom}(G_{\lambda_i}^{\mathbf{m}_i}, \mathbb{Z})$, there exist $\alpha < \lambda_i$ and $h \in \text{Hom}(\prod_{\eta \in \mathcal{T}_{i, \alpha}} \mathbb{Z}, \mathbb{Z})$ such that h is a factorization of f through $\pi_i \in \text{Hom}(G_{\alpha}^{\mathbf{m}_i}, \mathbb{Z}^{S_{\mathbf{m}_i}})$, defined by $\pi_i(x) = (f_{\mathbf{m}_i, s}(x) \mid s \in S_{\mathbf{m}_i})$.

Remark 2.9. For each $\alpha < \lambda_i$, the cardinality of $\mathcal{T}_{i, \alpha}$ is less than λ_i . This fact will be useful to show that

$$|G_{\alpha}^{\mathbf{m}_j}| < \lambda_{\mathbf{m}_j} \quad \text{for } \alpha < \lambda_{\mathbf{m}_j},$$

see the subsequent discussion in (§) below.

Assume that $i < \kappa$ and that the sequence $\langle \mathbf{m}_j : j < i \rangle$ has been defined. Fix a diamond sequence

$$\langle F_{i, \delta} : \delta \in S_i \rangle, \quad F_{i, \delta} : \delta \rightarrow \mathbb{Z}.$$

Notation 2.10. Let $\langle \beta_i(\gamma) : \gamma < \lambda_i \rangle$ be an increasing, continuous sequence of ordinals cofinal in λ_i , with $\beta_i(0) = 0$.

We proceed by setting $G_{< i} := \bigcup_{j < i} G_{\lambda_j}^{\mathbf{m}_j} \cup \{0\}$. Also, for $\eta \in \mathcal{T}_i$, define $f_{< i, \eta} : G_{< i} \rightarrow \mathbb{Z}$ by

$$f_{< i, \eta} := \bigcup_{j < i} f_{\mathbf{m}_j, \eta \upharpoonright (j+1)}.$$

In particular, $G_{< 0} = \{0\}$ and $f_{< 0, \eta} : G_{< 0} \rightarrow \mathbb{Z}$ is the zero map. We shall choose $\mathbf{m}_{i, \gamma}$ by induction on $\gamma < \lambda_i$ such that:

- (*) $^{\gamma}_C$: (a) $\mathbf{m}_{i, 0}$ is defined as
 - (α) $lg(\mathbf{m}_{i, 0}) = 0$,
 - (β) $\lambda_{\mathbf{m}_{i, 0}} := \chi + \sup_{j < i} \lambda_{\mathbf{m}_j}$ with the convention that $\sup_{j < 0} \lambda_{\mathbf{m}_j} = 0$
 - (γ) $G_0^{\mathbf{m}_{i, 0}} := G_{< i}$,
 - (δ) $S_{\mathbf{m}_{i, 0}} := \mathcal{T}_{i, \beta_i(0)}$,
 - (ϵ) for $\eta \in \mathcal{T}_{i, \beta_i(0)}$, $f_{\mathbf{m}_{i, 0}, \eta} := f_{< i, \eta}$.
- (b) $\langle \mathbf{m}_{i, \gamma} : \gamma < \lambda_i \rangle$ is an increasing and continuous sequence from $\mathbf{M}_{1, \lambda_i}$ with $S_{\mathbf{m}_{i, \gamma}} = \mathcal{T}_{i, \beta_i(\gamma)}$ and $lg(\mathbf{m}_{i, \gamma}) = \alpha_{i, \gamma} < \lambda_i$, which means:
 - (α) if $\rho < \gamma$, then $\mathbf{m}_{i, \rho} \leq \mathbf{m}_{i, \gamma}$,
 - (β) if γ is a limit ordinal, then $\mathbf{m}_{i, \gamma} := \bigcup_{\rho < \gamma} \mathbf{m}_{i, \rho}$. In particular, we have the following equalities:
 - (β_1) $\alpha_{i, \gamma} = \sup_{\rho < \gamma} \alpha_{i, \rho}$,
 - (β_2) $G_{\alpha_{i, \gamma}}^{\mathbf{m}_{i, \gamma}} = \bigcup_{\rho < \gamma} G_{\alpha_{i, \rho}}^{\mathbf{m}_{i, \rho}}$,

- (β_3) $S_{\mathbf{m}_{i,\gamma}} = \mathcal{T}_{i,\beta_i(\gamma)}$,
 (β_4) $f_{\mathbf{m}_{i,\gamma},\eta} = f_{i,<\eta} \cup \bigcup_{\rho < \gamma} f_{\mathbf{m}_{i,\rho},\eta \upharpoonright \rho+1}$ for any $\eta \in \mathcal{T}_{i,\beta_i(\gamma)}$.
 (c) If $\rho < \gamma$, and $\rho \notin S_i$, then $G_{\alpha_{i,\gamma}}^{\mathbf{m}_{i,\gamma}}/G_{\alpha_{i,\rho}}^{\mathbf{m}_{i,\rho}}$ is free.
 (d) $\bigcap \{\text{Ker}(f_{\mathbf{m}_{i,\gamma},\eta}) : \eta \in \mathcal{T}_{i,\beta_i(\gamma)}\} = \{0\}$.
 (e) $G_{\alpha_{i,\gamma}}^{\mathbf{m}_{i,\gamma}}$ has set of elements an ordinal $\delta_i(\gamma) < \lambda_i$.
 (f) Recall that $\langle F_{i,\delta} : \delta \in S_i \rangle$ denotes the diamond sequence. We now collect the following notations and assumptions:

- (α) $\gamma = \alpha_{i,\gamma} \in S_i$,
 (β) The underlying set of $G_{\alpha_{i,\gamma}}^{\mathbf{m}_{i,\gamma}}$ is γ ,
 (γ) $\text{Im}(F_{i,\gamma}) \subseteq \mathbb{Z}$ is nonzero; in particular, $\text{Im}(F_{i,\gamma}) = n\mathbb{Z} \cong \mathbb{Z}$ for some nonzero $n \in \mathbb{Z}$,
 (δ) $F_{i,\gamma}$ is a homomorphism from $G_{\alpha_{i,\gamma}}^{\mathbf{m}_{i,\gamma}}$ onto $\text{Im}(F_{i,\gamma})$,
 (ϵ) $F_{i,\gamma} \notin \langle f_{\mathbf{m}_{i,\gamma},\eta} : \eta \in \mathcal{T}_{i,\beta_i(\gamma)} \rangle$, where $\langle f_{\mathbf{m}_{i,\gamma},\eta} : \eta \in \mathcal{T}_{i,\beta_i(\gamma)} \rangle$ denotes the subgroup of $\text{Hom}(G_{\alpha_{i,\gamma}}^{\mathbf{m}_{i,\gamma}}, \mathbb{Z})$ generated by $\{f_{\mathbf{m}_{i,\gamma},\eta} : \eta \in \mathcal{T}_{i,\beta_i(\gamma)}\}$.

We then define $\mathbf{m}_{i,\gamma+1}$ so that $F_{i,\gamma}$ has no extension to a homomorphism from $G_{\alpha_{i,\gamma+1}}^{\mathbf{m}_{i,\gamma+1}}$ into K_1 . Namely, we have the following commutative diagram:

$$\begin{array}{ccc} G_{\alpha_{i,\gamma}}^{\mathbf{m}_{i,\gamma}} & \xrightarrow{\subseteq} & G_{\alpha_{i,\gamma+1}}^{\mathbf{m}_{i,\gamma+1}} \\ F_{i,\gamma} \downarrow & & \downarrow \nexists \\ \text{Im}(F_{i,\gamma}) & \xrightarrow{=} & \text{Im}(F_{i,\gamma}), \end{array}$$

where the right vertical arrow does not exist. To simplify notation, we let $G_{i,\gamma} := G_{\alpha_{i,\gamma}}^{\mathbf{m}_{i,\gamma}}$. We then define $G_{i,\gamma,\rho} := G_{i,\gamma}$ for any $\rho \leq \alpha_{i,\gamma}$. Also, we abbreviate the function $f_{\mathbf{m}_{i,\gamma},\eta}$ as $f_{i,\gamma,\eta}$ for any index $\eta \in \mathcal{T}_{i,\beta_i(\gamma)}$.

The starting case $\gamma = 0$ is trivial, since it can be defined as in $(*)_C^\gamma$ (a). By the induction hypothesis and the definition of $f_{i,0,\eta}$, we have:

- The sequence $\langle f_{i,0,\eta} : \eta \in \mathcal{T}_{i,\beta_i(0)} \rangle$ is an independent subset of $\text{Hom}(G_{i,0}, \mathbb{Z})$,
- Each $f_{i,0,\eta}$ extends $f_{<i,\eta}$,
- $\bigcap \{\text{Ker}(f_{i,0,\eta}) : \eta \in \mathcal{T}_{i,\beta_i(0)}\} = \{0\}$.

If γ is a limit ordinal, set $\alpha_{i,\gamma} = \sup_{\rho < \gamma} \alpha_{i,\rho}$ and define $\mathbf{m}_{i,\gamma}$ as in $(*)_C^\gamma$ (b)(β).

Suppose that $\mathbf{m}_{i,\gamma}$ has already been defined. We now proceed to define $\mathbf{m}_{i,\gamma+1}$. First, assume that one of the following cases occurs:

- (h_1) $\gamma \notin S_i$ or at least one of the hypotheses $(*)_C^\gamma$ (f)(α)-(δ) fails, or
 (h_2) $\gamma \in S_i$, the hypotheses $(*)_C^\gamma$ (i)(α)-(δ) hold, but either
 – $G_{i,\gamma}$ does not have domain γ , or
 – $F_{i,\gamma} \notin \text{Hom}(G_{i,\gamma}, \mathbb{Z})$, or
 – $F_{i,\gamma} \in \langle f_{i,\gamma,\eta} : \eta \in \mathcal{T}_{i,\beta_i(\gamma)} \rangle$.

In this case, define $\mathbf{m}_{i,\gamma+1}$ as follows:

- (1) $\alpha_{i,\gamma+1} := \alpha_{i,\gamma} + 1$,
- (2) $\mathbf{m}_{i,\gamma} \leq \mathbf{m}_{i,\gamma+1}$,

- (3) $S_{\mathbf{m}_{i,\gamma+1}} := \mathcal{T}_{i,\beta_i(\gamma+1)}$,
- (4) $G_{i,\gamma+1} := G_{i,\gamma+1,\alpha_{i,\gamma+1}} := G_{i,\gamma} \oplus \mathbb{Z}_{[u_{i,\gamma}]}$, where $u_{i,\gamma} := \mathcal{T}_{i,\beta_i(\gamma+1)}$,
- (5) For $\eta \in \mathcal{T}_{i,\beta_i(\gamma+1)}$, set $f_{i,\gamma+1,\eta} := f_{i,\gamma,\eta} \oplus \pi_\eta$, where $\pi_\eta : \mathbb{Z}_{[u_{i,\gamma}]} \rightarrow \mathbb{Z}x_\eta$ is the projection, and for $\eta \notin \mathcal{T}_{i,\beta_i(\gamma)}$, $f_{i,\gamma,\eta}$ is the zero map.

Then suppose that $\mathbf{m}_{i,\gamma}$ is defined and the following holds:

- (h_3) $\gamma \in S_i$, $G_{i,\gamma}$ has domain γ , hypotheses $(*)_C^\gamma(\mathbf{f})(\alpha)$ - (δ) hold, $F_{i,\gamma} \in \text{Hom}(G_{i,\gamma}, \mathbb{Z})$, and $F_{i,\gamma} \notin \langle f_{i,\gamma,\eta} : \eta \in \mathcal{T}_{i,\beta_i(\gamma)} \rangle$.

In this case, we define $\mathbf{m}_{i,\gamma+1}$ so as to satisfy $(*)_B^\gamma(\mathbf{i})$. Let $\alpha_{i,\gamma+1} := \alpha_{i,\gamma} + 1$ and set

$$G_\rho^{\mathbf{m}_{i,\gamma+1}} := G_\rho^{\mathbf{m}_{i,\gamma}} = G_{i,\gamma,\rho}, \quad \forall \rho \leq \alpha_{i,\gamma}.$$

It remains to define

$$G_{i,\gamma+1} = G_{\alpha_{i,\gamma+1}}^{\mathbf{m}_{i,\gamma+1}} \quad \text{and} \quad f_{i,\gamma+1,\eta} = f_{\mathbf{m}_{i,\gamma+1},\eta} : G_{i,\gamma+1} \rightarrow \mathbb{Z}, \quad \eta \in \mathcal{T}_{i,\beta_i(\gamma+1)}.$$

For each $\beta < \lambda_i$, define

$$G_{i,\gamma}^{[\beta]} := \{x \in G_{i,\gamma} : \eta \in \mathcal{T}_i \wedge \eta(i, 1) < \beta \implies f_{i,\gamma,\eta}(x) = 0\}.$$

Then $\langle G_{i,\gamma}^{[\beta]} : \beta < \lambda_i \rangle$ is increasing, and since $|G_{i,\gamma}| < \lambda_i$, there exists $\beta_{i,\gamma} < \lambda_i$ such that

$$G_{i,\gamma}^{[\beta]} = G_{i,\gamma}^{[\beta_{i,\gamma}]}, \quad \forall \beta \in (\beta_{i,\gamma}, \lambda_i).$$

Set

$$K_{i,\gamma} := \text{Im}(F_{i,\gamma} \upharpoonright G_{i,\gamma}^{[\beta_{i,\gamma}]})$$

and note that

$$\chi + \sum_{j < i} \lambda_j + |\mathcal{T}_{i,\beta_i(\gamma+1)}| + \aleph_0 < \lambda_i.$$

Consequently,

$$\chi + \sum_{j < i} \lambda_j + |\mathcal{T}_{i,\beta_i(\gamma+1)}| + \aleph_0 \leq \mu_i.$$

We set $\mu_{i,\gamma} := \mu_i$. Combining Lemma 1.9 with assumption 2.6(d), we obtain $\mu_{i,\gamma} \in \mathcal{K}$. Using Notation 1.1, define

$$(K_{i,\gamma})_{[u]} := \bigoplus_{\alpha \in u} K_{i,\gamma}x_\alpha \quad \text{for any index set } u.$$

Also, we define $(H_*, \phi_*) := (H_{K_{i,\gamma}}, \phi_{K_{i,\gamma}})$, so that

- H_* is a free abelian group of size $\mu_{i,\gamma}$ extending $(K_{i,\gamma})_{[u_{i,\gamma}]}$,
- $\phi_* : u_{i,\gamma} \rightarrow K_{i,\gamma} \setminus \{0\}$,
- $H_*/(K_{i,\gamma})_{[u_{i,\gamma}]}$ is $\mu_{i,\gamma}$ -free,
- there is no homomorphism $f : H_* \rightarrow K_{i,\gamma}$ such that $f(x_\eta) = \phi_*(\eta)$ for all $\eta \in \mathcal{T}_{i,\beta_i(\gamma+1)}$.

Finally, for each β with $\beta_{i,\gamma} \leq \beta < \lambda_i$ and $b \in K_{i,\gamma} = \text{Im}(F_{i,\gamma} \upharpoonright G_{i,\gamma}^{[\beta_{i,\gamma}]})$, there exists $y_{b,\beta} \in G_{i,\gamma}^{[\beta_{i,\gamma}]} \subseteq G_{i,\gamma}$ such that

- (*) _{β, b} (a) If $\eta \in \mathcal{T}_i$ and $\eta(i, 1) < \beta$, then $f_{i, \gamma, \eta}(y_{b, \beta}) = 0$,
 (b) $F_{i, \gamma}(y_{b, \beta}) = b$.

Since $|G_{i, \gamma}| < \lambda_i$, for each $b \in K_{i, \gamma}$ as above, there exists some fixed $y_b \in G_{i, \gamma}$ such that the set

$$X_b := \{\beta < \lambda_i : \beta_{i, \gamma} \leq \beta \text{ and } y_{b, \beta} = y_b\}$$

is stationary in λ_i .

The assignment $x_\eta \mapsto y_{\phi_*(\eta)}$ induces a morphism

$$g_{i, \gamma} : (K_{i, \gamma})_{[u_{i, \gamma}]} \longrightarrow G_{i, \gamma}.$$

Recall that $u_{i, \gamma} = \mathcal{T}_{i, \beta_i(\gamma+1)}$ and that

$$\text{id} : (K_{i, \gamma})_{[u_{i, \gamma}]} \longrightarrow H_{\mathbf{k}_{i, \gamma}, K_{i, \gamma}}$$

denotes the natural inclusion.

(‡) Define

$$G_{i, \gamma+1} := \frac{G_{i, \gamma} \times H_{\mathbf{k}_{i, \gamma}, K_{i, \gamma}}}{\langle (g_{i, \gamma}(k), -\text{id}(k)) : k \in (K_{i, \gamma})_{[u_{i, \gamma}]} \rangle}.$$

Recall that $H_{\mathbf{k}_{i, \gamma}, K_{i, \gamma}}$ is of size $\mu_{i, \gamma} < \lambda_i$. Combining this with an inductive argument along with (*), we conclude that $|G_{i, \gamma+1}| < \lambda_i$.

Notation 2.11. For $g \in G_{i, \gamma}$ and $h \in H_{\mathbf{k}_{i, \gamma}, K_{i, \gamma}}$, let $[(g, h)] \in G_{i, \gamma+1}$ denote the equivalence class of (g, h) .

By the definition of the push-out construction, this gives us two maps

$$h_{i, \gamma} : G_{i, \gamma} \longrightarrow G_{i, \gamma+1}, \quad k_{i, \gamma} : H_{\mathbf{k}_{i, \gamma}, K_{i, \gamma}} \longrightarrow G_{i, \gamma+1},$$

such that $h_{i, \gamma} \circ g_{i, \gamma} = k_{i, \gamma}$. In fact $h_{i, \gamma}$ is defined by $h_{i, \gamma}(x) := [(x, 0)]$. The situation is depicted in the following commutative diagram:

$$\begin{array}{ccc} H_{\mathbf{k}_{i, \gamma}, K_{i, \gamma}} & \xrightarrow{k_{i, \gamma}} & G_{i, \gamma+1} \\ \text{id} \uparrow & & \uparrow h_{i, \gamma} \\ (K_{i, \gamma})_{[u_{i, \gamma}]} & \xrightarrow{g_{i, \gamma}} & G_{i, \gamma} \end{array}$$

Claim 2.12. *Let $g \in G_{i, \gamma}$ and $h \in (K_{i, \gamma})_{[u_{i, \gamma}]}$. Then there exists an element $\tilde{g} \in G_{i, \gamma}$ such that $[(g, h)] = [(\tilde{g}, 0)]$ in $G_{i, \gamma+1}$.*

Proof. Set $\tilde{g} := g + g_{i, \gamma}(h) \in G_{i, \gamma}$. Then

$$(g, h) - (\tilde{g}, 0) = (g - \tilde{g}, h) = (-g_{i, \gamma}(h), h) \in \langle (g_{i, \gamma}(k), -\text{id}(k)) : k \in (K_{i, \gamma})_{[u_{i, \gamma}]} \rangle.$$

Hence $[(g, h)] = [(\tilde{g}, 0)]$ in $G_{i, \gamma+1}$. □_{2.12}

Clearly, the map $h_{i,\gamma} : G_{i,\gamma} \rightarrow G_{i,\gamma+1}$ is an embedding, and we may identify $G_{i,\gamma} \subseteq G_{i,\gamma+1}$ via $h_{i,\gamma}$. Let $g \in G_{i,\gamma}$ and $h \in H_{\mathbf{k}_{i,\gamma}, K_{i,\gamma}}$, and define

$$\psi : G_{i,\gamma+1} \longrightarrow H_{\mathbf{k}_{i,\gamma}, K_{i,\gamma}} / (K_{i,\gamma})_{[u_{i,\gamma}]}, \quad \psi([(g, h)]) := h + (K_{i,\gamma})_{[u_{i,\gamma}]}.$$

As consequences of well-known general facts, ψ is well-defined, surjective, and also

$$\ker(\psi) = \{[(g, h)] : g \in G_{i,\gamma}, h \in (K_{i,\gamma})_{[u_{i,\gamma}]}\} \stackrel{(2.12)}{=} \{[(\tilde{g}, 0)] : g \in G_{i,\gamma}\} \cong G_{i,\gamma}.$$

So, $G_{i,\gamma+1}/G_{i,\gamma} \cong H_{\mathbf{k}_{i,\gamma}, K_{i,\gamma}} / (K_{i,\gamma})_{[u_{i,\gamma}]}$. Since the latter quotient is $\mu_{i,\gamma}$ -free by assumption, the same holds for $G_{i,\gamma+1}/G_{i,\gamma}$. The next key observation is the non-extendability property of $F_{i,\gamma}$. Suppose, toward a contradiction, that $F : G_{i,\gamma+1} \rightarrow K_{i,\gamma}$ extends $F_{i,\gamma}$. Then the map

$$f := F \circ k_{i,\gamma} : H_{\mathbf{k}_{i,\gamma}, K_{i,\gamma}} \longrightarrow K_{i,\gamma}$$

satisfies

$$f(x_\eta) = F \circ k_{i,\gamma}(x_\eta) = F_{i,\gamma} \circ g_{i,\gamma}(x_\eta) = \phi_*(\eta)$$

for all $\eta \in u_{i,\gamma}$, contradicting the choice of $(H_{\mathbf{k}_{i,\gamma}, K_{i,\gamma}}, \phi_*)$.

Next, we define the map $f_{i,\gamma+1,\eta}$. Let $\eta \in \mathcal{T}_{i,\beta_i(\gamma+1)}$. For any $h \in H_{\mathbf{k}_{i,\gamma}, K_{i,\gamma}}$ and $g \in G_{i,\gamma}$, set

$$f_{i,\gamma+1,\eta}([(h, g)]) := f_{i,\gamma,\eta}(g).$$

This defines a homomorphism

$$f_{i,\gamma+1,\eta} = f_{\mathbf{m}_{i,\gamma+1,\eta}} : G_{i,\gamma+1} \longrightarrow \mathbb{Z}.$$

To verify that $f_{i,\gamma+1,\eta}$ is well-defined, it suffices to check that $f_{i,\gamma,\eta} \circ g_{i,\gamma} = 0$. Indeed, for a given $\eta \in \mathcal{T}_{i,\beta_i(\gamma+1)}$, choose $\beta \in X_{\phi_*(\eta)}$ with $\eta(i, 1) < \beta$. Then by $(*)_{\beta, \phi_*(\eta)}(\mathbf{a})$,

$$f_{i,\gamma,\eta} \circ g_{i,\gamma}(x_\eta) = f_{i,\gamma,\eta}(y_{\phi_*(\eta)}) = f_{i,\gamma,\eta}(y_{\phi_*(\eta), \beta}) = 0.$$

Also, the family $\{f_{i,\gamma+1,\eta} : \eta \in \mathcal{T}_{i,\beta_i(\gamma+1)}\}$ is independent, and

$$\bigcap_{\eta \in \mathcal{T}_{i,\beta_i(\gamma+1)}} \text{Ker}(f_{i,\gamma+1,\eta}) = \{0\}.$$

Finally, for $\eta \in \mathcal{T}_i$, define

$$f_{i,\eta} := \bigcup_{\gamma < \lambda_i} f_{i,\gamma,\eta}.$$

We are now prepared to state the following claim.

Claim 2.13. *The set $\{f_{i,\eta} : \eta \in \mathcal{T}_i\}$ generates $\text{Hom}(G_{\lambda_i}^{\mathbf{m}_i}, \mathbb{Z})$.*

Proof. Suppose, toward a contradiction, that there exists

$$f \in \text{Hom}(G_{\lambda_i}^{\mathbf{m}_i}, \mathbb{Z}) \setminus \langle f_{i,\eta} : \eta \in \mathcal{T}_i \rangle.$$

Choose $\gamma \in S$ such that $G_{i,\gamma}$ has domain γ , $f \upharpoonright \gamma = F_{i,\gamma}$, and

$$f \upharpoonright \gamma \notin \langle f_{i,\gamma,\eta} : \eta \in \mathcal{T}_i \rangle.$$

By construction, the map $f \upharpoonright \gamma$ cannot be extended to a homomorphism from $G_{i,\gamma+1}$ to \mathbb{Z} , yielding a contradiction. $\square_{2.13}$

We also note that clause $(*)_B^\gamma(i)$ holds by Claim 2.13. Indeed, given any $f \in \text{Hom}(G_{\lambda_i}^{\mathbf{m}_i}, \mathbb{Z})$, there exist $\eta_0, \dots, \eta_{n-1} \in \mathcal{T}_i$ and coefficients $\alpha_0, \dots, \alpha_{n-1} \in \mathbb{Z}$ such that $f = \sum_{k=0}^{n-1} \alpha_k f_{i,\eta_k}$. Pick $\alpha < \lambda_i$ large enough so that $\eta_k(i, 1) \leq \alpha$ for all $k < n$. By Fact 2.3(a), for each $i < \kappa$ we have

$$\text{Hom}\left(\prod_{\mathcal{T}_{i,\alpha}} \mathbb{Z}, \mathbb{Z}\right) \cong \bigoplus_{\eta \in \mathcal{T}_{i,\alpha}} \text{Hom}(\mathbb{Z}, \mathbb{Z}) \cong \bigoplus_{\eta \in \mathcal{T}_{i,\alpha}} \mathbb{Z}x_\eta.$$

In particular, there exists $h \in \text{Hom}(\prod_{\mathcal{T}_{i,\alpha}} \mathbb{Z}, \mathbb{Z})$ such that

$$h(\langle f_{i,\eta} : \eta \in \mathcal{T}_i \rangle) = \sum_{k < n} \alpha_k f_{i,\eta_k}.$$

Clause $(*)_C^\gamma(b)$ follows from the fact that S_i is non-reflecting. Hence there exists a club $C \subseteq \gamma$ with $\min(C) = \rho$ such that $C \cap S_i = \emptyset$. Then $G_{i,\gamma}/G_{i,\rho}$ is the union of the increasing and continuous sequence $\langle G_{i,\tau}/G_{i,\rho} : \tau \in C \rangle$. By the induction hypothesis, each $G_{i,\tau}/G_{i,\mu}$ is free for all $\mu < \tau$ in C , and therefore $G_{i,\gamma}/G_{i,\rho}$ is free as well.

Having defined the sequence $\langle \mathbf{m}_{i,\gamma} : \gamma < \lambda_i \rangle$ with properties $(*)_C^\gamma$, we set

$$\mathbf{m}_i := \bigcup_{\gamma < \lambda_i} \mathbf{m}_{i,\gamma}.$$

This completes the inductive construction of $\langle \mathbf{m}_i : i < \kappa \rangle$ as required by $(*)_B^i$.

Stage (C): In this step, we show that for each i , $\mathbf{m}_i \in \mathbf{M}_{2,\lambda_i}$ (see Definition 2.5(2)).

Items (a)–(e) of Definition 2.5(1) and the equalities $\alpha_{\mathbf{m}_i} = \lambda_i = \lambda_{\mathbf{m}_i}$ are immediate.

For clause (i), let $\alpha < \lambda_i$ and $0 \neq G' \subseteq G_\alpha^{\mathbf{m}_i}$. Then there exists $\gamma < \lambda_i$ such that $G' \subseteq G_{\alpha_i,\gamma}^{\mathbf{m}_i}$. Fix $0 \neq x \in G'$. By property $(*)_C^\gamma(d)$,

$$\bigcap \{ \text{Ker}(f_{i,s}) : s \in \mathcal{T}_i \} = \{0\}.$$

Hence there exists $s \in \mathcal{T}_i$ with $f_{i,s}(x) \neq 0$. In particular, the restriction $f_{i,s} \upharpoonright G' \in \text{Hom}(G', \mathbb{Z})$ is non-zero, as required.

Stage (D): In this stage we conclude the proof of Theorem 2.6. For each $i < \kappa$, set

$$G_i := G_{\mathbf{m}_i,\lambda_i}, \quad G_{<i} := \bigcup_{j < i} G_j.$$

Then $\langle G_i : i < \kappa \rangle$ is increasing and continuous, and for each i the quotient $G_i/G_{<i}$ is $(\chi + \sum_{j < i} \lambda_j)$ -free. Indeed, for all $\gamma < \lambda_i$ we have $\mu_{i,\gamma} \geq \chi + \sum_{j < i} \lambda_j$, and each $G_{i,\gamma+1}/G_{i,\gamma}$ is $\mu_{i,\gamma}$ -free.

Define

$$G := \bigcup_{i < \kappa} G_i.$$

Then G is an abelian group of size λ . We first show that G is χ -free. To this end, let $H \leq G$ be a subgroup of size $< \chi$. Then the sequence

$$\langle H \cap G_i : i < \kappa \rangle$$

is increasing and continuous. For each $i < \kappa$,

$$(H \cap G_i)/(H \cap G_{<i}) \cong ((H \cap G_i) + G_{<i})/G_{<i},$$

and this group is free since $G_i/G_{<i}$ is $(\chi + \sum_{j < i} \lambda_j)$ -free, and therefore χ -free. Hence $H = \bigcup_{i < \kappa} (H \cap G_i)$ is free.

Next, let $H \leq G$ be a nonzero subgroup of size $< \lambda$. Choose $i < \kappa$ such that $H \cap G_{<i} \neq \{0\}$ and $|H| < \lambda_i$. By Definition 2.5(2)(i),

$$\text{Hom}(H \cap G_i, \mathbb{Z}) \neq 0.$$

Moreover,

$$(H \cap G_i)/(H \cap G_{<i}) \cong ((H \cap G_i) + G_{<i})/G_{<i}$$

is free, and therefore $\text{Hom}(H, \mathbb{Z}) \neq 0$.

Finally, we show that $\text{Hom}(G, \mathbb{Z}) = 0$. Suppose, towards a contradiction, that $0 \neq f \in \text{Hom}(G, \mathbb{Z})$. By $(*)_B^i$ (i), for each $i < \kappa$ there exist $\alpha_i < \lambda_i$ and

$$h_i \in \text{Hom}\left(\prod_{\mathcal{T}_{i,\alpha_i}} \mathbb{Z}, \mathbb{Z}\right)$$

such that

$$f(x) = h_i(\langle f_{\mathbf{m}_i,\eta}(x) : \eta \in \mathcal{T}_{i,\alpha_i} \rangle), \quad x \in G_i.$$

By Fact 2.3(a),

$$\text{Hom}\left(\prod_{\mathcal{T}_{i,\alpha_i}} \mathbb{Z}, \mathbb{Z}\right) \cong \bigoplus_{\eta \in \mathcal{T}_{i,\alpha_i}} \mathbb{Z}x_\eta.$$

By [3, Corollary III.3.3], for each i there exists a finite set $u_i \subseteq \mathcal{T}_{i,\alpha_i}$ such that

$$f_{\mathbf{m}_i,\eta}(x) = 0 \text{ for all } \eta \in u_i \implies f(x) = 0, \quad x \in G_i.$$

Since $\kappa = \text{cf}(\kappa) > \aleph_0$, there exists n_* such that the set

$$\mathcal{V}_1 := \{i < \kappa : |u_i| = n_*\}$$

is unbounded in κ .

For $i < j < \kappa$, define

$$\text{pr}_{i,j} : \mathcal{T}_j \rightarrow \mathcal{T}_i, \quad \text{pr}_{i,j}(\eta) = \eta \upharpoonright (i+1).$$

If $\eta \in u_j$, then $\eta \upharpoonright (i+1) \in u_i$, since $G_i \subseteq G_j$ and $f_{\mathbf{m}_i,\eta \upharpoonright (i+1)} \subseteq f_{\mathbf{m}_j,\eta}$. Thus $\text{pr}_{i,j}$ maps u_j onto u_i .

Let \mathcal{T} be the tree of height κ whose i -th level is u_i . This is a well-defined tree of uncountable height with finite levels. By Fact 1.10, \mathcal{T} carries a cofinal branch, contradicting Claim 2.8. \square

ACKNOWLEDGEMENTS

The authors are grateful to the referees for a very careful reading of the paper and for helpful comments that have improved both its clarity and presentation.

REFERENCES

- [1] M. Asgharzadeh, M. Golshani and S. Shelah, *Many forcing axioms and incomactness for Ext*, work in progress.
- [2] P. C. Eklof and A. Mekler, *On constructing indecomposable groups in \mathbf{L}* , J. Algebra **40** (1977), 96-103.
- [3] P. C. Eklof and A. Mekler, *Almost free modules: Set theoretic methods*, Revised Edition, North-Holland Publishing Co., North-Holland Mathematical Library, **65**, 2002.
- [4] L. Fuchs, *Abelian groups*, Springer Monographs in Mathematics. Springer, Cham, 2015.
- [5] R. Göbel and J. Trlifaj, *Approximations and Endomorphism Algebras of Modules*, vol. i, ii, de Gruyter Expositions in Mathematics, Walter de Gruyter, 2012.
- [6] P. Hill, *On the freeness of abelian groups: a generalization of Pontryagin's theorem*, Bull. Amer. Math. Soc. **76** (1970), 1118–1120.
- [7] T. Jech, *Set theory*, The third millennium edition, revised and expanded. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003.
- [8] S. Shelah, *Infinite abelian groups, Whitehead problem and some constructions*, Israel J. Math. **18** (1974), 243–256.
- [9] S. Shelah, *A compactness theorem for singular cardinals, free algebras, Whitehead problem and transversals*, Israel Journal of Mathematics **21** (1975), 319–349.
- [10] S. Shelah, *Incompactness in singular cardinals*, Bulletin of Symbolic Logic 2008 sep. **14**(3) 2008, Annual Meeting in Irvine in 2008, available at <https://www.jstor.org/stable/pdf/20059993.pdf>, pp. 419-420.
- [11] S. Shelah, *Reflexive abelian groups and measurable cardinals and full MAD families*, Algebra Universalis **63** (2010), 351–366.
- [12] S. Shelah, *Quite free complicated abelian groups, pcf and black boxes*, Israel J. Math. **240** (2020), 1–64.

MOHSEN ASGHARZADEH, HAKIMIYEH, TEHRAN, IRAN.
E-mail address: mohsenasgharzadeh@gmail.com

MOHAMMAD GOLSHANI, SCHOOL OF MATHEMATICS, INSTITUTE FOR RESEARCH IN FUNDAMENTAL SCIENCES (IPM), P.O. Box: 19395–5746, TEHRAN, IRAN.
E-mail address: golshani.m@gmail.com

SAHARON SHELAH, EINSTEIN INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM, 91904, ISRAEL, AND DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW BRUNSWICK, NJ 08854, USA.
E-mail address: shelah@math.huji.ac.il