Paper Sh:1258, version 2026-09-29_3. See https://shelah.logic.at/papers/1258/ for possible updates.

CONSISTENCY OF SQUARE BRACKET PARTITION RELATION

SAHARON SHELAH

Abstract. Characteristic earlier results were of the form $\mathrm{CON}(2^{\aleph_0} \to [\lambda]^2_{n,2})$, with $2^{\aleph_0}$ an ex-large cardinal, in the best case the first weakly Mahlo cardinal.

Characteristic new results are $\mathrm{CON}\big((2^{\aleph_0} = \aleph_m) + \aleph_\ell \to [\aleph_k]^2_{n,2}\big)$, for suitable $k < \ell < m$. So we improve in three respects: the continuum may be small (e.g. not a weakly Mahlo), we use no large cardinal, and the cardinals λ involved are $< 2^{\aleph_0}$ after the forcing.

§ 0. Introduction

In their seminal list of problems [EH71], Erdös and Hajnal posed the question (15(a)): does $2^{\aleph_0} \not\to [\aleph_1]^2_3$? Recently, Komjáth [Kom25] provided a comprehensive update on this topic.

We continue here works which start with the problem above:[She88, §2], [She92], [She89], [She95] [She96], [She00] and the work with Rabus [RS00], but we try to be self-contained.

The simplest case of our result is (recall 0.3 below):

Theorem 0.1. *Assume* GCH *for transparency. Then for some ccc forcing notion of cardinality $\aleph_6$ in the universe $\mathbf{V}^{\mathbb{P}}$, we have $2^{\aleph_0} = \aleph_6$ and for any $n \geq 3$, $\aleph_5 \to [\aleph_2]^2_{n,2}$.*

Proof. Choose $(\mu, \theta, \partial, \lambda)$ as $(\aleph_6, \aleph_5, \aleph_2, \aleph_0)$ and apply Theorem 0.2 and Fact 1.12 with $\partial_0 = \aleph_1$. $\square_{0.1}$

For Hypothesis 1.1, the main case is:

Theorem 0.2. *Assume $\lambda = \lambda^{<\lambda} < \partial < \theta < \mu = \mu^\theta$, $\partial = \partial^\lambda$ and $2^{\partial^{+\ell}} = \partial^{+\ell+1}$ for $\ell = 0, 1, 2$ and $\partial^{+4} \leq \theta$. Then for some λ^+-cc, $(< \lambda)$-complete forcing notion $\mathbb{P}$ of cardinality μ (so the forcing does not collapse any cardinal and preserves cardinal arithmetic outside $[\lambda, \mu)$), in the universe $\mathbf{V}^{\mathbb{P}}$ we have, $2^\lambda = \mu$ and for every $\sigma < \lambda$, $\theta \to [\partial]^2_{\sigma,2}$*

Proof. All this paper is dedicated to proving this theorem. Pedantically, choose $\partial = \kappa^+$, notice that Hypothesis 1.1 holds (by Fact 1.12) so we can apply Conclusion 1.11. $\square_{1.11}$

We may weaken $\mu = \mu^\theta$ to $\mu = \mu^\partial$ and replace $\partial = \kappa$ by ∂ being a suitable limit cardinal.

Recall,

Date: December 24, 2025.

2020 *Mathematics Subject Classification.* 03E2, 03E35.

Key words and phrases. set theory, partition calculus, square bracket arrows.

First typed: January 7, 2024. The author would like to thank the Israel Science Foundation for partial support of this research by grant 2320/23. The author would like to thank Craig Falls for generously funding typing services, and the typist for his careful and beautiful work. Submitted to Yair Hayut in February 2025 for the IJM volume dedicated to Menachem Magidor. References like [Sh:950, Th0.2=Ly5] mean that the internal label of Th0.2 is y5 in Sh:950. The reader should note that the version in my website is usually more up-to-date than the one in arXiv. This is publication number 1258 on Saharon Shelah's list.

Paper Sh:1258, version 2026-09-29_3. See https://shelah.logic.at/papers/1258/ for possible updates.

Definition 0.3. For possibly finite cardinals θ, ∂, σ and κ, let $\theta \to [\partial]^2_{\sigma,\kappa}$ mean:

- if $\mathbf{c}$ is a function from $[\theta]^2 := \{u \subseteq \theta : |u| = 2\}$ into σ, then there exists some subset $\mathscr{U}$ of θ of cardinality ∂ such that $\{\mathbf{c}(u) : u \in [\mathscr{U}]^2\}$ has at most κ-many members.

We thank Yair Hayut and the referee for many helpful comments.

§ 0(A). **Preliminaries.**

Notation 0.4.

1) $\mathrm{cof}(\delta)$ is the class of ordinals of cofinality $\mathrm{cf}(\delta)$.

2) For a set x, let $\mathrm{trcl}(x)$ be the transitive closure of x, that is, the minimal set Y such that $x \in Y$ and $(\forall y)(y \in Y \Rightarrow y \subseteq Y)$.

3) Let $\mathscr{H}(\lambda) := \{x : |\mathrm{trcl}(x)| < \lambda\}$.

4) Let $\mathrm{trcl}_{\mathrm{Ord}}(x)$ be defined similarly to $\mathrm{trcl}(x)$ considering ordinals as atoms (= elements), equivalently, the minimal set Y such that $x \in Y$ and

$$(\forall y)[y \in Y \wedge (\text{if } y \text{ is not an ordinal, then } y \subseteq Y)].$$

5) Let $\mathscr{H}_{<\kappa}(x) = \{x : \mathrm{trcl}_{\mathrm{Ord}}(x) \subseteq \mathscr{H}(x)$ but has cardinality $< \kappa\}$.

Notation 0.5.

(1) $\mathbb{P}$, $\mathbb{Q}$ and $\mathbb{R}$ are forcing notions.

(2) p, q, r called *conditions* are members of a forcing notion.

(3) $\mathbf{q}$ is as in Definition 1.3, some kind of $(< \lambda)$-support iterated forcing with extra information.

Notation 0.6. We may write e.g. $N[\mathbf{q}, \beta, u]$ instead $N_{\mathbf{q},\beta,u}$ to help with sub-scripts (or super-script).

Definition 0.7. Let θ, ∂, κ and λ be infinite cardinals. We say that $\theta \to_{\mathrm{sq}} (\partial)^{\lambda,2}_{\kappa}$ when $\theta > \partial \geq \kappa \geq \lambda$ and:

$\boxplus$ If (a) then (b), where:

(a) $\mathscr{B}$ is an expansion of $(\mathscr{H}(\chi), \in, <_*)$, where $<_*$ is a well-ordering of $\mathscr{H}(\chi)$, $\chi > \theta$, and its vocabulary $\tau_{\mathscr{B}}$ has cardinality $\leq \lambda$.

(b) There is a tuple $\mathbf{s} = (\mathscr{U}, \bar{N}, \bar{\pi})$ solving $\mathbf{p} = (\mu, \theta, \partial, \kappa, \lambda, \mathscr{B})$, which means:

$\boxplus_{\mathbf{p},\mathbf{s}}$ for $u, v \in [\mathscr{U}]^{\leq 2}$,

- $\bullet_1$ $\bar{N} = \langle N_u : u \in [\mathscr{U}]^{\leq 2}\rangle$,
- $\bullet_2$ $\mathscr{U} \subseteq \theta$ is such that $\mathrm{otp}(\mathscr{U}) = \partial$,
- $\bullet_3$ $N_u \prec \mathscr{B}$, $[N_u]^{<\lambda} \subseteq N_u$,
- $\bullet_4$ $\varepsilon[\mathbf{s}] := \min(\mathscr{U})$,
- $\bullet_5$ $N_u \cap \mathscr{U} = u$,
- $\bullet_6$ $\|N_u\| = \kappa$ and $\kappa + 1 \subseteq N_u$,
- $\bullet_7$ $N_u \cap N_v \prec N_{u \cap v}$,
- $\bullet_8$ $\bar{\pi} = \langle \pi_{u,v} : u, v \in [\mathscr{U}]^{\leq 2}$ and $|u| = |v|\rangle$ such that if $|u| = |v|$, then $\pi_{u,v}$ is an isomorphism from N_v onto N_u mapping v onto u,
- $\bullet_9$ if $u_1 \subseteq u_2$ and $v_1 \subseteq v_2$ all from $[\mathscr{U}]^{\leq 2}$ and $|u_2| = |v_2|$, $\pi''_{u_2,v_2}(v_1) = u_1$ then π_{u_1,v_1}, π_{u_2,v_2} are compatible functions[1],
- $\bullet_{10}$ for $\ell = 1, 2$, the sets $N_u \cap \partial$ for $u \in [\mathscr{U}]^{\ell}$ are pairwise equal [2] and included in $N_{\emptyset}$.

[1] So e.g. it follows that: if $\zeta_1, \zeta_2 \in \mathscr{U}$ then $\pi_{\{\zeta_1\},\{\zeta_2\}} \restriction (N_{\emptyset} \cap N_{\{\zeta_2\}})$ is the identity map.

[2] Note that ∂ has two distinct roles: the size of $\mathscr{U}$ and the restriction on $N_u \cap \partial$. We may separate.

Paper Sh:1258, version 2026-09-29_3. See https://shelah.logic.at/papers/1258/ for possible updates.

Observation 0.8. *If $\bar{N} = \langle N_u : u \in [\mathscr{U}]^{\leq 2} \rangle$ satisfies 0.7(b)$\bullet_1 + \bullet_7$, then:*

($*$) *For every $x \in \cup\{N_u : u \in [\mathscr{U}]^{\leq 2}\}$ the set $\{u \in [\mathscr{U}]^{\leq 2} : x \in N_u\}$ has one of the following forms:*
- *(a)* $\{u\}$ *for some* $u \in [\mathscr{U}]^2$,
- *(b)* $\{\zeta\}$ *for some* $\zeta \in \mathscr{U}$,
- *(c)* $\{\{\zeta\}\} \cup \{\{\varepsilon, \zeta\} : \varepsilon \in \mathscr{U} \cap \zeta\}$ *for some* $\zeta \in \mathscr{U}$,
- *(d)* $\{\{\zeta\}\} \cup \{\{\zeta, \xi\} : \xi \in \mathscr{U} \setminus (\zeta + 1)\}$ *for some* $\zeta \in \mathscr{U}$,
- *(e)* $\{\emptyset\}$,
- *(f)* $\{\emptyset\} \cup \{\{\zeta\} : \zeta \in \mathscr{U}\}$,
- *(g)* $\{\emptyset\} \cup \{\{\zeta\} : \zeta \in \mathscr{U}\} \cup \{\{\varepsilon, \zeta\} : \varepsilon < \zeta$ *are from* $\mathscr{U}\}$.

§ 1. The forcing

Our aim here is to prove the consistency of the following configuration:

$$2 < \sigma < \lambda = \lambda^{<\lambda} < \partial = \partial^{<\lambda} < \theta < \mu = \mu^\theta = 2^\lambda,$$

and having $\theta \to [\partial]^2_{\sigma,2}$.

A continuation is in preparation [S$^+$], aiming to further develop the directions explored here, particularly for the case of superscript $\mathbf{n} > 2$, as dealt within [She92]. We also show there that we can weaken the requirements on the cardinals and have more pairs.

Hypothesis 1.1. The parameter $\mathbf{p} = (\mu, \theta, \partial, \lambda, \lambda, \mathscr{B})$ consists of the following:
- (a) $\lambda = \lambda^{<\lambda} < \partial < \theta < \mu = \mu^\theta$,
- (b) $\theta \to_{\rm sq} (\partial)^{\lambda,2}_\lambda$ (see Definition 0.7, a variant of [She89, 2.1]); in our case using λ twice in intentional.
- (c) σ will vary on the cardinal numbers from $(2, \lambda)$ and the "nice" μ-s are such that $\gamma < \mu \Rightarrow |\gamma|^\theta < \mu$.
- (d)
  - χ is e.g. $\beth_2(\mu)^+$,
  - let $\mathscr{B}$ be an expansion of $(\mathscr{H}(\chi), \in, <^*_\chi)$ with vocabulary of cardinality λ such that for any finite set $u \subseteq \mathscr{H}(\chi)$, the Skolem hull of u $N_u := {\rm Sk}(u, \mathfrak{C}_*)$ is of cardinality λ and $|N_u|^{<\lambda} \subseteq N$.

We intend to use $(<\lambda)$-support iterated forcing of quite a special kind but first, we define the iterand.

Definition 1.2.

(1) Let $\mathbf{A}$ be the set of objects $\mathbf{a}$ consisting of:
- (a)
  - $\gamma < \mu$ and $\sigma \in (2, \lambda)$,
  - $\mathbb{P}$ is a forcing notion such that:
    $$p \in \mathbb{P} \Rightarrow {\rm dom}(p) \in [\gamma]^{<\lambda} \wedge (\forall \alpha \in {\rm dom}(p))(p(\alpha) \in [\lambda \cup \gamma]^{<\lambda}),$$
  - $\mathbb{P}$ is λ^+-cc and $(<\lambda)$-complete,
  - the order $\leq_{\mathbb{P}}$ is: $p \leq_{\mathbb{P}} q$ iff:
    $${\rm dom}(p) \subseteq {\rm dom}(q) \wedge (\forall \alpha \in {\rm dom}(p))[p(\alpha) \subseteq q(\alpha)],$$
- (b)
  - $\underset{\sim}{\mathbf{c}}$ is a $\mathbb{P}$-name of a function from $[\theta]^2$ to σ, (we may write $\underset{\sim}{\mathbf{c}}(\alpha, \beta)$ instead $\underset{\sim}{\mathbf{c}}(\{\alpha, \beta\})$ for $\alpha \neq \beta < \theta$).
- (c) We have $(\mathscr{U}, \bar{N}, \bar{\pi})$ solving $\mathbf{p} = (\mu, \theta, \partial, \lambda, \lambda, \mathscr{B})$, (with $\mathscr{B}$ as in Definition 0.7⊞(b) and 1.1) such that $\mathbb{P}, \underset{\sim}{\mathbf{c}} \in N_u$ for every $u \in [\mathscr{U}]^{\leq 2}$.

(1A) In the context of Definition 1.2(1), $\mathbf{a} = (\gamma, \mathbb{P}, \underset{\sim}{\mathbf{c}}, \mathscr{U}, \bar{N}, \bar{\pi}) = (\gamma_{\mathbf{a}}, ...)$, so e.g. $N_{\mathbf{a},u} = N_u$.

(2) We say that the pair $(p, \bar{\iota})$ is a *solution* of $\mathbf{a} \in \mathbf{A}$, and write $(\mathbf{a}, p, \bar{\iota}) \in \mathbf{A}^+$, when:
- (a) $\bar{\iota} = (\iota_1, \iota_2) \in \sigma \times \sigma$,

Paper Sh:1258, version 2026-09-29_3. See https://shelah.logic.at/papers/1258/ for possible updates.

(b) $p \in \mathbb{P}_{\mathbf{a}} \cap N_{\mathbf{a},\{\varepsilon[\mathbf{a}]\}}$, recalling $\varepsilon(\mathbf{a}) = \min(\mathscr{U})$,

(c) if $p \le q \in \mathbb{P}_{\mathbf{a}} \cap N_{\mathbf{a},\{\varepsilon[\mathbf{a}]\}}$ and $\zeta_1 < \zeta_2$ are from $\mathscr{U}$ then there are q_1, q_2, r_1, r_2 such that for $\ell = 1, 2$, we have:

- $\bullet_0$ $q \le_{\mathbb{P}_{\mathbf{a}}} q_\ell$,
- $\bullet_1$ $q_\ell \in \mathbb{P}_{\mathbf{a}} \cap N_{\mathbf{a},\{\varepsilon[\mathbf{a}]\}}$ and $q_1 \restriction (N_{\mathbf{a},\emptyset} \cap \gamma_{\mathbf{a}}) = q_2 \restriction (N_{\mathbf{a},\emptyset} \cap \gamma_{\mathbf{a}})$,
- $\bullet_2$ $r_\ell \in \mathbb{P}_{\mathbf{a}} \cap N_{\mathbf{a},\{\zeta_1,\zeta_2\}}$,
- $\bullet_3$ $r_\ell \Vdash$ "$\underset{\sim}{\mathbf{c}}(\zeta_1, \zeta_2) = \iota_{\mathbf{a},\ell}$",
- $\bullet_4$ $r_\ell \restriction N_{\mathbf{a},\{\zeta_1\}}$ is $\le_{\mathbb{P}_{\mathbf{a}}}$-below $\pi^{\mathbf{a}}_{\{\zeta_1\},\{\varepsilon[\mathbf{a}]\}}(q_\ell)$,
- $\bullet_5$ $r_\ell \restriction N_{\mathbf{a},\{\zeta_2\}}$ is $\le_{\mathbb{P}_{\mathbf{a}}}$-below $\pi^{\mathbf{a}}_{\{\zeta_2\},\{\varepsilon[\mathbf{a}]\}}(q_{3-\ell})$.

(3) If $\mathbf{b} = (\mathbf{a}, p, \bar{\iota}) \in \mathbf{A}^+$ then let $\underset{\sim}{\mathbb{Q}}_{\mathbf{b}}$ be the $\mathbb{P}$-name of the following forcing notion:

($*$) For $\mathbf{G} \subseteq \mathbb{P}$ generic over $\mathbf{V}$,

(a) the set of elements of $\mathbb{Q}_{\mathbf{b}} = \underset{\sim}{\mathbb{Q}}_{\mathbf{b}}[\mathbf{G}]$ is:

$$\Big\{u \in [\mathscr{U}]^{<\lambda}\colon \text{ if } \zeta_1 < \zeta_2 \text{ in } \mathscr{U}, \text{ then } \underset{\sim}{\mathbf{c}}\{\zeta_1, \zeta_2\}[\mathbf{G}] \in \{\iota_1, \iota_2\}, \text{ moreover}$$

$$\text{for some } q_1, q_2, r_1, r_2 \text{ as in Definition 1.2(1)(c)}(\bullet_1)\text{-}(\bullet_5), \text{ we have } r_1 \in \mathbf{G} \text{ or } r_2 \in \mathbf{G}\Big\},$$

(b) the order of $\underset{\sim}{\mathbb{Q}}_{\mathbf{b}}[\mathbf{G}]$ is inclusion,

(c) the generic is $\underset{\sim}{\mathscr{Y}}_{\mathbf{b}} = \bigcup \underset{\sim}{\mathbf{G}}_{\mathbb{Q}_{\mathbf{b}}}$.

Definition 1.3.

(1) Let $\mathbf{Q} := \mathbf{Q}_{\mathbf{p}}$ be the class of $\mathbf{q}$ which consist of (below, $\alpha \le \lg(\mathbf{q})$ and $\beta < \lg(\mathbf{q})$ and e.g. $\mathbb{P}_\alpha = \mathbb{P}_{\mathbf{q},\alpha}$):

(a) $\lg(\mathbf{q})$ is an ordinal $\le \mu$,

(b) $\langle \mathbb{P}_\alpha, \underset{\sim}{\mathbb{Q}}_\beta\colon \alpha \le \lg(\mathbf{q}), \beta < \lg(\mathbf{q})\rangle$ is a $(<\lambda)$-support iteration,

(c) $\mathbb{P}_\beta$ satisfies the λ^+-cc,

(d) $\underset{\sim}{\mathbb{Q}}_\beta$ is $\underset{\sim}{\mathbb{Q}}_{\mathbf{b}_\beta}$, where:

- $\bullet_1$ $\mathbf{b}_\beta := (\mathbf{a}_\beta, p^*_\beta, \bar{\iota}^*_\beta) \in \mathbf{A}^+$,
- $\bullet_2$ $\mathbf{a}_\beta := (\gamma_\beta, \mathbb{P}^\bullet_\beta, \underset{\sim}{\mathbf{c}}_\beta, \mathscr{U}_\beta, \bar{N}_\beta, \bar{\pi}_\beta) \in \mathbf{A}$,
- $\bullet_3$ $\mathbb{P}^\bullet_\beta$ is equal to $\mathbb{P}'_{\xi(\beta)}$ for some $\xi(\beta) = \xi_{\mathbf{q}}(\beta) \le \beta$ (on $\mathbb{P}'_\beta$, see below),
- $\bullet_4$ The sequence $\langle(\mathbb{P}_\gamma, \mathbb{P}'_\gamma, \mathbf{a}_\gamma, \mathbf{b}_\gamma, \xi(\gamma)) : \gamma < \beta\rangle$ belongs to $N_{\beta,u}$ for every $u \in [\mathscr{U}_\beta]^{\le 2}$.
- $\bullet_5$ Let $\mathscr{W}_\beta = \bigcup\{N_{\beta,u} \cap \beta\colon u \in [\mathscr{U}_\beta]^{\le 2}\}$,
- $\bullet_6$ we [3] have: for every $\gamma \in \mathscr{W}_\beta$ the set $\mathscr{W}_\beta \cap \mathscr{W}_\gamma$ has cardinality $\le \lambda$,
- $\bullet_7$ For every $\gamma \in \mathscr{W}_\beta$, there is $u = u_{\beta,\gamma} \in [\mathscr{U}_\beta]^{\le 2}$ such that $\mathscr{W}_\beta \cap \mathscr{W}_\gamma \subseteq N_{\beta,u}$ and without loss of generality u is minimal with this property.

(e) $\mathbb{P}'_\alpha$ is a dense subset of $\mathbb{P}_\alpha$, where,

- $\mathbb{P}'_\alpha$ is $\mathbb{P}_\alpha$ restricted to the set of conditions $p \in \mathbb{P}_\alpha$ such that: if $\beta \in \mathrm{dom}(p)$ then $p(\beta)$ is a member of $\mathbf{V}$ (not just a $\mathbb{P}_\alpha$-name) and if $\zeta_1 < \zeta_2$ are in $p(\beta) \subseteq \mathscr{U}_\beta$, then there are q_1, q_2, r_1, r_2 as in Definition 1.2(2)(c)$(\bullet_1)$-$(\bullet_5)$ with $\mathbf{a}_\beta, \mathbf{b}_\beta$ here standing for $\mathbf{a}, \mathbf{b}$ there and
$$\bigvee_{\ell=1}^{2} (\forall \gamma \in \mathrm{dom}(r_\ell))[\gamma \in \mathrm{dom}(p) \wedge r_\ell(\gamma) \subseteq p(\gamma)].$$

(f) $\gamma_{\mathbf{q}} := \gamma(\mathbf{q}) := \sup\{\gamma_{\mathbf{q},\beta}\colon \beta < \lg(\mathbf{q})\}$, so $\mathbb{P}'_{\gamma(\mathbf{q})} \subseteq \mathscr{H}_{<\lambda}(\gamma_{\mathbf{q}})$; let $\mathbb{P}_{\mathbf{q}} := \mathbb{P}_{\lg(\mathbf{q})}$ and $\mathbb{P}'_{\mathbf{q}} := \mathbb{P}'_{\lg(\mathbf{q})}$.

(1A) We may write either $\mathbb{P}_{\mathbf{q},\alpha}$ or $\mathbb{P}_\alpha$ whenever $\mathbf{q}$ is clear and $(\iota_{\mathbf{q},\beta,1}, \iota_{\mathbf{q},\beta,2})$ is $\bar{\iota}_{\mathbf{b}_\beta}$.

[3] Why? By 0.7(b)$\bullet_{10}$.

Paper Sh:1258, version 2026-09-29_3. See https://shelah.logic.at/papers/1258/ for possible updates.

(2) Let $\leq_{\mathbf{p}}$ be the following two-place relation on $\mathbf{Q}_{\mathbf{p}}$:

$$\mathbf{q}_1 \leq_{\mathbf{p}} \mathbf{q}_2 \text{ iff } \mathbf{q}_1 = \mathbf{q}_2 \restriction \lg(\mathbf{q}_1), \text{ see below.}$$

(3) For $\mathbf{q}_2 \in \mathbf{Q}_{\mathbf{p}}$ and $\alpha_* \leq \lg(\mathbf{q}_2)$, we define $\mathbf{q}_1 := \mathbf{q}_2 \restriction \alpha_*$ by:

(a) $\lg(\mathbf{q}_1) = \alpha_*$,
(b) $(\mathbb{P}_{\mathbf{q}_1,\alpha}, \mathbb{P}'_{\mathbf{q}_1,\alpha}) = (\mathbb{P}_{\mathbf{q}_2,\alpha}, \mathbb{P}'_{\mathbf{q}_2,\alpha})$ for $\alpha \leq \alpha_*$,
(c) $(\mathbb{Q}_{\mathbf{q}_1,\beta}, \mathbf{b}_{\mathbf{q}_1,\beta}, \xi_{\mathbf{q}_1}(\beta)) = (\mathbb{Q}_{\mathbf{q}_2,\beta}, \mathbf{b}_{\mathbf{q}_2,\beta}, \xi_{\mathbf{q}_2}(\beta))$ for $\beta < \alpha_*$.

(4) We say that two conditions $p, q \in \mathbb{P}'_\alpha$ are *isomorphic*, when:

(a) $\mathrm{otp}(\mathrm{dom}(p)) = \mathrm{otp}(\mathrm{dom}(q))$, and
(b) if $\beta \in \mathrm{dom}(p) \cap \mathrm{dom}(q)$ then:
- $\bullet_1$ $\mathrm{otp}(p(\beta)) = \mathrm{otp}(q(\beta))$,
- $\bullet_2$ if $\varepsilon \in p(\beta) \cap q(\beta)$ then $\mathrm{otp}(\varepsilon \cap p(\beta)) = \mathrm{otp}(\varepsilon \cap q(\beta))$,
- $\bullet_3$ if $\varepsilon \in p(\beta), \zeta \in q(\beta)$ and $\mathrm{otp}(\varepsilon \cap p(\beta)) = \mathrm{otp}(\zeta \cap q(\beta))$ then:
$$\pi_{\beta,\{\zeta\},\{\varepsilon\}}(p \restriction N_{\beta,\{\varepsilon\}}) = q \restriction N_{\beta,\{\zeta\}}.$$
- $\bullet_4$ if $\varepsilon < \varepsilon_1$ belong to $p(\beta), \zeta < \zeta_1$ belong to $q(\beta)$, $\mathrm{otp}(\varepsilon \cap p(\beta)) = \mathrm{otp}(\zeta \cap q(\beta))$ and $\mathrm{otp}(\varepsilon_1 \cap p(\beta)) = \mathrm{otp}(\zeta_1 \cap q(\beta))$ then:
$$\pi_{\beta,\{\zeta,\zeta_1\},\{\varepsilon,\varepsilon_1\}}(p \restriction N_{\beta,\{\varepsilon,\varepsilon_1\}}) = q \restriction N_{\beta,\{\zeta,\zeta_1\}}.$$

Remark 1.4. If we prefer in clause (d) ($\bullet_3$) of Definition 1.3 (1) to have $\xi(\beta) = \beta$, i.e., $\mathbb{P}^\bullet_\beta = \mathbb{P}'_\beta$, we need to add, e.g. "$\mu$ is regular and e.g. use a preliminary forcing $(\{\mathbf{q} \in \mathbf{Q}_{\mathbf{p}} : \lg(\mathbf{q}) < \mu\}, \lhd)$".

Claim 1.5.

(0) For $\mathbf{q} \in \mathbf{Q}_{\mathbf{p}}$, we have: $\mathbb{P}'_{\mathbf{q}} \models$ "$p \leq q$" iff $\{p, q\} \subseteq \mathbb{P}'_{\mathbf{q}}$, $\mathrm{dom}(p) \subseteq \mathrm{dom}(q)$, and $\beta \in \mathrm{dom}(p) \Rightarrow p(\beta) \subseteq q(\beta)$.

(1) For $\mathbf{q} \in \mathbf{Q}_{\mathbf{p}}$, any increasing sequence of members of length $< \lambda$ of $\mathbb{P}'_{\mathbf{q}}$ has a lub, in fact, if $\delta < \lambda$, $\bar{p} = \langle p_i : i < \delta \rangle \in {}^\delta(\mathbb{P}'_{\mathbf{q}})$ is increasing, then the following $p \in \mathbb{P}'_{\mathbf{q}}$ is a lub of $\bar{p}$; defined by: $\mathrm{dom}(p) = \bigcup\{\mathrm{dom}(p_i) : i < \delta\}$, and if $\beta \in \mathrm{dom}(p)$ then

$$p(\beta) = \bigcup \{p_i(\beta) : i < \delta \text{ and } \beta \in \mathrm{dom}(p_i)\}.$$

We denote this p by $\lim(\bar{p})$.

(2) For $\mathbf{q} \in \mathbf{Q}_{\mathbf{p}}$, we have:

- *$p \in \mathbb{P}'_{\mathbf{q}}$ iff:*
  - *(a) p is a function with domain $\in [\lg(\mathbf{q})]^{<\lambda}$,*
  - *(b) if $\beta \in \mathrm{dom}(p)$ then $p(\beta)$ belongs to $[\mathscr{U}_\beta]^{<\lambda}$.*
  - *(c) If $\beta \in \mathrm{dom}(p)$ and $(\iota_1, \iota_2) = (\iota_{\mathbf{q},\beta,1}, \iota_{\mathbf{q},\beta,2})$ then for every $\zeta_1 < \zeta_2$ from $p(\beta)$, $(p \restriction \beta) \restriction N_{\mathbf{q},\beta,\{\zeta_1,\zeta_2\}} \Vdash_{\mathbb{P}_{\mathbf{q},\beta}}$ "$\mathfrak{c}\{\zeta_1, \zeta_2\} \in \{\iota_1, \iota_2\}$". Moreover, there are q_1, q_2, r_1, r_2 as in Definition 1.2(2)(c)($\bullet_1$)-($\bullet_5$) and*
$$\bigvee_{\ell=1}^{2} (\forall \gamma \in \mathrm{dom}(r_\ell))[\gamma \in \mathrm{dom}(p) \cap \beta \wedge r_\ell(\gamma) \subseteq p(\gamma)].$$

(3) If $\mathbf{q} \in \mathbf{Q}_{\mathbf{p}}$ and $\alpha \leq \lg(\mathbf{q})$ then $\mathbf{q} \restriction \alpha \in \mathbf{Q}_{\mathbf{p}}$.

(4) $\leq_{\mathbf{p}}$ is a partial order on $\mathbf{Q}_{\mathbf{p}}$.

(5) If $\bar{\mathbf{q}} = \langle \mathbf{q}_j : j < \delta \rangle$ is $\leq_{\mathbf{p}}$-increasing then it has a $\leq_{\mathbf{p}}$-lub, $\lim(\bar{\mathbf{q}})$, of length $\cup\{\lg(\mathbf{q}_j) : j < \delta\}$.

(6) If $\beta < \lg(\mathbf{q})$, $\mathbf{a} = \mathbf{a}_{\mathbf{q},\beta}$, $u \in [\mathscr{U}_{\mathbf{a},\beta}]^{\leq 2}$ and $N_u = N_{\mathbf{a},u}$, then:

- *($*$) if $p \in \mathbb{P}'_{\mathbf{q}}$ then $q = p \restriction N_{\mathbf{q},\beta,u}$ satisfies $q \in N_u$ and $q \leq_{\mathbb{P}_{\mathbf{q}}} p$ where q is defined by:*
  - *$\bullet_1$ $\mathrm{dom}(q) = \mathrm{dom}(p) \cap N_u \cap \beta$*
  - *$\bullet_2$ If $\gamma \in \mathrm{dom}(q)$ then $q(\gamma) = p(\gamma) \cap N_u$.*

(7) If (A) then (B), where:

Paper Sh:1258, version 2026-09-29_3. See https://shelah.logic.at/papers/1258/ for possible updates.

(A) (a) $i_* < \lambda$,
(b) $p_i \in \mathbb{P}'_{\mathbf{q}}$ *for* $i < i_*$,
(c) *if* $i < j < i_*$, *then* p_i *and* p_j *are essentially comparable, i.e.:*
- *if* $\beta \in \mathrm{dom}(p_i) \cap \mathrm{dom}(p_j)$ *then* $p_i(\beta) \subseteq p_j(\beta)$ *or* $p_j(\beta) \subseteq p_i(\beta)$.

(d) $\bar{p} = \langle p_i : i < i_* \rangle$.

(B) $\bar{p}$ *has a lub* p *called* $\lim(\bar{p})$ *or* $\lim(\{p_i : i < i_*\})$ *defined by:*
- $\mathrm{dom}(p) = \bigcup\{\mathrm{dom}(p_i) : i < i_*\}$,
- *if* $\beta \in \mathrm{dom}(p)$, *then*

$$p(\beta) = \bigcup\{p_i(\beta) : i < i_* \text{ satisfying } \beta \in \mathrm{dom}(p_i)\}.$$

Proof. Part (2) is crucial but easy to verify. Parts (0), (1), (3), and (4) are also easy.

(5) For this, define $\mathbf{q} := \lim(\bar{\mathbf{q}})$ naturally, but we elaborate.

(∗) (a) $\mathrm{lg}(\mathbf{q}) = \bigcup\{\mathrm{lg}(\mathbf{q}_i) : i < \delta\}$,
(b) if $i < \delta$ and $\alpha \le \mathrm{lg}(\mathbf{q}_i)$, then $(\mathbb{P}_{\mathbf{q},\alpha}, \mathbb{P}'_{\mathbf{q},\alpha}) = (\mathbb{P}_{\mathbf{q}_i,\alpha}, \mathbb{P}'_{\mathbf{q}_i,\alpha})$,
(c) if $i < \delta$ and $\beta < \mathrm{lg}(\mathbf{q}_i)$, then $(\mathbb{Q}_{\mathbf{q},\beta}, \mathbf{a}_{\mathbf{q},\beta}, \mathbf{b}_{\mathbf{q},\beta}) = (\mathbb{Q}_{\mathbf{q}_i,\beta}, \mathbf{a}_{\mathbf{q}_i,\beta}, \mathbf{b}_{\mathbf{q}_i,\beta})$,
(d) $(\mathbb{P}_{\mathbf{q},\mathrm{lg}(\mathbf{q})}, \mathbb{P}'_{\mathbf{q},\mathrm{lg}(\mathbf{q})})$ is $(\bigcup\{\mathbb{P}_{\mathbf{q}_i} : i < \delta\}, \bigcup\{\mathbb{P}'_{\mathbf{q}_i} : i < \delta\})$ when $\mathrm{cf}(\delta) \ge \lambda$,
(e) if $\mathrm{cf}(\delta) < \lambda$, then $(\mathbb{P}_{\mathbf{q},\mathrm{lg}(\mathbf{q})}, \mathbb{P}'_{\mathbf{q},\mathrm{lg}(\mathbf{q})})$ are defined as inverse limit. Then,
- $\mathbb{P}'_{\mathbf{q}} := \mathbb{P}'_{\mathbf{q},\mathrm{lg}(\mathbf{q})}$ is dense in $\mathbb{P}_{\mathbf{q}}$ because by Definition 1.2(3), for each $\beta < \mathrm{lg}(\mathbf{q}_j)$ with $j < \delta$, $\mathbb{Q}_{\mathbf{b}[\beta,\mathbf{q}_j]}$ is closed under increasing unions of length $< \lambda$.

Recalling that in Definition 1.3(1)(c), we use β and not α, "$\mathbb{P}_{\mathbf{q}}$ satisfies the λ^+-cc" is not required for proving 1.5 (5), only "if $\beta < \mathrm{lg}(\mathbf{q})$ then $\mathbb{P}_{\mathbf{q},\beta}$ satisfies the λ^+-cc", which is clear. Note that even though we formally do not need it here, the chain condition of $\mathbb{P}_{\mathbf{q}}$ will be proved in claim 1.6.

(6) Note that:

(a) If $\gamma \in \mathrm{dom}(q)$ then $\gamma \in N_u$ and $q(\gamma) \subseteq N_u$,
(b) As $\mathrm{dom}(q)$ and $q(\gamma)$ for $\gamma \in \mathrm{dom}(q)$ has cardinality $< \lambda$ and $[N_u]^{<\lambda} \subseteq N_u$ so recalling clause (a) obviously $q \in N_u$.
(c) To prove q is in $\mathbb{P}'_{\mathbf{q}}$ we need, for $\gamma \in \mathrm{dom}(q)$ and $\zeta_1 < \zeta_2$ from $q(\gamma) \subseteq \mathscr{U}_\gamma$ to verify the condition in 1.5(2)(c).
(d) But as $\gamma \in N_u$ hence $\mathbf{q} \restriction (\gamma + 1)$ and ζ_1, ζ_2 belong to N_u, also $N_{\mathbf{q},\gamma,\{\zeta_1\}}$, $N_{\mathbf{q},\gamma,\{\zeta_2\}}$, $N_{\mathbf{q},\gamma,\{\zeta_1,\zeta_2\}}$ belong to N_u hence are included in it so we can finish easily.

(7) Follows by our definitions. $\square_{1.5}$

We now arrive to the

Crucial Claim 1.6. *If* $\mathbf{q} \in \mathbf{Q}_{\mathbf{p}}$ *then* $\mathbb{P}_{\mathbf{q}}$ *satisfies* λ^+*-cc. Moreover* $\mathbb{P}_{\mathbf{q}}$ *is* λ^+*-Knaster.*

Proof. It suffices, by 1.3(1)(e), to prove that $\mathbb{P}'_{\mathbf{q}} = \mathbb{P}'_{\mathbf{q},\mathrm{lg}(\mathbf{q})}$ satisfies the λ^+-cc, so assume:

$(*)_1$ (a) Let $\bar{p} = \langle p_\xi : \xi < \lambda^+ \rangle$, where $p_\xi \in \mathbb{P}'_{\mathbf{q}}$,
(b) it suffices to prove that for some $\zeta < \xi < \lambda^+$, p_ζ and p_ξ are compatible.

[Why? By the definitions.]

$(*)_2$ For some stationary set $S \subseteq \mathrm{cof}(\lambda) \cap \lambda^+$, we have:
- $\bullet_1$ $\langle \mathrm{dom}(p_\xi) : \xi \in S \rangle$ is a Δ-system with heart $w_* \in [\mathrm{lg}(\mathbf{q})]^{<\lambda}$, and
- $\bullet_2$ if $\beta \in w_*$ then $\langle p_\xi(\beta) : \xi \in S \rangle$ is a Δ-system.

[Why? By the Delta system lemma, the proof using Fodor's lemma recalling $\lambda = \lambda^{<\lambda}$.]

Paper Sh:1258, version 2026-09-29_3. See https://shelah.logic.at/papers/1258/ for possible updates.

$(*)_3$ Without loss of generality, $\langle p_\xi : \xi \in S\rangle$ are pairwise isomorphic (see Definition 1.3(4)).

[Why? Easy because for every $\mathbf{a}, u$ the model $N_{\mathbf{a},u}$ has cardinality λ.]

$(*)_4$ For $\gamma < \beta$ from w_*, we have:

- $\bullet_1$ Let $\mathscr{W}_\beta, u_{\beta,\gamma}$ be as in 1.3(1)(d)$\bullet_5$.
- $\bullet_2$ Without loss of generality, $u_{\beta,\gamma}$ is disjoint to $N_{\mathbf{q},\beta,\{\zeta\}} \setminus N_{\mathbf{q},\beta,\emptyset} \cap \mu$ for every $\zeta \in \mathscr{U}_\beta$ and is disjoint to $N_{\mathbf{q},\beta,\{\varepsilon,\zeta\}} \setminus N_{\mathbf{q},\beta,\emptyset} \cap \mu$ for every $\varepsilon < \zeta$ from $\mathscr{U}_\beta$.

[Why? As for any $\gamma < \beta$ from w_* we have to omit from $\mathscr{U}_\beta$ at most two elements and w_* has cardinality $< \lambda$.]

$(*)_5$ We fix $\xi(1) \neq \xi(2)$ from S and we shall prove that $p_{\xi(1)}$ and $p_{\xi(2)}$ have a common upper bound; this suffices for proving the Crucial Claim 1.6.

$(*)_6$ For $\beta \in w_*$:

- (a) for $\ell \in \{1,2\}$, consider the sequence $\langle \alpha^\beta_{\xi(\ell),\varepsilon} : \varepsilon < \varepsilon_\beta\rangle$ listing the set $p_{\xi(\ell)}(\beta)$ in increasing order
- (b) Why ε_β and not $\varepsilon_{\beta,\ell}$? as the two sequences have the same length because $p_{\xi(1)}, p_{\xi(2)}$ are isomorphic, see Definition 1.3(4) $\bullet_1$.
- (c) Let $\mathscr{S}_\beta := \{\varepsilon < \varepsilon_\beta : \alpha^\beta_{\xi(1),\varepsilon} \neq \alpha^\beta_{\xi(2),\varepsilon}\}$,
- (d) so by Definition 1.3 (4) $\bullet_2$ the sets $\{\alpha^\beta_{\xi(1),\varepsilon} : \varepsilon \in \mathscr{S}_\beta\}$, $\{\alpha^\beta_{\xi(2),\varepsilon} : \varepsilon \in \mathscr{S}_\beta\}$ are disjoint and disjoint to $\{\alpha^\beta_{\xi(1),\varepsilon} : \varepsilon \in \varepsilon_\beta \setminus \mathscr{S}_\beta\} = \{\alpha^\beta_{\xi(2),\varepsilon} : \varepsilon \in \varepsilon_\beta \setminus \mathscr{S}_\beta\}$.

Let $\bar\beta = \langle \beta_i : i \le i_*\rangle$ list the closure of $\{\alpha, \alpha+1 : \alpha \in w_*\} \cup \{0, \lg(\mathbf{q})\}$ in increasing order, so necessarily $i_* < \lambda$ and clearly it suffices:

$(*)_7$ To choose $q_i \in \mathbb{P}'_{\mathbf{q},\beta_i}$ a common upper bound of $\{p_{\xi(1)} \restriction \beta_i, p_{\xi(2)} \restriction \beta_i\}$ increasing with $i \le i_*$ by induction on $i \le i_*$ such that:

- $(*)$ If $\beta \in w_* \setminus \{\beta_j : j < i\}$ and $\zeta(1), \zeta(2)$ are from $\mathscr{S}_\beta$ then:
  - $\bullet_1$ $\mathrm{dom}(q_j) \cap N_{\beta,\{\alpha_{\xi(1),\zeta(1)},\alpha_{\xi(2),\zeta(2)}\}}$ is a subset of
    $$N_{\beta,\{\alpha_{\xi(1),\zeta(1)}\}} \cup N_{\beta,\{\alpha_{\xi(2),\zeta(2)}\}} \cup N_{\beta,\emptyset},$$
  - $\bullet_2$ if $\ell = 1,2$ and $\gamma \in \mathrm{dom}(q_j) \cap N_{\beta,\{\alpha_{\xi(\ell),\zeta(\ell)}\}}$ then $q_i(\gamma) = p_{\xi(\ell)}(\gamma)$ or $\gamma \in N_{\beta,\emptyset}$

Let us carry the induction.

Case 1: $i = 0$. Clearly, this case is trivial, letting $q_0 = \emptyset$.

Case 2: i is a limit ordinal.

In this case, let $q_i := \lim\langle q_j : j < i\rangle$, so by Claim 1.5(1), q_i is well-defined and is as required by the definition of the order and satisfies $(*)_7$.

Case 3: $i = j+1$ and $\beta_j \notin w_*$.

In this case, $\mathrm{dom}(p_{\xi(1)}) \cap \mathrm{dom}(p_{\xi(2)}) \cap \beta_i \subseteq \beta_j$, hence the condition
$$q_i := q_j \cup \big(p_{\xi(1)} \restriction [\beta_j, \beta_i]) \cup (p_{\xi(2)} \restriction [\beta_j, \beta_i])\big)$$
is as promised.

Case 4: $i = j+1$ and $\beta_j \in w_*$.

By the choice of $\bar\beta$, clearly $\beta_i = \beta_j + 1$ and let $\mathscr{S} = \mathscr{S}_{\beta_j}$.

Recalling 1.3(1)(d) and 0.7(b)($\bullet_8$), we have:

$(*)_8$ $\mathbf{a}_{\beta_j} = \mathbf{a}_{\mathbf{q},\beta_j}$ determine:

- (a) $\bar\pi_{\beta_j} = \langle \pi_{u,v} : u, v \in [\mathscr{U}_{\beta_j}]^{\le 2}$ and $|u| = |v|\rangle$,
- (b) $\bar N_{\beta_j} = \langle N_u : u \in [\mathscr{U}_{\beta_j}]^{\le 2}\rangle$,
- (c) for $\varepsilon(1), \varepsilon(2) \in \mathscr{S}$, let:
  - $v[\varepsilon(1), \varepsilon(2)] = \{\alpha_{\xi(1),\varepsilon(1)}, \alpha_{\xi(1),\varepsilon(2)}\}$, and
  - $u[\varepsilon(1), \varepsilon(2)] = \{\alpha_{\xi(1),\varepsilon(1)}, \alpha_{\xi(2),\varepsilon(2)}\}$.

Paper Sh:1258, version 2026-09-29_3. See https://shelah.logic.at/papers/1258/ for possible updates.

(d) for $\varepsilon \in \mathscr{S}$, let $v[\varepsilon] = \{\alpha_{\xi(1),\varepsilon}\}$ and $u[\varepsilon] = \{\alpha_{\xi(2),\varepsilon}\}$,
(e) $\bar{\iota} = \bar{\iota}^*_{\beta_j}$, see 1.3 (1) (d) $\bullet_1$.
(f) $\gamma_j = \xi_{\mathbf{q}}(\beta_j)$; see 1.3(1)(d) $\bullet_3$.

We shall now define $p_{\varepsilon(1),\varepsilon(2)}$ for $\varepsilon(1), \varepsilon(2) \in \mathscr{S}$ such that:

$(*)_9$ (a) $p_{\varepsilon(1),\varepsilon(2)} \in \mathbb{P}_{\gamma_j} \cap N_{u[\varepsilon(1),\varepsilon(2)]}$, hence $\mathrm{dom}(p_{\varepsilon(1),\varepsilon(2)}) \subseteq \gamma_j \cap N_{u[\varepsilon(1),\varepsilon(2)]}$,
(b) if $\varepsilon(1) = \varepsilon(2)$, then $p_{\varepsilon(1),\varepsilon(2)} \restriction (\gamma_j \cap N_{v[\varepsilon(1)]})$, $p_{\xi(1)} \restriction N_{v[\varepsilon(1)]}$ are essentially comparable; see 1.5(7)(A)(c), moreover the first is $\leq_{\mathbb{P}_{\mathbf{q}}}$-above the second,
(c) if $\varepsilon(1) = \varepsilon(2)$, then $p_{\varepsilon(1),\varepsilon(2)} \restriction (\gamma_j \cap N_{u[\varepsilon(2)]})$, $p_{\xi(2)} \restriction N_{u[\varepsilon(2)]}$ are essentially comparable, moreover the first is $\leq_{\mathbb{P}_{\mathbf{q}}}$-above the second,
(d) $p_{\varepsilon(1),\varepsilon(2)}$ satisfies 1.3(1)(e)$\bullet$ with $(\gamma_j, \varepsilon(1), \varepsilon(2))$ here standing for $(\beta, \zeta_1, \zeta_2)$ there,
(e) $\{q_j \restriction N_\emptyset\} \cup \{p_{\varepsilon(1),\varepsilon(2)} \restriction N_\emptyset : \varepsilon(1), \varepsilon(2) \in \mathscr{S}\}$ are pairwise essentially comparable,
(f) if $\varepsilon(1) \neq \varepsilon(2)$ then $p_{\varepsilon(1),\varepsilon(2)} \restriction N_{\{\alpha_{\varepsilon(\ell)}\}} \leq p_{\xi(\ell)} \restriction N_{\{\alpha_{\varepsilon(\ell)}\}}$ for $\ell = 1, 2$.
(g) if $\mathscr{S}_* \subseteq \mathscr{S} \times \mathscr{S}$ then the lub $q_{\mathscr{S}_*}$ of $\{q_j[N_{u[\varepsilon(1),\varepsilon(2)]} : \varepsilon(1), \varepsilon(2) \in \mathscr{S}_*\}$ satisfies the condition in $(*)_7$.

We have to show two things: $\boxplus_1$ and $\boxplus_2$. The first says we can choose them (the $p_{\varepsilon(1),\varepsilon(2)}$-s), the second that this is enough.

$\boxplus_1$ we can choose $p_{\varepsilon(1),\varepsilon(2)}$ for $\varepsilon(1), \varepsilon(2) \in \mathscr{S}$ as required in $(*)_7$.

We consider two possible cases:

Case 4.1: $\varepsilon(1) \neq \varepsilon(2)$.

Let $p_{\varepsilon(1),\varepsilon(2)} = \pi(p_{\xi(1)} \restriction N_{v[\varepsilon(1),\varepsilon(2)]})$, where $\pi = \pi_{u[\varepsilon(1),\varepsilon(2)],v[\varepsilon(1),\varepsilon(2)]}$.

Why is $(*)_9$ preserved? Most clauses are obvious, but $(*)_9(g)$ deserve elaboration, recalling that we have to satisfy $(*)_7$.

So let $\beta \in \mathscr{W}_* \setminus \{\beta_\iota : \iota < i\}$, hence for some $j(*) < i_*$, we have $\beta = \beta_{j(*)}$, hence we have $\beta_{j(*)} \geq \beta_i$ hence $\beta_{j(*)} > \beta_j$ and we have $\mathscr{S}_* \subseteq \mathscr{S} \times \mathscr{S}$ and deal with $q_{\mathscr{S}_*}$.

For this, it is enough to consider the cases:

$\oplus_1$ $\mathscr{S}_* = \{\zeta(1), \zeta(2)\}$, where $\zeta(1) = \varepsilon(1)$ and $\zeta(2) = \varepsilon(2)$ hence from $\mathscr{S}$, so $\zeta(1) \neq \zeta(2)$,
$\oplus_2$ $\mathscr{S}_* = \{\zeta(1), \zeta(2)\}$ where $\zeta(1) \neq \zeta(2)$ are from $\mathscr{S}$ but $(\zeta(1), \zeta(2)) \neq (\varepsilon(1), \varepsilon(2))$.

Easy to check.

Case 4.2: $\varepsilon(1) = \varepsilon(2)$.

In this case, we pick some sequence $\langle p_{\varepsilon,\varepsilon} : \varepsilon \in \mathscr{S}\rangle$ by choosing $p_{\varepsilon,\varepsilon}$ by induction on $\varepsilon \in \mathscr{S}$. Now, $p_{\varepsilon,\varepsilon} \in \mathbb{P}'_{\beta_j} \cap N_{u[\varepsilon(1),\varepsilon(2)]}$ is such that:

$(*)$ (a) $p_{\varepsilon,\varepsilon}$ is $\leq_{\mathbb{P}'_{\mathbf{q},\beta_j}}$-above $p_{\xi(1)} \restriction N_{v[\varepsilon]}$ and above the restriction $p_{\xi(2)} \restriction N_{u[\varepsilon]}$,
(b) $\langle p_{\zeta,\zeta} \restriction N_\emptyset : \zeta \in (\varepsilon + 1) \cap \mathscr{S}\rangle$ is $\leq_{\mathbb{P}_{\beta[j]}}$-increasing, and
(c) there are q_1, q_2, r_1, r_2 as in Definition 1.3(2)(c) $(\bullet_1)$-$(\bullet_5)$ with $\mathbf{b}_{\mathbf{q},\beta_j}$ standing here for $(\mathbf{a}, p, \bar{\iota})$ there such that:

$$\bigvee_{\ell=1}^{2} (\forall \gamma \in \mathrm{dom}(r_\ell))\, [\gamma \in \mathrm{dom}(p_{\varepsilon,\varepsilon}) \wedge r_\ell(\gamma) \subseteq p_{\varepsilon,\varepsilon}(\gamma)]\,.$$

We can choose $p_{\varepsilon,\varepsilon}$ by the properties of $\mathbf{b}_{\beta_j}$

Having defined all the $p_{\varepsilon(1),\varepsilon(2)}$-s we can proceed.

$\boxplus_2$ The following set of members of $\mathbb{P}_{\beta_i}$ has a common upper bound q_*:
- $p_{\xi(1)}, p_{\xi(2)}$, and
- $p_{\varepsilon(1),\varepsilon(2)}$ for $\varepsilon(1), \varepsilon(2) \in \mathscr{S}$.

[Why? Recall Claim 1.5(2) and 1.2(1)(c)$(\bullet_1)$ by 1.5(7), clause (A) there holds, in particular sub-clause (A)(c). The main point is that:

Paper Sh:1258, version 2026-09-29_3. See https://shelah.logic.at/papers/1258/ for possible updates.

(∗) $\langle N_{v[\varepsilon(1),\varepsilon(2)]} \cap \gamma_j \setminus (N_{v[\varepsilon(1)]} \cup N_{u[\varepsilon(1)]})\colon \varepsilon(1),\varepsilon(2) \in \mathscr{S}\rangle$ is a sequence of pairwise disjoint sets.

Why? As "$N_u \cap N_v \subseteq N_{u\cap v}$ for $u,v \in [\mathscr{U}_{\beta_j}]^{<2}$ by $0.7\bullet_7$.
So q_* from $\boxplus_2$ is a common upper bound of $p_{\xi(1)}, p_{\xi(2)}$, as promised. $\square_{1.6}$

Remark 1.7. 1) No need so far, but we may add in $(*)_4$ of the proof of Crucial Claim 1.6 the following item:

(d) if $\beta \in w_*$ and $\langle \alpha_{\zeta,\beta,i}\colon i < \iota_{\zeta,\beta}\rangle$ lists in increasing order the members of $p_\zeta(\beta)$ for $\zeta \in S$, then:
- $\langle \iota_{\zeta,\beta}\colon \zeta \in S\rangle$ is constant called i_β,
- for $i < i_\beta$, the sequence $\langle \alpha_{\zeta,\beta,i}\colon \zeta \in S\rangle$ is constant or increasing,
- if $i,j < i_\beta$ the sequence of truth values

  $\langle \text{Truth value}(\alpha_{\zeta,\beta,i} < \alpha_{\xi,\beta,j})\colon \zeta < \xi \text{ are from } S\rangle$

  is constant, and
- if $i,j < i_\beta$, $\zeta \neq \xi$ are from S and $\alpha_{\zeta,\beta,i} = \alpha_{\xi,\beta,j}$ then $i = j$.

2) We can make our choice of q_1,q_2,r_1,r_2 canonical, that is:

(A) In 1.2(2) we replace $(\mathbf{a},p,\bar\iota)$ by $(\mathbf{a},p,\bar i,\mathbb{F})$, where:
- $\bullet_1$ $\mathbb{F}_{\zeta_1,\zeta_2}(q) = (q_1,q_2,r_1,r_2) = \langle \mathbb{F}_{\zeta_1,\zeta_2,\ell}(q) : \ell = 1,2,3,4\rangle$
- $\bullet_2$ if also $\zeta_3 < \zeta_4$ are from $\mathscr{U}$ then $\pi^{\mathbf{a}}_{\zeta_3,\zeta_4,\zeta_1,\zeta_2}\mathbb{F}_{\zeta_1,\zeta_2,\ell} = \mathbb{F}_{\zeta_3,\zeta_4,\ell}$, where if $p \le q \in \mathbb{P}_{\mathbf{a}} \cap N_{\mathbf{a},\{\varepsilon[\mathbf{a}]\}}$ and $\zeta_1 < \zeta_2$ are from $\mathscr{U}$, then $\langle \mathbb{F}_{\zeta_1,\zeta_1,\ell}(p,q)\colon \ell < \mu\rangle$ is the quadruple (q_1,q_2,r_1,r_2) as in 1.2(1)(c)($\bullet_1$)-($\bullet_5$).

(B) In 1.2(3) similarly and in 1.3(1)(d)

(C) In 1.5(1)(d) use $\mathbb{F}_\beta$,

(D) In the proof of 1.6, in $(*)_7\boxplus_1$, case $4.2(*)_{4.2}$ we use $\mathbb{F}_{\beta_j}$,

(E) Update the proof of 1.8 accordingly.

Claim 1.8. *If (A) then (B), where:*

(A) *(a)* $\mathbf{q} \in \mathbf{Q}_{\mathbf{p}}$,
(b) $2 < \sigma < \lambda$,
(c) $\underset{\sim}{\mathbf{c}}$ *is a* $\mathbb{P}_{\mathbf{q}}$*-name of a function from* $[\theta]^2$ *into* σ.
(d) $p_* \in \mathbb{P}_{\mathbf{q}}$.

(B) *There is some* $\mathbf{b} \in \mathbf{A}^+$ *such that* $\mathbb{P}_{\mathbf{b}} = \mathbb{P}'_{\mathbf{q}}$ *and* $\underset{\sim}{\mathbf{c}}_{\mathbf{b}} = \underset{\sim}{\mathbf{c}}$ *and* $p_* \le_{\mathbb{P}_{\mathbf{q}}} p_{\mathbf{b}}$.

Proof. Recalling Hypothesis 1.1(b), on the one hand, it is clear how to choose $\mathbf{a} \in \mathbf{A}$ such that $\mathbb{P}_{\mathbf{a}} = \mathbb{P}'_{\mathbf{q}}$ and $\underset{\sim}{\mathbf{c}}_{\mathbf{a}} = \underset{\sim}{\mathbf{c}}$. On the other hand, the choice of $p_{\mathbf{b}}$ and $\bar\iota_{\mathbf{b}}$ is similar to the proof of [She88, 2.1]. We now elaborate.

First, we can find $\mathbf{a}$ such that:

$(*)^1_{\mathbf{a}}$ (a) $\mathbf{a} \in \mathbf{A}$,
(b) $\mathbb{P}_{\mathbf{a}} = \mathbb{P}'_{\mathbf{q}}$,
(c) $\gamma = \lg(\mathbf{q})$,
(d) $\underset{\sim}{\mathbf{c}}_{\mathbf{a}} = \underset{\sim}{\mathbf{c}}$.

Why can we find? Because we have chosen $\mathbb{P}_{\mathbf{a}}$ as in $(*)^1_{\mathbf{a}}$(b), it is λ^+-cc by Claim 1.6; also $\gamma, \underset{\sim}{\mathbf{c}}_{\mathbf{a}}$ are as is required in Definition 1.2. Lastly we can choose $(\mathscr{U}_{\mathbf{a}}, \bar N)$ as required because $\theta \to_{\rm sq} (\partial)^{\lambda,2}_\lambda$ holds by Hypothesis 1.1 clause (b) and 0.7 in particular clause $(b)\bullet_{10}$.

We are left with choosing some appropriate $(p,\bar\iota)$ and then let $\mathbf{b} = (\mathbf{a},p,\bar\iota)$. Let

$$Y \coloneqq \{(q_1,q_2)\colon q_1,q_2 \in \mathbb{P}'_{\mathbf{a}} \cap N_{\mathbf{a},\{\varepsilon[\mathbf{a}]\}} \text{ are above } p_* \text{ and, } q_1 \restriction (N_{\mathbf{a},\emptyset} \cap \lg(\mathbf{q})) = q_2 \restriction (N_{\mathbf{a},\emptyset} \cap \lg(\mathbf{q}))\},$$

and let $\le_Y$ be the following two place relation on Y:

$(*)_2$ $(p_1,p_2) \le_Y (q_1,q_2)$ iff:

Paper Sh:1258, version 2026-09-29_3. See https://shelah.logic.at/papers/1258/ for possible updates.

(a) $(p_1,p_2)\in Y$ and $(q_1,q_2)\in Y$,
(b) $p_1\le_{\mathbb{P}'_{\mathbf{q}}} q_1$ and $p_2\le_{\mathbb{P}'_{\mathbf{q}}} q_2$.

Clearly,

$(*)_3$ $(Y,\le_Y)$ is a $(<\lambda)$-complete partial order.

[Why? Recalling 1.5(1).]

$(*)_4$ For $(p_1,p_2)\in Y$, let
(a) $\mathrm{solv}(p_1,p_2)$ be the set of pairs (ι_0,ι_1) such that for any $\zeta_1<\zeta_2$ from $\mathscr{U}_{\mathbf{a}}$, there are r_1,r_2 such that for $\ell=1,2$ clauses $\bullet_2$-$\bullet_5$ of Definition 1.2(2)(c) hold.
(b) $\mathrm{solv}^+(p_1,p_2):=\bigcap\{\mathrm{solv}(q_1,q_2)\colon (p_1,p_2)\le_Y(q_1,q_2)\in Y\}$.

$(*)_5$ (a) if $(p_1,p_2)\le_Y(q_1,q_2)$ then:

$$\mathrm{solv}(p_1,p_2)\supseteq\mathrm{solv}(q_1,q_2)\supseteq\mathrm{solv}^+(q_1,q_2)\supseteq\mathrm{solv}^+(p_1,p_2),$$

(b) if $(p_1,p_2)\in Y$ then $\mathrm{solv}(p_1,p_2)\ne\emptyset$.

[Why? The first inclusion in Clause (a) holds because $\le_{\mathbb{P}_{\mathbf{q}}}$ is transitive. The other inclusions are clear, and Clause (b) is easy too.]

$(*)_6$ If $(p_1,p_2)\in Y$ then for some (q_1,q_2) and $\bar\iota$, we have:
(a) $(p_1,p_2)\le_Y(q_1,q_2)\in Y$,
(b) if $(q_1,q_2)\le_Y(q'_1,q'_2)$ then $\bar\iota\in\mathrm{solv}(q'_1,q'_2)$, moreover, $\mathrm{solv}(q_1,q_2)=\mathrm{solv}(q'_2,q'_2)=\mathrm{solv}^+(q'_1,q'_2)=\mathrm{solv}^+(q_1,q_2)$.

[Why? Recalling $\sigma<\lambda$, hence $|\sigma\times\sigma|<\lambda$ and $(Y,\le_Y)$ is λ-complete by $(*)_3$.]

$(*)_7$ For $p\in\mathbb{P}'_{\mathbf{a}}\cap N_{\mathbf{a},\{\varepsilon[\mathbf{a}]\}}$, let $\mathrm{solv}(p)$ be the set of $\bar\iota\in\sigma\times\sigma$ such that there is (q_1,q_2) such that:
$\bullet_1$ $p\le_{\mathbb{P}_{\mathbf{q}}}q_1$, $p\le_{\mathbb{P}_{\mathbf{q}}}q_2$ and
$\bullet_2$ $(q_1,q_2)\in Y$,
$\bullet_3$ $\bar\iota\in\mathrm{solv}^+(q_1,q_2)$,
$\bullet_4$ $\mathrm{solv}(q_1,q_2)=\mathrm{solv}^+(q_1,q_2)$.

$(*)_8$ (a) if $p\in\mathbb{P}'_{\mathbf{a}}\cap N_{\mathbf{a},\{\varepsilon[\mathbf{a}]\}}$ then $\mathrm{solv}(p)\ne\emptyset$,
(b) if $p\le_{\mathbb{P}'_{\mathbf{a}}}q$ are from $\mathbb{P}'_{\mathbf{a}}\cap N_{\mathbf{a},\{\varepsilon[\mathbf{a}]\}}$ then $\mathrm{solv}(p)\supseteq\mathrm{solv}(q)$,
(c) if $p\in\mathbb{P}'_{\mathbf{a}}\cap N_{\mathbf{a},\{\varepsilon[\mathbf{a}]\}}$ then for some q and $\bar\iota$, for every q', we have $q\le_{\mathbb{P}'_{\mathbf{q}}}q'\wedge q'\in\mathbb{P}'_{\mathbf{a}}\cap N_{\mathbf{a},\{\varepsilon[\mathbf{a}]\}}\Rightarrow\bar\iota\in\mathrm{solv}(q')$.

[Why? Clause (a) follows by $(*)_6$, Clause (b) by the definitions, and Clause (c) holds as $\mathbb{P}'_{\mathbf{a}}$ and even $\mathbb{P}'_{\mathbf{a}}\cap N_{\mathbf{a},\{\varepsilon[\mathbf{a}]\}}$ is λ-complete and $|\sigma\times\sigma|<\lambda$.]

Now, applying $(*)_8$(c) to p_* finish the proof of 1.8. $\square_{1.8}$

Claim 1.9. *If (A) then (B), where:*

(A) *(a)* $\mathbf{q}\in\mathbb{Q}_{\mathbf{p}}$ *and* $\mathbf{q}_0<_{\mathbf{p}}\mathbf{q}$,
(b) $\gamma(\mathbf{q})<\mu$, *so* $\lg(\mathbf{q})<\mu$,
(c) $\mathbf{b}\in\mathbf{A}_{\mathbf{p}}$ *and* $\mathbb{P}_{\mathbf{b}}=\mathbb{P}_{\mathbf{q}_0}$.

(B) There exists some $\mathbf{q}_1$ *such that:*
(a) $\mathbf{q}\le_{\mathbf{p}}\mathbf{q}_1$,
(b) $\lg(\mathbf{q}_1)=\lg(\mathbf{q})+1$,
(c) $\mathbf{b}_{\lg(\mathbf{q})}[\mathbf{q}_1]=\mathbf{b}$.

Proof. Easy. $\square_{1.9}$

Lastly, before arriving at the main conclusion, we have to prove the following.

Claim 1.10.

(1) Assume $\mathbf{q}\in\mathbf{Q}_{\mathbf{p}}$, $\alpha<\lg(\mathbf{q})$ *and* $\mathbf{b}=\mathbf{b}_{\mathbf{q},\alpha}=(\mathbf{a}_\alpha,p_\alpha,\bar\iota_\alpha)=(\mathbf{a},p,\bar\iota)$, *then:*

- $\Vdash_{\mathbb{P}_{\mathbf{q},\alpha+1}}$ "$\mathcal{V}_{\mathbb{Q}_{\mathbf{b}}}\in[\mathscr{U}_{\mathbf{a}_\alpha}]^\partial$ *and for every* $\alpha\ne\beta\in\mathcal{V}_{\mathbb{Q}_{\mathbf{b}}}$, $\mathbf{c}_{\mathbf{a}_\alpha}\{\alpha,\beta\}\in\{\iota_1,\iota_2\}$".

Paper Sh:1258, version 2026-09-29_3. See https://shelah.logic.at/papers/1258/ for possible updates.

(2) If $\mathbf{b} = (\mathbf{a}, p, \bar{\iota}) \in \mathbf{A}^+$, $\mathrm{cf}(\partial) > \lambda$, *and in* $\mathbf{V}^{\mathbb{P}_{\mathbf{a}}}$, $\mathbb{Q}_{\mathbf{b}}$ *satisfies the* λ^+*-cc, then for some* $p \in \mathbb{Q}_{\mathbf{b}} \cap \mathbb{P}_{\mathbf{a}} \cap N_{\mathbf{a},\{\varepsilon[\mathbf{a}]\}}$ *we have*[4] $p \Vdash_{\mathbb{Q}_{\mathbf{b}}}$ "$\mathcal{V}_{\mathbb{Q}_{\mathbf{b}}} \in [\mathscr{U}_{\mathbf{a}}]^{\partial}$ *and for every* $\alpha \neq \beta \in \mathcal{V}_{\mathbb{Q}_{\mathbf{b}}}$, $\mathbf{c}_{\mathbf{a}}\{\alpha, \beta\} \in \{\iota_1, \iota_2\}$".

Proof. (1) The second phrase in both conclusion holds by the definitions of $\mathbb{Q}_{\mathbf{b}}$.

By the proof of "$\mathbb{P}_{\mathbf{q}}$ satisfies the λ^+-cc", we can show for $\varepsilon < \partial$, the density of the set

$$\mathscr{I}_\varepsilon := \{p \in \mathbb{P}'_{\mathbf{q}} \colon \alpha \in \mathrm{dom}(p) \text{ and there is } \beta \in p(\alpha) \text{ such that } \varepsilon < \mathrm{otp}(\mathscr{U}_{\mathbf{a}_\alpha} \cap \beta)\}.$$

(2) Easily, for every $\beta \in \mathscr{U}_{\mathbf{a}}$ we can choose $p^0_\beta = \{\beta\}$, $q_\beta = \{(p, p^0_\beta)\}$. Clearly, $q_\beta \in \mathbb{P}_\alpha * \mathbb{Q}_{\mathbf{b}}$ for $\beta \in \mathscr{U}_{\mathbf{a}}$. So by the λ^+-cc for some $\beta \in \mathscr{U}_{\mathbf{a}}$, $q_\beta \Vdash$"$\{\varepsilon \in \mathscr{U}_{\mathbf{a}} \colon q_\varepsilon \in \mathbb{Q}_{\mathbf{b}}\} \in [\mathscr{U}_{\mathbf{a}}]^{\partial}$; well assuming $\mathrm{cf}(\partial) > \lambda$. $\square_{1.10}$

Conclusion 1.11. *There exists a forcing notion* $\mathbb{P}$ *satisfying the following conditions:*

- *(a)* $\mathbb{P}$ *is* λ^+*-cc of cardinality* μ.
- *(b)* $\mathbb{P}$ *is* $(<\lambda)$*-complete; hence, it collapses no cardinals, changes no cofinalities, and preserves cardinal arithmetic outside the interval* $[\lambda, \mu)$.
- *(c)* $\Vdash_{\mathbb{P}}$ "$2^\lambda = \mu$".
- *(d)* $\Vdash_{\mathbb{P}}$ "$\theta \to [\partial]^2_{\sigma,2}$" *for every* $\sigma \in (2, \lambda)$.

Proof. Choose a $\leq_{\mathbf{p}}$-increasing continuous sequence $\langle \mathbf{q}_\alpha \colon \alpha < \mu \rangle \in {}^{\mu}(\mathbf{Q}_{\mathbf{p}})$ such that $\mathrm{lg}(\mathbf{q}_\alpha) = \alpha$, $\mathbb{P}_{\mathbf{q}_\alpha}$ has cardinality $\leq (|\alpha| + \lambda)^{<\lambda}$ and,

- if $\alpha < \mu$ and $\Vdash_{\mathbb{P}_{\mathbf{q}_\alpha}}$ "$\underset{\sim}{\mathbf{c}} \colon [\theta]^2 \to \sigma$", then for unboundedly many $\beta \in [\alpha, \mu)$, $\underset{\sim}{\mathbf{c}}_{\mathbf{q}_{\beta+1},\beta} = \underset{\sim}{\mathbf{c}}$.

The existence of $\mathbf{b}_\beta[\mathbf{q}_{\beta+1}]$ with $\underset{\sim}{\mathbf{c}}[\mathbf{b}_\beta[\mathbf{q}_{\beta+1}]] = \underset{\sim}{\mathbf{c}}$ as required holds by Claim 1.8 and Claim 1.9 below.

Clearly $\bigcup\{\mathbb{P}_{\mathbf{q}_\beta} \colon \beta < \mu\}$ is a forcing notion as is required. $\square_{1.11}$

Conclusion 1.11 is meaningful because:

Fact 1.12. Assume that $\lambda = \lambda^{<\lambda} < \partial < \theta < \mu = \mu^\theta$, and $[\alpha < \mu \Rightarrow |\alpha|^\lambda < \mu]$, $\theta > \beth_2(\kappa)$ and $\partial = \kappa^+$, $\kappa = \kappa^\lambda$. Then the demands in Hypothesis 1.1 hold.

Remark 1.13. To justify the assumption, notice that:

(A) Omitting $\kappa = \kappa^\lambda$ does not help.

(B) $\theta \to_{\mathrm{sq}} (\partial)^{2^{\leq\lambda}}_{\partial}$ implies $\theta \to (\partial)^2_{2^\partial}$, hence necessarily $\theta > 2^{2^\partial}$.

With stronger lower bound on θ, see [She89].

The main point is proving $\theta \to_{\mathrm{sq}} (\partial)^{\leq\lambda,2}_{\partial}$. For this, see [She89], $\theta = \beth_m(\partial)$ for some small m suffice, on this the bounds in 1.11 depends; we intend to return to this in [S$^+$]. Anyhow just $\theta < \partial^{+\omega}$ and GCH in $[\partial, \partial^{+\omega}]$ would suffice for me.

Proof. The point is to prove $\theta \to_{\mathrm{sq}} (\partial)^{\lambda,2}_{\partial}$. Let $\mathscr{B}$ be as in 0.7(a), $\partial_1 = 2^\kappa$, $\partial_2 = 2^{\partial_1}$, and $\theta > \partial_2$.

Let $\chi > 2^\mu$, and $\mathfrak{C}_*$ be an expansion of $(\mathscr{H}(\chi), \in, <^*_\chi, \mathscr{B})$ with vocabulary of cardinality λ such that for any finite set $u \subseteq \mathscr{H}(\chi)$, the Skolem hull of u, $N_u := \mathrm{Sk}(u, \mathfrak{C}_*)$ is of cardinality λ and $|N_u|^{<\lambda} \subseteq N_u$.

Let $\mathfrak{C}_2 \prec_{\mathbb{L}_{\partial(1)^+,\partial(1)^+}} \mathfrak{C}_*$ be of cardinality ∂_2 such that $\partial_2 + 1 \subseteq \mathfrak{C}_2$. Let $\beta_1 := \min(\theta \setminus \mathfrak{C}_2)$. Similarly, choose $\mathfrak{C}_1 \prec_{\mathbb{L}_{\partial,\partial}} \mathfrak{C}_*$ of cardinality ∂_1 such that $\partial_1 + 1 \subseteq \mathfrak{C}_1$ and $\{\mathfrak{C}_2, \beta_0\} \subseteq \mathfrak{C}_1$.

Let $\mathfrak{C}_0 = \mathfrak{C}_1 \cap \mathfrak{C}_2$ and choose $\beta_0 \in \beta_1 \cap \mathfrak{C}_2 \subseteq \theta \cap \mathfrak{C}_2$ realizing the $\mathbb{L}_{\partial,\partial}$-type which β_1 realizes over $\mathfrak{C}_0$.

Now,

[4] We may omit p but it does not matter.

Paper Sh:1258, version 2026-09-29_3. See https://shelah.logic.at/papers/1258/ for possible updates.

$(*)_1$ choose $\alpha_\varepsilon \in \mathfrak{C}_0 \cap \theta$ by induction on $\varepsilon < \partial$, such that:

- α_ε, β_1 realize the same first-order type in $\mathfrak{C}_*$ over the set $\{\beta_2\} \cup (A_\varepsilon \cap \mathfrak{C}_0)$, where:

$$A_\varepsilon = \mathrm{Sk}_{\mathfrak{C}}(\{\alpha_\zeta : \zeta < \varepsilon\} \cup \{\beta_1, \beta_0\}).$$

$(*)_2$ Let $N^\bullet_\emptyset = N_{\{\beta_0,\beta_1\}} \cap \mathfrak{C}_0$.

Note,

$(*)_3$ for $\varepsilon < \zeta < \partial$, the following pairs realize the same type over N^*_0 in $\mathfrak{C}_*$:

- $\bullet_1$ $(\alpha_\varepsilon, \alpha_\zeta)$,
- $\bullet_2$ $(\alpha_\varepsilon, \beta_0)$,
- $\bullet_3$ $(\alpha_\varepsilon, \beta_1)$,
- $\bullet_4$ (β_0, β_1).

[Why? For the equality of $\bullet_1$ and $\bullet_2$ note the choice of α_ε.
For the equality of $\bullet_2$ and $\bullet_3$, note the choice of β_0.
For the equality of $\bullet_3$ and $\bullet_4$ note the choice of α_ε.]

$(*)_4$

- $\bullet_1$ $N^*_{\{\varepsilon,\zeta\}} = N_{\{\alpha_\varepsilon,\alpha_\zeta\}}$, so for $\varepsilon < \zeta < \partial$,
- $\bullet_2$ $N^*_{\{\varepsilon,\zeta\}} \prec \mathfrak{C}_0$,
- $\bullet_3$ $N^\bullet_\emptyset \prec N^*_{\{\varepsilon,\zeta\}}$.

$(*)_5$ for $\varepsilon < \zeta < \partial$, let $f_{\{\varepsilon,\zeta\}}$ be the isomorphism from $N^*_{\{\varepsilon,\zeta\}}$ onto $N_{\{\beta_0,\beta_1\}}$.

[Why does it exist? by $(*)_3$.]

$(*)_6$ $f_{\{\varepsilon,\zeta\}}$ is the identity on $N^\bullet_\emptyset$ (and $N^\bullet_\emptyset \prec N_{\{\varepsilon,\zeta\}}$).

[Why? By $(*)_2$.]

$(*)_7$ if $\varepsilon(0) < \zeta(0) < \partial$, $\varepsilon(1) < \zeta(1) < \partial$ and $\{\varepsilon(0), \zeta(0)\} \cap \{\varepsilon(1), \zeta(1)\} = \emptyset$, then

$$N^*_{\{\varepsilon(0),\zeta(0)\}} \cap N^*_{\{\varepsilon(1),\zeta(1)\}} = N_{\{\beta_0,\beta_1\}} \cap \mathfrak{C}_0 = N^\bullet_\emptyset.$$

[Why? The second equality holds by $(*)_2$; without loss of generality $\zeta(0) < \zeta(1)$. Now,

- $\bullet_1$ $N^*_{\{\varepsilon(0),\zeta(0)\}} \cap N^*_{\{\varepsilon(1),\zeta(1)\}} = N^*_{\{\varepsilon(0),\zeta(0)\}} \cap N_{\{\alpha_{\varepsilon(1)},\beta_1\}}$ by the choice of $\zeta(1)$.
- $\bullet_2$ if $\zeta(0) < \varepsilon(1)$ then
  $N^*_{\{\varepsilon(0),\zeta(0)\}} \cap N_{\{\alpha_{\varepsilon(1)},\beta_1\}} = N^*_{\{\varepsilon(0),\zeta(0)\}\}} \cap N_{\{\beta_0,\beta_1\}} = N_{\{\varepsilon(0),\zeta(0)\}} \cap N^\bullet_\emptyset = N^\bullet_\emptyset$ because the first equality follows by the choice of $\alpha_{\varepsilon(1)}$ second equality by $(*)_4\bullet_2$ and $(*)_2$; the third equality by $(*)_3$.
- $\bullet_3$ if $\varepsilon(0) < \varepsilon(1) < \zeta(0)$, then:

$$\begin{aligned}
N^*_{\{\varepsilon(0),\zeta(0)\}} \cap N_{\{\alpha_{\varepsilon(1)},\beta_1\}} &= N_{\{\alpha_{\varepsilon(0)},\beta_0\}} \cap N_{\{\alpha_{\varepsilon(1)},\beta_1\}} \\
&= \Big(N^*_{\{\alpha_{\varepsilon(0)},\beta_0\}} \cap \mathfrak{C}_0\Big) \cap \Big(N_{\{\alpha_{\varepsilon(0)},\beta_1\}} \cap \mathfrak{C}_0\Big) \\
&= \Big(N_{\{\alpha_{\zeta(0)},\beta_1\}} \cap \mathfrak{C}_0\Big) \cap \Big(N_{\{\alpha_{\zeta(0)},\beta_1\}} \cap \mathfrak{C}_0\Big) \\
&= \Big(N_{\{\alpha_{\zeta(0)},\beta_1\}} \cap \mathfrak{C}_0\Big) \cap \big(N_{\{\beta_0,\beta_1\}} \cap \mathfrak{C}_0\big) \\
&= \Big(N_{\{\alpha_{\varepsilon(0)},\beta_1\}} \cap \mathfrak{C}_0\Big) \cap N^\bullet_\emptyset = N^\bullet_\emptyset.
\end{aligned}$$

[Why? The first equality holds by the choice of β_0. The second equality as $N_{\{\varepsilon(0),\zeta(0)\}} \subseteq \mathfrak{C}_0$ and the first equality. The third equality holds by the choice of β_0. The fourth equality holds by the choice of $\alpha_{\zeta(0)}$. The fifth equality holds by the choice of $N^\bullet_\emptyset$ i.e., $(*)_2$. Finally, the sixth equality holds as $N_{\{\alpha_{\varepsilon(1)},\beta_1\}} \supseteq N^\bullet_\emptyset$ by the choice of $\alpha_{\varepsilon(2)}$.]

Paper Sh:1258, version 2026-09-29_3. See https://shelah.logic.at/papers/1258/ for possible updates.

$\bullet_4$ If $\varepsilon(1) < \varepsilon(0)$, then:

$$\begin{aligned} N^*_{\{\varepsilon(0),\zeta(0)\}} \cap N_{\{\alpha_{\varepsilon(1)},\beta_1\}} &= N_{\{\alpha_{\varepsilon(0)},\beta_0\}} \cap N_{\{\alpha_{\varepsilon(1)},\beta_1\}} \\ &= \Big(N^*_{\{\alpha_{\varepsilon(0)},\beta_0\}} \cap \mathfrak{C}_0\Big) \cap \Big(N^*_{\{\alpha_{\varepsilon(1)},\beta_1\}} \cap \mathfrak{C}_0\Big) \\ &= \Big(N_{\{\alpha_{\varepsilon(0)},\beta_1\}} \cap \mathfrak{C}_0\Big) \cap \Big(N_{\{\alpha_{\varepsilon(1)},\beta_1\}} \cap \mathfrak{C}_0\Big) \\ &= \big(N_{\{\beta_0,\beta_1\}} \cap \mathfrak{C}\big) \cap \Big(N_{\{\alpha_{\varepsilon(1)},\beta_1\}} \cap \mathfrak{C}\Big) \\ &= N^*_\emptyset \cap \Big(N_{\{\alpha_{\varepsilon(1)},\beta_1\}} \cap \mathfrak{C}_0\Big) = N^\bullet_\emptyset. \end{aligned}$$

[Why? The first equality holds by the choice of β_0. The second one holds as $N_{\{\varepsilon(0),\zeta(0)\}} \subseteq \mathfrak{C}_0$ and the first equality. The third equality holds by the choice of β_0. The fourth equality holds by the choice of $\alpha_{\varepsilon(0)}$. The fifth equality holds by the choice of $N^\bullet_\emptyset$, i.e., by $(*)_2$. Finally, the sixth equality holds as $N_{\{\alpha_{\varepsilon(1)},\beta_1\}} \supseteq N^\bullet_\emptyset$ and by the choice of $\alpha_{\varepsilon(0)}$.

Recalling $\bullet_1$ and the division to cases in $\bullet_2, \bullet_3$ and $\bullet_4$, we are done proving $(*)_6$.]

$(*)_7$ if $\varepsilon < \zeta(1) < \zeta(2) < \partial$, then $N_{\{\varepsilon,\zeta(1)\}} \cap N_{\varepsilon,\zeta(2)} = N^\uparrow_{\{\varepsilon\}} := N_{\{\alpha_\varepsilon,\beta_1\}} \restriction \mathfrak{C}_0$.

[Why? By the choice of $\alpha_{\zeta(2)}$ and $\alpha_{\zeta(1)}$.]

$(*)_8$ if $\zeta_1 < \zeta_2 < \varepsilon < \partial$, then $N^*_{\{\zeta_1,\varepsilon\}} \cap N^*_{\{\zeta_2,\varepsilon\}} = N^\downarrow_{\{\varepsilon\}}$, where $N^\downarrow_{\{\varepsilon\}} := f^{-1}_{\varepsilon,\varepsilon+1}(N_{\{\beta_0,\beta_1\}})$.

[Why? For $\zeta < \partial$, $N_{\{\alpha_\zeta,\beta_0\}} \cap \mathfrak{C}_0 = N_{\{\alpha_{\zeta_3},\beta_1\}} \cap \mathfrak{C}_0$ by the choice of β_0, and $\alpha_\varepsilon, \beta_0$ realize the same type of $\mathfrak{C}_*$ over $\{\beta_1\} \cup (A_\varepsilon \cap \mathfrak{C}_0)$.]

$(*)_9$
- Let $N^*_{\{\varepsilon\}}$ be the $\mathrm{Sk}(N^\uparrow_{\{\varepsilon\}} \cup N^\downarrow_{\{\varepsilon\}}, \mathfrak{C}_*)$, and
- let $M^*_\varepsilon = \Big(\mathrm{Sk}(\bigcup_{\ell<5} N^*_{\{5\varepsilon+\ell\}} \cup \{N_{\{5\varepsilon+m,\delta\varepsilon+n\}} : m < n < 5\}^*)\Big)_{\ell<5}$,
- let M^+_ε be M^*_ε expanded by:
- $c^{M^+_\varepsilon}_\ell = \alpha_{5\varepsilon+\ell}$ for $\ell < 5$,
- $p^{M^+_\varepsilon}_\ell = |N^*_{\{5\varepsilon+\ell\}}|$ for $\ell < 5$,
- $P^{M^+_\varepsilon}_{m,n} = |N^*_{\{5\varepsilon+m,5\varepsilon+n:\, m<n<5\}}|$.

$(*)_{10}$ There is some $\mathscr{U}_1 \in [\partial]^\partial$ such that:
- $\langle M^*_\varepsilon : \varepsilon \in \mathscr{U}\rangle$ is a Δ-system with heart $N^*_\emptyset$,
- the M_ε are pairwise isomorphic.

[Why? Because $\partial = \partial_0$ and $\partial_0 = (\partial_0)^+$ by the Δ-system lemma.]

$(*)_{11}$ $\langle N^*_u : u \in \mathscr{U}_2\rangle$ is a required when $\mathscr{U}_2 = \{5\varepsilon+2 : \varepsilon \in \mathscr{U}_1\}$ and $N^*_{\{5\varepsilon+2\}} = M^*_\varepsilon$.

Pedantically, $\mathscr{U}_3 = \{\alpha_\zeta : \zeta \in \mathscr{U}_2\}$ and $N^*_{\{\alpha_\zeta :\, \zeta\in u\}} = N^*_u$ for $u \in [\mathscr{U}_3]^{\le 2}$. $\square_{1.12}$

References

[EH71] Paul Erdős and Andras Hajnal, *Unsolved problems in set theory*, Axiomatic Set Theory (Providence, R.I.), Proc. of Symp. in Pure Math., vol. XIII Part I, AMS, 1971, pp. 17–48.

[Kom25] Péter Komjáth, *The Erdős-Hajnal Problem List*, Bull. Symb. Log. **31** (2025), no. 3, 418–461.

[RS00] Mariusz Rabus and Saharon Shelah, *Covering a function on the plane by two continuous functions on an uncountable square—the consistency*, Ann. Pure Appl. Logic **103** (2000), no. 1-3, 229–240, arXiv: math/9706223. MR 1756147

[S$^+$] S. Shelah et al., *Tba*, In preparation. Preliminary number: Sh:F2407.

[She88] Saharon Shelah, *Was Sierpiński right? I*, Israel J. Math. **62** (1988), no. 3, 355–380. MR 955139

[She89] ———, *Consistency of positive partition theorems for graphs and models*, Set theory and its applications (Toronto, ON, 1987), Lecture Notes in Math., vol. 1401, Springer, Berlin, 1989, pp. 167–193. MR 1031773

Paper Sh:1258, version 2026-09-29_3. See https://shelah.logic.at/papers/1258/ for possible updates.

[She92] ———, *Strong partition relations below the power set: consistency; was Sierpiński right? II*, Sets, graphs and numbers (Budapest, 1991), Colloq. Math. Soc. János Bolyai, vol. 60, North-Holland, Amsterdam, 1992, arXiv: math/9201244, pp. 637–668. MR 1218224

[She95] ———, *Possibly every real function is continuous on a non-meagre set*, Publ. Inst. Math. (Beograd) (N.S.) **57(71)** (1995), 47–60, arXiv: math/9511220. MR 1387353

[She96] ———, *Was Sierpiński right? III. Can continuum-c.c. times c.c.c. be continuum-c.c.?*, Ann. Pure Appl. Logic **78** (1996), no. 1-3, 259–269, arXiv: math/9509226. MR 1395402

[She00] ———, *Was Sierpiński right? IV*, J. Symbolic Logic **65** (2000), no. 3, 1031–1054, arXiv: math/9712282. MR 1791363

Einstein Institute of Mathematics, The Hebrew University of Jerusalem, 9190401, Jerusalem, Israel; and, Department of Mathematics, Rutgers University, Piscataway, NJ 08854-8019, USA

URL: `https://shelah.logic.at/`