

## When automorphisms of $\mathcal{P}(\kappa)/[\kappa]^{<\aleph_0}$ are trivial off a small set

by

**Saharon Shelah** (Piscataway, NJ, and Jerusalem) and  
**Juris Steprāns** (Toronto)

**Abstract.** It is shown that if  $\kappa > 2^{\aleph_0}$  and  $\kappa$  is less than the first inaccessible cardinal then every automorphism of  $\mathcal{P}(\kappa)/[\kappa]^{<\aleph_0}$  is trivial outside of a set of cardinality  $2^{\aleph_0}$ .

**1. Introduction.** The study of automorphisms of  $\mathcal{P}(\omega)/[\omega]^{<\aleph_0}$  was initiated by W. Rudin [4, 5] who showed that the Continuum Hypothesis can be used to construct non-trivial autohomeomorphisms of  $\beta\mathbb{N}/\mathbb{N}$ , in other words, homeomorphisms from  $\beta\mathbb{N}/\mathbb{N}$  to  $\beta\mathbb{N}/\mathbb{N}$  that are not induced by any function from  $\mathbb{N}$  to  $\mathbb{N}$ . By Stone duality, this means that there are automorphisms of  $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$  that are not induced by a function from  $\mathbb{N}$  to  $\mathbb{N}$ . A major advance was provided by S. Shelah [6] who showed that it is consistent with set theory that every automorphism of  $\mathcal{P}(\omega)/[\omega]^{<\aleph_0}$  is induced by a function from  $\omega$  to  $\omega$ . Later, B. Velickovic [7] showed that the conjunction of OCA and MA implies that every automorphism of  $\mathcal{P}(\omega_1)/[\omega_1]^{<\aleph_0}$  is induced by a function from  $\omega_1$  to  $\omega_1$ . Moreover, he showed that PFA implies that if  $\kappa$  is uncountable then every automorphism of  $\mathcal{P}(\kappa)/[\kappa]^{<\aleph_0}$  is induced by a function from  $\kappa$  to  $\kappa$ .

However, finding extensions of Rudin's result on the existence on non-trivial automorphisms of  $\mathcal{P}(\kappa)/[\kappa]^{<\aleph_0}$  has proven to be much harder. This article will provide some reasons for this. In particular, a positive answer to the following question from [7] will be given for cardinals below the first inaccessible: Can it be shown from MA and OCA alone that for every uncountable  $\kappa$ , every automorphism of  $\mathcal{P}(\kappa)/[\kappa]^{<\aleph_0}$  is induced by a function from  $\kappa$  to  $\kappa$ ?

---

2010 *Mathematics Subject Classification*: 03E17, 03E35.

*Key words and phrases*: automorphism, dominating number.

Received 20 August 2015; revised 22 February 2016.

Published online 30 May 2016.

The main result to be established in this article is that if  $\kappa > 2^{\aleph_0}$  and  $\kappa$  is less than the first inaccessible cardinal then for every automorphism of  $\mathcal{P}(\kappa)/[\kappa]^{<\aleph_0}$  there is a set  $X \subseteq \kappa$  such that  $|\kappa \setminus X| \leq 2^{\aleph_0}$  and the restriction of the automorphism to  $\mathcal{P}(X)/[X]^{<\aleph_0}$  is induced by a function from  $\kappa$  to  $\kappa$ .

The question of cardinals not greater than the continuum has been dealt with by P. Larson and P. McKenney [3] who have shown the following.

**THEOREM 1.1** (Larson & McKenney). *If  $\kappa \leq 2^{\aleph_0}$  and  $\Psi$  is an automorphism of  $\mathcal{P}(\kappa)/[\kappa]^{<\aleph_0}$  such that  $\Psi \upharpoonright \mathcal{P}(X)/[X]^{<\aleph_0}$  is trivial for each  $X \in [\kappa]^{\aleph_1}$  then  $\Psi$  is trivial.*

Without too much extra work, these two results then provide a partial answer to Velickovic's question.

**COROLLARY 1.2.** *If MA and OCA both hold and if  $\kappa$  is less than the first inaccessible then every automorphism of  $\mathcal{P}(\kappa)/[\kappa]^{<\aleph_0}$  is trivial.*

*Proof.* Let  $\kappa$  be less than the first inaccessible cardinal and let  $\Phi$  be an automorphism of  $\mathcal{P}(\kappa)/[\kappa]^{<\aleph_0}$ . By Theorem 3.1 there is  $X \subseteq \kappa$  such that  $|X| = 2^{\aleph_0}$  and there is a function that induces  $\Phi \upharpoonright \mathcal{P}(\kappa \setminus X)/[\kappa \setminus X]^{<\aleph_0}$ . It therefore suffices to show that  $\Phi \upharpoonright \mathcal{P}(X)/[X]^{<\aleph_0}$  is trivial. To do so, it suffices by Theorem 1.1 to show that if  $Y \subseteq X$  and  $|Y| = \aleph_1$  then  $\Phi \upharpoonright \mathcal{P}(Y)/[Y]^{<\aleph_0}$  is trivial. But this follows from the theorem of Velickovic [7] that the conjunction of OCA and MA implies that every automorphism of  $\mathcal{P}(\omega_1)/[\omega_1]^{<\aleph_0}$  is trivial. Hence  $\Phi$  is trivial. ■

## 2. Terminology and notation

**NOTATION 2.1.** For  $X \subseteq \kappa$  let  $[X]$  be the equivalence class of  $X$  modulo the ideal  $[X]^{<\aleph_0}$ . For any infinite set  $X$  the notation  $\mathcal{P}(X)/\mathcal{F}in$  will be used in place of  $\mathcal{P}(X)/[X]^{<\aleph_0}$ , it being understood that  $\mathcal{F}in$  refers to  $[X]^{<\aleph_0}$  as appropriate. For any function  $f$  and any set  $x$  let  $f \langle x \rangle$  denote the image of  $x$  under  $f$ .

**DEFINITION 2.2.** A homomorphism  $\Phi : \mathcal{P}(\kappa)/\mathcal{F}in \rightarrow \mathcal{P}(\kappa)/\mathcal{F}in$  will be said to be  $\lambda$ -trivial if there is a set  $S \in [\kappa]^\lambda$  and a one-to-one function  $F : \kappa \setminus S \rightarrow \kappa$  such that  $F \langle X \rangle \in \Phi([X])$  for every  $X \subseteq \kappa \setminus S$ . If  $\Phi : \mathcal{P}(\kappa)/\mathcal{F}in \rightarrow \mathcal{P}(\kappa)/\mathcal{F}in$  is a homomorphism, let  $\hat{\Phi}$  be some lifting of  $\Phi$ . In other words,  $\hat{\Phi}(X)$  is a function from  $\mathcal{P}(\kappa)$  to  $\mathcal{P}(\kappa)$  such that  $\hat{\Phi}(X) \in \Phi([X])$  for each  $X \subseteq \kappa$  and  $\hat{\Phi}$  is constant on each  $[X]$ .

**THEOREM 2.3** (Balcar and Frankiewicz [1]). *For any cardinal  $\kappa$  and any automorphism  $\Psi$  of  $\mathcal{P}(\kappa)/\mathcal{F}in$ , if  $\Psi([X]) = [Y]$  and  $|X|, |Y| \geq \aleph_1$  then  $|X| = |Y|$ ; moreover, if  $|X| = \aleph_0$  then  $|Y| \leq \aleph_1$ .*

In particular, there is no isomorphism from  $\mathcal{P}(\kappa)/\mathcal{F}in$  to  $\mathcal{P}(\lambda)/\mathcal{F}in$  unless  $\kappa = \lambda$  or  $\{\kappa, \lambda\} = \{\aleph_0, \aleph_1\}$ . In this context, it is interesting to note that it is

shown in [2] that if there is an isomorphism from  $\mathcal{P}(\omega_1)/\mathcal{F}in$  to  $\mathcal{P}(\omega)/\mathcal{F}in$  then there is a non-trivial automorphism of  $\mathcal{P}(\omega)/\mathcal{F}in$  itself.

**DEFINITION 2.4.** If  $\mathfrak{B}$  is a subalgebra of  $\mathcal{P}(X)$  and  $\lambda$  is an infinite cardinal, define  $\mathcal{I}_\lambda(\mathfrak{B}) = [X]^{<\lambda} \cap \mathfrak{B}$  and  $I_\lambda(\mathfrak{B}) = \bigcup \mathcal{I}_\lambda(\mathfrak{B})$ . The subalgebra of  $\mathcal{P}(X \setminus I_\lambda(\mathfrak{B}))$  consisting of  $\{B \setminus I_\lambda(\mathfrak{B}) \mid B \in \mathfrak{B}\}$  will be denoted by  $\mathfrak{B}_\lambda$ .

Note that  $\mathfrak{B}_\lambda$  may be trivial if  $\lambda \geq |X|$  or  $[X]^{<\lambda} \subseteq \mathfrak{B}$ .

**DEFINITION 2.5.** If  $\Phi$  is an automorphism of  $\mathcal{P}(X)/\mathcal{F}in$  and  $\mathfrak{B}$  is a subalgebra of  $\mathcal{P}(X)$  then define  $\mathfrak{B}^\Phi$  to be the algebra generated by  $\hat{\Phi}(\mathfrak{B})$ , noting that  $\hat{\Phi}(\mathfrak{B})$  may not itself be a subalgebra. The notation  $\mathfrak{B}_\lambda^\Phi$  will be used instead of the more cumbersome  $(\mathfrak{B}^\Phi)_\lambda$ . If  $\lambda$  is an uncountable cardinal, define  $\Phi_\lambda : \mathfrak{B}_\lambda \rightarrow \mathfrak{B}_\lambda^\Phi$  by  $\Phi_\lambda(B \setminus I_\lambda(\mathfrak{B})) = \hat{\Phi}(B) \setminus I_\lambda(\mathfrak{B}^\Phi)$  for  $B \in \mathfrak{B}$ .

**LEMMA 2.6.** *If  $\Phi : \mathcal{P}(X)/\mathcal{F}in \rightarrow \mathcal{P}(X)/\mathcal{F}in$  is a one-to-one homomorphism,  $\mathfrak{B}$  is a subalgebra of  $\mathcal{P}(X)$ ,  $\lambda > \aleph_1$  and  $|\mathfrak{B}| < \mathbf{cof}(\lambda)$  then  $\Phi_\lambda$  is a well defined isomorphism from  $\mathfrak{B}_\lambda$  to  $\mathfrak{B}_\lambda^\Phi$ .*

*Proof.* To see that the mapping is well defined, begin by noting that  $|I_\lambda(\mathfrak{B})| < \lambda$ . Now suppose that  $B \setminus I_\lambda(\mathfrak{B}) = B' \setminus I_\lambda(\mathfrak{B})$ . Then  $|B \triangle B'| < \lambda$ , and hence, by Theorem 2.3,  $|\hat{\Phi}(B \triangle B')| < \lambda$ . However  $\hat{\Phi}(B) \triangle \hat{\Phi}(B') \equiv^* \hat{\Phi}(B \triangle B')$  and so  $\hat{\Phi}(B) \triangle \hat{\Phi}(B') \subseteq I_\lambda(\mathfrak{B}^\Phi)$ . In other words,  $\Phi_\lambda(B) = \Phi_\lambda(B')$ .

To see that  $\Phi_\lambda$  is a homomorphism suppose that  $\Phi_\lambda(B) \not\subseteq \Phi_\lambda(B')$ . Then  $|\hat{\Phi}(B) \setminus \hat{\Phi}(B')| \geq \lambda$ . Since  $\hat{\Phi}(B) \setminus \hat{\Phi}(B') \equiv^* \hat{\Phi}(B \setminus B')$ , it follows from Theorem 2.3 that  $|B \setminus B'| \geq \lambda$  and so  $B \setminus I_\lambda(\mathfrak{B}) \not\subseteq B' \setminus I_\lambda(\mathfrak{B})$ .

Since  $\Phi$  is a homomorphism, in order to see that  $\Phi_\lambda$  is one-to-one it suffices to show that if  $\Phi_\lambda(B) = \emptyset$  then  $B \subseteq I_\lambda(\mathfrak{B})$ . This is immediate from Theorem 2.3.

To see that  $\Phi_\lambda$  is onto let  $B \in \mathfrak{B}_\lambda^\Phi$ . By definition there is some  $B' \in \mathfrak{B}_\lambda$  such that  $\hat{\Phi}(B') \equiv^* B$ , and hence  $\hat{\Phi}(B') \triangle B \subseteq I_\lambda(\mathfrak{B}^\Phi)$ . In other words,  $\Phi_\lambda(B') \setminus I_\lambda(\mathfrak{B}^\Phi) = B \setminus I_\lambda(\mathfrak{B}^\Phi)$  as required. ■

**3. The lemmas needed for the main result.** The main result of this article is the following.

**THEOREM 3.1.** *If  $\kappa > 2^{\aleph_0}$  and  $\kappa$  is less than the first inaccessible cardinal then every automorphism of  $\mathcal{P}(\kappa)/\mathcal{F}in$  is  $2^{\aleph_0}$ -trivial.*

The proof proceeds by induction on  $\kappa$ , using several lemmas. The first is immediate.

**LEMMA 3.2.** *If every automorphism of  $\mathcal{P}(\kappa)/\mathcal{F}in$  is  $\lambda$ -trivial and every automorphism of  $\mathcal{P}(\lambda)/\mathcal{F}in$  is  $\mu$ -trivial then every automorphism of  $\mathcal{P}(\kappa)/\mathcal{F}in$  is  $\mu$ -trivial.*

The next lemma deals with the successors of cardinals with uncountable cofinality.

LEMMA 3.3. *If  $\kappa^{\aleph_0} = \kappa$  and  $\nu \leq 2^\kappa$  then every automorphism of  $\mathcal{P}(\nu)/\mathcal{F}in$  is  $\kappa$ -trivial.*

The singular cardinals of uncountable cofinality are easy to handle, as the next lemma demonstrates.

LEMMA 3.4. *If  $\kappa$  is singular of uncountable cofinality and if for every cardinal  $\lambda < \kappa$  every automorphism of  $\mathcal{P}(\lambda)/\mathcal{F}in$  is  $2^{\aleph_0}$ -trivial then so is every automorphism of  $\mathcal{P}(\kappa)/\mathcal{F}in$ .*

*Proof.* Let  $\Phi$  be an automorphism of  $\mathcal{P}(\kappa)/\mathcal{F}in$ . Let  $\{\kappa_\xi\}_{\xi \in \mathbf{cof}(\kappa)}$  be an increasing, cofinal sequence of cardinals in  $\kappa$ . Using the hypothesis, for each  $\xi \in \mathbf{cof}(\kappa)$  choose  $X_\xi$  and  $F_\xi : \kappa_\xi \setminus X_\xi \rightarrow \kappa$  such that

- $|X_\xi| \leq 2^{\aleph_0}$ ,
- $F_\xi \langle A \rangle \in \Phi([A])$  for each  $A \subseteq \kappa_\xi \setminus X_\xi$ .

Then let

$$X = \left( \bigcup_{\xi \in \mathbf{cof}(\kappa)} X_\xi \right) \cup \left( \bigcup_{\{\xi, \eta\} \in [\mathbf{cof}(\kappa)]^2} \{\zeta \in \kappa \mid F_\xi(\zeta) \neq F_\eta(\zeta)\} \right)$$

and note that  $|X| \leq \mathbf{cof}(\kappa) \cdot 2^{\aleph_0}$ . Let  $F = \bigcup_{\xi \in \mathbf{cof}(\kappa)} F_\xi \upharpoonright (\kappa \setminus X)$  and note that  $F$  is a function.

To see that  $F \langle A \rangle \in \Phi([A])$  for each  $A \subseteq \kappa \setminus X$  observe that otherwise there is some infinite  $A \subseteq \kappa \setminus X$  such that  $F \langle A \rangle \cap \Phi([A])$  is finite. Moreover,  $A$  can then be chosen to be countable, and hence  $A \subseteq \kappa_\xi \setminus X_\xi$  for some  $\xi \in \mathbf{cof}(\kappa)$ , contradicting  $F \langle A \rangle = F_\xi \langle A \rangle$ .

If  $\mathbf{cof}(\kappa) \leq 2^{\aleph_0}$  there is nothing else to do. Otherwise, it has been shown that  $\Phi$  is  $\mathbf{cof}(\kappa)$ -trivial. Since  $\mathbf{cof}(\kappa) < \kappa$ , the result now follows from Lemma 3.2. ■

The next two lemmas deal with the harder case of singular cardinals of countable cofinality.

LEMMA 3.5. *If  $\kappa$  has cofinality  $\omega$  and for every  $\lambda < \kappa$  every automorphism of  $\mathcal{P}(\lambda)/\mathcal{F}in$  is  $2^{\aleph_0}$ -trivial then every automorphism of  $\mathcal{P}(\kappa)/\mathcal{F}in$  is  $2^{\aleph_0}$ -trivial.*

LEMMA 3.6. *Suppose that*

- $\kappa$  has countable cofinality,
- $2^\mu < \kappa$  for each  $\mu < \kappa$ ,
- if  $\mu \leq \kappa$  then every automorphism of  $\mathcal{P}(\mu)/\mathcal{F}in$  is  $2^{\aleph_0}$ -trivial.

*Then every automorphism of  $\mathcal{P}(\rho)/\mathcal{F}in$  is  $\kappa$ -trivial for every cardinal  $\rho$  such that  $\kappa^+ \leq \rho \leq \kappa^{\aleph_0}$ .*

*Proof of Theorem 3.1.* If the result fails, let  $\lambda$  be the least cardinal such that there is an automorphism  $\Phi$  of  $\mathcal{P}(\lambda)/\mathcal{F}in$  that is not  $2^{\aleph_0}$ -trivial. By

Lemmas 3.4 and 3.5,  $\lambda$  must be a regular cardinal. Since  $\lambda$  is less than the first inaccessible, there must be some least  $\kappa < \lambda$  such that  $2^\kappa \geq \lambda$ . From Lemma 3.3 applied to  $\kappa$  and  $\lambda$ , it follows that  $\kappa^{\aleph_0} > \kappa$ . If  $\kappa^{\aleph_0} < \lambda$  then Lemma 3.3 applied to  $\kappa^{\aleph_0}$  and  $\lambda$  implies that  $\Phi$  is  $\kappa^{\aleph_0}$ -trivial. A contradiction then follows from the induction hypothesis and Lemma 3.2. It can therefore be assumed that  $\kappa^{\aleph_0} \geq \lambda$ .

Lemma 3.6 implies that either  $\kappa$  has uncountable cofinality or there is  $\nu < \kappa$  such that  $2^\nu \geq \kappa$ . (The third alternative cannot fail because of the minimality of  $\lambda$ .) If the second alternative holds then the minimality of  $\kappa$  implies that  $2^\nu < \lambda$ . Since  $2^\nu = (2^\nu)^{\aleph_0}$  and  $2^{2^\nu} \geq 2^\kappa \geq \lambda$  it is once again possible to apply Lemma 3.3 to  $2^\nu$  and  $\lambda$  to get the contradiction that  $\Phi$  is  $2^\nu$ -trivial and  $2^\nu < \lambda$ . So it suffices to consider the case that  $\kappa$  has uncountable cofinality and  $2^\nu < \kappa$  for each  $\nu < \kappa$ . This implies that  $\nu^{\aleph_0} < \kappa$  for each  $\nu < \kappa$ , and hence  $\kappa^{\aleph_0} = \kappa$ , which has already been ruled out. ■

#### 4. Automorphisms of $\mathcal{P}(2^\kappa)/\mathcal{F}in$ when $\kappa^{\aleph_0} = \kappa$

*Proof of Lemma 3.3.* Assume that  $\kappa^{\aleph_0} = \kappa$  and  $\nu \leq 2^\kappa$ , and note that there is nothing to prove if  $\nu \leq \kappa$ . Let  $\Psi$  be an automorphism of  $\mathcal{P}(\nu)/\mathcal{F}in$  and  $\hat{\Psi}$  a lifting of  $\Psi$ . Begin by noting that if  $(2^X)_\sigma$  denotes the space  $2^X$  with the topology generated by  $G_\delta$  sets in  $2^X$  with the usual product topology, then the density of  $(2^\nu)_\sigma$  is less than or equal to the density of  $(2^{2^\kappa})_\sigma$ , and this in turn is no greater than  $\kappa^{\aleph_0} = \kappa$ . Hence there is a family  $\mathcal{D} \subseteq \mathcal{P}(\nu)$  of cardinality  $\kappa$  such that for any two disjoint countable sets  $A \subseteq \nu$  and  $B \subseteq \nu$  there is  $D \in \mathcal{D}$  such that  $A \subseteq D$  and  $B \cap D = \emptyset$ .

Let  $\mathfrak{M}$  be an elementary submodel of  $(H((2^\kappa)^+), \mathcal{D}, \hat{\Psi}, \in)$  such that  $[\mathfrak{M}]^{\aleph_0} \subseteq \mathfrak{M}$  and  $|\mathfrak{M}| = \kappa$ . Let  $\mathfrak{A} = \mathcal{P}(\nu) \cap \mathfrak{M}$  and note that  $\mathfrak{A}$  is a Boolean  $\sigma$ -subalgebra of  $\mathcal{P}(\nu)$  containing  $\mathcal{D}$ . Moreover,  $\mathfrak{A}$  contains the Boolean  $\sigma$ -subalgebra generated by  $\hat{\Psi}[\mathfrak{A}]$ . By Lemma 2.6,  $\Psi_{\kappa^+} : \mathfrak{A}_{\kappa^+} \rightarrow \mathfrak{A}_{\kappa^+}^{\Psi}$  is an isomorphism, but it will be shown by using the following lemma that it is in fact a  $\sigma$ -isomorphism.

LEMMA 4.1. *If  $\mathcal{C} \subseteq \mathfrak{A}_{\kappa^+}$  is countable then*

$$\Psi_{\kappa^+} \left( \bigcup \mathcal{C} \right) = \bigcup \Psi_{\kappa^+} \langle \mathcal{C} \rangle.$$

*Proof.* Since  $\Psi_{\kappa^+}$  is a homomorphism, we have  $\Psi_{\kappa^+}(\bigcup \mathcal{C}) \supseteq \bigcup \Psi_{\kappa^+} \langle \mathcal{C} \rangle$ , so suppose that

$$(4.1) \quad \Psi_{\kappa^+} \left( \bigcup \mathcal{C} \right) \setminus \bigcup \Psi_{\kappa^+} \langle \mathcal{C} \rangle \neq \emptyset.$$

Then

$$(4.2) \quad Z = \hat{\Psi} \left( \bigcup \mathcal{C} \right) \setminus \bigcup \hat{\Psi} \langle \mathcal{C} \rangle \neq \emptyset,$$

and moreover  $Z \in \mathfrak{M}$  because  $\hat{\Psi}$  and  $\mathcal{C}$  both belong to  $\mathfrak{M}$ . From (4.1) it follows that  $|Z| \geq \kappa^+$ . By elementarity there is some  $Z^* \subseteq \nu$  such that  $\hat{\Psi}(Z^*) \equiv^* Z$ . By Theorem 2.3,  $|Z^*| \geq \kappa^+$  and so there is  $C \in \mathcal{C}$  such that  $|C \cap Z^*| \geq \kappa^+$ . Hence  $\hat{\Psi}(C \cap Z^*) \subseteq^* Z \cap \hat{\Psi}(C)$ . By Theorem 2.3 again,  $|\hat{\Psi}(C \cap Z^*)| \geq \kappa^+$ , contradicting  $Z \cap \hat{\Psi}(C) = \emptyset$ . ■

**COROLLARY 4.2.**  $\Psi_{\kappa^+}$  is a  $\sigma$ -isomorphism.

Now, for  $\xi \in \nu \setminus I_{\kappa^+}(\mathfrak{A})$  let  $U(\xi)$  be the ultrafilter on  $\mathfrak{A}_{\kappa^+}$  defined by  $U(\xi) = \{A \in \mathfrak{A}_{\kappa^+} \mid \xi \in A\}$ . Let  $U(\xi, \Psi)$  be the image of  $U(\xi)$  under  $\Psi_{\kappa^+}$ , and note that  $U(\xi, \Psi)$  is also an ultrafilter on  $\mathfrak{A}_{\kappa^+}^{\Psi}$ .

**LEMMA 4.3.** For all but finitely many  $\xi \in \nu \setminus I_{\kappa^+}(\mathfrak{A})$  the cardinality of  $\bigcap U(\xi, \Psi)$  is at most 1.

*Proof.* Let  $B = \{\xi \in \nu \setminus I_{\kappa^+}(\mathfrak{A}) \mid |\bigcap U(\xi, \Psi)| > 1\}$  and suppose that  $B$  is infinite. Choose a countable  $\bar{B} \subseteq B$  and then choose  $\{\beta_0, \beta_1\} \in [\bigcap U(\beta, \Psi)]^2$  for each  $\beta \in \bar{B}$ . Let  $B_i = \{\beta_i \mid \beta \in \bar{B}\}$ . Observe that since  $\mathcal{D} \subseteq \mathfrak{A}$ , it follows that if  $\xi \neq \eta$  then  $(\bigcap U(\xi, \Psi)) \cap \bigcap U(\eta, \Psi) = \emptyset$ , and hence  $\{\beta_0, \beta_1\} \cap \{\bar{\beta}_0, \bar{\beta}_1\} = \emptyset$  if  $\beta$  and  $\bar{\beta}$  are distinct elements of  $\bar{B}$ . Therefore  $B_0 \cap B_1 = \emptyset$ .

Then let  $C_0$  be such that  $\hat{\Psi}(C_0) \equiv^* B_0$ . Now, even though  $B_0$  is countable, Theorem 2.3 does not rule out the possibility that  $|C_0| = \aleph_1$ . In this case choose a countable  $C_0^* \subseteq C_0$  and let  $W = \{\beta \in \bar{B} \mid \beta_0 \in \hat{\Psi}(C_0^*)\}$ . Then let  $C_1$  be such that  $\hat{\Psi}(C_1) \equiv^* \{\beta_1 \mid \beta \in W\}$ . Once again,  $C_1$  might be uncountable. If this is the case, choose a countable  $B_1^* \subseteq C_1$  and let  $\bar{W} = \{\beta \in W \mid \beta_1 \in \hat{\Psi}(B_1^*)\}$ . Let  $\bar{B}_i = \{\beta_i \mid \beta \in \bar{W}\}$  and then let  $B_0^*$  be such that  $\hat{\Psi}(B_0^*) \equiv^* \bar{B}_0$ . It follows that  $\hat{\Psi}(B_i^*) \equiv^* \bar{B}_i$  for each  $i$ , and each  $B_i^*$  is countable.

Since  $\bar{B}_0 \cap \bar{B}_1 = \emptyset$ , a contradiction will be obtained if it can be shown that  $B_i^* \supseteq^* \bar{W}$  for each  $i$ . So suppose that  $Z \subseteq \bar{W} \setminus B_i^*$  is countably infinite for some  $i \in 2$ . Using the fact that  $\mathfrak{A} \supseteq \mathcal{D}$  and that  $Z$  and  $B_i^*$  are both countable we deduce that there is some  $D \in \mathcal{D}$  such that  $Z \subseteq D$  and  $D \cap B_i^* = \emptyset$ . It then follows that

$$|\bar{B}_i \cap \Psi_{\kappa^+}(D)| < \aleph_0,$$

contradicting, by Lemma 4.1, the inequality

$$(4.3) \quad |\bar{B}_i \cap \Psi_{\kappa^+}(D)| \geq \left| \bigcup_{\beta \in Z} \bar{B}_i \cap \bigcap U(\beta, \Psi) \right| \geq |Z| = \aleph_0$$

for each  $i \in 2$ . ■

**LEMMA 4.4.** For all but  $\kappa$  many  $\xi \in \nu \setminus I_{\kappa^+}(\mathfrak{A})$  the cardinality of  $\bigcap U(\xi, \Psi)$  is at least 1.

*Proof.* Let  $B = \{\xi \in \nu \setminus I_{\kappa^+}(\mathfrak{A}) \mid \bigcap U(\xi, \Psi) = \emptyset\}$  and suppose that  $|B| > \kappa$ . By Theorem 2.3 it follows that  $|\hat{\Psi}(B)| > \kappa$  also, and hence it is possible to

find a countable  $\bar{B} \subseteq B \setminus I_{\kappa^+}(\mathfrak{A})$  with  $\hat{\Psi}(\bar{B}) \cap I_{\kappa^+}(\mathfrak{A}^\Psi) \equiv^* \emptyset$ . As in the proof of Lemma 4.3, there is no loss of generality in assuming that  $\hat{\Psi}(\bar{B})$  is countable. Then observe that  $\bar{B} \subseteq \bigcup_{\beta \in \bar{B}} V(\beta)$  for all  $V \in \prod_{\beta \in \bar{B}} U(\beta)$ . Hence, using Corollary 4.2, we see that for any  $V \in \prod_{\beta \in \bar{B}} U(\beta)$ ,

$$\hat{\Psi}(\bar{B}) \subseteq^* \hat{\Psi}\left(\bigcup_{\beta \in \bar{B}} V(\beta)\right) \setminus I_{\kappa^+}(\mathfrak{A}^\Psi) = \Psi_{\kappa^+}\left(\bigcup_{\beta \in \bar{B}} V(\beta)\right) = \bigcup_{\beta \in \bar{B}} \Psi_{\kappa^+}(V(\beta))$$

and, since  $\hat{\Psi}(\bar{B})$  is countable, there are some  $\beta_1 \in \bar{B}$  and  $\beta_2 \in \hat{\Psi}(\bar{B})$  such that  $\beta_2 \in \Psi_{\kappa^+}(V(\beta_1))$  for cofinally many  $V \in \prod_{\beta \in \bar{B}} U(\beta)$ , where  $\prod_{\beta \in \bar{B}} U(\beta)$  is given the natural partial order of coordinatewise inclusion. But then, since each ultrafilter is countably closed by Corollary 4.2, it follows that  $\beta_2$  belongs to a cofinal subset of  $\Psi_{\kappa^+}(U(\beta_1)) = U(\beta_1, \Psi)$  and hence  $\beta_2 \in \bigcap U(\beta_1, \Psi)$ . This contradicts  $\beta_1 \in \bar{B} \subseteq B$ . ■

It now follows that if  $H(\xi)$  is defined to be the unique, if it exists, element of  $\nu$  such that  $\bigcap U(\xi, \Psi) = \{H(\xi)\}$  then  $H$  is defined for all but  $\kappa$  elements of  $\nu \setminus I_{\kappa^+}(\mathfrak{A})$ . The arguments of Lemmas 4.3 and 4.4 show that  $H^{-1}$  is defined for all but  $\kappa$  elements of  $\nu \setminus I_{\kappa^+}(\mathfrak{A}^\Psi)$ . Consequently,  $H$  is a bijection whose domain is  $E$  and range is  $E_\Psi$  such that  $|\nu \setminus E| \leq \kappa$  and  $|\nu \setminus E_\Psi| \leq \kappa$ . It may further be assumed that  $E_\Psi \cap I_{\kappa^+}(\mathfrak{A}^\Psi) = \emptyset$ .

All that remains to be shown is that  $[H\langle A \rangle] = \Psi([A])$  for each  $A \subseteq E$ . Since  $\Psi$  is an isomorphism and  $H$  induces an isomorphism, it suffices to show that if  $A, B \subseteq E$  and  $H\langle A \rangle \equiv^* \hat{\Psi}(B)$  then  $A \equiv^* B$ . So suppose that  $A$  and  $B$  provide a counterexample to this. It is then possible to find a countable  $C \subseteq H\langle A \rangle \cap \hat{\Psi}(B)$  such that there are countable  $A^* \subseteq A$  and  $B^* \subseteq B$  such that  $A^* \cap B^* = \emptyset$  and  $H\langle A^* \rangle \equiv^* \hat{\Psi}(B^*) \equiv^* C$ . That  $C$ ,  $A^*$  and  $B^*$  can be assumed countable uses an argument similar to that of Lemma 4.3.

Now use the density property of  $\mathcal{D}$  to find  $D \in \mathcal{D} \subseteq \mathfrak{A}$  such that  $A^* \subseteq D$  and  $D \cap B^* = \emptyset$ . Note that it is easy to see that  $H\langle F \cap E \rangle = \Psi_{\kappa^+}(F) \cap E_\Psi$  for any  $F \in \mathfrak{A}_{\kappa^+}$ , and in particular  $H\langle D \cap E \rangle = \Psi_{\kappa^+}(D) \cap E_\Psi$ . From this and the fact that  $I_{\kappa^+}(\mathfrak{A}^\Psi) \cap E_\Psi = \emptyset$ , it follows that

$$\begin{aligned} C &\subseteq^* H\langle A^* \rangle \cap \hat{\Psi}(B^*) \cap E_\Psi \\ &\subseteq^* H\langle D \cap E \rangle \cap \hat{\Psi}(\nu \setminus D) \cap E_\Psi = \Psi_{\kappa^+}(D) \cap \Psi_{\kappa^+}(\nu \setminus D) \cap E_\Psi, \end{aligned}$$

contradicting the fact that  $\Psi_{\kappa^+}$  is an isomorphism. ■

## 5. Automorphisms of $\mathcal{P}(\kappa)/\mathcal{F}in$ when $\text{cof}(\kappa) = \omega$

*Proof of Lemma 3.5.* Let  $\{\kappa_n\}_{n \in \omega}$  be an increasing sequence of cardinals cofinal in  $\kappa$ . Suppose that  $\Phi$  is an automorphism of  $\mathcal{P}(\kappa)/\mathcal{F}in$  and  $\hat{\Phi}$  is a lifting of it. Let  $\hat{\Phi}^{-1}$  be a lifting of  $\Phi^{-1}$  and note that  $\hat{\Phi}(\hat{\Phi}^{-1}(X)) \equiv^* X$  for all  $X$ .

Using the hypothesis, let  $F_n$  and  $S_n$  be such that

- (1)  $S_n \subseteq \kappa_n$  and  $|S_n| \leq 2^{\aleph_0}$ ,
- (2)  $F_n : \kappa_n \setminus S_n \rightarrow \kappa$ ,
- (3)  $F_n \langle X \rangle \in \Phi([X])$  for every  $X \subseteq \kappa_n \setminus S_n$ .

Let  $S = \bigcup_n S_n$ .

CLAIM 5.1. *For  $n \in m \in \omega$  there exists a finite set  $A_{n,m}$  such that  $F_n \upharpoonright (\kappa_n \setminus (S \cup A_{n,m})) \subseteq F_m$ .*

*Proof.* Suppose otherwise. There is then an infinite set  $A \subseteq \kappa_n \setminus S$  such that  $F_n \langle A \rangle \cap F_m \langle A \rangle = \emptyset$ . This contradicts the fact that  $F_n \langle A \rangle \in \Phi([A])$  and  $F_m \langle A \rangle \in \Phi([A])$ . ■

Now let  $\bar{S} = S \cup \bigcup_{n,m} A_{n,m}$  and  $F = \bigcup_n F_n \upharpoonright (\kappa \setminus \bar{S})$ . It suffices to show that  $F \langle X \rangle \in \Phi([X])$  for every  $X \subseteq \kappa \setminus \bar{S}$ .

To this end, suppose not and choose recursively  $X_\xi \subseteq \kappa \setminus \bar{S}$ , for  $\xi \in \omega_1$ , such that

- (4)  $|X_\xi| = \aleph_0$ ,
- (5)  $X_\xi \cap X_\eta = \emptyset$  if  $\xi \neq \eta$ ,
- (6)  $X_\xi \cap F^{-1}(\hat{\Phi}(\bigcup_{\eta \in \xi} X_\eta)) = \emptyset$ ,
- (7)  $X_\xi \cap \bigcup_{\eta \in \xi} \hat{\Phi}^{-1}(F \langle X_\eta \rangle) = \emptyset$ ,
- (8)  $F \langle X_\xi \rangle \cap \hat{\Phi}(X_\xi)$  is finite.

To see that this is possible note that if

$$S^\xi = \bar{S} \cup F^{-1} \left( \hat{\Phi} \left( \bigcup_{\eta \in \xi} X_\eta \right) \right) \cup \bigcup_{\eta \in \xi} \hat{\Phi}^{-1}(F \langle X_\eta \rangle) \cup \bigcup_{\eta \in \xi} X_\eta$$

then by Theorem 2.3,  $|S^\xi \setminus \bar{S}| \leq \aleph_1$  for every  $\xi \in \omega_1$ . Hence, if it is not possible to find  $X_\xi$ , then  $F$  and  $S^\xi$  witness that  $\Phi$  is  $2^{\aleph_0}$ -trivial.

Now let  $X = \bigcup_{\xi \in \omega_1} X_\xi$ .

CLAIM 5.2.  *$\hat{\Phi}(X) \cap F \langle X_\xi \rangle$  is finite for every  $\xi \in \omega_1$ .*

*Proof.* Let  $\xi \in \omega_1$  be given. Then

$$\hat{\Phi}(X) \equiv^* \hat{\Phi} \left( \bigcup_{\eta \in \xi} X_\eta \right) \cup \hat{\Phi}(X_\xi) \cup \hat{\Phi} \left( \bigcup_{\eta > \xi} X_\eta \right).$$

By (8) it follows that  $F \langle X_\xi \rangle \cap \hat{\Phi}(X_\xi)$  is finite, and by (6),

$$F \langle X_\xi \rangle \cap \hat{\Phi} \left( \bigcup_{\eta \in \xi} X_\eta \right) = \emptyset.$$



By (7) we see that  $\hat{\Phi}^{-1}(F\langle X_\xi \rangle) \cap (\bigcup_{\alpha > \xi} X_\alpha) = \emptyset$  and hence

$$\hat{\Phi}(\hat{\Phi}^{-1}(F\langle X_\xi \rangle)) \cap \hat{\Phi}\left(\bigcup_{\alpha > \xi} X_\alpha\right) \equiv^* \emptyset,$$

and so  $F\langle X_\xi \rangle \cap \hat{\Phi}(\bigcup_{\alpha > \xi} X_\alpha)$  is finite as well. ■

Now let  $g : \omega_1 \rightarrow \omega$  be such that  $\hat{\Phi}(X) \cap F\langle X_\xi \rangle \subseteq F\langle \kappa_{g(\xi)} \rangle$  for each  $\xi \in \omega_1$ . Then define  $g^+ : \omega_1 \rightarrow \omega$  such that  $F\langle X_\xi \cap \kappa_{g^+(\xi)} \rangle \not\subseteq F\langle \kappa_{g(\xi)} \rangle$  for each  $\xi \in \omega_1$ . To see that this is possible, note that the failure to find a suitable  $g^+(\xi)$  would imply that  $F\langle X_\xi \rangle \subseteq F\langle \kappa_{g(\xi)} \rangle$ , and hence  $X_\xi \subseteq \kappa_{g(\xi)}$ . Since  $F_{g(\xi)}$  induces  $\hat{\Phi}|\mathcal{P}(\kappa_{g(\xi)})/[\kappa_{g(\xi)}]^{<\aleph_0}$ , it would follow that  $F\langle X_\xi \rangle \equiv^* \hat{\Phi}(X_\xi)$ , contradicting (8). Now choose  $m, k \in \omega$  such that there is an infinite  $Z^* \subseteq \omega_1$  with  $g(\xi) = m$  and  $g^+(\xi) = k$  for  $\xi \in Z^*$ .

Now let  $Z = \bigcup_{\xi \in Z^*} X_\xi \cap \kappa_k$ . Since the  $X_\xi$  are pairwise disjoint and  $\hat{\Phi}(Z) \subseteq^* \hat{\Phi}(X)$ , it follows that

- $\hat{\Phi}(Z) \cap F\langle X_\xi \rangle \subseteq^* F\langle \kappa_m \rangle$  for all  $\xi \in Z^*$ ,
- $\hat{\Phi}(Z) \cap F\langle X_\xi \rangle \subseteq F\langle \kappa_m \rangle$  for all but finitely many  $\xi \in Z^*$ ,

and hence  $\hat{\Phi}(Z) \cap \bigcup_{\xi \in Z^*} F\langle X_\xi \rangle \subseteq^* F\langle \kappa_m \rangle$ . Furthermore,  $Z \subseteq \kappa_k \setminus \bar{S}$  and so  $\hat{\Phi}(Z) \equiv^* F\langle Z \rangle$ . Moreover  $F\langle Z \rangle \setminus F\langle \kappa_m \rangle$  is infinite because of the definition of  $g^+$  and the fact that the  $X_\xi$  are pairwise disjoint. This contradicts the fact that  $F\langle Z \rangle \setminus F\langle \kappa_m \rangle \subseteq^* \hat{\Phi}(Z) \cap F\langle Z \rangle \subseteq \hat{\Phi}(Z) \cap \bigcup_{\xi \in Z^*} F\langle X_\xi \rangle \subseteq^* F\langle \kappa_m \rangle$ . ■

## 6. Automorphisms of $\mathcal{P}(\kappa^{\aleph_0})/\mathcal{F}in$ when $\kappa$ has countable cofinality

*Proof of Lemma 3.6.* Suppose that  $\kappa^+ \leq \rho \leq \kappa^{\aleph_0}$  and that  $\rho$  is the least cardinal such that there is an automorphism of  $\mathcal{P}(\rho)/\mathcal{F}in$  which is not  $\kappa$ -trivial. Note that by Lemmas 3.4 and 3.5,  $\rho$  is a regular cardinal, and by hypothesis  $\kappa \neq 2^{\aleph_0}$  and so  $\rho > (2^{\aleph_0})^+$ . Let  $\{\kappa_n\}_{n \in \omega}$  be an increasing sequence of cardinals converging to  $\kappa$ . Let  $R \subseteq \prod_{n \in \omega} \kappa_n$  be such that  $|R| = \rho$  and suppose that there is an automorphism  $\Phi$  of  $\mathcal{P}(R)/\mathcal{F}in$  that is not  $\kappa$ -trivial.

For  $k \in \omega$  let  $T_k = \prod_{n \in k} \kappa_n$ , so that  $T_k$  consists of sequences  $t$  of length  $k$  such that  $t(j) \in \kappa_j$  for each  $j \in k = \mathbf{domain}(t)$ . For  $W \subseteq T_k$  let  $C(W) = \{f \in R \mid f|k \in W\}$ . Let  $\mathfrak{B}$  be the Boolean subalgebra of  $\mathcal{P}(R)$  generated by sets of the form  $C(W)$  where  $W \subseteq T_k$  for some  $k \in \omega$ , and note that  $|\mathfrak{B}| = \kappa$ . Let  $\hat{\Phi}$  be a lifting of  $\Phi$  and  $\Phi_{\kappa^+} : \mathfrak{B}_{\kappa^+} \rightarrow \mathfrak{B}_{\kappa^+}^{\hat{\Phi}}$ . This is a well defined isomorphism by Lemma 2.6. Since no confusion can arise because of it, the symbol  $C(W)$  will continue to be used to denote  $C(W) \setminus I_{\kappa^+}(\mathfrak{B})$  in the algebra  $\mathfrak{B}_{\kappa^+}$ . Note, however, that it may be that  $C(W) = \emptyset$  for some

non-empty  $W$ ; this depends on  $R$  of course. As well, for singletons  $\{t\}$  the notation  $C(t)$  will be used in place of the more cumbersome  $C(\{t\})$ .

Next observe that

$$(6.1) \quad (\forall m \in \omega)(\forall W \subseteq T_m) \quad \Phi_{\kappa^+}(C(W)) = \bigcup_{t \in W} \Phi_{\kappa^+}(C(t))$$

or, in other words,

$$\left| \hat{\Phi}(C(W)) \setminus \bigcup_{t \in W} \hat{\Phi}(C(t)) \right| \leq \kappa.$$

To see this, suppose not and let  $m \in \omega$ ,  $W \subseteq T_m$  and  $A \subseteq R$  be such that

$$\hat{\Phi}(A) \equiv^* \hat{\Phi}(C(W)) \setminus \bigcup_{t \in W} \hat{\Phi}(C(t)),$$

and note that  $|A| \geq \kappa_m^+$  by Theorem 2.3. Hence, since  $A \subseteq^* C(W)$ , there is  $t \in W$  such that  $|A \cap C(t)| \geq \kappa_m^+$ . Since  $\Phi$  is a homomorphism, this contradicts  $\hat{\Phi}(A) \cap \hat{\Phi}(C(t)) \equiv^* \emptyset$ . Furthermore,

$$(6.2) \quad \text{if } s \subseteq t \text{ then } \Phi_{\kappa^+}(C(s)) \supseteq \Phi_{\kappa^+}(C(t)).$$

For each  $f \in R \setminus I_{\kappa^+}(\mathfrak{B})$  let  $U(f)$  be the ultrafilter  $\{B \in \mathfrak{B}_{\kappa^+} \mid f \in B\}$  and let  $V(f)$  be the ultrafilter  $\{B \in \mathfrak{B}_{\kappa^+}^\Phi \mid f \in B\}$ . Let  $U_\Phi(f)$  be the ultrafilter  $\{\Phi_{\kappa^+}(B) \in \mathfrak{B}_{\kappa^+}^\Phi \mid B \in U(f)\}$ .

CLAIM 6.1. *Let*

$$S = \{f \in R \setminus I_{\kappa^+}(\mathfrak{B}) \mid (\exists h \in R \setminus I_{\kappa^+}(\mathfrak{B})) V(h) = U_\Phi(f)\}.$$

*Then*

$$|R \setminus (S \cup I_{\kappa^+}(\mathfrak{B}))| \leq 2^{\aleph_0}.$$

*Proof.* If the claim fails then let  $A \subseteq R \setminus (S \cup I_{\kappa^+}(\mathfrak{B}))$  be such that  $|A| > 2^{\aleph_0}$ . There must then be some  $k \in \omega$  such that  $|\{f \upharpoonright k \mid f \in A\}| > 2^{\aleph_0}$ . Hence there is  $A_1 \subseteq A$  of cardinality greater than  $2^{\aleph_0}$  such that the mapping  $f \mapsto f \upharpoonright k$  is one-to-one on  $A_1$ .

Since  $A_1 \cap S = \emptyset$  it follows that  $\bigcap_{n \in \omega} \Phi_{\kappa^+}(C(f \upharpoonright n)) = \emptyset$  for each  $f \in A_1$ . Since  $\Phi_{\kappa^+}(C(f \upharpoonright k)) \cap \hat{\Phi}(A_1)$  is finite for each  $f \in A_1$  it follows that for each  $f \in A_1$  there is  $m_f \geq k$  such that

$$\Phi_{\kappa^+}(C(f \upharpoonright m_f)) \cap \Phi_{\kappa^+}(C(f \upharpoonright k)) \cap \hat{\Phi}(A_1) = \emptyset,$$

and hence

$$(6.3) \quad \Phi_{\kappa^+}(C(f \upharpoonright m_f)) \cap \hat{\Phi}(A_1) = \emptyset$$

because  $m_f \geq k$  and (6.2) holds.

Now let  $A_2 \subseteq A_1$  be infinite and  $m$  be such that  $m_f = m$  for all  $f \in A_2$ , and let  $Z = \{f \upharpoonright m \mid f \in A_2\}$ . Then  $A_2 \subseteq C(Z)$  and  $A_2 \cap I_{\kappa^+}(\mathfrak{B}) = \emptyset$ , and

so  $C(Z) \neq \emptyset$ . Moreover, from (6.1) it follows that

$$\Phi_{\kappa^+}(C(Z)) = \bigcup_{f \in A_2} \Phi_{\kappa^+}(C(f \upharpoonright m)),$$

and hence  $\Phi_{\kappa^+}(C(Z)) \cap \hat{\Phi}(A_1) = \emptyset$  by (6.3). However, this contradicts  $\hat{\Phi}(A_2) \subseteq^* \hat{\Phi}(A_1)$  and  $\hat{\Phi}(A_2) \subseteq^* \Phi_{\kappa^+}(C(Z))$ . ■

CLAIM 6.2. *Let*

$$\tilde{S} = \{f \in R \setminus I_{\kappa^+}(\mathfrak{B}) \mid (\forall h \in R \setminus I_{\kappa^+}(\mathfrak{B})) \text{ if } V(h) = V(f) \text{ then } h = f\}.$$

Then  $|R \setminus (\tilde{S} \cup I_{\kappa^+}(\mathfrak{B}))| \leq 2^{\aleph_0}$ .

*Proof.* Suppose that  $\{(f_\xi, g_\xi)\}_{\xi \in \mathfrak{c}^+}$  are disjoint pairs such that  $V(f_\xi) = V(g_\xi)$  and  $\{f_\xi, g_\xi\} \cap I_{\kappa^+}(\mathfrak{B}) = \emptyset$  for each  $\xi \in \mathfrak{c}^+$ . Let  $A$  be such that  $\hat{\Phi}(A) = \{f_\xi\}_{\xi \in \mathfrak{c}^+}$ . Using arguments as in Claim 6.1, find  $k \in \omega$  and  $A_1 \subseteq A$  and  $Z \subseteq \mathfrak{c}^+$  such that

- $|A_1| = |Z| = \mathfrak{c}^+$ ,
- the mapping  $f \mapsto f \upharpoonright k$  is one-to-one on  $A_1$ ,
- $\hat{\Phi}(A_1) \equiv^* \{f_\xi\}_{\xi \in Z}$ .

Then find  $B, Z^* \subseteq Z$  and  $k^* \geq k$  such that

- $|B| = |Z^*| = \mathfrak{c}^+$ ,
- the mapping  $f \mapsto f \upharpoonright k^*$  is one-to-one on  $B$ ,
- $\hat{\Phi}(B) \equiv^* \{g_\xi\}_{\xi \in Z^*}$ .

Since  $A_1$  and  $B$  are almost disjoint, it is possible to find  $m \geq k^*$  and an infinite  $\bar{A} \subseteq A_1$  such that if  $f \in \bar{A}$  and  $g \in B$  then  $f \upharpoonright m \neq g \upharpoonright m$ . Let  $W = \{f \upharpoonright m \mid \xi \in \bar{A}\}$ .

Hence  $\hat{\Phi}(\bar{A}) \subseteq^* \Phi_{\kappa^+}(C(W))$  and  $\hat{\Phi}(B) \cap \Phi_{\kappa^+}(C(W))$  is finite. It follows that for all but finitely many  $\xi$  such that  $f_\xi \in \hat{\Phi}(\bar{A})$  it must be the case that  $\Phi_{\kappa^+}(C(W)) \in V(f_\xi)$  but  $\Phi_{\kappa^+}(C(W)) \notin V(g_\xi)$ , contradicting  $V(f_\xi) = V(g_\xi)$ . ■

A similar argument shows the following.

CLAIM 6.3. *Let*

$$S^* = \{f \in R \setminus I_{\kappa^+}(\mathfrak{B}) \mid (\forall h \in R \setminus I_{\kappa^+}(\mathfrak{B})) \text{ if } U_\Phi(h) = U_\Phi(f) \text{ then } f = h\}.$$

Then  $|R \setminus (S \cup I_{\kappa^+}(\mathfrak{B}))^*| \leq 2^{\aleph_0}$ .

It follows from Claims 6.1–6.3 that letting  $E = S \cap \tilde{S} \cap S^* \setminus I_{\kappa^+}(\mathfrak{B})$  we can find a well defined, one-to-one function  $F : E \rightarrow \prod_{n \in \omega} \kappa_n$  such that  $V(F(f)) = U_\Phi(f)$  for every  $f \in E$ . Observe that

$$(6.4) \quad (\forall A \in \mathfrak{B}_{\kappa^+}) F \langle A \cap \tilde{E} \rangle = \Phi_{\kappa^+}(A) \cap F \langle \tilde{E} \rangle \equiv^* \hat{\Phi}(A) \cap F \langle \tilde{E} \rangle$$

for any  $\tilde{E} \subseteq E$  by the definition of  $F$ .

CLAIM 6.4. *Suppose that  $W \subseteq E$  and  $\Psi : W \rightarrow W$  are such that  $\Psi \langle A \rangle \equiv^* \hat{\Phi}(A)$  for each  $A \subseteq W$ . Then  $|\{w \in W \mid \Psi(w) \neq F(w)\}| \leq 2^{\aleph_0}$ .*

*Proof.* If the claim fails then it is possible to find  $k \in \omega$  and

$$Z \subseteq \{w \in W \cap E \mid \Psi(w) \neq F(w)\}$$

such that  $|Z| > 2^{\aleph_0}$  and the mapping  $z \mapsto z \upharpoonright k$  is one-to-one. For each  $z \in Z$  there is some  $m_z \geq k$  and  $t_z^0 : m_z \rightarrow \kappa$  and  $t_z^1 : m_z \rightarrow \kappa$  such that  $t_z^0 \neq t_z^1$  and

$$(6.5) \quad \Psi(z) \in \Phi_{\kappa^+}(C(t_z^0)),$$

$$(6.6) \quad F(z) \in \Phi_{\kappa^+}(C(t_z^1)).$$

Observe that the definition of  $F$  implies that  $t_z^1 = z \upharpoonright m_z$ . Let  $\bar{Z} \in [Z]^{\aleph_1}$  and  $m$  be such that  $m_z = m$  for all  $z \in \bar{Z}$ .

Since the  $t_z^1$  are all distinct, there is an infinite  $\bar{\bar{Z}} \subseteq \bar{Z}$  such that if  $W_i = \{t_z^i \mid z \in \bar{\bar{Z}}\}$  then  $W_0 \cap W_1 = \emptyset$ , and hence  $C(W_0) \cap C(W_1) = \emptyset$ . The construction of  $\Phi_{\kappa^+}$  guarantees that

$$(6.7) \quad \Phi_{\kappa^+}(C(W_0)) \cap \Phi_{\kappa^+}(C(W_1)) = \emptyset.$$

By (6.4) and the hypothesis on  $\Psi$  it follows that

$$(6.8) \quad F \langle C(W_i) \cap W \cap E \rangle \equiv^* \hat{\Phi}(C(W_i)) \cap F \langle W \cap E \rangle \equiv^* \Psi \langle C(W_i) \cap W \cap E \rangle.$$

Since  $\bar{\bar{Z}} \subseteq C(W_1)$  it follows that

$$\Psi \langle \bar{\bar{Z}} \rangle \subseteq^* \Psi \langle C(W_1) \cap W \cap E \rangle \equiv^* \hat{\Phi}(C(W_1) \cap W \cap E) \subseteq^* \hat{\Phi}(C(W_1)).$$

On the other hand,

$$\Psi \langle \bar{\bar{Z}} \rangle \subseteq \bigcup_{z \in \bar{\bar{Z}}} \Phi_{\kappa^+}(C(t_z^0)) = \Phi_{\kappa^+}(C(W_0)) \subseteq \hat{\Phi}(C(W_0))$$

by (6.1) and (6.5) and the definition of  $\Phi_{\kappa^+}$ . This, of course, contradicts  $\hat{\Phi}(C(W_0)) \cap \hat{\Phi}(C(W_1)) \equiv^* \emptyset$ . ■

Now let  $\{r_\xi\}_{\xi \in \rho}$  be an enumeration of  $E \subseteq R$  and define  $E_\alpha = \{r_\xi\}_{\xi \in \alpha}$ . Two cases need to be considered, the first one being that  $\rho = \kappa^+$ . Observe that if  $\mathfrak{M}$  is an elementary submodel of  $(H(\rho^+), \{r_\xi\}_{\xi \in \rho}, \hat{\Phi}, \in)$  and  $\mathfrak{M} \cap \rho^+$  is an ordinal of uncountable cofinality then  $\hat{\Phi}(\{r_\xi \mid \xi \in \mathfrak{M}\}) \equiv^* \{r_\xi \mid \xi \in \mathfrak{M}\}$ . Recalling that  $\rho$  is a regular cardinal greater than  $\mathfrak{c}^+$ , we deduce that

$$Y = \{\xi \in \rho \mid \hat{\Phi}(E_\xi) \equiv^* E_\xi \text{ and } \mathbf{cof}(\xi) > \mathfrak{c}\}$$

is a stationary set. The third hypothesis of the lemma then implies that for each  $\xi \in Y$  the restriction of  $\Phi$  to  $\mathcal{P}(E_\xi)/\mathcal{F}in$  is an automorphism of  $\mathcal{P}(E_\xi)/\mathcal{F}in$  that must be  $2^{\aleph_0}$ -trivial.

The other case is that  $\rho > \kappa^+$ . In this case

$$Y = \{\xi \in \rho \mid \hat{\Phi}(E_\xi) \equiv^* E_\xi \text{ and } \mathbf{cof}(\xi) = \kappa^+\}$$

is a stationary set. Therefore the minimality of  $\rho$  shows that for each  $\xi \in Y$  the restriction of  $\Phi$  to  $\mathcal{P}(E_\xi)/\mathcal{F}in$  is an automorphism of  $\mathcal{P}(E_\xi)/\mathcal{F}in$  that must be  $\kappa$ -trivial.

In either case there is some  $\alpha(\xi) \in \xi$  and a function  $\Psi$  such that  $\hat{\Phi}(A) \equiv^* \Psi\langle A \rangle$  for each  $A \subseteq E_\xi \setminus \alpha(\xi)$ . By Claim 6.4 there is then  $\beta(\xi) \in \xi \setminus \alpha(\xi)$  such that  $\Psi(\eta) = F(\eta)$  provided that  $\beta(\xi) \in \eta \in \xi$ . The stationarity of  $Y$  yields a  $\beta \in \rho$  such that  $F\langle A \rangle \equiv^* \hat{\Phi}(A)$  for every bounded  $A \subseteq \rho \setminus \beta$ . This is enough to conclude that  $\Phi$  is  $|\beta|$ -trivial, and Lemma 3.2 and the minimality of  $\rho$  imply that  $\Phi$  is  $\kappa$ -trivial. ■

**7. Remarks and questions.** Observe that Theorems 1.1 and 3.1 imply that if  $\kappa$  is less than the first inaccessible cardinal and  $\Phi$  is a non-trivial automorphism of  $\mathcal{P}(\kappa)/\mathcal{F}in$  then there is  $X \in [\kappa]^{N_1}$  such that  $\Phi\upharpoonright\mathcal{P}(X)/\mathcal{F}in$  is also non-trivial. Hence it is of interest to understand the non-trivial automorphisms of  $\mathcal{P}(\omega_1)/\mathcal{F}in$ . Of course, among them there are ones that are obtained by taking a non-trivial automorphism of  $\mathcal{P}(\omega)/\mathcal{F}in$  and extending it to all of  $\mathcal{P}(\omega_1)/\mathcal{F}in$  by the identity on sets disjoint from  $\omega$ . Hence the following definition is natural in this context.

**DEFINITION 7.1.** An automorphism  $\Phi$  of  $\mathcal{P}(\omega_1)/\mathcal{F}in$  will be called *non-trivially non-trivial* if it is non-trivial (as defined in §1) but  $\Phi\upharpoonright Z$  is trivial for every countable  $Z \subseteq \omega_1$ .

**QUESTION 7.2.** Is it consistent with set theory that there is an automorphism  $\Phi$  of  $\mathcal{P}(\omega_1)/\mathcal{F}in$  that is non-trivially non-trivial?

Recall from §1 that Velickovic [7] showed that the conjunction of OCA and MA implies that every automorphism of  $\mathcal{P}(\omega_1)/\mathcal{F}in$  is trivial. However it might be that there is a stronger result answering the following question.

**QUESTION 7.3.** Does it follow from the fact that every automorphism of  $\mathcal{P}(\omega)/\mathcal{F}in$  is trivial that every automorphism of  $\mathcal{P}(\omega_1)/\mathcal{F}in$  is trivial?

Note that in any model with an automorphism  $\Phi$  of  $\mathcal{P}(\omega_1)/\mathcal{F}in$  providing a positive answer to Question 7.2 there must be a family of injections  $f_\xi : \xi \rightarrow \omega_1$  for each  $\xi \in \omega_1$  such that if  $\xi \in \eta$  then  $f_\xi \equiv^* f_\eta\upharpoonright\xi$  and  $\Phi\upharpoonright\mathcal{P}(\xi)/\mathcal{F}in$  is induced by  $f_\xi$  but there is no  $f : \omega_1 \rightarrow \omega_1$  threading the  $f_\xi$ —in other words, there is no  $f$  such that  $f_\xi \equiv^* f\upharpoonright\xi$  for all  $\xi$ .

So given a potential family of functions  $\{f_\xi\}_{\xi \in \omega_1}$ , what needs to be done is to define the values of  $\Phi(X)$  for uncountable  $X \subseteq \omega_1$  without adding a function  $f$  threading the  $f_\xi$ . The natural partial order for adding the unique element of  $\mathcal{P}(\omega_1)/\mathcal{F}in$  that must be equal to  $\Phi(X)$  will preserve  $\omega_1$  for certain, carefully constructed families of functions  $\{f_\xi\}_{\xi \in \omega_1}$ . However, the countable support iteration poses several problems. Larson and McKenney

and, independently, Rinot and Steprāns have shown that certain instances of this partial order can be iterated, with countable support, to yield a model where there is a non-trivial automorphism of the Boolean subalgebra of  $\mathcal{P}(\omega_1)/\mathcal{F}in$  generated by the countable sets and a maximal independent family of subsets of  $\omega_1$  which is trivial when restricted to any countable set.

On the other hand, there is also the question of inaccessible  $\kappa$ .

**QUESTION 7.4.** Is it consistent, relative to the consistency of an inaccessible cardinal, that there is a non-trivially non-trivial automorphism of  $\mathcal{P}(\kappa)/\mathcal{F}in$  where  $\kappa$  is inaccessible?

Observe that if  $\kappa$  provides a positive answer to Question 7.4 then  $\kappa$  cannot be too large a cardinal. For example, if  $\kappa$  is the least cardinal such that there is a non-trivially non-trivial automorphism  $\Phi$  of  $\mathcal{P}(\kappa)/\mathcal{F}in$  then it cannot carry a normal  $\kappa$ -additive ultrafilter. To see this, note that for each  $\xi \in \kappa$  there is some  $f_\xi : \xi \rightarrow \kappa$  such that  $\Phi \upharpoonright \mathcal{P}(\xi)/\mathcal{F}in$  is induced by  $f_\xi$ . Of course,  $f_\xi \equiv^* f_\eta \upharpoonright \xi$  for  $\xi \in \eta$ . In other words, for each  $\xi \in \eta \in \kappa$  there is  $a_{\xi,\eta} \in [\xi]^{<\aleph_0}$  such that  $f_\eta \upharpoonright \xi \setminus a_{\xi,\eta} = f_\xi \upharpoonright \xi \setminus a_{\xi,\eta}$ . The normal  $\kappa$ -additive ultrafilter yields a set  $X \subseteq \kappa$  of cardinality  $\kappa$  and an  $a$  such that  $a_{\xi,\eta} = a$  for all  $\{\xi, \eta\} \in [X]^2$ . Hence  $\bigcup_{\xi \in X} f_\xi \upharpoonright \xi \setminus a$  induces  $\Phi$ .

Theorem 3.1 suggests an alternative version of non-trivial non-triviality.

**DEFINITION 7.5.** An automorphism  $\Phi$  of  $\mathcal{P}(\omega_1)/\mathcal{F}in$  will be called *very non-trivially non-trivial* if  $\Phi \upharpoonright Z$  is non-trivial for every co-countable  $Z \subseteq \omega_1$ .

Results about the existence or non-existence of very non-trivially non-trivial would also be of interest.

**Acknowledgements.** The first author's research for this paper was partially supported by the United States–Israel Binational Science Foundation (grant no. 2006108), and by the National Science Foundation (grant no. NSF-DMS 0600940). The second author's research for this paper was partially supported by NSERC of Canada. This is Number 1071 in the first author's list of publications that can be found at <http://shelah.logic.at>.

## References

- [1] B. Balcar and R. Frankiewicz, *To distinguish topologically the spaces  $m^*$ . II*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. 26 (1978), 521–523.
- [2] D. Chodounský, A. Dow, K. P. Hart and H. de Vries, *The Katowice problem and autohomeomorphisms of  $\omega_0^*$* , arXiv:1307.3930 (2013).
- [3] P. Larson and P. McKenney, *Automorphisms of  $\mathcal{P}(\lambda)/\mathcal{I}_\kappa$* , Fund. Math. 233 (2016), 271–291.
- [4] W. Rudin, *Homogeneity problems in the theory of Čech compactifications*, Duke Math. J. 23 (1956), 409–419.
- [5] W. Rudin, *Note of correction*, Duke Math. J. 23 (1956), 633.

- [6] S. Shelah, *Proper Forcing*, Lecture Notes in Math. 940, Springer, Berlin, 1982.
- [7] B. Velickovic, *OCA and automorphisms of  $\mathcal{P}(\omega)/\text{fin}$* , *Topology Appl.* 49 (1993), 1–13.

Saharon Shelah  
Department of Mathematics  
Rutgers University  
Hill Center  
Piscataway, NJ 08854-8019, U.S.A.

*Current address:*  
Institute of Mathematics  
Hebrew University  
Givat Ram, Jerusalem 91904, Israel  
E-mail: shelah@math.rutgers.edu

Juris Steprāns  
Department of Mathematics  
York University  
4700 Keele Street  
Toronto, ON, Canada M3J 1P3  
E-mail: steprans@yorku.ca

