

Set Mappings on 4-Tuples

Shahram Mohsenipour and Saharon Shelah

Abstract In this article, we study set mappings on 4-tuples. We continue a previous work of Komjath and Shelah by getting new finite bounds on the size of free sets in a generic extension. This is obtained by an entirely different forcing construction. Moreover, we prove a ZFC result for set mappings on 4-tuples. Also, as another application of our forcing construction, we give a consistency result for set mappings on triples.

1 Introduction

In this article, we continue the work of Komjath and Shelah [9]. Our main objects of study here are set mappings that are, for our current purpose, functions of the type $f: [\lambda]^k \rightarrow [\lambda]^{<\mu}$ for some natural number $k \geq 1$ and cardinals λ, μ , which satisfy $f(\bar{x}) \cap \bar{x} = \emptyset$ for every $\bar{x} \in [\lambda]^k$. The motivation in this part of combinatorial set theory is to know how large free sets exist. A subset H of λ is called *free* if $f(\bar{x}) \cap H = \emptyset$ for every $\bar{x} \in [H]^k$. The case $k = 1$ was settled by Hajnal in [6] where he showed that if $\mu < \lambda$, then there is a free set of size λ (we call it *Theorem A* for later references). The case $k > 1$ came to attention when Kuratowski and Sierpinski proved that for set mappings on $[\lambda]^k$, there always exists a free set of size $k + 1$ if and only if $\lambda \geq \mu^{+k}$ (see Erdős, Hajnal, Máté, and Rado [1, Section 45]). It is interesting to know that, assuming GCH, the authors of [1] showed that if $\lambda \geq \mu^{+k}$, then there is a free set of cardinality μ^+ . Coming back to ZFC, Hajnal and Máté [7] (and later [1, Section 46]) managed to improve Kuratowski and Sierpinski's result for the cases $k = 2, 3$ by showing that if $\lambda \geq \mu^{+k}$, then f has arbitrary large finite sets (we call it *Theorem B* for the case $k = 3$). The case $k = 4$ remained open until Komjath and Shelah in [9] showed that it is consistent with ZFC that there are set mappings with finite bounds on their size of free sets. More precisely, suppose that μ is a regular cardinal, that $\lambda = \mu^{+n}$, $n \in \omega$, and that GCH holds for every μ^{+l} ($l < n$), that

Received September 19, 2015; accepted March 21, 2016

First published online June 26, 2018

2010 Mathematics Subject Classification: Primary 03E05, 03E35

Keywords: combinatorial set theory, set mappings

© 2018 by University of Notre Dame 10.1215/00294527-2018-0002

is, $2^{\mu^{+l}} = \mu^{+(l+1)}$. Then there exist a natural number t_n and a cardinal-preserving generic extension in which there is a set mapping $f: [\lambda]^4 \rightarrow [\lambda]^{<\mu}$ with no free set of size t_n . The bound t_n is a Ramsey number which is defined inductively to be the least natural number satisfying the Ramsey relation $t_n \rightarrow (t_{n-1}, 7)^5$, $t_0 = 5$; hence $n \mapsto t_n$ is essentially the tower function, that is, iterating exponentiation n -times. In general, the Ramsey relation $a \rightarrow (b, c)^r$ means that the following statement is true. Whenever the r -element subsets of an a -element set are colored with two colors, say, 0 and 1, then either there exists a b -element with all its r -tuples colored with 0 or there exists a c -element subset whose r -tuples are all colored with 1. Subsequently, Komjath and Shelah [9] proved another independence result to show that the results in [7] and [1] for cases $k = 2, 3$ are optimal. In precise terms, let μ be a regular cardinal, let $\lambda = \mu^{+n}$, $n \in \omega$, and assume that GCH holds for every μ^{+l} ($l < n$). Then there is a cardinal-preserving generic extension in which there exists a set mapping $f: [\lambda]^2 \rightarrow [\lambda]^{<\mu}$ with no free set of size ω . This easily implies the similar result for $k = 3$. Coming back to the case $k = 4$, we should say that this is not the end of the story. Now we definitely have the task of improving the bound t_n in front of us, which would be more serious if we notice that the following question is open.

Question 1.1 (Gillibert and Wehrung [4]) Suppose that μ is an infinite cardinal and that $\lambda = \mu^{+4}$. Does any set mapping $f: [\lambda]^4 \rightarrow [\lambda]^{<\mu}$ have a free set of size 7?

Recall that the Kuratowski–Sierpinski theorem guarantees the existence of a free set of size 5. In fact, Hajnal and Máté [7] had asked in 1975 whether the above set mapping $f: [\lambda]^4 \rightarrow [\lambda]^{<\mu}$ has a free set of size 6. This question had remained open until 2008, when Gillibert [3], by using algebraic tools of a completely different nature, gave a positive answer to it. We refer the reader to [4] for a self-contained proof of this fact and other interesting results.

Now we are ready to briefly describe how this article is organized. In Section 2 we obtain another bound s_n instead of t_n with a different forcing construction. The new bound s_n is also a Ramsey number and is defined to be the least natural number satisfying $s_n \rightarrow (5)_{3^{n+1}}^3$ and essentially is triple exponentiation. In general, for natural numbers a, b, c , and r , the relation $a \rightarrow (b)_c^r$ means that the following assertion is true. Whenever the r -element subsets of an a -element set are colored with c colors, there is a b -element subset whose r -element subsets have the same color. Note that by the Erdős and Hajnal theorem quoted above, GCH must fail in the generic extension. We manage to control our construction in such a way that in the generic extension we will have $2^\mu = 2^{\mu^+} = \dots = 2^{\mu^{+(n-1)}} = \lambda$. In Section 3 we will compare the two bounds s_n and t_n asymptotically by using Ramsey theory to show that s_n is much better than t_n when n tends to infinity. Motivated by our main results in Section 2, we consider set mappings on 4-tuples with a restriction on the location of the image in Section 4 and get a ZFC result. In fact, in both of the forcing constructions in this article and in [9], we find that the constructed set mappings $f: [\lambda]^4 \rightarrow [\lambda]^{<\mu}$ in the generic extension have this property: for all $x_0 < x_1 < x_2 < x_3$ in $[\lambda]^4$, $f\{x_0, x_1, x_2, x_3\} \subset (x_1, x_2)$. By modifying Hajnal's proof of Theorem B, we show that this is necessary; more precisely, we show that if $\lambda \geq \mu^{+3}$, then any set mapping $f: [\lambda]^4 \rightarrow [\lambda]^{<\mu}$ with the additional property, for all $x_0 < x_1 < x_2 < x_3$ in $[\lambda]^4$, $f\{x_0, x_1, x_2, x_3\} \cap (x_1, x_2) = \emptyset$, will have arbitrary large finite free sets. In Section 5 we consider a similar situation for set mapping on triples. This time we are motivated by Komjath and Shelah's second construction

in [9], and then we will deal with set mappings on triples with a restriction on the location of the image. As an application of our forcing in Section 2, we obtain a negative consistency result.

Notation. Cardinals are identified with initial ordinals. If S is a set and κ is a cardinal, then $[S]^\kappa = \{X \subset S : |X| = \kappa\}$, $[S]^{<\kappa} = \{X \subset S : |X| < \kappa\}$, $[S]^{\leq\kappa} = \{X \subset S : |X| \leq \kappa\}$. If A and B are subsets of an ordered set, then $A < B$ means that $x < y$ whenever $x \in A$ and $y \in B$. Let $F: [A]^k \rightarrow B$ be a function on finite subsets; we always write $F\{a_1, \dots, a_k\}$ rather than $F(\{a_1, \dots, a_k\})$. If λ is a cardinal and $\alpha \in \lambda$, then by $(\alpha, +\infty)$ we mean $\{x \in \lambda : x > \alpha\}$.

2 The Main Theorems

We begin with a definition and two simple lemmas.

Definition 2.1 Suppose that A is a set, that $k \geq 2$, $F: [A]^k \rightarrow \mathcal{P}(A)$ is a set mapping, and that Γ is a set of functions of the form $\rho: [A]^2 \rightarrow L_\rho$, where L_ρ is a linear ordering with the order $<_\rho$; pedantically Γ is the set of pairs (ρ, L_ρ) for $\rho \in \Gamma$. We say that Γ k -generates F if for all $\bar{\gamma} \in [A]^k$, we have $x \in F(\bar{\gamma})$ if and only if $x \in A$ and

$$(\forall \rho \in \Gamma)(\forall \gamma \in \bar{\gamma})(\exists \gamma', \gamma'' \in \bar{\gamma}) \quad [\gamma' \neq \gamma'' \wedge \rho\{x, \gamma\} \leq_\rho \rho\{\gamma', \gamma''\}].$$

We note that in this article all the linear orderings L_ρ will be some cardinals with their natural \in -orderings. It is evident from the above definition that for any Γ as above and any $k \geq 2$ there is a unique set mapping $F: [A]^k \rightarrow \mathcal{P}(A)$ such that Γ k -generates F , so we may denote it by $F = F_\Gamma$.

Definition 2.2 In Definition 2.1, suppose $B \subset A$. By $\Gamma|_B$ we mean $\{\rho|_B : \rho \in \Gamma\}$.

The following two very easy lemmas will be very useful later in this article.

Lemma 2.3 Let $F_1: [A_1]^k \rightarrow \mathcal{P}(A_1)$, $F_2: [A_2]^k \rightarrow \mathcal{P}(A_2)$ ($k \geq 2$) be set mappings with $A_1 \subset A_2$. Let Γ_1, Γ_2 be sets of functions such that Γ_i k -generates F_i ($i = 1, 2$) and $\Gamma_2|_{A_1} \subset \Gamma_1$. Then $\forall \bar{\gamma} \in [A_1]^k$, and we have $F_1(\bar{\gamma}) \subset F_2(\bar{\gamma})$.

Lemma 2.4 Let $F_1: [A_1]^k \rightarrow \mathcal{P}(A_1)$, $F_2: [A_2]^k \rightarrow \mathcal{P}(A_2)$ ($k \geq 2$) be set mappings with $A_1 \subset A_2$. Let Γ_1, Γ_2 be sets of functions such that Γ_i k -generates F_i ($i = 1, 2$) and $\Gamma_2|_{A_1} = \Gamma_1$. Then $\forall \bar{\gamma} \in [A_1]^k$, and we have $F_1(\bar{\gamma}) = F_2(\bar{\gamma}) \cap A_1$.

Theorem 2.5 Assume that $n < \omega$, $k \geq 2$, and $\lambda = \mu^{+n}$ for some regular cardinal $\mu = \mu^{<\mu}$. Suppose

$$2^\mu = \mu^+, 2^{\mu^+} = \mu^{++}, \dots, 2^{\mu^{+(n-1)}} = \mu^{+n} \quad (\text{when } n > 0).$$

Then there is a notion of forcing $\mathbb{P}_n = \mathbb{P}_n^{\mu, \lambda}$ such that $|\mathbb{P}_n| = \lambda$, \mathbb{P}_n is $(< \mu)$ -complete and collapses no cardinal, and in $V^{\mathbb{P}_n}$:

(i) there are functions

$$\rho_0: [\lambda]^2 \rightarrow \mu^{+n}, \rho_1: [\lambda]^2 \rightarrow \mu^{+(n-1)}, \dots, \rho_n: [\lambda]^2 \rightarrow \mu$$

such that if $f: [\lambda]^k \rightarrow \mathcal{P}(\lambda)$ is k -generated by $\Gamma = \{\rho_0, \dots, \rho_n\}$, then $\forall \bar{\gamma} \in [\lambda]^k$ and we have $|f(\bar{\gamma})| < \mu$, which means that $f: [\lambda]^k \rightarrow [\lambda]^{<\mu}$;

(ii) $2^\mu = 2^{\mu^+} = \dots = 2^{\mu^{+(n-1)}} = \lambda$.

Proof This is by induction on n . We first prove the case $n = 0$ in ZFC. Set $\rho_0: [\lambda]^2 \rightarrow \lambda$ by $\rho_0\{x, y\} = \max\{x, y\}$ for all $x, y \in \lambda$. Let $f: [\lambda]^k \rightarrow \mathcal{P}(\lambda)$ be k -generated by $\{\rho_0\}$. It is very easy to check that when $\forall \bar{\gamma} \in [\lambda]^k$, we have $|f(\bar{\gamma})| < \lambda$.

We are also able to prove the case $n = 1$ in ZFC. Set ρ_0 as in the previous case, $\rho_0\{x, y\} = \max\{x, y\}$. It is well known that there is a function $\rho_1: [\lambda]^2 \rightarrow \mu$ such that if $\alpha < \beta < \mu^+ = \lambda$ and $\nu < \mu$, then $|\{\xi \leq \alpha : \rho_1\{\xi, \alpha\} \leq \nu\}| < \mu$ (\clubsuit). Now let $\bar{\gamma} \in [\lambda]^k$. Set $v^* = \max\{\rho_1\{\gamma, \gamma'\} : \gamma, \gamma' \in \bar{\gamma}\}$, $\gamma^* = \max \bar{\gamma}$. Since f is k -generated by $\{\rho_0, \rho_1\}$, we have $f(\bar{\gamma}) \subset \{x \leq \gamma^* : \rho_1\{x, \gamma^*\} \leq v^*\}$. Now (\clubsuit) implies that $|\{x \leq \gamma^* : \rho_1\{x, \gamma^*\} \leq v^*\}| < \mu$, and thus $|f(\bar{\gamma})| < \mu$.

Now assume that the theorem is true for $n \geq 1$. We prove it for $n + 1$. Recall that μ is a regular cardinal, $\mu = \mu^{<\mu}$, $2^\mu = \mu^+, \dots, 2^{\mu^{+n}} = \mu^{+(n+1)} = \lambda$. Obviously $\mu^+ = (\mu^+)^{<(\mu^+)}$, and so by the induction hypothesis there is a notion of forcing $\mathbb{P}_n = \mathbb{P}_n^{\mu^+, \lambda}$ such that \mathbb{P}_n is $(< \mu^+)$ -complete, of cardinality λ , and collapses no cardinal, and in $W = V^{\mathbb{P}_n}$:

(i) there are functions

$$\rho_0: [\lambda]^2 \rightarrow \mu^{+(n+1)}, \rho_1: [\lambda]^2 \rightarrow \mu^{+n}, \dots, \rho_n: [\lambda]^2 \rightarrow \mu^+$$

such that if $f: [\lambda]^k \rightarrow \mathcal{P}(\lambda)$ is k -generated by $\{\rho_0, \dots, \rho_n\}$, then $\forall \bar{\gamma} \in [\lambda]^k$ and we have $|f(\bar{\gamma})| < \mu^+$;

(ii) $2^{\mu^+} = \dots = 2^{\mu^{+n}} = \lambda$.

We define a forcing notion \mathbb{P} in $W = V^{\mathbb{P}_n}$ which is $(< \mu)$ -complete and has the μ^+ -c.c. such that for a generic $G \subset \mathbb{P}$ over W , we can find $\rho_{n+1}: [\lambda]^2 \rightarrow \mu$ in the generic extension $W[G]$ so that if $f: [\lambda]^k \rightarrow \mathcal{P}(\lambda)$ is k -generated by $\{\rho_0, \dots, \rho_n, \rho_{n+1}\}$, then $\forall \bar{\gamma} \in [\lambda]^k$ and we have $|f(\bar{\gamma})| < \mu$. Let (\mathbb{P}, \leq) consist of triples of the form $\langle s, g, \varrho \rangle$ when $s \in [\lambda]^{<\mu}$, $\varrho: [s]^2 \rightarrow \mu$, $g: [s]^k \rightarrow \mathcal{P}(s)$ and $\{\rho_0|_s, \rho_1|_s, \dots, \rho_n|_s, \varrho\}$ k -generates g . For ease of notation, we sometimes denote the condition $p = \langle s, g, \varrho \rangle$ by $p = \langle s_p, g_p, \varrho_p \rangle$.

For $p = \langle s, g, \varrho \rangle$ and $p' = \langle s', g', \varrho' \rangle$ in (\mathbb{P}, \leq) , we say that p' extends p ($p' \leq p$) if and only if $s' \supset s$, $g'|_{[s]^k} = g$ and $\varrho'|_{[s]^2} = \varrho$.

Claim 1 (\mathbb{P}, \leq) is $(< \mu)$ -complete.

Proof of Claim 1 Let η be a cardinal $< \mu$, and let $\langle p_i = \langle s_i, g_i, \varrho_i \rangle; i < \eta \rangle$ be a decreasing sequence of conditions in \mathbb{P} . Set

$$s = \bigcup_{i < \eta} s_i, \quad g = \bigcup_{i < \eta} g_i, \quad \varrho = \bigcup_{i < \eta} \varrho_i.$$

We show that $p = \langle s, g, \varrho \rangle$ is a condition in \mathbb{P} extending all $p_i, i < \eta$. By regularity of μ , we have $|s| < \mu$. Now suppose that $g' : [s]^k \rightarrow \mathcal{P}(s)$ is the function k -generated by $\{\rho_0|_s, \dots, \rho_n|_s, \varrho\}$. We prove that $g = g'$. Let $\bar{\gamma} \in [s]^k$, and also let $i_0 < \eta$ be such that $\bar{\gamma} \in [s_{i_0}]^k$. Lemma 2.4 says that for all $i_0 \leq i < \eta$, $g'(\bar{\gamma}) \cap s_i = g_i(\bar{\gamma})$. Thus

$$\begin{aligned} g'(\bar{\gamma}) &= g'(\bar{\gamma}) \cap s = g'(\bar{\gamma}) \cap \left(\bigcup_{i_0 \leq i < \eta} s_i \right) = \bigcup_{i_0 \leq i < \eta} (g'(\bar{\gamma}) \cap s_i) \\ &= \bigcup_{i_0 \leq i < \eta} g_i(\bar{\gamma}) = g(\bar{\gamma}). \end{aligned}$$

Also by definition of p , it is clear that p extends all $p_i, i < \eta$. This proves Claim 1. \square

Claim 2 *We have that (\mathbb{P}, \leq) has the μ^+ -c.c.*

Proof of Claim 2 Assume that $p_i = \langle s_i, g_i, \varrho_i \rangle \in (\mathbb{P}, \leq)$ for $i < \mu^+$. Using a standard Δ -system argument, we can suppose that $s_i = a \cup b_i$ for the pairwise disjoint sets $a, b_i, i < \mu^+$. Let $f: [\lambda]^k \rightarrow \mathcal{P}(\lambda)$ be k -generated by $\{\rho_0, \dots, \rho_n\}$. By Lemma 2.3, for any $\bar{\gamma} \in [a]^k$, $g_i(\bar{\gamma})$ is a subset of $f(\bar{\gamma})$, and by definition, $g_i(\bar{\gamma})$ is of cardinality $< \mu$. As $|f(\bar{\gamma})| \leq \mu$ (by the induction hypothesis) and $|a| < \mu$, we can assume, by $\mu^{<\mu} = \mu$, that $g_i|_{[a]^k}$ is the same for $i < \mu^+$. Also, considering the fact that $\text{range}(\varrho_i) \subset \mu$ for $i < \mu^+$, we can assume that $\varrho_i|_{[a]^2}$ is the same for $i < \mu^+$.

Now we will show that for any $i, j < \mu^+$, p_i and p_j are compatible. Set $s = s_i \cup s_j = a \cup b_i \cup b_j$. We intend to find $q = \langle s, g, \varrho \rangle \in \mathbb{P}$ such that q extends both p_i, p_j . Define $\varrho: [s]^2 \rightarrow \mu$ by $\varrho \supset \varrho_i, \varrho_j$ and for $\{\alpha_1, \alpha_2\} \in [s]^2 - [s_i]^2 - [s_j]^2$,

$$\varrho\{\alpha_1, \alpha_2\} = \sup(\varrho''_i[s_i]^2 \cup \varrho''_j[s_j]^2) + 1. \quad (1)$$

Note that since μ is regular and $|s_i|, |s_j| < \mu$, the sets $\varrho''_i[s_i]^2, \varrho''_j[s_j]^2$ are bounded in μ , so the above definition is well defined. Now let $g: [s]^k \rightarrow \mathcal{P}(s)$ be the function k -generated by $\{\rho_0|_s, \dots, \rho_n|_s, \varrho\}$. This completes the definition of $q = \langle s, g, \varrho \rangle$. Clearly $q \in \mathbb{P}$. We must show that q extends p_i, p_j . By symmetry, it is enough to show that q extends p_i . As $s_i = a \cup b_i \subset s$ and $\varrho_i \subset \varrho$, this would be done if we could show that $\forall \bar{\gamma} \in [s_i]^k$ $g_i(\bar{\gamma}) = g(\bar{\gamma})$. Lemma 2.3 tells us that $g_i(\bar{\gamma}) \subset g(\bar{\gamma})$. By definition, $g(\bar{\gamma}) \subset s$, $g_i(\bar{\gamma}) \subset s_i$ and from Lemma 2.4 it follows that $g_i(\bar{\gamma}) = g(\bar{\gamma}) \cap s_i$. So it remains to show that $g(\bar{\gamma}) \cap (s_j \setminus s_i) = \emptyset$. By way of contradiction, assume there is $x \in g(\bar{\gamma}) \cap b_j$. Note that if $\bar{\gamma} \in [a]^k$, then $x \in g_j(\bar{\gamma}) = g_i(\bar{\gamma}) \subset s_i$, which implies that $x \in b_j \cap s_i$. This contradicts the disjointness of s_i, b_j . So suppose that there exists $\gamma \in \bar{\gamma}$ such that $\gamma \notin a$. This means that

$$\{x, \gamma\} \in [s]^2 - [s_i]^2 - [s_j]^2.$$

Then by (1), we have $\varrho\{x, \gamma\} > \sup \varrho''_i[s_i]^2$. In other words, for all $\{\gamma', \gamma''\} \in [\bar{\gamma}]^2$ we have $\varrho\{x, \gamma\} > \varrho\{\gamma', \gamma''\}$, implying that $x \notin g(\bar{\gamma})$, which is a contradiction. Thus we have proved that $\forall \bar{\gamma} \in [s_i]^k$ $g_i(\bar{\gamma}) = g(\bar{\gamma})$ and this shows that q extends p_i , which completes the proof of Claim 2. \square

Now let G be a \mathbb{P} -generic filter over W . \square

Claim 3 $\bigcup_{p \in G} s_p = \lambda$.

Proof of Claim 3 Obviously $\bigcup_{p \in G} s_p \subset \lambda$. For any $\alpha \in \lambda$, let $D_\alpha = \{p \in \mathbb{P} : \alpha \in s_p\}$. We will show that D_α is dense in \mathbb{P} . Assume that $q = \langle s, g, \varrho \rangle \in \mathbb{P}$ with $\alpha \notin s$. Let $s' = s \cup \{\alpha\}$. We define $\varrho': [s_1]^2 \rightarrow \mu$ as follows: $\varrho'|_s = \varrho$ and for $\{\alpha, \gamma\} \in [s']^2 - [s]^2$, set $\varrho'\{\alpha, \gamma\} = \sup(\varrho''[s]^2) + 1$. By regularity of μ and $|s| < \mu$, this is well defined. Now let $g': [s']^k \rightarrow \mathcal{P}(s')$ be the function k -generated by $\{\rho_0|_{s'}, \dots, \rho_n|_{s'}, \varrho'\}$. So $p = \langle s', g', \varrho' \rangle \in \mathbb{P} \cap D_\alpha$. We show that $p \leq q$. Obviously $s \subset s', \varrho \subset \varrho'$. Suppose that $\bar{\gamma} \in [s]^k$; we must prove that $g(\bar{\gamma}) = g'(\bar{\gamma})$. By Lemmas 2.3 and 2.4, we have $g(\bar{\gamma}) \subset g'(\bar{\gamma})$, $g(\bar{\gamma}) = g'(\bar{\gamma}) \cap s$. Thus it remains to show that $\alpha \notin g'(\bar{\gamma})$. This is so because for each $\gamma \in \bar{\gamma}$ and each $\{\gamma', \gamma''\} \in [\bar{\gamma}]^2$, the definition of ϱ' implies that $\varrho'\{\alpha, \gamma\} > \varrho'\{\gamma', \gamma''\}$. This proves the density of

D_α , from which it follows that $\exists r \in D_\alpha \cap G$. So $\alpha \in s_r \subset \bigcup_{p \in G} s_p$. Therefore $\bigcup_{p \in G} s_p = \lambda$, and Claim 3 is proved.

Now we are ready to introduce $\rho_{n+1}: [\lambda]^2 \rightarrow \mu$ and $f: [\lambda]^k \rightarrow \mathcal{P}(\lambda)$. Set

$$\rho_{n+1} = \bigcup_{p \in G} \rho_p, \quad f = \bigcup_{p \in G} g_p.$$

We show that

- (1) $\forall \bar{\gamma} \in [\lambda]^k \ |f(\bar{\gamma})| < \mu$,
- (2) f is k -generated by $\{\rho_0, \dots, \rho_{n+1}\}$.

Let $\bar{\gamma} \in [\lambda]^k$ and $p \in G$ be such that $\bar{\gamma} \in [s_p]^k$. Then $f(\bar{\gamma}) = g_p(\bar{\gamma}) \subset s_p \in [\lambda]^{<\mu}$. This proves (1). Now let $f': [\lambda]^k \rightarrow \mathcal{P}(\lambda)$ be the function k -generated by $\{\rho_0, \dots, \rho_{n+1}\}$. Let $\bar{\gamma} \in [\lambda]^k$ and $p \in G$ be such that $\bar{\gamma} \in [s_p]^k$. By Lemma 2.4, we have $f'(\bar{\gamma}) \cap s_p = g_p(\bar{\gamma})$. But since G is a filter, for every such $p, q \in G$, $g_p(\bar{\gamma}) = g_q(\bar{\gamma})$. Putting this together with $\bigcup_{p \in G} s_p = \lambda$, we can deduce (2). In other words,

$$\begin{aligned} f'(\bar{\gamma}) &= f'(\bar{\gamma}) \cap \lambda = f'(\bar{\gamma}) \cap \bigcup_{p \in G} s_p = \bigcup_{p \in G} (f'(\bar{\gamma}) \cap s_p) \\ &= \bigcup_{p \in G} g_p(\bar{\gamma}) = f(\bar{\gamma}). \end{aligned}$$

Now let $\dot{\mathbb{P}}$ be a \mathbb{P}_n -name for \mathbb{P} . Then $\mathbb{P}_n * \dot{\mathbb{P}}$ is ($< \mu$)-complete, collapses no cardinal, and has cardinality λ (recall the definition of \mathbb{P} in $V^{\mathbb{P}_n}$, and note that $V^{\mathbb{P}_n} \models \lambda^{<\lambda} = \lambda$), and by what we have done so far, $V^{\mathbb{P}_n * \dot{\mathbb{P}}}$ satisfies requirement (i) of the theorem for $n + 1$. Now observe that for $l = 1, \dots, n$, we have $V^{\mathbb{P}_n} \models (\lambda)^{\mu^{+l}} = (2^{\mu^+})^{\mu^{+l}} = 2^{\mu^+} = \lambda$. Since \mathbb{P} has the μ^+ -c.c. in $V^{\mathbb{P}_n}$, it is well known that we have $V^{\mathbb{P}_n * \dot{\mathbb{P}}} \models (2)^{\mu^{+l}} = \lambda$ (\heartsuit_1). This implies that for $k = 0, \dots, n$, $V^{\mathbb{P}_n * \dot{\mathbb{P}}} \models (\lambda)^{\mu^{+k}} = (2^{\mu^+})^{\mu^{+k}} = \lambda$ (\heartsuit_2). Also, by ($< \mu$)-completeness of \mathbb{P} in $V^{\mathbb{P}_n}$, we have $V^{\mathbb{P}_n * \dot{\mathbb{P}}} \models 2^{<\mu} = \mu$ (\heartsuit_3). Assume that \mathbb{Q} is the poset of all functions $p: \lambda \times \mu \rightarrow 2$ in $V^{\mathbb{P}_n * \dot{\mathbb{P}}}$ such that $|p| < \mu$. Let $\dot{\mathbb{Q}}$ be a $(\mathbb{P}_n * \dot{\mathbb{P}})$ -name for \mathbb{Q} . It is well known that from \heartsuit_2 for $k = 0$ together with \heartsuit_3 , we can deduce $V^{\mathbb{P}_n * \dot{\mathbb{P}} * \dot{\mathbb{Q}}} \models 2^\mu = \lambda$. Since \mathbb{Q} has the μ^+ -c.c. in $V^{\mathbb{P}_n * \dot{\mathbb{P}}}$, from \heartsuit_2 it follows that $V^{\mathbb{P}_n * \dot{\mathbb{P}} * \dot{\mathbb{Q}}} \models (2)^{\mu^{+l}} = \lambda$ for $l = 1, \dots, n$ (as in the case of \heartsuit_1). Now we define $\mathbb{P}_{n+1} = \mathbb{P}_{n+1}^{\mu, \lambda} = \mathbb{P}_n^{\mu^+, \lambda} * \dot{\mathbb{P}} * \dot{\mathbb{Q}}$. Surely \mathbb{P}_{n+1} is ($< \mu$)-complete, collapses no cardinal, has cardinality λ , and satisfies requirements (i) and (ii) of the theorem for $n + 1$. So we have finished the proof. \square

It is interesting to note that the referee had suggested an alternate exposition of the proof of Theorem 2.5. In fact, the proof of Theorem 2.5 can be split into two steps, first proving that if $(\mu = \mu^{<\mu})$, $2 \leq k < \omega$, $F: [\lambda]^k \rightarrow [\lambda] \leq \mu$, then a ($< \mu$)-complete cardinal-preserving forcing adds a function $h^*: [\lambda]^2 \rightarrow \mu$ such that if for $x \in [\lambda]^k$

$$F^*(x) = \{\xi \in F(x): h^*(\xi, \gamma) \leq h^*(x)(\gamma \in x)\},$$

then $|F^*(x)| < \mu$. Here $h^*(x) = \max\{h^*(y): y \in [x]^2\}$. We also define a forcing notion (\mathcal{Q}, \leq) as follows. The elements are tuples (s, h) such that $s \in [\lambda]^{<\mu}$, $h: [s]^2 \rightarrow \mu$. Also, $(s', h') \leq (s, h)$ if $s' \supset s$, $h = h' \upharpoonright [s]^2$, and there are no $x \in [s]^k$, $\xi \in (F(x) \cap s') - s$ such that for each $\gamma \in x$, $h'(\xi, \gamma) \leq h(x)$. Then

we argue as in the proof of Theorem 2.5. Finally, Theorem 2.5 can be obtained by iteration. We leave it to the reader to reproduce the proof via this approach.

The following definition is useful in presenting the proof of the next theorem.

Definition 2.6 Assume that A is a set with $|A| > 3$, that $\langle L, < \rangle$ is a linear order, and that $\rho: [A]^2 \rightarrow L$. Let $B \subset A$, $x \in A \setminus B$. We say that x is ρ -close to B if

$$(\forall \gamma \in B)(\exists \gamma', \gamma'' \in B) \quad [(\gamma' \neq \gamma'') \wedge \rho\{x, \gamma\} \leq \rho\{\gamma', \gamma''\}].$$

Theorem 2.7 Assume that $n < \omega$, $k = 4$, and $\lambda = \mu^{+n}$ for some regular cardinal μ and that s_n is the least number satisfying the Ramsey relation $s_n \rightarrow (5)_{3^{n+1}}^3$. Let $\mathbb{P}_n = \mathbb{P}_n^{\mu, \lambda}$ be as in Theorem 2.5. Then in $V^{\mathbb{P}_n}$ there is a set mapping $F: [\lambda]^4 \rightarrow [\lambda]^{<\mu}$ such that F has no free set of cardinality s_n .

Proof By Theorem 2.5, in $V^{\mathbb{P}_n}$ there are functions $F: [\lambda]^4 \rightarrow [\lambda]^{<\mu}$ and

$$\rho_0: [\lambda]^2 \rightarrow \mu^{+n}, \dots, \rho_n: [\lambda]^2 \rightarrow \mu$$

such that F is 4-generated by $\Gamma = \{\rho_0, \dots, \rho_n\}$. We will show that F has no free set of cardinality s_n . By way of contradiction, suppose not. Let $A = \{\gamma_1, \dots, \gamma_{s_n}\}$ be a free set for F with s_n elements. We are going to define a partition relation \sim on $[A]^3$. Let $\rho \in \Gamma$, $\{\alpha, \beta, \gamma\} \in [A]^3$ with $\alpha < \beta < \gamma$. At least one of the following three possibilities will occur:

- (1) $\max\{\rho\{\alpha, \beta\}, \rho\{\beta, \gamma\}\} \leq \rho\{\alpha, \gamma\}$,
- (2) $\max\{\rho\{\alpha, \beta\}, \rho\{\alpha, \gamma\}\} \leq \rho\{\beta, \gamma\}$,
- (3) $\max\{\rho\{\beta, \gamma\}, \rho\{\alpha, \gamma\}\} \leq \rho\{\alpha, \beta\}$.

We say that $\{\alpha, \beta, \gamma\}$ has ρ -type l ($l = 1, 2, 3$) if the possibility (l) occurs, and l is minimal. Now for $\bar{\gamma}_1, \bar{\gamma}_2 \in [A]^3$, we put $\bar{\gamma}_1 \sim \bar{\gamma}_2$ if and only if $\forall \rho \in \Gamma$, $\bar{\gamma}_1, \bar{\gamma}_2$ have the same ρ -type. The number of equivalence classes is at most 3^{n+1} , so from $|A| = s_n$, $s_n \rightarrow (5)_{3^{n+1}}^3$ it follows that there is a 5-element homogenous set $B = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5\}$. Assume that $\alpha_1 < \alpha_2 < \alpha_3 < \alpha_4 < \alpha_5$.

Claim For every $\rho \in \Gamma$, α_3 is ρ -close to $\{\alpha_1, \alpha_2, \alpha_4, \alpha_5\}$.

Proof of Claim Let $\rho \in \Gamma$. There are three cases to deal with.

Case 1: For all $\bar{\gamma} \in [B]^3$, $\bar{\gamma}$ has ρ -type 1. In this case, we have the following relations:

$$\begin{aligned} \rho\{\alpha_3, \alpha_1\} &\leq \rho\{\alpha_1, \alpha_4\}, & \rho\{\alpha_3, \alpha_2\} &\leq \rho\{\alpha_2, \alpha_4\}, \\ \rho\{\alpha_3, \alpha_4\} &\leq \rho\{\alpha_4, \alpha_1\}, & \rho\{\alpha_3, \alpha_5\} &\leq \rho\{\alpha_5, \alpha_1\}. \end{aligned}$$

Case 2: For all $\bar{\gamma} \in [B]^3$, $\bar{\gamma}$ has ρ -type 2. In this case, we have the following relations:

$$\begin{aligned} \rho\{\alpha_3, \alpha_1\} &\leq \rho\{\alpha_4, \alpha_5\}, & \rho\{\alpha_3, \alpha_2\} &\leq \rho\{\alpha_4, \alpha_5\}, \\ \rho\{\alpha_3, \alpha_4\} &\leq \rho\{\alpha_4, \alpha_5\}, & \rho\{\alpha_3, \alpha_5\} &\leq \rho\{\alpha_4, \alpha_5\}. \end{aligned}$$

Case 3: For all $\bar{\gamma} \in [B]^3$, $\bar{\gamma}$ has ρ -type 3. In this case, we have the following relations:

$$\begin{aligned} \rho\{\alpha_3, \alpha_1\} &\leq \rho\{\alpha_1, \alpha_2\}, & \rho\{\alpha_3, \alpha_2\} &\leq \rho\{\alpha_1, \alpha_2\}, \\ \rho\{\alpha_3, \alpha_4\} &\leq \rho\{\alpha_1, \alpha_2\}, & \rho\{\alpha_3, \alpha_5\} &\leq \rho\{\alpha_1, \alpha_2\}. \end{aligned}$$

Therefore, in all cases, we have shown that α_3 is ρ -close to $\{\alpha_1, \alpha_2, \alpha_4, \alpha_5\}$. This proves the claim. \square

Now from the claim it follows that $\alpha_3 \in F\{\alpha_1, \alpha_2, \alpha_4, \alpha_5\}$, which violates the freeness of A . This is a contradiction. \square

3 Some Ramsey Considerations

In this section we show that when n tends to infinity, s_n gives us a better bound than t_n . First, we fix our notation for representing Ramsey numbers:

$$\begin{aligned} R_k(l_1, \dots, l_r) &= \min\{n_0 : \text{for } n \geq n_0 \ n \rightarrow (l_1, \dots, l_r)^k\}, \\ R_k(l; r) &= \min\{n_0 : \text{for } n \geq n_0 \ n \rightarrow (l)_r^k\}, \\ R_k(l) &= \min\{n_0 : \text{for } n \geq n_0 \ n \rightarrow (l)^k\}. \end{aligned}$$

So in terms of the above notation, we have $s_n = R_3(5; 3^{n+1})$ and $t_{n+1} = R_5(t_n, 7)$. Now we state Erdős and Rado's upper bound for $R_k(l; r)$. For this we need to define a binary operation $*$ on positive integers as follows:

$$a * b = a^b.$$

Also, for $n \geq 3$, we put

$$a_1 * a_2 * \dots * a_n = a_1 * (a_2 * (\dots * (a_{n-1} * a_n) \dots)).$$

Then if $1 \leq m < n$, we have

$$a_1 * a_2 * \dots * a_m * (a_{m+1} * \dots * a_n) = a_1 * a_2 * \dots * a_n.$$

Erdős and Rado [2] proved the following.

Theorem 3.1 (Erdős and Rado) For $r \geq 2$ and $l \geq k \geq 2$, we have

$$R_k(l; r) \leq r * r^{k-1} * r^{k-2} * \dots * r^2 * [r(l-k) + 1].$$

Therefore,

$$s_n \leq 3^{n+1} * 3^{2n+1} * (2 \cdot 3^{n+1} + 1). \quad (2)$$

On the other hand, by using Erdős and Hajnal's stepping-up lemma (see Graham, Rothschild, and Spencer [5]), we get a lower bound for t_n . In fact, what we want is an off-diagonal version of the stepping-up lemma which can be obtained by easily modifying the proof of the original version mentioned in [5]. Thus we have the following.

Theorem 3.2 (Off-diagonal stepping-up lemma) Suppose that $k \geq 3$ and $n \rightarrow (l_1, l_2)^k$. Then $2^n \rightarrow (2l_1 + k - 4, 2l_2 + k - 4)^{k+1}$.

By a simple coloring argument, we can show that $R_3(l, 4) > 2l$. Then the stepping-up lemma implies that $R_5(2l - 1, 7) > 2^{2l}$. So we have $t_{n+1} = R_5(t_n, 7) > 2^{t_n}$. Hence, for $n > 1$,

$$t_n > \text{Tower}_n(7), \quad (3)$$

where $\text{Tower}_1(x) = x$, $\text{Tower}_{n+1}(x) = 2^{\text{Tower}_n(x)}$. Now an easy computation through (2) and (3) would show that t_n exceeds s_n when n is large enough.

4 More on Set Mappings on 4-Tuples

In this section we deal with a set mappings $f: [\lambda]^4 \rightarrow [\lambda]^{<\mu}$ with the additional property

$$\forall x_0 < x_1 < x_2 < x_3 \in \lambda \ [f\{x_0, x_1, x_2, x_3\} \cap (x_1, x_2) = \emptyset]. \quad (4)$$

As mentioned in the Introduction, the above property excludes both the set mappings in Section 2 and in Komjath and Shelah's work. By slightly modifying Hajnal's proof for Theorem B (see the Introduction), at some points, we show the following result.

Theorem 4.1 *Let $\lambda \geq \mu^{+3}$, and assume that the set mapping $f: [\lambda]^4 \rightarrow [\lambda]^{<\mu}$ satisfies (4). Then f has arbitrary large finite free sets.*

Proof We assume that the reader is familiar with Hajnal's proof from [1, Section 46] and for convenience follow the same terminology here. We construct the sets $A_m = \{x_{m\xi} : \xi < \mu^+\}$ ($m < \omega$), where $x_{m\xi}$ are pairwise distinct elements of λ , and a sequence $H_0 \supset H_1 \supset \dots \supset H_m \supset \dots$ ($m \in \omega$) with the following properties:

- (i) $|H_m| = \lambda$, $\{x_{m\xi} : \xi < \mu^+\} \subset H_m$,
- (ii) $A_m < A_{m+1}$,
- (iii) $u \notin f\{x_{l_1\xi_1}, x_{l_2\xi_2}, x_{m\eta}, v\}$ holds whenever $l_1 < l_2 < m < \omega$ and $\xi_1, \xi_2, \eta < \mu^+$ and $u, v \in H_{m+1}$.

This can be done exactly as in Hajnal's proof. The extra condition (ii) can easily be fixed by putting A_m as the set of the first μ^+ elements of H_m . After doing this, we construct an n -element set $\{x_i : i < n\}$ that is free with respect to f such that $x_i \in A_i$. Simultaneously with the construction of this set, we also construct sequences of sets $E_j^i \subset A_j$ for $i \leq n-3$ and $j \leq i$, and we would have $x_i \in E_j^i$ for $i \leq n-3$. For $i = n-1, n-2, n-3$ pick $x_i \in A_i$ arbitrarily, and write $E_j^{n-3} = A_j$ for $j \leq n-3$. Given $m < n-3$, if x_i and E_j^i has already been defined in the case $m < i < n$, $j \leq i$, then define the set mapping h_m on $E^{m+1} = \bigcup_{j \leq m} E_j^{m+1}$ by putting

$$h_m(u) = \bigcup_{m+1 < i_1 < i_2 < n} f\{u, x_{m+1}, x_{i_1}, x_{i_2}\}.$$

Clearly $|h_m(u)| < \mu$, and so by using a lemma quoted in Hajnal's proof from [1, Section 46], there is a set X_m free with respect to h such that $X_m \cup E_j^{m+1}$ has cardinality μ for each $j \leq m$. Put

$$E_j^m = X_m \cup E_j^{m+1}$$

for $j \leq m$. Pick an arbitrary element of E_m^m as x_m . This finishes the construction. Now we see that this set is free with respect to f . By (ii), we have $x_0 < x_1 < \dots < x_{n-1}$. Let

$$\{x_{i_0}, x_{i_1}, x_{i_2}, x_{i_3}, x_{i_4}\} \subset \{x_i : i < n\},$$

where $i_0 < i_1 < i_2 < i_3 < i_4$. Since $x_{i_3}, x_{i_4} \in H_{i_2+1}$, it follows from (i) and (iii) that

$$x_{i_3} \notin f\{x_{i_0}, x_{i_1}, x_{i_2}, x_{i_4}\}, \quad x_{i_4} \notin f\{x_{i_0}, x_{i_1}, x_{i_2}, x_{i_3}\}.$$

Moreover, we have

$$x_{i_0} \notin f\{x_{i_1}, x_{i_2}, x_{i_3}, x_{i_4}\}, \quad x_{i_1} \notin f\{x_{i_0}, x_{i_2}, x_{i_3}, x_{i_4}\},$$

since $x_{i_1}, x_{i_2} \in E^{i_2-1}$ and this latter is free with respect to h_{i_2} . Finally,

$$x_{i_2} \notin f\{x_{i_0}, x_{i_1}, x_{i_3}, x_{i_4}\}$$

follows from $x_{i_0} < x_{i_1} < x_{i_2} < x_{i_3} < x_{i_4}$ and condition (4) imposed on f . Now the proof is complete. \square

5 Set Mappings on Triples

Komjath and Shelah's second construction in [9] implies that for infinite cardinals λ, μ , where μ is regular, $\lambda = \mu^{+n}$, $n \in \omega$, and GCH holds for μ^{+l} ($l < n$), there is a cardinal-preserving generic extension in which there exists a set mapping $f: [\lambda]^3 \rightarrow [\lambda]^{<\mu}$ with no free set of size ω . The set mapping in question essentially has the following property:

$$\forall x_0, x_1, x_2 \in \lambda \quad [x_0 < x_1 < x_2 \rightarrow f\{x_0, x_1, x_2\} \subset x_0].$$

Also, it is easy to see that the proofs of Theorems 4.1 and 4.2 in Hajnal and Máté's paper [7] for set mappings on pairs can be adapted for set mappings on triples. We then have the following theorem.

Theorem 5.1 *Let μ, λ be infinite cardinals, and consider the set mapping $f: [\lambda]^3 \rightarrow [\lambda]^{<\mu}$.*

- (i) *If λ is regular, $\lambda = \mu$, and for all $x_0 < x_1 < x_2 \in \lambda$, $f\{x_0, x_1, x_2\} \subset (x_2, +\infty)$, then f has a free set of cardinality λ .*
- (ii) *If $\lambda = \mu^{+n}$ ($n \geq 1$) and for all $x_0 < x_1 < x_2 \in \lambda$, $f\{x_0, x_1, x_2\} \subset (x_1, x_2)$, then f has a free set of size ω .*

It is interesting to note that, in [8], Komjath proved several results of these kinds for a general set mapping on k -tuples.

Now it remains to see what happens if a set mapping $f: [\lambda]^3 \rightarrow [\lambda]^{<\mu}$ satisfies the following property:

$$\forall x_0 < x_1 < x_2 \in \lambda \quad [f\{x_0, x_1, x_2\} \subset (x_0, x_1)].$$

The following theorem can be proved by an adaptation of Komjath and Shelah's second construction. Here we show that our forcing construction in Section 2 will enable us to give another proof for this case. In fact, we show that it is consistent that the above f has no free set of size ω . In precise terms, we have the following.

Theorem 5.2 *Assume that $n < \omega, k = 3$, and $\lambda = \mu^{+n}$ for some regular cardinal μ . Let $\mathbb{P}_n = \mathbb{P}_n^{\mu, \lambda}$ be as in Theorem 2.5. Then in $V^{\mathbb{P}_n}$ there is a set mapping $F: [\lambda]^3 \rightarrow [\lambda]^{<\mu}$ such that F has no infinite free set and for all $x_0 < x_1 < x_2 \in \lambda$ we have $F\{x_0, x_1, x_2\} \subset (x_0, x_1)$.*

Proof By Theorem 2.5, in $V^{\mathbb{P}_n}$ there are functions $F: [\lambda]^3 \rightarrow [\lambda]^{<\mu}$ and

$$\rho_0: [\lambda]^2 \rightarrow \mu^{+n}, \dots, \rho_n: [\lambda]^2 \rightarrow \mu$$

such that F is 3-generated by $\Gamma = \{\rho_0, \dots, \rho_n\}$. We show that F has no infinite free set. By way of contradiction, suppose not. Let $A \subset \lambda$, $|A| = \omega$, be a free set for F . For every $\{\alpha, \beta, \gamma\} \in [A]^3$ with $\alpha < \beta < \gamma$ and every $\rho \in \{\rho_0, \dots, \rho_n\}$, we have two possibilities:

$$(Ia) \quad \rho\{\alpha, \beta\} \leq \rho\{\beta, \gamma\}, \quad (Ib) \quad \rho\{\alpha, \beta\} > \rho\{\beta, \gamma\}.$$

According to which one of the above possibilities occurs, we say that the 4-tuple $\langle \rho; \alpha, \beta, \gamma \rangle$ satisfies that possibility. Now we define an equivalence relation on $[A]^3$. For $\alpha < \beta < \gamma, \alpha' < \beta' < \gamma'$ in A , we put $\{\alpha\beta\gamma\} \sim \{\alpha'\beta'\gamma'\}$ when for each $\rho \in \Gamma$, either both of $\langle \rho; \alpha, \beta, \gamma \rangle, \langle \rho; \alpha', \beta', \gamma' \rangle$ satisfy (Ia) or both of them satisfy (Ib). By the Ramsey theorem $\omega \rightarrow (\omega)^3$, there exists an infinite subset

$$B = \{\beta_0, \beta_1, \beta_2 \dots\} \subset A, \quad \beta_0 < \beta_1 < \beta_2 < \dots$$

such that either all of $\langle \rho; \alpha, \beta, \gamma \rangle$ in question satisfy (Ia) or all of them satisfy (Ib). We claim this cannot be (Ib), since in this case we would have

$$\rho\{\beta_0, \beta_1\} > \rho\{\beta_1, \beta_2\} > \dots,$$

which is an infinite sequence of strictly decreasing ordinals.

Again, for every $\{\alpha, \beta, \gamma\} \in [B]^3$ with $\alpha < \beta < \gamma$ and every $\rho \in \{\rho_0, \dots, \rho_n\}$, we have two possibilities:

$$(IIa) \quad \rho\{\alpha, \beta\} \leq \rho\{\alpha, \gamma\}, \quad (IIb) \quad \rho\{\alpha, \beta\} > \rho\{\alpha, \gamma\}.$$

We define an equivalence relation on $[B]^3$. For $\alpha < \beta < \gamma, \alpha' < \beta' < \gamma'$ in B , we put $\{\alpha\beta\gamma\} \sim \{\alpha'\beta'\gamma'\}$, when, for each $\rho \in \Gamma$, either both of $\langle \rho; \alpha, \beta, \gamma \rangle, \langle \rho; \alpha', \beta', \gamma' \rangle$ satisfy (IIa) or both of them satisfy (IIb). As in the previous case by the Ramsey theorem $\omega \rightarrow (\omega)^3$, we deduce that there is an infinite subset

$$C = \{\gamma_0, \gamma_1, \gamma_2 \dots\} \subset B, \quad \gamma_0 < \gamma_1 < \gamma_2 < \dots$$

such that either all of $\langle \rho; \alpha, \beta, \gamma \rangle$ in question satisfy (IIa) or all of them satisfy (IIb). This cannot be (IIb), since we would have the following infinite sequence of strictly decreasing ordinals:

$$\rho\{\gamma_0, \gamma_1\} > \rho\{\gamma_1, \gamma_2\} > \dots.$$

So far we have obtained that for every $\{\alpha\beta\gamma\} \in [C]^3$ and every $\rho \in \Gamma$, both of the following two items occur:

$$(Ia) \quad \rho\{\alpha, \beta\} \leq \rho\{\beta, \gamma\}, \quad (IIa) \quad \rho\{\alpha, \beta\} \leq \rho\{\alpha, \gamma\}.$$

Now, for every $\{\alpha, \beta, \gamma\} \in [C]^3$ with $\alpha < \beta < \gamma$ and every $\rho \in \{\rho_0, \dots, \rho_n\}$, we have two possibilities:

$$(IIIa) \quad \rho\{\beta, \gamma\} > \rho\{\alpha, \gamma\}, \quad (IIIb) \quad \rho\{\beta, \gamma\} \leq \rho\{\alpha, \gamma\}.$$

As in the two previous cases, we define an equivalence relation on $[C]^3$. For $\alpha < \beta < \gamma, \alpha' < \beta' < \gamma'$ in C , we put $\{\alpha\beta\gamma\} \sim \{\alpha'\beta'\gamma'\}$ when for each $\rho \in \Gamma$, both of $\langle \rho; \alpha, \beta, \gamma \rangle$ and $\langle \rho; \alpha', \beta', \gamma' \rangle$ satisfy the same possibility. Using the Ramsey theorem $\omega \rightarrow (\omega)^3$ for the third time, we obtain an infinite homogenous subset $D \subset C$. Now let $\{\alpha, \beta, \gamma, \gamma'\} \subset D, \alpha < \beta < \gamma < \gamma'$. We show that for every $\rho \in \Gamma$, β is ρ -close to $\{\alpha, \gamma, \gamma'\}$. Fix an arbitrary $\rho \in \Gamma$. By (Ia) we have $\rho\{\beta, \gamma\} \leq \rho\{\gamma, \gamma'\}$. Observe that if (IIIa) occurs, then $\rho\{\beta, \gamma'\} < \rho\{\gamma, \gamma'\}$, and if (IIIb) occurs, then $\rho\{\beta, \gamma'\} \leq \rho\{\alpha, \gamma'\}$. This shows that in both cases of (IIIa) and (IIIb), β is ρ -close to $\{\alpha, \gamma, \gamma'\}$. Since ρ was arbitrary, we conclude that $\beta \in F\{\alpha, \gamma, \gamma'\}$, which violates the freeness of A , a contradiction. \square

References

- [1] Erdős, P., A. Hajnal, A. Máté, and R. Rado, *Combinatorial Set Theory: Partition Relations for Cardinals*, vol. 106 of *Studies in Logic and the Foundations of Mathematics*, North-Holland, Amsterdam, 1984. [Zbl 0573.03019](#). [MR 0795592](#). [405](#), [406](#), [413](#)
- [2] Erdős, P., and R. Rado, “Combinatorial theorems on classifications of subsets of a given set,” *Proceedings of the London Mathematical Society, Third Series*, vol. 3 (1952), pp. 417–39. [Zbl 0048.28203](#). [MR 0065615](#). [DOI 10.1112/plms/s3-2.1.417](#). [412](#)
- [3] Gillibert, P., “Points critiques de couples de varietes d’algebras,” Ph.D. dissertation, University of Caen, Normandy, France, 2008. [406](#)
- [4] Gillibert, P., and F. Wehrung, “An infinite combinatorial statement with a poset parameter,” *Combinatorica*, vol. 31 (2011), pp. 183–200. [Zbl 1274.03065](#). [MR 2848250](#). [DOI 10.1007/s00493-011-2602-y](#). [406](#)
- [5] Graham, R. L., B. L. Rothschild, and J. H. Spencer, *Ramsey Theory*, second edition, Wiley, New York, 1980. [Zbl 0455.05002](#). [MR 0591457](#). [412](#)
- [6] Hajnal, A., “Proof of a conjecture of S. Ruziewicz,” *Fundamenta Mathematicae*, vol. 50 (1961), pp. 123–28. [Zbl 0100.28003](#). [MR 0131986](#). [405](#)
- [7] Hajnal, A., and A. Máté, “Set mappings, partitions, and chromatic numbers,” pp. 347–79 in *Logic Colloquium ’73 (Bristol, 1973)*, vol. 80 of *Studies in Logic and the Foundations of Mathematics*, North-Holland, Amsterdam, 1975. [Zbl 0324.04004](#). [MR 0424569](#). [405](#), [406](#), [414](#)
- [8] Komjath, P., “A note on set mappings,” in preparation. [414](#)
- [9] Komjath, P., and S. Shelah, “Two consistency results on set mappings,” *Journal of Symbolic Logic*, vol. 65 (2000), pp. 333–38. [Zbl 0948.03048](#). [MR 1782123](#). [DOI 10.2307/2586540](#). [405](#), [406](#), [407](#), [414](#)

Acknowledgments

We would like to thank Peter Komjath for letting us know about his paper as well as those of Gillibert and Wehrung. We would also like to thank the referee for his interesting comment.

Mohsenipour
 School of Mathematics
 Institute for Research in Fundamental Sciences (IPM)
 Tehran
 Iran
sh.mohsenipour@gmail.com

Shelah
 Einstein Institute of Mathematics
 Hebrew University of Jerusalem
 Jerusalem
 Israel
 and
 Department of Mathematics
 Rutgers University
 New Brunswick
 New Jersey
 USA
shelah@math.huji.ac.il