

## THE LAZY MODEL-THEORETICIAN'S GUIDE TO STABILITY

by

Saharon SHELAH \*

The Hebrew University of Jerusalem  
Université Catholique de Louvain

## § 0. INTRODUCTION

The main aim of this article is to make propaganda for [S 1]; hence novelty is not the intention; there are many explanations, definitions and theorems and few proofs. No previous knowledge of stability is required, but then you have to take some statements on faith. We have in mind mainly those who are interested in algebraically-minded model theory, i.e. in generic models, the class of e-closed (= existentially closed) models and universal-homogeneous models rather than elementary classes and saturated models. So our main point is that though stability theory was developed for the latter context, almost everything goes through in the wider context (with suitable changes in the definitions). We also examine some specific algebraic theories (\*). An exact list of the theorems from [S 1] which generalize or do not generalize, will be included in the Ph. D. thesis of M. Abramsky.

At this stage I will define stability theory as an attempt to give a classification of, and structure/non-structure theorems for elementary classes, and other related classes. An ideal *structure theorem* is a characterization (up to isomorphism) of

(\*) I would like to thank Paul HENRARD for his hospitality and for his endless efforts to organize the conference. I thank also MAKOWSKI, WILKIE, HODGES, CHERLIN and MAKOWSKI again, for writing notes on my lectures, on which this paper is based.

This research has been partially supported by the Israel Academy of Sciences.

(<sup>1</sup>) It seems that we get new information on the number of modules over a fixed ring (see 8.7).

each model in the class, by invariants, which are cardinals or sets of cardinals etc. (e.g. the dimension for algebraically closed fields, or the Ulm theorem for countable torsion groups (see [Fu]), or the theory of one equivalence relation (a model  $M = (|M|, E)$  is characterized by the function  $f, f(\lambda) = |\{a/E : |a/E| = \lambda\}|$ ).

An ideal *structure/non-structure theorem* is the characterization of the classes which have a structure theorem, together with a proof of the complexity of the other classes. More explicitly the problem presents itself as determining the possible functions  $I(\lambda, K) =$  the number of models  $M \in K$ , of cardinality  $\lambda$  up to isomorphism. We have now a complete solution only for the family of the classes of  $\aleph_\alpha$ -saturated (\*) models of a complete theory (the theorem essentially says that any  $T$  which does not satisfy  $I(\lambda, T) = 2^\lambda$ , is like some  $T_\alpha$ ;  $T_\alpha$  has equivalence relations  $E_i (i < \alpha)$  where  $E_i$  refines  $E_j$  when  $i < j$ ).

Notice that the question may degenerate when  $2^{\aleph_\alpha} = 2^{|\alpha|} + \aleph_\alpha$ , which may hold for every  $\aleph_\alpha$ .

We consider a classification meaningful if (but not only if) it helps in proving theorems not mentioning it, or has applications or appears in a characterization. For example  $\aleph_\alpha$ -stability helps in proving Morley's categoricity theorem and also in dealing with the differential closure of a differential field of characteristic zero. Stability helps to investigate Keisler's order ([K 1], [S 4]), to prove the uniqueness of prime differential closure of fields (8.7), to characterize  $\{\lambda : K \text{ has a homogeneous-universal [saturated] model in } \lambda\}$  and so forth.

Notice, that it is usually not hard to find properties implying complexity or simplicity; the point is to have both. For example  $\aleph_\alpha$ -stability is a very strong property; but there are theo-

(\*) This notion is slightly stronger than  $\aleph_\alpha$ -saturated. It means that every type almost over a finite set is realized, see § 6.

ries  $T$  which are not  $\aleph_\alpha$ -stable, but  $I(\lambda, T) \leq 2^{2^\lambda}$ , and have a structure theorem (the example is due to Morley). Hence we use superstability or stability more than  $\aleph_\alpha$ -stability. When we deal with  $|T|^{+}$ -saturated models stability is more interesting; but for elementary classes superstability is better. By § 8 we can have a structure theorem for the class of  $R$ -modules, iff the theory of the class is superstable; but if we assume  $|R|^{+}$ -saturation, or compactness of appropriate kind, we can have it in general, as all modules are stable.

A *non-structure theorem* is a little vague; specifically we are proving  $I(\lambda, K) = 2^\lambda$ . However this does not *a priori* exclude the existence of a complete set of invariants. The main non-structure theorem of [S 1] is for  $K = \text{Mod } T$  when  $T$  is not superstable and it is presented in [S 3]. However from this proof it follows that we should use invariants of the form  $A/D_\lambda$  (where  $\lambda$  is regular,  $A \subseteq \lambda, D_\lambda$  the filter on  $\lambda$  generated by the closed unbounded subsets of  $\lambda$ ) and even this is not sufficient (e.g. when  $V = L$ ) (explained in [S 2] for separable reduced  $p$ -groups).

Eklof [Ek] devised a more rigorous version of the thesis that there is no structure theorem here, by Feferman-Sabbagh generalized  $\kappa$ -functors. Note that in general, the  $L_{\infty, \lambda}$ -theory of models (of cardinality  $\lambda$ ) does not characterize them up to isomorphism when  $T$  is unsuperstable,  $\lambda > \aleph_\alpha$  is regular (\*); and if  $\mu < \kappa(T)$  ( $\mu$  regular) also  $L_{\infty, \lambda, G(\mu)}$  is not sufficient ( $G(\mu)$  is the game quantifier  $(\dots \forall x_\alpha \exists y_\alpha \dots)_{\alpha < \mu}$ ).

We can apply the non-structure theorem itself; but usually we have to check directly that the appropriate construction works. This is done in [S 2] for reduced separable  $p$ -groups and in [MS] for universal locally finite groups which are not ele-

(\*) Added in Proof June 76: Meanwhile it was proved also for singular

$\lambda = \aleph_\alpha$ .

mentary classes). Usually we get the existence of  $2^\lambda$  models, no one embeddable in another (the kind of embedding is determined by the kind of formulas exhibiting nonsuperstability). The problem of many non-isomorphic models is closely related to the existence of *rigid models*;  $[M]$  is rigid iff  $a \neq b \in M \Rightarrow (M, a) \not\equiv (M, b)$  so we can apply the method to (complete) order and (complete) Boolean algebras (see [S3]) (improving previous partial results [Jn] [Lz] [Mc] [Mm]). Another related problem is indecomposability. Here unsuperstability of the theory of modules over a ring  $R$  gives us for each regular  $\lambda > |R|$ , an  $R$ -module  $M$  of cardinality  $\lambda$  which is not the free sum of modules of smaller cardinality (see 8.6).

We could have looked at the problem differently: instead of building models from sets, i.e. cardinals, we could build them from ordered sets; e.g. to characterize a real closed field by its order. Unfortunately we know essentially nothing in this direction. We can reinterpret the problem as classifying the unstable theories by complexity. The concrete problem we have is the classification of countable (first-order) theories by Keisler's order (see [K1], [S4]).

Here the theory of linear order is maximal ([S1], VI), the stable theories are classified ([S4]), and the problem is to classify the unstable theories without the strict order property; (see [S5] § 4, [S1] II § 4, III § 7, VI). There are at least two classes, and it is clear they should have some positive properties exemplifying their simplicity.

Another related problem is e.g. categoricity over a predicate (see [G1], [G2]; this is quite complicated, (see [S6]).

*Notation:*  $L$  will be a first-order language;  $T$  a first-order theory in  $L$ ;  $\varphi, \psi, \emptyset$  formulas;  $\varphi(\bar{x})$  means all free variables of  $\varphi$  are in  $\bar{x}$ .

We identify  $L$  with its set of first-order formulas.

Let  $\Delta, \Phi, \Psi$  be sets of formulas  $\varphi(\bar{x})$ . Let  $\bigwedge \Phi = \{ \bigwedge_{i=1}^n \varphi_i : \varphi_i \in \Phi \}$  etc., and e.g.  $\exists \bigwedge \Phi = \exists (\bigwedge \Phi)$ . Sub  $\Phi$  is the set of subformulas of  $\Phi$ , including the formulas of  $\Phi$ .  $M, N$  are mo-

dels,  $|M|$  the universe of  $M$ ,  $a, b, c, d$  elements,  $\bar{a}, \dots$  finite sequences. We write  $\bar{a} \in A$  instead of  $\text{Range}(\bar{a}) \subseteq A$ ; do not differentiate between  $\bar{a}$  and its range, and write  $\bar{a} \in M$  instead of  $\bar{a} \in |M|$ .

$$\text{Let } \lambda^\alpha = \{f : \alpha \rightarrow \lambda\}, \lambda^{\alpha > \beta} = \bigcup_{\beta < \alpha} \lambda^\alpha, \lambda^{\alpha < \kappa} = \sum_{\omega \leq \mu < \kappa} \lambda^\mu$$

*Sketch Proof of the Nonstructure Theorem* ( $\lambda$  regular)

*First Approximation:* Unsuperstability gives us many types  $p_\eta$  over a small set. In fact for each  $\lambda$  there is a tree  ${}^\omega \lambda$ , points  $\{a_\nu : \nu \in {}^\omega \lambda\}$ ,  $\{b_\eta : \eta \in {}^\omega \lambda\}$  and formulas  $\varphi_n$  ( $n < \omega$ ) such that for each  $n$  the formulas  $\{\varphi_n(x, a_\nu) : \nu \in {}^\omega \lambda\}$  are pairwise inconsistent but  $\models \varphi_n(b_\eta : a_\eta \upharpoonright n)$  ( $\eta \in {}^\omega \lambda$ ). (we have  $a_\eta$  and not  $\bar{a}_\eta$ , for notational simplicity only).

The types  $p_\eta = \{\varphi_n(\bar{x}, \bar{a}_\eta) : n < \omega\}$  ( $\eta \in {}^\omega \lambda$ ) are mutually contradictory and there are  $\lambda$  of them. If we take Skolem Hulls of various sets of the  $\{b_\eta\}$  we can hopefully realize various collections of  $p_\eta$ 's while omitting others, and so get many non-isomorphic models. How shall we do this?

*Second Approximation:* Our problem would seem simpler in the unstable case, where the tree  ${}^{\omega \geq} \lambda$  would be replaced by a linear ordering containing all the points  $b_\eta$ . Then we could build Ehrenfeucht-Mostowski models over the  $\{a_\nu\}$  and

Now for  $W \subseteq \lambda$  let  $X_W = \bigcup \{S_\alpha : \alpha \in W\}$ , then the various  $M_{X_W}$  are non-isomorphic (this should be proved more carefully.)

**Fourth Approximation:** But if  $\aleph_0 > \lambda$  then  $|M_{X_W}| > \lambda$ , hence for each  $\gamma < \lambda$ ,  $cf \gamma = \aleph_0$ ; choose an increasing  $\eta_\gamma$  converging to  $\gamma$ , and redefine  $B_\alpha$  as

$$\{b_\eta\} \cup \{a_\nu : \nu \in \lambda\}$$

Now we summarize the content of each section.

§ 1. Here we review the results from [S7] on stability for homogeneous models including the stability spectrum theorem, and the equivalence between unstability and order. A worthwhile new theorem proved here is

**Theorem 1.13:** *There is a  $\mathcal{D}$ -homogeneous model of cardinality*

$$\lambda \text{ if } \lambda = \lambda^{<\lambda} + |\mathcal{D}| \text{ or } \mathcal{D} \text{ is stable in } \lambda \text{ (remember that } \mathcal{D} \text{ is stable in } \lambda \text{ iff } \lambda = \lambda(\mathcal{D}) + \lambda^{<\kappa(\mathcal{D})}).$$

This generalizes a similar theorem for saturated models (formulated in [S5] by combining results of Morley and Vaught [MV], Shelah [S7], Harnik [H1] and Shelah [S1]. Some specific instances were proved by: Eklof and Fisher [EF] (abelian groups), Boffa [Bo 1], [Bo 2], [Bo 3] (infinite generic skew fields).

§ 2. Here we discuss two additional assumptions, which make the stability theory more like that of an elementary class; prove that a class of infinite generic models satisfies one of them; generalize this; and show that the more general

some of the  $\{b_\eta\}$ , and try in this way to realize some types and to omit some others. What we must do here is:

1. Develop a theory of trees of indiscernibles and Ehrenfeucht-Mostowski models over such trees.

This can be done in a very natural way and then  $p_\eta$  is realized in the Skolem Hull of  $\{b_\eta : \eta \in X\} \cup \{a_\nu : \nu \in \lambda\}$  iff  $\eta \in X$ .

Now we start again with a tree of indiscernibles

$$\{a_\nu \in \lambda\} \cup \{b_\eta : \eta \in \lambda\}$$

to sequences  $\eta \in \lambda$  which are increasing, and we let

$$B_\alpha = \{b_\eta : \lim_{\eta \rightarrow \alpha} \eta(n) = \alpha\} \cup \{a_\nu : \nu \in \lambda\}$$

for  $\alpha$  a limit ordinal. We will build Ehrenfeucht-Mostowski models over sets of the form  $B = \bigcup_{\alpha \in X} B_\alpha$  for  $X \subseteq \lambda$  a set

of limit ordinals. Let  $M_X = EM(B_X)$  (an Ehrenfeucht-Mostowski model). We would like to have the various models  $M_X$  non-isomorphic. This requires more care.

**Third Approximation:** All we have to do now is:

2. Choose the sets  $X$  so that the models  $M_X$  are non-isomorphic.

The trick is to use stationary subsets of  $\lambda$ .

**Fact (Solovay [Sl])** ( $\lambda$  regular). There are  $\lambda$  disjoint stationary sets  $S_\alpha \subseteq \lambda$  ( $\alpha < \lambda$ ).

**Fact (Solovay [Sl]).** For  $\lambda$  regular, each stationary set can be partitioned into  $\lambda$  disjoint stationary sets, hence there are disjoint stationary  $S_\alpha \subseteq \{\gamma < \lambda : cf \gamma = \aleph_0\}$  ( $\alpha < \lambda$ ).

cases are not so far from the main example. This provides us also with a tool to construct examples of classes of infinite generic models.

*Historical Note:* The context 2.1-2.5 is of Robinson [Rb 2]; a generalization to infinitary languages  $L_{\lambda, \omega}^{\kappa}$  is due to Wood [W 3], and a definition and investigation of  $K^n$  is in Cherlin [Cr] (for the category of embeddings), sketching the generalization to infinitary languages  $L_{\lambda, \kappa}$ ; so in 2.6-2.10(1) there is nothing essentially new. (\*)

§ 3. Here we define appropriate ranks, prove some of their properties, and use them to investigate the spectrum of stability.

As the treatment of Morley [Mr] of ranks and types is related to Boolean algebras, our treatment is related to lattices (see Simmons paper in this volume).

H. Priestly [Pr] develops the right kind of Stone space for lattices. Fisher, Simmons and Wheeler [F S W] and McKenzie and Shelah [McS] prove some model theoretic results essentially equivalent to the fact that countable lattices have  $\leq \aleph_0$  or  $2^{\aleph_0}$  ultrafilters (in [F S W] this is done for the number of theories of e. closed models of T, finite-generic models of T, and infinite generic models of T, T countable; in [McS] for the number of ep-types realized over a countable set in appropriate  $\aleph_1$ -homogeneous models.

§ 4. Here we deal with  $\Psi$ -prime models. We describe the natural way to construct prime models, prove their existence. They are unique when our class is  $\aleph_0$ -stable. The most important theorem is that for a countable stable T, if over A there is a strictly prime model, it is unique; we prove it for  $|A| \leq \aleph_1$  only.

§ 5 Here we show that Morley's categoricity theorem generalizes easily.

(\*) Added in proof June 76: Independently, W. Forrset generalized some theorems on  $\aleph_0$ -stable theories to the context of assumption II (see § 1).

§ 6. In the context of assumption II (i.e. infinite generic model for a universal T with the amalgamation property) we develop «forking» and prove its important properties.

Some of this, mainly the symmetry lemma, was developed later and independently, from a different point of view by Lascar (see this volume and [Ls]). He concentrates on super-stable T and complete types over models.

§ 7. We deal with some «negative» results. We quote Macintyre that any infinite field with an  $\aleph_{\alpha}$ -stable theory is algebraically closed; and then prove that  $\aleph_1$ -categorical division rings are fields, using Baldwin theorem ( $\alpha_T < \omega$ ). We then show how our general results include a theorem of Boffa [Bo 1], [Bo 2], [Bo 3], (on  $\{\lambda: \text{there is an infinite generic division ring of card } \lambda\}$ ) and solve one of his problems (on the number of non-isomorphic infinite generic division rings).

*Historical Note:* (Lemma 7.5 (1) is from Macintyre [Mc 2] and 7.3-7.6 are from letters to Macintyre distributed in summer 73. Later Baldwin and Saxel [BS] reproved 7.5. There exist papers of Zilberg on  $\aleph_1$ -categoricity of rings, partially overlapping Cherlin and Reineke [CR] (\*).

§ 8. We mainly deal with modules. They are always stable (Baur [Br] Fisher [Fi]) and we characterize  $\kappa(T)$ , and in particular superstability. We also give essentially, a structure/non-structure theorem for the class of R-modules over a fixed R, and show that formulas in this theory are not too complicated (8.3).

We also deal with differential fields and with separably closed fields and conclude with a criterion for an operation on models to preserve stability.

§ 1. HOMOGENEITY.

We shall deal in this article with four kinds of classes, for which we have stability theory. They are essentially:

Kind I: Elementary classes.

(\*) Added in Proof June 76: Zilberg also proved independently that  $\aleph_1$ -categorical division rings are fields.

Kind II: The e.c. (existentially closed) models of an universal theory with the J.E.P. (joint embedding property) and the amalgamation property.

Kind III: The e.c. models of a universal theory with the J.E.P.

Kind IV: The elementary submodels of some  $(\mathcal{D}, \lambda)$  — homogeneous model. (\*)

More complicated cases, like the class of models of some  $\psi \in L_{\omega_1, \omega}$  or abstract elementary classes will be dealt with elsewhere. In kind II (but not III) we can deal with the class of models of T. (†)

In kind I embeddings are elementary.

Kind I will be the subject of the book [S1], and kind IV was the subject of [S7]; the other two constitute the (small) novelty of this article. Here we shall review the necessary facts from [S7], in order to use them and to see how restricting the context adds to the information. We shall also note that kinds II and III can be replaced by a somewhat more general condition. Some of the results will be reproved in the context of kinds II and III.

We prove here one new theorem: there is a  $(\mathcal{D}, \lambda)$  — homogeneous model of cardinality  $\lambda$  iff  $\lambda = \lambda^{<\lambda} + |\mathcal{D}|$  or  $\mathcal{D}$  is stable in  $\lambda$  (for good  $\mathcal{D}$ ).

Definition 1.1.

We define the  $\Delta$ -type of the sequence  $\bar{a}$  (usually finite) in  $M$  over  $A$  by

$$tp_{\Delta}(\bar{a}, A, M) = \{ \varphi(\bar{x}, \bar{b}) : \bar{b} \in A, M \models \varphi(\bar{a}, \bar{b}), \varphi(\bar{x}, \bar{y}) \in \Delta \}$$

(\*) We assume there is a  $(\mathcal{D}, \lambda)$ -homogeneous model of cardinality  $\geq \lambda$  for every  $\lambda$ .

(†) Note that in kind III, for every model  $M$  of the theory, the type of every  $\bar{a} \in M$  in any universal - homogeneous model  $N, M \subseteq N$ , is uniquely determined iff  $M$  is e-closed; whereas in kind II this holds for any  $M$ . Hence it is natural in kind II to deal with the class of models of the theory and in kind III with the e-closed ones.

When  $\Delta$  is  $L(M)$  = the set of first order formulas in the language of  $M$ , we omit it.

Definition 1.2.

The finite diagram of  $M, \mathcal{D}(M)$ , is  $\{ tp(\bar{a}, \emptyset, M) : \bar{a} \in M, (\bar{a} \text{ finite}) \}$ .

Let  $\mathcal{D}$  denote such sets.

Definition 1.3.

$M$  is  $\lambda$ -homogeneous if for every  $a_i, b_i \in M, i < \alpha < \lambda$  and  $a \in M$  such that

$$tp(\langle a_i : i < \alpha \rangle, \emptyset, M) = tp(\langle b_i : i < \alpha \rangle, \emptyset, M)$$

there is a  $b \in M$  such that

$$tp(\langle a_i : i \leq \alpha \rangle, \emptyset, M) = tp(\langle b_i : i \leq \alpha \rangle, \emptyset, M).$$

Definition 1.4.

$M$  is  $(\mathcal{D}, \lambda)$  — homogeneous if it is  $\lambda$ -homogeneous and  $\mathcal{D}(M) = \mathcal{D}$ .  $M$  is  $\mathcal{D}$ -homogeneous if it is  $(\mathcal{D}, \|M\|)$  — homogeneous.

Assumption 1.  $\mathfrak{C}$  is some fixed  $(\mathcal{D}, \lambda_0)$ - homogeneous model, for a fixed  $\mathcal{D}$  and  $\lambda_0$  sufficiently large. All models are elementary submodels of  $\mathfrak{C}$ , all elements, sequences of elements and sets  $(a, \dots, \bar{a}, \dots, A, \dots)$  are from  $\mathfrak{C}$ . Satisfaction  $\models$  will mean «in  $\mathfrak{C}$ ».

Definition 1.5.

$$S^m_{\mathcal{D}}(A) = \{ tp(\bar{a}, A, \mathfrak{C}) : \bar{a} \in \mathfrak{C}, l(\bar{a}) = m \}. \text{ (If } A \subset M < \mathfrak{C}, S^m_{\mathcal{D}}(A) \text{ depends only on } m, M, A \text{ and } \mathcal{D} \text{.)}$$

Lemma 1.6.

(1) A set  $p$  of formulas with parameters from  $A$  is in  $S^m_{\mathcal{D}}(A)$

iff:

- (i) the variables are  $\bar{x} = \langle x_0, \dots, x_{m-1} \rangle$
- (ii)  $p$  is complete finitely satisfiable
- (iii) for each  $\bar{b} \in A, \{\varphi(\bar{x}; \bar{y}) : \varphi(\bar{x}; \bar{b}) \in p\} \in \mathcal{D}$
- (2)  $M$  is  $(\mathcal{D}, \lambda)$  — homogeneous iff whenever  $A \subseteq M, |A| < \lambda$ , and  $p \in S^m_{\mathcal{D}}(A)$  then  $p$  is realized

(3) If  $M, N$  have cardinality  $\lambda, \mathcal{D}(M) = \mathcal{D}(N)$ , and  $M, N$  are homogeneous then  $M \cong N$ . If only  $M$  is  $\lambda$  — homogeneous,  $N$  can be elementarily embedded into  $M$ .  
 Lemma 1.6, (2), stresses the similarity of homogeneity and saturation.

**Definition 1.7.**

- (1)  $\mathcal{D}$  is  $\lambda$ -stable iff  $m < \omega$  and  $|A| \leq \lambda$  implies  $|S^m_{\mathcal{D}}(A)| \leq \lambda$ .
- (2)  $\mathcal{D}$  is stable if it is  $\lambda$ -stable for some  $\lambda$ .

**Definition 1.8.**

$\mathcal{D}$  has the  $\lambda$ -order property if there are  $\bar{a}_i (i < \lambda)$  and  $\varphi(\bar{x}, \bar{y})$  such that  
 $\models \varphi(\bar{a}_i, \bar{a}_j)$  iff  $i < j$ .

**Theorem 1.9. (Stability vs. Order)**

- (1) If  $\mathcal{D}$  has the  $\lambda$ -order property for all  $\lambda$  then  $\mathcal{D}$  is unstable (in all  $\lambda$ ).
- (2) If  $\mathcal{D}$  is not  $\lambda$ -stable,  $\lambda = \lambda^x + 2^{2^x}, x \geq |L|$  then  $\mathcal{D}$  has

the  $x^+$ -order property.

- (3) If  $\mathcal{D}$  has the  $x$ -order property for every  $x < \beth(|L| + 1)$ , then  $\mathcal{D}$  has the  $x$ -order property for all  $x$ .

**Remark:** Essentially this theorem says that unstability and order are equivalent. See [S7], [S8].  
**Theorem 1.10. (The stability spectrum theorem).**  
 Always (1)  $\mathcal{D}$  is unstable

- or (2) there exist cardinals  $\kappa(\mathcal{D}), \lambda(\mathcal{D}) < \beth(|L| + 1)$

such that:  $\mathcal{D}$  is  $\lambda$ -stable iff  $\lambda = \lambda(\mathcal{D}) + \lambda^{<\kappa(\mathcal{D})}$ .  $(\lambda^{<\kappa(\mathcal{D})} = \sum_{\mu < \kappa} \lambda^\mu)$ .

**Remarks:** (1) In (1) we stipulate  $\lambda(\mathcal{D}) = \kappa(\mathcal{D}) = \infty$ ; in (2) if we demand « $\kappa(\mathcal{D})$  is regular» then  $\kappa(\mathcal{D})$  is unique.

(2) In [S7] we prove more; that if there is a  $(\mathcal{D}, \lambda^+)$  — homogeneous model of cardinality  $\geq \lambda^+$  and  $\mathcal{D}$  is  $\lambda$ -stable then there are  $(\mathcal{D}, \kappa)$  — homogeneous models for all  $\kappa$ .

(3) For theories, and even for kind III,  $\kappa(\mathcal{D}) \leq |T|^+$  and  $\lambda(\mathcal{D}) \leq 2^{|L|}$ .

**Definition 1.11.**

Let  $I$  be a set of sequences of fixed (finite) length from  $\mathcal{C}$ .  $I$  is indiscernible over  $A$  iff for all distinct  $\bar{a}_i \in I (i < n)$ ,  $\text{tp}(\bar{a}_0, \dots, \bar{a}_{n-1}, A, \mathcal{C})$  is fixed.

**Theorem 1.12.**

- (1) If  $\mathcal{D}$  is  $\lambda$ -stable,  $|A| \leq \lambda < |I|$ ,  $I$  a set of sequences from  $\mathcal{C}$ , then there exists  $I' \subseteq I, |I'| = \lambda^+, I'$  indiscernible over  $A$ .
- (2) If  $I$  is indiscernible over  $A$ , for any  $\bar{a}$  there is  $J \subseteq I, |J| < \kappa(\mathcal{D})$  such that  $I - J$  is indiscernible over  $A \cup J \cup \bar{a}$ .

**Remark.**

By 1.12 (2) for any indiscernible set  $I(\bar{a} \in I \Rightarrow \ell(\bar{a}) = m), |I| \geq \kappa(\mathcal{D})$ , and  $A$ , we can define  $\text{Av}(I, A) = \{\varphi(\bar{x}, \bar{b}) : \text{for } \kappa(\mathcal{D}) \text{ } \bar{a}'\text{'s in } I \models \varphi(\bar{a}, \bar{b})\} \in S^m_{\mathcal{D}}(A)$ .

For theories and even kind II it suffices to assume  $I$  is infinite. Question: what about kind III and IV?

**Theorem 1.13. (Homogeneity spectrum theorem). (\*)**

There is a  $(\mathcal{D}, \lambda)$  — homogeneous model of cardinality  $\lambda$  iff

$$\lambda = \lambda^{<\lambda} + |\mathcal{D}| \text{ or } \mathcal{D} \text{ is stable in } \lambda.$$

(\*) In fact, it suffices to assume on  $\mathcal{D}$  that there is a  $(\mathcal{D}, \lambda)$ -homogeneous model of cardinality  $\geq \lambda$  and replace « $\mathcal{D}$  stable in  $\lambda$ » by « $M$  stable in  $\lambda$ ».

*Remark.*

Naturally this should have appeared in [S7] but I did not know it at the time. The reader can skip it easily.

*Proof.* The proof is split into cases.

case 1:  $\lambda = \lambda^{<\lambda} + |\mathcal{D}|$

It is easy to prove the existence (by the classical [MV]).

case 2:  $\lambda < |\mathcal{D}|$ .

The non-existence is trivial.

case 3:  $\lambda < \lambda^{<\lambda}$ ,  $\mathcal{D}$  unstable.

See [S7], 6.5 (2), using the order property.

case 4:  $\mathcal{D}$  stable in  $\lambda$ .

If  $\lambda$  is regular the proof is trivial.

So assume  $\lambda$  is singular. Now we start using [S7] notions and theorems heavily.

As  $\mathcal{D}$  is stable in  $\lambda$ ,  $\lambda = \lambda^{<\kappa(\mathcal{D})}$  hence  $\kappa(\mathcal{D}) \leq \text{cf } \lambda$ .

We define an increasing sequence  $M_i$  ( $i \leq \lambda^2$ ),  $\|M_i\| = \lambda$  such that each  $q \in S^m_{\mathcal{D}}(M_i)$  is realized in  $M_{i+1}$  and  $M_\delta = \bigcup_{i < \delta} M_i$  for limit  $\delta$ .

Let  $A_0 \subseteq M_{\lambda^2}$ ,  $|A_0| < \lambda$ ,  $p_0 \in S^m_{\mathcal{D}}(A_0)$  and we should prove that  $p_0$  is realized in  $M_{\lambda^2}$ .

Let  $p_0 \subseteq p \in S^m_{\mathcal{D}}(M_{\lambda^2})$ , and choose  $C \subseteq M_{\lambda^2}$ ,  $|C| < \kappa(\mathcal{D})$

such that  $p$  does not split strongly over  $C$  (see [S7] §4). As  $\text{cf } \lambda \geq \kappa(\mathcal{D})$  for some  $\alpha_0 < \lambda^2$ ,  $C \subseteq M_{\alpha_0}$ .

As  $\mathcal{D}$  is stable in  $\lambda$ ,  $C$  is stable in  $\lambda$ , hence  $\{i < \lambda : p \upharpoonright M_{\lambda(i+1)}$  splits over  $M_{\lambda^1}\} < \lambda$ , hence for some  $\beta, \alpha_0 < \beta < \lambda^2$  and  $p \upharpoonright M_{\beta+\lambda}$  does not split over  $M_\beta$ .

Now we shall prove that  $p$  does not split over  $M_{\beta+\lambda}$ . Otherwise

wise suppose  $\bar{b}, \bar{c} \in M_{\lambda^2}$ ,  $\text{tp}(\bar{b}, M_{\beta+\lambda}) = \text{tp}(\bar{c}, M_{\beta+\lambda})$ ,  $\varphi(\bar{x}; \bar{b}) \wedge \varphi(\bar{x}; \bar{c}) \in p$ . As before there is  $\gamma, \beta < \gamma < \beta + \lambda$  such that  $\text{tp}(\bar{b}, M_{\gamma+\omega})$  does not split over  $M^\lambda$  ( $\lambda > \omega$  as  $\lambda$  is singular).

Choose  $\bar{b}_n \in M_{\gamma+n+1}$  which realizes  $\text{tp}(\bar{b}, M_{\gamma+n})$ , so by [S7], p. 83,  $\{\bar{b}_n : n < \omega\}$  is indiscernible over  $M_\lambda$ , and so are  $\{\bar{b}_n : n < \omega\} \cup \{\bar{c}_n : n < \omega\} \cup \{\bar{c}\}$ . Hence  $p$  splits strongly over  $M_\gamma$ ; hence over  $C$ , contradiction. So  $p$  does not split over  $M_{\beta+\lambda}$ .

For each  $\gamma, \beta + \lambda < \gamma < \lambda^2$  choose  $\bar{a}_\gamma \in M_{\gamma+1}$  realizing  $p \upharpoonright M_\gamma$ , hence, again by [S7] p. 83,  $I = \{\bar{a}_\gamma : \beta + \lambda < \gamma < \lambda^2\}$  is indiscernible.

For each  $\bar{c} \in M_{\lambda^2}$ ,  $I \cup \{\bar{c}\}$  realizes  $p \upharpoonright \bar{c}$  has cardinality  $\lambda$ , hence  $|I - I_{\bar{c}}| < \kappa(\mathcal{D})$ , hence  $p \upharpoonright A_0 = p_0$  is realized by some  $\bar{a}_\gamma \in I$ .

Before treating the final case, we prove

Lemma 1.14. Suppose  $A \subseteq B$ , and each  $p \in S^m_{\mathcal{D}}(A)$  ( $m < \omega$ ) is realized in  $B$ . Then  $|\Gamma| \leq |U S^m_{\mathcal{D}}(A)|^{\aleph_0} \leq |\mathcal{D}|^{|A| + \aleph_0}$  where  $\Gamma = \{p : p \in S^m_{\mathcal{D}}(B), m < \omega, p \text{ does not split over } A\}$ .

*Remark:* This improves [S7], 2.5.

*Proof:* For each  $p \in \Gamma$  we define by induction  $\bar{a}_p^i$  ( $i < \kappa(\mathcal{D})$ ) such that  $\text{tp}(\bar{a}_p^i, B \cup \{\bar{a}_p^j : j < i\})$  extends  $p$  and does not split over  $A$ .

This is possible by [S7] 2.2.(2), and  $I_p = \{\bar{a}_p^i : i < \kappa(\mathcal{D})\}$  is

some  $M_i$  (If  $\lambda$  is regular because  $\lambda_*(\mathcal{D}) \leq \lambda$ ; if  $\lambda$  is singular choose for each  $i < \text{cf} \lambda$   $\bar{a}_i, \bar{b}_i \in M_i$ ,  $\text{tp}(\bar{a}_i, M_i) = \text{tp}(\bar{b}_i, M_i)$  but  $\varphi_i(\bar{x}, \bar{a}_i) \wedge \neg \varphi_i(\bar{x}, \bar{b}_i) \in p$ . Now let  $A = \bigcup_{i < \text{cf} \lambda} \bar{a}_i \wedge \bar{b}_i \subseteq |M|$ , so  $|A| < \lambda$  and  $p \upharpoonright A \in S^m_{\mathcal{D}}(A)$  is not realized in  $M_i$ ; contradiction.)

So for some  $\alpha < \text{cf} \lambda$ ,  $m < \omega$ ,  $\Gamma = \{p \in S^m_{\mathcal{D}}(M) : p \text{ does not split over } M\}$  has cardinality  $> \lambda$ . As  $\bigcup_{m < \omega} S^m(M_i) \leq \lambda$  (because  $\|M_i\| < \lambda$  and the assumption), by lemma 1.14, we have necessarily that  $\lambda^{N_0} > \lambda$ ; hence by assumption  $\kappa(\mathcal{D}) = N_0$ . Let  $\bar{a}^i_p (i \leq \omega)$  be defined as in lemma 1.14. for each  $p \in \Gamma$

with  $M_\alpha$  for  $A$  and  $M$  for  $B$ . So  $p \neq q \in \Gamma$  implies that for some  $n$ ,  $\text{tp}(\langle \bar{a}^i_p : i < n \rangle, M_i) \neq \text{tp}(\langle \bar{a}^i_q : i < n \rangle, M_i)$ . Now we define by induction on  $i \leq \omega$ ,  $\bar{b}^i_p \in M$  for each  $p \in \Gamma$  such that

- (i)  $\text{tp}(\langle \bar{b}^j_p : j \leq i \rangle, M_\alpha) = \text{tp}(\langle \bar{a}^j_p : j \leq i \rangle, M_\alpha)$
- (ii) If  $\text{tp}(\langle \bar{a}^j_p : j \leq i \rangle, M_\alpha) = \text{tp}(\langle \bar{a}^j_q : j \leq i \rangle, M_\alpha)$  then  $\bar{b}^j_p = \bar{b}^j_q$  for  $j \leq i$ .

Clearly this is possible (as  $\lambda > N_0$  by the hypothesis  $\lambda < \lambda^{< \lambda}$ , and  $p \neq q$  implies  $\bar{b}^{\omega}_p \neq \bar{b}^{\omega}_q$  because there is a minimal  $n < \omega$  such that  $\text{tp}(\langle \bar{a}^j_p : j \leq n \rangle, M_\alpha) \neq \text{tp}(\langle \bar{a}^j_q : j \leq n \rangle, M_\alpha)$ , hence  $\bar{b}^j_p = \bar{b}^j_q$  for  $j < n$  but for  $r = p, q$   $\text{tp}(\langle \bar{a}^r_0 \wedge \dots \wedge \bar{a}^{n-1}_r, \bar{a}^n_r \rangle, M_\alpha) = \text{tp}(\langle \bar{a}^r_j : j \leq n \rangle, M_\alpha)$

indiscernible over  $B$ . Now (\*) if  $\bar{b} \in B$ ,  $\text{tp}(\bar{b}_i, A) = \text{tp}(\bar{b}, A)$ ,  $q = \text{tp}(\bar{b} \wedge \bar{a}^{\omega}_p, A)$  then  $\{i < \kappa(\mathcal{D}) : q \neq \text{tp}(\bar{b}_i \wedge \bar{a}^{\omega}_p, A)\} < \kappa(\mathcal{D})$ .

(because we can define  $\bar{a}^i_p (\kappa(\mathcal{D}) \leq i < \kappa(\mathcal{D})^+)$  such that  $\text{tp}(\bar{a}^i_p, B \cup \{\bar{a}^j : j < i\} \cup \bar{b}_i)$  extends  $p$  and does not split over  $A$ . Again this is possible and  $\Gamma = \{\bar{a}^i_p : i < \kappa(\mathcal{D})^+\}$  is indiscernible. Clearly for  $i \geq \kappa(\mathcal{D})$ ,  $\text{tp}(\bar{b}_i \wedge \bar{a}^i_p, A) = \text{tp}(\bar{b} \wedge \bar{a}^i_p, A) = \text{tp}(\bar{b} \wedge \bar{a}^{\omega}_p, A)$ . So  $\{i \in \Gamma : q = \text{tp}(\bar{b}_i \wedge \bar{a}^i_p, A)\} > \kappa(\mathcal{D})$ ; hence, by 1.12(2),  $\{\bar{a} \in \Gamma : \text{tp}(\bar{b} \wedge \bar{a}, A) \neq \text{tp}(\bar{b}_i \wedge \bar{a}, A)\} < \kappa(\mathcal{D})$ . So (\*) follows.)

So if  $p \neq q \in \Gamma$ , there is no automorphism  $F$  of  $\mathbb{C}$ , with  $F \upharpoonright A =$  the identity and  $F(\bar{a}^i_p) = \bar{a}^i_q (i < \kappa(\mathcal{D}))$ . Hence  $\text{tp}(\langle \bar{a}^i_p : i < \kappa(\mathcal{D}) \rangle, A) \neq \text{tp}(\langle \bar{a}^i_q : i < \kappa(\mathcal{D}) \rangle, A)$  hence by the indiscernibility  $\text{tp}(\langle \bar{a}^i_p : i < \omega \rangle, A) \neq \text{tp}(\langle \bar{a}^i_q : i < \omega \rangle, A)$ . Simple computation gives our result.

Continuation of the proof of 1.13. Case 5.  $\mathcal{D}$  is not stable in  $\lambda$ , but  $\mathcal{D}$  is stable,  $\lambda \geq |\mathcal{D}|$  and  $\lambda < \lambda^{< \lambda}$ .

If  $\lambda^{< \kappa(\mathcal{D})} > \lambda$ , this is easy by [S7] §6. So we assume  $\lambda =$

$< \kappa(\mathcal{D})$ . Similarly we can assume that  $2^\mu > \lambda$  implies not (\* $\mu$ ), and let  $\lambda_*(\mathcal{D})$  be the first cardinal  $\mu$  such that not (\* $\mu$ ): so  $2^{< \lambda_*(\mathcal{D})} \leq \lambda$ . We can also assume  $|A| < \lambda$  implies  $|S^m_{\mathcal{D}}(A)| \leq \lambda$ .

Suppose  $M$  is a  $(\mathcal{D}, \lambda)$  — homogeneous model of cardinality  $\lambda$ , so by hypothesis  $|S^m_{\mathcal{D}}(M)| > \lambda$ . Let  $M = \bigcup_{i < \text{cf} \lambda} M_i$ ,  $M_i$  is increasing,  $\|M_i\| < \lambda$ . Now each  $p \in S^m_{\mathcal{D}}(M)$  does not split over

So we get a contradiction:  $\lambda \geq \|M\| = |\{\bar{v}_p : p \in \Gamma\}| > \lambda$ . Similarly we can prove

Theorem 1.15.

Suppose  $M_i$  is  $(\mathcal{D}, \lambda)$  — homogeneous,  $i < \delta$ , cf  $\delta \geq \kappa(\mathcal{D})$ ,  $\lambda_* (\mathcal{D}) \leq \lambda$ ,  $M_i$  increasing. Then  $\bigcup_{i < \delta} M_i$  is  $(\mathcal{D}, \lambda)$  — homogeneous.

The categoricity theorem is

Theorem 1.16.

Suppose the class  $\{M : M < \mathfrak{C}\}$  is categorical in one  $\lambda > |L|$ . Then there is  $\lambda_1 < \beth_{\omega_1} |L|$  such that

- (1) the class is categorical in every  $\lambda \geq \lambda_1$ ; moreover  $M < \mathfrak{C}$ ,  $\|M\| \geq \lambda_1$  implies  $M$  is  $\mathcal{D}$ -homogeneous.
- (2) For every  $\mu, |L| < \mu < \lambda_1$  the class is not categorical in  $\mu$ ; moreover there is a non —  $|L|^+$  — homogeneous model  $M < \mathfrak{C}$  of cardinality  $\mu$ .

Remark: See [S7] for better results, i.e. applicable to more cases. But we do not know whether the bound on  $\lambda_1$  can be improved. Related questions: suppose there is a  $(\mathcal{D}, \lambda^+)$  — homogeneous model of cardinality  $\lambda$ ; is  $\lambda \leq 2^{|L|}$ ?

For kind IV, can  $\kappa(T)$  be  $> (2^{|L|})^+$ ?

For which  $\lambda$  does the existence of a  $(\mathcal{D}, \lambda)$  — homogeneous model imply the existence of a  $(\mathcal{D}, \mu)$  — homogeneous model for every  $\mu$ ?

§ 2. GENERIC MODELS

Now we can state more exactly what extra assumptions we use in kinds II, III, and explain why the generic models satisfy them.

(\*) Notice that 1.14 has new consequences concerning the question of Keisler and Morley, on the number of homogeneous models of  $T$  in cardinality  $\lambda$ , when G.C.H. fails.

Assumption for kind III: there exists a set  $\Phi$  of formulas such that (where  $\mathfrak{C}, \mathcal{D}$ , are as before)

- (i) if  $p_1, p_2$  are distinct  $m$ -types in  $\mathcal{D}$ , then there are  $\varphi_1(\bar{x}) \in p_1, \varphi_2(\bar{x}) \in p_2$  which are in  $\Phi$  and are contradictory, that is  $\mathfrak{C} \models \neg (\exists \bar{x}) (\varphi_1(\bar{x}) \wedge \varphi_2(\bar{x}))$
- (ii) any set of formulas from  $\Phi$ ; with parameters from  $\mathfrak{C}$ , of cardinality  $< |\mathfrak{C}|$  which is finitely satisfiable in  $\mathfrak{C}$ , is satisfiable in  $\mathfrak{C}$ .

Assumption for kind II: as the previous one, when we add

- (iii) every  $\varphi(\bar{x}) \in \Phi$  has a negation in  $\Phi$ , that is a  $\psi(\bar{x}) \in \Phi$  such that  $\mathfrak{C} \models (\forall \bar{x}) (\varphi(\bar{x}) \equiv \neg \psi(\bar{x}))$ .

Remark: In the assumption for kind III, we can assume  $\Phi$  is closed under existential quantification, and for kind II, under negation. In any case we can assume it is closed under conjunctions and disjunctions.

We deviate from the assumption of § 1, by letting  $M$  be any submodel of  $\mathfrak{C}$ , such that for any formula  $\varphi(\bar{x})$  which is in sub  $\Phi, \bar{a} \in M \Rightarrow \mathfrak{C} \models \varphi(\bar{a}) \leftrightarrow M \models \varphi(\bar{a})$ . ( $M$  is called  $\Phi$ -closed; or  $M <_{\text{sub } \Phi} \mathfrak{C}$ ).

In this section we prove that the assumptions defined above are satisfied by the appropriate kinds.

We let now  $T$  denote a fixed universal theory with the JEP.

Definition 2.1. (A)  $T^\omega$  is the unique class of models of  $T$  such that

- (1)  $T^\omega$  is cofinal in  $T$  i.e. in  $\text{Mod}(T)$  (= the class of models of  $T$ ); that is every model of  $T$  is embeddable in a model of  $T^\omega$

(the usual notation is  $G_T$ , Cherlin [Cr] uses  $T^\infty$ ).

- (2)  $T^\omega$  is model complete, i.e. if  $M_1, M_2 \in T^\omega$  and  $M_1 \subseteq M_2$ , then  $M_1 < M_2$ .

- (3) If  $M_1 < M_2 \in T^\omega$ , then  $M_1 \in T^\omega$ .

Note that  $T^\omega$  is closed under unions of increasing chains.

Definition 2.1. (B)  $M$  is infinitely generic (for  $T$ ) if  $M \in T^\omega$ .

Definition 2.2.

(A)  $\mathcal{D}_\Delta(T) = \{tp_\Delta(\bar{a}, \emptyset, M) : \bar{a} \in M, M \text{ a model of } T\}$ ,

$\mathcal{D}_T = \mathcal{D}_L(T)$ .

(B)  $M <_\Delta N$  if  $\bar{a} \in M, \varphi \in \Delta$ , implies  $M \models \varphi[\bar{a}] \leftrightarrow N \models \varphi[\bar{a}]$

and  $M \subseteq N$ .

(C) In  $\mathcal{D}_\Delta$ ,  $tp_\Delta$ , etc., when  $\Delta$  is the set of existential, positive-

existential, quantifier-free, atomic formulas, we write instead of  $\Delta, e, ep (= pe), qf, a$  resp. and denote  $\Delta$  itself by  $\Phi_e, \Phi_{ep}, \Phi_{qf}, \Phi_a$  resp.

Lemma 2.3. (1) If  $M$  is a model of  $T, \alpha$  an ordinal,

$$tp_\alpha(\langle a_i : i < \alpha \rangle, \emptyset, M) \subseteq tp_\alpha(\langle b_i : i < \alpha \rangle, \emptyset, M)$$

then for every  $a_\alpha \in M$  there are  $N$  and  $b_\alpha \in N$  such that  $M \subseteq N$ , and

$$tp_\alpha(\langle a_i : i \leq \alpha \rangle, \emptyset, N) \subseteq tp_\alpha(\langle b_i : i \leq \alpha \rangle, \emptyset, N)$$

We can replace  $\subseteq$  by  $=$  (for the types).

(2) For any model  $M$  of  $T$  and  $\kappa$ , there is a model  $N, M \subseteq N$  such that

(\*) For any  $a_i, b_i \in N (i < \alpha < \kappa)$  such that

$$tp_\alpha(\langle a_i : i < \alpha \rangle, \emptyset, N) = tp_\alpha(\langle b_i : i < \alpha \rangle, \emptyset, N)$$

for any  $a_\alpha \in N$  there is  $b_\alpha \in N$  such that

$$tp_\alpha(\langle a_i : i \leq \alpha \rangle, \emptyset, N) = tp_\alpha(\langle b_i : i \leq \alpha \rangle, \emptyset, N).$$

(3) If in (2) in  $M$  every  $p \in \mathcal{D}_\alpha(T)$  is realized, then (\*) is equivalent to " $N$  is  $(\mathcal{D}^\alpha(T), \kappa)$ -homogeneous" where  $\mathcal{D}^\alpha(T) = \{tp(\bar{a}, \emptyset, M) : \bar{a} \in M_\alpha\}$ , where  $M_\alpha$  realizes every  $p \in \mathcal{D}_\alpha(T)$  and satisfies (\*).

- (4) Every  $(\mathcal{D}^\alpha(T), \aleph_\alpha)$ -homogeneous model belongs to  $T^\omega$ .
- (5) If  $T$  has the amalgamation property, we can replace in (1) - (4),  $e$  by  $qf$ .

Conclusion 2.4.  $\mathcal{D}^\alpha(T)$  satisfies the assumption for kind III (with  $\Phi_e$ ). If  $T$  has the amalgamation property, then  $\mathcal{D}^\alpha(T), \Phi_{qf}$  satisfy the assumption for kind II.

Proof: Easy.

Remark: Note that by using  $\mathcal{C}$ , the existential type of each  $\bar{a}$  is determined. Note

Claim 2.5.  $M \in T^\omega$  iff  $M < N$  for some  $(\mathcal{D}^\alpha(T), \lambda)$ -homogeneous models iff  $\mathcal{D}(M) \subseteq \mathcal{D}^\alpha(T)$ .

\* \* \*

We can present genericity in a more general way.

Context: let  $L$  be a language, and  $K$  a category of  $L$ -models such that

- (1) the morphisms are some functions from one model to another
- (2) direct limits exist
- (3) for every atomic  $R(\bar{x}) \in L$  and  $\bar{a} \in M \in K$  there are  $g, N \in K, g : M \rightarrow N$  such that for all  $g' : N \rightarrow N'$  in  $K$ , the truth value of  $R(g'g(\bar{a}))$  (in  $N'$ ) is the same.

We say  $K$  has the J M P (joint mapping property) if for every  $M_1, M_2 \in K$  there are  $g : M_1 \rightarrow N (f = 1, 2)$  in  $K$ .

Definition 2.6. Let  $\Psi$  be a set of formulas from  $L_{\infty, \omega}$  closed under subformulas, and let  $\Psi_\alpha = \{\varphi \in \Psi : \varphi \text{ has quantifier depth } \leq \alpha\}$ .

We define full subcategories  $K^\alpha (= K^\alpha(\Psi))$  of  $K$ , and for each  $M \in K, \varphi(\bar{x}) \in \Psi$ , when  $M \models \varphi[\bar{a}]$ . (If  $\Psi$  is not mentioned, we let  $\Psi = L_{\infty, \omega}$ ).

(1) for  $\alpha = 0, M \in K^\alpha$  iff for each atomic formula  $R(\bar{x})$  and  $g : M \rightarrow N$  in  $K$ , and  $\bar{a} \in M, M \models R[\bar{a}] \leftrightarrow N \models R(g(\bar{a}))$

(2) we define  $M \Vdash \varphi [a]$  by induction on  $\varphi$ , where  $M \in K$ ,  $\varphi \in \Psi$ ,  $a \in M$

- (A) for atomic  $\varphi$ ,  $M \Vdash \varphi [a]$  iff  $M_1 \models \varphi[g(a)]$  for each  $g : M \rightarrow M_1$  in  $K$
- (B)  $M \Vdash \bigwedge \varphi_i$  iff  $M \Vdash \varphi_i$  for each  $i$
- (C)  $M \Vdash \exists x \varphi [x, a]$  iff there is  $b \in M$  such that  $M \Vdash \varphi [b, a]$
- (D)  $M \Vdash \neg \varphi (a)$  iff there are no  $g : M \rightarrow N$  in  $K$  such that  $N \Vdash \varphi [g(a)]$

(3)  $M \in K^\alpha$  iff for each  $\varphi \in \Psi$ ,  $a \in M$ ,  $M \models \varphi [a] \Leftrightarrow M \Vdash \varphi [a]$

(4)  $K^\omega$  is called the class of  $K$ -generic models.

Lemma 2.7.

- (1)  $K^\alpha$  is decreasing.
- (2) Each  $K^\alpha$  is unbounded in  $K$ , i.e. for every  $M \in K$  there is  $g : M \rightarrow N$  in  $K$ ,  $N \in K^\alpha$ . Also each  $K^\alpha$  is closed under direct limits.
- (3) If  $g : M \rightarrow N$  is in  $K$ ,  $a \in M$ ,  $M \Vdash \varphi [a]$  then  $N \Vdash \varphi [a]$ .
- (4) If  $K$  has the amalgamation property,  $\Psi$  closed under negation then for every  $\varphi(x) \in \Psi$ ,  $a \in M \in K$ ,  $K \Vdash \neg \varphi(x)$  or  $K \Vdash \neg \neg \varphi(x)$ .

Remark: We could do the same for  $\Psi \subseteq L_{\infty, \aleph}$  but then  $K^\alpha$  is closed only for direct limits of cofinality  $\geq \aleph$ .

Definition 2.8. (A) Let  $T$  be a theory in  $L$ ,  $\Phi$  a set of  $L$ -formulas.  $K(T, \Phi)$  will be the following category:

- (i)  $M \in K(T, \Phi)$  iff  $M$  is a model of  $T$  (an  $L$ -model, of course)
- (ii)  $g : M \rightarrow N$  is a morphism if  $\varphi(x) \in \Phi$ ,  $a \in M$ ,  $M \models \varphi [a]$  implies  $N \models \varphi [g(a)]$  (in short  $g$  is a  $\Phi$ -morphism) <sup>(10)</sup>

<sup>(10)</sup> Note that  $M < N$  is not equivalent to he identity on  $|M|$  being a  $\Phi$ -morphism into  $N$ .

(B)  $K^\alpha(T, \Phi)$  is  $K^\alpha(\text{sub } \Phi)$  where  $K = K(T, \Phi)$ .

Definition 2.9.  $\mathcal{D}(T, \Phi) = \{tp(\bar{a}, \Phi, M) : \bar{a} \in M \in K^\omega(T, \Phi)\}$ .

Lemma 2.10.

- (1)  $K(T, \Phi)$  satisfies the assumptions before definition 2.6 when
  - (A)  $T \subseteq \forall \exists \Phi$ .
  - (B)  $\Phi$  is closed under subformulas except possibly the subformulas of a negation of an atomic subformula.
  - (C) For every atomic  $R(\bar{x})$ ,  $R(\bar{x}) \in \Phi$  or  $\neg R(\bar{x}) \in \Phi$ .
- (2) If in addition  $K(T, \Phi)$  has the JMP, then  $\mathcal{D} = \mathcal{D}(T, \Phi)$  satisfies the assumption from § 1, which corresponds to kind IV.
- (3) Assumption III is satisfied in  $\mathcal{D}(T, \Phi)$  for  $\exists \wedge \Phi$ .
- (4) If  $K(T, \Phi)$  has the amalgamation property,  $T$  universal,  $\Phi_{qt} = \Phi$ , then in (3) we can replace  $\exists \wedge \Phi$  by  $\Phi_a$ , and assumption II is satisfied.
- (5) For a universal theory with the JEP,  $\mathcal{D}^*(T) = \mathcal{D}(T, \Phi_{qt})$ :

$T^\omega = K^\omega(T, \Phi_{qt})$ . If  $T \subseteq \forall \exists \Phi_{qt}$ ,  $T_1, T_2$  have the same universal consequences, then for each  $\alpha > 0$ ,  $K^\alpha(T_1, \Phi_{qt}) = K^\alpha(T_2, \Phi_{qt})$ . <sup>(11)</sup>

Remarks: Having proved that kinds II, III satisfy the appropriate assumptions, we will now prove conversely that any example (of kind IV) which satisfies assumption II or III is essentially of kind II or III. In 2.11 by changing the language, we get to «infinite positive forcing», and in 2.12 by also adding more elements we get to «infinite forcing».

Theorem 2.11. In assumption III we can restrict ourselves to  $\mathcal{D}(T, \Phi_a)$  for a universal-negative  $T$  with the JMP,  $K(T^\omega, \Phi_a)$  having the amalgamation property. More exactly, let  $\mathcal{E}, \Phi$  be as in assumption III, and let

<sup>(11)</sup> We can prove it as before, or add names for the appropriate formulas.

- (i)  $L^* = \{R_{\varphi(\bar{x})} : \varphi(\bar{x}) \in \Phi\}$  where  $R_{\varphi(\bar{x})}$  is a predicate with  $\ell(\bar{x})$ -places and  $R_{x_1, \dots, x_l, Y_1}$  is  $x_1 = Y_1$ .
  - (ii)  $\mathcal{C}^* = (|\mathcal{C}|, \dots, R_{\varphi}, \dots)$  where  $R_{\varphi}^{\mathcal{C}^*} = \{\bar{a} \in \mathcal{C} : \mathcal{C} \models \varphi[\bar{a}]\}$
  - (iii)  $T^*$  is the negative universal theory of  $\mathcal{C}^*$ .
- Then
- (1)  $\mathcal{C}^*$  is  $\mathcal{D}(T^*, \Phi_a)$ -homogeneous,  $K(T^*, \Phi_a)$  has the JMP.
  - (2)  $\text{tp}(\bar{a}, \emptyset, \mathcal{C}) = \text{tp}(\bar{b}, \emptyset, \mathcal{C})$  iff  $\text{tp}(\bar{a}, \emptyset, \mathcal{C}^*) = \text{tp}(\bar{b}, \emptyset, \mathcal{C}^*)$ .
  - (3)  $A \subseteq |\mathcal{C}|$  is the universe of a  $(\mathcal{D}, \lambda)$ -homogeneous elementary submodel of  $\mathcal{C}$ , iff it is the universe of a  $(\mathcal{D}(T^*, \Phi_a, \lambda)$ -homogeneous elementary submodel of  $\mathcal{C}^*$ .
  - (4)  $\mathcal{D}, \Phi$  satisfy assumption II iff  $\mathcal{D}(T^*, \Phi_a), \Phi_a$  satisfy assumption II.
  - (5) If  $\Phi = \exists \wedge \Phi_1$ , we can replace  $\mathcal{C}^*$  by  $\mathcal{C}_1^* = (\mathcal{C}, \dots, R_{\varphi}, \dots)$ .

... $\}_{\varphi \in \Phi_1}$ . If, in addition,  $\mathcal{D} = \mathcal{D}(T, \Phi_1)$  as in lemma 2.10 (1), then  $\mathcal{C} \upharpoonright A < \exists \wedge \Phi_1 \mathcal{C}$  iff  $\mathcal{C}_1^* \upharpoonright A < \mathcal{C}_1^*$  iff  $\mathcal{C}^* \upharpoonright A < \mathcal{C}^*$ .

**Theorem 2.12.** In 2.10 (1) the case « $T$  universal,  $\Phi = \Phi_a$ » is a particular case of « $T$  universal,  $\Phi = \Phi_{qt}$ », except that equality is not standard. More exactly, let  $T$  be universal,  $\Phi = \Phi_a$ ,  $\mathcal{D} = \mathcal{D}(T, \Phi)$ ,  $\mathcal{C}$  is  $(\mathcal{D}, \lambda_0)$ -homogeneous ( $\lambda_0 = \|\mathcal{C}\|$ ). We define  $\mathcal{C}^+$  as follows:

- (i)  $|\mathcal{C}^+| = |\mathcal{C}| \times \lambda_0 \cup \{\langle \bar{a}, R, \bar{\alpha} \rangle : R(\bar{x}) \text{ is an atomic formula, } \mathcal{C} \models R[\bar{a}], \bar{\alpha} \in \ell(\bar{a})_{\lambda_0} \cup \lambda_0 \text{ (remember } \lambda_0 = \{ \alpha : \alpha < \lambda_0 \})\}$
- (ii)  $P_{\mathcal{C}^+} = |\mathcal{C}| \times \lambda_0$  (a one place relation)
- (iii)  $P_{\mathcal{R}}^{\mathcal{C}^+} = \{\langle \bar{a}, R, \bar{\alpha} \rangle : \mathcal{C} \models R[\bar{a}], \bar{\alpha} \in \ell(\bar{a})_{\lambda_0}\}$  (a one place relation)

- (iv)  $Q_{\mathcal{R}}^{\mathcal{C}^+} = \{\langle \langle \bar{a}_0, \alpha_0 \rangle, \dots, \langle \bar{a}_{n-1}, \alpha_{n-1} \rangle, \langle \bar{a}_0, \dots, \bar{a}_{n-1} \rangle, R, \bar{\alpha} \rangle : \alpha_0 < \lambda_0, \mathcal{C} \models R[\bar{a}_0, \dots, \bar{a}_{n-1}], \bar{\alpha} = \langle \alpha_0, \dots, \alpha_{n-1} \rangle\}$
- (v) let  $T^+$  be the universal theory of  $\mathcal{C}^+$ .

Then (1)  $\mathcal{C}^+$  is  $(\mathcal{D}(T^+, \Phi_{qt}), \lambda_0)$ -homogeneous  
 (2) for  $\bar{x}, \bar{y} \in |\mathcal{C}|$ ,  $\text{tp}(\bar{x}, \emptyset, \mathcal{C}) = \text{tp}(\bar{y}, \emptyset, \mathcal{C})$  iff  $\text{tp}(\bar{x}^*, \emptyset, \mathcal{C}^+) = \text{tp}(\bar{y}^*, \emptyset, \mathcal{C}^+)$  where  $\bar{x}^* = \langle \bar{x}, \alpha \rangle$ , and  $\langle \bar{a}_0, \dots \rangle^* = \langle \bar{a}_0^*, \dots \rangle$ .  
 (3) for  $A \subseteq \mathcal{C}^+$ , the following are equivalent  
 (a)  $\mathcal{C}^+ \upharpoonright A < \mathcal{C}^+$   
 (b)  $A' = \{a : \langle a, \alpha \rangle \in A \text{ for some } \alpha < \lambda_0, \text{ and } a \in \mathcal{C}\}$  is the universe of an elementary submodel of  $\mathcal{C}$  and if  $\bar{a}_1^* = \langle \bar{a}_1, \alpha_1 \rangle \in A$ ,  $\mathcal{C} \models R[\bar{a}_0, \dots]$  then  $\langle \langle \bar{a}_0, \dots \rangle, R, \bar{\alpha} \rangle \in A$  for some  $\bar{\alpha}$  and  $\langle \langle \bar{a}_0, \dots \rangle, R, \bar{\alpha} \rangle \in A$  implies  $\bar{a}_0, \dots \in A'$ .  
 (4) In (3) we can replace  $\langle by \langle \epsilon$  in (a) and  $\langle \epsilon_p$  in (b).

**Remark:** Hence we can give examples for  $K(T, \Phi_a)$  instead  $K(T, \Phi_{qt})$ , and almost always the relevant properties are preserved (except e.g. categoricity).

**Remark:** Normed spaces can be treated in this context. Let  $T_{NS}$  say that under  $+$  the model is an abelian group, and letting  $R_a^b(x)$  to mean  $a \leq \|x\| \leq b$ , and  $F_a(x)$  be  $ax$  (scalar multiplication) ( $a, b$  real numbers) that the natural conditions hold (e.g.  $R_a^b(x) \wedge R_c^d(y) \rightarrow R_{a+b}^{c+d}(x+y)$  and  $[a, b] = [a_1, b_1] \cup [a_2, b_2]$  implies  $R_a^b(x) \equiv R_{b_1}^{a_1}(x) \vee R_{b_2}^{a_2}(x)$ ). Then the normed spaces are essentially  $K^0(T_{NS}, \Phi_a)$ . For the class of  $L_p$ -spaces we get stability (by an unpublished result of Krivine), and «forgetting completeness», categoricity for Hilbert space. However, for Banach spaces, we should look only at, essentially, algebraically closed models, and replace cardinality by

density character; and then for  $L_p$ -spaces we get  $\aleph_0$ -stability. Somewhat more complicated is the case of injective hull. Those variations will be dealt with in Abramsky Ph. D. thesis. Notice that  $(\mathcal{D}(T_{NS}, \Phi_a), \aleph_1)$ -homogeneous models are Banach spaces.

*Remark.* Note that if  $T, \Phi$  ( $\ell = 1, 2$ ) satisfies the hypothesis of 2.10 (1), (2), and  $\Phi_1 \subseteq \Phi_2$ , then: if  $\mathcal{D}(T, \Phi_2)$  is stable in  $\lambda$ , then  $\mathcal{D}(T, \Phi_1)$  is stable in  $\lambda$ . So if (every completion of  $T$ ) is  $\lambda$ -stable,  $\mathcal{D}(T, \Phi_1)$  is  $\lambda$ -stable.

§ 3. Ranks and Stability

We work here in the context of assumption III, so  $L, \mathcal{E}, \mathcal{D}, \Phi$  are given and are fixed.

We let  $\varphi(\bar{x}; \bar{y})$  denote a pair  $\langle \varphi_0(\bar{x}; \bar{y}), \varphi_1(\bar{x}; \bar{y}) \rangle$  where  $\varphi_0, \varphi_1 \in \Phi$  are contradictory. Let  $\varphi(\bar{x}; \bar{y}) = \varphi(\bar{x}; \bar{y})$  for  $\ell = 0, 1$  and  $M \models \varphi[a, b]$  mean  $M \models \varphi_0[a, b]$ .

Let  $\Delta$  denote a set of pairs  $\varphi$ . A  $\Delta$ - $m$ -type is a (onsistent) set of formulas  $\varphi(\bar{x}, \bar{a})$ , ( $t \in \{0, 1\}, \bar{x} = \langle x_0, \dots, x_{m-1} \rangle$ .) Let  $\Phi^1 = \{\varphi : \varphi \in \Phi^1 \text{ contradictory, } \ell = 0, 1\}^{(12)}$ .

*Definition 3.1.* We define the rank  $R^m(p, \Delta, \lambda)$  as an ordinal or  $\infty$  or  $-1$ , where  $p$  is an  $m$ -type (or a set of  $m$ -formulas), and  $\Delta$  a set of pairs  $\varphi$  and  $\lambda \geq 2$  is a cardinal or  $\infty$ . We stipulate  $-1 < \alpha < \infty$  for any ordinal  $\alpha$ , and define the rank as follows:

- (i)  $R^m(p, \Delta, \lambda) \geq 0$  iff  $p$  is consistent, or equivalently finitely satisfiable in  $\mathcal{E}$ .
- (ii)  $R^m(p, \Delta, \lambda) \geq \delta$  ( $\delta$  a limit ordinal) iff for any  $\alpha < \delta$   $R^m(p, \Delta, \lambda) \geq \alpha$ .
- (iii)  $R^m(p, \Delta, \lambda) \geq \alpha + 1$  iff for any finite  $q \subseteq p$  and any  $\mu < \lambda$  there are  $\Delta$ - $m$ -types  $r_j$  ( $j \leq \mu$ ) such that:
  - (a)  $R^m(q \cup r_j, \Delta, \lambda) \geq \alpha$

<sup>(12)</sup> For kind II there is no need for  $\Phi^1$ , and we do not distinguish strictly between it and  $\Phi$ .

and (b) the  $r_j$  are pairwise explicitly contradictory, that is for any distinct  $i, j \leq \mu$  there is  $\varphi^i \in r_i$  such that  $\varphi^i \vdash \neg \varphi^j$ .

*Remarks.* (1)  $j \leq \mu$  occurs in (iii) (rather than  $j < \mu$ ) so that  $\lambda = 2$  (and not  $\lambda = 3$ ) is the first interesting case.

(2) The most important cases for  $\Delta$  are  $\underline{\Delta} = \overline{\Phi}$  (all possible pairs) and  $\underline{\Delta}$  finite or with one pair (which are essentially equivalent). The latter case is interesting because the splitting types  $r_j$  will consist only of instances (using different parameters) of finitely many formulas of  $\Phi$ , and hence we can apply compactness arguments.

(3) The most interesting cases for  $\lambda$  in what follows will be  $\lambda = 2$  and  $\lambda = \infty$ . When we assume Kind II (section 6, on forking),  $\lambda = \aleph_0$  is interesting too. Notice that in these cases the definition is absolute.

*Lemma 3.2.* (1) If  $p_1 \subseteq p_2$  [or even  $p_2 \vdash p_1$ ; i.e. for any finite  $q_1 \subseteq p_1$  there is a finite  $q_2 \subseteq p_2$  such that  $\mathcal{E} \models (\forall \bar{x}) (\bigwedge q_2 \rightarrow \bigwedge q_1)$ ] and  $\underline{\Delta}_1 \supseteq \underline{\Delta}_2$  and  $\lambda_1 \leq \lambda_2$ , then  $R^m(p_1, \underline{\Delta}_1, \lambda_1) \geq (R^m(p_2, \underline{\Delta}_2, \lambda_2))$ .

(2) For every  $p, \Delta, \lambda$  there is a finite  $q \subseteq p$  such that  $R^m(q, \Delta, \lambda) = R^m(p, \Delta, \lambda)$ .

*Proof.* (1) Check the definition.

(2) If  $R^m(p, \Delta, \lambda) = \infty$ , choose  $q = \emptyset$ . If  $R^m(p, \Delta, \lambda) = -1$ ,  $p$  is inconsistent, so choose some finite inconsistent  $q \subseteq p$ . So let  $R^m(p, \Delta, \lambda) = \alpha < \infty$ ; then  $R^m(p, \Delta, \lambda) \not\geq \alpha + 1$ , hence by the definition there is a finite  $q \subseteq p$  for which there are no suitable  $r_j$ 's. As  $q$  is a finite subset of  $p$ , also  $R^m(q, \Delta, \lambda) \not\geq \alpha + 1$ ; but, by (1),  $R^m(q, \Delta, \lambda) \geq R^m(p, \Delta, \lambda)$ , so the conclusion follows.

We note the following trivial but important claim:

*Claim 3.3.* If  $R^m(p, \underline{\Delta}, 2) = \alpha < \infty$ , then there is no  $\varphi \in \underline{\Delta}$  and  $\bar{a}$  such that  $R^m(p \cup \{\varphi(\bar{x}; \bar{a})\}, \underline{\Delta}, 2) = \alpha$  for  $t = 0, 1$ .

*Remark.* This claim explains why the rank for  $\lambda = 2$  is important. The extension property (i.e. for each  $p$  which is realized and  $A$ , there is a complete type  $q$  over  $A$  such that  $R^m(p, \underline{\Delta}, \lambda) = R^m(p \cup q, \underline{\Delta}, \lambda)$ ) and  $p \cup q$  is realized) fails in general, but it holds for  $\underline{\Delta}$  finite,  $\lambda = \infty$ , or (as noted by Hinkis) for  $\lambda \geq \aleph_0$ ,  $p$  a  $\Phi$ - $m$ -type, when  $\mathcal{D}, \Phi$  satisfy assumption II. If we

waive the demand « $p \cup q$  is realized in  $\mathfrak{C}$ », there will be no problem, but not much interest too. Notice that for  $\Phi$ -types, being consistent is equivalent to being realized.

*Remark.* There are quite accurate results specifying when  $R^m(p, \Delta, \lambda) \geq \alpha \Rightarrow R^m(p, \Delta, \lambda) = \infty$  and when  $R^m(p, \Delta, \lambda) = R^m(p, \Delta, \mu)$ .

*Lemma 3.4.* For every finite  $p$ ,  $R^m(p, \varphi, 2) \geq n$  iff there are  $\bar{\alpha} (\eta \in^{n >} 2)$  such that for any  $\eta \in 2$ ,  $p \cup \{\varphi(\bar{x}, \bar{\alpha}, \eta \upharpoonright \ell) : \ell < n\}$  is consistent (notice we write  $\varphi$  for  $\{\varphi\}$ ).

*Proof.*  $\Leftarrow$ : let  $p_\eta = p \cup \{\varphi(\bar{x}, \bar{\alpha}, \eta \upharpoonright \ell) : \ell < n(\eta)\}$  and now prove that  $R^m(p_\eta, \varphi, 2) \geq n - \ell(\eta)$  by downward induction on  $\ell(\eta)$ .

$\Rightarrow$ : merely define  $\bar{\alpha}_\eta$  by induction on  $\ell(\eta)$  such that  $\eta \in 2$  implies  $R^m(p_\eta, \varphi, 2) \geq n - \ell(\eta)$ .

*Remark.* The requirement that  $p$  be finite can be eliminated by a compactness argument when  $p$  is a  $\Phi$ - $m$ -type.

*Lemma 3.5.* If  $R^m(\emptyset, \varphi, 2) \geq \omega$ , then  $\mathcal{D}$  is unstable (in every cardinal  $\lambda$ ).

*Proof.* Let  $\mu = \min\{\mu : 2^\mu > \lambda\}$ , so  $\sum_{\alpha < \mu} 2^{|\alpha|} \leq \lambda$ .

Let  $\Gamma_\mu = \{\varphi(\bar{x}; \bar{y} \upharpoonright_\eta \upharpoonright_\alpha) : \eta \in 2, \alpha < \mu\}$ . Now  $\Gamma_\mu$  is finitely satisfiable in  $\mathfrak{C}$  as it is sufficient to check each  $\Gamma_n$ , which is consistent by the previous lemma and the hypothesis. <sup>(\*)</sup> Let

$A = \cup \{\bar{y}_\eta : \eta \in^{> \mu} 2\}$ , and then the  $\bar{x}_\eta (\eta \in 2)$  realize differ-

<sup>(\*)</sup> As  $\Gamma$  is a set of  $\Phi$ -formulas, some assignment satisfies it.

ent types over  $A$  (in  $\mathfrak{C}$ ). Now  $|A| \leq \aleph_\alpha \sum_{\alpha < \mu} 2^{|\alpha|} \leq \lambda <$

$$2^\mu \leq |\{tp(\bar{x}, A) : \eta \in 2\}| \leq |S^m_{\mathcal{D}}(A)|$$

*Theorem 3.6.* The following are equivalent:

- (1)  $\mathcal{D}$  is stable,
- (2)  $\mathcal{D}$  is stable in every  $\lambda = \lambda^{< L}$ ,
- (3) For every  $\varphi$ ,  $R^m(\emptyset, \varphi, 2) < \omega$ ,
- (4) There are no  $\varphi$  and  $\bar{\alpha}_n$  such that  $\models \varphi[\bar{\alpha}_n, \bar{\alpha}_n]$  where  $\text{if } (n < m) = \begin{cases} 0 & \text{when } n < m \\ 1 & \text{otherwise.} \end{cases}$

*Remark.* Only after this theorem stability makes sense: as by (1), (2) it is not an accident (i.e. «a priori») it is possible that  $\mathcal{D}$  is stable just in one cardinal, and this will not give enough information), (3) enables us to prove theorems on stable  $\mathcal{D}$ , and (4) on unstable  $\mathcal{D}$ .

*Proof.* (1)  $\Rightarrow$  (4)  $\Rightarrow$  (2) : by 1.9.

(2)  $\Rightarrow$  (1) : trivial

(1)  $\Rightarrow$  (3) : by the previous lemma

(3)  $\Rightarrow$  (2) : Suppose  $|A| \leq \lambda = \lambda^{< L}$ . For each  $p \in S^m_{\mathcal{D}}(A)$

and  $\varphi$  choose a finite  $p \subseteq p$  such that  $R^m(p, \varphi, 2) = R^m(p, \varphi, 2)$ .

Let  $p^* = \cup_{\varphi} p$ , so  $|p^*| \leq |L|$ . Clearly for every  $\varphi$ ,  $R^m(p, \varphi, 2) = R^m(p^*, \varphi, 2)$ .

Now suppose  $p, q \in S^m_{\mathcal{D}}(A)$ ,  $p \neq q$ , then there is  $\bar{a} \in A$  such

that  $p \upharpoonright \bar{a} \neq q \upharpoonright \bar{a}$ ; hence  $p_1 \neq q_1$  where  $p_1 = \{\varphi(\bar{x}, \bar{y}) : \varphi(\bar{x}, \bar{a}) \in p\}$ ,  $q_1 = \{\varphi(\bar{x}, \bar{y}) : \varphi(\bar{x}, \bar{a}) \in q\}$ . So there are contradictory  $\varphi \in \Phi$  such that  $\varphi_0(\bar{x}, \bar{y}) \in p_1$ ,  $\varphi_1(\bar{x}, \bar{y}) \in p_2$ . So letting

$\varphi = \langle \varphi_0(\bar{x}, \bar{y}); \varphi_1(\bar{x}, \bar{y}) \rangle$ ,  $\varphi(x, \bar{a})^0 \in p$ ,  $\varphi(x, \bar{a})^1 \in q$ . Hence, by

There is an axiomatic way (see [S1] ch. IV) to describe the construction of a model M over a set A, by adding step by step, each time one element realizing a type of a specific sort (usually isolated in a proper sense).

Tractable interesting cases here are e.g. the e.c. models of T (T from § 2), the infinite generic models, the (D<sup>0</sup>(T), λ)-homogeneous models, the λ-∃<sub>n</sub>-compact structures (i.e. the set of models M, such that any ∃<sub>n</sub>-type of cardinality < λ over M, is realized in M). To prove existence, uniqueness etc., we have to assume usually, some stability assumptions.

This axiomatic treatment not only includes all previous cases and in it we can prove the known results (even the uniqueness of the constructed model) and see what hypothesis is needed for which theorem, but also shows that we can replace isolation by e.g. definability (e.g. the anti-prime models). See remarks to 6.10 ("v").

Definition 4.2.

Let p be a type over A. We say that a formula φ(x; b) isolates p if φ(x; b) ∈ p and ⊨ (∀ x) ((φ(x; b) → ψ) for all ψ ∈ p. We say p is isolated if some formula isolates it.

Lemma 4.3.

Suppose D is totally transcendental. Then, for each Ψ (\*)<sub>ψ</sub> for every B, b̄ ∈ B and φ ∈ L, if ⊨ (∃ x) φ(x, b̄) then there is φ<sub>1</sub>(x, b̄<sub>1</sub>) (φ<sub>1</sub> ∈ Φ, b̄<sub>1</sub> ∈ B) and a\* such that ⊨ (∃ x)[φ(x, b̄) ∧ φ<sub>1</sub>(x, b̄<sub>1</sub>)] and φ(x, b̄) ∧ φ<sub>1</sub>(x, b̄<sub>1</sub>) ⊢ tp<sub>Φ</sub>(a\*, B).

Proof.

Choose φ<sub>1</sub>(x, b̄<sub>1</sub>), a Φ-formula, such that

(14) Added in proof (June 76): This can be applied also to the following. Let M be an e.c. group such that: if H ⊆ L, K are finitely generated subgroups of M, the free product with amalgamation L \*<sub>H</sub> K is a subgroup of M. Let N be a group, all whose finitely generated subgroups "appear" in M. We can find a "canonical" (e.g. unique) closure of N to a group L<sup>∞, (10)</sup>-equivalent to M, by the above method.

claim 3.3, it is impossible that p\* = q\*. Hence |S<sup>m</sup><sub>D</sub>(A)| ≤ |{p\* : p ∈ S<sup>m</sup><sub>D</sub>(A)}| ≤ |A|<sup>|L|</sup> ≤ λ<sup>|L|</sup> = λ.

We can prove similarly

Theorem 3.7. The following conditions are equivalent:

- (1) D is stable in every λ ≥ 2<sup>|L|</sup>,
(2) D is stable in some λ, λ<sup>ℵ<sub>0</sub></sup> > λ,
(3) R<sup>m</sup>(∅, Φ, ∞) < ∞.

If D satisfies any of the above, it is called superstable.

Theorem 3.8. The following conditions are equivalent when

- |L| < 2<sup>ℵ<sub>0</sub></sup>,
(1) D is stable in every λ ≥ |L|,
(2) D is stable in some λ < 2<sup>ℵ<sub>0</sub></sup>,
(3) R<sup>m</sup>(∅, Φ, 2) < ∞.

If D satisfies (3), it is called totally transcendental.

§ 4. Prime models

We again work in the context of assumption III.

Définition 4.1.

Let K be a class of models M ⊆ C.

We call M K-prime over A if

- (1) A ⊆ M ⊆ K
(2) If A ⊆ M' ∈ K then there is an embedding f : M → M' such that f ↗ A is the identity and f is elementary in C (that is can be extended to an automorphism of C).

Remark

The natural way to build a prime model over A is to add to it elements a<sub>i</sub> one-by-one, so that each a<sub>i</sub> realizes over A<sub>i</sub> = A ∪ {a<sub>j</sub> : j < i} a type which is realized in every M, A<sub>i</sub> ⊆ M ∈ K. In the tractable cases this will be an isolated type.

(i)  $\models (\exists x) [\varphi(x, \bar{b}) \wedge \varphi_1(x, \bar{b}_1)]$   
(ii) relative to (i),  $\alpha = R^1(\varphi_1(x, \bar{b}_1), \Phi, 2)$  is minimal.  
By 3.8  $\alpha < \infty$ , and by 3.3  $\varphi(x, \bar{b}) \wedge \varphi_1(x, \bar{b}_1)$  isolates some type  $tp_\Phi(a^*, B)$ .

**Definition 4.4.**  
Let  $A \subseteq B$ . Let  $\Psi \subseteq L$  be closed under subformulas,  $\Psi = \exists \Psi$ .  
By a  $\Psi$ -isolating sequence of  $B$  over  $A$ , we mean a sequence  $\langle \langle a_j, \varphi_j(x, \bar{b}_j) \rangle : j < \alpha \rangle$  such that  $B = A \cup \{a_j : j < \alpha\}$ ,  $b_j \subseteq A_j = A \cup \cup_{i < j} a_i$  and  $\varphi_j(x, \bar{b}_j)$  is a  $\Psi$ -formula such that  $\varphi_j(x, \bar{b}_j) \vdash tp_\Phi(a_j, A_j)$  and  $\models \varphi_j[a_j, \bar{b}_j]$  (equivalently  $\varphi_j(x, \bar{b}_j)$  isolates  $tp_\Phi(a_j, A_j)$ , since  $tp_\Phi(a_j, A_j)$  determines  $tp(a_j, A_j)$ ).

We say  $M$  is strictly  $\Psi$ -prime over  $A$  if  $M$  has a  $\Psi$ -isolating sequence over  $A$  and  $M$  is  $\Psi$ -closed: i.e. for every  $\varphi(\bar{x}) \in \Psi = \text{Sub } \Psi, M \models \varphi[\bar{a}] \Leftrightarrow \mathcal{C} \models \varphi[\bar{a}]$  for each  $\bar{a} \in M$ .  
 $M$  is  $\Psi$ -prime over  $B$ , if it is prime among the  $\Psi$ -closed models. If  $\Psi = \exists \wedge \Phi$  we omit it.

**Theorem 4.5.** (The existence of prime models)  
(1) If  $M$  is strictly  $\Psi$ -prime over  $A$ , then it is  $\Psi$ -prime over  $A$ , i.e. prime among the  $\Psi$ -closed  $M \subseteq \mathcal{C}$ .  
(2) If  $\mathcal{D}$  is totally transcendental, then over any  $A$  there is a strictly  $\Psi$ -prime model.  
(3) In (2), it suffices to assume that (\*) $_\Psi$  of 4.3 holds.

**Proof.**  
(1) If  $\langle \langle a_i, \varphi_i(x, \bar{b}_i) \rangle : \langle \alpha \rangle$  is a  $\Psi$ -isolating sequence of  $M$  over  $A$ ,  $N$  is  $\Psi$ -closed,  $A \subseteq N$ , then we define  $c_i \in N$ , by induction such that  $tp(\langle a_j : j \leq i \rangle, A) = tp(\langle c_j : j \leq i \rangle, A)$ . If we have defined for each  $j < i$ ,  $b_i = \bar{b}_i \wedge \langle a_{j(i), \dots} \rangle$ ,  $\bar{b}_i^1 \in A$ ,  $j(i), \ell < i$ , choose  $c_i$  to satisfy  $\varphi_i(x, \bar{b}_i^1, c_{j(i), \dots})$ .  
(2), (3) Use 4.3 (\*) repeatedly.

Lemma 4.6.

If  $L$  is countable and  $\Psi = \exists \wedge \Phi$  then (\*) $_\Psi$  in lemma 4.3 is actually equivalent to:  
(\*\*) $_\Psi$  over every  $A$  there is a  $\Psi$ -prime model of  $T$ .

Proof.

(\*) $_\Psi \Rightarrow$  (\*\*) $_\Psi$  essentially this is the previous theorem.  
(\*\*) $_\Psi \Rightarrow$  (\*) $_\Psi$  : suppose (\*) $_\Psi$ -fails and  $B, \varphi(x, \bar{a})$  ( $\bar{a} \in B, \varphi \in \Psi$ ) is a counterexample; that is no  $\varphi_1(x, \bar{b}_1)$  is suitable. So for every  $\varphi_1 \in \Psi, \bar{b}_1 \in B, \models \neg(\exists x) [\varphi_1(x, \bar{b}_1) \wedge \varphi(x, \bar{a})]$  or there are  $\vartheta_1, \vartheta_2 \in \Phi, \bar{a}_1, \bar{a}_2 \in B$  such that  $\models (\exists x) (\varphi(x, \bar{b}) \wedge \varphi_1(x, \bar{b}_1) \wedge \vartheta_1(x, \bar{a}_1))$  ( $\ell = 1, 2$ ) and  $\vartheta_\ell(x, \bar{a}_\ell)$  ( $\ell = 1, 2$ ) are contradictory.

By a Löwenheim-Skolem argument, we can assume  $B$  is countable.  
Suppose  $M$  is  $\Psi$ -prime over  $B$ , so  $M$  is countable.  
Let  $\Gamma = \{tp_\Phi(a^*, B) : a^* \in M, a^* \notin B, \models \varphi(a^*, \bar{b})\}$ .

It is easy to find a  $\Psi$ -closed model  $M_1, B \subseteq M_1$ , in which no  $p \in \Gamma$  is realized.

Lemma 4.7.

If  $M$  is strictly  $\Psi$ -prime over  $A$  then  
(1) For every  $\bar{c} \in M$ , there is an  $(\exists \wedge \Psi)$ -formula  $\psi(\bar{x}, \bar{a})$ , ( $\bar{a} \in A$ ), which isolates  $tp_\Phi(\bar{c}, A)$  and which  $\bar{c}$  satisfies.

(2) We can find a  $\Psi$ -isolating sequence of  $M$  over  $A$  of length  $\|M\|$ .

Remark.

For  $\mathcal{D}$  totally transcendental we can characterize the  $\Psi$ -prime models as in [S9], hence

Theorem 4.8.

For  $\mathcal{D}$  totally transcendental  $\Phi \subseteq \Psi = \exists \Psi$  closed under sub-formulas, the  $\Psi$ -closed prime model over  $A$  is unique over  $A$ . If  $L = \Phi = \Psi$ ,  $|L| = \aleph_0$ ,  $\mathcal{D}$  is stable (so  $\mathcal{D} = \mathcal{D}(T, L)$ ) then we can prove uniqueness when there is a strictly  $\Psi$ -prime model, but we have no characterization. But even for countable theories (i.e.  $\mathcal{D} = \mathcal{D}(T, L)$ ) which satisfy 4.3 (\*), we do not know if uniqueness necessarily holds (\*\*). We think the answer is negative, and the following proof may give us an idea how a counterexample might look. The situation is the same for uncountable  $T$ .

Aid.

The proof of the next theorem will use stationary sets. The main facts we need are as follows. Let  $\kappa$  be a fixed cardinal of cofinality  $> \aleph_0$ . Then  $X \subseteq \kappa$  is closed unbounded if  $\kappa = \sup X$  and for each  $\alpha < \kappa$ ,  $\alpha = \sup (X \cap \alpha)$  implies  $\alpha \in X$ . We call  $S \subseteq \kappa$  stationary if  $S$  meets every closed unbounded set.

Fact A: The intersection of two (or even  $\lambda < \text{cf}(\kappa)$ ) closed unbounded subsets of  $\kappa$ , is closed and unbounded.

Fact B: If  $X \subseteq \kappa$  is closed unbounded and  $F: X \rightarrow \kappa$  is normal (i.e. strictly increasing and continuous), then the set of fixed points of  $F$  (i.e.  $\{\alpha: F(\alpha) = \alpha\}$ ) is a closed unbounded set.

Fact C (Fodor's Theorem): If  $S$  is a stationary subset of  $\kappa$ , and  $F: S \rightarrow \kappa$  is regressive (i.e.  $\alpha \in S \Rightarrow F(\alpha) < \alpha$ ) and  $\kappa$  is regular (i.e.  $\text{cf } \kappa = \kappa$ ) then  $F$  is constant on some stationary  $S' \subseteq S$ . (See e.g. Juhász [J 1]). We use those facts freely.

The Uniqueness Theorem 4.9.

Suppose  $L$  is countable,  $\mathcal{D}$  is stable and  $\Phi = L$ . If over  $A$  there is a strictly prime model  $M$ , then any prime model over  $A$  is isomorphic to it.

(\*\*) Added in proof June 76: There are countable (complete) theories  $T$  which satisfy (\*), but the uniqueness fails.

Proof:

We prove it only when  $|A| \leq \aleph_1$ . If  $|A| < \aleph_0$ , then Vaught's argument works. Henceforth let  $|A| = \aleph_1$ , and let  $\langle \langle a_j, \varphi_j(x, \bar{a}_j) \rangle : j < \omega_1 \rangle$  be an isolating sequence for  $M$  over  $A$ , and let  $N$  be prime over  $A$ , so we can assume  $A \subseteq N \subseteq M$ . Let  $M_i$  ( $i < \omega_1$ ) be countable models such that  $\bigcup_{i < \omega_1} M_i = M$

There is a closed unbounded  $X \subseteq \omega_1$  such that for each  $i \in X$

- (i)  $M_i < M, M_i \cap N < N$
- (ii)  $M_i \cap \{a_j : j < \omega_1\} = \{a_j : j < i\}$
- (iii)  $j < i$  implies  $\bar{a}_j \in M_i$
- (iv) for each  $\bar{c} \in M_i$ ,  $q = \text{tp}(\bar{c}, A)$  is isolated by some  $\varphi(\bar{x}) \in q \upharpoonright (A \cap M_j)$ .

We define

$S_1 = \{i \in X : \text{for some } \bar{b} \in N, q = \text{tp}(\bar{b}, A \cup (N \cap M_j)) \text{ is not isolated}\}$ .

Case 1:  $S_1$  is not stationary.

We shall show that in this case,  $N$  is isomorphic to  $M$  over  $A$ . There is by assumption a closed unbounded set  $X_1$  disjoint from  $S_1$ . Hence there is a closed unbounded  $X^* \subseteq X_1 \cap X$ , such that for each  $i, j \in X^*$ ,  $i < j$  and  $\bar{b} \in N \cap M_j$ ,  $q = \text{tp}(\bar{b}, A \cup (N \cap M_j))$  is isolated by some  $\varphi(\bar{x}, \bar{c}) \in q$ ,  $\bar{c} \in (A \cap M_j) \cup (N \cap M_j)$ .

We shall define, by induction on  $i \in X^*$ , elementary maps  $F_i: A \cup (N \cap M_i) \rightarrow A \cup M_i$  such that:

- $F_i \upharpoonright A$  is the identity
- $i < j; i, j \in X^*$  implies that  $F_j$  extends  $F_i$ .

For  $i$  a limit (in  $X^*$ )  $F_i = \bigcup_{j < i} F_j$ . Now suppose  $i, j \in X^*$ ,  $j$  the successor of  $i$  in  $X^*$  and  $F_i$  is defined. We shall define  $F_j$ , just by Vaught's argument, because  $M_j \cap N, M_j$  are prime over  $(M_j \cap A) \cup (M_i \cap N), (M_j \cap A) \cup M_i$  resp.. For the first  $i \in X^*$  we define  $F_i$  similarly.

Clearly  $\bigcup_{i < \omega_1} F_i$  gives the required isomorphism from  $N$  onto  $M$  over  $A$ .

Case 2:  $S_1$  is stationary  
 We shall show that this implies a contradiction. For all  $i \in S_1$  there is  $\bar{b}_i \in N$  such that  $tp(\bar{b}_i, A \cup (N \cap M_i))$  is not isolated. However, since  $M$  has an L-isolating sequence, for each  $i \in S_1$   $q_i = tp(\bar{b}_i, A \cup M_i)$  is isolated by some  $\varphi_i(\bar{x}, \bar{a}_i, \bar{c}_i) \in q_i$  where  $\bar{a}_i \in A, \bar{c}_i \in M_i$ . By Fodor's theorem there are a stationary set  $S_2 \subseteq S_1$ , a formula  $\varphi$  and a sequence  $\bar{c}$  such that for each  $i \in S_2$ ,  $\varphi_i = \varphi$  and  $\bar{c}_i = \bar{c}$ .

Choose  $i_0 < i_1 < \dots < i_k < \dots$  ( $k < \omega$ ) in  $S_2$  such that  $a_{i_k}, b_{i_k} \in M_{i_{k+1}}$ .

For each  $n < \omega$  we can find  $\bar{c}_n^* \in N \cap M_{i_n}$  such that  $tp(\bar{c}_n^*, A \cup \bigcup_{l < n} \bar{b}_{i_l}) = tp(\bar{c}, A \cup \bigcup_{l < n} \bar{b}_{i_l})$  (if we replace  $A$  by  $A \cap M_{i_n}$ , this holds as  $M_{i_n}$  is strictly prime over  $A \cap M_{i_n}$  and the same  $\bar{c}_n^*$  is good by the choice of  $X$ , and as  $S_2 \subseteq S_1 \subseteq X$ ).

Now note.

First Fact: If  $k \leq n$  then  $\models \varphi[\bar{b}_{i_k}, \bar{a}_{i_k}, \bar{c}_n^*]$ .

This holds by the choice of  $\bar{c}_n^*$ .

Second Fact: If  $k > n$  then  $\models \neg \varphi[\bar{b}_{i_k}, \bar{a}_{i_k}, \bar{c}_n^*]$ . As in case 1 there is an elementary map  $F$ ,  $\text{Dom } F = A \cup M_{i_k}, F \upharpoonright A =$

the identity,  $F$  maps  $\bar{b}_{i_l}$  to  $\bar{b}_{i_l}$ ,  $\bar{c} (l < n)$  and  $F$  maps  $M_{i_k}$  onto  $M_{i_k}$ .

Hence  $\varphi(\bar{x}, \bar{a}_{i_k}, \bar{c}_n^*)$  isolates a complete type over  $A \cup M_{i_k}$ . So if  $\models \varphi[\bar{b}_{i_k}, \bar{a}_{i_k}, \bar{c}_n^*]$  then it isolates  $tp(\bar{b}_{i_k}, A \cup M_{i_k})$  hence  $tp(\bar{b}_{i_k},$

$A \cup (N \cap M_{i_k}))$ , contradicting the choice of  $\bar{b}_{i_k}$ .  
 From the two facts we infer that for  $k, n < \omega, \models \varphi[\bar{b}_{i_k}, \bar{a}_{i_k}, \bar{c}_n^*]$  iff  $k \leq n$ . We get a contradiction.  $(\mathcal{D})$  is stable by hypothesis, but we just proved it has the order property (see 3.6).

§ 5. Categoricity

We work in the context of assumption III. We prove the parallel of Morley's categoricity theorem which is quite similar to the standard case. We could have used 1.15 to shorten the proof. For simplicity, we restrict ourselves to countable  $L$ .

Definition 5.1.

- (1) Let  $K$  be a class of structures and  $\lambda$  a cardinal.  $K$  is  $\lambda$ -categorical (or categorical in  $\lambda$ ) iff any two structures in  $K$  of cardinality  $\lambda$  are isomorphic
- (2)  $\Psi$  is categorical in  $\lambda$  if the class of  $\Psi$ -closed models is categorical in  $\lambda$  (where  $\Phi \subseteq \Psi$ )

Theorem 5.2. Let  $L$  be countable. If  $\Psi$  is categorical in one  $\lambda > \aleph_0$ , then it is categorical in every  $\lambda > \aleph_0$ ; assuming  $\Psi =$  Sub  $\Psi = \exists \Psi$ .

Proof: we prove the theorem in three steps:

- Step 1:  $\mathcal{D}$  is  $\aleph_0$ -stable.
- Step 2: In every  $\lambda$  there is a  $(\mathcal{D}, \lambda)$ -homogeneous model
- Step 3: If there is a  $\Psi$ -closed model of cardinality  $\lambda > \aleph_0$  which is not  $(\mathcal{D}, \lambda)$ -homogeneous, then in every  $\mu > \aleph_0$  there is such a model.

We can conclude that every  $\Psi$ -closed model of cardinality  $\lambda > \aleph_0$  is  $(\mathcal{D}, \lambda)$ -homogeneous. By 1.6(3), it follows that  $\Psi$  is categorical in  $\lambda$ .

Step 1.

The idea is the following: first build one model of cardinality  $\aleph_1$  in which there are few types realized over each countable subset. If  $\mathcal{D}_T$  is  $\aleph_0$ -unstable, we build a second model of cardi-

nality  $\lambda$  in which there are many types realized over some countable set. This contradicts the  $\lambda$ -categoricity, so  $\mathcal{D}_T$  must be  $\aleph_0$ -stable.

We will first build the second model, which is easy: If  $\mathcal{D}$  is  $\aleph_0$ -unstable, this means there is some countable subset  $A$  of  $\mathcal{C}$  and  $\mathcal{C}$  realizes  $\geq \aleph_1$  different types over  $A$ . We may choose an elementary substructure  $M$  of  $\mathcal{C}$  satisfying:

$$\|M\| = \lambda \text{ and } M \text{ realizes } \geq \aleph_1 \text{ types over } A \subseteq |M|$$

Next we will build a model  $N$  of cardinality  $\lambda$  which realizes just countably many types over every countable subset of  $N$ . We use Ehrenfeucht-Mostowski models  $EM(I)$  generated by a sequence  $I$  of indiscernibles. We will review this notion in more detail after finishing the proof.

We expand  $\mathcal{C}$  to  $\mathcal{C}^s$  by adding Skolem-functions. Because  $K' = \{M \mid M \models \text{Th } \mathcal{C}^s, M \text{ is } \Psi\text{-closed}\}$  has models of arbitrarily large cardinality, it is possible to find a structure in  $K'$  generated by indiscernibles  $I$  (because  $K'$  is definable in  $L^{\aleph_0}$ )<sup>+</sup>  $\aleph_0$ .

this requires an argument based on the Erdős-Rado theorem and compactness). Replacing  $I$  by the ordinal  $\lambda$  and taking  $N$  to be the corresponding Ehrenfeucht-Mostowski model  $N = EM(I)$  we see that  $N$  has cardinality  $\lambda$ , and we claim:

(\*) for any countable subset  $A$  of  $N$ ,  $N$  realizes only countably many types over  $A$ .

The argument proving (\*) may be reconstructed from the observation that the ordered set  $\lambda$  realizes only countably many types over any countable subset. Now  $M, N$  cannot be isomorphic, contradicting the  $\lambda$ -categoricity. Thus  $M$  cannot exist, so  $\mathcal{D}_T$  is  $\aleph_0$ -stable.

*Review of Ehrenfeucht-Mostowski Models:*

Whenever  $A \subseteq \mathcal{C}^s$  we define the hull  $H(A)$  as the closure of  $A$  under Skolem functions in  $\mathcal{C}^s$ , considered as a submodel of

$\mathcal{C}$ . Then  $H(A)$  is determined by  $A$  and the types of  $n$ -tuples of elements of  $A$  in  $\mathcal{C}$ .

If  $I$  is a sequence of indiscernibles in  $\mathcal{C}^s$ , then we write  $EM(I)$  instead of  $H(I)$ .  $EM(I)$  can be reconstructed from

1. the order type of  $I$ ;
2. the types  $p_n$  of increasing  $n$ -tuples from  $I$  (there is one  $p_n$  for each  $n$ ). The set  $\{p_n : \text{all } n\}$  is called the character  $\Gamma$ .

If  $I$  is a sequence of indiscernibles in  $\mathcal{C}^s$  with character  $\Gamma$  and if  $J$  is any ordered set, we may construct an Ehrenfeucht-Mostowski model  $EM^1(J, \Gamma)$  generated by  $J$  as a sequence of indiscernibles of character  $\Gamma$ , and let  $EM(J, \Gamma)$  be its  $L$ -reduct. Obviously the models  $EM(J, \Gamma)$  are infinitely generic (Once indiscernibles have been constructed, they may be taken to lie in  $\mathcal{C}$ ).

Step 2.

Obvious by 1.13.

The main point is:

Step 3.

If  $\lambda_1, \lambda_2 > \aleph_0$  and we have a  $\Psi$ -closed non- $\mathcal{D}$ -homogeneous model  $M_2$  of cardinality  $\lambda_2$ , then we can find such a model in  $M_1$ .

*Proof:* This is a Lowenheim-Skolem theorem. We will shrink  $M_2$  to a model  $M_0$  and expand  $M_0$  to the desired model  $M_1$ . Fix any  $M_2$  of cardinality  $\lambda_2$  which is not  $(\mathcal{D}_T, \lambda_2)$ -homogeneous. Fix a subset  $A_2 \subseteq |M_2|$  of cardinality less than  $\lambda_2$  such that some type  $p_2$  in  $S_{\mathcal{D}}(A_2)$  is not realized in  $M_2$ .

*Going down:* We will make a countable structure  $M_0$  resembling  $M_2$ . By the  $\aleph_0$ -stability of  $\mathcal{D}$  and 1.10, 1.12, there is in  $M$  an indiscernible set  $I_2$  of cardinality  $(\aleph_0 + |A_2|)^+$  over  $A_2$ . Let  $M'_2$  be  $\Psi$ -prime over  $A_2 \cup I_2$ , and embed  $M'_2$  in  $M_2$ . In particular

$M'_2$  omits  $p_2$ . Thus:

(\*) There is no  $\Psi_1$ -formula  $\varphi$  with parameters in  $A_2 \cup I_2$  which isolates  $p_2$ . ( $\Psi_1 = \exists \wedge \Psi$ ).

That is:

(\*\*) for every  $\Psi_1$ -formula  $\varphi$  over  $A_2 \cup I_2$ , there is a formula  $\varphi \in \Phi$ ,  $\varphi' \in p_2$  and  $\varphi \wedge \varphi'$  is consistent.

We may choose countable sets  $A_0 \subseteq A_2$ ,  $I_0 \subseteq I_2$  such that the type  $p_0 = p_2 \upharpoonright A_0$  satisfies the analog of (\*\*), hence of (\*), and  $I_0$  is infinite. Of course,  $I_0$  is indiscernible over  $A_0$ . Let  $M_0$  be  $\Psi$ -prime over  $A_0 \cup I_0$ . Then  $M_0$  omits  $p_0$ . The next task is to expand  $M_0$  to a large model omitting  $p_0$ .

Going up: We assume that  $A_0, I_0, p_0$  are countable and satisfy

(\*) Let  $I_1$  be a set of indiscernibles over  $A_0$  such that  $I_0 \subseteq I_1$  and  $|I_1| = \lambda_1$ . Let  $M_1$  be  $\Psi$ -prime over  $A_0 \cup I_1$ . We still have

(\*\*)' There is no  $\Psi_1$ -formula  $\varphi$  with parameters in  $A_0 \cup I_1$  which isolates  $p_0$ .

Hence  $M_1$  omits  $p_0$ . Since  $A_0$  is countable,  $M_1$  is not even  $(\mathcal{D}, \aleph_1)$ -homogeneous. This proves the lemma, and completes the proof of the Los' Conjecture.

Remark.

To see that this theorem is interesting, we will give some examples of theories  $T$  such that  $E_T$  (the class of e-closed models of  $T$ ) is categorical in  $\aleph_1$  without being elementary. We want three such examples:

- 1.  $E_T$  categorical in  $\aleph_1$  and not in  $\aleph_0$ ;
- 2.  $E_T$  categorical in  $\aleph_0$  and not in  $\aleph_1$ ;
- 3.  $E_T$  categorical both in  $\aleph_0$  and  $\aleph_1$ .

Example 1. (Macintyre)

We consider models which consist of pairs of algebraically closed fields  $F_0 \subseteq F_1$ , of fixed characteristic. The existentially complete models are pairs  $(F_0, F_1)$  in which  $F_1$  has transcendence degree 1 over  $F_0$ . Clearly this is an example of type 1. It is also an example of a complete inductive non model-complete theory.

Example 2.

$T$  is the theory of undirected graphs without cycles. The exist

tentially complete models are the connected graphs without cycles having infinite branching at each point.

Example 3.

Adjoin to example 2 a function symbol  $f(x_1, x_2, x_3)$ . We want  $f(a_1, a_2, x)$  to be an automorphism  $f_{a_1, a_2}$  of our graph carrying  $a_1$  to  $a_2$ . Take as a further axiom:

$$(A) f_{a_2, a_3} f_{a_1, a_2} = f_{a_1, a_3}.$$

$K^1 = K^1(T, \Phi_{qt}) = E_T$  consists of connected graphs without cycles having constant infinite branching at each point. Because we have axiom (A), the function symbol  $f$  does not affect the number of isomorphism types. The computation of  $E_T$  requires some care.

### § 6. Forking

In this section we work in the context of assumption II, assume  $\mathcal{D}$  is stable, and for simplicity let  $\Phi = \Phi_{qt}$ , hence  $\text{tp}_{qt}(\bar{a}, A)$  determines  $\text{tp}(\bar{a}, A)$ ; so here a type will mean a q.f. type (though many theorems are true more generally). The important and positive notion will be «not forking over» which is closely related to «definable over», and sometimes plays a role similar to «isolated».

Forking was invented to deal with

A. A union of an increasing elementary chain of  $\kappa$ -saturated models is  $\kappa$ -saturated, when the cofinality of the length of the sequence is  $\geq \kappa(T)$ .

The difficult case is  $\kappa(T) < \kappa < |T|$ .

Another use of it is in proving the stability spectrum theorem. For 1.10 we can use also strong splitting, but it is essential for

B.  $\lambda(\mathcal{D})$  is  $|\mathcal{D}|$  or  $2^{\aleph_0}$ .

Another use is

- C. Constructing anti-prime models [S 1] (ch. IV). See remark to 6.10.
- D. Proving that if a countable T has only universal models in  $\aleph_1$ , then it is categorical in  $\aleph_1$  (affirming a conjecture of Keisler).

The relation of forking and ranks is determined in [S 1, III, § 4] and has applications to ranks.

If T is superstable, we can, in some cases, use ranks instead of forking (when p is complete over B,  $A \subseteq B$ ). This is natural, as «p does not fork over A» is equivalent to «p and p f A have the same rank  $R(-, L, \infty)$ »; and there are equivalent formulations.

All theorems here are from [S 1], most of them from ch. III. See also Lascar [Ls], and this volume.

The notion of «tp(a, B) does not fork over A,  $A \subseteq B$ » is similar to «the free algebras generated by A, B,  $A \cup \{a\}$  are freely amalgamated in  $\mathbb{C}$ », and many times in algebraic contexts this is what we get by explicating.

«Forking» was invented as a version of strong splitting (see [S 7]). However it can be developed only under more restricted assumptions.

*Definition 6.1.* A type p forks over A if there exists a set of indiscernibles  $I_j = \{\bar{a}_j^\alpha \mid \alpha < \omega\}$  over A and q.f. formulas  $\varphi_j$

$$\text{such that } p \vdash \bigvee_{j < n} [\varphi_j(\bar{x}, \bar{a}_0^j) \leftrightarrow \neg \varphi_j(\bar{x}, \bar{a}_j^j)].$$

When  $A = \emptyset$ , we omit it.

The notion is not interesting for unstable theories with this definition.

*Example:* Let T be the theory of an equivalence relation E;  $p = \{x E a\}$  does not fork if E has finitely many equivalence classes, otherwise yes.

*Theorem 6.2.*

- (1) If  $q \subseteq p$  (or even  $p \vdash q$ ) are types and p does not fork over A, q does not fork over A.
- (2) If p is over B and p does not fork over A, we can com-

plete p to a q.f. complete type over B which does not fork over A. (This we call the «extension property» of (not-forking) types).

(3) If p is over A, p does not fork over A. (Remember  $\mathcal{D}_T$  is stable).

*Proof of (3).* (as a sample proof)

Assume p forks over A. Then  $p \vdash \bigvee_{j < n} [\varphi_j(\bar{x}, \bar{a}_0^j) \leftrightarrow \neg \varphi_j(\bar{x}, \bar{a}_j^j)]$ .

Let  $\{\varphi_j : j \leq n\} \subseteq \Delta$ ,  $\Delta$  finite. By stability we have  $R^m(p, \Delta, \aleph_0) < \omega$ . Extend p to a q.f. complete type q over  $A \cup \bigcup_{m, j} \bar{a}_j^m$

of the same rank. So there is a j such that (w.l.o.g.)  $p \vdash$

$\varphi_j(\bar{x}_j, \bar{a}_0^j) \wedge \neg \varphi_j(\bar{x}_j, \bar{a}_j^j)$ . By stability again  $[R^m(p, \Delta, \aleph_0) < \omega$  is enough]  $\varphi_j$  divides  $I_j$  in two sets, one big and one small. W.l.o.g.

for most  $\bar{a} \in I_j$ ,  $\neg \varphi_j(\bar{x}, \bar{a}) \in p$ . Now for all  $m_0 < \omega$ ,  $p \cup \{\varphi_j(\bar{x}_j, \bar{a}_j^m) \mid m = m_0\}$

contradicts the definition of rank.

*Remark:* we need quantifier-free for completing p preserving the rank.

*Definition 6.3.*

$\varphi(\bar{x}, \bar{a})$  is almost over A iff  $\{\varphi(\bar{x}, F(\bar{a})) : F \text{ is an automorphism of } \mathbb{C} \text{ over } A\}$  has cardinality  $< |\mathbb{C}|$  (here this amounts to being finite).

*Examples:* 1. Algebraically closed fields,  $A = \emptyset$ .

$$x^2 + 17x - 3 = 0 \text{ is over } \emptyset$$

$$x^2 + \pi x - 3 = 0 \text{ is not almost over } \emptyset$$

$$x^2 + \sqrt{3}x - \sqrt{7} \text{ is almost over } \emptyset$$

(in fact it has 21 twinbrothers)

2. E is an equivalence relation with finitely many equivalence classes. Then  $x E a$  is almost over  $\emptyset$ .

Definition 6.4.

p is stationary over A iff

- (i) p does not fork over A and
- (ii) p has no two contradictory extensions which do not fork over A.

Definition 6.5.

The strong type  $stp_{qt}(\bar{a}, A) = \{\varphi(\bar{x}, \bar{b}) : \models \varphi(\bar{a}, \bar{b}) \text{ and } \varphi(\bar{x}, \bar{b}) \text{ is almost over } A \text{ and } \varphi \text{ is } qf\}$ .

(There is no cardinality problem in the definition: we can choose representatives for equivalent formulas in  $stp_{qt}(\bar{a}, A)$ , there are  $\leq |A| + |T|$  such equivalence classes). What we want to understand is: what are the possibilities of type extensions which do not fork; how much do we need for a type to be stationary?

Theorem 6.6. (The finite equivalence relation theorem)

A strong type  $stp_{qt}(\bar{a}, A)$  is stationary over A.

Intuitively this says: for each equivalence relation  $\varphi(x, y, \bar{b})$ ,  $b \in A$ , with finitely many equivalence classes,  $stp_{qt}(\bar{a}, A)$  says to which class it belongs i.e. to the class of  $\bar{a}$ .

Hence

Conclusion 6.7. If  $tp_{qt}(\bar{a}, B)$  does not fork over  $A \subseteq B$ ,  $\bar{b}, \bar{c} \in B$ ,  $stp_{qt}(\bar{b}, A) = stp_{qt}(\bar{c}, A)$ , then

$$tp_{qt}(\bar{a} \wedge \bar{b}, A) = tp_{qt}(\bar{a} \wedge \bar{c}, A).$$

Theorem 6.8. Let  $B \subseteq A$ , a sequence,  $p = tp_{qt}(\bar{a}, A)$ ; p does not fork over B iff  $stp_{qt}(\bar{a}, A)$  does not fork over B.

Proof ( $\Leftarrow$  easy) Suppose  $stp_{qt}(\bar{a}, A)$  does fork over B. As strong types are closed under conjunction some  $\varphi(\bar{x}, \bar{b}) \in stp(\bar{a}, A)$  forks over B.

Hence  $\varphi(\bar{x}, \bar{b}) \models \bigvee_{j \leq n} \varphi_j(x, \bar{a}_j) \Leftrightarrow \neg \varphi_1(x, \bar{a}_1)$  and  $\{\bar{a}_j \mid j < \omega\}$

indiscernible over B.

Let  $\varphi(\bar{x}, F_j(\bar{b}))$  ( $j < k < \omega$ ) be all «sisters» of  $\varphi(\bar{x}, \bar{b})$  up to equivalence, where  $F_j$  is an automorphism of  $\mathbb{C}$  and  $F_j \upharpoonright A = id_A$ .

Then  $p \vdash \bigvee_{i,j} \varphi(\bar{x}, F_j(\bar{b})) \vdash \bigvee_{i,j} [\varphi_1(x, F_j(\bar{a}_i)) \Leftrightarrow \neg \varphi_1(x, F_j(\bar{a}_i))]$

and  $\{F_i(\bar{a}_i) : i < \omega\}$  is indiscernible over B.

So  $tp(\bar{a}, A)$  forks over B, a contradiction.

Lemma 6.9. If p is a complete qf-m-type over A, it is stationary over  $B \subseteq A$  iff it is stationary over A and does not fork over B.

Remarks:

- (1) For complete types this amounts to being stationary.
- (2) If M is existentially closed,  $tp_{qt}(a, M)$  is stationary. Some important facts are summarized by

Theorem 6.10.

- (i) If  $B \subseteq A$ ,  $tp_{qt}(\bar{a} \wedge \bar{b}, A)$  does not fork over B, then  $tp_{qt}(\bar{b}, A)$  does not fork over B.
- (ii) (Symmetry)  $tp_{qt}(\bar{a}, A \cup \{b\})$  forks over A iff  $tp_{qt}(\bar{b}, A \cup \{\bar{a}\})$  forks over A.
- (iii) (Transitivity) Let  $C \subseteq B \subseteq A$ . If  $tp_{qt}(\bar{a}, A)$  does not fork over B and  $tp_{qt}(\bar{a}, B)$  does not fork over C, then  $tp_{qt}(\bar{a}, A)$  does not fork over C. (Also the converse holds).
- (iv) Let  $B \subseteq A$ . Then  $tp_{qt}(\bar{a} \wedge \bar{b}, A)$  does not fork over B iff  $tp_{qt}(\bar{a}, A)$  does not fork over B and  $tp_{qt}(\bar{b}, A \cup \{\bar{a}\})$  does not fork over  $B \cup \{\bar{a}\}$ .

Proof (i) trivial

(ii) Define by induction on  $i < \omega$  sequences  $\bar{a}_i, \bar{b}_i$ . For  $i = 0$ ,  $\bar{a}_i = \bar{a}, \bar{b}_i = \bar{b}$ ; for  $i > 0$ , choose  $\bar{a}_i \wedge \bar{b}_i$  such that  $stp_{qt}(\bar{a}_i \wedge \bar{b}_i, A \cup \bigcup_{j < i} \bar{a}_j \wedge \bar{b}_j)$  does not fork over A and extends  $stp_{qt}(\bar{a} \wedge \bar{b}, A)$ . Suppose  $tp_{qt}(\bar{a}, A \cup \bar{b})$  forks over A but  $tp_{qt}(\bar{b}, A \cup \bar{a})$  does not fork over A.

By 6.7 if  $j < i$  then  $tp_{qt}(\bar{a}_j \wedge \bar{b}_j, A) = tp_{qt}(\bar{a}_0 \wedge \bar{b}_0, A)$ .

By 6.6 if  $j \geq i$  then  $\bar{a}_j \wedge \bar{b}_j$  realizes

$$tp_{qt}(\bar{a}_i \wedge \bar{b}_i, A \cup \bigcup_{\alpha < i} \bar{a} \wedge \bar{b}).$$

Easily for all  $j < i$ ,  $tp_{qt}(\bar{a}_j \wedge \bar{b}_j, A) = tp_{qt}(\bar{a} \wedge \bar{b}, A)$ .

Similarly if  $j > i > 0$   $tp_{qt}(\bar{a}_j \wedge \bar{b}_j, A) = tp_{qt}(\bar{a}_j \wedge \bar{b}_0, A) = tp_{qt}(\bar{a}_0 \wedge \bar{b}_0, A) \neq tp_{qt}(\bar{a}_0 \wedge \bar{b}_0, A)$

(as  $tp_{qt}(\bar{a}, A \cup \{\bar{b}_0\})$  forks over  $A$  iff  $\ell = 0$ ).

This shows  $\mathcal{D}$  has the order property, contradiction.  
(iii) There is  $\bar{a}$  realizing  $stp_{qt}(\bar{a}, C)$  such that  $stp_{qt}(\bar{a}, A)$  does not fork over  $C$  (by 6.6, 6.8).

So  $stp_{qt}(\bar{a}, B)$ ,  $stp_{qt}(\bar{a}, C)$  extend  $stp_{qt}(\bar{a}, A)$  and do not fork over  $C$ ; hence they are equal. By trivial monotonicity properties  $stp_{qt}(\bar{a}, A)$  does not fork over  $B$ , so similarly  $stp_{qt}(\bar{a}, A) = stp_{qt}(\bar{a}, A)$ , so the latter too does not fork over  $C$ ; so we finish.

(iv)  $\Rightarrow$  : So suppose  $tp_{qt}(\bar{a} \wedge \bar{b}, A)$  does not fork over  $B$ . Then by (i)  $tp_{qt}(\bar{a}, A)$  does not fork over  $B$ . Let  $\bar{c} \in A$ , then  $tp_{qt}(\bar{a}, \bar{b}, B \cup \{\bar{c}\})$  does not fork over  $B$  (by monotonicity of forking) hence  $tp_{qt}(\bar{c}, B \cup \{\bar{a} \wedge \bar{b}\})$  does not fork over  $B$  (by (ii)), hence over  $B \cup \bar{a}$ , hence  $tp_{qt}(\bar{b}, B \cup \bar{a} \cup \bar{c})$  does not fork over  $B \cup \bar{a}$  (by (ii) again). As this holds for every  $\bar{c}$ ,  $tp_{qt}(\bar{b}, A \cup \bar{a})$  does not fork over  $B \cup \bar{a}$ .

$\Leftarrow$  : So assume  $tp_{qt}(\bar{a}, A)$ ,  $tp_{qt}(\bar{b}, A \cup \bar{a})$  does not fork over  $B$ ,  $B \cup \bar{a}$  respectively. Let  $\bar{b}_i (i < i_0)$  be a maximal list such that  $i \neq j \Rightarrow stp_{qt}(\bar{b}_i, B \cup \bar{a}) \neq stp_{qt}(\bar{b}_j, B \cup \bar{a})$ .

We can find (by a simple version of 6.2(2))  $\bar{b}_i, \bar{a}'$  such that  $stp_{qt}(\bar{a}' \wedge \dots \wedge \bar{b}_i \wedge \dots \wedge \bar{a})$  extends  $stp_{qt}(\bar{a}' \wedge \dots \wedge \bar{b}_i \wedge \dots, B)$  and

does not fork over  $B$  (i.e. every finite subtype satisfies this). Then w.l.o.g.  $\bar{a}' = \bar{a}$ , hence for some  $i$ ,  $stp_{qt}(\bar{b}_i, B \cup \bar{a}) =$

$stp_{qt}(\bar{b}_i, B \cup \bar{a})$ , and as  $stp_{qt}(\bar{b}_i, A \cup \bar{a})$  does not fork over  $B \cup \bar{a}$  (by the "only if part" of (iv)), they are equal, so  $stp_{qt}(\bar{a}' \wedge \bar{b}_i, A)$  does not fork over  $B$ .

Remark: If  $\mathcal{D}$  is stable, then for every  $A$  there is a model  $M$  (say existentially closed) with  $A \subseteq |M|$ , such that for every  $\bar{a} \in |M|$ ,  $tp_{qt}(\bar{a}, A)$  does not fork over some finite subset of  $A$ . If  $\mathcal{D}$  is superstable, the above is trivially true: every  $M$  does it. This helps us to show that for unsuperstable  $T$ ,  $\lambda = |T| + \aleph_1$ .

$I(\lambda, T) = 2^\lambda$ , and to build models with absolute indiscernibles. The facts from the theorem above fit well into an

axiomatic treatment of prime models. This is done in chapter IV of [S 1].

Definition 6.11.

$\kappa(\mathcal{D})$  is the first cardinal  $\kappa$  such that there are no  $A_i (i < \omega)$  with  $A_i \subseteq \neq A_j$  for  $i < j$ ) and  $tp_{qt}(\bar{a}, A_{i+1})$  forks over  $A_i$  (cf. the  $\kappa(\mathcal{D})$  in theorem 12).

Theorem 6.12. (i)  $\mathcal{D}$  is superstable iff  $\kappa(\mathcal{D}) = \aleph_0$ .

(ii) For stable  $\mathcal{D}$  and  $p$  over  $A$  there is a  $B \subseteq A$  with  $|B| < \kappa(T)$  s.t.  $p$  does not fork over  $B$ .

Forking is defined only for stable theories. For unstable theories there are troubles in defining it generally, at least it doesn't make sense. But if we have Skolem-functions, the following lemma may well serve as an alternative definition.

Lemma 6.13.  $p$  does not fork over  $M$  iff  $p$  is finitely satisfiable in  $M$  (though  $p$  is not necessarily over  $M$ ).

This lemma, as a definition for unstable theories, is useful in the following problem: how can we show that there are  $\lambda$ -universal but not  $\lambda^+$ -universal models for some unstable theory. If we are not interested in the sharp calculation of this  $\lambda$  (i.e. if  $\lambda_1 < \lambda_2$  we are happy with  $\lambda_1$ -universal but not  $\lambda_2$ -universal) an ultrapower construction with Erhenfeucht-Mostowski model will do. If we are interested in the sharp  $\lambda$ , we need forking as defined by the previous lemma.

In particular, for every  $T$  we have either

(i)  $T$  is categorical in all  $\lambda > |T|$  after adding  $2^{|T|}$  constants (and hence all models are saturated and hence universal),

or

(ii) for all  $\lambda, \mu, \mu \geq 2^\lambda, \lambda \geq 2^{|T|}$ ,  $T$  has a  $\lambda$ -universal but not  $\lambda^+$ -universal model of cardinality  $\mu$ .

From this one deduces, answering a question due to J. Keisler,

Theorem 6.14. If  $|T| = \aleph_0$  and every model of  $T$  of cardinality  $\aleph_1$  is universal then  $T$  is categorical in  $\aleph_1$ .

A natural question is whether restricting ourselves to kind II is merely incompetence.

*Lemma 6.15.* *There is a countable universal  $T$ , such that  $\mathcal{D} = \mathcal{D}(T, \Phi_a)$  is superstable and there is a countable  $A \subseteq \mathbb{C}$  such that for each  $a \in \mathbb{C}$ ,  $tp_a(a, A)$  splits strongly over  $\emptyset$ , i.e. there is an infinite indiscernible set  $I \subseteq A$ , and  $b, c \in I$  and  $q\bar{\varphi}$  such that  $\models \varphi[a, b] \wedge \neg \varphi[a, c]$ .*

*Remark:* By 2.12 we can find such an example for  $\mathcal{D} = \mathcal{D}(T, \Phi_{q\bar{\varphi}})$ , i.e. infinite generic models.

*Proof.* Let us define a model  $M$ . Its universe  $|M|$  is the set

of functions  $f$  such that (i)  $\text{Dom } f = [n, \infty] = \{m < \omega \mid n \leq m\}$ ,  
 Range  $f \subseteq \omega$   
 (ii)  $f$  is eventually zero

The relations are  $P_n = \{f : \text{Dom } f = [n, \infty]\}$

$R_n = \{ \langle f_1, f_2 \rangle : f_1, f_2 \in P_n, f_1 \upharpoonright [n+1, \infty] = f_2 \upharpoonright [n+1, \infty] \text{ and } f_1(n) \neq f_2(n) \}$

The functions are  $G_n$ , defined by  $G_n(f) = f \upharpoonright [n, \infty]$ .

Let  $T$  be the universal theory of  $M$ .

Now if  $\mathcal{D} = \mathcal{D}(T, \Phi_a)$ ,  $\mathbb{C} = \mathcal{D}(\mathcal{D}, \lambda_0)$ -homogeneous model, then for any  $a, b \in \mathbb{C}$  for some  $n, m$

$$G_n(a) = G_m(b).$$

The rest is left to the reader.

§ 7. Algebra: non-structure theorems

The problem here is to show that various algebraic theories are unstable etc... In § 7, § 8, for simplicity we concentrate on assumption I. Notice that by [S1], e.g.

Theorem 7.1.

(1) *If  $T$  is not superstable,  $\lambda \geq \aleph_1 + |T|$ , then  $I(\lambda, T) = 2$ . In most cases this is true for pseudo-elementary classes, and*

we get  $2^\lambda$  models, no one elementarily embedded into another. (We can omit "elementarily" e.g. if the formulas showing unsuperstability are q.f..)

(2) *If  $T$  is countable (for simplicity) but not  $\aleph_0$ -stable, then for each  $\lambda > \aleph_0$ ,  $T$  has  $\min\{2^\lambda, 2^{\aleph_0}\}$  non isomorphic models, no one elementarily embedded into another.*

(3) *If  $T$  is  $\aleph_0$ -stable not  $\aleph_1$ -categorical then  $I(\aleph_\alpha, T) \geq |\alpha + 1|$*

(4) (Baldwin [Ba]) *If  $T$  is categorical,  $\alpha_T = R(\emptyset, L, \aleph_0) < \omega$ .*

Note: Bokut asked on the number of e.c. algebras (associative, non-associative) over a field  $F$ . This class is unstable, so for  $\lambda \geq |F| + \aleph_1$ , the number is  $2^\lambda$ . Similar results holds of course for division rings. For e.c. groups Ziegler proved that for each e.c. group  $M$ , there is  $N \equiv_{\infty, \omega} M$  in each cardinal  $\geq ||M||$ ; we improve this to: for each  $\lambda \geq ||M|| + \aleph_1$ , there are  $2^\lambda$  such  $N$ 's (both results should appear in Springer Lecture Notes by Ziegler).

FIELDS

Theorem 7.2.

(Macintyre [Mc2]) *The only infinite fields with  $\aleph_0$ -stable theory are the algebraically closed fields.*

We still do not know whether there are stable or superstable fields or  $\aleph_0$ -stable division rings except the algebraically [or separably] closed fields (discarding the finite ones). [See 7.2, 8, 9] There is a small hope (hope, because then we can apply theorems on stability to such fields).

Macintyre, using 7.5 and elaborations of [Mc2] proved: if a field  $F$  is superstable then for each  $n$  it has finitely many extensions of degree  $n$ .<sup>(19)</sup>

<sup>(19)</sup> Added in Proof June 76: Note that if  $\tilde{F}$  is the algebraic closure of the field  $F$ , then  $\text{Th}(\tilde{F})$  is  $\lambda$ -stable iff  $\text{Th}(F, F)$  is  $\lambda$ -stable.



is finite. Hence  $D$  is finite, contradiction. If there is no such  $a$ , assume w.l.o.g.  $D$  is  $\aleph_1$ -saturated. Then for some  $n$ ,  $a^n = a$  for every  $a \in D$ , so  $D$  satisfies an identity and we finish as before. So we finish 7.3

**Rings**

The results on stability and  $\aleph_1$ -categoricity (7.7 - 7.9) are summed up in Cherlin and Reineke [CR] (see (1), (2), (3)), which contains more material. Macintyre and Rosenstein [MR] classifying the  $\aleph_0$ -categorical rings with no nilpotent elements Baldwin and Rose [BR] proving (4) (on  $\aleph_0$ -categoricity), give alternative proofs for some results of [CR], and some other results. Theorem 7.7 - 7.11 summarise some of their results. Baur, Macintyre and Cherlin, [\*1], have characterized  $\aleph_0$ -categorical  $\aleph_0$ -stable theories of rings and groups with  $\alpha_T < \omega$  (e.g. they are abelian by finite). Felgner, [\*2], characterizes  $\aleph_1$ -categorical semi-simple rings which are not necessarily commutative; in [\*3] and [\*4] he makes considerable advance for  $\aleph_0$ -categorical stable groups.

Let  $J = J(R)$  be the Jacobson radical of  $R$ .

**Theorem 7.7:**

If  $R$  is a ring with at least one non zero divisor, and  $R$  is  $(\aleph_1)$ -stable, then  $R$  has a unit,  $J$  is nilpotent, and  $R/J$  is a finite product of matrix rings of the form  $M_n(D)$ ,  $D$  a  $(\aleph_1)$ -stable division ring.

**Theorem 7.8:**

Suppose  $R$  is a commutative noetherian ring, with at least one nonzero divisor, then  $R$  is  $(\aleph_1)$ -stable iff  $J$  is nilpotent and  $R/J$  is a finite product of  $(\aleph_1)$ -stable fields.

**Theorem 7.9:**

Suppose  $R$  is a commutative ring with at least one non-zero divisor and  $R/J$  is infinite; then  $R$  is  $\aleph_1$ -categorical iff  $R = R' \oplus H$  where:

- i)  $H$  is finite
- ii)  $R'$  is a noetherian local ring (local means having a unique maximal ideal  $M$ ), this ideal  $M$  is nilpotent and  $R'/M$  is an algebraically closed field.

**Remark:**

- (1) If  $R$  has characteristic zero,  $R$  stable, then  $R/J$  is necessarily infinite.
- (2) The case  $R/J$  finite is open, but if in addition  $R$  is stable and noetherian, then  $R$  is finite.

**Theorem 7.10:**

Suppose  $R$  is noetherian, with at least one non-zero divisor. Then  $R$  is  $\aleph_0$ -categorical iff  $R$  is finite.

**Theorem 7.11:**

Suppose  $R$  has 1 and no nilpotent element. Then  $R$  is  $\aleph_0$ -categorical iff  $R$  is a finite direct product  $\prod_{1 \leq i \leq n} R_i$  where each  $R_i$  is of the form

$$C(X_i, F; X_1 \dots X_n, F_1 \dots F_n) \text{ where:}$$

- i)  $X$  is a Boolean space with finitely many isolated points,  $F$  is a finite field,  $X_1 \dots X_n$  are closed subsets of  $X$ ,  $F_1 \dots F_n$  are subfields of  $F$ ,
- ii)  $C$  is a ring of continuous functions  $f: X \rightarrow F$  such that  $f(X_i) \subseteq F_i$  and the dual structure  $(\hat{X}, \hat{X}_1, \dots, \hat{X}_n)$  (in the sense of Stone duality) (a Boolean algebra with ideals  $\hat{X}_1 \dots \hat{X}_n$ ) is  $\aleph_0$ -categorical. (Those structures are classified in [MR].)

**Generic division rings**

Let  $T$  be the (universal) theory of division rings. Cohn [CO] prove the JEM (for a fixed characteristic) and the amalgamation property. Boffa [BO1], [BO2], [BO3] want to find in which cardinal  $T$  has a universal homogeneous model. By 1.13 it suffices to note that  $\mathcal{D}(T, \Phi_{qt})$  is unstable (which, in fact, he

does). He asked for the number of infinite generic division rings, so 7.1 solves his problem (in cardinality  $\lambda > \aleph_0$ , there are  $2^\lambda$  non-isomorphic ones)

**Question:**

Suppose  $D$  is a division ring, and  $T$  is the universal theory of  $(D, a)_{a \in D}$ , and it has the JEM. In which cases is the class of infinite generic models of  $T$  stable? superstable?  $\aleph_0$ -stable?  $\aleph_1$  categorical?

**§ 8. ALGEBRA AND STRUCTURE THEOREMS**

*Algebraically closed fields*

It is well known that for each  $p$ , the theory of algebraically closed fields of characteristic  $p$  is  $\aleph_1$ -categorical. In fact this was the motivation of Loś' conjecture.

**Modules.**

For each ring  $R$ , we consider (always left) modules over  $R$  as additive groups with one-place functions  $F_a$  for each  $a \in R$ . Let  $R$  be fixed,  $L = L_R$  the corresponding language and  $T = T_R$  be the theory of such modules. By  $|R|$  we mean  $|R| + \aleph_0$ . The first order theories of abelian groups were classified by Szmielew [Sz]. Macintyre [Mc 1] classified the  $\aleph_1$ -categorical theories of abelian groups. Sabbagh [Sb 2] reproved Szmielew's theorem as well as Eklof and Fisher [EF] who further found in which cardinals a given complete theory of abelian groups has a saturated model. By 1.13 (and as mentioned in [S 5]) they essentially classified abelian groups by stability. In particular every abelian group is stable.

**Theorem 8.1.** (Mycielski [My], Sabbagh [Sb 2]) *If  $M$  is  $|R|$ -saturated,  $M < N$ , or even  $M < N$ , then  $M$  is a direct summand of  $N$ .*

*Remark.* In fact, we prove  $M$  is atomically compact (see e.g. Taylor [T 1]).

*Proof.* It suffices to show there is a homomorphism from  $N$  into  $M$ , which is the identity over  $M$  (as  $N = M \oplus \text{Ker } f$ ). For this it suffices to prove:

(\*) If  $\Gamma$  is a set of ep-formulas, with parameters from  $M$ , finitely satisfiable in  $M$ , then  $\Gamma$  is satisfiable in  $M$ .

(Because let  $|N| = \{a_i : i < \alpha\}$ ,  $\Gamma = \{\varphi(x_1, \dots, a_{j_1}, \dots) : \varphi \text{ is qf, } N \models \varphi [a_{i_1}, \dots, a_{i_j}, \dots]\}$  and  $a_{j_1}, \dots \in M$ ).

Clearly for proving (\*) it suffices to prove it for  $\Gamma$  with one variable (replace variables appearing in  $\Gamma$  by elements in  $M$  one by one; in limit stages remember that «finite satisfiability» has a finite character; for a variable  $x$  let

$$\Gamma' = \{ \exists y \bigwedge_{i=1}^n \varphi_i(x, \bar{y}) : \varphi_i(x, \bar{y}) \in \Gamma \}$$

and apply the assumption).

So let  $\Gamma = \{\varphi_1(x) : i < \alpha\}$  and let, w.l.o.g.

$$\varphi_1(x) = (\exists y_1 \dots y_{n(i)}) \bigwedge_{m < m(i)} \sum_{j=1}^{n(i)} r^{j,m} y_j + r^{i,m} x = a^{i,m}$$

where

$r^{j,m}, r^{i,m} \in R$ ,  $a^{j,m} \in M$ . ( $\varphi_j$  could have a disjunction inside, but by extending  $\Gamma$  we can get rid of it). Let us call  $\varphi_{(1)}$ ,  $\varphi_{(2)}$  similar if  $n(i(1)) = n(i(2))$ ,  $m(i(1)) = m(i(2))$ ,  $r^{j(i(1)),m} = r^{j(i(2)),m}$ ,

$r^{(1),m} = r^{(2),m}$ . Clearly similarity is an equivalence relation, so let  $\varphi_i (i < \alpha_0 \leq |R| + \aleph_0)$  be a set of representatives. So  $\Gamma_0 = \{\varphi_i(x) : i < \alpha_0\}$  is finitely satisfiable in  $M$ ,  $|\Gamma_0| \leq |R|$ ; so there is a  $b_0$  realizing it. We shall show that  $b_0$  realizes  $\Gamma$ . For  $\varphi_j$  there is  $i < \alpha_0$  such that  $\varphi_i, \varphi_j$  are similar. So let  $b \in M$

satisfies  $\varphi_\beta \wedge \varphi_i$ . So we can find  $y_1, \dots, y_j$  such that:

$$(i) \sum_{j=1}^{\beta(m)} r^{j,m} y_j + r^{i,m} b = a^{i,m} \text{ for } m < m(\beta)$$

$$(ii) \sum_{j=1}^{n(\beta)} \beta_{j,m} \gamma_j + \tau_j = a \quad \text{for } m < m(\beta)$$

(remember the similarity). Subtracting, we get

$$\sum_{j=1}^{n(\beta)} \beta_{j,m} (\gamma_j - \tau_j) = a - a$$

Hence if  $\gamma_j$  ( $j < n(i)$ ) exemplifies  $\varphi_1(b_0)$ , then  $\gamma_j + \tau_j - \tau_j$  exemplifies  $\varphi_\beta(b_0)$ .

Theorem 8.2. (Baur [Br], Fisher [Fi 1] [Fi 2])

Every module has a stable theory; so clearly also  $\mathcal{D}(T_R, \Phi a)$  is stable.

Proof. If not, there are R-modules  $M_1, M_2, A \subseteq M_1 \subseteq M_2, |A| = 2^{|R|}$ ,

$|\{tp(a, A, M_2) : a \in M_2\}| > ||M_1||$ . W.l.o.g.,  $M_2$  is  $|R|^{+-}$ -saturated, and  $A = |M_1|, M_1 < M_2, M_1$  is  $|R|^{+-}$ -saturated too. Now

by Feferman-Vaught if  $a_1, a_2, \dots \in M_2, f : M_2 \rightarrow M_1$  is a homomorphism,  $f \upharpoonright M_1 =$  the identity,  $M_0 = \text{Ker } f$ , and  $f(a_i) = f(a_i^2), tp(\langle \dots, a_i^1 - f(a_i^1), \dots \rangle, \emptyset, M_0) = tp(\langle \dots, a_i^2 - f(a_i^2), \dots \rangle, \emptyset, M_0)$ , then  $tp(\langle \dots, a_i^1, \dots \rangle, \emptyset, M_2) = tp(\langle \dots, a_i^2, \dots \rangle, \emptyset, M_2)$ . So we get a contradiction easily. The second part is left to the reader.

The following lemma is much related to Sabbagh [Sb 3]:

Lemma 8.3. Every formula in  $L_R$  is equivalent (modulo  $T_R$ ) to a Boolean combination of pe-formulas where  $L'_R = L_R \cup \{c_i : i < |R| + \aleph_0\}, T'_R$  is a  $\Pi_2$ -theory in  $L'_R$ , extending  $T_R$  such that every  $L_R$ -model of  $T_R$  can be expanded to a model of  $T'_R$ .

Proof. Left to the reader, remembering

Claim 8.4. (1) Let  $\Psi$  be a set of atomic formulas in  $L_R, \Psi_n$  the set of first-order formulas whose atomic subformulas belong to  $\Psi$  and their quantifier depth is  $\leq n$ . Let  $th_n(M) = \{\psi \in \Psi_n : M \models \psi\}$  and note  $|\Psi| < \aleph_0 \Rightarrow |\Psi_n| < \aleph_0$ . Then for each  $n$  there is  $k(n) < \omega$  such that : if  $M = \bigoplus_{i \in I} M_i$  (free sum of mo-

dules) then  $th_n(M)$  is determined by the truth value of the following statements : for each  $t$  which is a formally possible  $th_n(N)$  and cardinal  $\lambda, \lambda \in \{0, 1, \dots, k(n)\} \cup \{\aleph_0\}, \ll \{i \in I : t \models th_n(M_i)\} \geq \lambda \gg$ .

(2) Part (1) can be generalized to similar classes (as abelian structures of Fisher [Fi 1]).

Remark. For such theorems see Feferman-Vaught [FV], Galvin [Ga], Eklof and Fisher [EF] proved :  $th(M_i) = th(N_i) \Rightarrow th(\bigoplus M_i) = th(\bigoplus N_i)$  and Barwise and Eklof [BE] had similar results for infinitary languages (which we can too). The case  $\aleph_1 = \aleph_0$  is needed above, for if  $B_i$  are Boolean rings,  $\bigoplus_{i < \alpha} B_i$  has one iff  $\alpha < \omega$ . See also Waszkiewicz and Weglorz [WW].

The following theorem characterizes  $\kappa_r(T_R)$  except when  $\kappa(T_R) = \aleph_0$ .

Theorem 8.5. The following conditions on  $R, \kappa$  (where  $cf \kappa > \aleph_0$ ) are equivalent

- (1)  $\kappa \leq \kappa(T_R)$  (where  $\kappa(T_R) = \sup \{\kappa(T) : T \text{ a completion of } T_R\}$ ),
- (2) For each  $n < \omega$ , every submodule of  ${}_R(R^n)$  is generated by  $< \kappa$  elements.
- (3) Every ideal of  $R$  is generated by  $< \kappa$  elements. (17)

Proof. (2)  $\Rightarrow$  (1). Essentially just like 8.1, 8.2 ( $M$  will be  $\kappa$ -compact). In the proof of 8.1 we have to show that we can re-

(17) It seems that (2)  $\Rightarrow$  (1) does not work, so 8.5 is proved with  $qf$ -stability in (1). But 8.7 can be generalized for charactering  $\kappa < \kappa(\tau)$ .

place  $\Gamma_k = \{\varphi_i(x) : \langle \alpha_0, n(i), m(i) \rangle \leq k\}$  ( $k < \omega$ ) by a subset of cardinality  $< \kappa$ , which is easy by our assumption.

not (2)  $\Rightarrow$  not (1). (Here cf  $\kappa > \aleph_0$  is not used). We can find

$n < \omega$ , and  $\bar{a}^\alpha = (a_1, \dots, a_n) \in R^n$  ( $\alpha < \kappa$ ) such that  $\bar{a}^\alpha$  does not belong to the submodule of  $R^n$  generated by  $\{\bar{a}^\beta : \beta < \alpha\}$ .

For each  $\lambda$ , let  $M_\lambda$  be the module generated by:  $Y_\eta^\lambda$  ( $\eta \in \lambda$ ,

$\ell = 1, \dots, n$ ),  $z_\nu$  ( $\nu \in \lambda, \alpha < \kappa$ ) and subject only to the restric-

tion: whenever  $\nu$  is an initial segment of  $\eta$ ,  $\alpha \in \lambda$ ,

$$\sum_{\ell=1}^n a_\ell Y_\eta^\alpha = z_\nu.$$

It suffices to show that

$$\sum_{\ell=1}^n a_\ell Y_\eta^\alpha = z_\nu \text{ holds iff } \nu \text{ is an initial segment of } \eta.$$

Suppose  $\eta_0, \nu$  is a counter-example, and let  $h$  be the following endomorphism of  $M_\lambda$ , where

$$\alpha = \min \{\alpha : \eta_0(\alpha) \neq \nu(\alpha)\} : \eta_1 \in \lambda, \nu \text{ an initial segment of } \eta_1.$$

$$(i) \text{ for } \varrho \in \lambda, \varrho \uparrow (\alpha + 1) = \nu \uparrow (\alpha + 1), h(y_\varrho^\ell) = y_{\eta_1}^\ell$$

$$(ii) \text{ for } \varrho \in \lambda, \varrho \uparrow (\alpha + 1) \neq \nu \uparrow (\alpha + 1), h(y_\varrho^\ell) = y_{\eta_0}^\ell$$

$$(iii) \text{ for } \varrho \in \lambda, \varrho \uparrow (\alpha + 1) = \nu \uparrow (\alpha + 1), h(z_\varrho) = z_{\eta_1 \uparrow \ell(\varrho)}$$

$$(iv) \text{ for } \varrho \in \lambda, \varrho \uparrow (\alpha + 1) \neq \nu \uparrow (\alpha + 1), h(z_\varrho) = z_{\eta_0 \uparrow \ell(\varrho)}$$

Clearly  $h$  is a projection onto the submodule generated by

$Y_{\eta_0}^\ell, Y_{\eta_1}^\ell$  ( $\ell = 1, \dots, n$ ) subject to the conditions:

$$\sum a_\ell Y_{\eta_0}^\beta = \sum a_\ell Y_{\eta_1}^\beta \text{ (for } \beta < \alpha).$$

Easily this contradicts the assumption on  $\bar{a}^\gamma$  ( $\gamma < \kappa$ ).  
(2)  $\Rightarrow$  (3). Trivial.

(3)  $\Rightarrow$  (1). We prove by induction on  $n$ .  $R^{n+1} = R^n \oplus R$ , so if  $M$  is a submodule of  $R^{n+1}$ , then  $R \oplus M/R$  (considering  $R$  as submodule of  $R^{n+1}$ ) and  $M \cap R^n$  are  $< \kappa$ -generated; from this the conclusion follows.

Remark. Here we get the unstability by qf-formulas. In fact we proved (we can even refine 8.5 by adding in (2)  $\ell(a) = n_0$  and restrict (2) to  $n_0$ ):

Theorem 8.6. The following conditions on  $\kappa$  are equivalent (when cf  $\kappa > \aleph_0$ ):

- (1) If  $\lambda^{<\kappa} = \lambda \geq 2$ ,  $|R| \leq \lambda$ ,  $A \subseteq M \models T_R$ , then  $\{\{tp_{cf}(\bar{a}, A, M) : \bar{a} \in M\} \leq \lambda_i$
- (2) For each  $n < \omega$ , every submodule of  ${}_R(R^n)$  is generated by  $< \kappa$  elements; and when  $\kappa > \aleph_0$ ,
- (3) The conditions of 8.4.

Remark. We can infer now that if  $R$  is not noetherian,  $T_R$  is not superstable (which can be inferred from «not artinian»); moreover, there are qf-formulas, hence (by 7.1) for every regular

$\lambda > \aleph_0$  there are  $2^\lambda$   $R$ -modules, no one embeddable in another. Of course this conclusion holds under more general conditions.

Theorem 8.7. The following conditions are equivalent:

- (1)  $T_R$  is superstable,
- (2) not: for every regular  $\lambda > |R|$  there is an  $R$ -module of cardinality  $\lambda$  which is not the direct sum of  $R$ -modules of cardinality  $< \lambda$ ,
- (3) every  $R$ -module is the direct sum of countably generated  $R$ -modules,
- (4) if  $M < N$  then there is a projection from  $N$  onto  $M$ ,
- (5) every e p-1-type over  $M$  which is finitely satisfiable in

$M$ , has an equivalent finite subtype, or equivalently is realized in  $M$ .

Note: It seems that the number of e.g.  $|R|^{+}$ -saturated modules in  $\aleph$  is  $\leq 2^{|\alpha|}$ , so we have a structure theorem.

*Remark.* Fisher [Fi] generalizes this (to abelian structures) and strengthens it to having a unique decomposition. <sup>(19)</sup>

*Proof (5)  $\Rightarrow$  (4).* Just like the proof of 8.1.

(4)  $\Rightarrow$  (3). This is a standard fact. Indeed (4) means that every  $R$ -module is pure-injective, hence that every module is pure-projective, hence that every module is a direct summand of a direct sum of finitely presented  $R$ -modules (see for all those implications Warfield [Wf]). A classical theorem of Kaplansky concludes the proof.

For completeness, let us prove that every  $R$ -module  $M$  is the direct sum of  $R$ -modules of cardinality  $\leq |R|$ . We prove it by induction on  $\|M\|$ . For  $\|M\| \leq |R|$  there is nothing to prove. Otherwise let  $\|M\| = \lambda > |R|$ , and we define by induction on  $\alpha < \lambda$  an increasing continuous sequence  $M_\alpha <_{ep} M$ ,  $\|M_\alpha\| < \alpha$ .

Let  $U_\alpha = M_\alpha$ , and projections  $h_\alpha$  from  $M$  onto  $M_\alpha$ . (We define by induction on  $\alpha$ ,  $M_\alpha$  and then  $h_\alpha$  using (3)). Let  $N_\alpha = \text{Ker } h_\alpha \upharpoonright M_\alpha$ , so  $M_\alpha = M_\alpha \oplus N_\alpha$ , hence  $M = \sum_{\alpha < \lambda} N_\alpha$  so using the induction hypothesis on  $\lambda$  the conclusion of (3) follows.

(3)  $\Rightarrow$  (2). Trivial.

*not (1)  $\Rightarrow$  not (2).* Easy by the construction in [S3] 1.1, see here § 0,7.1.

*not (5)  $\Rightarrow$  not (1).* So there is an  $R$ -module  $M$  and ep-fol-

<sup>(19)</sup> In a proper sense. We can add: (6)  $I(\aleph_\alpha, \aleph_\alpha) < 2^\alpha$  for some  $\aleph_\alpha > |T_R|$ , (7)  $I(\aleph_\alpha, \aleph_\alpha) \leq |\alpha|^z$  (and we can get the exact value, see [S1], IX, § 2).

mulas  $\varphi_n(x) = \varphi_n(x, a_1, \dots, a_k)$  ( $a_i \in M$ ) such that  $M \models (\exists x) \bigwedge_{i < n} \varphi_i(x)$  but  $M \models (\exists x) (\bigwedge_{i \leq n} \varphi_i(x) \wedge \neg \varphi_n(x))$ . By notational

changes we can assume  $\varphi_n(x) = (\exists y_1 \dots y_{k(n)}) \bigwedge_{i < k(n)} \sum_{j=1}^i y_j + s_i x = a_j$  and that for each  $n$ ,  $M \models \neg \varphi_{n+1}(c_n) \wedge \bigwedge_{i < k(n)} \sum_{j=1}^i y_j + s_i c_n = a_j$ . For notational simplicity let  $k(n) = n$ .

Let us find  $M^*$ ,  $c^*$ ,  $b_j$  ( $j < \omega$ ) such that  $M < M^*$   $\models \sum_{j=1}^i y_j b_j + s_i c^* = a_j$ . Now for each  $\lambda$ , let  $N$  be an  $R$ -module generated by  $x, y, z, (\eta \in \omega, \lambda, \nu \in \omega > \lambda, \ell < \omega)$  subject only to the fol-

lowing conditions: when  $\eta \in \lambda$ ,  $\bigwedge_{i < k(n)} \sum_{j=1}^i y_j y_j + s_i x =$

$\eta \uparrow i$ . Clearly  $N \models \varphi_n(x, z, \eta \uparrow 0, \dots, z, \eta \uparrow k(n))$ , and it is sufficient to prove that if  $\eta \in \lambda$ ,  $n < \omega$ ,  $\nu = \eta \uparrow n \wedge \langle \alpha \rangle$ ,  $\eta(n) \neq \alpha$ , then  $N \models \neg \varphi_{n+1}(x, z, \nu \uparrow 0, \dots, z, \nu)$ . Suppose  $\eta, \nu$  are a counterexample, and we shall define a homomorphism  $h$  from  $N$  into  $M^*$ :

- (i) if  $\eta^1 \in \lambda$ ,  $\eta^1 \uparrow (n+1) \neq \nu$  then  $h(x) = c^*$  and  $h(y_j) = b_j$ ,
- (ii) if  $\eta^1 \in \lambda$ ,  $\eta^1 \uparrow (n+1) = \nu$  then  $h(x) = c_n$  and  $h(y_j) = b_j$  for  $j \leq k(n)$ ,
- (iii) if  $\eta^1 \in \lambda$ ,  $\eta^1 \uparrow (n+1) = \nu$ ,  $j > k(n)$ , then  $h(y_j) = b_j$ ,
- (iv) if  $\eta \in \aleph_\lambda$ ,  $m \leq n$  or  $\eta \uparrow (n+1) \neq \nu$ , then  $h(z) = a_m$ ,
- (v) for other  $\eta \in \aleph_\lambda$ , choose any  $\eta^1 \in \omega$ ,  $\eta \in \eta^1 \uparrow m$  and

$$\text{let } h(z) = \sum_{j=1}^m r_j h(Y_j) + s_m h(x).$$

The contradiction we get is clear (by the choice of  $\eta$ ).

$N \models \varphi_{n+1}(x, z, \bigvee_{i=0}^{\eta} z_i, \dots, z_\eta)$ , so as  $h$  is ep-formula,  $M^* \models$

$\varphi_{n+1}(c_n, a_0, \dots, a_{n+1})$  contradicting the choice of  $c_n$ .

Sabbagh noticed that our results imply that if  $R$  is commutative and countable, then  $T_R$  is superstable iff  $T_R$  is  $\aleph_0$ -stable.

*Differentially closed fields.*

A differential field is a natural generalization of a field: we add to the theory of fields of characteristic  $p$  a one-place function  $D$ , and the axioms  $D(x + y) = Dx + Dy$ ,  $D(xy) = (Dx)y + x(Dy)$ , and get  $T_{df}^p$ . Using Seidenberg [Si], Robinson [Rb-1] showed that  $T_{df}^0$  has a model completion  $T_{dc}^0$  (the theory of differentially closed fields).

Blum [Bl 1] gives a nice axiomatization: if  $P_1, P_2$  are differential polynomials with one variable  $Y$ , of orders  $m_1, m_2, m_1 > m_2 \neq 0$ , and for  $P_2 = 0$ . (<sup>(9)</sup>)

Later Wood [W 1] [W 2], again using [Si 1], proves similar results for  $T_{dc}^p, p > 0$ , when we add the  $p$ -th root as an operation; i.e. we add the axioms:  $r(x) \neq 0 \rightarrow r(x)^p = x, r(x) = 0 \rightarrow (\forall y) y^p \neq y$  (note that  $x = y^p \rightarrow Dx = 0$ ) and we get the theory of radical fields  $T_{rd}^p$  and its model completion the theory  $T_{rdc}^p$  of the radically differentially closed fields.

Blum [Bl 1] shows  $T_{dc}^0$  is  $\aleph_0$ -stable: quite naturally, the Morley rank of a 1-type is  $\leq$  the minimal  $m$  such that it implies  $x^m = 0$ .

(<sup>9</sup>) Meanwhile S. Fakir in a mimeograph asserts there is a mistake, and suggests a correction.

is a root of a differential polynomial of order  $m$ , and  $\omega$  if there is no such  $m$ ; hence by Morley [M 1], over any differential field, there is a prime differentially closed field, hence by [S 9] (here 4.5), it is unique up to isomorphism (but it is not minimal, see [K1 3] [Rs 1] [S 10]).

Shelah [S 10] and Wood [W 2] prove, independently, that over any radical differential field there is a prime radical differentially closed field. In [S 10] it is proved that  $T_{rdc}^p$  is stable (it is not superstable as noted by [S 10] from a proof in [W 1] that it is not  $\aleph_0$ -stable).

Hence by 4.6,

*Theorem 8.8.* Over any radical differential field (in fact a field such that  $Dx = 0 \rightarrow (\exists y) y^p = x$ ), there is a unique prime radical differentially closed field.

We should note that though  $T_{dc}^0$  is  $\aleph_0$ -stable, by [S 10] it has the dimensional order property (e.g. by adding the cardinality quantifier  $\exists > \aleph_0$  we get long orders in some models), hence it has non-structure theorem.

Macintyre and Shelah note

*Theorem 8.9.* If  $F$  is a separably closed but not algebraically closed field, then  $\text{Th}(F)$  is stable but not superstable.

*Proof.* In some cases it follows from the theorem on the stability of  $T_{rdc}^p$  ([S 10]) and always similarly. The unsuperstability follows from 7.5(2) as  $F^{p^n}$  is a strictly decreasing sequence of definable additive subgroups and  $F^p/F^{p^{n+1}}$  is infinite (quotient as additive groups).

C. Wood, [<sup>\*</sup>5], has written the proof in details. It seems that those theories has the dimensional order property, hence  $2^\lambda$  modules in each  $\lambda > \aleph_0$ .  
*Operations preserving stability.*

For a wide class of operations on models, we can prove that they preserve stability

Lemma 8.10. If  $Op(\dots, M_i, \dots)_{i < \alpha}$  is an operation on models

- (i) preserving elementary equivalence (i.e.  $M_i \equiv N_i \Rightarrow Op(\dots, M_i, \dots) \equiv Op(\dots, N_i, \dots)$ ),
- (ii) for each  $\lambda$ , it preserves  $\lambda^+$ -saturation (i.e. if each  $M_{i_j}$  is  $\lambda^+$ -saturated then  $Op(\dots, M_i, \dots)$  is),
- (iii) if each  $M_i$  is  $\lambda_0$ -stable model then  $Op(\dots, M_i, \dots)$  is  $\lambda_0$ -stable for each  $i$  implies  $Th Op(\dots, M_i, \dots)$  is  $\lambda_0$ -stable.

Proof. Trivial (in fact, (i) is not needed).

The simplest application is that reducts preserve stability. Similar theorems hold for stability of classes we discuss here (generic,  $\mathcal{D}$ -homogeneous, etc...). The proof of the hypothesis is in many cases easy by Feferman-Vaught [FV], or maybe the refinements to autonomous systems of Galvin [G1].

J. Wierzejewski [W1] used a particular instance of 8.10 (for products using the lemma of J. Waszkiewicz and B. Weglorz [WW]) that finite products preserve saturation). So he proved  $\aleph_0$ -stability, superstability, stability and also the negation of the finite-cover-property, the negation of the independence property, and the negation of the strict order property.

He also proved in [W1] that the negations of those properties are not preserved even by squares. The author found when  $M = M_1 + M_2$ ,  $M$  is stable iff  $M_1$  and  $M_2$  are stable; and similarly for the other properties.

REFERENCES

- [Ba] J.T. Baldwin,  $\sigma_T$  is finite for  $\aleph_1$ -categorical T, *Trans. A.M.S.* 161 (1973), 37-51.
- [BE] J. Barwise and P. Eklof, Infinitary properties of abelian torsion groups, *Annals of Math. Logic* 2 (1970), 25-68.
- [BL 1] L. Blum, Generalized algebraic structures, a model theoretic approach, Ph. D. Thesis, M.I.T. 1968.
- [Bo 1] M. Boffa, Corps  $\lambda$ -clos, *C. R. Acad. Sc. Paris, sér. A* 275 (1972) 881-882.
- [Bo 2] M. Boffa, Sur l'existence des corps universels homogènes, *Acad. Sc. Paris, sér. A* 275 (1972), 1267-1268.

- [Bo 3] M. Boffa, Generic skew fields, *Logical seminar in Warsaw*, 1973, preprint.
- [Bf] W. Baur,  $\aleph_0$ -categorical modules, *J. Symb. Logic*, 40, (1975), 213-220.
- [BS] J.T. Baldwin and J. SaxeL, Logical Stability in Group Theory, *J. of the Australian Math. Soc.*, XXV (Ser. A), Part 3, May 1976, 267-276.
- [Co] P.M. Cohn, The embedding of firs in skew fields, *Proc. London Math. Soc.* (3) 23 (1971), 193-213.
- [Cr] G. Cherlin, The model-companion of a class of structures, *J. Symb. Logic* 37 (1972), 546-556.
- [CR] G. Cherlin and J.M. Reinecke, Categoricity and stability of commutative rings, *Annals of Math. Logic*.
- [EF] P.C. Eklof and E.R. Fisher, The Elementary properties of abelian groups, *Annals of Math. Logic*, 4 (1972), 115-171.
- [EHR] P. Erdős, A. Hajnal and R. Rado, Partition relations for cardinal numbers, *Acta. Math. Acad. Scientiarum Hungaricae* 16 (1965), 93-196.
- [Ek] P. C. Eklof, Categories of local functors, *ROBINSON Memorial Volume*, Springer.
- [EM] A. Ehrenfeucht and A. Mostowski, Models of axiomatic theories admitting automorphisms, *Fund. Math.* 43 (1956), 50-68.
- [Fi 1] E. Fisher, Powers of saturated modules, abstract, *J. Symb. Logic*, 37 (1972), 777.
- [Fi 2] E. Fisher, Abelian Structures.
- [Fu] L. Fuchs, *Infinite abelian groups*, Vol. I 1970, Vol. II 1973, Academic Press, N.Y.
- [FV] S. Feferman and R.L. Vaught, The first order properties of algebraic systems, *Fund. Math.*, 47 (1959), 57-103.
- [G 1] H. Gaifman, Operations and relations on structures, functors and classes, I, *Proc. of the Tarski Symp. Berkeley* 71. *Proc. of Symp. in Pure Math.* vol. XXV, ed. L. Henkin, A.M.S. 1974, 21-41.
- [G 2] H. Gaifman, *Definability and many sorted logic*, mimeographed notes of a lecture given at the Robinson memorial Coll. at Yale 1975.
- [Ga] F. Galvin, Horn sentences, *Annals of Math. Logic* 1 (1970), 389-422.
- [Ha] V. Harnik, On the existence of saturated models of stable theories, *Israel J. of Math.*
- [J 1] I. Juhasz, Cardinal function in topology (in collaboration with A. Verbeek and S. Koenenberc), *Math. Centr. Trans.* 34.
- [Jn] B. Jonsson, A Boolean algebra without proper automorphism, *Proc. A.M.S.* 2 (1951), 770-776.
- [K 1] H.J. Keisler, Ultraproducts which are not saturated, *J. Symb. Logic* 32 (1967), 23-47.

SAHARON SHELAH

[K1 2] E.K. KOŁCHIN, *Differential algebra and algebraic groups*, N.Y. Academic Press 1973 (Pure and Applied Math. 54).

[K1 3] E.K. KOŁCHIN, *Constrained extensions of differential fields, Advances in Math.*, 12 (1974), 141-170.

[Ls] D. LASCAR, *Ranks and definitability in superstable theories; Israel J. of Math.*

[Lz] F.W. LOZIER, *A class of compact rigid 0-dimensional spaces, Canadian J. Math.* 21 (1969), 817-821.

[MA] K. MC ALOON, *Les algèbres de Boole rigides et minimales, C.R. Acad. Sc. Paris sér. A* 272 (1971), 89-91.

[Mc 1] A. MACINTYRE, *On  $\aleph_1$ -categorical theories of abelian groups, Fund. Math.* 70 (1971), 253-270.

[Mc 2] A. MACINTYRE, *On  $\aleph_1$ -categorical theories of fields, Fund. Math.* 71 (1971), 1-25.

[McS] R. MCKENZIE and S. SHELAH, *The cardinal of simple models for universal theories, Proc. of Berkeley 71 Symp. in honour of TARSKI seventieth birthday; Proc. Symp. in Pure Math.*, Vol. XXV, A.M.S. 1974; 53-74.

[M.M] R. MCKENZIE and J.D. MONK, *On automorphism groups of Boolean algebras, Proc. of the International Colloquium on Infinite and Finite sets; in honour of P. ERDŐS sixtieth birthday, Keszthely Hungary, 1973.*

[MR] A. MACINTYRE and J. ROSENSTEIN,  $\aleph_0$ -categorical rings without nilpotent elements, and Boolean Structures, *J. of Algebra*, submitted.

[MS] A. MACINTYRE and S. SHELAH, *Uncountable universal locally finite groups, J. of Algebra.*, submitted.

[MV] M.D. MORLEY and R.L. VAUGHT, *Homogeneous universal models; Math. Scandinavica* 11 (1962), 37-57.

[My] J. MYCIELSKI

[Pr] H. PRIESTLEY, *Ordered topological spaces and the representation of distributive lattices, Proc. London Math. Soc.* 24 (1972), 507-530.

[Rb 1] A. ROBINSON, *Forcing in model theory, Actes du Congrès International des Mathématiciens, Nice 1970. Gauthier-Villars 1971* 245-250.

[RS 1] M. ROSENBLUTH, *The non-minimality of differential closure, Pacific J. of Math.*

[Rt] J.F. RITT, *Differential algebras, Amer. Math. Soc. Colloquium Publ.*, 33 (1950).

[Sb 1] G. SABBAGH, *Sur la pureté dans les modules, C.R. Acad. Sci. Paris sér. A* 271 (1970), 865-867.

[Sb 2] G. SABBAGH, *Aspects logiques de la pureté dans les modules, C.R. Acad. Sci. Paris sér. A* 271 (1971), 865-867.

[Sb 3] G. SABBAGH, *Sous-modules purs existentiellement clos et élémentaires, C.R. Acad. Sci. Paris sér. A* 272 (1971), 1289-1272.

[Sc] G. SACKS, *Saturated model theory*, Benjamin, Reading Mass., 1972

[Sd] A. SEIDENBERG, *An elimination theory for differential algebra, Univ. California Publ. Math.* (new serie) 3 (1964), 31-66.

[S 1] S. SHELAH, *Stability and the number of non-isomorphic models, North-Holland Publ. Co.*, in preparation.

[S 2] S. SHELAH, *Infinite abelian groups, Whitehead Problem and some Constructions, Israel J. of Math.* 18 (1974), 243-256.

[S 3] S. SHELAH, *Why there are many non-isomorphic models for unstable theories, Proc. of the 1974 International Congress of Mathematicians, Vancouver.*

[S 4] S. SHELAH, *Saturation of ultrapowers and Keisler's order, Annals of Math. Logic* 4 (1972), 75-114.

[S 5] S. SHELAH, *Stability, the f.c.p. and superstability; model theoretic properties of formulas in the first order theory, Annals of Math. Logic* 3 (1971), 271-362.

[S 6] S. SHELAH, *On categoricity and stability over a predicate, in preparation.*

[S 7] S. SHELAH, *Finite diagrams stable in power, Annals of Math. Logic* 2 (1970), 69-118.

[S 8] S. SHELAH, *A combinatorial problem, stability and order for models in infinitary languages, Pacific J. of Math.* 41 (1972), 247-261.

[S 9] S. SHELAH, *Uniqueness and characterization of prime models over sets for totally transcendental first order theories, J. of Symbolic Logic* 37 (1972), 107-113.

[S 10] S. SHELAH, *Differentially closed fields, Israel J. of Math.* 16 (1973), 314-328.

[S1] R.M. SOLOVAY, *Real-valued measurable cardinals, Proc. of Symp. in Pure Math.*, XIII part I, Ed. D. Scott, A.M.S., Providence R.I.

[T] W. TAYLOR, *Some constructions of compact algebra, Annals of Math. Logic* 3 (1971), 395-346.

[W 1] C. WOOD, *The model theory of differential fields of characteristic  $p \neq 0$ , Proc. Amer. Math. Soc.* 40 (1973), 577-584.

[W 2] C. WOOD, *Prime model extensions for differential fields of characteristic  $p \neq 0$ , J. Symb. Logic.*

[W 3] C. WOOD, *Forcing for infinitary languages, Z. Math. Logic und Grundlagen Math.* 18 (1972), 385-402.

[WF] R.B. WARFIELD, *Purity and algebraic compactness for modules, Pacific J. Math.* 28 (1969), 699-719.

[Wt] J. WIERZEJSKI, *On stability and products, Fund. Math.*, to appear.

[WW] J. WASZKIEWICZ and B. WĘGIŁORZ, *On  $\omega_0$ -categoricity of powers, Bull. Acad. Pol. Sci. sér. de Math., Astron. et Physique.*

[\*1] W. BAUR, G. CHERLIN and A. MACINTYRE, *On totally categorical groups and rings, to appear.*

[\*2] U. FELCNER,  *$\aleph_1$ -kategorische Theorien nicht-kommutativer Ringe, Fund. Math.*