

Non-trivial automorphisms of $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$ from variants of small dominating number

Saharon Shelah^{1,2} · Juris Steprāns³

Received: 15 May 2014 / Revised: 9 January 2015 / Accepted: 28 May 2015 /

Published online: 21 July 2015

© Springer International Publishing AG 2015

Abstract It is shown that if various cardinal invariants of the continuum related to \mathfrak{d} are equal to \aleph_1 then there is a non-trivial automorphism of $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$. Some of these results extend to automorphisms of $\mathcal{P}(\kappa)/[\kappa]^{<\kappa}$ if κ is inaccessible.

Keywords Boolean algebra · Automorphism · Cardinal invariant · Dominating number · Forcing

Mathematics Subject Classification 03E35 · 03E50 · 03G05

Originally considered by the Central European Journal of Mathematics but withdrawn due to imposition of publishing fees and resubmitted to the European Journal of Mathematics.

The first author's research for this paper was partially supported by the United States–Israel Binational Science Foundation (Grant No. 2006108), and by the National Science Foundation (Grant No. NSF-DMS 0600940). (It is number 990 on his list of papers.) The second author's research for this paper was partially supported by NSERC of Canada.

✉ Juris Steprāns
steprans@yorku.ca

Saharon Shelah
shelah@math.huji.ac.il

- ¹ Institute of Mathematics, Hebrew University of Jerusalem, Givat Ram, 9190401 Jerusalem, Israel
- ² Department of Mathematics, Rutgers University, Hill Center – Busch Campus, 110 Frelinghuysen Road, Piscataway, NJ 08854-8019, USA
- ³ Department of Mathematics, York University, N520 Ross, 4700 Keele Street, Toronto, ON M3J 1P3, Canada

1 Introduction

A fundamental result in the study of the Čech–Stone compactification, due to Rudin [7, 8], is that, assuming the Continuum Hypothesis, there are 2^c autohomeomorphisms of $\beta\mathbb{N} \setminus \mathbb{N}$ and, hence, there are some that are non-trivial in the sense that they are not induced by any one-to-one function on \mathbb{N} . While Rudin established his result by showing that for any two P-points of weight \aleph_1 there is an autohomeomorphism sending one to the other, Parovičenko [6] showed that non-trivial autohomeomorphisms could be found by exploiting the countable saturation of the Boolean algebra of clopen subsets of $\beta\mathbb{N} \setminus \mathbb{N}$ —this is isomorphic to the algebra $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$. Indeed, the duality between Stone spaces of Boolean algebras and algebras of regular open sets shows that the existence of non-trivial autohomeomorphisms of $\beta\mathbb{N} \setminus \mathbb{N}$ is equivalent to the existence of non-trivial isomorphisms of the Boolean algebra $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$ to itself.

Notation 1.1 If A and B are subsets of κ let \equiv_κ denote the equivalence relation defined by $A \equiv_\kappa B$ if and only if $|A \Delta B| < \kappa$ and $A \subseteq_\kappa B$ will denote the assertion that $|A \setminus B| < \kappa$. Let $[A]_\kappa$ denote the equivalence class of A modulo \equiv_κ and let $\mathcal{P}(\kappa)/[\kappa]^{<\kappa}$ denote the quotient algebra of the $\mathcal{P}(\kappa)$ modulo the congruence relation \equiv_κ . If $\kappa = \omega$ it is customary to use \equiv^* instead of \equiv_ω and \subseteq^* instead of \subseteq_ω .

Notation 1.2 If f is a function defined on the set A and $X \subseteq A$ then the notation $f(X)$ will be used to denote $\{f(x) : x \in X\}$ in spite of the potential for ambiguity.

Definition 1.3 An isomorphism $\Phi : \mathcal{P}(\kappa)/[\kappa]^{<\kappa} \rightarrow \mathcal{P}(\kappa)/[\kappa]^{<\kappa}$ will be said to be *trivial* if there is a one-to-one function $\varphi : \kappa \rightarrow \kappa$ such that $\Phi([A]_\kappa) = [\varphi(A)]_\kappa$ for each $A \subseteq \kappa$. The isomorphism Φ will be said to be *somewhere trivial* if there is some $B \in [\kappa]^\kappa$ and a one-to-one function $\varphi : B \rightarrow \kappa$ such that $\Phi([A]_\kappa) = [\varphi(A)]_\kappa$ for each $A \subseteq B$ and Φ will be said to be *nowhere trivial* if it is not somewhere trivial.

The question of whether the Continuum Hypothesis, or some other hypothesis, is needed in order to find a non-trivial isomorphism of $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$ to itself was settled in the affirmative by Shelah in [9]. The argument of [9] relies on an iterated oracle chain condition forcing to obtain a model where $2^{\aleph_0} = \aleph_2$ and every isomorphism of $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$ to itself is induced by a one-to-one function from \mathbb{N} to \mathbb{N} . The oracle chain condition requires the addition of cofinally many Cohen reals and so $\mathfrak{d} = \aleph_2$ in this model. Subsequent work has shown that it is also possible to obtain that every isomorphism of $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$ is trivial by other approaches [2, 10, 14] but these have always required $\mathfrak{d} > \aleph_1$ as well. However, it was shown in [11] that this cardinal inequality is not entailed by the non existence of nowhere trivial isomorphisms from $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$ to itself—in the model obtained by iterating ω_2 times Sacks reals there are no nowhere trivial isomorphisms yet $\mathfrak{d} = \aleph_1$.

On the other hand, while we now know that the Continuum Hypothesis cannot be completely eliminated from Rudin's result, perhaps it can be weakened to some other cardinal equality such as $\mathfrak{d} = \aleph_1$. It will be shown in this article that non-trivial isomorphisms of $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$ to itself can indeed be constructed from hypotheses on cardinal arithmetic weaker than $2^{\aleph_0} = \aleph_1$ and reminiscent of $\mathfrak{d} = \aleph_1$. However, it is shown in [3] that it is consistent with set theory that $\mathfrak{d} = \aleph_1$ yet all isomorphisms of $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$ are trivial so some modification of the equality $\mathfrak{d} = \aleph_1$ will be required.

It will also be shown that natural generalizations of the arguments can be applied to the same question for $\mathcal{P}(\kappa)/[\kappa]^{<\kappa}$ where κ is inaccessible. The chief interest here is that, unlike $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$, the algebra $\mathcal{P}(\kappa)/[\kappa]^{<\kappa}$ is not countably saturated if $\kappa > \omega$ —to see this, simply consider a family $\{A_n\}_{n \in \omega} \subseteq [\kappa]^\kappa$ such that $\bigcap_{n \in \omega} A_n = \emptyset$. In other words, Parovičenko's transfinite induction argument to construct non-trivial isomorphisms from $\mathcal{P}(\kappa)/[\kappa]^{<\kappa}$ to itself is not available and some other technique is needed.

The statement and proof of Lemma 2.1 is provided for all κ and we will apply both to the case that $\kappa = \omega$ and to the case that κ is inaccessible. However, Lemma 3.2 deals only with the case that κ is inaccessible. It is somewhat simpler than the case for ω and so is dealt with first because the general approach is similar in the $\kappa = \omega$ case, but this case requires some technical details not needed in the inaccessible case.

2 A sufficient condition for a non-trivial isomorphism

The following lemma provides sufficient conditions for the existence of a non-trivial isomorphism of $\mathcal{P}(\kappa)/[\kappa]^{<\kappa}$ to itself. The set theoretic requirements for the satisfaction of these conditions will be examined later. The basic idea of the lemma is that an isomorphism of $\mathcal{P}(\kappa)/[\kappa]^{<\kappa}$ can be approximated by partitioning κ into small sets I_ν and constructing isomorphisms from subalgebras of $\mathcal{P}(I_\nu)$ and taking the union of these. Unless the subalgebras of $\mathcal{P}(I_\nu)$ are all of $\mathcal{P}(I_\nu)$, this union will only be a partial isomorphism. Hence a κ^+ length sequence of ever larger families of subalgebras of $\mathcal{P}(I_\nu)$ is needed to obtain a full isomorphism. In order to guarantee that this isomorphism is not trivial the prediction principle described in hypothesis (H₃) of the lemma is needed.

Lemma 2.1 *There is a non-trivial automorphism of $\mathcal{P}(\kappa)/[\kappa]^{<\kappa}$ provided that there is a partition of κ by $\{I_\nu\}_{\nu \in \kappa}$ such that*

- (H₁) $|I_\nu| < \kappa$ for each $\nu \in \kappa$.
- (H₂) For each $\xi \in \kappa^+$ and $\nu \in \kappa$ there is a Boolean subalgebra $\mathfrak{B}_{\xi, \nu}$ of $\mathcal{P}(I_\nu)$ and an automorphism $\Phi_{\xi, \nu}$ of $\mathfrak{B}_{\xi, \nu}$.
- (H₃) If $\xi \in \eta$ then there is $\iota \in \kappa$ such that $\mathfrak{B}_{\xi, \nu} \subseteq \mathfrak{B}_{\eta, \nu}$ and $\Phi_{\xi, \nu} = \Phi_{\eta, \nu} \upharpoonright \mathfrak{B}_{\xi, \nu}$ for all $\nu \in \kappa \setminus \iota$.
- (H₄) For any one-to-one $F: \kappa \rightarrow \kappa$ there are $\xi \in \kappa^+$ and cofinally many $\nu \in \kappa$ for which there is an atom $a \in \mathfrak{B}_{\xi, \nu}$ and $\iota \in a$ such that $F(\iota) \notin \Phi_{\xi, \nu}(a)$.
- (H₅) For any $A \subseteq \kappa$ there are $\xi \in \kappa^+$ and ι in κ such that $A \cap I_\nu \in \mathfrak{B}_{\xi, \nu}$ for all $\nu \in \kappa \setminus \iota$.

Proof Define

$$\Phi([A]_\kappa) = \lim_{\xi \rightarrow \kappa^+} \left[\bigcup_{\nu \in \kappa} \Phi_{\xi, \nu}(A \cap I_\nu) \right]_\kappa$$

and begin by observing that this is well defined. To see this, it must first be observed that given A and B such that $|A \Delta B| < \kappa$ there is $\alpha \in \kappa^+$ such that for all $\xi > \alpha$ and for all ν in a final segment of κ the equation

$$\Phi_{\xi,v}(A \cap I_v) = \Phi_{\xi,v}(B \cap I_v)$$

is defined and valid by hypothesis (H₄). From hypothesis (H₂) it then follows that if ξ and η are greater than α then

$$\bigcup_{v \in \kappa} \Phi_{\xi,v}(A \cap I_v) \equiv_{\kappa} \bigcup_{v \in \kappa} \Phi_{\eta,v}(B \cap I_v)$$

and, hence, $\Phi([A]_{\kappa})$ is well defined. Since each $\Phi_{\xi,v}$ is an automorphism it follows that Φ is an automorphism of $\mathcal{P}(\kappa)/[\kappa]^{<\kappa}$.

To see that Φ is non-trivial, suppose that there is a one-to-one function $F: \kappa \rightarrow \kappa$ such that $F(A) \in \Phi([A]_{\kappa})$ for all $A \subseteq \kappa$. Using hypothesis (H₃) choose $\xi \in \kappa^+$ for which there is $Z \in [\kappa]^{\kappa}$ and atoms $a_v \in \mathfrak{B}_{\xi,v}$ and $j_v \in a_v$ such that $F(j_v) \notin \Phi_{\xi,v}(a_v)$ for each $v \in Z$. Let $W \in [Z]^{\kappa}$ be such that for each $v \in W$, if $F(j_v) \in I_{\mu}$ and $\mu \neq v$ then $\mu \notin W$. Let $A = \bigcup_{v \in W} a_v$. It follows from hypothesis (H₂) that for any $\eta \geq \xi$

$$\{F(j_v) : v \in W\} \cap \bigcup_{v \in W} \Phi_{\eta,v}(a_v) \equiv_{\kappa} \{F(j_v) : v \in W\} \cap \bigcup_{v \in W} \Phi_{\xi,v}(a_v) \equiv_{\kappa} \emptyset$$

and, hence, $F(A) \notin \Phi([A]_{\kappa})$. \square

3 When are the hypotheses of Lemma 2.1 satisfied?

In answering a question of A. Blass concerning the classification of cardinal invariants of the continuum based on the Borel hierarchy Goldstern and Shelah introduced a family of cardinal invariants called $\mathbf{c}(f, g)$ defined to be the least number of uniform trees with g -splitting needed to cover a uniform tree with f -splitting [4] and showed that uncountably many of these can be distinct simultaneously. The following definition is very closely related to this as well as to the notion of a *slalom* found in [1].

Definition 3.1 Given functions f and g from $\mathbf{cof}(\kappa)$ to κ such that $g(\xi)$ is a cardinal for each $\xi \in \mathbf{cof}(\kappa)$, let $\mathfrak{d}_{f,g}$ be the least cardinal of a family $\mathcal{D} \subseteq \prod_{v \in \kappa} [f(v)]^{g(v)}$ such that for every $F \in \prod_{v \in \kappa} f(v)$ there is $G \in \mathcal{D}$ such that $F(v) \in G(v)$ for all v . Given a uniform filter \mathcal{F} on κ define $\mathfrak{d}_{f,g}(\mathcal{F})$ to be the least cardinal of a family $\mathcal{D} \subseteq \prod_{v \in \kappa} [f(v)]^{g(v)}$ such that for every $F \in \prod_{v \in \kappa} f(v)$ there is $G \in \mathcal{D}$ and $X \in \mathcal{F}$ such that $F(v) \in G(v)$ for all $v \in X$.

Lemma 3.2 Let κ be inaccessible and $f: \kappa \rightarrow \kappa$ and $g: \kappa \rightarrow \kappa$ be functions such that:

- both f and g take their values in the set of cardinals,
- $\lim_{v \rightarrow \kappa} g(v) = \kappa$,
- $2^{g(v)} < |f(v)|$ for all $v \in \kappa$,
- $\mathfrak{d}_{2^f,g}(\mathcal{F}) = \kappa^+$ for some filter \mathcal{F} generated by a \subseteq_{κ} -descending tower of length κ^+ ,

then the hypotheses of Lemma 2.1 hold.

Proof Given the hypothesis, it may be assumed that there are \subseteq_{κ} -descending sets $\{X_{\xi}\}_{\xi \in \kappa^+} \subseteq \mathcal{F}$ and functions $\{G_{\xi}\}_{\xi \in \kappa^+} \subseteq \prod_{v \in \kappa} [2^{f(v)}]^{g(v)}$ such that for every $F \in$

$\prod_{\nu \in \kappa} 2^{f(\nu)}$ there is $\xi \in \kappa^+$ such that $F(\nu) \in G_\xi(\nu)$ for all ν in a final segment of X_ξ . (This is done simply by reindexing so that for all $\xi \in \kappa^+$ there are cofinally many $\eta \in \kappa^+$ such that $G_\xi = G_\eta$.) Moreover, by a diagonal argument using the fact that $\lim_{\nu \rightarrow \kappa} g(\nu) = \kappa$ it can be assumed that if $\xi \in \eta$ then $G_\xi(\nu) \subseteq G_\eta(\nu)$ for a final segment of $\nu \in X_\eta$. (This is the part of the argument that does not extend to the case $\kappa = \omega$.)

Now let $\{I_\nu\}_{\nu \in \kappa}$ partition κ so that $|I_\nu| = f(\nu)$ and let $\{\theta_{\iota, \nu}\}_{\iota \in 2^{f(\nu)}}$ enumerate all permutations of I_ν . Let $A_\nu : 2^{f(\nu)} \rightarrow \mathcal{P}(I_\nu)$ be a bijection. Let $A_{0, \nu}$ and $A_{1, \nu}$ partition I_ν into two sets of cardinality $f(\nu)$ and let $\varphi_{0, \nu}$ be an involution of I_ν interchanging $A_{0, \nu}$ and $A_{1, \nu}$. For $\nu \in X_0$ let $\mathfrak{B}_{0, \nu} = \{\emptyset, I_\nu, A_{0, \nu}, A_{1, \nu}\}$ and let $\Phi_{0, \nu}$ be the automorphism of $\mathfrak{B}_{0, \nu}$ induced by $\varphi_{0, \nu}$. For $\nu \in \kappa \setminus X_0$ let $\mathfrak{B}_{0, \nu} = \mathcal{P}(I_\nu)$ and let $\Phi_{0, \nu}$ be the identity. As the induction hypothesis assume condition (H₂) of Lemma 2.1 holds and that, in addition,

- $\mathcal{A}_{\xi, \nu}$ are the atoms of $\mathfrak{B}_{\xi, \nu}$ and $\mathcal{A}_{\xi, \nu}$ generate $\mathfrak{B}_{\xi, \nu}$,
- $\mathcal{A}_{\xi, \nu}$ is a partition of I_ν ,
- $|\mathcal{A}_{\xi, \nu}| \leq 2^{g(\nu)}$ provided that $\nu \in X_\xi$,
- for $\nu \in X_\xi$ there are involutions $\varphi_{\xi, \nu}$ of I_ν that induce $\Phi_{\xi, \nu}$.

If $\mathfrak{B}_{\xi, \nu}, \mathcal{A}_{\xi, \nu}, \varphi_{\xi, \nu}$ and $\Phi_{\xi, \nu}$ have been defined for all ξ less than the limit ordinal η then a standard diagonalization yields $\mathfrak{B}_{\eta, \nu}, \mathcal{A}_{\eta, \nu}, \varphi_{\eta, \nu}$ and $\Phi_{\eta, \nu}$.

Therefore assume that $\mathfrak{B}_{\xi, \nu}, \mathcal{A}_{\xi, \nu}, \varphi_{\xi, \nu}$ and $\Phi_{\xi, \nu}$ have been defined. Let $\mathcal{A}_{\xi+1, \nu}^*$ be the atoms generated by $\mathcal{A}_{\xi, \nu}$ and $\{A_\nu(\iota), \varphi_{\xi, \nu}(A_\nu(\iota))\}_{\iota \in G_{\xi+1}(\nu)}$ —in other words, $\mathcal{A}_{\xi+1, \nu}^*$ consists of intersections of maximal centred subfamilies of

$$\mathcal{A}_{\xi, \nu} \cup \left\{ A_\nu(\iota) \cap \varphi_{\xi, \nu}(A_\nu(\iota)), I_\nu \setminus (A_\nu(\iota) \cup \varphi_{\xi, \nu}(A_\nu(\iota))), \right. \\ \left. \varphi_{\xi, \nu}(A_\nu(\iota)) \setminus A_\nu(\iota), A_\nu(\iota) \setminus \varphi_{\xi, \nu}(A_\nu(\iota)) \right\}_{\iota \in G_{\xi+1}(\nu)}.$$

Observe that since the elements of $\mathcal{A}_{\xi, \nu}$ are pairwise disjoint, at most one of them can belong to a centred family and so $|\mathcal{A}_{\xi+1, \nu}^*| \leq |\mathcal{A}_{\xi, \nu}| 2^{g(\nu)} \leq 2^{g(\nu)}$ for all ν in a final segment of X_ξ . Moreover, $\mathcal{A}_{\xi+1, \nu}^*$ is a partition of I_ν . Since $f(\nu) > 2^{g(\nu)}$ there must be for each $\nu \in X_\xi$ some $a_\nu \in \mathcal{A}_{\xi+1, \nu}^*$ such that $|a_\nu| > g(\nu)$. For each $\nu \in X_{\xi+1}$ let $\varphi : a_\nu \rightarrow \varphi_{\xi, \nu}(a_\nu)$ be any bijection such that for each $\iota \in G_{\xi+1}(\nu)$ there is some $k_{\iota, \nu} \in a_\nu$ such that $\varphi(k_{\iota, \nu}) \neq \theta_{\iota, \nu}(k_{\iota, \nu})$. Now for $\nu \in X_{\xi+1}$ let

$$\mathcal{A}_{\xi+1, \nu} = (\mathcal{A}_{\xi+1, \nu}^* \setminus \{a_\nu\}) \cup \{\{k_{\iota, \nu}\} : \iota \in G_{\xi+1}(\nu)\} \cup \{a_\nu \setminus \{k_{\iota, \nu}\}_{\iota \in G_{\xi+1}(\nu)}\}$$

and note that $\mathcal{A}_{\xi+1, \nu}$ is also a partition. Let $\varphi_{\xi+1, \nu}$ be defined by

$$\varphi_{\xi+1, \nu}(z) = \begin{cases} \varphi_{\xi, \nu}(z) & \text{if } z \notin a_\nu \cup \varphi_{\xi, \nu}(a_\nu), \\ \varphi(z) & \text{if } z \in a_\nu, \\ \varphi^{-1}(z) & \text{if } z \in \varphi_{\xi, \nu}(a_\nu) \end{cases}$$

and let $\Phi_{\xi+1, \nu}$ be induced by $\varphi_{\xi+1, \nu}$. Let $\mathfrak{B}_{\xi+1, \nu}$ be the Boolean algebra generated by the atoms $\mathcal{A}_{\xi+1, \nu}$. On the other hand, for $\nu \in \omega \setminus X_{\xi+1}$ let $\mathfrak{B}_{\xi+1, \nu} = \mathcal{P}(I_\nu)$ and let

$\Phi_{\xi+1, \nu}$ be induced by $\varphi_{\xi, \nu}$. It is immediate for each $\nu \in \kappa$ that $\mathfrak{B}_{\xi, \nu} \subseteq \mathfrak{B}_{\xi+1, \nu}$ and that $\Phi_{\xi+1, \nu} \upharpoonright \mathfrak{B}_{\xi, \nu} = \Phi_{\xi, \nu}$. Moreover,

$$|\mathcal{A}_{\xi+1, \nu}| \leq |\mathcal{A}_{\xi+1, \nu}^*| + g(\nu) \leq 2^{g(\nu)}$$

for all ν in a final segment of $X_{\xi+1}$ as required.

To see that hypothesis (H₃) of Lemma 2.1 holds let $F: \kappa \rightarrow \kappa$ be one-to-one. If there are cofinally many ν such that there is $z_\nu \in I_\nu$ such that $F(z_\nu) \notin I_\nu$ then let $\xi = 0$ and, without loss of generality, it may be assumed that z_ν belongs to the atom $A_{0, \nu}$ of $\mathfrak{B}_{0, \nu}$ for cofinally many ν . Since $\varphi_{0, \nu}(A_{0, \nu}) = A_{1, \nu} \subseteq I_\nu$ it is clear that $F(z_\nu) \notin \varphi_{0, \nu}(A_{0, \nu})$. If, on the other hand, $F(I_\nu) \subseteq I_\nu$ for all ν in a final segment of κ then $F \upharpoonright I_\nu = \theta_{J(\nu), \nu}$ for some $J(\nu)$ also for a final segment. There is then some $\xi \in \kappa^+$ such that $J(\nu) \in G_\xi(\nu)$ for all ν in a final segment of X_ξ and it may as well be assumed that ξ is a successor. By construction, for all ν in a final segment of X_ξ there is a singleton $\{l\} \in \mathcal{A}_{\xi, \nu}$ such that $\Phi_{\xi, \nu}(\{l\}) = \{\varphi_{\xi, \nu}(k)\}$ and $\varphi_{\xi, \nu}(l) \neq \theta_{J(\nu), \nu}(l) = F(l)$.

Finally, to see that hypothesis (H₄) of Lemma 2.1 holds let $A \subseteq \kappa$. Let $F \in \prod_{\nu \in \kappa} 2^{f(\nu)}$ be such that $A \cap I_\nu = A_\nu(F(\nu))$ for all ν . By the hypothesis of Lemma 3.2 there is $\xi \in \kappa^+$ such that $F(\nu) \in G_\xi(\nu)$ for all $\nu \in X_\xi$. It follows that $A \cap I_\nu = A_\nu(F(\nu)) \in \mathfrak{B}_\xi(\nu)$ for all ν in a final segment of X_ξ . Since $\mathfrak{B}_{\xi, \nu} = \mathcal{P}(I_\nu)$ if $\nu \notin X_\xi$ it follows that there is some $\iota \in \kappa$ such that $A \cap I_\nu \in \mathfrak{B}_{\xi, \nu}$ for all $\nu \geq \iota$. \square

4 The special case of $\kappa = \omega$

The proof of Lemma 3.2 does not apply to $\kappa = \omega$ because it relies on the fact that $\mu \cdot \mu = \mu$ if μ is an infinite cardinal. This is used to reduce to the case $G_\xi(\nu) \subseteq G_\eta(\nu)$ for most ν if $\xi \in \eta$. The first part of the proof of Lemma 4.1 corrects this—the function f in its hypothesis can be thought of as yielding a sequence of integers approximating infinite cardinals—but the general outline of the proof is the same as that of Lemma 3.2.

Lemma 4.1 *If there are functions $f: \mathbb{N} \rightarrow \mathbb{N}$ and $g: \mathbb{N} \rightarrow \mathbb{N}$ such that for all $k \in \mathbb{N}$*

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)k^{g(n)}} = \infty$$

and if $\mathfrak{d}_{f, g}(\mathcal{F}) = \aleph_1$ for some filter \mathcal{F} generated by a \subseteq^ -descending tower of length ω_1 then the hypotheses of Lemma 2.1 hold.*

Proof Given the hypothesis, it may be assumed that there are \subseteq^* -descending sets $\{X_\xi\}_{\xi \in \omega_1} \subseteq \mathcal{F}$ and functions $\{G_\xi\}_{\xi \in \omega_1} \subseteq \prod_{n \in \omega} [f(n)!]^{g(n)}$ such that for every $F \in \prod_{n \in \omega} f(n)!$ there is $\xi \in \omega_1$ such that $F(n) \in G_\xi(n)$ for all but finitely many $n \in X_\xi$. (This is done simply by reindexing so that for all $\xi \in \omega_1$ there are cofinally many $\eta \in \omega_1$ such that $G_\xi = G_\eta$.)

It will first be shown that it can be assumed that there are functions, $\bar{g}, h_\xi: \mathbb{N} \rightarrow \mathbb{N}$ and $H_\xi \in \prod_{n \in \mathbb{N}} [f(n)!]^{h_\xi(n)}$ for $\xi \in \omega_1$ such that

(a) $\lim_{n \rightarrow \infty} f(n)/\bar{g}(n)2^{\bar{g}(n)} = \infty,$

- (b) if $\xi \in \eta \in \omega_1$ then $4h_\xi \leq^* h_\eta \leq \bar{g}$,
 (c) if $\xi \in \eta \in \omega_1$ then $H_\xi(n) \subseteq H_\eta(n)$ for all but finitely many n ,
 (d) if $F \in \prod_{n \in \mathbb{N}} f(n)!$ and $F(n) \in G_\xi(n)$ for all but finitely many $n \in X_\xi$ then also $F(n) \in H_\xi(n)$ for all but finitely many $n \in X_\xi$.

To see this note that the hypothesis that $\lim_{n \rightarrow \infty} f(n)/g(n)k^{g(n)} = \infty$ for all k makes it possible to choose $h: \mathbb{N} \rightarrow \mathbb{N}$ such that

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)h(n)2^{g(n)h(n)}} = \infty$$

and $\lim_{n \rightarrow \infty} h(n) = \infty$. Then find $\bar{h}_\xi: \mathbb{N} \rightarrow \mathbb{N}$ for $\xi \in \omega_1$ such that if $\xi \in \eta$ then $4\bar{h}_\xi(n) < \bar{h}_\eta(n) < h(n)$ for all but finitely many $n \in \mathbb{N}$. Let $\bar{g}(n) = h(n)g(n)$ and note that it can be assumed that $f(n) > \bar{g}(n)2^{\bar{g}(n)}$ for all $n \in \mathbb{N}$. Define $h_\xi(n) = g(n)\bar{h}_\xi(n)$ and observe that if $\xi \in \eta \in \omega_1$ then

$$4h_\xi(n) = 4\bar{h}_\xi(n)g(n) \leq \bar{h}_\eta g(n) \leq h(n)g(n) = \bar{g}(n)$$

for all but finitely many n .

Let $H_0(n) = G_0(n)$. Given H_ξ satisfying conditions (c) and (d), define $H_{\xi+1}(n) = H_\xi(n) \cup G_{\xi+1}(n)$ and note that

$$\begin{aligned} |H_{\xi+1}(n)| &\leq |H_\xi(n)| + |G_{\xi+1}(n)| \leq h_\xi(n) + g(n) \\ &\leq (\bar{h}_\xi(n) + 1)g(n) \leq \bar{h}_{\xi+1}(n)g(n) = h_{\xi+1}(n). \end{aligned}$$

On the other hand, if η is a limit ordinal and H_ξ satisfying the desired requirements have been chosen for $\xi \in \eta$, then a diagonalization argument yields H_η such that $|H_\eta(n)| = h_\eta(n)$ and $H_\xi(n) \subseteq H_\eta(n)$ for all but finitely many n for each $\xi \in \eta$. Hence h_ξ and H_ξ satisfy conditions (b), (c) and (d).

Now let $\{I_n\}_{n \in \omega}$ partition \mathbb{N} so that $|I_n| = f(n)$ and let $\{\theta_{j,n}\}_{j \in f(n)!}$ enumerate all permutations of I_n . Let $A_n: f(n)! \rightarrow \mathcal{P}(I_n)$ be a surjection. Without loss of generality, $f(n)$ is even for each n . So let $A_{0,n}$ and $A_{1,n}$ partition I_n into two equal sized sets and let $\varphi_{0,n}$ be an involution of I_n interchanging $A_{0,n}$ and $A_{1,n}$. For $n \in X_0$ let $\mathfrak{B}_{0,n} = \{\emptyset, I_n, A_{0,n}, A_{1,n}\}$ and let $\Phi_{0,n}$ be the automorphism of $\mathfrak{B}_{0,n}$ induced by $\varphi_{0,n}$. For $n \in \omega \setminus X_0$ let $\mathfrak{B}_{0,n} = \mathcal{P}(I_n)$ and let $\Phi_{0,n}$ be the identity. As the induction hypothesis assume condition (H₃) of Lemma 2.1 holds and that, in addition,

- (e) $\mathcal{A}_{\xi,n}$ are the atoms of $\mathfrak{B}_{\xi,n}$ and that $|\mathcal{A}_{\xi,n}| \leq 2^{4h_\xi(n)}$ provided that $n \in X_\xi$,
 (f) for $n \in X_\xi$ there are involutions $\varphi_{\xi,n}$ of I_n that induce $\Phi_{\xi,n}$.

If $\mathfrak{B}_{\xi,n}$, $\mathcal{A}_{\xi,n}$, $\varphi_{\xi,n}$ and $\Phi_{\xi,n}$ have been defined for all ξ less than the limit ordinal η then a standard diagonalization yields $\mathfrak{B}_{\eta,n}$, $\mathcal{A}_{\eta,n}$, $\varphi_{\eta,n}$ and $\Phi_{\eta,n}$.

Therefore assume that $\mathfrak{B}_{\xi,n}$, $\mathcal{A}_{\xi,n}$, $\varphi_{\xi,n}$ and $\Phi_{\xi,n}$ have been defined. Let $\mathcal{A}_{\xi+1,n}^*$ be the atoms generated by $\mathcal{A}_{\xi,n}$ and $\{A_n(j), \varphi_{\xi,n}(A_n(j))\}_{j \in H_{\xi+1}(n)}$ —in other words, $\mathcal{A}_{\xi+1,n}^*$ consists of intersections of maximal centred subfamilies of $\mathcal{A}_{\xi,n} \cup \{A_n(j), \varphi_{\xi,n}(A_n(j)), I_n \setminus A_n(j), I_n \setminus \varphi_{\xi,n}(A_n(j))\}_{j \in H_{\xi+1}(n)}$. Then

$$|\mathcal{A}_{\xi+1,n}^*| \leq |\mathcal{A}_{\xi,n}| 4^{h_{\xi+1}(n)} \leq 2^{4h_\xi(n)} 2^{2h_{\xi+1}(n)} \leq 2^{3h_{\xi+1}(n)} \leq 2^{\bar{g}(n)}$$

for all but finitely many $n \in X_\xi$. Since $f(n) > \bar{g}(n)2^{\bar{g}(n)}$ there must be some $a_n \in \mathcal{A}_{\xi+1,n}^*$ such that $|a_n| > \bar{g}(n)$ for each $n \in X_\xi$. Let $\varphi: a_n \rightarrow \varphi_{\xi,n}(a_n)$ be any bijection such that for each $n \in X_{\xi+1}$ and each $j \in H_{\xi+1}(n)$ there is some $k_{j,n} \in a_n$ such that $\varphi(k_{j,n}) \neq \theta_{j,n}(k_{j,n})$. Now for $n \in X_{\xi+1}$ let $\mathcal{A}_{\xi+1,n} = \mathcal{A}_{\xi+1,n}^* \cup \{k_{j,n} : j \in H_{\xi+1}(n)\}$ and let $\varphi_{\xi+1,n}$ be defined by

$$\varphi_{\xi+1,n}(z) = \begin{cases} \varphi_{\xi,n}(z) & \text{if } z \notin a_n \cup \varphi_{\xi,n}(a_n), \\ \varphi(z) & \text{if } z \in a_n, \\ \varphi^{-1}(z) & \text{if } z \in \varphi_{\xi,n}(a_n) \end{cases}$$

and let $\Phi_{\xi+1,n}$ be induced by $\varphi_{\xi+1,n}$. Let $\mathfrak{B}_{\xi+1,n}$ be the Boolean algebra whose atoms are $\mathcal{A}_{\xi+1,n}$. On the other hand, for $n \in \omega \setminus X_{\xi+1}$ let $\mathfrak{B}_{\xi+1,n} = \mathcal{P}(I_n)$ and let $\Phi_{\xi+1,n}$ be induced by $\varphi_{\xi,n}$. It is immediate for each $n \in \omega$ that $\mathfrak{B}_{\xi,n} \subseteq \mathfrak{B}_{\xi+1,n}$ and that $\Phi_{\xi+1,n} \upharpoonright \mathfrak{B}_{\xi,n} = \Phi_{\xi,n}$. Moreover,

$$|\mathcal{A}_{\xi+1,n}| \leq |\mathcal{A}_{\xi+1,n}^*| + h_{\xi+1}(n) \leq 2^{3h_{\xi+1}(n)} + h_{\xi+1}(n) \leq 2^{4h_{\xi+1}(n)}$$

for all but finitely many $n \in X_{\xi+1}$ as required.

To see that hypothesis (H₄) of Lemma 2.1 holds let $F: \mathbb{N} \rightarrow \mathbb{N}$ be one-to-one. If there are infinitely many n such that there is $z_n \in I_n$ such that $F(z_n) \notin I_n$ then let $\xi = 0$ and, without loss of generality, it may be assumed that z_n belongs to the atom $A_{0,n}$ of $\mathfrak{B}_{0,n}$ for infinitely many n . Since $\varphi_{0,n}(A_{0,n}) = A_{1,n} \subseteq I_n$ it is clear that $F(z_n) \notin \varphi_{0,n}(A_{0,n})$. If, on the other hand, $F(I_n) \subseteq I_n$ for all but finitely many n then $F \upharpoonright I_n = \theta_{J(n),n}$ for some $J(n)$ also for all but finitely many n . By condition (d) there is some $\xi \in \omega_1$ such that $J(n) \in H_\xi(n)$ for all but finitely many $n \in X_\xi$ and condition (c) allows the assumption that ξ is a successor. By construction, for all but finitely many $n \in X_\xi$ there is a singleton $\{k\} \in \mathcal{A}_{\xi,n}$ such that $\Phi_{\xi,n}(\{k\}) = \{\varphi_{\xi,n}(k)\}$ and $\varphi_{\xi,n}(k) \neq \theta_{J(n),n}(k) = F(k)$.

Finally, to see that hypothesis (H₅) of Lemma 2.1 holds let $A \subseteq \mathbb{N}$. Let $F \in \prod_{n \in \mathbb{N}} f(n)!$ be such that $A \cap I_n = A_n(F(n))$ for all n . By the hypothesis of Lemma 3.2 there is $\xi \in \omega_1$ such that $F(n) \in G_\xi(n)$ for all $n \in X_\xi$. From condition (d) it follows that $F(n) \in H_\xi(n)$ for all but finitely many $n \in X_\xi$. Since $\mathfrak{B}_{\xi,n} = \mathcal{P}(I_n)$ if $n \notin X_\xi$ it follows that $A \cap I_n \in \mathfrak{B}_{\xi,n}$ for all but finitely many $n \in \mathbb{N}$. \square

Corollary 4.2 *If $\mathfrak{d}_{f^1,g} = \aleph_1$ then there is a non-trivial isomorphism of $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$.*

Proof In this case let \mathcal{F} be the co-finite filter and note it is generated by the constant \subseteq^* -descending sequence all of whose terms are ω . \square

Corollary 4.3 *If there is an \aleph_1 -generated filter \mathcal{F} such that $\mathfrak{d}_{f^1,g}(\mathcal{F}) = \aleph_1 \neq \mathfrak{d}$ then there is a non-trivial isomorphism of $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$.*

Proof Let \mathcal{F} be generated by $\{X_\xi\}_{\xi \in \omega_1}$. Use Rothberger's argument and $\aleph_1 \neq \mathfrak{d}$ to construct a \subseteq^* -descending sequence $\{Y_\xi\}_{\xi \in \omega_1}$ all of whose terms are \mathcal{F} positive and such that $Y_\xi \subseteq X_\xi$. Let \mathcal{F}' be generated by $\{Y_\xi\}_{\xi \in \omega_1}$ and note that $\mathfrak{d}_{f^1,g}(\mathcal{F}') = \aleph_1$. \square

It has to be noted that the hypothesis of Corollary 4.3 is not vacuous in the sense that there are models of set theory in which it holds. For example, in the model obtained by iterating Miller reals ω_2 times the following hold:

- $\mathfrak{d} = \aleph_2$ because the Miller reals themselves are unbounded by the ground model,
- $\mathfrak{d}_{f,g} = \aleph_1$ for appropriate f and g because the Miller partial order satisfies the Laver property,
- $\mathfrak{u} = \aleph_1$ because P-points from the ground model generate ultrafilters in the extension.

However there does not seem to be any model demonstrating that the assumption that $\aleph_1 \neq \mathfrak{d}$ in Corollary 4.3 is essential. It is shown in [3] that it is consistent with set theory that $\mathfrak{d} = \aleph_1$ yet all automorphisms of $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$ are trivial. However, $\mathfrak{u} = \aleph_2$ in that model because random reals are added cofinally often.

5 Remarks and questions

The first thing to note is that there are models where $\mathfrak{d}_{f!,g} = \aleph_1 < 2^{\aleph_0}$ for f and g satisfying the hypotheses of Lemma 4.1—for example, this is true in the model obtained by either iteratively adding ω_2 Sacks reals¹ or adding any number of Sacks reals, greater than \aleph_1 of course, side-by-side. Of course $\mathfrak{d} = \aleph_1$ also in these models. It is therefore of interest to note that the Laver property implies that $\mathfrak{d}_{f!,g} = \aleph_1$ in the Laver model as well, yet $\mathfrak{d} = \aleph_2$ in this model. It should also be observed that it is possible for $\mathfrak{d}_{f,g}$ to be larger than \mathfrak{d} . For example, iteratively forcing ω_2 times with perfect trees T that are cofinally f branching will yield such a model.

To be a bit more precise, given $f: \omega \rightarrow \omega$ define $\mathbb{S}(f)$ to consist of all trees $T \subseteq \bigcup_{n \in \omega} \prod_{j \in n} f(j)$ such that for each $t \in T$ there is $s \supseteq t$ such that $s \cap j \in T$ for all $j \in f(|s|)$. So Sacks forcing is just $\mathbb{S}(2)$ where 2 is the constant 2 function. The same proof as for Sacks forcing shows that $\mathbb{S}(f)$ is proper and adds no reals unbounded by the ground model. Iterating $\mathbb{S}(f)$ with countable support ω_2 times then yields model in which $\mathfrak{d} = \aleph_1$. However, if $g: \omega \rightarrow \omega$ and $\mathcal{H} \subseteq \prod_{n \in \omega} [f(n)]^{g(n)}$ has cardinality \aleph_1 then there is some model containing g and \mathcal{H} and there is $\Gamma \in \prod_{n \in \omega} f(n)$ which is generic over this model. This generically ensures that for all $h \in \mathcal{H}$ there is some j such that $\Gamma(j) \notin h(j)$.

The generalization of Sacks reals to uncountable cardinals in [5] establishes that the hypotheses of Lemma 3.2 can be satisfied for uncountable cardinals. Alternatively, one could iteratively force κ^+ times with the partial order $\mathbb{P}(f, g)$ consisting of pairs (G, \mathcal{F}) where $|G| < \kappa$ and G is a function whose domain is a subset of κ and $G(\xi) \in [f(\xi)]^{g(\xi)}$ and $\mathcal{F} \subseteq \prod_{\xi \in \kappa} f(\xi)$ and $|\mathcal{F}| < \kappa$ with the ordering defined by $(G, \mathcal{F}) \leq (G', \mathcal{F}')$ if

- $G \supseteq G'$,
- $\mathcal{F} \supseteq \mathcal{F}'$,
- if ξ is in the in the domain of $G \setminus G'$ and $f \in \mathcal{F}'$ then $f(\xi) \in G(\xi)$.

¹ See [1] for definitions of terms not defined in this section as well as for details of proofs.

It is easy to see that $\mathbb{P}(f, g)$ is κ -closed and that if $\Gamma \subseteq \mathbb{P}(f, g)$ is generic then the domain of $\bigcup_{(G, \mathcal{F}) \in \Gamma} G$ has cardinality κ and the family of such domains added iteratively will generate the necessary filter.

However, the following question does not seem to be answered.

Question 5.1 Is it consistent for an inaccessible cardinal κ that $\mathfrak{d}_{f,g} = \kappa^+$ where $f(\alpha) = (2^{\aleph_\alpha})^+$ and $g(\alpha) = \aleph_\alpha$ yet $\mathfrak{d}(\kappa) > \kappa^+$ where $\mathfrak{d}(\kappa)$ is the generalization of \mathfrak{d} to κ ?

It is worth observing that the isomorphism of Lemma 2.1 is trivial on some infinite sets—indeed, if $\xi \in \kappa^+$ and $X \subseteq \mathbb{N}$ are such that $\{x\}$ belongs to some $\mathfrak{B}_{\xi,v}$ for each $x \in X$ then Φ is trivial on $\mathcal{P}(X)$. However, if $\mathcal{I}(\Phi)$ is defined to be the ideal $\{X \subseteq \mathbb{N} : \Phi \upharpoonright \mathcal{P}(X) \text{ is trivial}\}$ then $\mathcal{I}(\Phi)$ is a small ideal in the sense that the quotient algebra $\mathcal{P}(\mathbb{N})/\mathcal{I}(\Phi)$ has large antichains, even modulo the ideal of finite sets—in the terminology of [2], the ideal $\mathcal{I}(\Phi)$ is not ccc by fin. To see this, simply observe that the proof of Lemma 4.1 actually shows that hypothesis (H₄) of Lemma 2.1 can be strengthened to: For any one-to-one $F: \mathbb{N} \rightarrow \mathbb{N}$ there is $\xi \in \kappa^+$ such that for all but finitely many $v \in \omega$ there is an atom $a \in \mathfrak{B}_{\xi,v}$ and $\iota \in a$ such that $F(\iota) \notin \Phi_{\xi,v}(a)$. It follows that if $Z \subseteq \mathbb{N}$ is infinite then $Z^* = \bigcup_{v \in Z} I_v \notin \mathcal{I}(\Phi)$. Hence, if \mathcal{A} is an almost disjoint family of subsets of \mathbb{N} then $\{A^* : A \in \mathcal{A}\}$ is an antichain modulo the ideal of finite sets.

One should not, therefore, expect to get a nowhere trivial isomorphism by these methods. It is nevertheless, conceivable that there are some other cardinal invariants similar to $\mathfrak{d}_{f,g}$ that would, when small, imply the existence of nowhere trivial isomorphisms of $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$. In this context it is interesting to note that it is at least consistent with small \mathfrak{d} that there are nowhere trivial isomorphisms.

Proposition 5.2 *It is consistent that $\aleph_1 = \mathfrak{d} \neq 2^{\aleph_0}$ and there is a nowhere trivial isomorphism of $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$.*

Sketch of proof The partial order defined in [12, Section 2, Definition 2.1] will be used². Begin with a model V satisfying $2^{\aleph_0} > \aleph_1$ and construct a tower of permutations $\{(A_\xi, F_\xi, \mathfrak{B}_\xi)\}_{\xi \in \text{Lim}(\omega_1)}$ such that, letting $\mathfrak{S}_\eta = \{(A_\xi, F_\xi, \mathfrak{B}_\xi)\}_{\xi \in \text{Lim}(\eta)}$ and \mathbb{P}_η be the finite support iteration of partial orders that are $\mathbb{Q}(\mathfrak{S}_\xi)$ for $\xi \in \text{Lim}(\eta)$ and Hechler forcing if ξ is a successor, the following holds for each η and G that is \mathbb{P}_{ω_1} generic over V :

- $A_\eta = A_{\mathfrak{S}_\eta}[G \cap \mathbb{Q}(\mathfrak{S}_\eta)]$,
- $F_\eta = F_{\mathfrak{S}_\eta}[G \cap \mathbb{Q}(\mathfrak{S}_\eta)]$,
- $\mathfrak{B}_\eta = \mathcal{P}(\mathbb{N}) \cap V[G \cap \mathbb{P}_\eta]$.

The proof of [12, Theorem 2.1] shows that there is a nowhere trivial isomorphism of $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$ in this model and, since \mathbb{P}_{ω_1} is ccc, it is also true that 2^{\aleph_0} remains larger than \aleph_1 in the generic extension. The Hechler reals guarantee that $\mathfrak{d} = \aleph_1$. \square

It should also be noted that Lemma 2.1 actually yields $2^{(\kappa^+)}$ isomorphisms. It is shown in [13] that it is possible to have non-trivial isomorphisms of $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$

² The reader is warned that the word “finite” should be removed from of [12, Definition 2.1 (3)].

without having 2^c such isomorphisms. This motivates the following, somewhat vague, question.

Question 5.3 Can there be some variant of $\mathfrak{d}_{f,g}$ which, when small, yields a non-trivial isomorphism of $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$ without yielding the maximal possible number of such?

Given the remarks following Corollary 4.3 it is natural to ask the following.

Question 5.4 Is it consistent that $\mathfrak{d}_{f^!,g} = \mathfrak{d}$ for f and g satisfying the hypothesis of Lemma 4.1 and to have $\mathfrak{u} = \aleph_1$ and to have that all isomorphisms of $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$ are trivial?

As a final remark it will be noted that Corollary 4.2 shows that [12, Theorem 3.1] cannot be improved to show that in models obtained by iterating Sacks or Silver reals all isomorphisms of $\mathcal{P}(\mathbb{N})/[\mathbb{N}]^{<\aleph_0}$ are trivial because the equality $\mathfrak{d}_{f^!,g} = \aleph_1$ holds in these models for the necessary f and g .

References

1. Bartoszyński, T., Judah, H.: Set Theory. On the Structure of the Real Line. A K Peters, Wellesley (1995)
2. Farah, I.: Analytic Quotients: Theory of Liftings for Quotients over Analytic Ideals on the Integers. *Memoirs of the American Mathematical Society*, vol. 148(702). American Mathematical Society, Providence (2000)
3. Farah, I., Shelah, S.: Trivial automorphisms. *Israel J. Math.* **201**(2), 701–728 (2014)
4. Goldstern, M., Shelah, S.: Many simple cardinal invariants. *Arch. Math. Logic* **32**(3), 203–221 (1993)
5. Kanamori, A.: Perfect-set forcing for uncountable cardinals. *Ann. Math. Logic* **19**(1–2), 97–114 (1980)
6. Parovičenko, I.I.: On a universal bicomactum of weight \aleph . *Dokl. Akad. Nauk SSSR* **150**, 36–39 (1963)
7. Rudin, W.: Homogeneity problems in the theory of Čech compactifications. *Duke Math. J.* **23**(3), 409–419 (1956)
8. Rudin, W.: Note of correction: “Homogeneity problems in the theory of Čech compactifications,” vol. 23 (1956), pp. 409–419. *Duke Math. J.* **23**(6), 633 (1956)
9. Shelah, S.: Proper Forcing. *Lecture Notes in Mathematics*, vol. 940. Springer, Berlin (1982)
10. Shelah, S., Steprāns, J.: PFA implies all automorphisms are trivial. *Proc. Amer. Math. Soc.* **104**(4), 1220–1225 (1988)
11. Shelah, S., Steprāns, J.: Somewhere trivial autohomeomorphisms. *J. London Math. Soc.* **49**(3), 569–580 (1994)
12. Shelah, S., Steprāns, J.: Martin’s axiom is consistent with the existence of nowhere trivial automorphisms. *Proc. Amer. Math. Soc.* **130**(7), 2097–2106 (2002)
13. Steprāns, J.: The autohomeomorphism group of the Čech–Stone compactification of the integers. *Trans. Amer. Math. Soc.* **355**(10), 4223–4240 (2003)
14. Veličković, B.: OCA and automorphisms of $\mathcal{P}(\omega)/\text{fin}$. *Topology Appl.* **49**(1), 1–13 (1993)