

## TEST GROUPS FOR WHITEHEAD GROUPS

PAUL C. EKLOF, LÁSZLÓ FUCHS AND SAHARON SHELAH

**ABSTRACT.** We consider the question of when the dual of a Whitehead group is a test group for Whitehead groups. This turns out to be equivalent to the question of when the tensor product of two Whitehead groups is Whitehead. We investigate what happens in different models of set theory.

**1. Introduction.** All groups in this note are abelian. A *Whitehead group*, or *W-group* for short, is defined to be an abelian group  $A$  such that  $\text{Ext}(A, \mathbf{Z}) = 0$ . We are looking for groups  $C$  other than  $\mathbf{Z}$  such that a group  $A$  is a  $W$ -group if and only if  $\text{Ext}(A, C) = 0$ ; such a  $C$  will be called a *test group for Whitehead groups*, or a *W-test group* for short. Notice that, if  $C$  is a non-zero separable torsion-free group, then  $\text{Ext}(A, C) = 0$  implies  $A$  is a  $W$ -group (since  $\mathbf{Z}$  is a summand of  $C$ ), but the converse may not hold, that is,  $\text{Ext}(A, C)$  may be non-zero for some  $W$ -group  $A$ .

Among the separable torsion-free groups are the dual groups, where by a dual group we mean one of the form  $\text{Hom}(B, \mathbf{Z})$  for some group  $B$ . We call  $\text{Hom}(B, \mathbf{Z})$  the ( $\mathbf{Z}$ -)dual of  $B$  and denote it by  $B^*$ . We shall call a group a *W\*-group* if it is the dual of a  $W$ -group. The principal question we will consider is whether every  $W^*$ -group is a  $W$ -test group. This turns out to be equivalent to a question about tensor products of  $W$ -groups. (See Corollary 2.4.)

As is almost always the case with problems related to Whitehead groups, the answer depends upon the chosen model of set theory. We have an easy affirmative answer if every  $W$ -group is free (for example in a model of  $V = L$ ); therefore, we will focus on models where there are non-free  $W$ -groups. We will exhibit models with differing results about whether  $W^*$ -groups are  $W$ -test groups, including information about

---

2010 AMS *Mathematics subject classification.* Primary 20K20, Secondary 03E35, 20A15, 20K35, 20K40.

*Keywords and phrases.* Whitehead group, dual group, tensor product.

The third author would like to thank the United States-Israel Binational Science Foundation for their support. Publication 879.

Received by the editors on December 3, 2009, and in revised form on March 15, 2010.

DOI:10.1216/RMJ-2012-42-6-1863 Copyright ©2012 Rocky Mountain Mathematics Consortium

some of the “classical” models where there are non-free  $W$ -groups. In particular, we will show that the answer to the question is independent of  $\text{ZFC} + \text{GCH}$ .

**2. Theorems of ZFC.** We begin with some theorems of ZFC.

**Theorem 2.1.** *A group  $A$  is a  $W^*$ -group if and only if it is the kernel of an epimorphism  $\mathbf{Z}^\kappa \rightarrow \mathbf{Z}^\lambda$  for some cardinals  $\kappa$  and  $\lambda$ .*

*Proof.* If  $A = B^*$  where  $B$  is a  $W$ -group, choose a free resolution

$$0 \longrightarrow K \longrightarrow F \longrightarrow B \longrightarrow 0$$

where  $F$  and  $K$  are free groups. Taking the dual yields

$$0 \longrightarrow B^* = A \longrightarrow F^* \longrightarrow K^* \longrightarrow \text{Ext}(B, \mathbf{Z}) = 0.$$

Since  $F^*$  and  $K^*$  are products, this proves the claim in one direction.

Conversely, if  $H$  is the kernel of an epimorphism  $\phi : \mathbf{Z}^\kappa \rightarrow \mathbf{Z}^\lambda$ , then we have an exact sequence

$$0 \longrightarrow H \longrightarrow F^* \longrightarrow K^* \longrightarrow 0$$

for certain free groups  $F$  and  $K$ . Dualizing, we obtain an exact sequence

$$0 \longrightarrow K \longrightarrow F \longrightarrow B \longrightarrow 0$$

for a subgroup  $B$  of  $H^*$ . When we take the dual of the last sequence, we obtain the original epimorphism  $\phi$  and conclude that its kernel is  $B^*$  and  $\text{Ext}(B, \mathbf{Z}) = 0$ .  $\square$

**Theorem 2.2.** *For  $W$ -groups  $A$  and  $B$ , there are natural isomorphisms*

$$\text{Ext}(A, B^*) \cong \text{Ext}(B, A^*)$$

and

$$\text{Ext}(A, B^*) \cong \text{Ext}(A \otimes B, \mathbf{Z}).$$

*Proof.* Recall that, for any pair of groups  $A$  and  $B$ , there are natural isomorphisms

$$\mathrm{Hom}(A, B^*) \cong \mathrm{Hom}(A \otimes B, \mathbf{Z}) \cong \mathrm{Hom}(B \otimes A, \mathbf{Z}) \cong \mathrm{Hom}(B, A^*).$$

Now let  $A$  and  $B$  be  $W$ -groups. Using these isomorphisms as vertical maps and a free resolution  $0 \rightarrow K \rightarrow F \rightarrow A \rightarrow 0$ , we can form a commutative diagram with exact rows as follows:

$$\begin{array}{ccccccccc} 0 \rightarrow & \mathrm{Hom}(A, B^*) \rightarrow & \mathrm{Hom}(F, B^*) \rightarrow & \mathrm{Hom}(K, B^*) \rightarrow & \mathrm{Ext}(A, B^*) & \rightarrow & 0 \\ & \downarrow & \downarrow & \downarrow & & & \\ 0 \rightarrow & \mathrm{Hom}(A \otimes B, \mathbf{Z}) \rightarrow & \mathrm{Hom}(F \otimes B, \mathbf{Z}) \rightarrow & \mathrm{Hom}(K \otimes B, \mathbf{Z}) \rightarrow & \mathrm{Ext}(A \otimes B, \mathbf{Z}) & \rightarrow & 0 \\ & \downarrow & \downarrow & \downarrow & \downarrow & & \\ 0 \rightarrow & \mathrm{Hom}(B \otimes A, \mathbf{Z}) \rightarrow & \mathrm{Hom}(B \otimes F, \mathbf{Z}) \rightarrow & \mathrm{Hom}(B \otimes K, \mathbf{Z}) \rightarrow & \mathrm{Ext}(B \otimes A, \mathbf{Z}) & \rightarrow & 0 \\ & \downarrow & \downarrow & \downarrow & & & \\ 0 \rightarrow & \mathrm{Hom}(B, A^*) \rightarrow & \mathrm{Hom}(B, F^*) \rightarrow & \mathrm{Hom}(B, K^*) \rightarrow & \mathrm{Ext}(B, A^*) & \rightarrow & 0 \end{array}$$

The 0's at the end of the four rows are justified by the freeness of  $F$  and by the fact that  $F \otimes B$ ,  $B \otimes F$  and  $B$  are  $W$ -groups, respectively. All of the horizontal and vertical maps are natural isomorphisms. Consequently, the diagram can be completed, preserving commutativity, by natural maps between the two top Exts and the two bottom Exts. These maps provide the desired natural isomorphisms.  $\square$

As an immediate corollary we obtain:

**Corollary 2.3.** *Let  $A$  and  $B$  be  $W$ -groups. The following are equivalent:*

- (i)  $A \otimes B$  is a  $W$ -group;
- (ii)  $\mathrm{Ext}(A, B^*) = 0$ ;
- (iii)  $\mathrm{Ext}(B, A^*) = 0$ .  $\square$

Hence, we have

**Corollary 2.4.** *In any model of ZFC, the following are equivalent:*

- (i) the tensor product of any two  $W$ -groups is a  $W$ -group;
- (ii) every  $W^*$ -group is a  $W$ -test group.  $\square$

The following proposition demonstrates that the hypothesis that  $A$  and  $B$  are  $W$ -groups is necessary for the equivalent conditions of Corollary 2.3 to hold.

**Proposition 2.5.** *For any abelian groups  $A$  and  $B$ , if either*

- (i)  $A \otimes B$  is a non-zero  $W$ -group, or
- (ii)  $\text{Ext}(A, B^*) = 0$  and  $B^*$  is non-zero,

*then  $A$  is a  $W$ -group.*

*Proof.* For (ii), the conclusion follows since  $B^*$  is separable. For (i), we use the fact ([2, page 116]) that there is a (non-natural) isomorphism

$$\begin{aligned} \text{Ext}(A, \text{Hom}(B, \mathbf{Z})) \oplus \text{Hom}(A, \text{Ext}(B, \mathbf{Z})) \\ \cong \text{Ext}(A \otimes B, \mathbf{Z}) \oplus \text{Hom}(\text{Tor}(A, B), \mathbf{Z}). \end{aligned}$$

In our case the right-hand side reduces to  $\text{Ext}(A \otimes B, \mathbf{Z})$ , so the hypothesis implies that  $\text{Ext}(A, B^*) = 0$ . It suffices then to show that  $B^* \neq 0$ , but this follows from the fact that  $A \otimes B$  is a  $W$ -group, hence is separable, and is also assumed to be non-zero.  $\square$

**3. An independence result.** In this section we will show that there are: (1) a model of ZFC where the equivalent conditions of Corollary 2.4 fail; and (2) a model of ZFC where the conditions of Corollary 2.4 hold. In particular, we will exhibit

(1) a model of ZFC + GCH such that there is a  $W$ -group  $B$  of cardinality  $\aleph_1$  such that  $B^*$  is not a test group for  $W$ -groups of cardinality  $\aleph_1$ , and

(2) another model of ZFC + GCH in which there are non-free  $W$ -groups and for every  $W$ -group  $B$  (of arbitrary cardinality),  $B^*$  is a test group for  $W$ -groups (of arbitrary cardinality).

**Model (1).** For the first model, we use the model in [6, Theorem 0.5] in which a non-reflexive  $W$ -group exists. In fact, the following theorem is proved there:

**Theorem 3.1.** *It is consistent with ZFC + GCH that there is a non-free  $W$ -group  $B$  of cardinality  $\aleph_1$  such that  $B^*$  is free.*

*Proof.* Let  $B$  be as in the theorem. Note that  $B^*$  must be of infinite rank, since  $B$  is not free, but there is a monomorphism  $: B \rightarrow B^{**}$ . Since  $B$  is a non-reflexive  $W$ -group, a result of Huber (see [8] or [5, XI.2.7]) implies that  $B$  is not  $\aleph_1$ -coseparable, i.e.,  $\text{Ext}(B, \mathbf{Z}^{(\omega)}) \neq 0$ . Thus,  $\text{Ext}(B, B^*) \neq 0$ .  $\square$

**Model (2).** We shall work in a model of  $\text{Ax}(S) + \diamond^*(\omega_1 \setminus S)$  plus  $(\ddagger) \diamond_\kappa(E)$  holds for every regular cardinal  $\kappa > \aleph_1$  and every stationary subset  $E$  of  $\kappa$ .

For the definition and implications of  $\text{Ax}(S) + \diamond^*(\omega_1 \setminus S)$ , see [5, pages 178–179 and Theorem XII.2.1]. This is the “classical” model of ZFC + GCH in which there are non-free  $W$ -groups of cardinality  $\aleph_1$ ; in fact, whether an  $\aleph_1$ -free group of cardinality  $\aleph_1$  is a  $W$ -group is determined by its  $\Gamma$ -invariant.

**Theorem 3.2.** *Assuming  $\text{Ax}(S) + \diamond^*(\omega_1 \setminus S)$ , if  $A$  and  $B$  are  $W$ -groups of cardinality  $\leq \aleph_1$ , then  $A \otimes B$  is a  $W$ -group.*

*Proof.* The case when either  $A$  or  $B$  is countable (hence free) is trivial so we can assume that  $A$  and  $B$  have cardinality  $\aleph_1$ . By Theorem XII.2.1 of [5],  $\Gamma(A) \leq \tilde{S}$  and  $\Gamma(B) \leq \tilde{S}$  since  $A$  and  $B$  are  $W$ -groups. It suffices to prove that  $\Gamma(A \otimes B) \leq \tilde{S}$ . Fix  $\omega_1$ -filtrations  $\{A_\nu : \nu < \omega_1\}$  and  $\{B_\nu : \nu < \omega_1\}$  of  $A$  and  $B$ , respectively, such that

$$\{\nu < \omega_1 : \text{there exists a } \mu > \nu \text{ such that } A_\mu/A_\nu \text{ is not free}\} \subseteq S,$$

and similarly for  $B$ . Then  $\{A_\nu \otimes B_\nu : \nu < \omega_1\}$  is an  $\omega_1$ -filtration of  $A \otimes B$ . For each  $\mu > \nu$ , there is an exact sequence

$$0 \longrightarrow A_\nu \otimes B_\nu \longrightarrow A_\mu \otimes B_\mu \longrightarrow (A_\mu \otimes (B_\mu/B_\nu)) \oplus ((A_\mu/A_\nu) \otimes B_\mu)$$

(cf. [2, Proposition 4.3a (c), page 25]). If  $\nu \notin S$ , then the two summands on the right are free, and hence the quotient  $A_\mu \otimes B_\mu/A_\nu \otimes B_\nu$  is free. Thus, we have proved that  $\Gamma(A \otimes B) \leq \tilde{S}$ .  $\square$

The following is proved by the methods of proof of Theorem 3.1 in [1] (see also Corollary XII.1.13 of [5]).

**Lemma 3.3.** *In any model of  $(\ddagger)$ , every  $W$ -group  $A$  is the union of a continuous chain of subgroups  $\{A_\nu : \nu < \sigma\}$  such that  $A_0 = 0$  and for all  $\nu < \sigma$ ,  $A_\nu$  is a  $W$ -group and  $A_{\nu+1}/A_\nu$  has cardinality  $\leq \aleph_1$  and is a  $W$ -group.  $\square$*

The hypothesis  $\text{Ax}(S) + \diamond^*(\omega_1 \setminus S)$  implies that the second assumption of the following theorem holds. Thus, we will show that our model has the desired property if we prove the following theorem.

**Theorem 3.4.** *Assume that  $(\ddagger)$  holds and that the tensor product of two W-groups of cardinality  $\leq \aleph_1$  is again a W-group. Then the tensor product of any two W-groups (of arbitrary cardinality) is again a W-group.*

*Proof.* Let  $A$  and  $B$  be W-groups. It suffices to prove that  $A \otimes B$  is a W-group when at least one of the groups, say  $B$ , has cardinality  $> \aleph_1$ . Let  $\{B_\nu : \nu < \sigma\}$  be a continuous chain as in the conclusion of Lemma 3.3. We shall first prove that, if  $|A| = \aleph_1$ , then  $A \otimes B$  is a W-group. Now  $\{A \otimes B_\nu : \nu < \sigma\}$  is a continuous filtration of  $A \otimes B$ . To show that  $A \otimes B$  is a W-group, it suffices to show that, for all  $\nu < \sigma$ ,  $A \otimes B_{\nu+1}/A \otimes B_\nu$  is a W-group (cf. [5, XII.1.5]). Now, as above, there is an exact sequence

$$0 \longrightarrow A \otimes B_\nu \longrightarrow A \otimes B_{\nu+1} \longrightarrow A \otimes (B_{\nu+1}/B_\nu) \longrightarrow 0$$

since  $A$  is torsion-free. By hypothesis, the right-hand term is a W-group (of cardinality  $\aleph_1$ ); hence  $A \otimes B_{\nu+1}/A \otimes B_\nu$  is a W-group.

Next, suppose that  $|A| \geq \aleph_1$ . Again we use the continuous filtration  $\{A \otimes B_\nu : \nu < \sigma\}$  and the displayed exact sequence above. Now the right-hand term  $A \otimes (B_{\nu+1}/B_\nu)$  is a W-group by the first case (and the symmetry of the tensor product) because it is a tensor product of two W-groups one of which has cardinality  $\aleph_1$ .  $\square$

*Remark.* As mentioned above, Martin Huber has shown that every  $\aleph_1$ -coseparable group is reflexive. In Model (2) above and also in the model of the next section, all W-groups are  $\aleph_1$ -coseparable. This raises the question of whether (provably in ZFC) every  $\aleph_1$ -coseparable group, or every reflexive group,  $B$  satisfies  $\text{Ext}(B, B^*) = 0$ , or even satisfies:  $B^*$  is a W-test group.

**4. Martin's axiom.** A model of Martin's Axiom (MA) was the first model in which it was proved (in [9]) that there are non-free W-groups. So it is of interest to see what happens in that model. In fact,

the conclusion is the same as that of Theorem 3.2. But we prove it by proving the following theorem:

**Theorem 4.1.** *(MA +  $\neg$ CH) If  $A$  and  $B$  are W-groups of cardinality  $\leq \aleph_1$ , then  $\text{Ext}(A, B^*) = 0$ .*

*Proof.* The proof does not make use directly of the definition of Whitehead groups, but rather of the identification of W-groups of cardinality  $\aleph_1$  as Shelah groups, which holds under the set-theoretic hypotheses here (but not in all models of ZFC). See [10] or [5, XII.2.5, XIII.3.6]. We fix a short exact sequence

$$0 \rightarrow B^* \xrightarrow{\iota} N \xrightarrow{\pi} A \rightarrow 0$$

and proceed to prove that it splits. We may assume that  $\iota$  is the inclusion map. We also fix a set function  $\gamma : A \rightarrow N$  such that, for all  $a \in A$ ,  $\pi(\gamma(a)) = a$ . We will show that the short exact sequence splits by proving the existence of a function  $h : A \rightarrow B^*$  such that the function

$$\gamma - h : A \rightarrow N : a \mapsto \gamma(a) - h(a)$$

is a homomorphism. The function  $h$  will be obtained via a directed subset  $\mathcal{G}$  of a c.c.c. poset  $P$ ; the directed subset (which will be required to intersect certain dense subsets of  $P$ ) will exist as a consequence of MA.

We define  $P$  to consist of all triples  $p = (A_p, B_p, h_p)$  where  $A_p$  (respectively,  $B_p$ ) is a finitely generated summand of  $A$  (respectively,  $B$ ) and  $h_p$  is a function from  $A_p$  to  $B_p^*$ . Moreover, we require that the function which takes  $a \in A_p$  to  $\gamma(a) - h_p(a)$  is a homomorphism from  $A_p$  into  $N/\{f \in B^* : f \upharpoonright B_p \equiv 0\}$ . (Strictly speaking, this is an abuse of notation: by  $\gamma(a) - h_p(a)$  we mean the coset of  $\gamma(a) - \eta_a$  where  $\eta_a$  is any element of  $B^*$  such that  $\eta_a \upharpoonright B_p = h_p(a)$ .)

The partial ordering on  $P$  is defined as follows:  $p = (A_p, B_p, h_p) \leq p' = (A'_p, B'_p, h'_p)$  if and only if  $A_p \subseteq A'_p$ ,  $B_p \subseteq B'_p$  and  $h'_p(a) \upharpoonright B_p = h_p(a)$  for all  $a \in A_p$ . The dense subsets that we use are:

$$D_a^1 = \{p \in P : a \in A_p\}$$

for all  $a \in A$  and

$$D_b^2 = \{p \in P : b \in B_p\}$$

for all  $b \in B$ . Assuming that these sets are dense and that  $P$  is c.c.c., the axiom  $\text{MA} + \neg\text{CH}$  yields a directed subset  $\mathcal{G}$  which has non-empty intersection with each of these dense subsets. We can then define  $h$  by:  $h(a)(b) = h_p(a)(b)$  for some (all)  $p \in \mathcal{G}$  such that  $a \in A_p$  and  $b \in B_p$ . It is easy to check that  $h$  is well-defined and has the desired properties.

For use in proving both density and the c.c.c. property, we state the following claim, whose proof we defer to the end.

(1) *Given a basis  $\{x_i : i = 1, \dots, n\}$  of a finitely generated pure subgroup  $A_0$  of  $A$ , a basis  $\{y_j : j = 1, \dots, m\}$  of a finitely generated pure subgroup  $B_0$  of  $B$ , and an indexed set  $\{e_{ij} : i = 1, \dots, n, j = 1, \dots, m\}$  of elements of  $\mathbf{Z}$ , there is one and only one  $p \in P$  such that  $A_p = A_0$ ,  $B_p = B_0$  and  $h_p(x_i)(y_j) = e_{ij}$  for all  $i = 1, \dots, n, j = 1, \dots, m$ .*

Assuming this claim, we proceed to prove the density of  $D_a^1$ . Given  $p$ , there is a finitely generated pure subgroup  $A'$  of  $A$  which contains  $A_p$  and  $a$ . Now  $A_p$  is a summand of  $A'$  so we can choose a basis  $\{x_i : i = 1, \dots, n\}$  of  $A'$  which includes a basis  $\{x_i : i = 1, \dots, k\}$  of  $A_p$ ; choose a basis  $\{y_j : j = 1, \dots, m\}$  of  $B_p$ . Then by the claim there is an element  $p'$  of  $P$  such that  $A_{p'} = A'$ ,  $B_{p'} = B_p$  and  $h_{p'}(x_i)(y_j) = h_p(x_i)(y_j)$  for all  $i = 1, \dots, k$ . Clearly  $p'$  extends  $p$  and belongs to  $D_a^1$ . The proof of the density of  $D_b^2$  is similar.

Next we prove that  $P$  is c.c.c. We will make use of the following fact, which is proved in [3, Lemma 7.5]. (It is proved there for strongly  $\aleph_1$ -free groups, there called groups with Chase's condition, but the proof may be adapted for Shelah groups; cf. [4, Theorem 7.1].)

(2) *If  $G$  is a Shelah group of cardinality  $\aleph_1$  and  $\{S_\alpha : \alpha \in \omega_1\}$  is a family of finitely generated pure subgroups of  $G$ , then there is an uncountable subset  $I$  of  $\omega_1$  and a pure free subgroup  $G'$  of  $G$  such that  $S_\alpha \subseteq G'$  for all  $\alpha \in I$ .*

Suppose that  $\{p_\nu = (A_\nu, B_\nu, h_\nu) : \nu \in \omega_1\}$  is a subset of  $P$ . We must prove that there are indices  $\mu \neq \nu$  such that  $p_\mu$  and  $p_\nu$  are compatible. By claim (2), passing to a subset, we can assume that there is a pure free subgroup  $A'$  of  $A$  and a pure free subgroup  $B'$  of  $B$  such that



$A_\nu \subseteq A'$  and  $B_\nu \subseteq B'$  for all  $\nu \in \omega_1$ . (Now we follow the argument in [3] for property (7.1.3).) Choose a basis  $X$  of  $A'$  and a basis  $Y$  of  $B'$ . By density, we can assume that each  $A_\nu$  is generated by a finite subset  $X_\nu$  of  $X$  and each  $B_\nu$  is generated by a finite subset,  $Y_\nu$ , of  $Y$ . Moreover, we can assume that there is a (finite) subset  $T$  of  $X$  (respectively  $W$  of  $Y$ ) which is contained in each  $X_\nu$  (respectively each  $Y_\nu$ ) and is maximal with respect to the property that it is contained in uncountably many  $X_\nu$  (respectively  $Y_\nu$ ). Passing to a subset, we can assume that  $h_\nu(x)(y)$  has a value independent of  $\nu$  for each  $x \in T$  and  $y \in W$ . By a counting argument, we can find  $\nu > 0$  such that  $X_\nu \cap X_0 = T$  and  $Y_\nu \cap Y_0 = W$ . We define a member  $q \in P$  which extends  $p_0$  and  $p_\nu$ , as follows. Let  $A_q = \langle X_0 \cup X_\nu \rangle$  and  $B_q = \langle Y_0 \cup Y_\nu \rangle$ . Clearly, these are pure subgroups of  $A$  (respectively,  $B$ ). Then, by claim (1), there is a homomorphism  $h_q : A_q \rightarrow B_q^*$  such that  $q \geq p_0$  and  $q \geq p_\nu$ . In particular, for  $x \in T$ ,

$$h_q(x)(y) = \begin{cases} \text{the common value} & \text{if } y \in W \\ h_0(x)(y) & \text{if } y \in Y_0 - W \\ h_\nu(x)(y) & \text{if } y \in Y_\nu - W \end{cases}$$

and, for  $x \in X_0 - T$ ,

$$h_q(x)(y) = \begin{cases} h_0(x)(y) & \text{if } y \in Y_0 \\ \text{arbitrary} & \text{if } y \in Y_\nu - W \end{cases}$$

and similarly for  $x \in X_\nu - T$ .

Thus we have shown the existence of  $p_\mu$  and  $p_\nu$ , which are compatible, and it remains to prove claim (1). We will prove uniqueness first. Suppose that, with the notation of (1), there are  $p_1$  and  $p_2$  in  $P$  such that for  $\ell = 1, 2$ ,  $h_{p_\ell}(x_i)(y_j) = e_{ij}$  for all  $i = 1, \dots, n$ ,  $j = 1, \dots, m$ . It suffices to prove that for all  $a \in A_0$ ,

$$h_{p_1}(a)(y_j) = h_{p_2}(a)(y_j)$$

for all  $j = 1, \dots, m$ . Write  $a$  as a linear combination of the basis:  $a = \sum_{i=1}^n d_i x_i$ . By the definition of  $P$

$$\gamma(a) - \sum_{i=1}^n d_i \gamma(x_i) = h_{p_\ell}(a) - \sum_{i=1}^n d_i h_{p_\ell}(x_i)$$

for  $\ell = 1, 2$ . Note that the right-hand side is, by definition, a function on  $B_0$ . Applying both sides to  $y_j \in B_0$ , we obtain that

$$h_{p_\ell}(a)(y_j) = \sum_{i=1}^n d_i e_{ij} + (\gamma(a) - \sum_{i=1}^n d_i \gamma(x_i))(y_j)$$

for  $\ell = 1, 2$ . Since the right-hand side is independent of  $\ell$ , we can conclude the desired identity. For existence, we use the last displayed equation to define  $h_p$  and easily check that it gives an element of  $P$ .  $\square$

By starting with a base model of  $\text{MA} + 2^{\aleph_0} = \aleph_2$  and doing the standard iterated forcing to force all instances of diamond above  $\aleph_1$  one obtains

**Model (3).** A model of  $\text{MA} + 2^{\aleph_0} = \aleph_2$  plus

( $\dagger$ )  $\diamond_{\kappa}(E)$  holds for every regular cardinal  $\kappa > \aleph_1$  and every stationary subset  $E$  of  $\kappa$ .

Just as for Model (2), in this model the tensor product of any two Whitehead groups of arbitrary cardinality is again a W-group.

## REFERENCES

1. T. Becker, L. Fuchs and S. Shelah, *Whitehead modules over domains*, Forum Math. **1** (1989), 53–68.
2. H. Cartan and S. Eilenberg, *Homological algebra*, Princeton University Press, Princeton, 1956.
3. P.C. Eklof, *Whitehead's problem is undecidable*, Amer. Math. Month. **83** (1976), 775–788.
4. ———, *Set theoretic methods in homological algebra and abelian groups*, Les Presses de L'Université de Montréal, Montreal, 1980.
5. P.C. Eklof and A.H. Mekler, *Almost free modules*, revised edition, North-Holland, Amsterdam, 2002.
6. P.C. Eklof and S. Shelah, *The structure of  $\text{Ext}(A, \mathbf{Z})$  and GCH: Possible co-Moore spaces*, Math. Z. **239** (2002), 143–157.
7. L. Fuchs, *Infinite Abelian groups*, vols. I and II, Academic Press, New York, 1970, 1973.
8. M. Huber, *On reflexive modules and abelian groups*, J. Algebra **82** (1983), 469–487.

9. S. Shelah, *Infinite abelian groups, Whitehead problem and some constructions*, Israel J. Math. **18** (1974), 243–256.

10. ———, *Whitehead groups may not be free even assuming CH, II*, Israel J. Math. **35** (1980), 257–285.

MATHEMATICS DEPT, UCI, IRVINE, CA 92697

**Email address:** peklof@pacbell.net

DEPARTMENT OF MATHEMATICS, TULANE UNIVERSITY, NEW ORLEANS, LA 70118

**Email address:** fuchs@tulane.edu

INSTITUTE OF MATHEMATICS, HEBREW UNIVERSITY, JERUSALEM 91904, ISRAEL

**Email address:** shelah@math.huji.ac.il