



ELSEVIER

Annals of Pure and Applied Logic 109 (2001) 117–129

---



---

ANNALS OF  
PURE AND  
APPLIED LOGIC

---



---

www.elsevier.com/locate/apal

# Fallen cardinals

Menachem Kojman<sup>a,\*</sup>, Saharon Shelah<sup>b,1</sup>

<sup>a</sup>*Department of Mathematics and Computer Science, Ben Gurion University of the Negev, Beer Sheva, Israel*

<sup>b</sup>*Institute of Mathematics, The Hebrew University of Jerusalem, Jerusalem 91904, Israel*

---

## Abstract

We prove that for every singular cardinal  $\mu$  of cofinality  $\omega$ , the complete Boolean algebra  $\text{Comp } \mathcal{P}_\mu(\mu)$  contains a complete subalgebra which is isomorphic to the collapse algebra  $\text{Comp Col}(\omega_1, \mu^{\aleph_0})$ . Consequently, adding a generic filter to the quotient algebra  $\mathcal{P}_\mu(\mu) = \mathcal{P}(\mu)/[\mu]^{<\mu}$  collapses  $\mu^{\aleph_0}$  to  $\aleph_1$ . Another corollary is that the Baire number of the space  $U(\mu)$  of all uniform ultrafilters over  $\mu$  is equal to  $\omega_2$ . The corollaries affirm two conjectures of Balcar and Simon. The proof uses pcf theory. © 2001 Elsevier Science B.V. All rights reserved.

MSC: 03G05; 04A20; 04A10; 54D80; 54A25; 54F65

Keywords: Boolean algebra; Forcing; Distributivity; Infinite cardinals; Uniform ultrafilters; Baire number; pcf theory

---

## 1. Introduction

### 1.1. Forcing and distributivity of complete Boolean algebras

Every separative poset  $P$  which may be used as a forcing notion, is embedded as a dense subset of a (unique) complete Boolean algebra, called the *completion of  $P$*  and denoted by  $\text{Comp } P$ . The properties of the forcing extension  $V^P$  of the universe  $V$  of set theory, which is obtained by forcing with  $P$ , are tightly related to the Boolean-algebraic properties of  $\text{Comp } P$ , in particular to the *distributivity* properties of  $\text{Comp } P$ . The least cardinality of a new set in  $V^P$ , for example, is equal to the *distributivity number* of  $\text{Comp } P$ , denoted  $\mathfrak{h}(\text{Comp } P)$ , which should really be called the “non-distributivity number”, since it is the least cardinality of a product of sums which violates distributivity (see [9] for more information). Finer properties of non-distributivity determine

---

\* Corresponding author.

*E-mail addresses:* kojman@math.bgu.ac.il (M. Kojman), shelah@math.huji.ac.il (S. Shelah).

<sup>1</sup> Partially supported by the Israeli Foundation for Basic Science. This paper has number 720 in the Shelah’s list of publications.

which cardinals of  $V$  are preserved and which are collapsed in the extension  $V^P$ . The non-distributivity property which is important for our context is the following:

**Definition 1.1.** A complete Boolean algebra  $B$  is  $(\kappa, \cdot, \lambda)$ -nowhere distributive iff  $B$  contains partitions of unity  $P_\alpha$  for  $\alpha < \kappa$  (namely,  $\sum P_\alpha = 1$  and  $p \wedge q = 0$  for  $p \neq q$  in  $P_\alpha$ ) so that for every  $b \in B - \{0\}$  there exists  $\alpha < \kappa$  so that  $b \wedge p \neq 0$  for  $\geq \lambda$  members  $q \in P_\alpha$ .

Clearly, if  $\lambda_1 < \lambda_2$  and  $B$  is  $(\kappa, \cdot, \lambda_2)$  nowhere distributive, it is also  $(\kappa, \cdot, \lambda_1)$ -nowhere distributive. The systematic study of distributivity in Boolean algebras was pursued by the Czech school of set theory ever since the discovery of Forcing in 1963.

### 1.2. The quotient algebra $\mathcal{P}_\mu(\mu)$

In 1972 Balcar and Vopěnka began the study non-distributivity in quotient algebras  $\mathcal{P}_\kappa(\kappa)$  for infinite cardinals  $\kappa$ . For every infinite cardinal  $\kappa$ , the algebra  $\mathcal{P}_\kappa(\kappa)$  is obtained as the quotient of the power set algebra  $\mathcal{P}(\kappa)$  over the ideal  $[\kappa]^{<\kappa}$  of all subsets of  $\kappa$  whose cardinality is strictly smaller than  $\kappa$ . It was first shown that for every singular cardinal  $\mu$  of countable cofinality the distributivity number of  $\mathcal{P}_\mu(\mu)$  is  $\omega_1$  and that for every cardinal  $\kappa$  of uncountable cofinality the distributivity number of  $\mathcal{P}_\kappa(\kappa)$  is  $\omega$  [6]. The distributivity number of  $\mathcal{P}_\omega(\omega)$  was discussed separately in [5]. The exact nature of distributivity in various  $\mathcal{P}_\kappa(\kappa)$  was addressed in a series of papers [1, 3, 2], usually under additional set theoretic assumptions. The optimal ZFC non-distributivity properties of  $\mathcal{P}_\kappa(\kappa)$  were obtained in [2], from which we quote:

**Theorem 1.2** (Balcar and Simon [2]). 1. For every singular  $\mu$  of countable cofinality  $\mathcal{P}_\mu(\mu)$  is  $(\omega_1, \cdot, \mu^{\aleph_0})$ -nowhere distributive

2. For every singular  $\kappa$  of uncountable cofinality  $\mathcal{P}_\kappa(\kappa)$  is  $(\omega, \cdot, \kappa^+)$ -nowhere distributive.

### 1.3. The collapse algebra $\text{Col}(\omega_1, \mu^{\aleph_0})$

It was in [1] that it was first shown that under certain set theoretic assumptions  $\mathcal{P}_\kappa(\kappa)$  (and some other factor algebras of  $\mathcal{P}(\kappa)$ ) have completions which are isomorphic to suitable *collapse algebras*. Let us introduce collapse algebras. For cardinals  $\kappa < \lambda$ ,  $\kappa$  regular, the poset  $\text{Col}(\kappa, \lambda)$  is the natural  $\kappa$ -complete poset for introducing a function  $\varphi$  from  $\kappa$  onto  $\lambda$ , namely for “collapsing”  $\lambda$  to  $\kappa$ .

$$\text{Col}(\kappa, \lambda) = \{f : \text{for some } \alpha < \kappa, f \text{ is a functions,} \\ \text{dom } f = \alpha \text{ and } \text{ran } f \subseteq \lambda\}. \quad (1)$$

The completion  $\text{CompCol}(\kappa, \lambda)$  is the collapse algebra for  $(\kappa, \lambda)$ . The cardinality of  $\text{Col}(\kappa, \lambda)$  is clearly  $\lambda^{<\kappa}$ , and therefore, since  $\text{Col}(\kappa, \lambda)$  is dense in its completion, the density  $\pi(\text{CompCol}(\kappa, \lambda))$  is equal to  $\lambda^{<\kappa}$ . For each  $\alpha < \kappa$  let  $P_\alpha$  be a maximal

antichain in  $\text{Col}(\kappa, \lambda)$  composed of conditions which decide  $\varphi \upharpoonright \alpha$ . The set  $\{P_\alpha : \alpha < \kappa\}$  (which is, really, a *name* for  $\varphi$ ) is also a witness to the fact that  $\text{Comp Col}(\kappa, \lambda)$  is  $(\kappa, \cdot, \lambda)$ -nowhere distributive. The following characterization of  $\text{Comp Col}(\omega_1, \mu^{\aleph_0})$  for a singular  $\mu$  of countable cofinality is a particular instance of a general characterization theorem for collapse algebras ([3, 1.15]):

**Theorem 1.3.** *Let  $B$  be a complete  $(\omega_1, \cdot, \mu)$  nowhere distributive Boolean algebra containing an  $\aleph_1$ -closed dense subset. If  $\pi(B) = \mu^{\aleph_0}$ , then  $B$  is isomorphic to  $\text{Comp Col}(\omega_1, \mu^{\aleph_0})$ .*

Thus  $\text{Comp Col}(\omega_1, \mu^{\aleph_0})$  is characterized by the existence of an  $\aleph_1$ -closed dense subset, density  $\mu^{\aleph_0}$  and  $(\omega_1, \cdot, \mu)$ -nowhere distributivity (which is a weaker condition than  $(\omega_1, \cdot, \mu^{\aleph_0})$ -nowhere distributivity which is actually satisfied).

#### 1.4. The problem

Let  $\mu$  be a singular cardinal of countable cofinality. Two of the properties which characterize  $\text{Comp Col}(\omega_1, \mu^{\aleph_0})$  hold also in the quotient algebra  $\mathcal{P}_\mu(\mu)$ :  $\aleph_1$ -completeness (easily) and  $(\omega_1, \cdot, \lambda)$ -nowhere distributivity (by Theorem 1.2). Could it be true that  $\text{Comp } \mathcal{P}_\mu(\mu)$  and  $\text{Comp Col}(\omega_1, \mu^{\aleph_0})$  are isomorphic? An old independence result of Baumgartner's rules that out. Baumgartner forced an almost disjoint family in  $\mathcal{P}(\mu)$  of size  $2^\mu > \mu^{\aleph_0}$ , showing thus that it is consistent with ZFC that the cellularity, hence density, of  $\mathcal{P}_\mu(\mu)$  strictly exceeds  $\mu^{\aleph_0}$  ([7, 6.1]). In Baumgartner's model  $\text{Comp } \mathcal{P}_\mu(\mu)$  cannot be isomorphic to  $\text{Comp Col}(\omega_1, \mu^{\aleph_0})$ , whose density is exactly  $\mu^{\aleph_0}$ .

However, if one assumes that  $\mu^{\aleph_0} = 2^\mu$ , it follows trivially that the density of  $\mathcal{P}_\mu(\mu)$  is  $\mu^{\aleph_0}$ , and hence, by the aforementioned characterization of  $\text{Comp Col}(\omega_1, \mu^{\aleph_0})$ , it is isomorphic to  $\text{Comp } \mathcal{P}_\mu(\mu)$ . In particular, denoting by  $\vdash$  provability in ZFC, we have [2]:

$$\mu^{\aleph_0} = 2^\mu \vdash \mu^{\aleph_0} \text{ collapses to } \aleph_1 \text{ in } V^{\mathcal{P}_\mu(\mu)}. \quad (2)$$

What, then, is the precise relation between  $\mathcal{P}_\mu(\mu)$  and  $\text{Col}(\omega_1, \mu^{\aleph_0})$ ? Most importantly, does forcing with  $\mathcal{P}_\mu(\mu)$  *always* collapse  $\mu^{\aleph_0}$  to  $\aleph_1$ ?

Balcar and Simon conjectured in [4] that the answer is “yes”, namely, that the cardinal arithmetic assumption  $\mu^{\aleph_0} = 2^\mu$  could be removed from (2). In the same paper they advance towards an affirmative solution of their conjecture by proving in ZFC that forcing with  $\mathcal{P}_\mu(\mu)$  collapse the  $2^{\aleph_0}$  to  $\omega_1$ . Since for  $\mu < 2^{\aleph_0}$  it holds trivially that  $\mu^{\aleph_0} = 2^{\aleph_0}$ , that proves their conjecture for all countably cofinal singular cardinals  $\mu$  which are below the continuum:

$$\mu < 2^{\aleph_0} \vdash \mu^{\aleph_0} \text{ collapses to } \aleph_1 \text{ in } V^{\mathcal{P}_\mu(\mu)}. \quad (3)$$

Finally, there was the problem of computing the *Baire number* of the space  $U(\mu)$  of all uniform ultrafilters over  $\mu$ . An ultrafilter  $u$  over  $\mu$  is *uniform* if it does not

contain a set of cardinality  $< \mu$ . With the usual topology, in which the basic open sets are  $\hat{p} = \{u \in U(\mu) : p \in u\}$  for  $p \in [\mu]^\mu$ , the space  $U(\mu)$  is a compact Hausdorff space and is therefore not coverable by  $\omega_1$  nowhere-dense sets. The *Baire number* of a space with no isolated points is the least number of nowhere-dense sets needed to cover the space. In [4] it was proved that the Baire number of  $U(\mu)$  is  $\omega_2$  under any of the following assumptions: (i)  $2^{\aleph_0} > \aleph_1$ , (ii)  $2^\mu = \mu^{\aleph_0}$  or (iii)  $2^{\omega_1} = \omega_2$ . It was conjectured that the Baire number of  $U(\mu)$  could be shown to equal  $\omega_2$  in ZFC alone.

### 1.5. The solution

The main result in the present paper determines the precise relation between  $\text{Comp } \mathcal{P}_\mu(\mu)$  and  $\text{Comp Coll}(\omega_1, \mu^{\aleph_0})$ . The collapse algebra is isomorphic to a complete subalgebra of the quotient algebra (Theorem 2.1 below):

$$\vdash \text{Comp Coll}(\omega_1, \mu^{\aleph_0}) \triangleleft \text{Comp } \mathcal{P}_\mu(\mu). \quad (4)$$

This implies that the universe  $V^{\mathcal{P}_\mu(\mu)}$  contains  $V^{\text{Coll}(\omega_1, \mu^{\aleph_0})}$  as a subuniverse. Therefore,

$$\vdash \mu^{\aleph_0} \text{ collapses to } \omega_1 \text{ in } V^{\mathcal{P}_\mu(\mu)}, \quad (5)$$

which proves the conjecture. An easy corollary of (4) is that the Baire number of  $U(\mu)$  is equal to  $\omega_2$  (Theorem 2.15 below).

Balcar and Simon stated in [4] another ZFC conjecture concerning singular cardinals of uncountable cofinality. The authors will present a solution of that conjecture in a sequel paper.

### 1.6. History

Balcar presented this conjecture to the authors during a meeting in Hattingen, Germany, in June of 1999. Shelah then proved, using the Erdős–Rado theorem, that

$$\mu > 2^{\aleph_0} \vdash \mu^{\aleph_0} \text{ collapses to } 2^{\aleph_0} \text{ in } V^{\mathcal{P}_\mu(\mu)}. \quad (6)$$

This affirmed the ZFC conjecture, since (3) and (6) together give (5) ( $2^{\aleph_0} = \mu$  is of course impossible by König's Lemma). In August of 1999 Kojman found a direct ZFC proof of (4) by replacing Shelah's use of the Erdős–Rado theorem (which requires cardinal arithmetic assumptions) by a use of a pcf theorem. This proof is presented below.

### 1.7. Description of the proof

Let  $P = \langle [\mu]^\mu, \leq \rangle$  where, for  $p_1, p_2 \in P$ ,  $p_1 \leq p_2 \Leftrightarrow |p_1 - p_2| < \mu$ . For every  $D \subseteq \mathcal{P}_\mu(\mu)$ ,  $D$  is dense in  $\mathcal{P}_\mu(\mu)$  if and only if  $\bigcup D$  is dense in  $P$  and  $D$  is a filter

in  $\mathcal{P}_\mu(\mu)$  if and only if  $\bigcup D$  is a filter in  $P$ . Therefore,  $G \subseteq \mathcal{P}_\mu(\mu)$  is a generic filter over  $\mathcal{P}_\mu(\mu)$  if and only if  $\bigcup G$  is a generic filter over  $P$ . Hence  $V^{\mathcal{P}_\mu(\mu)} = V^P$ . For convenience, we work with  $P$  rather than with  $\mathcal{P}_\mu(\mu)$ . Let  $\lambda$  denote  $\mu^{\aleph_0}$ .

The main point in finding a complete copy of  $\text{Comp Col}(\omega_1, \lambda)$  inside  $\text{Comp } \mathcal{P}_\mu(\mu)$  is to overcome the large cellularity that  $\text{Comp } P$  may possess, e.g. in Baumgartner's model. This is achieved by forcing only with  $Q \subseteq P$ , which contains all *closed* conditions of  $P$ . It is not hard to verify that  $\text{Comp } Q$  is isomorphic to a complete subalgebra of  $\text{Comp } P$ . Then it is shown that  $\text{Comp } Q \cong \text{Comp Col}(\omega_1, \lambda)$ . To that end one needs to prove that  $\pi(Q) = \lambda$ . This fact is achieved by an old trick: club guessing. Once density is out of the way, it remains to establish  $(\omega_1, \cdot, \mu)$ -nowhere distributivity of  $\text{Comp } Q$ , to facilitate the use of Theorem 1.3 above. Here another pcf tool is used: the Trichotomy Theorem.

### 1.8. Notation and preliminaries

Our notation is mostly standard. One exception is that when the relations  $f <_U g$ ,  $f \leq_U g$  for ordinal functions  $f, g$  where  $U$  an ultrafilter over  $\omega$  is extended to *partial* functions. We recall that if  $P$  and  $Q$  are posets and for some  $P$ -name  $\mathcal{G}$  it holds that  $\Vdash_P$  “ $\mathcal{G}$  is a generic filter over  $Q$ ” and for every  $q \in Q$  there exists  $p \in P$  such that  $p \Vdash q \in \mathcal{G}$ , then  $\text{Comp } Q$  is isomorphic to a complete subalgebra of the  $\text{Comp } P$  via the embedding  $b \mapsto \sum \{p \in P : p \Vdash “b \in \mathcal{G}”\}$ .

The following two theorems from pcf theory will be used:

**Theorem 1.4 (Club Guessing).** *If  $\kappa^+ < \lambda$  and  $\kappa, \lambda$  are regular cardinals, then there exists a sequence  $\bar{c} = \langle c_\delta : \delta < \lambda \wedge \text{cf } \delta = \kappa \rangle$  so that:*

1. *For every  $\delta < \lambda$  with  $\text{cf } \delta = \kappa$ ,  $c_\delta$  is closed and unbounded in  $\delta$  and  $\text{otp } c_\delta = \kappa$ .*
2. *For every club  $E$  of  $\lambda$  there exists  $\delta \in S_\kappa^\lambda$  so that  $c_\delta \subseteq E$ .*

**Theorem 1.5 (The Trichotomy).** *Suppose  $A$  is an infinite set,  $I$  an ideal over  $A$  and  $\lambda > |A|^+$  a regular cardinal. If  $\bar{f} = \langle f_\alpha : \alpha < \lambda \rangle$  is a  $<_I$ -increasing sequence of ordinal functions on  $A$ , then one of the following conditions holds:*

- (Good)  $\bar{f}$  has an exact upper bound  $f$  with  $\text{cf } f(a) > |A|$  for all  $a \in A$ ,
- (Bad) there are sets  $S(a)$  for  $a \in A$  satisfying  $|S(a)| \leq |A|$  and an ultrafilter  $D$  over  $A$  extending the dual of  $I$  so that for all  $\alpha < \lambda$  there exists  $h_\alpha \in \prod S(a)$  and  $\beta < \lambda$  such that  $f_\alpha <_D h_\alpha <_D f_\beta$ .
- (Ugly) there is a function  $g : A \rightarrow \text{On}$  such that letting  $t_\alpha = \{a \in A : f_\alpha(a) > g(a)\}$ , the sequence  $\bar{t} = \langle t_\alpha : \alpha < \lambda \rangle$  does not stabilize modulo  $I$ .

A proof of the club guessing theorem can be found in [17, III, Section 1], [10] or the appendix to [12]. The Trichotomy Theorem is Lemma 3.1 in [17], and a shorter proof of it is available in the appendix to [11].

## 2. The proof

Throughout this section let  $\mu$  be a fixed singular cardinal of countable cofinality, and let  $\mu_n$  be a fixed strictly increasing sequence of regular cardinals with  $\sum \mu_n = \mu$ . Let  $P = \{p \subseteq \mu : |p| = \mu\}$  and for  $p_1, p_2 \in P$  let  $p_1 \leq p_2$  iff  $|p_1 - p_2| < \mu$ . As pointed out in the introduction, forcing with  $P$  is equivalent to adding a generic filter to  $\mathcal{P}_\mu(\mu)$ .

Denote  $\lambda = \mu^{\aleph_0} = |\prod \mu_n|$ .

**Theorem 2.1.**  $\text{Comp Col}(\omega_1, \lambda) \ll \text{Comp } \mathcal{P}_\mu(\mu)$ .

**Proof.** Let  $Q = \{q \in P : q \text{ is closed}\}$ . For a condition  $p \in P$ , let  $\text{acc } p$  be the set of all accumulation points of  $p$ . Clearly,  $\text{acc } p \in Q$

**Lemma 2.2.** *Let  $G \subseteq P$  be a generic filter. Then  $G_1 = \{q \in Q : (\exists p \in G)(\text{acc } p \leq q)\}$  is generic in  $Q$ .*

**Proof.** If  $p_1, p_2 \in G$  then there is some  $p_3 \in G$  so that  $p_3 \subseteq p_1 \cap p_2$ . So  $\text{acc } p_3 \leq \text{acc } p_1, \text{acc } p_2$  and  $\text{acc } p_3 \in \{\text{acc } p : p \in G\}$ . Thus  $G_1$  is closed under finite intersections. Clearly,  $G_1$  is upwards closed. Thus  $G_1$  is a filter.

Suppose that  $D \subseteq Q$  is dense and downwards closed. Let  $p \in P$  be arbitrary, and consider  $q = \text{acc } p$ . Let  $q_1 \leq q$  be chosen in  $D$ .

For  $\alpha \in q_1$  define  $\beta_\alpha = \min(p - (\alpha + 1))$ , and let  $p_1 = \{\beta_\alpha : \alpha \in q_1\}$ .  $q_2 = \text{acc } p_1 \subseteq \text{acc } q_1 \leq q_1$  so there is some  $p_1 \leq p$  with  $\text{acc } p_1 \in D$ .  $\square$

By this lemma it follows that

$$\text{Comp } Q \ll \text{Comp } P. \quad (7)$$

We aim now to show that

$$\text{Comp } Q \cong \text{Comp Col}(\omega_1, \lambda). \quad (8)$$

First, we shall see that  $\pi(Q) = \lambda$ .

Let  $q \in Q$  be arbitrary. Let  $a(p) = \{n : p \cap [\mu_n, \mu_{n+1}) \neq \emptyset\}$  and let  $\{m_n : n < \omega\}$  be the increasing enumeration of  $a(p)$ .

**Definition 2.3.** A condition  $q \in Q$  is *normal* if it satisfies

$$\text{otp}[q \cap [\mu_{m_n}, \mu_{m_{n+1}})] = \mu_n + 1. \quad (9)$$

**Lemma 2.4.** *The set of normal conditions is dense in  $Q$ .*

**Proof.** Given a condition  $q$ , let  $m_n$  be the least so that  $q \cap [\mu_{m_n}, \mu_{m_{n+1}}) \geq \mu_n$  and choose  $c_n \subseteq q \cap [\mu_{m_n}, \mu_{m_{n+1}})$  of order type  $\mu_n + 1$ . Let  $q' = \bigcup_n c_n$ . Thus  $q' \leq q$ , and is of the form (9) above.  $\square$

Let  $M$  be a fixed elementary submodel of  $\langle H(\Omega), \in \rangle$  for a sufficiently large regular cardinal  $\Omega$  so that  $\mu \subseteq M$ ,  $[M]^{\aleph_0} \subseteq M$  and the cardinality of  $M$  is  $\lambda$ . Let  $Q^M = Q \cap M$ . Clearly,  $|Q^M| = \lambda$ .

**Lemma 2.5.**  $Q^M$  is dense in  $Q$ .

**Proof.** Let  $q$  be a condition in  $Q$  and assume, without loss of generality, that it is normal. Let  $c_n = q \cap [\mu_{m_n}, \mu_{m_n+1})$ .

**Claim 2.6.** For every  $n$ , there exists a closed subset of  $c_{n+2}$  of order type  $\mu_n + 1$  which belongs to  $M$ .

**Proof.** Let  $\gamma = \sup c_{n+2}$ . In  $M$ , fix an increasing and continuous function  $f : \mu_{n+2} \rightarrow \gamma$  with  $\sup \text{ran} f = \gamma$ . Let  $E = \{i < \mu_{n+2} : f(i) \in c_{n+2}\}$ . Thus  $E \subseteq \mu_{n+2}$  is a club in  $\mu_{n+2}$ .

The club  $E$  itself may not belong to  $M$  (because  $c_{n+2}$  may not belong to  $M$ ). But since  $\mu_n^+ < \mu_{n+2}$  and both (regular) cardinals belong to  $M$ ,  $M$  contains some club guessing sequence  $\langle c_\delta : \delta < \mu_{n+2} \wedge \text{cf} \delta = \mu_n \rangle$  by the club guessing Theorem 1.4 above. Thus there is some  $\delta < \mu_{n+2}$  so that  $c_\delta \subseteq E$ . Clearly,  $c_\delta \in M$ . Since  $f \in M$ , also  $\text{ran}(f \upharpoonright c_\delta) \cup \{\sup \text{ran}(f \upharpoonright c_\delta)\} \subseteq E$  belongs to  $M$ , and is a closed subset of  $c_{n+2}$  of order type  $\mu_n + 1$ .  $\square$

Using the claim, choose, for every  $n$ , a closed set  $b_n$  so that  $b_n \subseteq c_{n+2}$ ,  $\text{otp} b_n = \mu_n + 1$  and  $b_n \in M$ . Since  $M$  is closed under countable sequences,  $q' = \bigcup b_n \in Q^M$ , and clearly  $q' \leq q$  is a normal condition in  $Q$ .  $\square$

This has established that  $\pi(Q) = \lambda$ .

We need the following simple fact about  $Q$  and  $Q^M$ :

**Fact 2.7.**  $Q$  is  $\aleph_1$ -complete and  $Q^M$  is  $\aleph_1$ -complete.

**Proof.** Suppose that  $q_0 \geq q_1 \geq \dots$  is a decreasing sequence of conditions in  $Q$ . By induction on  $n$ , let  $m_n$  be chosen so that  $\text{otp}[q_n \cap [\mu_{m_n}, \mu_{m_n+1})] > \sum_{i < n} |q_n - q_i|^+$ , and choose a closed subset  $c_{n+1}$  of  $\bigcap_{i \leq n} q_i \cap [\mu_{m_n}, \mu_{m_n+1})$  with  $\text{otp} c_n = \mu_n + 1$ . The condition  $\bigcup c_n$  belongs to  $Q$  and  $q \leq q_n$  for all  $n$ . If each  $q_n$  belongs to  $M$  then the sequence itself belongs to  $M$  because  $M$  is closed under taking  $\omega$ -sequences, and hence some  $q$  which satisfies  $q \leq q_n$  for all  $n$  belongs to  $M$ , by elementarity. (Alternatively, one can do the induction for proving completeness of  $Q$  inside  $M$ ).  $\square$

Thus,  $\text{Comp} Q$  contains an  $\aleph_1$ -complete dense set of size  $\lambda$ . To prove (8) from Theorem 1.3 it remains to show that  $\text{Comp} Q$  is  $(\omega_1, \cdot, \mu)$ -nowhere distributive. For this purpose we inspect the generic cut which  $Q$  creates in  $\prod \mu_n / U$ , where  $U$  is the generic ultrafilter over  $\omega$  which forcing with  $Q$  introduces.

**Fact 2.8.** *Suppose  $G \subseteq \mathcal{Q}$  is a generic filter. Then  $\{a(q) : q \in G\}$  is an ultrafilter over  $\omega$ .*

**Proof.** If  $q_1 \leq q_2$  are normal conditions, then  $a(q_1) \setminus a(q_2)$  is finite. Thus  $a : \{q \in \mathcal{Q} : q \text{ is normal}\} \rightarrow \mathcal{P}(\omega)$  is an order preserving map onto  $\langle \mathcal{P}(\omega), \subseteq^* \rangle$ . Furthermore, if  $t \leq a(q)$ , then there is some  $q' \leq q$  such that  $a(q') = t$ . Therefore, the image of a generic  $G \subseteq \mathcal{Q}$  under  $a$  is an ultrafilter over  $\omega$ .  $\square$

Given a normal condition  $q \in \mathcal{Q}$ , define the following two functions on  $a(q)$  by letting, for each  $n \in a(q)$ ,

$$\chi_q^+(n) = \sup[q \cap [\mu_n, \mu_{n+1})],$$

$$\chi_q^-(n) = \min[q \cap [\mu_n, \mu_{n+1})].$$

The set of conditions  $q$  for which  $\chi_q^+ \in \prod \mu_n$  is clearly dense in  $\mathcal{Q}$ , so we always assume that  $\chi_q^+ \in \prod \mu_n$ .

Since  $\chi_q^-(n) < \chi^+ + q(n)$  for every  $n \in a(q)$ , and  $q \Vdash "a(q) \in U"$ , it follows that  $q \Vdash "\chi_q^- <_U \chi_q^+"$ .

Let  $G \subseteq \mathcal{Q}$  be a generic filter. Define

$$D_0^+ = \{\chi_q^+ : q \in G \text{ and } q \text{ is normal}\},$$

$$D_0^- = \{\chi_q^- : q \in G \text{ and } q \text{ is normal}\}.$$

Now for each normal  $q \in \mathcal{Q}$ ,  $q \Vdash "\chi_q^+ \in D_0^+ \wedge \chi_q^- \in D_0^-"$ .

Let

$$D^+ = \left\{ f \in \prod \mu_n : (\exists g \in D_0^+) (g \leq_U f) \right\},$$

$$D^- = \left\{ f \in \prod \mu_n : (\exists g \in D_0^-) (f \leq_U g) \right\}.$$

**Lemma 2.9.** *Suppose that  $f \in \prod \mu_n$  and  $q \in \mathcal{Q}$  is a normal condition. Then there exists a normal condition  $q' \leq q$  so that*

$$q' \Vdash \chi_{q'}^+ <_U f \vee f <_U \chi_{q'}^-.$$

**Proof.** If for some infinite set  $B \subseteq A(q)$ ,  $m_n \in a(q) \Rightarrow f(m_n) < \chi_q^+(m_n)$  then  $q' := \bigcup_{m_n \in B} c_n - (f(m_n) + 1)$  is a normal condition and for all  $n \in a(q')$  it holds that  $\chi_{q'}^-(n) > f(n)$ . Since  $q' \Vdash a(q') \in U$ , the second alternative holds for  $q'$ .

If  $\{n \in a(q) : f(n) < \chi_q^+(n)\}$  is finite, let  $n_0$  be fixed so that for every  $n > n_0$  it holds that  $\sup c_n \leq f(n)$  and let, for  $n > n_0$ ,  $b_n \subseteq c_n$  be the initial segment of  $c_n$  whose order type is  $\mu_{n-1} + 1$ . Now  $\bigcup b_n$  is a normal condition and  $q' \Vdash \chi_{q'}^+ <_U f$ .  $\square$

Since  $\mathcal{Q}$  is  $\aleph_1$ -complete, no new members are added to  $\prod \mu_n$  after forcing with  $\mathcal{Q}$ . Therefore, by Lemma 2.9, it holds that  $D^-$  is a lower half of a Dedekind cut in  $\prod \mu_n/U$  whose upper half is  $D^+$ ; that  $D^-$  has no last element and that  $D^+$  has no first element. By the definition of  $D^-$ , it is clear that  $D_0^-$  is cofinal in  $(D^-, <_U)$ . Furthermore, if  $\{f_i : i < \omega\}$  is a set of functions,  $q \in \mathcal{Q}$  and for all  $i < \omega$  it holds that  $q \Vdash f_i \in D^-$  then by iterated use of Lemma 2.9 and  $\aleph_1$ -completeness there exists  $q' \leq q$  so that  $q' \Vdash \bigwedge_i f_i <_U \chi_{q'}^-$ . As a consequence, the cofinality of  $D^-$  is uncountable.

We shall need the following strengthening of Lemma 2.9, which says that the generic cut  $(D^-, D^+)$  is not trapped by any product of countable sets.

**Lemma 2.10.** *Suppose  $A_n \subseteq [\mu_n, \mu_{n+1})$  is a countable set for each  $n < \omega$ , and  $q \in \mathcal{Q}$  is a normal condition. Then there is a condition  $q' \leq q$  in  $\mathcal{Q}$  so that for every  $n \in a(q')$  it holds that  $A_n \cap (\chi_{q'}^-, \chi_{q'}^+) = \emptyset$ .*

**Proof.** Let  $\varepsilon_n < \omega_1$  be the order type of  $A_n$  and let  $\langle \alpha_i^n : i < \varepsilon_n \rangle$  be the increasing enumeration of  $A_n$ . Partition  $[\mu_n, \mu_{n+1})$  to the intervals  $[\mu_n, \alpha_0^n)$ ,  $\{[\alpha_i^n, \alpha_{i+1}^n) : i + 1 < \varepsilon_n\}$  and  $[\sup A_n, \mu_{n+1})$ . For every  $n > 0$ , choose an interval  $I_n$  in the partition of  $[\mu_n, \mu_{n+1})$  so that  $|I_n \cap q| = \mu_n$ , and let  $c_n \subseteq (I_n \cap q)$  be closed of order type  $\mu_{n-1} + 1$ . Let  $q' = \bigcup I_n \cap q$ .  $\square$

**Lemma 2.11.** *The cofinality of  $D^-$  is  $\omega_1$ .*

**Proof.** We have seen that  $\text{cf}(D^-) > \aleph_0$ . Suppose now, to the contrary, that  $\kappa > \aleph_1$  is regular and that  $q \Vdash \bar{f} = \langle f_i : i < \kappa \rangle$  is  $<_U$  increasing and cofinal in  $D^-$ . The Trichotomy Theorem applies to  $\bar{f}$ , but:

The third condition (“Ugly”) cannot hold, since  $U$  is an ultrafilter.

The first condition (“Good”) cannot hold, because in  $D^+$  there is no first element.

Let us see now that the second condition (“Bad”) cannot hold either. Suppose that  $q \Vdash \langle A_n : n < \omega \rangle$ , witnesses ‘Bad’ for  $\bar{f}$ . Then,

$$q \Vdash (\forall \alpha < \kappa) \left( \exists h \in \prod A_n \right) (\exists \beta < \kappa) [f_\alpha <_U h <_U f_\beta]. \quad (10)$$

By  $\aleph_1$ -completeness, we may assume that  $\langle A_n : n < \omega \rangle$  and each  $A_n$  belong to the ground model. By Lemma 2.10 there is a condition  $q' \leq q$  so that for all  $n \in a(q)$ ,

$$A_n \cap (\chi_{q'}^-(n), \chi_{q'}^+(n)) = \emptyset. \quad (11)$$

Since  $q$  forces that  $\bar{f}$  is cofinal in  $D^-$  and  $q' \Vdash \chi_{q'}^- \in D^- \wedge \chi_{q'}^+ \in D^+$ , there is some  $\alpha < \kappa$  and  $q'' \leq q'$  so that

$$q'' \Vdash \chi_{q'}^- <_U f_\alpha <_U \chi_{q'}^+. \quad (12)$$

By strengthening  $q''$  we may assume that for some  $\beta < \kappa$  and  $h \in \prod A_n$ ,

$$q'' \Vdash \chi_{q'}^- <_U f_\alpha <_U h <_U f_\beta <_U \chi_{q'}^+. \quad (13)$$

So there is some  $n$  (in fact, infinitely many) so that

$$\chi_{q'}^-(n) < f_\alpha(n) < h(n) < f_\beta(n) < \chi_{q'}^+(n). \quad (14)$$

This is a contradiction to (11), since  $h(n) \in A_n$ .

Thus, the cofinality of  $D^-$  is at least  $\omega_1$  and no more than  $\omega_1$ ; so it is exactly  $\omega_1$ .  $\square$

Since  $\text{cf}(D^-, <_U) = \omega_1$  and  $D_0^-$  is cofinal in  $(D^-, <_U)$ , using  $\aleph_1$ -completeness of  $Q$  it is easy to find a sequence of conditions  $\langle q(i) : i < \omega_1 \rangle \subseteq G$  such that  $i < j < \omega_1 \Rightarrow q(i) \geq q(j)$  and  $\langle \chi_{q(i)}^- : i < \omega_1 \rangle$  is ( $<_U$ -increasing and) cofinal in  $(D^-, <_U)$ . Fix a  $Q$ -name  $\underline{q}$  for such a sequence. Observe that if  $q_1, q_2$  are incompatible, then  $\Vdash \neg(q_1 \in \text{ran } \underline{q} \wedge q_2 \in \text{ran } \underline{q})$ , since any two conditions in  $\text{ran } \underline{q}$  are compatible.

**Lemma 2.12.** *For every  $q \in Q$  there is a set  $\{q'_\alpha : \alpha < \lambda\}$  of pairwise incompatible conditions below  $q$ , so that for each  $\alpha < \lambda$  there is  $q'_\alpha \leq q$  and  $i(\alpha)$  so that  $q'_\alpha \Vdash \underline{q}(i(\alpha)) = q''_\alpha$ , and  $\{q'_\alpha : \alpha < \lambda\}$  are pairwise incompatible.*

**Proof.** Let  $q \in Q$  be a normal condition, and let  $c_n = q \cap [\mu_{m_n}, \mu_{m_n+1})$ .  $c_n$  is a closed set of order type  $\mu_n$ . For each  $n$  let  $b_n$  be the initial segment of  $c_{n+1}$  of order type  $(\prod_{i \leq n} \mu_i, <_{\text{lex}})$ , the lexicographic ordering of all sequences  $(x_0, x_1, \dots, x_n)$  in the product  $\mu_0 \times \mu_1 \times \dots \times \mu_n$ . Identify each member in  $b_n$  with the sequence in  $\prod_{i \leq n} \mu_i$  it corresponds to via the order isomorphism, and define a projection  $\pi_{m,n} : b_n \rightarrow b_m$  for  $m < n$  by mapping a sequence of length  $n$  to its initial segment of length  $m$ . The inverse limit of this system is the set of all functions  $g \in \prod b_n$  with the property that for all  $m < n$ ,  $\pi_{m,n}g(n) = g(m)$ . Denote this set of functions by  $L \subseteq \prod b_n$ .

Choose a set of  $\lambda$  different functions  $\langle g_\alpha : \alpha < \lambda \rangle \subseteq L \cap \prod \text{acc } b_n$  and for each  $\alpha$  let  $g'_\alpha(n) = g_\alpha(n) + 1$ . Let  $q_\alpha = \bigcup_{n > 0} b_n \cap [g_\alpha(m_n), g'_\alpha(m_n))$ . Thus each  $q_\alpha$  is a condition below  $q$ . Furthermore, if  $\alpha \neq \beta$  then from some point  $n_0$  on, either  $\chi_{q_\alpha}^+(n) < \chi_{q_\beta}^-(n)$  or  $\chi_{q_\beta}^+(n) < \chi_{q_\alpha}^-(n)$ . Thus  $\{q_\alpha : \alpha < \lambda\}$  is a set of pairwise incompatible conditions below  $q$ .

For each  $\alpha < \lambda$ ,

$$q_\alpha \Vdash \chi_{q_\alpha}^- \in D^- \wedge \chi_{q_\alpha}^+ \in D^+,$$

$$q_\alpha \Vdash \langle (\exists i < \omega_1)(\forall j < \omega_1)[i < j \Rightarrow \chi_{q_\alpha}^- <_U \chi_{q(j)}^-] \rangle.$$

Fix  $q'_\alpha \leq q_\alpha$  so that for some  $i(\alpha) < \omega_1$  and  $q''_\alpha \leq q_\alpha$ ,  $q'_\alpha \Vdash \underline{q}(i(\alpha)) = q''_\alpha$ . For  $\alpha < \beta < \lambda$ , since  $q''_\alpha \leq q'_\alpha \leq q_\alpha$ ,  $q''_\beta \leq q'_\beta \leq q_\beta$  and  $q_\alpha, q_\beta$  are incompatible,  $q'_\alpha$  is incompatible with  $q'_\beta$  and  $q''_\alpha$  is incompatible with  $q''_\beta$ .  $\square$

Fix, for each  $i < \omega_1$ , a maximal antichain  $P_i \subseteq Q$  of conditions that decides  $\dot{q}(i)$ .

**Claim 2.13.** *For every condition  $q \in Q$  there exists some  $i < \omega_1$  so that  $q$  is compatible with  $\geq \mu$  members of  $P_i$ .*

**Proof.** Let  $q \in Q$  be an arbitrary condition. By Lemma 2.12 there are  $\lambda$  pairwise incompatible conditions  $\{q''_\alpha : \alpha < \lambda\}$  below  $q$ , each of which is forced to be  $\dot{q}(i(\alpha))$  for some  $i(\alpha) < \omega_1$ , by some extension  $q'_\alpha \leq q$ , and  $\{q'_\alpha : \alpha < \lambda\}$  are pairwise incompatible extensions of  $q$ . Since  $\lambda > \mu$ , there is necessarily some fixed  $i < \omega_1$  so that  $|\{\alpha < \lambda : i(\alpha) = i\}| > \mu$ . Since different  $q'_\alpha, q'_\beta$  in this set force different values for  $\dot{q}(i)$ , they cannot be compatible with the same member of  $P_i$ . Thus  $q$  is compatible with  $\geq \mu$  members of  $P_i$ .  $\square$

The last claim established  $(\omega_1, \cdot, \mu)$ -nowhere distributivity of  $\text{Comp } Q$ . By Theorem 1.3  $\text{Comp Col}(\omega_1, \lambda) \cong \text{Comp } Q$ , and since  $\text{Comp } Q < \text{Comp } \mathcal{P}_\mu(\mu)$ , the proof is complete.

**Corollary 2.14.**  $V^{\mathcal{P}_\mu(\mu)} \models |\lambda| = \aleph_1$ .

**Proof.** Since  $\text{Comp Col}(\omega_1, \lambda)$  is a complete subalgebra of  $\text{Comp } \mathcal{P}_\mu(\mu)$ , the universe  $V^{\text{Col}(\omega_1, \lambda)}$  is contained in  $V^{\mathcal{P}_\mu(\mu)}$ . Therefore, there is an onto function  $\varphi : \omega_1 \rightarrow \lambda$  in  $V^{\mathcal{P}_\mu(\mu)}$ . Since  $\mathcal{P}_\mu(\mu)$  is  $\omega_1$ -complete,  $\omega_1$  is preserved in  $V^{\mathcal{P}_\mu(\mu)}$ . Thus, the cardinality of  $\lambda$  in  $V^{\mathcal{P}_\mu(\mu)}$  is  $\aleph_1$ .  $\square$

**Corollary 2.15.** *For every singular cardinal  $\mu$  with  $\text{cf } \mu = \aleph_0$  the Baire number of  $U(\mu)$ , the space of uniform ultrafilters over  $\mu$ , is equal to  $\omega_2$ .*

**Proof.** By Theorem 2.1 there exists a dense subset of  $Q$  which is isomorphic to the dense subset  $D = \{f : \exists i < \omega_1 [f : (i+1) \rightarrow \lambda]\}$  of  $\text{Col}(\omega_1, \lambda)$ , namely, there are conditions  $\{q_f : f \in D\} \subseteq Q$  so that  $q_f \leq q_g \Leftrightarrow g \subseteq f$ .

Let  $W_{i,\alpha} = \{q_f : \text{dom } f = i+1 \wedge f(i) = \alpha\}$ . Define  $V_{i,\alpha} = \{u \in U(\mu) : (\exists p \in u)(\exists q_f \in W_{i,\alpha})[\text{acc } p \subseteq q_f]\}$ . It should be clear that for each  $i < \omega_1$  the collection  $\{V_{i,\alpha} : \alpha < \lambda\}$  is a family of pairwise disjoint (non-empty) open sets in  $U(\mu)$ .

For each  $\beta < \omega_2$  define  $O_\beta = \bigcup \{V_{i,\beta} : i < \omega_1, \beta \leq \alpha < \omega_2\}$ . Since  $O_\beta$  is a union of open sets in  $U(\mu)$ , it is open. We show that  $O_\beta$  is also dense in  $U(\mu)$  for each  $\beta < \omega_2$ . Let  $\beta < \omega_2$  be arbitrary and let  $\hat{p}$  for  $p \in P$  be an arbitrary basic open set in  $U(\mu)$ . By density of  $\{q_f : f \in D\}$  there exists some  $f \in \text{Col}(\omega_1, \lambda)$  with domain  $i+1 < \omega$  so that  $q_f \leq \text{acc } p$ . By 2.2, there exists some  $p' \leq p$  so that  $\text{acc } p' \leq q_{f \cup (i+1, \beta)}$ . Thus  $\hat{p}' \subseteq \hat{p} \cap O_\beta$ .

Finally, we show that  $\bigcap \{O_\beta : \beta < \omega_2\} = \emptyset$ . Let  $u \in U(\mu)$  be arbitrary and let  $\alpha(u) = \sup\{\alpha < \omega_2 : (\exists i < \omega_1)(u \in V_{i,\alpha})\}$ . Since for every  $u \in U(\mu)$  and  $i < \omega_1$  there is at most one  $\alpha < \omega_2$  for which  $u \in V_{i,\alpha}$ ,  $\alpha(u) < \omega_2$ . Now  $u \notin O_{\alpha(u)+1}$ .

Thus, by passing to the complements of  $O_\beta$ ,  $\beta < \omega_2$ , it is seen that  $U(\mu)$  is coverable by  $\omega_2$  nowhere-dense sets. Since it is known [4] that  $U(\mu)$  cannot be covered by fewer than  $\omega_2$  nowhere-dense sets, its Baire number is equal to  $\omega_2$ .  $\square$

### 3. Concluding remarks

We first remark that the part of the proof between Lemmas 2.7 and Lemma 2.12 can be applied verbatim to  $P$  instead of to  $Q$  to show that  $\text{Comp } \mathcal{P}_\mu(\mu)$  is  $(\omega_1, \cdot, \mu^{\aleph_0})$ -nowhere distributive, and constitutes thus an alternative ZFC proof of  $(\omega_1, \cdot, \mu^{\aleph_0})$ -nowhere distributivity of  $\mathcal{P}_\mu(\mu)$  from the Trichotomy theorem.

Next, we remark that Corollary 2.14 can be derived directly, without invoking Theorem 1.3, as follows: fix a 1–1 function  $f : Q^M \rightarrow \lambda$  and apply Lemma 2.12 to  $Q^M$ , observing that the set  $\{q''_\alpha : \alpha < \lambda\}$  belongs to  $M$ . Now fix a function  $h : \lambda \rightarrow \lambda$  such that for every  $A \in [\lambda]^\lambda \cap M$ ,  $\text{ran}(h \upharpoonright A) = \lambda$ . The function  $h \circ \underline{q}$  is a collapsing function by a simple density argument.

We devote now a few words to the role of pcf theory in this proof and in several other proofs. Pcf theory was developed to provide bounds on powers of strong limit singular cardinals, or, better, on the covering numbers of singular cardinals. The most well known discovery of the theory is that poset  $\langle \mathcal{P}_{\aleph_0}(\aleph_\omega), \supseteq \rangle$  of countable subsets of  $\aleph_\omega$  ordered by reverse inclusion has a dense subset of size  $< \aleph_{\omega_4}$ . In other words: the cardinality of this poset may be arbitrarily large, but its density is bounded.

From the point of view of pcf theory, powers of regular cardinals are the “soft” part of cardinal arithmetic, which envelopes the hard “skeleton” of powers of singulars that pcf theory addresses—the revised power set function  $\text{pp}$ . To read more about this philosophy the reader is referred to [17] (especially the analytical index, Section 14), [16] and [10].

The proof above is yet another example of the same theme: a complete subalgebra of density  $\mu^{\aleph_0}$  is uncovered inside  $\text{Comp } \mathcal{P}_\mu(\mu)$ , whose own density may be  $2^\mu > \mu^{\aleph_0}$  in case the power function at regular cardinals assumes large values. The powers of regular uncountable cardinals may be “peeled off” from  $\text{Comp } \mathcal{P}_\mu(\mu)$  by the club-guessing technique to get to the “skeleton”  $\text{Comp Col}(\omega_1, \mu^{\aleph_0})$ .

Pcf methods are used also in other contexts to show that various structures on the power set of a singular cardinal contain “skeletons” of bounded cardinality. We quote the example [13] of a Dowker subspace of cardinality  $\aleph_{\omega+1}$  inside Rudin’s Dowker space [15], whose cardinality is  $(\aleph_\omega)^{\aleph_0}$ .

Pcf techniques were used for studying collapses of cardinals by Cummings [8] (see also [11]).

Lastly, we remark that while the role of closed unbounded subsets of *regular* cardinals in combinatorial set theory is so central that one could not imagine uncountable combinatorics without them, the proof above shows that also closed subsets of a *singular* cardinal may be sometimes useful.

## Acknowledgements

The authors wish to thank Bohuslav Balcar for presenting this problem to them during the ESF meeting in Hattingen, June 1999, and take the opportunity to thank Rüdiger Göbel and Simone Pabst for their wonderful work in organizing this meeting. The result in this paper was presented by Kojman in the last of five lectures on pcf theory which were delivered at the Winter School on General Topology held in the Czech republic in January 2000. His thanks for the hospitality and for the opportunity to present this material to an interested audience are extended here to the organizers.

## References

- [1] B. Balcar, F. Franěk, Completion of factor algebras of ideal, *Proc. Amer. Math. Soc.* 100 (2) (1987) 205–212.
- [2] B. Balcar, P. Simon, On collections of almost disjoint families, *Comment. Math. Univ. Carolinae* 29 (4) (1988) 631–646.
- [3] B. Balcar, P. Simon, Disjoint refinements, in: *Handbook of Boolean Algebras*, eds. J.D. Monk, R. Bonnet, vol. 2. North-Holland, Amsterdam, 1989, pp. i–xx and 313–716.
- [4] B. Balcar, P. Simon, Baire number of the space of uniform ultrafilters, *Israel J. Math.* 92 (1995) 263–272.
- [5] B. Balcar, J. Pelant, P. Simon, The space of ultrafilters on  $\mathbb{N}$  covered by nowhere dense sets, *Fund. Math.* 110 (1980) 11–24.
- [6] B. Balcar, P. Vopěnka, On Systems of almost disjoint sets, *Bull. Acad. Polon. Sci. Ser. Sci Math* 20 (1972) 421–424.
- [7] J.E. Baumgartner, Almost-disjoint sets, the dense set problem and the partition calculus, *Ann. Math. Logic* 9 (4) (1976) 401–439.
- [8] J. Cummings, Collapsing successors of singulars, *Proc. Amer. Math. Soc.* 125 (1997) 2703–2709.
- [9] T. Jech, Distributive laws, in: *Handbook of Boolean Algebra*, eds. J.D. Monk, R. Bonnet, vol. 2, North-Holland, Amsterdam, 1989, pp. i–xx and 313–716.
- [10] M. Kojman, The abc of pcf, *Circulated Notes*, 1995.
- [11] M. Kojman, Exact upper bounds and their uses in set theory, *Ann. Pure Appl. Logic* 92 (3) (1998) 267–282.
- [12] M. Kojman, S. Shelah, Non-existence of universal orders in many cardinals, *J. Symbolic Logic* 57 (1992) 875–891.
- [13] M. Kojman, S. Shelah, A ZFC Dowker space in  $\aleph_{\omega+1}$ : an application of PCF theory to topology, *Proc. Amer. Math. Soc.* 126 (8) (1998) 2459–2465.
- [14] M. Kojman, S. Shelah, The condition in the trichotomy theorem is optimal, *Arch. Math. Logic*, in press.
- [15] M.E. Rudin, A normal space  $X$  for which  $X \times I$  is not normal, *Fund. Math.* 73 (1971) 179–186.
- [16] S. Shelah, Cardinal arithmetic for skeptics, *Bull. Amer. Math. Soc. (N.S.)* 26 (1992) 197–210.
- [17] S. Shelah, *Cardinal Arithmetic*, Oxford Logic Guides, vol. 29, Oxford University Press, 1994.